1 **Diversity for commonality in the evolutionary changes of the reduced**
2 **genome to regain the growth fitness**
3

4 Kenya Hitomi[1], Yoichiro Ishii[1], Bei-Wen Ying[1,*]
5
[1]School of Life and Environmental Sciences, University of Tsukuba, Tennodai 1-1-1,
Tsukuba, Ibaraki 305-8572, Japan
[*]Correspondence: ying.beiwen.gf@u.tsukuba.ac.jp

Abstract

As the genome encodes the information crucial for cell growth, a sizeable genomic

deficiency often causes a significant decrease in growth fitness. Whether and how the

decreased growth fitness caused by genome reduction could be compensated by

evolution was investigated here. Experimental evolution with an *Escherichia coli* strain

carrying a reduced genome was conducted in multiple lineages for approximately 1,000

generations. The growth rate, which was largely declined due to genome reduction, was

considerably recovered, associated with the improved carrying capacity. Genome

mutations accumulated during evolution were significantly varied across the

evolutionary lineages and were randomly localized on the reduced genome.

Transcriptome reorganization showed a common evolutionary direction and conserved

the chromosomal periodicity, regardless of highly diversified gene categories, regulons,

and pathways enriched in the differentially expressed genes. Genome mutations and

transcriptome reorganization caused by evolution, which were found to be dissimilar to

those caused by genome reduction, must have followed divergent mechanisms in

individual evolutionary lineages. Gene network reconstruction successfully identified

three gene modules functionally differentiated, which were responsible for the

evolutionary changes of the reduced genome in growth fitness, genome mutation, and

gene expression, respectively. The diversity in evolutionary approaches improved the

growth fitness associated with the homeostatic transcriptome architecture as if the

evolutionary compensation for genome reduction was like all roads leading to Rome.

Keywords: genome reduction, experimental evolution, growth fitness, transcriptome,

genome mutation, chromosomal periodicity, gene expression, network analysis

Introduction

The genome encodes the information for cell growth, and its size is likely the evolutionary consequence [1, 2, 3]. To determine the essential genomic sequence of modern cells, removing redundant DNA sequences from the wild-type genome in bacteria, so-called genome reduction, has been challenged to a large extent [4, 5, 6, 7]. These efforts have been made to discover the minimal genetic requirement for a free-living organism growing under the defined conditions [8, 9, 10]. It resulted in the finding of the coordination of genome with cell growth, i.e., genome reduction significantly decreased the growth rate of *Escherichia coli (E. coli)* cells independent of culture media or growth forms [11, 12, 13]. Slow growth or fitness decline was commonly observed in the genetically reduced [13, 14] and chemically synthesized genomes [10, 15].

The growth decrease could be recovered by experimental evolution. Growth fitness generally represents the adaptiveness of the living organism to the defined environment [16]. Although the reduced genomes somehow showed differentiated evolvability compared to the wild-type genomes [17, 18, 19], their evolutionary adaptation to the environmental changes has been successfully achieved [20, 21, 22]. Experimental evolution under the defined culture condition successfully increased the decreased growth rates of the reduced genomes [17, 20, 21] and fastened the slow-growing synthetic genome [23]. The evolutionary rescue of the growth rate must be associated with the changes in genomic sequence and gene expression benefited for the growth fitness, as what happened in nature adaptive evolution [24, 25, 26]. Our previous study showed that the decreased growth rate caused by the absence of a sizeable genomic sequence could be complemented by introducing the mutations elsewhere in the genome [20, 21]. Additionally, the changes in growth rate caused by either genome reduction or experimental evolution were dependent on the genome size but not the specific gene function [20].

A genome-wide understanding of the evolutionary compensated fitness increase of the reduced genome is required. The experimental evolution compensated for the genome reduction-mediated growth changes were considered stringently related to transcriptome reorganization. Our previous studies found the conserved features in transcriptome reorganization, although the gene expression patterns were significantly disturbed by either genome reduction or experimental evolution [27, 28, 29]. It indicated that

experimental evolution rescued the growth rate disturbed by genome reduction *via* the
underlying mechanisms to maintain homeostasis as a growing cell. How the growth
recovery of reduced genomes was achieved and whether it was in general are unclear.
To address the questions, we conducted the experimental evolution with a reduced
genome in multiple lineages and analyzed the evolutionary changes of the genome and
transcriptome in the present study.
70
Results
Fitness recovery of the reduced genome by experimental evolution
Experimental evolution of the reduced genome was conducted to regain the growth
fitness, which was decreased due to genome reduction. Serial transfer was performed
with multiple dilution rates to keep the bacterial growth within the exponential phase
(Fig. S1), as described [17, 20]. Nine evolutionary lineages were conducted independently
(Fig. S2, Table S1). A gradual increase in growth rate was observed along with the
generation passage, which was combined with a rapid increase in the early evolutionary
phase and a slow increase in the later phase (Fig. 1A). Most evolved populations (Evos)
showed improved growth fitness. Nevertheless, the final growth rates were considerably
varied (Fig. 1B, upper), and the evolutionary dynamics of the nine lineages were
somehow divergent (Fig. S2). It indicated the diversity in the evolutionary approaches
for improved fitness. Eight of nine Evos achieved faster growth than the
genome-reduced ancestor (Anc), whereas all Evos decreased in their saturated
population densities (Fig. 1B, bottom). In comparison to the wild-type strain carrying
the full-length genome (WT), the primarily decreased growth rate caused by genome
reduction was significantly improved by experimental evolution (Fig. 1C, upper),
associated with the considerable reduction in saturated density (Fig. 1C, bottom). It
demonstrated that the experimental evolution could compensate for the genome
reduction with a trade-off in population size (carrying capacity), consistent with the
previous findings [20, 21].
Intriguingly, a positive correlation was observed between the growth fitness and the
carrying capacity of the Evos (Fig. 1D). It seemed that the present experimental
evolution did not obey the *r/K* selection theory[30, 31], which was known as the trade-off

relationship (negative correlation) between the growth rate and the carrying capacity [32,

33]. Taking account of our previous finding that the colony growth rates of

genome-reduced strains were proportional to the colony sizes [11], the collapse of the

trade-off law likely resulted from genome reduction. As the trade-off between growth

fitness and carrying capacity was proposed to balance the cellular metabolism that

resulted from the cost of enzymes involved [33], the deleted genomic sequences might

play a role in maintaining the metabolism balance.

Significant variation and random localization of genome mutations

Genome resequencing (Table S2) identified a total of 65 mutations fixed in the nine

Evos (Table 1). The number of mutations largely varied among the nine Evos, from two

to 13, and no common mutation was detected (Table S3). 51 out of 65 mutations

occurred in the genes, and 45 out of 51 were SNPs. As 36 out of 45 SNPs were

nonsynonymous, the mutated genes might benefit the fitness increase. In addition, the

abundance of mutations was unlikely to be related to the magnitude of fitness increase.

For instance, A2 accumulating only two mutations presented a highly increased growth

rate compared to F2 of eight mutations (Table 1), which poorly improved the growth

(Fig. 1B). B2, D2, and E2 all succeeded in fitness increase to an equivalent degree (Fig.

1B), whereas they fixed 13, 7, and 3 mutations, respectively (Table 1). The mutated

genes were varied in 14 gene categories, somehow more frequently in the gene

categories of Transporter, Enzyme, and Unknown function (Fig. S3). They seemed

highly related to essentiality, as 11 out of 49 mutated genes were essential (Table S3).

As the essential genes were known to be more conserved than nonessential ones [34, 35],

the high frequency of the mutations fixed in the essential genes suggested the mutation

in essentiality for fitness increase was the evolutionary strategy for reduced genome.

The large variety in genome mutations and no correlation of mutation abundance to

fitness improvement strongly suggested that no mutations were specifically responsible

or crucially essential for recovering the growth rate of the reduced genome.

Additionally, there were no overlapped genomic positions for the 65 mutations in the

nine Evos (Fig. 2A). Random simulation was performed to verify whether there was any

bias or hotspot in the genomic location for mutation accumulation (Fig. 2B). The

simulation of 65 mutations randomly occurred on the reduced genome was performed

1,000 times. The distance of the mutation (mutated genomic location) to the nearest

genomic scar caused by genome reduction was calculated. Welch's t-test was performed

to evaluate the significance of the locational bias between the random mutations and the

mutations accumulated in the Evos (Fig. 2C). As the mean of p values from simulations

was insignificant ($\mu_p > 0.05$), there was no locational bias for mutation accumulation

caused by genome reduction.

Common evolutionary direction and homeostasis in transcriptome reorganization

Since no specificity was detected in the genome mutations, whether these mutations

disturbed the genome-wide gene expression pattern was investigated. Hierarchical

clustering and principal component analysis (PCA) showed that the evolved

transcriptomes were directed to similar patterns but divergent to the WT transcriptome

(Fig. S4). The evolutionary direction of the transcriptomes of the reduced genome was

not approaching the wild-type transcriptome. As a global feature of gene expression, the

chromosomal periodicity of the transcriptome was analyzed using the Fourier transform

as previously described [28, 36]. As a result, the transcriptomes of all Evos presented a

common six-period with statistical significance, equivalent to those of the wild-type and

ancestral reduced genomes (Fig. 3A, Table S4). It demonstrated that the chromosomal

architecture of gene expression patterns remained highly conserved, regardless of the

considerably varied mutations. The homeostatic periodicity was consistent with our

previous findings that the chromosomal periodicity of the transcriptome was

independent of genomic or environmental variation [27, 28].

In addition, the genomic locations of the mutations seemed irrelevant to

chromosomal periodicity (Fig. 3A, red lines). No mutagenesis hotspot was observed

even if these mutations were accumulated on a single genome (Fig. 2A). Alternatively,

the expression levels of the mutated genes were somehow higher than those of the

remaining genes (Fig. 3B), which sounded reasonable. The ratio of the essential genes

in the mutated genes was ~ 22% (Table S3), much higher than the ratio (~9%) of

essential to all genes (302 out of 3,290) in the reduced genome. As the essential genes

showed higher expression levels than the nonessential ones [37], the high essentiality of

the mutated genes might result in a higher mean expression level. On the other hand, the

high frequency of the mutations fixed in the essential genes was unexpected because the

essential genes were generally more conserved than nonessential ones [34]. It strongly

suggested the essentiality mutation for homeostatic transcriptome architecture happened

in the reduced genome.

Diversified functions and pathways of the differentially expressed genes

As the evolved transcriptomes were differentiated from those of the WT and reduced

genomes (Fig. S4), the differentially expressed genes (DEGs) were further identified to

discover the gene functions or biological processes contributing to the fitness changes.

The abundance of DEGs among the Evos varied from 333 to 1,130 genes and of few

overlaps (Fig. 4A). Most DEGs were unique to each evolutionary lineage, and the

common DEGs across all Evos were only 108 genes (Table S5). The number of DEGs

partially overlapped among the Evos declined significantly along with the increased

lineages of Evos (Fig. 4B). Enrichment analysis showed that only the histidine-related

pathways were significantly enriched in the common DEGs (Fig. 4C). Functional

enrichment of the DEGs in individual Evos showed that the amino acid metabolism

considerably participated and that the enriched pathways were poorly overlapped (Fig.

S5). These findings strongly suggested no universal rule for evolutionary changes of the

reduced genome.

In comparison, 1,226 DEGs were induced by genome reduction. The common DEGs

of genome reduction and evolution varied from 168 to 540, fewer than half of the DEGs

responsible for genome reduction in all Evos (Fig. 5A). The conclusion remained

consistent even if the DEGs were determined with an alternative method, RankProd

(Fig. S6). Functional enrichment of the DEGs observed the specific transcriptional

regulation and metabolic pathways participating in the transcriptome reorganization in

response to genome reduction and evolution. Only σ38 was enriched in the genome

reduction-mediated DEGs, which was the only regulon that partially overlapped with

the Evos (Fig. 5B). No regulon in common was enriched, besides a few partially

overlapped regulons, i.e., GadW, GadX, and RcsB (Fig. 5B). In addition, both the

number of enriched pathways and their overlaps were significantly differentiated among

the nine Evos or between Evos and reduced genome (Fig. 5C). No common pathways

were commonly enriched between the genome reduction-mediated DEGs and Evos, no

matter annotated the metabolic pathways with GO or KEGG (Fig. S7). The flagellar

assembly, the only enriched pathway in genome reduction-mediated DEGs, was absent

in all Evos (Fig. 5D). Alternatively, the amino acids-related metabolisms were

frequently detected in the Evos, e.g., histidine metabolism, biosynthesis of amino acids,

etc. (Fig. 5D, Fig. S7). The variable pathways in the Evos indicated that evolution

compensated for the genome reduction in various ways, which differed from how the

genome reduction was caused.

Gene modules responsible for the evolutionary changes of the reduced genome

Genome mutation analysis and transcriptome analysis failed to identify the common

gene categories or pathways that correlated to the evolution of the reduced genome; thus,

the gene modules correlated to evolution were newly evaluated. The weighted gene

co-expression network analysis (WGCNA) [38] was performed toward the evolved

transcriptomes as tested previously [27]. Reconstruction of the 3,290 genes in the reduced

genome led to 21 gene modules, comprising 8 to 320 genes per module (Fig. S8).

Hierarchical clustering of these modules showed that roughly three major classes could

be primarily divided (Fig. 6A). Functional correlation analysis showed that three of 21

modules (M2, M10, and M16) were highly significantly correlated to the growth fitness,

the number of DEGs, and mutation frequency, respectively (Fig. 6A). It indicated that

the three modules were highly essential and functionally differentiated for growth

control, transcriptional change, and mutagenesis.

Enrichment analyses further identified the gene categories, regulons, and

metabolisms that significantly appeared in the three modules (Fig. 6B). Two gene

categories of nonessential function were enriched in M10, and the module correlated to

the number of DEGs; in contrast, no gene category was enriched in M2 and M16 (Fig.

6B, left). All three modules successfully enriched the regulons without overlaps (Fig.

6B, middle). It indicates that the main regulatory mechanisms participating in the three

modules for growth control, transcriptional change, and mutagenesis were divergent.

GO enrichment resulted in various biological processes in the three modules, roughly

related to transport, transposition, and translation in M2, M10, and M16, respectively

(Fig. 6B, right). Compared to the enriched functions of the genes that disappeared due

to genome reduction (Fig. 6B, bottom), the gene categories of phage and unknown

function and the biological processes related to DNA transposition and integration were

commonly identified in M10. It strongly indicated that M10 was responsible for

genome reduction. The newly constructed gene networks successfully identified three

modules correlated to mutation, DEGs, and growth, revealing the functional

differentiation responsible for evolution to maintain the homeostatic transcriptome

architecture for a growing cell.

Discussion

The evolutionary compensation for genome reduction was directed toward an

identical goal of increased fitness but differentiated in the manner of genomic changes.

Firstly, various genetic functions seemed to trigger the increased growth rates of the

Evos. A few mutations could compensate for the sizeable genomic deficiency. The

considerable variation in the fixed mutations without overlaps among the nine Evos

(Table 1) implied no common mutagenetic strategy for the evolutionary improvement of

growth fitness. It was supported by the fact that no genomic locational bias for

mutations fixed in the evolution (Fig. 2). Secondly, the transcriptomes presented

conserved chromosomal architectures (Fig. 3) and universal directional changes (Fig.

S4), regardless of the significant and differentiated changes in gene expression in

response to evolution (Fig. 4, Fig. S5). Although the periodicity of chromosomal

architecture was well known [39, 40] and coordinated to the bacterial growth rate[28, 41],

employing its homeostasis as the evolutionary consequence provided a conceptual and

unique understanding of growing cells.

It's unclear whether the differentiation in evolutionary paths was particularly

significant for the reduced genome used in the present study. Evolution studies often

focus on finding the common mutations accumulated in multiple evolutionary lineages

to obtain the reasonable mechanism responsible for the adaptation to the defined

condition (Fig. 7, i). Common mutations[22, 42] or identical genetic functions[43] were

reported in the experimental evolution with different reduced genomes. Nevertheless,

divergent evolutionary mechanisms were proposed as not all mutations contribute to the

evolved fitness [22, 43]. The present study accentuated the variety of mutations fixed

during evolution. Considering the high essentiality of the mutated genes (Table S3),

most or all mutations were assumed to benefit the fitness increase, partially

demonstrated previously [20]. The differentiated mutations and DEGs guided to a

homeostatic (Fig. 7, iii) rather than a variable consequence (Fig. 7, ii). Multiple

evolutionary paths for the reduced genome to improve growth fitness were likely all

roads leading to Rome.

In addition, the transcriptome reorganization for fitness increase triggered by

evolution differed from that for fitness decrease caused by genome reduction. General

analyses failed to detect the regulatory network or genetic function mediated by genome

reduction and evolution in common. Instead, the newly constructed gene modules

successfully enriched the gene categories of mobile elements and unknown functions

(Fig. 6B, left) as the evolutionary compensation for genome reduction. The represented

mobile elements, flagella, were known to be responsive to environmental stresses such

as hypoosmotic pressure or pH [44, 45]. Genome reduction and evolution seemed

equivalent to the stress response in *E. coli*. These findings were reasonable as

enterobactin protected *E. coli* from oxidative stress [46, 47], and enterobactin biosynthesis

was upregulated for biofilm formation in genome-reduced *E. coli* [48]. The compensation

of evolution to genome reduction not only verified the known function and mechanism

from a global regulatory viewpoint but also revealed a novel understanding of the

molecular mechanisms and gene functions.

The discriminated functions of gene modules might play a crucial role in response to

genomic and evolutionary changes. WGCNA was conducted to discover the potential

correlation of gene expression to growth fitness. It succeeded in finding the genes

participating in the evolutionary changes of the reduced genome to regain growth

fitness. Three enriched gene modules were assumed separately responsible for

replication, transcription, and population dynamics (Fig. 6B). The growth-correlated

gene module significantly enriched the iron-related biological functions (M2). Although

the translation was commonly reported to be correlated to the growth rate [49, 50], it was

enriched in the gene module coordinated to transcriptional changes (M10). Such

functional differentiation of the gene modules might connect with the differentiated

medium components responsible for varied bacterial growth phases, which was

observed using the high-throughput growth assay in hundreds of medium combinations

combined with machine learning [51, 52]. We assumed that the various chemicals disturbed

different metabolic fluxes in which different gene modules might have participated. The

biological meaningfulness of the gene modules suggested an alternative genetic

classification besides the commonly used clustering criteria, such as Gene Orthology [53]

and Regulon [54]. In summary, the present study provided a representative example

showing multiple evolutionary paths (i.e., gene mutation and expression) directed the

reduced genome to the improved fitness with the homeostatic transcriptome.

Materials and Methods

E. coli strains and culture medium

The *E. coli* K-12 W3110 wild-type and its genome-reduced strains were initially

distributed by the National BioResource Project of the National Institute of Genetics.

The reduced genome was approximately 21% smaller than the wild-type genome. The

minimal medium M63 was used as described in detail elsewhere [13, 55].

Experimental evolution

The genome-reduced *E. coli* strain was evolved in 2 mL of the M63 medium by serial

transfer, as previously described [17, 20]. Nine evolutionary lineages were all initiated from

the identical culture stock prepared in advance. The 24-well microplates specific for

microbe culture (IWAKI) were used, and every four wells of four tenfold serial dilutions,

e.g., 10^3~10^6, were used for each lineage. The microplates were incubated in a

microplate bioshaker (Deep Well Maximizer, Taitec) at 37°C, with rotation at 500 rpm.

The serial transfer was performed at ~24-h intervals. Only one of the four wells

(dilutions) showing growth in the early exponential phase (OD_{600} = 0.01-0.1) was

selected and diluted into four wells of a new microplate using four dilution ratios. Serial

transfer was repeated until all evolutionary lineages reached approximately 1,000

generations, which required approximately two months per lineage. A total of nine

lineages were conducted, and the daily records of the nine lineages were summarized in

Table S1. The evolutionary generations (*G*) and the growth rates (*μ*) were calculated

according to the following equations (Eq. 1 and Eq. 2).

$$G = \log_2\left(\frac{N_t}{N_0}\right) \qquad \text{(Eq. 1)}$$

$$\mu = \frac{\ln\left(\frac{N_t}{N_0}\right)}{\Delta t} \qquad \text{(Eq. 2)}$$

Fitness assay

Both the ancestor and the evolved *E. coli* populations of the reduced genome were

subjected to the fitness assay, as previously described [13, 55]. In brief, every 200 μl of

culture was dispensed to each well in the 96-well microplate (Coaster). The microplate

was incubated at 37°C in a plate reader (EPOCH2, BioTek), shaking at 567 cpm (cycles

per minute) for 48 h. The temporal changes in OD_{600} were measured at 30-min intervals.

The growth fitness (*r*) was calculated using the following equation between any two

consecutive points (Eq. 3).

$$r_i = \frac{\ln\left(\frac{C_{i+1}}{C_i}\right)}{t_{i+1}-t_i} \qquad \text{(Eq. 3)}$$

Here, t_i and t_{i+1} are the culture times at the two consecutive measurement points, and C_i

and C_{i+1} are the OD_{600} at time points t_i and t_{i+1}. The growth rate was determined as the

average of three consecutive r_i, showing the largest mean and minor variance. The mean

of the biological triplicates was defined as the growth fitness and used in the analyses.

Genome resequencing and mutation analysis

The *E. coli* cells were collected at the stationary growth phase (i.e., $OD_{600} > 1.0$) and

subjected to genome resequencing, as described previously [20]. In brief, the bacterial

culture was stopped by adding rifampicin at 300 μg/mL. The cell pellet was collected

for genomic DNA purification using a Wizard Genomic DNA Purification Kit

(Promega) under the manufacturer's instructions. The sequencing library was prepared

using the Nextera XT DNA Sample Prep Kit (Illumina), and the paired-end sequencing

(300 bp × 2) was performed using the Illumina HiSeq platform. The raw datasets were

deposited in the DDBJ Sequence Read Archive under the accession number

DRA013662. The sequencing reads were aligned to the reference genome *E. coli*

W3110 (AP009048.1, GenBank), and the mutation analysis was performed with the
Breseq pipeline [56, 57]. The DNA sequencing and mapping statistics are summarized in
Table S2. The mutations fixed in the coding regions due to evolution are shown in Table
S3.

Calculation and simulation of the genomic positions of mutations

The distances from the genomic positions of the genome mutations fixed in the nine
Evos to the nearest genomic scars caused by the genome reduction were calculated as
described previously [13, 36]. Random mutations that occurred on the reduced genome
were simulated 1,000 times, and the distances of these mutations to the nearest genomic
scars were calculated. Note that the number of mutations in the simulation remained
equivalent to that detected in the Evos. Welch t-tests were performed to evaluate the
statistical significance (*p* values) of the bias in the distances observed in the Evos and
the simulations. The distribution and the mean of 1,000 *p* values acquired from the
simulation were calculated to evaluate the locational bias of the mutations fixed in
Evos.

RNA sequencing

The *E. coli* cells were collected at the exponential growth phase (i.e., $5 \times 10^7 \sim 2 \times 10^8$
cells/mL) and subjected to RNA sequencing, as described previously [27, 41]. In brief, the
bacterial growth was stopped by mixing with the iced 10% phenol ethanol solution. The
cell pellet was collected to purify the total RNAs using the RNeasy Mini Kit (QIAGEN)
and RNase-Free DNase Set (QIAGEN) according to product instructions. The
paired-end sequencing (150 bp $\times$ 2) was performed using the Novaseq6000
next-generation sequencer (Illumina). The rRNAs were removed from the total RNAs
using the Ribo-Zero Plus rRNA Depletion Kit (Illumina), and the mRNA libraries were
prepared using the Ultra Directional RNA Library Prep Kit for Illumina (NEBNext).
Biological replicates were performed for all conditions (N=2~4). The raw datasets were
deposited in the DDBJ Sequence Read Archive under the accession number
DRA013662.

 Data processing and normalization

The FASTQ files were mapped to the reference genome W3110 (accession number

AP009048.1, GenBank) using the mapping software Bowtie2 [58], as described

previously [27, 41]. The obtained read counts were converted to FPKM values according to

the gene length and total read count values. Global normalization of the FPKM values

was performed to reach an identical mean value in the logarithmic scale in all datasets.

The gene expression level was determined as the logarithmic value of FPKM, and the

mean values of biological replicates were used for the following analyses (Table S6).

Computational analysis

The normalized datasets were subjected to the computational analyses performed

with the R statistical analysis software. A total of 3290 genes were used for hierarchical

clustering and principal component analysis (PCA), which were performed with the R

functions of "hclust" and "prcomp", respectively, as described previously[27]. The

corresponding parameters of "method" and "scale" were set as "average" and "F",

respectively. The R package of DESeq2[59] was used to determine the differentially

expressed genes (DEGs), based on the false discovery rate (FDR < 0.05) [60]. The read

counts were used as the input data for DESeq2, in which the data normalization was

performed at each run for pair comparison.

Enrichment analysis

Functional enrichments were performed according to the features of gene category[61],

transcriptional regulation[54], gene ontology[53], and metabolic pathways[62, 63]. Twenty-one

gene categories and 46 regulons, which comprised more than 15 genes and 15

regulatees, were subjected to the enrichment, respectively. The statistical significance

was evaluated by the binomial test with Bonferroni correction. The enrichment analysis

of gene ontology (GO terms) [53, 64] and metabolic pathway (Kyoto encyclopedia of genes

and genomes, KEGG) [62, 63] was performed using DAVID, a web-based tool for

visualizing the characteristics of gene clusters with expression variation [65] [66]. The

statistical significance was according to FDR.

 Chromosomal periodicity analysis

    Fourier transform was used to evaluate the chromosomal periodicity of the

 transcriptome, as previously described [28, 41]. The genome was divided into

 compartments of 1 kb each, and the mean expression level of the genes within the

 corresponding sections was calculated. Gene expression levels were smoothed with a

 moving average of 100 kb and subjected to the periodicity analysis using the function

 "periodogram" in R. The max peak (periodic wavelength) of the periodogram was fitted

 to the gene expression data using the function "nls" in R by the least squares method

 according to the following equation (Eq. 4).

$$\text{Exp}(x) = a * \sin\left(x + \frac{b}{T}\right) + c \quad \text{(Eq. 4)}$$

 Here, a, b, T, and c represent the periodic amplitude, the periodic phase, the periodic

 wavelength indicated by the max peak, and the mean expression level of the whole

 transcriptome as a constant, respectively. The statistical significance of the periodicity

 was assessed with Fisher's g test (Table S4), as described previously [28, 41]. The genomic

 position of *ori* was according to the previous reports [41, 67]. In addition, the function

 "abline" in R was used to point out the genomic positions of the mutations.

 Gene network analysis

    The weighted gene co-expression network analysis (WGCNA) of the nine Evos was

 performed with the R package of WGCNA [38], as described previously [27]. A

 Step-by-Step method was used to determine the parameters for constructing the gene

 networks. The soft threshold was set at 12, where the R^2 of Scale Free Topology Model

 Fit was approximately 0.9 recommended by the developer's instruction. The resultant

 gene networks were clustered with the "hclust" function (method=average) and

 reconstructed by merging similar modules using the "mergeCloseModules" function

 with a height cut of 0.25 in the "dynamic tree cut" method. Finally, a total of 21

 modules were determined for 3,290 genes. The correlation coefficients and p-values

 between the expression of the gene modules and the other global features (growth rates,

 DEGs, and mutations) were evaluated using the functions "cor" and "corPvalueStudent"

 in R, respectively. FDR correction was applied to the p-values to account for

 multiplicity. Functional enrichment of the gene modules was performed, and the

statistical significance was evaluated by the binomial test with Bonferroni correction as
described above.

Acknowledgments
We thank NBRP for providing the *E. coli* strains carrying the wild-type and reduced
genomes (W3110 and KHK collection). This work was supported by the JSPS
KAKENHI Grant-in-Aid for Scientific Research (B) (grant number 19H03215) and
partially by Grant-in-Aid for Challenging Exploratory Research (grant number
21K19815).

Competing interests
The authors declare that there are no competing interests.

References
1.   Koonin EV. Evolution of genome architecture. *Int J Biochem Cell Biol* **41**,
298-306 (2009).
2.   Lynch M. Streamlining and simplification of microbial genome architecture.
Annu Rev Microbiol **60**, 327-349 (2006).
3.   Lynch M, Conery JS. The origins of genome complexity. *Science* **302**,
1401-1404 (2003).
4.   Kotaka Y, Hashimoto M, Lee KI, Kato JI. Mutations identified in engineered
Escherichia coli with a reduced genome. *Front Microbiol* **14**, 1189877 (2023).
5.   Kato J, Hashimoto M. Construction of consecutive deletions of the Escherichia
coli chromosome. *Mol Syst Biol* **3**, 132 (2007).
6.   Pósfai G*, et al.* Emergent properties of reduced-genome Escherichia coli.
Science **312**, 1044-1046 (2006).
7.   Hashimoto M*, et al.* Cell size and nucleoid organization of engineered
Escherichia coli cells with a reduced genome. *Mol Microbiol* **55**, 137-149
(2005).
8.   Aida H, Ying B-W. Efforts to Minimise the Bacterial Genome as a Free-Living
Growing System. *Biology* **12**, 1170 (2023).
9.   Breuer M*, et al.* Essential metabolism for a minimal cell. *Elife* **8**,   (2019).
10.   Hutchison CA, 3rd*, et al.* Design and synthesis of a minimal bacterial genome.
Science **351**, aad6253 (2016).

11. Hitomi K, Weng J, Ying BW. Contribution of the genomic and nutritional
differentiation to the spatial distribution of bacterial colonies. *Front Microbiol*
13, 948657 (2022).

12. Xue H, Kurokawa M, Ying BW. Correlation between the spatial distribution and
colony size was common for monogenetic bacteria in laboratory conditions.
BMC Microbiol **21**, 114 (2021).

13. Kurokawa M, Seno S, Matsuda H, Ying BW. Correlation between genome
reduction and bacterial growth. *DNA Res* **23**, 517-525 (2016).

14. Karcagi I*, et al.* Indispensability of Horizontally Transferred Genes and Its
Impact on Bacterial Genome Streamlining. *Mol Biol Evol* **33**, 1257-1269 (2016).

15. Gibson DG*, et al.* Creation of a bacterial cell controlled by a chemically
synthesized genome. *Science* **329**, 52-56 (2010).

16. Orr HA. Fitness and its role in evolutionary genetics. *Nat Rev Genet* **10**, 531-539
(2009).

17. Nishimura I, Kurokawa M, Liu L, Ying BW. Coordinated Changes in Mutation
and Growth Rates Induced by Genome Reduction. *mBio* **8**, (2017).

18. Csorgo B, Feher T, Timar E, Blattner FR, Posfai G. Low-mutation-rate,
reduced-genome Escherichia coli: an improved host for faithful maintenance of
engineered genetic constructs. *Microbial cell factories* **11**, 11 (2012).

19. Umenhoffer K*, et al.* Reduced evolvability of Escherichia coli MDS42, an
IS-less cellular chassis for molecular and synthetic biology applications.
Microbial cell factories **9**, 38 (2010).

20. Kurokawa M, Nishimura I, Ying BW. Experimental Evolution Expands the
Breadth of Adaptation to an Environmental Gradient Correlated With Genome
Reduction. *Front Microbiol* **13**, 826894 (2022).

21. Choe D*, et al.* Adaptive laboratory evolution of a genome-reduced Escherichia
coli. *Nat Commun* **10**, 935 (2019).

22. Suzuki S, Horinouchi T, Furusawa C. Prediction of antibiotic resistance by gene
expression profiles. *Nat Commun* **5**, 5792 (2014).

23. Moger-Reischer RZ*, et al.* Evolution of a minimal cell. *Nature* **620**, 122-127
(2023).

24. Ishikawa A*, et al.* Different contributions of local- and distant-regulatory
changes to transcriptome divergence between stickleback ecotypes. *Evolution* **71**,
565-581 (2017).

25. Rozen DE, de Visser JA, Gerrish PJ. Fitness effects of fixed beneficial mutations
in microbial populations. *Curr Biol* **12**, 1040-1045 (2002).

26.   Lynch M. Evolution of the mutation rate. *Trends Genet* **26**, 345-352 (2010).

27.   Matsui Y, Nagai M, Ying BW. Growth rate-associated transcriptome
506   reorganization in response to genomic, environmental, and evolutionary
507   interruptions. *Front Microbiol* **14**, 1145673 (2023).

508   28.   Nagai M, Kurokawa M, Ying BW. The highly conserved chromosomal
509   periodicity of transcriptomes and the correlation of its amplitude with the growth
510   rate in Escherichia coli. *DNA Res* **27**,   (2020).

511   29.   Ying BW, Yama K. Gene Expression Order Attributed to Genome Reduction and
512   the Steady Cellular State in Escherichia coli. *Front Microbiol* **9**, 2255 (2018).

513   30.   Engen S, Saether BE. r- and K-selection in fluctuating populations is determined
514   by the evolutionary trade-off between two fitness measures: Growth rate and
515   lifetime reproductive success. *Evolution* **71**, 167-173 (2017).

516   31.   Luckinbill LS. r and K Selection in Experimental Populations of Escherichia
517   coli. *Science* **202**, 1201-1203 (1978).

518   32.   Molenaar D, van Berlo R, de Ridder D, Teusink B. Shifts in growth strategies
519   reflect tradeoffs in cellular economics. *Mol Syst Biol* **5**, 323 (2009).

520   33.   Wortel MT, Noor E, Ferris M, Bruggeman FJ, Liebermeister W. Metabolic
521   enzyme cost explains variable trade-offs between microbial growth rate and
522   yield. *PLoS Comput Biol* **14**, e1006010 (2018).

523   34.   Jordan IK, Rogozin IB, Wolf YI, Koonin EV. Essential genes are more
524   evolutionarily conserved than are nonessential genes in bacteria. *Genome
525   research* **12**, 962-968 (2002).

526   35.   Zhang J. Important genomic regions mutate less often than do other regions.
527   *Nature* **602**, 38-39 (2022).

528   36.   Ying BW, Seno S, Kaneko F, Matsuda H, Yomo T. Multilevel comparative
529   analysis of the contributions of genome reduction and heat shock to the
530   Escherichia coli transcriptome. *BMC Genomics* **14**, 25 (2013).

531   37.   Ying BW, Seno S, Matsuda H, Yomo T. A simple comparison of the extrinsic
532   noise in gene expression between native and foreign regulations in Escherichia
533   coli. *Biochem Biophys Res Commun* **486**, 852-857 (2017).

534   38.   Langfelder P, Horvath S. WGCNA: an R package for weighted correlation
535   network analysis. *BMC Bioinformatics* **9**, 559 (2008).

536   39.   Mathelier A, Carbone A. Chromosomal periodicity and positional networks of
537   genes in Escherichia coli. *Mol Syst Biol* **6**, 366 (2010).

538   40.   Krogh TJ, Moller-Jensen J, Kaleta C. Impact of Chromosomal Architecture on
539   the Function and Evolution of Bacterial Genomes. *Front Microbiol* **9**, 2019

(2018).

41.   Liu L, Kurokawa M, Nagai M, Seno S, Ying BW. Correlated chromosomal
periodicities according to the growth rate and gene expression. *Sci Rep* **10**,
15531 (2020).

42.   Choe D*, et al.* Adaptive laboratory evolution of a genome-reduced Escherichia
coli. *Nature Communications* **10**,   (2019).

43.   Lu H*, et al.* Primordial mimicry induces morphological change in Escherichia
coli. *Commun Biol* **5**, 24 (2022).

44.   Ikeda T*, et al.* Hypoosmotic stress induces flagellar biosynthesis and swimming
motility in Escherichia albertii. *Commun Biol* **3**, 87 (2020).

45.   Maurer LM, Yohannes E, Bondurant SS, Radmacher M, Slonczewski JL. pH
regulates genes for flagellar motility, catabolism, and oxidative stress in
Escherichia coli K-12. *J Bacteriol* **187**, 304-319 (2005).

46.   Adler C, Corbalan NS, Peralta DR, Pomares MF, de Cristóbal RE, Vincent PA.
The alternative role of enterobactin as an oxidative stress protector allows
Escherichia coli colony development. *PLoS One* **9**, e84734 (2014).

47.   Peralta DR, Adler C, Corbalán NS, Paz García EC, Pomares MF, Vincent PA.
Enterobactin as Part of the Oxidative Stress Response Repertoire. *PLoS One* **11**,
e0157799 (2016).

48.   May T, Okabe S. Enterobactin is required for biofilm development in
reduced-genome Escherichia coli. *Environ Microbiol* **13**, 3149-3162 (2011).

49.   Scott M, Hwa T. Shaping bacterial gene expression by physiological and
proteome allocation constraints. *Nat Rev Microbiol*,   (2022).

50.   Dai X*, et al.* Reduction of translating ribosomes enables Escherichia coli to
maintain elongation rates during slow growth. *Nat Microbiol* **2**, 16231 (2016).

51.   Aida H, Hashizume T, Ashino K, Ying BW. Machine learning-assisted discovery
of growth decision elements by relating bacterial population dynamics to
environmental diversity. *Elife* **11**,   (2022).

52.   Ashino K, Sugano K, Amagasa T, Ying BW. Predicting the decision making
chemicals used for bacterial growth. *Sci Rep* **9**, 7251 (2019).

53.   Ashburner M*, et al.* Gene Ontology: tool for the unification of biology. *Nature
Genetics* **25**, 25-29 (2000).

54.   Salgado H*, et al.* RegulonDB v8.0: omics data sets, evolutionary conservation,
regulatory phrases, cross-validated gold standards and more. *Nucleic Acids Res*
41, D203-213 (2013).

55.   Kurokawa M, Ying BW. Precise, High-throughput Analysis of Bacterial Growth.

J Vis Exp,   (2017).

56.   Deatherage DE, Traverse CC, Wolf LN, Barrick JE. Detecting rare structural
variation in evolving microbial populations from new sequence junctions using
breseq. *Front Genet* **5**, 468 (2014).

57.   Barrick JE*, et al.* Identifying structural variation in haploid microbial genomes
from short-read resequencing data using breseq. *BMC Genomics* **15**, 1039
(2014).

58.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**, 357-359 (2012).

59.   Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550 (2014).

60.   Storey JD. A direct approach to false discovery rates. *Journal of the Royal*
Statistical Society: Series B (Statistical Methodology) **64**, 479-498 (2002).

61.   Riley M*, et al.* Escherichia coli K-12: a cooperatively developed annotation
snapshot--2005. *Nucleic Acids Res* **34**, 1-9 (2006).

62.   Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a
reference resource for gene and protein annotation. *Nucleic Acids Res* **44**,
D457-462 (2016).

63.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes.
Nucleic Acids Res **28**, 27-30 (2000).

64.   Carbon S*, et al.* AmiGO: online access to ontology and annotation data.
Bioinformatics **25**, 288-289 (2009).

65.   Huang DW*, et al.* DAVID Bioinformatics Resources: expanded annotation
database and novel algorithms to better extract biology from large gene lists.
Nucleic Acids Res **35**, W169-175 (2007).

66.   Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57
(2009).

67.   Bryant JA, Sellars LE, Busby SJ, Lee DJ. Chromosome position effects on gene
expression in Escherichia coli K-12. *Nucleic Acids Res* **42**, 11383-11392 (2014).

Table 1 Overview of fixed genome mutations. The number of mutations in the nine
Evos are shown separately and summed**.** SNP, N and S indicate single nucleotide
substitution, nonsynonymous and synonymous SNP, respectively.

Evos	All	Intergenic	Genic		Genic SNP	
			Indel	SNP	N	S
A1	3	0	0	2	1	1
A2	2	1	0	1	1	0
B2	13	0	1	11	9	2
C2	11	1	0	9	8	1
D2	7	0	1	5	4	1
E2	3	1	0	1	1	0
F2	8	1	0	6	5	1
G2	12	2	2	6	4	2
H2	6	0	2	4	3	1
Sum	65	6	6	45	36	9

Figure legends

Figure 1 Fitness increase of the reduced genome mediated by experimental

evolution. A. Temporal changes in growth rate. Color variation indicates the nine

evolutionary lineages. **B.** Growth rate and maximal population size of the reduced

genome. Blue and pink indicate the common ancestor and the nine evolved populations,

respectively. Standard errors are shown according to the biological replications (N=4~6).

C. Boxplots of growth rate and maximum. Cross and open circles indicate the mean and

individual values, respectively. Statistic significance evaluated by Mann-Whitney U

tests is indicated. **D.** Correlation between growth rate and maximum. Spearman's rank

correlation coefficient and p-value are indicated.

Figure 2 Genomic localization of mutations. A. Normalized genomic positions of all

mutations. The vertical lines highlight the total 65 mutations fixed in the nine Evos.

Color variation indicates the nine Evos. WT and Reduced represent the wild-type and

reduced genomes used in the present study. **B.** Normalized genomic positions of random

mutations. The simulation of 65 mutations randomly fixed in the reduced genome was

performed 1,000 times. As an example of the simulation, the genomic positions of 65

random mutations are shown. The vertical lines in purple indicate the mutations. **C.**

Statistic significance of the genome locational bias of mutations. The distance from the

mutated location to the nearest genomic scar caused by genome reduction was

calculated. The mutations accumulated in the nine Evos and the 1,000-time random

simulation were all subjected to the calculation. The significance of genome locational

bias of the mutations in Evos was evaluated by Welch's t-test. The histogram of 1,000

tests for 1,000 simulated results is shown. The mean of *p*-values (μ_p) is indicated, which

is within the 95% confidence interval ($0.07 < \mu_p < 0.09$).

Figure 3 Chromosomal periodicity of transcriptome and mutated gene expression.

A. Chromosomal periodicity of transcriptomes. The transcriptomes of the nine Evos are

shown. Black lines, red curves, and red vertical lines indicate the gene expression levels,

fitted periods, and locations of mutations, respectively. *Ori* and *dif* are indicated with

the vertical broken lines. **B.** Boxplot of gene expression levels. Gene expression levels

of the 49 mutated genes in the nine Evos and the remaining 3,225 genes are shown.

Statistic significance evaluated by Welch's t-test is indicated.

Figure 4 Differentially expressed genes (DEGs) and their enriched functions. A.

Commonality of DEGs in the nine Evos. Closed circles represent the combinations of

the Evos. Vertical and horizontal bars indicate the number of the overlapped DEGs in

the combinations and the number of all DEGs in each Evos, respectively. The

combinations with more than 20 DEGs in common are shown. **B.** The number of DEGs

overlapped among the Evos. The numbers of DEGs overlapped across 2 to 9 Evos are

shown. The number of Evos detected in the single Evos is indicated as 1. **C.** Enriched

function in common. The KEGG and GO terms enriched in the common DEGs across

the nine Evos are shown. The statistical significance (FDR) of the enriched pathways

and biological processes is shown on a logarithmic scale represented by color gradation.

Figure 5 Transcriptome comparison between genome reduction and evolution. A.

Venn diagrams of DEGs induced by genome reduction and evolution. The numbers of

individual and overlapped DEGs are indicated. **B.** Heatmap of enriched regulons.

Statistically significant regulons are shown with the FDR values on a logarithmic scale.

C. Number of enriched functions in common. Left and right panels indicate the numbers

of enriched GO terms and KEGG pathways caused by genome reduction and evolution,

respectively. **D.** Enriched functions in common. The overlapped GO terms enriched in

the nine Evos and genome reduction are shown. Blue and pink represent genome

reduction and evolution, respectively.

Figure 6 Reconstructed gene modules. A. Cluster dendrogram of the gene modules

reconstructed by WGCNA. A total of 21 gene modules (M1~M21) were reconstructed.

The significance of the correlation coefficients of the gene modules to growth, mutation,

and expression is represented in purple gradation. From light to dark indicates the

logarithmic p-values from high to low. **B.** Enriched functions of gene modules and

deletion. Enriched gene categories, regulons, and GO terms are shown from left to right.

The numbers of the genes assigned in the three gene modules and the genomic deletion

for genome reduction are indicated in the brackets. Color gradation indicates the

normalized p values on a logarithmic scale.

Figure 7 Schematic drawing of evolutionary approaches for the reduced genome.

Figure 1

Figure 3

Figure 4

Figure 5

Figure 6

Genome reduction   Experimental evolution

i

Similar Paths for Universality

ii

Differentiated Paths for Diversity

iii

Differentiated Paths for Universality

Figure 7