Divergent Protein Kinase A contributes to the regulation of flagellar waveforms in
Leishmania mexicana

Running Head: PKA modulates flagellar waveforms in *Leishmania*

Authors

Sophia Fochler [1, 2], Benjamin J Walker [3], Richard J Wheeler [4,5], Eva Gluenz [1,2,6]

Affiliations

[1] Sir William Dunn School of Pathology, University of Oxford, Oxford, UK

[2] Institute of Cell Biology, University of Bern, Switzerland

[3] Department of Mathematics, University College London, London, UK

[4] Medawar Building for Pathogen Research, Nuffield Department of Medicine, University of
Oxford, Oxford, UK.

[5] Institute for Immunology and Infection Research, School of Biological Sciences, University
of Edinburgh, Ashworth Laboratories, Edinburgh, UK.

[6] Wellcome Centre for Integrative Parasitology, School of Infection and Immunity, University
of Glasgow

* To whom correspondence should be addressed: eva.gluenz@unibe.ch

Abstract

The second messenger cyclic adenosine monophosphate (cAMP) and protein kinase A (PKA)
have been implicated in the regulation of flagellar motility in diverse eukaryotes, yet the
specific contributions of PKA isoforms remain unclear. Moreover, the kinetoplastid PKA is
insensitive to cAMP. Here we investigated subcellular localisations of PKA subunits in
Leishmania and study the flagellar waveform parameters in PKA deletion mutants. Expansion

microscopy localised *Lmx*PKAR1 between the paraflagellar rod and the axoneme and
*Lmx*PKAC1, *Lmx*PKAC2 depend on *Lmx*PKAR1 for enrichment in the flagellum.
*Lmx*PKAC3 requires *Lmx*PKAR3 for co-localisation at the cell cortex. Deletion of
*Lmx*PKAR3 resulted in increased flagellar *Lmx*PKAC3 signal and increased swimming speed.
Conversely, deletion of *Lmx*PKAC3 resulted in reduced flagellar beat frequencies and
swimming speed. The absence of *Lmx*PKAC1 resulted in a lower incidence of high frequency
tip-to-base waveforms, and consequently reduced swimming speed. *Lmx*PKAC1 dissociated
from cytoskeletons upon addition of guanosine, adenosine or inosine, consistent with a novel
nucleoside-based PKA activation mechanism. These data suggest that this divergent PKA
pathway regulates the motility of *Leishmania* through the modulation of flagellar waveforms.

Introduction

Motile cilia and flagella are dynamic, specialised organelles used by many cells for movement
through the environment. Over 700 proteins (Broadhead et al. 2006; Nakachi et al. 2011;
Pazour et al. 2005; Vasquez, van Dam, and Wheway 2021) form the flagellar cytoskeleton,
the axoneme, which typically consists of nine doublet microtubules and two central singlet
microtubules (called the central pair, CP). Radial spoke complexes (RS) project from the
doublets to the CP and the nexin links/dynein regulatory complex (N-DRC) project from one
doublet to its neighbour radially. Two rows of dynein motor protein complexes, the outer
dynein arms (ODA) and inner dynein arms (IDA), align in a 96 nm repeating pattern along
the doublet microtubules (Lindemann and Lesich 2010). Dyneins are a family of ATPases
which hydrolyse ATP to iteratively bind, pull and release adjacent microtubules (Heale and
Alisaraie 2020). This sliding is converted to bending by sliding restrictions imposed by the
basal body and sliding resistance from the N-DRC (Heuser et al. 2009). While eukaryotic
flagella have the same conserved core structures, important open questions remain about how
flagellar waveforms and frequency are modulated. Moreover, little is known of how different
cells translate cues from the local environment into appropriate flagellar waveforms.

Calcium and cAMP are the two main intracellular signalling molecules (called second
messengers) that have been linked to changes in flagellar waveforms in diverse types of cilia
and flagella, including mammalian respiratory cilia (Nlend et al. 2007), human sperm (Chung
et al. 2017; Balbach et al. 2018), bull sperm (Alonso et al. 2017; Ishijima 2015), hamster
sperm (Aoki, Sakai, and Kohmoto 1999; Kinukawa et al. 2006) and sea urchin sperm (Tash

1989; Vacquier et al. 2014; Beltrán, Zapata, and Darszon 1996), the green algae

Chlamydomonas (Hyams and Borisy 1978; Saegusa and Yoshimura 2015; Schmidt and

Eckert 1976), *Paramecium* (Lodh et al. 2016; N M Bonini, D L Nelson 1988), Euglena (Nasir

et al. 2022) and *Leishmania* (Mukhopadhyay and Dey 2016; 2017). Intraciliary cAMP

concentration correlates with beat frequency in mammalian cilia (Schmid et al. 2006; Ma,

Silberberg, and Priel 2002). The second messenger signal is transmitted through kinases,

phosphatases and calcium-binding proteins that are have been found to be incorporated into

core axoneme structures (Tash 1989; Christensen et al. 2001; Brokaw and Kamiya 1987;

Kutomi et al. 2012; Hamasaki et al. 1991). A key transducer of cAMP signals is cAMP-

dependent protein kinase (PKA), a highly conserved protein sharing similarities with cGMP-

dependent protein kinase (PKG) and protein kinase C (PKC) (Vacquier et al. 2014),

collectively named the AGC kinase subgroup. The paradigm for eukaryotic PKA signal

transduction involves a heterotetrameric holoenzyme composed of two PKA catalytic

subunits (PKAC) and two PKA regulatory subunits (PKAR), (C_2R_2). In vertebrates, A-kinase

anchoring proteins (AKAPs) bind the PKA heterotetrameric holoenzyme via the RII domains

in the PKAR, creating local micro-signalosomes (Torres-Quesada, Mayrhofer, and Stefan

2017). However, most recently, mammalian PKA has been shown to also exist in homodimer

form (C_1R_1) localising to radial spoke 3 within bovine sperm axonemes (Leung et al. 2025).

In its inactive state, unbound to cAMP, the autoinhibitory domain (AI) of PKAR occupies and

inhibits the catalytic core of the PKAC subunit (Anand et al. 2007). Upon binding of a cAMP

molecule to each of the PKAR subunits, the PKAC is then released to phosphorylate its

substrates. Flagellar ODAs are thought to represent an important target of PKA (Peng et al.

2015a; Salathe 2007).

Flagellate protozoa of the genera *Leishmania* and *Trypanosoma* are well-studied due to their

importance as parasites of humans and livestock. They also provide genetically tractable

study models for eukaryotic flagella in a lineage that is distinct from the three lineages in

which humans, green algae (*Chlamydomonas*) or ciliates (e.g. *Tetrahymena*, *Paramaecium*)

evolved. Motile *Leishmania* use a symmetric tip-to base flagellar waveform to move forward,

and switch to an asymmetric base-to-tip waveform to rotate the cell and change direction

(Gadelha, Wickstead, and Gull 2007). Trypanosomatid signal transduction pathways remain

poorly defined and, whilst cAMP causes flagellar waveform switching in *Leishmania*

(Mukhopadhyay and Dey 2016), there is considerable uncertainty in how signals are

transduced within trypanosomatid flagella.

All kinetoplastid genomes available on VEuPathDB (Alvarez-Jarreta et al. 2024) asides from

the aflagellate endosymbiont *Perkinsela*, encode three PKAC genes that share some canonical

features of eukaryotic PKAs, such as the conserved phosphoresidue located in the activation

loop (Cheng, Zhang, and McCammon 2006) along with a conserved catalytic core sequence

(Bachmaier et al. 2019a). Intriguingly, PKA in trypanosomes forms C_1R_1 heterodimers and

does not respond to cAMP (Bachmaier et al. 2019a; Shalaby, Liniger, and Seebeck 2001).

Instead, nucleosides and nucleoside analogues, including inosine, 7-deazaadenosine

(tubercidin) and toyocamycin, bind trypanosomatid PKAR with high affinity (Ober et al.

2024). Interestingly, when the known mammalian PKA phosphorylation substrate VASP was

exogenously expressed in *T. brucei,* it was phosphorylated on serine-157 by endogenous

parasite PKA (Bachmaier et al. 2019a), suggesting that, although *Trypanosoma* PKA

homologues are not regulated by cyclic nucleotides, their target site specificity may still be

shared across eukaryotic species.

Several lines of evidence point towards the trypanosomatid flagellum as a key site for PKA

function. First, flagellar localisation has been demonstrated for PKAC1/2 and PKAR subunits,

in flagellar proteomes of *T. brucei* and *L. mexicana* (Oberholzer et al. 2011; Beneke et al.

2019), by labelling *L. donovani* promastigote forms and *T. brucei* with PKAR1-specific

antibodies (Bachmaier et al. 2016) and in protein tagging screens (Beneke et al. 2019; Dean,

Sunter, and Wheeler 2017; Billington et al. 2023). Moreover, the flagellar localisation of

TbPKAR1 was shown to depend on IFT (Fort et al. 2016). Second, loss-of-function mutations

altered parasite motility patterns: depletion of TbPKAR1 by RNAi in bloodstream forms

reduced the cells' capacity for forward motility (Oberholzer et al. 2011). A complete deletion

of PKAR1 or a partial deletion of PKAC1 slowed the swimming speeds of *L. mexicana*

promastigotes (Beneke et al. 2019). Finally, the developmental regulation of transcripts and

activity in *Leishmania* spp. reflects the presence or absence of a motile flagellum: *L. major*

promastigotes, which possess a motile flagellum, showed much higher transcript levels

(Siman-Tov et al. 1996) than the amastigote form, which only have a short, non-motile cilium

(Bachmaier et al. 2016; Kobe and Kajava 2001). Proteome-wide, the phosphorylation kinetics

of the PKA target motif (RXXS*/T*) in *L. donovani* during promastigote to amastigote

differentiation consistently detected the highest signal in promastigote forms (Bachmaier et al.

2016), suggesting a promastigote-specific function for PKA activity. Still, the precise PKA

localisation, the mechanism of upstream physiological activation, and downstream biological

function remain to be fully explored.

Here we present a systematic analysis of three PKAC and two PKAR isoforms of *L. mexicana*,
using gene deletions and microscopy to discover how gene loss affects flagellar beat
dynamics. Localisation of fluorescently tagged proteins showed distinct subcellular
localisation patterns for PKAC and PKAR subunits, and specific co-dependencies, from
which we infer pairing preferences for *Lmx*PKAC1 and *Lmx*PKAC2 with *Lmx*PKAR1 in the
flagellum, and *Lmx*PKAC3 with *Lmx*PKAR3 along the cell cortex. Ultrastructure expansion
microscopy showed that LmxPKAR1 is localised adjacent to the axonemal structure, within
the inner/middle domain of the paraflagellar rod. Targeted gene deletion of PKA subunits
individually and in combination showed that none are essential for promastigote growth.
Assessment of swimming speed and measurement of flagellar waveform parameters showed
that LmxPKAR3/C3 seem to modulate beat frequency and swimming speed, whereas
*Lmx*PKAC1 is essential for the generation of a symmetric tip-to base flagellar waveform.

Results

Leishmania spp. have three PKAC and two PKAR subunits.

BLASTp searches with the three PKAC and the single PKAR protein sequences from
Trypanosoma brucei (Bachmaier et al. 2019a) identified syntenic *Leishmania mexicana*
orthologues for all four encoding genes (Table 1, Supplementary Figure 1A-C) (Amos et al.
2022). Analysis of the *L. mexicana* PKAC sequences confirmed overall conservation of the
trypanosomatid PKAC repertoire: the highly similar *Lmx*PKAC1 and *Lmx*PKAC2 are located
on chromosome 34, not in a tandem array but separated by four other genes. This locus is
syntenic with the locus of *T. brucei* and *T. cruzi* PKAC1 and PKAC2, but the designation of
these PKACs as 1 or 2, respectively, varies between species (Table 1, Supplementary Figure
1A). *Lmx*PKAC1 and *Lmx*PKAC2, but not *Lmx*PKAC3, have variable N-terminal extensions
that harbour a predicted disordered domain (Supplementary Figure 2A, B) and both the active
sites and peptide binding sites within all *Leishmania* PKAC1 and PKAC2 are identical
(Supplementary Figure 2C, D). Catalytic activity of eukaryotic PKAC is modulated by
phosphorylation of several residues, such as threonine-197 in the activation loop, which is
particularly involved in regulating activity (Steinberg et al. 1993; Steichen et al. 2012), and is
conserved across trypanosomatid PKAC (Supplementary Figure 2E). At the C-terminus,
trypanosomatid PKAC1 and PKAC3 have a conserved FXXF sequence, where mutating
either phenylalanine was shown to diminish catalytic activity (Etchebehere et al. 1997; Chan

et al. 2020). *LmjF*PKAC2 lacks the first of these phenylalanines (Supplementary Figure 2F).

All trypanosomatid PKAC3, including *Lmx*PKAC3, have an 8 amino acid C-terminal

extension of unknown function (Siman-Tov et al. 1996) (Supplementary Figure 2F).

Several reports have highlighted the existence of a second PKAR-like protein within the

trypanosomatids (Baker et al. 2021; Jäger et al. 2014), which was given the name PKAR3 in

L. donovani (Fischer Weinberger et al. 2024); for consistency, we adopt this terminology here.

L. mexicana has a syntenic orthologue of PKAR3 (Table 1) (Supplementary Figure 1D). All

trypanosomatid PKAR3s share a similar protein architecture with canonical PKARs (Figure

1A), with two CNBDs towards the C-terminus and an autoinhibitory (AI) domain that mimics

the phosphorylation targets of PKAC and inhibits the catalytic kinase subunits. This critical

sequence is typically RRXA(G) in Type I PKARs (PKAR-I) and RRXS(T) in Type II PKARs

(PKAR-II) (Figure 1B). In Type I PKARs, the alanine (A) or Glycine (G) replaces the

phosphorylatable serine (S) or threonine (T) residues found in substrate sites (C. E. Poteet-

Smith et al. 1997; Scott et al. 1985). Trypanosomatid PKAR1 sequences all have an RRXT

sequence that aligns with the autophosphorylatable RRXS motif in human PKAR-II. The

Leishmania PKAR3 sequences share the RGAIS motif of the canonical mammalian

RRRRGAISAE sequence of the PKAR-Iα AI, but notably are missing the first arginine

residue of the Type I PKAR motif RRXA (Figure 1B).

Deviations from the canonical PKAR architecture are found at the N-termini (Figure 1A):

trypanosomatid PKAR1 proteins lack the dimerization/docking domain (DD; PF02197) that

holds canonical eukaryotic PKA heterotetramer together and anchors it to the amphipathic

helix region of an A-kinase anchoring protein (AKAP) (Banky, Huang, and Taylor 1998).

Instead, they contain a predicted leucine-rich repeat (LRR; IPR032675). PKAR3 sequences

lack a recognisable DD domain or an LRR and, instead, there is a short coil domain predicted

near the N-terminus of all examined PKAR3 sequences. This suggests that the family of

divergent kinetoplastid PKAR have evolved subcellular anchoring mechanisms distinct from

AKAP.

The cyclic nucleotide binding domains of Leishmania PKAR subunits deviate from the

canonical cAMP binding domain of mammalian PKAR.

Studies using crystal structures showed that, in all trypanosomatid PKAR1s, there are

substitutions to key residues in the phosphate binding cassette (PBC) that impede docking of

the cAMP molecule (Bachmaier et al. 2019a): a conserved alanine (A) is replaced by a

negatively charged glutamate (E), which is predicted to clash with the phosphate group on the

cAMP molecule, and the arginine (R) residue required to dock the exocyclic phosphate has

been switched to polar uncharged (R/TN) or aliphatic (R/AV) residues (Bachmaier et al.

2019a; Ober et al. 2024). These changes have been experimentally demonstrated to cause a

ligand switch from cAMP to nucleoside analogues in trypanosomatid PKAR1s, and they are

conserved in *Leishmania* PKAR1 sequences (Figure 1C). These findings prompted us to

examine in detail the nucleotide binding sites of the PKAR3 subunits. Key residue switches

are evident in the parasite PKAR3 CNBDs but with some notable deviations from the PKAR1

patterns (Figure 1C). In the place of the first alanine (A) in CNBD-A, PKAR3 has a glycine

(G), which, assuming a local neutral pH environment would have a net zero charge. Thus, it is

hard to predict if this residue would clash with cAMP docking in the same manner as the

glutamate in PKAR1, defined by (Bachmaier et al. 2019a). The first alanine in CNBD-B has

been substituted for a glutamate in PKAR3, identical to PKAR1 sequences, and the arginine

residues have also been substituted in PKAR3 CNBD-A and CNBD-B with polar uncharged

(R/STN) or hydrophobic side chains (R/A). Most surprisingly is that the conserved initial

glutamate within the PBC (G**E**LAL), which forms a hydrogen bond with the 2'-OH moiety on

the ribose sugar of cAMP (Berman et al. 2005), has been switched to an aspartic acid (D) in

the *Leishmania* PKAR3 CNBD-A and aspartic acid or glutamine (D/Q) in all PKAR3 CNBD-

B asides from *Lmx*PKAR3. These changes suggest that PKAR3, like PKAR1, has diverged

from cAMP binding, but raise the question of whether the same or different ligands bind to

each of these sites.

All parasite sequences for PKAR1 and PKAR3 have an extended C-terminus end compared to

the canonical mammalian PKAR reference sequence (Figure 1A). Two tyrosine (Y) and two

valine (V) residues were implicated in forming a parasite specific hydrophobic pocket

documented in *Tc*PKAR1 (V444, Y486, Y485 and V489 denoted by orange asterisks in

Figure 1C and D), interactions with which were shown to increase the binding affinity and

activation potential of certain nucleoside analogues (Bachmaier et al. 2019a; Ober et al. 2024).

All parasite PKAR1 sequences, contain at least two of these YV residues. Contrastingly, in

PKAR3 sequences, *Leishmania* only have one Y and one V, where *Tc*PKAR3 has three of the

four hydrophobic residues. Taken together these data indicate that a divergent PKAR1 is

conserved in all trypanosomatids, and an additional PKAR3 is found in *Leishmania* spp. and

T. cruzi, with an unknown ligand specificity, that shares in its CNBD sequences some of the
divergent features of PKAR1.

L. mexicana PKACs and PKARs show distinct localisation patterns to the flagellum and
cell body cortex.

To establish the subcellular localisation of the different PKAC and PKAR subunits in *L.*
mexicana in different cell- and life-cycle stages, we inserted fluorescent protein fusion tags
(mNeonGreen, mNG) at the endogenous gene loci, producing mutants expressing either N- or
C-terminal fusion products for each PKA subunit.

mNG::PKAC1 localised to the flagellum in non-dividing cells (1 kinetoplast, 1 nucleus, 1
flagellum (1K1N1F) (Wheeler, Gluenz, and Gull 2011), a signal that remained strong in 0.1%
NP-40 detergent-extracted cytoskeletons (Figure 2A). Through the cell division cycle, the
signal abundance diminished in the flagellum and increased within the nucleus
(Supplementary Figure 3A), corroborating the nuclear localisation that was reported for this
protein in a *L. mexicana* kinase localisation screen (Baker et al. 2021). Strikingly, within the
nucleus, the *Lmx*PKAC1 signal intensity was consistently strongest in areas of low Hoechst
staining (Supplementary Figure 3A'), indicating localisation to the nucleolus. The
*Lmx*PKAC1 signal re-appeared in both the old and the newly growing flagella at cytokinesis.
In axenic amastigotes, the mNG::PKAC1 signal appeared greatly diminished, with weak
variable cytoplasmic signal (Figure 2A). A C-terminal tag (PKAC1::mNG) disrupted the
localisation to the flagellum (Supplementary Figure 4A), as previously reported (Beneke et al.
2019).

The mNG::PKAC2 signal was partitioned between the flagellum, the cell body and cortex
throughout in the promastigote, with variations in relative intensity in the different
compartments at different cell cycle stages (Figure 2B and Supplementary Figure 3B).
Notably, in early promastigote division (2F1N1K), the *Lmx*PKAC2 signal appeared enriched
in the new growing flagellum and a strong cell cortex signal was seen during mitosis
(Supplementary Figure 3B). The *Lmx*PKAC2 axonemal signal was lost from detergent-
extracted cytoskeletons, and only a weak variable cytoplasmic signal was seen in axenic
amastigotes (Figure 2B). Fusion of mNG to the C-terminus hindered protein localisation to
the flagellum (Supplementary Figure 4B).

The mNG::PKAC3 signal was associated with the cell body cortex in non-dividing cells, with

a signal that was strongest at the central region of the cell body and tapering off towards the

posterior pole (Figure 2C). In 95% of non-dividing cells, a distinctive 'edge effect' was

evident (Figures 2C and 3A) (Halliday et al. 2019). A faint signal was also present in the

flagellum, with an intensity gradient decreasing towards the distal flagellum (Figure 2C). This

signal was lost upon membrane extraction with 0.1% NP-40 (Figure 2C, middle panel). In

dividing promastigotes, the *Lmx*PKAC3 signal appeared more uniform across the cell body

cortex, with an accumulation at the anterior end of the developing cleavage furrow and a

reduced flagellar signal (Supplementary Figure 3C). The promastigote cortex signal

disappeared in detergent extracted cells and showed a weak variable cytoplasmic signal in the

cell body of axenic amastigotes (Figure 2C).

PKAR1::mNG showed a strong flagellar signal in promastigote forms, with a fainter diffuse

cell body signal throughout all stages of promastigote cell division (Figure 2D, 2F). Upon

detergent extraction, the fluorescent signal remained in the axonemal cytoskeleton. Fusion of

mNG to the N-terminus of PKAR1 resulted in a weak reticulated cell body signal, suggesting

that blocking the LRR domain may affect protein targeting (Supplementary Figure 4D). After

induction of axenic amastigote differentiation for three days, a strong PKAR1::mNG signal

was observed in the cytoplasm of axenic amastigotes (Figure 2D).

Both N- and C-terminally tagged *Lmx*PKAR3 localised exclusively to the cell body cortex in

live promastigotes (Figure 2E and Supplementary Figure 4E). This signal was detergent

extractable (Figure 2E), suggesting *Lmx*PKAR3 is anchored weakly to the cortical

cytoskeleton, or is associated with the plasma membrane. In axenic amastigotes, the

*Lmx*PKAR3 signal persisted at the cell body cortex (Figure 2E). Closer examination of the

different cell cycle stages showed that, in non-dividing (1F1N1K), and early dividing

(2F1N1K) promastigote cells, the cortical *Lmx*PKAR3 signal was most intense in the cell

body mid-zone and tapered off towards the posterior pole (Figure 2G, quantified in Figure

3B). In late division (2F2K1N / 2F2K2N), the *Lmx*PKAR3 signal intensity became more

uniform along the cortex, occasionally appearing enriched at the posterior pole (Figure 2G).

Overall, these data show that PKA concentrations in *L. mexicana* vary across sub-cellular

space and over time, with two dominant PKA structures being the promastigote flagellum

(dominated by *Lmx*PKAC1, -C2 and -R1) and the cell cortex of promastigotes and

amastigotes (*Lmx*PKAC3 and -R3). Within these dominant structures, there were cell-cycle

stage dependent changes in signal intensity and also a transient shift of the *Lmx*PKAC1 signal
in dividing cells into the nucleus and away from the flagellum.

Identification of localisation dependencies between PKAR and PKAC subunits to the
flagellum and cell cortex

Given the striking similarities in the localisation of *Lmx*PKAC3 and *Lmx*PKAR3 at the cell
cortex (Figure 2C, 2E), and the strong flagellar localisation of *Lmx*PKAC1 and *Lmx*PKAR1,
the question arose whether there were specific PKAC-PKAR pairings in different cellular
compartments. To test whether the localisation of any PKAC subunit was dependent on the
expression of specific PKARs, we deleted the *Lmx*PKAR1 or *Lmx*PKAR3 genes in cells
expressing tagged *Lmx*PKAC1, *Lmx*PKAC2 or *Lmx*PKAC3. Following deletion of
*Lmx*PKAR1, promastigotes no longer showed the strong axonemal *Lmx*PKAC1 signal that
was seen in the parental cell line (Figure 3C). Similarly, the flagellar signal from *Lmx*PKAC2
was also diminished upon deletion of *Lmx*PKAR1 (Figure 3D). By contrast, deletion of
*Lmx*PKAR1 had no visible effect on *Lmx*PKAC3 localisation at the cell cortex (Figure 3E).
We then tested whether loss of *Lmx*PKAR3 would affect *Lmx*PKAC3 localisation and found
that the distinct cortical signal of *Lmx*PKAC3 became diminished and the signal appeared to
become more widely distributed throughout the cell body (Figure 3F). This change in signal
distribution was quantified over a cross-section of the cell body between the kinetoplast and
the nucleus. This showed a complete loss of the edge effect for *Lmx*PKAC3 signal after
*Lmx*PKAR3 was deleted (Figure 3A, iii). Secondly, the change in signal distribution along the
axis of the cell was assessed by measuring fluorescent intensity at six positions along the cell
(Figure 3B). In a parental background, the signal profile of *Lmx*PKAC3 was at its lowest at
the distal flagellum, gradually increasing in intensity towards the pocket region (Figure 3B).
Following deletion of *Lmx*PKAR3, the *Lmx*PKAC3 signal increased in the flagellum and
decreased in the cell body (Figure 3B). Taken together, these data suggest pairing preferences
between PKAC and PKAR subunits, whereby in non-dividing promastigotes, *Lmx*PKAR1
anchors *Lmx*PKAC1 and *Lmx*PKAC2 in the flagellum, whilst *Lmx*PKAR3 docks *Lmx*PKAC3
at the cell body cortex.

PKAR1 is located in the zone between the axoneme and paraflagellar rod.

We next tested whether the *Lmx*PKAR1 was located at the radial spokes of the *Leishmania*
axoneme, since it was suggested that PKA associated with the AKAP domain of RSP3 in
Chlamydomonas spokes 1 and 2 (Gaillard et al. 2001; Yang and Yang 2006; Gaillard et al.

2006). More recently, a cryo-EM structure directly showed mammalian PKA associated with

malate dehydrogenase in the head of radial spoke 3 in human cilia (Leung et al. 2025). We

deleted proteins of the head (RSP9) and stalk (RSP3 and RSP11) of *L. mexicana* radial spokes

1 and 2 (Dahlin, Zheng, and Scott 2021) in the PKAR1::mNG expressing cell line to see if the

flagellar signal disappeared. A PKAR1::mNG signal was still observed in the axoneme of the

examined RSP deletion mutants, including the short flagella that are characteristic of RSP3

deletion mutants (Supplementary Figure 5A-D). We next employed ultrastructure expansion

microscopy to visualise the PKAR1 localisation patterns within *Leishmania* flagella (Figure 4)

at higher resolution. Cells were stained with antibodies against the c-myc epitopes (contained

in the PKAR1::mNG fusion protein), alpha-tubulin and PFR2 (a marker for the outer domain

of the PFR (Kohl, Sherwin, and Gull 1999)). In the expanded cells, microtubules of the

axonemes were clearly resolved and the PFR2 signal was on one side of the axoneme,

adjacent to the tubulin signal. The PKAR1 signal was sandwiched between these two

reference structures (Figure 4A, 4B). A recent systematic study of paraflagellar rod (PFR)

sub-domains defined in *T. brucei* predicted PKAC1, PKAC2 and PKAR1 to reside within the

inner/middle domain of the PFR (Gabriel et al. 2024). The observed fluorescence pattern is

consistent with this prediction. The PKAR1 signal initiated in the flagellar pocket, before that

of PFR2, and extended beyond PFR2, tapering off shortly before the flagellar tip (Figure 4C,

4D). Finally, we asked whether the PKAR1 localisation in this PFR domain depended on

PFR2, PFC21 or PFR assembly factor 1 (Alves et al. 2020), which are required for formation

of the outer PFR domain. Deletion of these genes in cells expressing PKAR1::mNG had no

effect on PKAR1 localisation to the flagellum (Supplementary Figure Fig 5E-J). Together,

these data suggest that flagellar PKAR1 is localised to the middle/inner PFR subdomain or at

the PFR-axoneme connectors (PACs) with close proximity to doublet microtubules 4-7

(Sivadas et al. 2012; Zhang et al. 2022) (Figure 4E).

Inosine dissociates PKAC1 signal from the flagellar cytoskeleton.

To determine whether *Lmx*PKAR1 could be pharmacologically activated by purine

nucleosides as was shown in *Trypanosoma* spp. (Bachmaier et al. 2019a; Ober et al. 2024),

we used an indirect assay that measured solubilisation of mNG::PKAC1 from flagella treated

with different compounds. Cytoskeletons from mNG::PKAC1 expressing cells were

incubated with cAMP, toyocamycin, adenosine, guanosine and inosine prior to imaging and

quantification of axonemal fluorescence (Figure 5).

The flagellar cytoskeleton mNG::PKAC1 signal remained visible after treatment with cAMP,

at 100 µM (Figure 5A-B). However, quantification revealed a significant decrease in

fluorescent signal relative to untreated mNG::PKAC1 cytoskeleton controls (Figure 4F).

Toyocamycin was more efficacious at dissociating the mNG::PKAC1 signal at 25 µM than

100 µM cAMP (Figure 5F). Adenosine treatment at 10 µm had a similar effect on

mNG::PKAC1 signal solubilisation than toyocamycin treatment. Guanosine and inosine were

the most efficacious at reducing mNG::PKAC1 signal, with inosine treatment resulting in

flagellar cytoskeletons with similar signal intensity to untagged negative controls (Figure 5C-

F). Assessing a range of inosine concentrations on mNG::PKAC1 signal dissociation

indicated an EC_{50} of ~9 nM (Figure 5G-5H). This is consistent with recently published

biochemical assays highlighting inosine as the highest affinity activator for trypanosomatid

PKAR1, with guanosine and adenosine following at lower efficacies (Ober et al. 2024).

Together, this adds to growing evidence that the physiological activating ligand is

biochemically similar to purine nucleoside analogues like inosine, or perhaps inosine itself.

LmxPKAC1 plays an important role in flagellar motility.

We previously showed that partial deletion of *Lmx*PKAC1 (*ΔLmxM.34.4010*) reduced

promastigote swimming speed (Beneke et al. 2019). Here, we employed the same CRISPR-

Cas9 gene deletion strategy to generate the cell lines *ΔLmxPKAC1*, *ΔLmxPKAC2* and

ΔLmxPKAC3 as well as *ΔLmxPKAR1* and *ΔLmxPKAR3* (Supplementary Figure 6) to measure

precisely how the loss of each of these proteins affected flagellar motility. The deletion

mutants were all viable as promastigotes, with comparable *in vitro* growth rates and overall

cell morphology as the parental cell line (Supplementary Figure 7). The swimming behaviour

of deletion mutant populations was measured from darkfield videos (Wheeler 2017) and the

results (Figure 6A, Supplementary Table 1) confirmed that promastigotes lacking *Lmx*PKAC1

swam significantly slower (average 4.9 µm/s ± 0.035 standard deviation, unpaired t-test

assuming equal variance ($p < 3.5 \times 10^{-14}$) than the parental cell line (7.4 µm/s ± 0.917).

ΔLmxPKAC3 and *ΔLmxPKAR1* also swam slower (5.7 µm/s ± 0.032 and 5.4 µm/s ± 0.273,

respectively); deletion of *LmxPKAR3* had the opposite effect and swam faster (8.4 µm/s ±

0.775). By contrast, deletion of axonemal *Lmx*PKAC2 did not significantly alter parasite

swimming speed (average 6.4µm/s ± 0.650, unpaired t-test, p = 0.093) (Figure 6A). To test

what the combined effect was of deleting *Lmx*PKAC1 and either *Lmx*PKAR3 or *Lmx*PKAC2

in the same cells, the respective double deletion mutants were generated (Supplementary

Figure 6B). These showed the same reduced swimming speed as the *ΔLmxPKAC1* population

(Figure 6A), indicating that loss of *Lmx*PKAC1 was the dominant cause of the observed slow

motility phenotype, which could not be rescued or further compounded by secondary gene

deletion of *Lmx*PKAR3 or *Lmx*PKAC2.

LmxPKAC1 is required for the generation of symmetric flagellar waves.

Since swimming speed measurements report the average for a whole population, we next used

high speed video microscopy, alongside semi-automated tracing scripts (Walker, Ishimoto,

and Wheeler 2019; Walker et al. 2019), to further dissect how deficiency of each PKA

subunit was affecting the flagellar waveform of individual cells. First, we captured 0.5

seconds of high speed (200 Hz) low magnification (×20) videos to assess the relative

proportions of the different waveforms that *L. mexicana* are known to generate: the tip-to-

base symmetric flagellar waveform, the base-to-tip asymmetric ciliary waveform (Figure 6B),

or, an aperiodic movement referred to as uncoordinated.

ΔLmxPKAC3 and *ΔLmxPKAR3* showed similar proportions of cells performing symmetric

tip-to-base waveforms as the parental cells (~75%). However, all PKA deletion mutants

appeared to display fewer instances of ciliary waveforms and a greater proportion of cells (>

25%) moving in an uncoordinated manner (Figure 6C). *ΔLmxPKAC2* and *ΔLmxPKAR1* both

displayed a lesser proportion of cells propagating the flagellar wave (< 65%). Strikingly,

ΔLmxPKAC1 cells were completely unable to propagate the typical tip-to-base symmetric

flagellar waveform and, instead, showed completely uncoordinated flagellum movements

(Figure 6C, D, Supplementary movies 1 and 2). The uncoordinated movement and slow

population swimming speed were rescuable by the addition of an episomal addback copy of

*Lmx*PKAC1 (Supplementary Figure S8). This observation demonstrates that *Lmx*PKAC1 is

essential for producing a symmetric and rhythmic flagellar waveform.

All PKAC subunits contribute in different ways to beat amplitude and frequency.

In order to dissect how the loss or dysregulation of PKA contributes to the overall waveform,

we used cell tracing and Fourier analysis to extract the dominant beat frequency, amplitude

and waves per flagellum comprising the tip-to-base symmetric waves for the parental control

cells each PKA deletion cell line (Figure 7A-F). *L. mexicana* tip-to-base waves usually have a

frequency of 20-35 Hz and an amplitude between 40 to 60 degrees (Walker et al. 2019; Wang

et al. 2020). This was confirmed here for the parental control cell line (Figure 7, p-values

listed in Supplementary Table 2). *ΔLmxPKAC1* were never observed propagating a rhythmic

symmetric flagellar wave (Figure 6C). Our best effort to fit a dominant frequency of

attempted tip-to-base movements showed a greatly diminished frequency (< 10 Hz), with an

average shear angle amplitude of 26 degrees (Figure 7A), indicating that *Lmx*PKAC1 is

required for the generation of a normal flagellar waveform. The flagellar length and

wavelength are coupled in *Leishmania* and relates species, leading to tendency of ~1

wavelength per flagellum (Gadelha, Wickstead, and Gull 2007). We saw here that these

aforementioned parameters showed a positive correlation in all PKA deletion cell lines,

except *ΔLmxPKAC1* cells (Supplementary Figure 9). Flagellar waves from *ΔLmxPKAC2*

displayed a slight but significantly diminished beat amplitude (Figure 7B). We also observed

that *ΔLmxPKAC3* had significantly diminished beat frequencies (averaging 22 Hz), but

otherwise displayed normal beat amplitudes (Figure 7C). *ΔLmxPKAR1* retained comparable

flagellar wave parameters relative to parental control cells (Figure 7D). Lastly, although

ΔLmxPKAR3 and *ΔLmxPKAC2* mutants showed a comparable average beat frequency for tip-

to-base symmetric waves to parental control populations, the measured values appear to show

a bimodal distribution, with a sub-population with low beat frequency (~15 Hz) and one with

an abnormally high beat frequency (> 40Hz) (Figure 7B, 7E). The normal average population

speed for *ΔLmxPKAC2* and elevated population swimming speed for *ΔLmxPKAR3* (Figure

6A) could have arisen from the averaging of measurements from a population displaying

these extreme frequency values. More cells would need to be captured, and perhaps observed

over longer time periods, to ascertain the true relative proportions of these phenotypes in the

population and assess whether individual flagella can switch between abnormally low and

high frequencies.

Taken together, these measurements show that removal of different PKA subunits perturbed

the flagellar waveforms of *L. mexicana* in specific ways. Removal of the catalytic subunit

PKAC1, which is normally docked to the flagellar cytoskeleton through association with

PKAR1, had the biggest effect; without PKAC1 the flagellar waveforms lost their symmetry

and this uncoordinated beating reduced the speed at which the cells swam.

Discussion

Protein kinase A is one of the most studied eukaryotic signal transducers, which is highly

conserved throughout the eukaryotes (with the exception of most plant lineages) (Peng et al.

2015a) and fine-tunes a multitude of cellular processes. Within motile cilia, PKA activity modulates sperm beating behaviours, capacitation and hyperactivation, which are both essential for reproduction (Alonso et al. 2017; Lefièvre et al. 2002). Despite this importance, there is a lack of clear understanding of how PKA activity is spatially controlled and how its activity fine-tunes flagellar waveform properties. In kinetoplastids, converging evidence from proteomics and transcriptomics datasets and mutant phenotypes suggested their divergent PKA may also have a role in flagellar motility regulation. Here, we used a genetic approach combined with fluorescence-, expansion- and video-microscopy, to dissect the localisations and paring preferences of the three *L. mexicana* PKAC and two PKAR subunits and studied how gene deletions impacted cell swimming and generation of symmetric waveforms. Our results corroborate recent reports of PKAC/R pairings in *Leishmania* (Fischer Weinberger et al. 2024) and extend previous analyses on PKA function by providing multiple lines of evidence for a necessary function for *Lmx*PKAC1 in flagellar waveform regulation.

We found that there are two regions of PKA concentration in promastigote forms: *Lmx*PKAC1, *Lmx*PKAC2 and *Lmx*PKAR1 showed enrichment in the flagellum, while a strong cell body cortex signal was detected for *Lmx*PKAC3 and *Lmx*PKAR3. These localisations were neither exclusive nor static and examination of cells at different stages in the division cycle indicated relocalisation of all three *Lmx*PKAC subunits and of *Lmx*PKAR3; only *Lmx*PKAR1 maintained its dominant flagellar signal in all examined cell cycle stages, and withstood detergent extraction. Upon deletion of *Lmx*PKAR1, the flagellar signal of *Lmx*PKAC1 was greatly diminished and we conclude that *Lmx*PKAR1-*Lmx*PKAC1 interaction is necessary for *Lmx*PKAC1 localisation to the flagellum.

Characterising the phenotypes of gene deletion mutants showed that *Lmx*PKAC1 was essential for propagation of symmetric flagellar waveforms used for forward motility. All PKA deletion cell lines were viable and remained motile (including the double deletion of *Lmx*PKAC1 and *Lmx*PKAC2), but measurement of swimming speeds showed differences compared to the parental control line for all deletion mutants except *ΔLmxPKAC2* (Figure 6A). *ΔLmxPKAC1* showed the greatest reduction in swimming speed and this mutant was unique in its complete inability to generate any symmetric tip-to-base flagellar waves (Figure 6C), instead beating its flagella in an uncoordinated manner characterised by a lack of periodicity, symmetry, a very low dominant beat frequency and reduced amplitude compared to normal tip-to-base waveforms of promastigotes (Figure 7). These data suggest *Lmx*PKAC1

467 is required within the flagellum to maintain the appropriate spatio-temporal activation of
468 axonemal dyneins, most likely as a component of a signal transduction cascade that leads to
469 the phosphorylation of either core axonemal proteins that regulate dyneins, or of dynein
470 subunits directly. The ODAs, which are important for the frequency of base-to-tip asymmetric
471 waveform in *Chlamydomonas* (Brokaw and Kamiya 1987), are thought to be a key target of
472 cAMP-dependent protein kinase (PKA) (models, reviewed in (Tash 1989; Cheng, Zhang, and
473 McCammon 2006)). Outer dynein light chain proteins from the ciliates *Paramecium*
474 *tetraurelia* (p29 (Barkalow, Hamasaki, and Satir 1994)) and *Tetrahymena* (p34 (Christensen
475 et al. 2001)) were shown to be phosphorylated upon treatment with cAMP and RNAi
476 knockdown of *P. tetraurelia* outer dynein light chain 1 (ODLC1) resulted in loss of
477 phosphorylated p29 and rendered cilia unresponsive to cAMP (Kutomi et al. 2012). It will be
478 interesting to test whether ODAs are targets for the divergent cAMP-independent PKA in
479 kinetoplastids. Phosphoproteomic data from *T. brucei* treated with a synthetic PKA activator,
480 7-CN-7-C-ino, identified phosphopeptides for several proteins with known functions in
481 normal motility including several dynein intermediate chains (e.g. IC138), distal docking
482 complex 2 (dDC2) and nexin dynein regulatory complex-2 (DRC2) (Bachmaier et al. 2019a).
483 The specificity and functional significance of any of these phosphosites remains to be tested.
484 Comparisons of phosphorylation patterns in axonemes of *L. mexicana* PKA mutants and wild
485 type controls could further corroborate candidates for *Lmx*PKAC1 phosphorylation substrates
486 and allow for further dissection of the mechanism by which phosphorylation contributes to
487 flagellar waveform modulation.

488 Cells lacking *Lmx*PKAR1 also swam significantly slower, and this mutant lost most of its
489 flagellar *Lmx*PKAC1 signal. This is consistent with a model where axonemal *Lmx*PKAR1
490 binds and concentrates PKAC1 along the flagellum, placing it in proximity to axonemal
491 proteins which are phosphorylated upon receipt of an activating signal. The expansion
492 microscopy data place the *Lmx*PKAR1 at the interface between axoneme and PFR structure
493 (Figure 4). This suggests a model where *Lmx*PKAC1 activity is concentrated asymmetrically
494 within the motile axoneme with proximity to doublets 4-7. It is tempting to speculate that the
495 different placement of PKAR in different types of motile cilia is related to their distinct
496 waveforms and preferred wave propagation direction: localisation to distinct radial spokes is
497 observed in *Chlamydomonas* and mammalian cilia (Leung et al. 2025), with exclusively base-
498 to-tip propagation, while the asymmetric extra-axonemal PKAR localisation in kinetoplastids
499 occurs in flagella with a need to modulate both near-planar tip-to-base and asymmetric base-

to-tip beat types (Gadelha, Wickstead, and Gull 2007; Wang et al. 2020). *Leishmania* lacking

the outer PFR domain due to deletion of PFR2 swim slower (Santrich et al. 1997) but still

propagate predominantly tip-to-base waveforms that tend to remain near symmetric, with

variable frequency and lower amplitude (Wang et al. 2020). These mutants do not fully

phenocopy the defects in the *ΔLmxPKAC1* mutants. Deletion of PFR assembly factors or

major components of the outer and middle/inner domains (PFR2 array and PFC21,

respectively) leave a remnant PFR structure intact (Santrich et al. 1997; Alves et al. 2020;

Gabriel et al. 2024), which still retains PKA in *Leishmania* (Supplementary Figure 5E-I).

Future identification of the PAC components necessary for tethering the PFR to the axoneme

could provide a route to a clean removal of the entire PFR structure, which would allow

further studies to observe the *Leishmania* wave shape upon complete removal of the

asymmetric PFR structure.

Of course, it is not unusual for kinases to have pleotropic function and moreover, localisation

of proteins does not necessarily indicate function within that domain. Likewise with the

observed dual localisation of *Lmx*PKAC1, its localisation in the flagellum in non-dividing

cells could be, in part, a sequestration away from other targets within the cell body or nucleus,

which may be involved in replication events. However, while the mutant analysis revealed a

clear motility phenotype upon loss of *Lmx*PKAC1, cell growth remained unaffected, arguing

against a key role in the regulation of cell division. Alternatively, during division, where

forward swimming may need to be dampened, *Lmx*PKAC1 may be partitioned to the nucleus

until returning to the new and old flagella. Experiments involving pulse-chase labelling of

proteins would be required to assess the relationship between the different *Lmx*PKAC1

populations fractions over the course of the cell cycle. In any case, the activation status of

*Lmx*PKAC1 in the nucleus remains an open question. *Lmx*PKAR1 was never observed

alongside its catalytic subunit in the nucleolus during division and this spatial uncoupling

would presumably populate the nucleolus with active *Lmx*PKAC1 or, if sequestration was the

purpose, require an alternative regulatory mechanism for *Lmx*PKAC1 inactivation.

*Lmx*PKAC1 and *Lmx*PKAC2 share 86% sequence identity and localisation data suggest

*Lmx*PKAR1 is required for their enrichment in the flagellum. Unlike *Lmx*PKAC1, however,

*Lmx*PKAC2 was solubilised by treatment with 0.1% NP-40 while *Lmx*PKAC1 remained in

the flagellar cytoskeletons. Furthermore, loss of *Lmx*PKAC2 had only a small effect on

flagellar waveforms when observed at the level of populations: *ΔLmxPKAC2* displayed

comparable population swimming speeds to parental controls, with only a small increase in the proportion of cells exhibiting uncoordinated flagellar waveforms (Figure 6C). The observation that the frequencies of symmetric waveforms of individual cells segregated into two groups, low frequency or high frequency, suggests that *Lmx*PKAC2 may have a role in maintaining a constant beat frequency (Figure 7B). The population overview waveform analyses represent snapshot observations of flagella over a short time period (0.5 seconds). To study this further, it would be informative to observe individual cells for a longer time to determine if periods of low frequency waves alternate with high frequency waves in single flagella.

*Lmx*PKAC3 had a clearly distinct localisation, concentrated at the cell cortex, similar to *Lmx*PKAR3. The peripheral enrichment was lost upon deletion of *Lmx*PKAR3, suggestive of a *Lmx*PKAC3/R3 complex. A faint flagellar signal was also observed for *Lmx*PKAC3, which was increased in the *ΔLmxPKAR3* deletion mutant. Interestingly, the motility phenotype of gene deletion mutants correlated with changes of *Lmx*PKAC3 abundance in the flagellum: the *ΔLmxPKAC3* population swam more slowly and *ΔLmxPKAR3* swam faster than parental controls. The slower swimming speed for *ΔLmxPKAC3* was likely the combined effect of the increased proportion of cells showing uncoordinated waveforms and the significant decrease in the beat frequency of symmetric waves (Figure 7C). Additionally, there was a small reduction in average flagellum length (Supplementary Figure 7D). Interestingly, loss of *Lmx*PKAR3, which led to a re-distribution of *Lmx*PKAC3 away from the cell cortex, resulted in more cells propagating flagellar waveforms exceeding 40 Hz (Figure 7E) and increased mean population swimming speed (Figure 6A). It remains an open question whether *Lmx*PKAC3 has bona-fide phosphorylation targets within the flagellum, or whether the increased amount of soluble catalytic subunit in the *Lmx*PKAR3 deletion mutant causes off-target phosphorylation events in the axoneme that affect motility, secondary to its main role elsewhere. As deletion of *Lmx*PKAR3 (shown to increase flagellar *Lmx*PKAC3 Figure 3B) in *ΔLmxPKAC1* flagella did not rescue the motility defect of *ΔLmxPKAC1* (Figure 5A), it seems probable that these catalytic subunits have distinct phosphorylation substrates within the motile flagellum. A recent report linked *Lmx*PKAR3 and *Lmx*PKAC3 to processes involved in cell morphology in *L. donovani* (Fischer Weinberger et al. 2024). This may explain the shortened flagella in our *ΔLmxPKAC3* cells; however, we did not observe any morphological changes in *ΔLmxPKAR3*. Moreover, all of our *L. mexicana* PKA deletion mutant cell lines

here showed normal cell growth and maintained the same mean cell pseudo-diameter as the

parental controls (Supplementary Figure 7).

Owing to the high conservation of PKA, it has long been presumed that its activation

mechanism through cAMP was equally as conserved, until the recent discovery that

trypanosome PKA is insensitive to cAMP and instead activated by purine nucleoside

analogues (Ober et al. 2024; Bachmaier et al. 2019a). Multiple lines of evidence indicate that

this alternative PKA activation pathway also operates in *L. mexicana*: The sequences of the

*Lmx*PKAR1 and *Lmx*PKAR3 phosphate binding cassettes showed some of the characteristic

residue switches that were shown in *Trypanosoma* to prevent cAMP binding. Moreover,

treatment of cytoskeletons with cAMP or nucleosides to test for dissociation of the

*Lmx*PKAC1 signal revealed inosine as the most potent (EG$_{50}$ 9 nM), followed closely by

guanosine (Figure 5). This is an indirect assay for PKA activation and the results are

consistent with the model of the *T. brucei* PKAR1/C2 complex, where inosine binding to

PKAR resulted in the release of the catalytic subunit and subsequent substrate

phosphorylation (Bachmaier et al. 2019a; Ober et al. 2024). As inosine is an intermediate

compound of the purine metabolism pathway (Boitz et al. 2012), it may be that the

endogenous ligand for divergent PKA signalling is a derivative of this pathway, or inosine

itself.

Although the role of cAMP-responsive PKA in flagellar signalling has been established

(Soriano-Úbeda et al. 2019; Lefièvre et al. 2002), detailed structural maps of ciliary axonemes

revealed variations in PKA positioning between *Chlamydomonas* and humans. Our study

adds to this emerging picture of lineage-specific positional variation, and shows that second

messengers outside of cAMP and calcium may be used for flagellar waveform modulation

through PKA. Despite this diversity of signal transducers, it will be interesting to discover

whether the signals converge at conserved regulatory structures to orchestrate the spatio-

temporal activity of axonemal dyneins.

Methods

Cell culture

All cell lines used in this project were derived from *Leishmania mexicana* (WHO strain

MNYC/BZ/62/M379) cell line expressing Cas9 and T7 RNA polymerase (Beneke et al. 2017).

Promastigote form cells were cultured exactly as described in (Beneke et al. 2017). In brief,

cells were incubated at 28^{o}C and maintained in M199 (Life Technologies) medium

supplemented with 2.2 g/L sodium bicarbonate between 1×10^5 and 2×10^7 cells/ml with

regular sub-culturing.

Endogenous fluorescent tagging and gene deletion

Gene deletions and cell lines expressing endogenously fused fluorescent gene products were

essentially created as outlined by Beneke *et al*., (Beneke et al. 2017). An online resource for

automated primer design (www.LeishGEdit.net) was used to obtain primer sequences for the

replacement of target gene ORFs with drug resistance cassettes (pT plasmids used to create

donor products) or insertion of mNeonGreen (pPLOT plasmids used to create donor products)

to the endogenous gene locus (Beneke et al. 2017). Promastigote populations transfected with

DNA donor and guide constructs were maintained in M199 medium supplemented with

relevant selection drugs (either 20 µg/ml puromycin, 5 µg/ml Blasticidin or 40 µg/ml G418)

for 3 culture passages before performing a PCR based gene deletion verification.

Diagnostic PCR for gene deletion validation.

Primers sequences targeting the open reading frames (ORF) of genes of interest (GOI) were

taken from the LeishGEdit primer database (www.LeishGEdit.net) and sourced from

ThermoFisher. Due to high sequence similarity, *LmxPKAC1* and *LmxPKAC2* gene deletion

verification, primers were manually designed using Primer3 (see Supplementary Table 3A for

primer sequences and corresponding PCR amplicon sizes). To test for presence or absence of

a GOI, the relevant forward and reverse (F/R) primers were used to amplify a fragment of the

ORF from genomic DNA from the parental cell line (positive control) and genomic DNA

from the deletion mutant. A PCR reaction targeting a "housekeeping" gene, here PF16

(LmxM.20.1400) or IFT88 (LmxM.27.1130), was run in parallel as a technical control.

Master mixes for the PCR reactions constituted the FastGene Optima HotStart Ready Mix

sourced from Nippon Genetics containing 4 ng/µl of the relevant gDNA and 2.5 µM each of

the forward and reverse primers. For all diagnostic PCRs, the following PCR program steps

were used. Step 1: First denaturation at 95^{o}C for 3 minutes. Step 2: denaturation at 95^{o}C for

15 seconds. Step 3: 58^{o}C for 15 seconds. Step 4: 72^{o}C for 30 seconds. Step 5: repeat steps 2-4

for 30 cycles. Step 6: 72^{o}C for 1 minute.

Detergent extraction of cell membranes.

Promastigote cytoskeletons were prepared from 1 – 5 ml (depending on the cell culture density assessed by eye) of mid-log cells. Cells were washed once with PBS and the washed pellet was resuspended in 100 µl (or 1 ml if > 1E7 cells are in the sample) PEME [100 mM Pipes, 2 mM EGTA, 1 mM MgSO4 and 100 nM EDTA] with 0.1% NP-40. Once homogenous and the solution turns clear, the sample was spun at 2ºC at 10,000g for 5 minutes. The supernatant was removed and the pellets were washed with PEME and then resuspended in PEME or PBS with 10 µg/ml Hoechst 33342. For the experiments assessing the fluorescent *Lmx*PKAC1 signal in cytoskeletons, the treatment substance was added to the PEME used during the final wash step at a 1:1000 ratio from a stock solution at 1000x final concentration (cAMP (Sigma-Aldrich A6885), at 1 and 100 mM in water; toyocamycin (Sigma-Aldrich T3580) at 1 and 100 mM in DMSO; inosine (ThermoFisher Scientific A14459); 1 and 100 mM in water). If the substance was dissolved in DMSO, then the respective amount of DMSO (0.1%) was added to the control cytoskeletons in PEME buffer.

Light microscopy.

For live cell fluorescence microscopy of promastigote cells, 1 ml of a mid-logarithmic culture (between 1×10^6 – 1×10^7 cells/ml) was spun at 800g for 5 minutes and the pellet resuspended in 100 µl of PBS and [10 µg/ml] Hoechst 33342 (ThermoFischer). A volume of 0.7 µl was spread onto a poly-lysine coated glass slide, bordered (~ 5 x 2 cm rectangle) with a hydrophobic fat pen (Dako). A glass coverslip, thickness 1 or 1.5 was used to cover the slide. Live cell imaging was typically done with a Zeiss Axioimager.Z2 microscope with a Hamamatsu ORCA-Flash4.0 camera, x63 oil objective lens with numerical aperture (NA) 1.4, and phase contrast illumination. Fluorescence channel was acquired with between 0.5 and 3 seconds of excitation depending on the brightness of the fluorescent signal.

For live cell imaging of axenic amastigotes, 500 µl of mid-log promastigote culture was inoculated into a vented 25 cm^3 flask (total 5 ml volume) containing Schneider's Drosophila medium (Life Technologies) supplemented with 20% heat-inactivated foetal bovine serum (FBS) (Gibco) and adjusted to pH 5.5 with concentrated HCl. Cultures were incubated at 34ºC with 5% CO_2 for 72 hours prior to imaging. 1 ml of sample culture was pelleted by centrifugation at 800g for 5 minutes, pellet washed with PBS, spun again, then the washed pellet resuspended in 50 µl of PBS plus Hoechst 33342 (final 10 µg/ml). A volume of 20 µl was placed on the poly-lysine coated glass slide and left inside a humid chamber for 10

minutes to settle. Excess liquid was gently removed by pipetting and the cover slide placed on

top.

Expansion microscopy

The protocol for ultrastructural expansion microscopy was adapted from (Gambarotto, Hamel,

and Guichard 2021). Five million exponentially growing *Leishmania* promastigotes were

washed twice in PBS before settling on a 12 millimetre glass coverslip in a 24-well plate for

10 minutes. Excess liquid was removed, and cells were fixed with 2 % formaldehyde (FA)

(ThermoFisher, 28906) for 10 minutes. Samples were permeabilized using 0.1% Triton X-100

for 1 minute before washing twice with PBS. Samples were blocked with 2% blocking buffer

(2% bovine albumin in PBS) for 10 minutes with gentle shaking before being incubated with

primary antibodies (Supplementary Table 3B) in blocking buffer overnight at room

temperature. After three washes with 0.1% Tween in PBS (PBS-T), samples were incubated

with secondary antibodies in blocking buffer for 2 hours at room temperature with gentle

shaking, and all subsequent steps were performed in the dark. For samples destined for

microscopy quality control check, Hoechst 33342 (Thermofisher 62249) was added with the

secondary antibody at 20 µg/ml. After staining was confirmed to be high quality and specific

(i.e. no corresponding signal observed in the untagged parental control), samples were

anchored in 2% acrylamide (Merck, A4058), 1.4 % FA for 2 hours at 37 °C before being

washed with PBS. Sample gelation was performed in an ice-filled humid chamber on top of a

parafilm layer. Ammonium persulfate (APS) (Merk, A3678) and N,N,N´,N´-

tetramethylethylenediamine (TEMED) (Merk, T9281) were added to monomer solution to a

final concentration of 0.5 % each, vortexed briefly, then the sample coverslip placed sample-

side-down onto 35 µl gelation solution atop the parafilm. After 5 minutes incubation on ice,

samples were transferred to 37 °C for 30 minutes before removed from the parafilm and

added to 1 ml of denaturation buffer (200 mM SDS, 200 mM NaCl, 50 mM Tris in water,

balanced to pH = 9 using NaOH) in a 6 well-plate. After 10 minutes of gentle shaking, the

glass coverslips were discarded and the gelated sample transferred to a 2 ml Eppendorf with 1

ml fresh denaturation buffer and heated to 95 °C for 30 minutes. The gel was washed thrice in

5 ml PBS before a quarter was cut out and incubated at room temperature overnight in

guineapig anti alpha-tubulin primary antibody (ABCD Antibodies, ABCD_AA345) in

blocking buffer for post staining. The gel sample was then washed thrice in PBS-T then

incubated for 2 hours at room temperature with the corresponding anti guineapig secondary

antibody (Fisher Scientific, 10624773) in blocking buffer. A final three washes with PBS

were done prior to placing the sample in 50 ml of Mili-Q water. The water was changed every
20 minutes until the gel had reached maximum expansion (estimated to be ~4.5 fold based on
axoneme width). Lastly the gel was mounted onto a poly-L-lysine coated round sample holder
and imaged with a Nikon Ti2 Cicero spinning disk microscope using the 100X objective (NA
1.45) and the Kinetix sCMOS Camera. Images were further deconvolved using Huygens
software version 3.10 (Scientific Volume Imaging, The Netherlands, http://svi.nl) using the
following parameters (Acuity: 0, SNR:28,27,20, 30 iterations, background estimation: auto),
then the maximum intensity z-projection taken unless otherwise indicated.

Cell swimming speed and directionality

Cell swimming speed and directionality were measured from 512 frame-long darkfield videos
of cell cultures in a 0.1 mm deep chamber, captured at 5 frames per second (fps) using a
ZeissAxioimager.Z2 microscope with 10× objective and an Andor Neo 5.5 camera, exactly
as described in (Wheeler 2017) with the modification described in (Beneke et al. 2019).

Population overview waveform analysis (POWA).

To quantify the proportion of cells undergoing different waveforms, 1 ml of exponential
growth phase promastigote cells (1×10^6 – 1×10^7 cells/ml) was centrifuged for 5 minutes at
800g. Between 600 – 900 µl supernatant was removed and the cells resuspended in the
remaining 400 – 100 µl of M199 medium, effectively concentrating the cells. 1 µl of 1:100
bead mix (Sigma 79633 polystyrene microparticle beads (5 µm size) in M199 medium) was
added to cell suspension and mixed gently until homogenous. 1 µl of cell sample was added
to the centre of a glass microscope slide bordered by a Dako hydrophobic pen (~5 cm by 2 cm
rectangle). A glass coverslip of 1.0 or 1.5 thickness bordered by a similar Dako hydrophobic
pen rectangle was placed gently on top of the sample slide, effectively creating a sealed
chamber. Videomicrographs of swimming cells under phase contrast illumination were
captured with an Andor Neo 5.5 camera at 200 frames/s for 0.5 seconds, using a ×20 NA 0.3
objective lens on a Leica Inverted microscope. Manual categorisation of waveforms was
performed in ImageJ or Fiji (Collins 2007; Schindelin et al. 2012) for non-dividing cells only.
Categories include symmetric wave tip-to-base (continuous and interrupted), asymmetric
wave base-to-tip and uncoordinated which is neither symmetric tip-to-base nor asymmetric
base-to-tip. Flagellar length was measured from stills obtained from these videomicrographs,
using the Fiji freehand line measuring tool.

Waveform parameter analysis.

The capture of videomicrographs for waveform analysis was exactly as above (POWA), with the exception of using a ×100 NA 1.4 oil immersion objective and capturing for 5 seconds duration. All videomicrographs were captured within 10 minutes of making sample slides. Tip-to-base beat waveforms were analysed in ImageJ, using custom scripts (https://github.com/zephyris/flagellar-beat-fourier-ij). Flagella were traced using the midline method as described in (Walker, Ishimoto, and Wheeler 2019), except the cell body was manually identified. Data was represented as a tangent angle kymograph, plotting tangent angle at each position in the flagellum over time. The resulting tangent-angle kymographs were manually cropped to a length of either 128 (0.64 seconds) or 256 (1.28 seconds) frames. Each datapoint on the plots represent one kymograph (cropped to the lengths described above), where several datapoints (up to three) could derive from one cell.

Tangent angle kymographs were analysed using a Fourier transform-based method to extract beat parameters, using the concepts described in (Walker et al. 2019). The Fourier power spectrum of the tangent angle was calculated and summed over the length of the flagellum. The dominant beat frequency was obtained by determining the frequency with the greatest power from the power spectrum using a sliding window with a width of 10% of the measurable frequency range.

Data obtained from Fourier analysis were compiled in Excel and plotted using GraphPad Prism 9. Amplitude measurements along each flagellum (computed for the identified dominant frequency from the FFT and defined in terms of the tangent angle) were averaged and plotted against the dominant frequency. The number of complete sinusoidal waves present along the flagellum (equivalent to the wavelength divided by the flagellum length) was calculated in Excel by taking the difference in phase of the identified dominant frequency between the proximal and distal ends of the flagellum, then dividing the phase difference by 2π. This analysis effectively approximates the flagellar waveform as a single sinusoidal mode (when expressed in terms of tangent angle), as justified by observations of the wild type (Walker et al. 2019).

Acknowledgements

A part of this work was carried out at the Wellcome Centre for Integrative Parasitology (WCIP) at the University of Glasgow and we thank the WCIP for hosting SF as a visiting

student. We thank Keith Gull (University of Oxford) for the gift of antibody L8C4 and access

to laboratory equipment at the Dunn School, Clirim Jetishi (Ochsenreiter lab, University of

Bern) for help with ultrastructure expansion microscopy and Michael Boshart for stimulating

discussions.

Funding statement

EG was supported by a Royal Society University Research Fellowship (UF160661).

SF was supported by a Royal Society Enhancement Award (RGF\EA\180189) and

Studentship funding from the Sir William Dunn School of Pathology, University of Oxford.

RJW was supported by the Sir Henry Dale Fellowship (Nuffield Department of Medicine)

[211075/Z/18/Z].

BJW was supported by the Royal Commission for the Exhibition of 1851.

We acknowledge support for this project for EG and SF through the WCIP core Wellcome

Centre Award [104111/Z/14/Z], through the Wellcome Trust grant [104627/Z/14/Z] to Keith

Gull; https://wellcome.org) and a project grant from the Swiss National Science foundation

(project number: 320030L-227939).

This research was funded in whole, or in part by the Wellcome Trust [Grant number

211075/Z/18/Z, 104111/Z/14/Z, 104627/Z/14/Z]. For the purpose of open access, the author

has applied a CC BY public copyright licence to any Author Accepted Manuscript version

arising from this submission.

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Table 1. Accession numbers for all PKAR and PKAC orthologues found encoded in the

genomes of *L. mexicana, L. major, L. donovani, T. brucei* and *T. cruzi*. Each row represents

genes found in a syntenic locus across kinetoplastid species, as illustrated in Supplementary

Figure 1A-D.

	L. mexicana (Lmx)	*L. major* (Lmj)	*L. donovani* (Ld)		*T. brucei* (Tb)	*T. cruzi* (Tc)
PKAR1	LmxM.13.0160	LmjF.13.0160	LdCL_130006600	**PKAR1**	Tb927.11.4610	TcCLB.506227.150
PKAR3	LmxM.33.2820	LmjF.34.2820	LdCL_340034800	**PKAR3**	n/a	TcCLB.510879.50
PKAC1	LmxM.34.4010	LmjF.35.4010	LdCL_350045700	**PKAC2**	Tb927.9.11100	TcCLB.508461.310
PKAC2	LmxM.34.3960	LmjF.35.3960	LdCL_350045200	**PKAC1**	Tb927.9.11030	TcCLB.508461.280
PKAC3	LmxM.18.1080	LmjF.18.1080	LdCL_180016100	**PKAC3**	Tb927.10.13010	TcCLB.509805.10

Figure Legends

Figure 1. Alignments of PKAR sequences.

(A) Cartoon representation of trypanosomatid PKAR proteins *Ld*PKAR1
(LdBPK.13.2.000160), *Lmj*PKAR1 (LmjF.13.0160), *Lmx*PKAR1 (LmxM.13.0160),
*Tb*PKAR1 (Tb927.11.4610), *Tc*PKAR1 (TcCLB.506227.150), *Ld*PKAR3
(LdBPK.34.2.002680), *Lmj*PKAR3 (LmjF.34.2820), *Lmx*PKAR3 (LmxM.33.2820),
*Tc*PKAR3 (TcCLB.510879.50), and Homo sapiens cAMP-dependent protein kinase type I-
alpha regulatory subunit PKAR-Iα (P10644, KAP0_HUMAN) and type II-alpha regulatory
subunit PKAR-IIα (P13861|KAP2_HUMAN). LRR, leucine rich repeat; DD, dimerization
docking domain; CNBD, cyclic nucleotide binding domains (PFAM ID: PF00027); AI,
autoinhibitory domain sequence (Celeste E. Poteet-Smith et al. 1997). **(B)** AI sequence
alignment. Bold letters highlight the RRxA and RRxT motifs which form a pseudo-substrate
or are autophosphorylated by docked catalytic subunits in mammalian PKA heterotetramers.
A universally conserved arginine residue is highlighted in red. The conserved RGAIS
residues in *Leishmania* PKAR3 proteins and the human AI sequence are underlined. **(C)**
CNBD: Underlined is the canonical phosphate binding cassette (PBC) motif
FGELAL[ILVM]xxxPRAA (Peng et al. 2015b); residues matching the PBC are in bold black
type. Highlighted in blue (asterisks) are residues associated with altered ligand specificity in
trypanosomatid PKAR1 (Bachmaier et al. 2019b): the switch from alanine (position 5) in
human PKA to glutamate in trypanosomatid and the switch from arginine (position 12) in
human PKA to a alanine, valine, serine, threonine, or asparagine in the trypanosomatid PKAR
sequences. Orange asterisk indicate a valine residue reported to form part of the
trypanosomatid specific hydrophobic pocket. **(D)** Hydrophobic pocket: orange asterisk
indicates a residue reported to form part of the *Trypanosoma* specific hydrophobic pocket.
Highlighted in orange type is a lysine residue reported to increase affinity to binding synthetic
nucleoside analogue 7-cyano-7-deazainosine (Bachmaier et al. 2019b).

Figure 2. Localisations of tagged *L. mexicana* PKAC and PKAR subunits.

Micrographs of cells expressing mNG-tagged PKAC or PKAR subunits. **(A)**
mNG::*Lmx*PKAC1 (LmxM.34.4010), **(B)** mNG::*Lmx*PKAC2 (LmxM.34.3960), **(C)** mNG::
*Lmx*PKAC3 (LmxM.18.1080), **(D)** *Lmx*PKAR1::mNG (LmxM.13.0160), **(E)**

*Lmx*PKAR3::mNG (LmxM.33.2820). Panels in A-E show live promastigotes (left), detergent-

extracted promastigote cytoskeletons (middle) and live axenic amastigotes (right). For each

sample type a merged fluorescence with phase contrast image is shown, and the combined

mNG and Hoechst fluorescence. **(F)** Promastigotes in different stages of the cell cycle,

expressing *Lmx*PKAR1::mNG and **(G)** mNG::*Lmx*PKAR3. White arrows indicate dynamic

localization of mNG::*Lmx*PKAR3 to posterior cell body pole. F, flagella; N, nuclei; N*,

dividing nucleus, K, kinetoplasts. Scale bar indicates 5 µm.

Figure 3. Localisation dependencies of *L. mexicana* PKAC and PKAR subunits.

(A) Measurement of the mNG fluorescent signal intensity across a line though the cell body

(cartoon) show a clear edge effect for the majority of mNG::*Lmx*PKAC3 cells (i) (black lines,

n = 11); the red lines show a minority of cells (n = 3) with a diffuse cell body signal. An edge

effect was detected for all measured *Lmx*PKAR3::mNG cells (ii) (n = 12). The edge effect for

mNG::*Lmx*PKAC3 was lost following *Lmx*PKAR3 deletion (iii) (n = 10). **(B)** Quantitation of

fluorescence signal intensity in cells expressing *Lmx*PKAR3::mNG, mNG::*Lmx*PKAC3 and

in cells expressing mNG::*Lmx*PKAC3 following *Lmx*PKAR3 deletion. Measurements were

taken at six points along the flagellum and cell body as shown in the cartoon.

(C-F) Micrographs showing the localisation of mNG-tagged *Lmx*PKAC subunits following

deletion of *Lmx*PKAR subunits: **(C)** mNG::*Lmx*PKAC1 *ΔLmxPKAR1*; **(D)**

mNG::*Lmx*PKAC2 *ΔLmxPKAR1*; **(E)** mNG::*Lmx*PKAC3 *ΔLmxPKAR1*; **(F)**

mNG::*Lmx*PKAC3 *ΔLmxPKAR3*.

Figure 4. Localisation of LmxPKAR1 to the paraflagellar rod.

Ultrastructure expansion microscopy of promastigotes expressing LmxPKAR1::mNG::c-myc.

(A) A field of view depicting expanded promastigote cells with fluorescent signal

corresponding to *Lmx*PKAR1 (anti-c-myc antibody, pseudocoloured green), PFR2 (anti-PFR2

antibody L8C4, pseudocoloured red), alpha-tubulin (anti-alpha-tubulin antibody,

pseudocoloured grey). Insets show close-up views of flagellar segments where the

LmxPKAR1 signal is seen between the PFR2 and tubulin signals. Abbreviations: CB, cell

body; F, flagellum. **(B)** Flagellar cross sections showing the *Lmx*PKAR1 signal sandwiched

between the PFR2 and tubulin signals. **(C)** Selection of micrographs showing the start of the

*Lmx*PKAR1 signal before the PFR2 signal in the flagellar pocket. **(D)** Views showing the

*Lmx*PKAR1 signal extending beyond the PFR2 signal in the distal flagellum. These were

computationally straightened in ImageJ. Open arrows point to the distal end of the

*Lmx*PKAR1 signal, filled arrows to the end of the PFR2 signal. **(E)** Cartoon of a promastigote

flagellum cross-section showing the location of *Lmx*PKAR1, which may be anchored in the

inner/middle domain of the PFR and/or at the PFR-axonemal connectors (PACs).

Figure 5. Pharmacological displacement of *Lmx*PKAC1 signal from the flagellum.

Micrographs showing detergent-extracted promastigote cytoskeletons from

mNG::*Lmx*PKAC1-expressing cells **(A)** without treatment (DMSO only control), **(B)** after

treatment with 100 µM cAMP, and 10 µM **(C)** adenosine, **(D)** guanine and **(E)** inosine,

respectively. Left panels, bright field images; right panels, merged fluorescence images of the

mNG signal (green) and the signal from the DNA stain Hoechst (magenta). Scale bar, 20 µm.

(F) Quantification of the cytoskeleton-associated mNG::*Lmx*PKAC1 signals from cells

treated with 0.1% DMSO, 100 µM cAMP, 25 µM toyocamycin, 10 µM adenosine, 10 µM

guanine, 10 µM inosine; untagged parental cells served as negative control. **(G)**

Quantification of the cytoskeleton-associated mNG::*Lmx*PKAC1 signals in cells treated with

inosine concentrations ranging from 10 µM to 10 nM. For all flagellar signal quantification,

background fluorescence was subtracted from maximum intensity values. Each data point

represents one cell measured across the flagellum midpoint. Horizontal bar indicates mean

corrected fluorescent intensity. Significance was determined using an unpaired t-test

assuming equal variance. *** represents a p-value < 0.001. n > 50 cell measured for each

condition. **(H)** Normalisation of datapoints in (G) with a non-linear regression and calculated

EC_{50}. See Supplementary Table 4 for raw data.

Figure 6. Mutant population motility. **(A)** Scatter plot of mean population swim speed and

directionality (velocity/speed) of parental controls and deletion mutants for *Lmx*PKAC and

*Lmx*PKAR subunits and of dual knock outs of *Lmx*PKAC1/PKAC2 and *Lmx*PKAC1/PKAR3.

Swim speed measurement data from at least three replicates per population. Error bars

indicate one standard deviation of the speed and directionality measurements. Raw data with

p-values from unpaired t-tests for significance are summarised in Supplementary Table 1. **(B)**

Phase contrast micrographs overlaid with real traces of flagellum position over one beat cycle,

from a cell propagating either the (upper) symmetric tip-to-base flagellar wave for forward

motility, or (lower) the asymmetric base-to-tip ciliary waveform for tumbling/turning. On the

right are cartoon summaries of symmetric and asymmetric waves with a hollow arrow

indicating the direction of cell displacement, and a filled grey arrow indicating direction of

wave travel. **(C)** Relative instances of the waveforms observed in *ΔLmxPKAC* and

ΔLmxPKAR mutants and parental control populations. **(D)** Computer-generated waveform

reconstruction exemplified for each mutant, highlighting the lack of symmetry in

ΔLmxPKAC1 flagellar movement.

Figure 7. Symmetric wave amplitude and frequency parameters. Fourier decomposition

of the tip-to-base uncoordinated wave attempts of *ΔLmxPKAC1,* and the symmetric flagellar

waves of *ΔLmxPKAC2, ΔLmxPKAC3, ΔLmxPKAR1 and ΔLmxPKAR3*. **(A-E)** Scatter plots

of the dominant beat frequency in hertz (Hz) plotted against beat amplitude in degrees for **(A)**

ΔLmxPKAC1, **(B)** *ΔLmxPKAC2,* **(C)** *ΔLmxPKAC3,* **(D)** *ΔLmxPKAR1* **(E)** *ΔLmxPKAR3*

populations (purple diamonds) overlaid with measurements from parental control cells (black

filled circles). Each point represents one unit of time observed (see Methods). **(F)** All

datapoints across (A-E) representing beat frequency and amplitude represented in one scatter

plot. **(G)** Tangent kymograph (i) and flagellar traces (ii) of a parental control cell; distance is

on the x-axis, time one the y-axis. **(H)** Tangent kymograph (i) and flagellar traces (ii) of a

ΔLmxPKAC1 cell. Tangent kymographs follow 1 second of flagellar beating, and z-

projections of 150 ms of beating with overlayed traces of flagellum position every 25 ms

where rainbow coloration indicates start (red) to end (indigo) of projected time. **(I)** Summary

numbers for each deletion mutant cell line: Number (n) of timepoints captured by high-speed

video microscopy, where each timepoint represents 256 video frames, and n cells captured in

total; the total time observed for each mutant dataset in seconds. The mean angular amplitude

(amp) of the waveform in degrees (deg); mean beat frequency (beats per second, bps; Hz);

number of waves per flagellum (w/f); flagellum length in µm and number (n) of cells for

which length was measured. Mann-Whitney **U** test was used to ascertain significance of each

dataset against that of the parental control measurements. Significant p-values are indicated as,

*** = p < 0.001, ** = p < 0.005, * = p < 0.05. All p-values can be found in Supplementary

Figure 7. See Supplementary Table 5 for raw data.

References

Alonso, Carlos Agustín I., Claudia E. Osycka-Salut, Luciana Castellano, Andreína Cesari, Nicolás Di Siervi, Adrián Mutto, Anders Johannisson, Jane M. Morrell, Carlos Davio, and Silvina Perez-Martinez. 2017. 'Extracellular cAMP Activates Molecular Signalling Pathways Associated with Sperm Capacitation in Bovines'. *Molecular Human Reproduction* 23 (8): 521–34. https://doi.org/10.1093/molehr/gax030.

Alvarez-Jarreta, Jorge, Beatrice Amos, Cristina Aurrecoechea, Saikou Bah, Matthieu Barba, Ana Barreto, Evelina Y. Basenko, et al. 2024. 'VEuPathDB: The Eukaryotic Pathogen, Vector and Host Bioinformatics Resource Center in 2023'. *Nucleic Acids Research* 52 (D1): D808–16. https://doi.org/10.1093/nar/gkad1003.

Alves, Aline A., Heloisa B. Gabriel, Maria J. R. Bezerra, Wanderley de Souza, Sue Vaughan, Narcisa L. Cunha-E-Silva, and Jack D. Sunter. 2020. 'Control of Assembly of Extra-Axonemal Structures: The Paraflagellar Rod of Trypanosomes'. *Journal of Cell Science* 133 (10): jcs242271. https://doi.org/10.1242/jcs.242271.

Amos, Beatrice, Cristina Aurrecoechea, Matthieu Barba, Ana Barreto, Evelina Y. Basenko, Wojciech Bażant, Robert Belnap, et al. 2022. 'VEuPathDB: The Eukaryotic Pathogen, Vector and Host Bioinformatics Resource Center'. *Nucleic Acids Research* 50 (D1): D898–911. https://doi.org/10.1093/nar/gkab929.

Anand, Ganesh S., Matthew Hotchko, Simon H.J. Brown, Lynn F. Ten Eyck, Elizabeth A. Komives, and Susan S. Taylor. 2007. 'R-Subunit Isoform Specificity in Protein Kinase A: Distinct Features of Protein Interfaces in PKA Types I and II by Amide H/2H Exchange Mass Spectrometry'. *Journal of Molecular Biology* 374 (2): 487–99. https://doi.org/10.1016/j.jmb.2007.09.035.

Aoki, F., S. Sakai, and K. Kohmoto. 1999. 'Regulation of Flagellar Bending by cAMP and Ca2+ in Hamster Sperm'. *Molecular Reproduction and Development* 53 (1): 77–83. https://doi.org/10.1002/(SICI)1098-2795(199905)53:1<77::AID-MRD9>3.0.CO;2-C.

Bachmaier, Sabine, Yuri Volpato Santos, Susanne Kramer, George Boniface Githure, Thomas Klöckner, Julia Pepperl, Cordula Baums, et al. 2019a. 'Nucleoside Analogue Activators of Cyclic AMP-Independent Protein Kinase A of Trypanosoma'. *Nature Communications* 10 (March):1421. https://doi.org/10.1038/s41467-019-09338-z.

Bachmaier, Sabine, Ronit Witztum, Polina Tsigankov, Roni Koren, Michael Boshart, and Dan Zilberstein. 2016. 'Protein Kinase A Signaling during Bidirectional Axenic Differentiation in Leishmania'. *International Journal for Parasitology* 46 (2): 75–82. https://doi.org/10.1016/j.ijpara.2015.09.003.

Baker, N., C. M. C. Catta-Preta, R. Neish, J. Sadlova, B. Powell, E. V. C. Alves-Ferreira, V. Geoghegan, et al. 2021. 'Systematic Functional Analysis of Leishmania Protein Kinases Identifies Regulators of Differentiation or Survival'. *Nature Communications* 12 (1): 1244. https://doi.org/10.1038/s41467-021-21360-8.

Balbach, Melanie, Vera Beckert, Jan N. Hansen, and Dagmar Wachten. 2018. 'Shedding Light on the Role of cAMP in Mammalian Sperm Physiology'. *Molecular and Cellular Endocrinology* 468 (June):111–20. https://doi.org/10.1016/j.mce.2017.11.008.

Banky, Poopak, Lily Jun-Shen Huang, and Susan S. Taylor. 1998. 'Dimerization/Docking Domain of the Type Iα Regulatory Subunit of cAMP-Dependent Protein Kinase: REQUIREMENTS FOR DIMERIZATION AND DOCKING ARE DISTINCT BUT OVERLAPPING *'. *Journal of Biological Chemistry* 273 (52): 35048–55. https://doi.org/10.1074/jbc.273.52.35048.

Barkalow, K., T. Hamasaki, and P. Satir. 1994. 'Regulation of 22S Dynein by a 29-kD Light Chain'. *The Journal of Cell Biology* 126 (3): 727–35. https://doi.org/10.1083/jcb.126.3.727.

Beltrán, C., O. Zapata, and A. Darszon. 1996. 'Membrane Potential Regulates Sea Urchin
Sperm Adenylylcyclase'. *Biochemistry* 35 (23): 7591–98.
https://doi.org/10.1021/bi952806v.
Beneke, Tom, François Demay, Edward Hookway, Nicole Ashman, Heather Jeffery, James
Smith, Jessica Valli, et al. 2019. 'Genetic Dissection of a Leishmania Flagellar
Proteome Demonstrates Requirement for Directional Motility in Sand Fly Infections'.
PLoS Pathogens 15 (6): e1007828. https://doi.org/10.1371/journal.ppat.1007828.
Beneke, Tom, Ross Madden, Laura Makin, Jessica Valli, Jack Sunter, and Eva Gluenz. 2017.
'A CRISPR Cas9 High-Throughput Genome Editing Toolkit for Kinetoplastids'.
Royal Society Open Science 4 (5): 170095. https://doi.org/10.1098/rsos.170095.
Berman, Helen M., Lynn F. Ten Eyck, David S. Goodsell, Nina M. Haste, Alexandr Kornev,
and Susan S. Taylor. 2005. 'The cAMP Binding Domain: An Ancient Signaling
Module'. *Proceedings of the National Academy of Sciences of the United States of*
America 102 (1): 45–50. https://doi.org/10.1073/pnas.0408579102.
Billington, Karen, Clare Halliday, Ross Madden, Philip Dyer, Amy Rachel Barker, Flávia
Fernandes Moreira-Leite, Mark Carrington, et al. 2023. 'Genome-Wide Subcellular
Protein Map for the Flagellate Parasite Trypanosoma Brucei'. *Nature Microbiology* 8
(3): 533–47. https://doi.org/10.1038/s41564-022-01295-6.
Boitz, Jan M., Buddy Ullman, Armando Jardim, and Nicola S. Carter. 2012. 'Purine Salvage
in Leishmania: Complex or Simple by Design?' *Trends in Parasitology* 28 (8): 345–
52. https://doi.org/10.1016/j.pt.2012.05.005.
Broadhead, Richard, Helen R. Dawe, Helen Farr, Samantha Griffiths, Sarah R. Hart, Neil
Portman, Michael K. Shaw, et al. 2006. 'Flagellar Motility Is Required for the
Viability of the Bloodstream Trypanosome'. *Nature* 440 (7081): 224–27.
https://doi.org/10.1038/nature04541.
Brokaw, C. J., and R. Kamiya. 1987. 'Bending Patterns of *Chlamydomonas* Flagella: IV.
Mutants with Defects in Inner and Outer Dynein Arms Indicate Differences in Dynein
Arm Function'. *Cell Motility* 8 (1): 68–75. https://doi.org/10.1002/cm.970080110.
Chan, Tung O., Roger S. Armen, Santosh Yadav, Sushrut Shah, Jin Zhang, Brian C. Tiegs,
Nikhil Keny, et al. 2020. 'A Tripartite Cooperative Mechanism Confers Resistance of
the Protein Kinase A Catalytic Subunit to Dephosphorylation'. *The Journal of*
Biological Chemistry 295 (10): 3316–29. https://doi.org/10.1074/jbc.RA119.010004.
Cheng, Yuhui, Yingkai Zhang, and J. Andrew McCammon. 2006. 'How Does Activation
Loop Phosphorylation Modulate Catalytic Activity in the cAMP-Dependent Protein
Kinase: A Theoretical Study'. *Protein Science: A Publication of the Protein Society*
15 (4): 672–83. https://doi.org/10.1110/ps.051852306.
Christensen, S. T., C. Guerra, Y. Wada, T. Valentin, R. H. Angeletti, P. Satir, and T.
Hamasaki. 2001. 'A Regulatory Light Chain of Ciliary Outer Arm Dynein in
Tetrahymena Thermophila'. *The Journal of Biological Chemistry* 276 (23): 20048–54.
https://doi.org/10.1074/jbc.M008412200.
Chung, Jean-Ju, Kiyoshi Miki, Doory Kim, Sang-Hee Shim, Huanan F. Shi, Jae Yeon Hwang,
Xinjiang Cai, Yusuf Iseri, Xiaowei Zhuang, and David E. Clapham. 2017. 'CatSperζ
Regulates the Structural Continuity of Sperm Ca2+ Signaling Domains and Is
Required for Normal Fertility'. *eLife* 6 (February):e23082.
https://doi.org/10.7554/eLife.23082.
Collins, Tony J. 2007. 'ImageJ for Microscopy'. *BioTechniques* 43 (1S): S25–30.
https://doi.org/10.2144/000112517.
Dahlin, Heather R., Ning Zheng, and John D. Scott. 2021. 'Beyond PKA: Evolutionary and
Structural Insights That Define a Docking and Dimerization Domain Superfamily'.

The Journal of Biological Chemistry 297 (2): 100927. https://doi.org/10.1016/j.jbc.2021.100927.

Dean, Samuel, Jack D. Sunter, and Richard J. Wheeler. 2017. 'TrypTag.Org: A Trypanosome Genome-Wide Protein Localisation Resource'. *Trends in Parasitology* 33 (2): 80–82. https://doi.org/10.1016/j.pt.2016.10.009.

Etchebehere, L. C., M. X. Van Bemmelen, C. Anjard, F. Traincard, K. Assemat, C. Reymond, and M. Véron. 1997. 'The Catalytic Subunit of Dictyostelium cAMP-Dependent Protein Kinase -- Role of the N-Terminal Domain and of the C-Terminal Residues in Catalytic Activity and Stability'. *European Journal of Biochemistry* 248 (3): 820–26. https://doi.org/10.1111/j.1432-1033.1997.t01-2-00820.x.

Etchebehere, Lilian C., Miguel X. P. Van Bemmelen, Christophe Anjard, François Traincard, Karine Assemat, Christophe Reymond, and Michel Veron. 1997. 'The Catalytic Subunit of Dictyostelium cAMP-Dependent Protein Kinase'. *European Journal of Biochemistry* 248 (3): 820–26. https://doi.org/10.1111/j.1432-1033.1997.t01-2-00820.x.

Fischer Weinberger, Renana, Sabine Bachmaier, Veronica Ober, George B. Githure, Ramu Dandugudumula, Isabelle Q. Phan, Michal Almoznino, et al. 2024. 'A Divergent Protein Kinase A Regulatory Subunit Essential for Morphogenesis of the Human Pathogen Leishmania'. *PLoS Pathogens* 20 (3): e1012073. https://doi.org/10.1371/journal.ppat.1012073.

Fort, Cécile, Serge Bonnefoy, Linda Kohl, and Philippe Bastin. 2016. 'Intraflagellar Transport Is Required for the Maintenance of the Trypanosome Flagellum Composition but Not Its Length'. *Journal of Cell Science* 129 (15): 3026–41. https://doi.org/10.1242/jcs.188227.

Gabriel, Heloisa B., Kelly, Ruby., Moreira-Leite, Flavia., and Sunter, Jack D. bioRxiv. 'Dramatic Reduction in Trypanosome Motility Occurs without Large-Scale Changes to Paraflagellar Rod Ultrastructure | bioRxiv'. bioRxiv. https://www.biorxiv.org/content/10.1101/2024.04.19.590284v1.

Gadelha, Catarina, Bill Wickstead, and Keith Gull. 2007. 'Flagellar and Ciliary Beating in Trypanosome Motility'. *Cell Motility and the Cytoskeleton* 64 (8): 629–43. https://doi.org/10.1002/cm.20210.

Gaillard, A. R., D. R. Diener, J. L. Rosenbaum, and W. S. Sale. 2001. 'Flagellar Radial Spoke Protein 3 Is an A-Kinase Anchoring Protein (AKAP)'. *The Journal of Cell Biology* 153 (2): 443–48. https://doi.org/10.1083/jcb.153.2.443.

Gaillard, Anne R., Laura A. Fox, Jeanne M. Rhea, Branch Craige, and Winfield S. Sale. 2006. 'Disruption of the A-Kinase Anchoring Domain in Flagellar Radial Spoke Protein 3 Results in Unregulated Axonemal cAMP-Dependent Protein Kinase Activity and Abnormal Flagellar Motility'. *Molecular Biology of the Cell* 17 (6): 2626–35. https://doi.org/10.1091/mbc.E06-02-0095.

Gambarotto, Davide, Virginie Hamel, and Paul Guichard. 2021. 'Ultrastructure Expansion Microscopy (U-ExM)'. *Methods in Cell Biology* 161:57–81. https://doi.org/10.1016/bs.mcb.2020.05.006.

Halliday, Clare, Karen Billington, Ziyin Wang, Ross Madden, Samuel Dean, Jack Daniel Sunter, and Richard John Wheeler. 2019. 'Cellular Landmarks of Trypanosoma Brucei and Leishmania Mexicana'. *Molecular and Biochemical Parasitology* 230 (June):24–36. https://doi.org/10.1016/j.molbiopara.2018.12.003.

Hamasaki, T., K. Barkalow, J. Richmond, and P. Satir. 1991. 'cAMP-Stimulated Phosphorylation of an Axonemal Polypeptide That Copurifies with the 22S Dynein Arm Regulates Microtubule Translocation Velocity and Swimming Speed in

Paramecium'. *Proceedings of the National Academy of Sciences of the United States of America* 88 (18): 7918–22. https://doi.org/10.1073/pnas.88.18.7918.

Heale, Kali A., and Laleh Alisaraie. 2020. 'C-Terminal Tail of β-Tubulin and Its Role in the Alterations of Dynein Binding Mode'. *Cell Biochemistry and Biophysics* 78 (3): 331–45. https://doi.org/10.1007/s12013-020-00920-7.

Heuser, Thomas, Milen Raytchev, Jeremy Krell, Mary E. Porter, and Daniela Nicastro. 2009. 'The Dynein Regulatory Complex Is the Nexin Link and a Major Regulatory Node in Cilia and Flagella'. *The Journal of Cell Biology* 187 (6): 921–33. https://doi.org/10.1083/jcb.200908067.

Hyams, J. S., and G. G. Borisy. 1978. 'Isolated Flagellar Apparatus of Chlamydomonas: Characterization of Forward Swimming and Alteration of Waveform and Reversal of Motion by Calcium Ions in Vitro'. *Journal of Cell Science* 33 (October):235–53. https://doi.org/10.1242/jcs.33.1.235.

Ishijima, Sumio. 2015. 'Ca2+ and cAMP Regulations of Microtubule Sliding in Hyperactivated Motility of Bull Spermatozoa'. *Proceedings of the Japan Academy. Series B, Physical and Biological Sciences* 91 (3): 99–108. https://doi.org/10.2183/pjab.91.99.

Jäger, Adriana V., Javier G. De Gaudenzi, Jesica G. Mild, Bárbara Mc Cormack, Sergio Pantano, Daniel L. Altschuler, and Martin M. Edreira. 2014. 'Identification of Novel Cyclic Nucleotide Binding Proteins in Trypanosoma Cruzi'. *Molecular and Biochemical Parasitology* 198 (2): 104–12. https://doi.org/10.1016/j.molbiopara.2015.02.002.

Kinukawa, Masashi, Shoji Oda, Yoshiyuki Shirakura, Masaki Okabe, Junko Ohmuro, Shoji A. Baba, Masao Nagata, and Fugaku Aoki. 2006. 'Roles of cAMP in Regulating Microtubule Sliding and Flagellar Bending in Demembranated Hamster Spermatozoa'. *FEBS Letters* 580 (5): 1515–20. https://doi.org/10.1016/j.febslet.2006.01.078.

Kobe, B., and A. V. Kajava. 2001. 'The Leucine-Rich Repeat as a Protein Recognition Motif'. *Current Opinion in Structural Biology* 11 (6): 725–32. https://doi.org/10.1016/s0959-440x(01)00266-4.

Kohl, L., T. Sherwin, and K. Gull. 1999. 'Assembly of the Paraflagellar Rod and the Flagellum Attachment Zone Complex during the Trypanosoma Brucei Cell Cycle'. *The Journal of Eukaryotic Microbiology* 46 (2): 105–9. https://doi.org/10.1111/j.1550-7408.1999.tb04592.x.

Kutomi, Osamu, Manabu Hori, Masaki Ishida, Takashi Tominaga, Hiroyuki Kamachi, France Koll, Jean Cohen, Norico Yamada, and Munenori Noguchi. 2012. 'Outer Dynein Arm Light Chain 1 Is Essential for Controlling the Ciliary Response to Cyclic AMP in Paramecium Tetraurelia'. *Eukaryotic Cell* 11 (5): 645–53. https://doi.org/10.1128/EC.05279-11.

Lefièvre, Linda, Kula N. Jha, Eve de Lamirande, Pablo E. Visconti, and Claude Gagnon. 2002. 'Activation of Protein Kinase A during Human Sperm Capacitation and Acrosome Reaction'. *Journal of Andrology* 23 (5): 709–16.

Leung, Miguel Ricardo, Chen Sun, Jianwei Zeng, Jacob R. Anderson, Qingwei Niu, Wei Huang, Willem E. M. Noteborn, Alan Brown, Tzviya Zeev-Ben-Mordehai, and Rui Zhang. 2025. 'Structural Diversity of Axonemes across Mammalian Motile Cilia'. *Nature* 637 (8048): 1170–77. https://doi.org/10.1038/s41586-024-08337-5.

Lindemann, Charles B., and Kathleen A. Lesich. 2010. 'Flagellar and Ciliary Beating: The Proven and the Possible'. *Journal of Cell Science* 123 (Pt 4): 519–28. https://doi.org/10.1242/jcs.051326.

Lodh, Sukanya, Junji Yano, Megan S. Valentine, and Judith L. Van Houten. 2016. 'Voltage-Gated Calcium Channels of Paramecium Cilia'. *The Journal of Experimental Biology* 219 (19): 3028–38. https://doi.org/10.1242/jeb.141234.

Ma, Weiyuan, Shai D. Silberberg, and Zvi Priel. 2002. 'Distinct Axonemal Processes Underlie Spontaneous and Stimulated Airway Ciliary Activity'. *The Journal of General Physiology* 120 (6): 875–85. https://doi.org/10.1085/jgp.20028695.

Mukhopadhyay, Aakash Gautam, and Chinmoy Sankar Dey. 2016. 'Reactivation of Flagellar Motility in Demembranated Leishmania Reveals Role of cAMP in Flagellar Wave Reversal to Ciliary Waveform'. *Scientific Reports* 6 (1): 37308. https://doi.org/10.1038/srep37308.

Mukhopadhyay, Aakash Gautam, and Chinmoy Sankar Dey. 2017. 'Role of Calmodulin and Calcineurin in Regulating Flagellar Motility and Wave Polarity in Leishmania'. *Parasitology Research* 116 (11): 3221–28. https://doi.org/10.1007/s00436-017-5608-6.

N M Bonini, D L Nelson. 1988. 'Differential Regulation of Paramecium Ciliary Motility by cAMP and cGMP'. *The Journal of Cell Biology* 106 (5): 1615–23. https://doi.org/10.1083/jcb.106.5.1615.

Nakachi, Mia, Ayako Nakajima, Mamoru Nomura, Kouki Yonezawa, Keisuke Ueno, Toshinori Endo, and Kazuo Inaba. 2011. 'Proteomic Profiling Reveals Compartment-Specific, Novel Functions of Ascidian Sperm Proteins'. *Molecular Reproduction and Development* 78 (7): 529–49. https://doi.org/10.1002/mrd.21341.

Nasir, Adeel, Peter Rolf Richter, Aude Le Bail, Viktor Daiker, Julia Stoltze, Binod Prasad, Sebastian Michael Strauch, and Michael Lebert. 2022. 'Molecular Cross-Talk between Gravity- and Light-Sensing Mechanisms in Euglena Gracilis'. *International Journal of Molecular Sciences* 23 (5): 2776. https://doi.org/10.3390/ijms23052776.

Nlend, Marie-Christine, Andreas Schmid, Zoltan Sutto, George A. Ransford, Gregory E. Conner, Nevis Fregien, and Matthias Salathe. 2007. 'Calcium-Mediated, Purinergic Stimulation and Polarized Localization of Calcium-Sensitive Adenylyl Cyclase Isoforms in Human Airway Epithelia'. *FEBS Letters* 581 (17): 3241–46. https://doi.org/10.1016/j.febslet.2007.06.015.

Ober, Veronica Teresa, George Boniface Githure, Yuri Volpato Santos, Sidney Becker, Gabriel Moya Munoz, Jérôme Basquin, Frank Schwede, Esben Lorentzen, and Michael Boshart. 2024. 'Purine Nucleosides Replace cAMP in Allosteric Regulation of PKA in Trypanosomatid Pathogens'. *eLife* 12 (March):RP91040. https://doi.org/10.7554/eLife.91040.

Oberholzer, Michael, Gerasimos Langousis, HoangKim T. Nguyen, Edwin A. Saada, Michelle M. Shimogawa, Zophonias O. Jonsson, Steven M. Nguyen, James A. Wohlschlegel, and Kent L. Hill. 2011. 'Independent Analysis of the Flagellum Surface and Matrix Proteomes Provides Insight into Flagellum Signaling in Mammalian-Infectious Trypanosoma Brucei'. *Molecular & Cellular Proteomics : MCP* 10 (10): M111.010538. https://doi.org/10.1074/mcp.M111.010538.

Pazour, Gregory J., Nathan Agrin, John Leszyk, and George B. Witman. 2005. 'Proteomic Analysis of a Eukaryotic Cilium'. *The Journal of Cell Biology* 170 (1): 103–13. https://doi.org/10.1083/jcb.200504008.

Peng, Mao, Thin Thin Aye, Berend Snel, Bas van Breukelen, Arjen Scholten, and Albert J. R. Heck. 2015a. 'Spatial Organization in Protein Kinase A Signaling Emerged at the Base of Animal Evolution'. *Journal of Proteome Research* 14 (7): 2976–87. https://doi.org/10.1021/acs.jproteome.5b00370.

Poteet-Smith, C. E., J. B. Shabb, S. H. Francis, and J. D. Corbin. 1997. 'Identification of Critical Determinants for Autoinhibition in the Pseudosubstrate Region of Type I

Alpha cAMP-Dependent Protein Kinase'. *The Journal of Biological Chemistry* 272 (1): 379–88. https://doi.org/10.1074/jbc.272.1.379.

Poteet-Smith, Celeste E., John B. Shabb, Sharron H. Francis, and Jackie D. Corbin. 1997. 'Identification of Critical Determinants for Autoinhibition in the Pseudosubstrate Region of Type Iα cAMP-Dependent Protein Kinase'. *Journal of Biological Chemistry* 272 (1): 379–88. https://doi.org/10.1074/jbc.272.1.379.

Saegusa, Yu, and Kenjiro Yoshimura. 2015. 'cAMP Controls the Balance of the Propulsive Forces Generated by the Two Flagella of Chlamydomonas'. *Cytoskeleton (Hoboken, N.J.)* 72 (8): 412–21. https://doi.org/10.1002/cm.21235.

Salathe, Matthias. 2007. 'Regulation of Mammalian Ciliary Beating'. *Annual Review of Physiology* 69:401–22. https://doi.org/10.1146/annurev.physiol.69.040705.141253.

Santrich, C., L. Moore, T. Sherwin, P. Bastin, C. Brokaw, K. Gull, and J. H. LeBowitz. 1997. 'A Motility Function for the Paraflagellar Rod of Leishmania Parasites Revealed by PFR-2 Gene Knockouts'. *Molecular and Biochemical Parasitology* 90 (1): 95–109. https://doi.org/10.1016/s0166-6851(97)00149-7.

Schindelin, Johannes, Ignacio Arganda-Carreras, Erwin Frise, Verena Kaynig, Mark Longair, Tobias Pietzsch, Stephan Preibisch, et al. 2012. 'Fiji: An Open-Source Platform for Biological-Image Analysis'. *Nature Methods* 9 (7): 676–82. https://doi.org/10.1038/nmeth.2019.

Schmid, Andreas, Ge Bai, Nathalie Schmid, Manuela Zaccolo, Lawrence E. Ostrowski, Gregory E. Conner, Nevis Fregien, and Matthias Salathe. 2006. 'Real-Time Analysis of cAMP-Mediated Regulation of Ciliary Motility in Single Primary Human Airway Epithelial Cells'. *Journal of Cell Science* 119 (Pt 20): 4176–86. https://doi.org/10.1242/jcs.03181.

Schmidt, J. A., and R. Eckert. 1976. 'Calcium Couples Flagellar Reversal to Photostimulation in Chlamydomonas Reinhardtii'. *Nature* 262 (5570): 713–15. https://doi.org/10.1038/262713a0.

Scott, J D, E H Fischer, J G Demaille, and E G Krebs. 1985. 'Identification of an Inhibitory Region of the Heat-Stable Protein Inhibitor of the cAMP-Dependent Protein Kinase.' *Proceedings of the National Academy of Sciences of the United States of America* 82 (13): 4379–83. https://doi.org/10.1073/pnas.82.13.4379.

Shalaby, T., M. Liniger, and T. Seebeck. 2001. 'The Regulatory Subunit of a cGMP-Regulated Protein Kinase A of Trypanosoma Brucei'. *European Journal of Biochemistry* 268 (23): 6197–6206. https://doi.org/10.1046/j.0014-2956.2001.02564.x.

Siman-Tov, M. M., R. Aly, M. Shapira, and C. L. Jaffe. 1996. 'Cloning from Leishmania Major of a Developmentally Regulated Gene, c-Lpk2, for the Catalytic Subunit of the cAMP-Dependent Protein Kinase'. *Molecular and Biochemical Parasitology* 77 (2): 201–15. https://doi.org/10.1016/0166-6851(96)02601-1.

Sivadas, Priyanka, Jennifer M. Dienes, Martin St Maurice, William D. Meek, and Pinfen Yang. 2012. 'A Flagellar A-Kinase Anchoring Protein with Two Amphipathic Helices Forms a Structural Scaffold in the Radial Spoke Complex'. *The Journal of Cell Biology* 199 (4): 639–51. https://doi.org/10.1083/jcb.201111042.

Soriano-Úbeda, Cristina, Jon Romero-Aguirregomezcorta, Carmen Matás, Pablo E. Visconti, and Francisco A. García-Vázquez. 2019. 'Manipulation of Bicarbonate Concentration in Sperm Capacitation Media Improvesin Vitro Fertilisation Output in Porcine Species'. *Journal of Animal Science and Biotechnology* 10:19. https://doi.org/10.1186/s40104-019-0324-y.

Steichen, Jon M., Michael Kuchinskas, Malik M. Keshwani, Jie Yang, Joseph A. Adams, and Susan S. Taylor. 2012. 'Structural Basis for the Regulation of Protein Kinase A by

Activation Loop Phosphorylation'. *The Journal of Biological Chemistry* 287 (18): 14672–80. https://doi.org/10.1074/jbc.M111.335091.

Steinberg, R. A., R. D. Cauthron, M. M. Symcox, and H. Shuntoh. 1993. 'Autoactivation of Catalytic (C Alpha) Subunit of Cyclic AMP-Dependent Protein Kinase by Phosphorylation of Threonine 197'. *Molecular and Cellular Biology* 13 (4): 2332–41. https://doi.org/10.1128/mcb.13.4.2332-2341.1993.

Tash, J. S. 1989. 'Protein Phosphorylation: The Second Messenger Signal Transducer of Flagellar Motility'. *Cell Motility and the Cytoskeleton* 14 (3): 332–39. https://doi.org/10.1002/cm.970140303.

Torres-Quesada, Omar, Johanna E. Mayrhofer, and Eduard Stefan. 2017. 'The Many Faces of Compartmentalized PKA Signalosomes'. *Cellular Signalling* 37 (September):1–11. https://doi.org/10.1016/j.cellsig.2017.05.012.

Vacquier, Victor D., Arlet Loza-Huerta, Juan García-Rincón, Alberto Darszon, and Carmen Beltrán. 2014. 'Soluble Adenylyl Cyclase of Sea Urchin Spermatozoa'. *Biochimica Et Biophysica Acta* 1842 (12 Pt B): 2621–28. https://doi.org/10.1016/j.bbadis.2014.07.011.

Vasquez, Suly Saray Villa, John van Dam, and Gabrielle Wheway. 2021. 'An Updated SYSCILIA Gold Standard (SCGSv2) of Known Ciliary Genes, Revealing the Vast Progress That Has Been Made in the Cilia Research Field'. *Molecular Biology of the Cell* 32 (22): br13. https://doi.org/10.1091/mbc.E21-05-0226.

Walker, Benjamin J., Kenta Ishimoto, and Richard J. Wheeler. 2019. 'Automated Identification of Flagella from Videomicroscopy via the Medial Axis Transform'. *Scientific Reports* 9 (1): 5015. https://doi.org/10.1038/s41598-019-41459-9.

Walker, Benjamin J., and Richard J. Wheeler. 2019. 'High-Speed Multifocal Plane Fluorescence Microscopy for Three-Dimensional Visualisation of Beating Flagella'. *Journal of Cell Science* 132 (16): jcs231795. https://doi.org/10.1242/jcs.231795.

Walker, Benjamin J., Richard J. Wheeler, Kenta Ishimoto, and Eamonn A. Gaffney. 2019. 'Boundary Behaviours of Leishmania Mexicana: A Hydrodynamic Simulation Study'. *Journal of Theoretical Biology* 462 (February):311–20. https://doi.org/10.1016/j.jtbi.2018.11.016.

Wang, Ziyin, Tom Beneke, Eva Gluenz, and Richard John Wheeler. 2020. 'The Single Flagellum of Leishmania Has a Fixed Polarisation of Its Asymmetric Beat'. *Journal of Cell Science* 133 (20): jcs246637. https://doi.org/10.1242/jcs.246637.

Wheeler, Richard J., Eva Gluenz, and Keith Gull. 2011. 'The Cell Cycle of Leishmania: Morphogenetic Events and Their Implications for Parasite Biology'. *Molecular Microbiology* 79 (3): 647–62. https://doi.org/10.1111/j.1365-2958.2010.07479.x.

Wheeler, Richard John. 2017. 'Use of Chiral Cell Shape to Ensure Highly Directional Swimming in Trypanosomes'. *PLOS Computational Biology* 13 (1): e1005353. https://doi.org/10.1371/journal.pcbi.1005353.

Yang, Chun, and Pinfen Yang. 2006. 'The Flagellar Motility of Chlamydomonas Pf25 Mutant Lacking an AKAP-Binding Protein Is Overtly Sensitive to Medium Conditions'. *Molecular Biology of the Cell* 17 (1): 227–38. https://doi.org/10.1091/mbc.e05-07-0630.

Zhang, Jinyao, Jie Xu, Huijie Li, Yongyong Zhang, Zheng Ma, Andreas Bechthold, and Xiaoping Yu. 2022. 'Enhancement of Toyocamycin Production through Increasing Supply of Precursor GTP in Streptomyces Diastatochromogenes 1628'. *Journal of Basic Microbiology* 62 (6): 750–59. https://doi.org/10.1002/jobm.202100622.

A

LRR   AI   CNBD-A   CNBD-B

LdPKAR1-LdBPK.13.2.000160   502

LmjPKAR1-LmjF.13.0160   502

LmxPKAR1-LmxM.13.0160   503

TbPKAR1-Tb927.11.4610   499

TcPKAR1-TcCLB.506227.150   503

Coil

LdPKAR3-LdBPK.34.2.002680   647

LmjPKAR3-LmjF.34.2820   648

LmxPKAR3-LmxM.33.2820   646

TcPKAR3-TcCLB.510879.50   558

DD

HsPKAR-Iα –P10644   381

HsPKAR-IIα –P13861   404

B   AI

```
            **  *
LdPKAR1    RARRQTVRVEGI 212
LmjPKAR1   RARRQTVRVEGI 212
LmxPKAR1   RARRQTVRVEGI 212
TbPKAR1    RKKRRTTVRGEGI 211
TcPKAR1    RNRRRTVRSEGI 212
LdPKAR3    RPSRGAISDSSL 334
LmjPKAR3   RPSRGAISDSSL 334
LmxPKAR3   RPSRGAISDSSL 332
TcPKAR3    GHSRLGISARSI 245
HsPKAR-Iα  RRRRGAISAEVY 105
HsPKAR-IIα FNRRVSVCAETY 105
```

C   CNBD-A   CNBD-B

```
       *        * *               *       * *
AVGELELLYDTPAVAT 322   HIGELEFLNEHRTVADIVAVT 451
AVGELELLYDTPAVAT 322   HIGELEFLNKHRTVADIVAVT 451
AVGELELLYDTPAVAS 322   HFGELEFLNEHRTVADIVAVT 451
AVGELELMYQTPTVAT 321   HVGELEFLNNHANVADVVAKT 451
AVGELELMYDTPVVAT 322   HIGELEFLNNHRTVADVVATT 451
TFGDLGLMYEISNGAA 446   FAGQIEFLYHHVAVADVVAVS 452
TFGDLGLMYEISNGAA 447   FAGQIEFLYHHVAVADVVAVS 452
TFGDLGLMYEVSNGGA 445   FAGELEFLHHHVAAADVVAVS 452
VFGELGLLYETQSEVS 359   CFGDLEFLYHHETVADVVATS 451
SFGELALIYGTPRAAT 214   YFGEIALLMNRPRAATVVARG 343
SFGELALMYNTPRAAT 220   YFGELALVTNKPRAASAYAVG 355
```

D   HP

```
          **  *
AKYDYYQQVL 490
AKYDYYQQVL 490
AKYDYYQQVL 490
PNYEYYQSKL 489
PKYEYYQNVL 490
ETYATYQKRM 616
ETYATYQKRM 617
ETYAAYQERM 615
ASYDYYRRTM 528
--YNSFVSLS 380
--YEEQLVKM 392
```

Figure 1

Figure 2

A

(i) mNG::PKAC3

(ii) PKAR3::mNG

fluorescent intensity (a.u.)

(iii) mNG::PKAC3 *ΔPKAR3*

fractional distance

B

PKAR3::mNG(n=38)
mNG::PKAC3(n=54)
mNG::PKAC3*ΔPKAR3*(n=57)

fluorescent intensity (a.u.)

F tip   F mid   F prox   FP   N   CB tip

live promastigotes

merged   mNG

C   mNG::PKAC1 *ΔPKAR1*

D   mNG::PKAC2 *ΔPKAR1*

E   mNG::PKAC3 *ΔPKAR1*

F   mNG::PKAC3 *ΔPKAR3*

Figure 3

Figure 4

Figure 5

Figure 6

mutant	n datapoints (n cells)	total time obs (s)	mean Hz (bps)	mean amp (degrees)	waves / flagellum (w/f)	F-length (µm) (n cells)
ΔLmxPKAR1	24 (17)	32.6	28	37	0.84	15.5 (57)
ΔLmxPKAR3	15 (10)	20.4	33	39	0.91	15 (88)
ΔLmxPKAC1	25 (17)	34	4***	26***	1.01	14.6 (93)
ΔLmxPKAC2	13 (7)	17.7	32	34*	1.08*	13.9 (97)
ΔLmxPKAC3	25 (15)	34	22***	41	0.79	13*** (143)
parental	35 (21)	48	27	41	0.85	15.2 (75)

Figure 7