Insights on the effect of mega-carcass abundance on the population dynamics of a

facultative scavenger predator and its prey

Mellina Sidous[1,2], Sarah Cubaynes[2], Olivier Gimenez[2], Nolwenn Drouet-Hoguet[3,4,5], Stéphane

Dray[3], Loïc Bollache[1], Daphine Madhlamoto[6], Nobesuthu Adelaide Ngwenya[6], Hervé

Fritz[3,4,7,8,*], Marion Valeix[2,3,4,*]

[1] Chrono-environnement UMR6249, CNRS, Université Bourgogne Franche-Comté, F-25000, Besançon, France

[2] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

[3] CNRS, Université de Lyon, Université de Lyon 1, VetAgroSup, LBBE, UMR 5558, F-69622, Villeurbanne,

France

[4] Long-Term Socio-Ecological Research Site (LTSER) France, Zone Atelier "Hwange", Hwange National Park,

Bag 62, Dete, Zimbabwe

[5] OFB, 5 All. de Bethléem, 38610 Gières, France

[6] Zimbabwe Parks and Wildlife Management Authority, Main Camp Research, Hwange National Park, Zimbabwe

[7] REHABS International Research Laboratory, CNRS – Université Lyon – Nelson Mandela University, George

Campus, Madiba Drive, 6531 George, South Africa

[8] Sustainability Research Unit, Nelson Mandela University, George Campus, Madiba Drive, 6531 George, South

Africa

[*] shared last-authorship

ORCIDId:

Mellina Sidous : **0009-0009-3003-5080**
Sarah Cubaynes : **0000-0002-3935-9825**
Olivier Gimenez : **0000-0001-7001-5142**
Nolwenn Drouet Hoguet : **0000-0002-2610-6917**
Stéphane Dray : **0000-0003-0153-1105**
Loic Bollache : **0000-0003-0316-6746**
Daphine Madhlamoto **: 0000-0002-5856-0153**
Nobesuthu Adelaide Ngwenya : **0000-0002-0152-1364**
Marion Valeix : **0000-0002-9732-1737**
Hervé Fritz : **0000-0002-7106-3661**
32

Abstract

The interplay between facultative scavenging and predation has gained interest in the last decade. The prevalence of scavenging induced by the availability of large carcasses may modify predator density or behaviour, potentially affecting prey. In contrast to behavioural mechanisms through which scavenging affects predation, the demographic effects of facultative scavenging on predator and prey populations remain poorly studied. We used the semi-natural experimental opportunity in Hwange National Park, Zimbabwe, where contrasted management measures (culling and artificial supply of water) have led to fluctuations in elephant carrion abundance, to identify the consequences of facultative scavenging on the population dynamics of a large mammalian carnivore, the spotted hyaena (*Crocuta crocuta*), and its prey. Using a 50-year dataset and Multivariate Autoregressive State Space models, we estimated hyaena and prey densities over four periods contrasted in elephant carrion availability due to management practices. Models that allow hyaena and their prey populations' growth rate to vary depending on these four periods contributed significantly to explain variations in their density, which is consistent with an effect of management measures on the population dynamics of hyaena and its prey. Although our results support a predominant role of bottom-up mechanisms, whereby hyaena density is driven by herbivore density, itself driven by resources availability, some subtle patterns of densities could be interpreted as consequences of changes in predation pressure following changes in scavenging opportunities. We discuss why signals of prey and predator population dynamics decoupling are less likely to be observed in systems with a high diversity of prey, such as African savannas, and why inputs of mega-carcasses as pulsed resources hardly impacted top-down relationships in the long run. This study represents a first investigation of the long-term effects of carrion pulses, whose frequency may increase with climate changes, on the classical predator-prey coupling for large mammals.

Introduction

Predation is the most studied ecological process amongst those linking predator and prey populations (Abrams 2000, Guiden et al. 2019). Although overlooked, scavenging (i.e., when animals feed on dead animals and carrion they did not kill) recently gained increased recognition (Wilson and Wolkovich 2011, Moleón and Sánchez-Zapata 2015, Luna et al. 2021). So far, these two processes have been little investigated together. However, because carnivores are often facultative scavengers (Fallows et al. 2013, Pereira et al. 2014, Gomo et al. 2017), and scavenging is energetically less costly than hunting, carnivores are distracted from their live prey in the presence of carrion (DeVault et al. 2003). Consequently, carrion and live prey are linked by indirect interactions. On the one hand, live prey can be subject to a higher predation pressure, or hyper-predation, if carrion pulses lead to a higher abundance of carnivores (through demography or local immigration) but carrions are not available (because of interference or decay of the carrion; Courchamp et al. 2000, Cortés-Avizanda et al. 2009a , Cortés-Avizanda et al. 2009b, Moleón et al. 2014). On the other hand, live prey can be subject to a reduced predation pressure, or hypo-predation, if carnivore abundance remains stable but the rate of consumption of live prey is reduced and compensated by scavenging (Bate and Hilker 2012, Fallows et al. 2013, Moleón et al. 2014, Mellard et al. 2021). These indirect interactions linking carrion and live prey, through predators, have been studied theoretically (Andrén et al. 2011, Mellard et al. 2021), experimentally (Cortés-Avizanda et al. 2009a), and empirically (Cortés-Avizanda et al. 2009b, Fallows et al. 2013). These previous works globally conclude about an impact of scavenging on predation processes, either through an alteration of prey and predator space use (Cortés-Avizanda et al. 2009b), or through a direct modification of predation rate (Cortés-Avizanda et al. 2009a, Andrén et al. 2011, Fallows et al. 2013, Mellard et al. 2021). However, there is a dearth of long-term, demographic studies to assess whether regular pulses of carrion interfere with the population dynamics of predators and their prey (Moleón et al.

2014). Still, in some systems, population dynamics of a meso-predator was affected by

facultative scavenging with associated consequences on prey populations (Roth 2003), and

regular anthropogenic food subsidies seem able to alter the classical predator-prey coupling of

population dynamics (Rodewald et al. 2011). The megafauna is unparalleled in the extent of

carrion they provide after death and mega-carcasses (from whales, elephants, rhinoceros) can

be considered as important pulsed resources for scavengers.

In this study, we explored the effect of elephant (*Loxodonta Africana*) carrion

availability on the numerical response of a large opportunist scavenging predator, the spotted

hyaena (*Crocuta crocuta*; hyaena hereafter; Figure 1 ; Kruuk 1972) and its prey. Their relative

abundances were monitored over almost 50 years in Hwange National Park, Zimbabwe. As a

consequence of management practices, this ecosystem witnessed large variations in the

abundance of the elephant population over the years, and four periods characterized by different

elephant carcass availability emerged.

From 1960 to 1986, massive culls were undertaken in the area in response to an elephant

abundance perceived as "too high" regarding its effects on biodiversity (Child 2004, Slotow et

al. 2008). During this time, at least 18,000 elephants were killed (Cumming 1981, ZPWMA

1998, Child 2004, Slotow et al. 2008). This period corresponds to our first study period (1972-

1986) when hundreds to a few thousand carcasses were available annually and clearly provided

an important source of food for scavengers. In contrast, in the second period (1987-1992),

elephant carcasses were rare due to a steady and continuous increase of the population triggered

by surface water management (pumping of underground water) and the cessation of culling

(Chamaillé-Jammes et al. 2008). This lasted until the third period (1993-2008) when elephant

carcasses were present in large numbers in dry years (Dudley et al. 2001), after the elephant

population reached density-dependence, fluctuating around 30,000 individuals (Chamaillé-

Jammes et al. 2008). Finally, during the fourth period (2009-2020), natural mortality (density-

dependence driven) still existed but was likely to be much lower than during period 3 due to

the creation of additional waterholes (Appendix 1), allowing the elephant population to pass

the equilibrium reached previously and start growing again.

The existence of these four periods contrasted in terms of elephant carcass availability,

together with the long-term monitoring of hyaena and prey populations covering these different

periods, provide a unique semi-natural experimental opportunity to assess the impacts of

scavenging on predation at the level of population dynamics. Here, we hypothesised that if

scavenging modified predation with further demographic consequences, we would detect that

the Hwange hyaena population is driven by its live prey in periods of low elephant carcass

abundance (periods 2 and 4), but less dependent of its live prey in periods of high elephant

carcass abundance (periods 1 and 3).

Figure 1: Hyaena scavenging from an elephant carcass (credit: S. Périquet).

Methods

Study area

Hwange National Park covers ~15,000 km^2 of dystrophic semi-arid (~600 mm annual rainfall)

savanna in western Zimbabwe (19°00' S, 26°30' E). This protected area is not fenced, which

allows wildlife to move freely. Surface water is primarily found in natural depressions that hold

rainwater. However, most dry up as the dry season progresses, from May to September. To

guarantee sufficient drinking water for animals in the dry season, water has been managed since

1935 through the pumping of underground water into waterholes distributed mostly in the

northern sector of the park. The number of pumped waterholes has varied over the past 50 years,

from about 20 to more than 80 (Appendix 1). In this study, we focus on a 3,000 km^2 study area

in the north-eastern sector of the park (Figure 2), the most touristic area of the park, with most

pumped waterholes.

Data

We used a long-term data set resulting from the regular monitoring of waterholes from 1972 to

2020 by the Wildlife Environment Zimbabwe (WEZ). This monitoring consists of annual 24h

counts during full moon at the end of the dry season. At this time of the year, it is mostly the

pumped waterholes that provide drinking water to animals, but in years with a good rainy season

a few natural pans can hold water too. Because this monitoring is largely influenced by annual

rainfall, counts are characterized by high inter-annual variance and cannot be considered as

total counts. However, abundance trends detected with these counts are the same as those

obtained with other more traditional methods (aerial census; Valeix et al. 2008), and thus we

consider that long-term trends inferred from counts at waterholes reliably reflect actual patterns

of abundance changes in populations (as in Valeix et al. 2008). Additionally, to control for the

proximity of some waterholes and the differential surfaces considered as their attraction zones,

monitored waterholes were grouped in new spatial units created using a centroidal Voronoi

tessellation based on pumped waterholes (Okabe et al. 2009) (Figure 2). Indeed, if we consider

a pumped waterhole, its attraction is reduced when new pumped waterholes are added in its

vicinity or in years when neighbouring natural pans still hold water. In other words, we expect

less animals to be observed at a focused pumped waterhole since animals should split between

this waterhole and the neighbouring pumped or natural waterholes. This could lead to conclude

that species abundance has decreased when in reality the patterns observed come from this

change in attraction rather than an actual decrease in regional abundance of species. By

grouping neighbouring waterholes into new spatial units, we expect to reduce this bias arising

from the fluctuation in the number of waterholes over years. For each year, we summed the

animals counted at the different waterholes of each spatial unit and divided it by the number of

waterholes surveyed, and then by the surface of the spatial unit to assess densities. For political

(civil war) and economic reasons, counts were not performed in 1974, 1983, and from 1976 to

1981. For the analyses, we selected the spatial units that were monitored for at least 50% of the

years composing each period, except for period 1 (the period impacted by the civil war) for

which we kept units with at least 25% of the years monitored.

One spatial unit was excluded for the analyses because it is known for the presence of a hyaena

den close to a waterhole and thus presented singularly high hyaena counts. Counts were log-

transformed and we dealt with 0 values by adding 0.0001 to the whole dataset, which

corresponds to one order of magnitude lower than the smallest value of the data.

We used data on hyaena and its prey. For prey, we pooled all (males, females, adults,

sub-adults and juveniles) preferred and common prey species of hyaena in Hwange National

Park (Périquet et al. 2015) ('prey' hereafter). This includes plains zebra (*Equus quagga*), greater

kudu (*Tragelaphus strepsiceros*), sable (*Hippotragus niger),* roan *(Hippotragus equinus*), blue

wildebeest (*Connochaetes taurinus),* waterbuck (*Kobus ellipsiprymnus*), impala (*Aepyceros*

melampus), and warthog (*Phacochoerus africanus*).

<u>Figure 2:</u> Hwange National Park and location of the study area. Black dots are the waterholes of the
park. Voronoi polygons used for the analysis are coloured in blue.

Statistical analyses

We fitted a Multivariate Autoregressive State-Space Model (Holmes et al. 2012) using R

package MARSS (Holmes et al. 2021, R Core Team 2021) to estimate prey and hyaena densities

from 1972 to 2020 from counts data at the waterholes, while accounting for observation and

process errors. We considered each spatial unit to be a replicate of a common dynamic, and

therefore estimated one density for prey and one density for hyaena for the entire study area.

MARSS separates the state process, capturing variations in species density, from the

observation process explaining variations in the number of individuals observed at the

waterholes. More precisely, the model uses an autoregressive process, which means that the

state x of the system at time t (x_t) depends on its value at time t-1 (x_{t-1}). In our case, x is the

densities of a guild (either prey or hyaena) and we have:

$x_t = x_{t-1} + u_t + w_t; \ w_t \sim Normal(0, Q)$ (1)

Where x_t, x_{t-1}, and Q are 1x1 matrices. x_t and x_{t-1} represent the abundance estimated at time t

and t-1 respectively for the focus guild. The parameter w_t is the error term of the state process

and represents environmental stochasticity. This environmental stochasticity is modelled as a

normal distribution with mean 0 and a variance Q that takes the same values for the whole time

series of each guild. The parameter u_t is the growth rate of the population at time t. This is the

parameter that was constrained differently in the three models we tested. For each time step,

the growth rate has a unique value and thus u_t is a 1x1 matrix.

Equation (1) models the state process. Here, it represents the changes in densities of each guild

over time. We have only access to a noisy signal of these dynamics through their observation

at the different waterholes of our study area. The densities of each guild can thus be considered

as "hidden states" and state-space models aim at retrieving access to these states through their

signals. This is done by incorporating a second equation that models the observation process:

$\mathbf{y_t} = \mathbf{Z_t}x_t + \mathbf{D}d_t + a + \mathbf{v_t}; \mathbf{v_t} \sim \text{MVN}(0, \mathbf{R})$ (2)

For n waterholes, $\mathbf{y_t}$ has a dimension nx1, where n is the number of waterholes. $\mathbf{Z}$ is a nx1

matrix that links the observations $\mathbf{y_t}$ to the state to be estimated (x_t). Since we only estimated

the density of each species separately and not in the same model, Z matrix is full of 1s. The bias

parameter, a, is set to be equal between sites. The parameter v_t is the error term of the

observation process equation, and comes from a multivariate normal distribution that share a

common variance over time and site. The variance is the parameter $\mathbf{R}$ a variance covariance

matrix of dimension *nxn*. Finally, d_t is a vector with annual rainfall value at year t, and $\mathbf{D}$ further

links the effect of d_t to the $\mathbf{y_t}$ that are inputs of the model. In this observation model, we included

annual rainfall as a covariate to account for bias on counts at waterholes (Appendix 2). During

dry years, animals need to visit waterholes more often than during wet years when more

ephemeral natural ponds can be found in the park. Thus, when annual rainfall is not incorporated

into the model, estimated densities are overestimated in dry years, and underestimated in wet

year (Appendix 2).

We fitted and compared (using AICc ;Burnham et al. 2011) three different models: (i) a *constant*

model with a constant growth rate, (ii) a *time-varying* model estimating a growth rate per year,

and (iii) a *period* model estimating one growth rate per period. The period model aims at testing

a signal in densities caused by the fluctuations of elephant carrion availability. We used an

analysis of deviance (ANODEV) to test the significance and assess the proportion of the

deviance explained by a period-dependent growth rate for hyaena and its prey (Grosbois et al.

2008).

<u>Results</u>

Observed counts data and estimated population densities for prey and hyaena are provided in

Figure 3. The period model best described the variations in density of prey (AICc=1862.05,

1890.89 and 1850.95 for the constant, time-varying and period models respectively) and hyaena

(AICc=3039.18, 3089.02 and 3029.70 for the constant, time-varying and period models

respectively). ANODEV tests were significant for both prey (F = 0.009; r^2 = 0.22; df= 3) and

hyaena (F= 0.002; r^2 = 0.28; df = 3). Hyaena and prey densities showed similar trends across

the 4 study periods: the two populations increased during period 1, stabilized during period 2,

decreased during period 3, and increased again during period 4 (Figure 3; period models'

growth rate estimates in Table 1). Differences between growth rate of hyaena and prey are

similar in both periods when elephant carcasses were less abundant (0.069 in the second and

0.063 in the fourth period; Table 1). During periods 1 and 3 characterized by increased

abundance of elephant carrion in the landscape, this difference was increased or decreased

(0.123 during the first and 0.004 during the third period). However, confidence intervals of the

two guilds partially overlap and are large.

Figure 3: **Top:** Observed population densities of prey (**A**) and hyaena (**B**). Dots correspond to the
densities observed at the spatial units (Voronoi polygons), each colour representing a spatial unit (95%
of the data kept for visual clarity). Fluctuating thin lines represent the median of the observed densities
each year. Bold line and ribbon represent the smooth of the median and its 95% confidence interval.
Bottom: Estimated population densities of prey (**C**) and hyaena (**D**). Model estimates are given for the
constant, time-varying, and period models. The ribbon represents the 95% of these estimates, but is very
close to the estimates for the time-varying and period models. In all panels, the dotted vertical lines
delimitate the four study periods; elephant pictogram indicates periods of high elephant carcass
availability.

Table 1: Estimation of growth rate (with 95% confidence interval) from the period model for prey and
hyaena populations.

	Prey	**Hyaena**
Period 1 (1972-1986)	0.052 (0.017; 0.092)	0.175 (0.079; 0.269)
Period 2 (1987-1992)	-0.010 (-0.073; 0.049)	0.059 (-0.094; 0.204)
Period 3 (1993-2008)	-0.042 (-0.063; -0.021)	-0.058 (-0.107; -0.008)
Period 4 (2009-2020)	0.025 (-0.009; 0.056)	0.090 (0.013; 0.168)

Discussion

A megafauna's carcass, which represents several tons of resource, undoubtedly

influences the foraging and ranging behaviour of facultative scavengers, which can use such a

carrion for long periods. In marine ecosystems, sharks can use whale carrion for several days

(Fallows et al. 2013). In African savannas, hyaenas definitely modify their foraging and ranging

habits in the presence of an elephant carcass, which they can use for up to two weeks (Cozzi et

al. 2015, Périquet et al. 2021). During this time, hyaenas are diverted from their prey. In periods

of high elephant carcass availability, these carrion-induced behavioural changes should affect

interactions between carnivores and their prey. Surprisingly, little is known about the

cumulative long-term effects of these changes and the ultimate implication for population

dynamics of predators able of facultative scavenging and their prey. Using almost 50 years of

data, and a semi-natural experiment involving carrion of the largest terrestrial mammal, we do

not detect patterns supporting effects of high carcass availability on the population dynamics

of hyaena and its prey (i.e. we do not detect that hyaena population is less driven by its live

prey in periods of high elephant carcass abundance, as we initially hypothesized). The main

result is that hyaena and its prey showed the same population trends over all periods, with

hyaena population dynamics following that of its prey independently of elephant carcass

availability, as in classical predator-prey fluctuations (Table 1 and Figure 3 ; Krebs et al. 2001,

Hebblewhite et al. 2002, Wang et al. 2009).

The periods contributed significantly to explain variations in prey and hyaena density

(~ 30% of the deviance). Thus, management measures (elephant culling and water supply) and

the resulting elephant population dynamics used to define the periods partly explained

population trends. Prey abundance increased in the first and last periods, when artificial

waterholes (determining the amount of resource available for herbivores) were added in the

landscape, and decreased in periods 2 and 3 coinciding with an increase in the elephant

population, but also to a decade of low dry season rainfall (Valeix et al. 2008). These patterns

are coherent with the hypothesis that bottom-up mechanisms drive the patterns of densities we

observe. Note that we did not explore other factors underlying the population trends of hyaena

and its prey, such as the role of other large carnivores or climatic conditions, which was beyond

the scope of this study. For example, the African lion, *Panthera leo,* which population has

undergone large fluctuations, has the potential to influence herbivore populations and hyaena

foraging behaviour (Grange et al. 2015, Périquet et al. 2015, Loveridge et al. 2016). Moreover,

even though the hyaena population seems to decline 2-3 years after its prey population (Figure

3), we could not test for lags between their densities because periods were too short.

287   The main result of this study does not allow to conclude that a mechanism of hyper- or

hypo-predation interfered with the population dynamics of predators and their prey on the long-

term. Nonetheless some depicted patterns, although subtle, could be interpreted in the light of

a role of scavenging. In period 1, both prey and hyaena population abundances increased, but

this increase was stronger for hyaena (Table 1), which could be explained by hyaenas benefiting

from both the increase in prey populations and the additional energy-rich resources provided

by elephant culling. In period 2, growth rate confidence intervals are large and encompass 0 for

both prey and hyaenas (Table 1), showing a general stabilization for the two populations.

However, estimated growth rates suggest that prey abundance tended to decrease while hyaena

abundance continued increasing in period 2, which is consistent with a scenario whereby the

hyaena population, then abundant, compensated for the loss of the energy-rich resources

provided by elephant carrion in period 1 by a renewed predation pressure in period 2. In period

3, hyaena population abundance did not decrease at a rate lower than its prey population, but

maintains a higher density than at the start of period 1 despite a strong decrease of prey density,

which could be explained by the presence of the additional food source provided by elephant

carcasses. Finally, during period 4, both hyaena and prey densities increased, but the rate of

increase is slightly lower for prey than hyaena, possibly indicating again a renewed predation

of hyaena on prey populations. Differences between estimated growth rates of each guild are

also potentially suggesting a change in the link between hyaena and prey densities caused by

elephant carrion abundance : the differences are similar in periods of lower abundance of

elephant carrion (about 0.06), but higher and lower in periods of high elephant carrion

abundance. Testing these hypotheses would require further investigations in the future, by

comparing hyaena diet and hunting behaviour in drought years (when elephant mortality will

be high) and in years with good annual rainfall (when elephant mortality will be low).

Overall, these patterns are indicative of a numerical response of hyaena to the increased

elephant carcass availability, through a rapid increase in density in period 1 and the maintenance

of densities higher than at the start of our study period in period 3. If they are not hunted,

mammalian predators' density are generally associated with prey biomass (Fuller and Sievert

2001, Hayward et al. 2007) and this was verified for hyaena in Kruger National Park, South

Africa (Ferreira and Funston 2016). Thus, a large increase of exploitable resources though

inputs of additional food sources such as mega-carcasses should allow the survival and

reproduction of more individuals; in other words, results in an increased carrying capacity (Oro

et al. 2013). Contrary to our study, in Manitoba (Canada), the increased arctic fox (*Vulpes*

lagopus) abundance after increased scavenging on carcasses of polar bear (*Usrus maritimus*)

kills was accompanied by a signal in the population dynamics of its usual main prey, the

collared lemming (*Dicrostonyx richardsoni*) (Roth 2003). The classical cyclic dynamics of

lemming was delayed, probably due to an increased predation pressure arising from the higher

predator abundance, which is akin to a phenomenon of hyper-predation (Courchamp et al. 2000,

Moleón et al. 2014). However, arctic and African savannas ecosystems differ fundamentally.

While arctic predator-prey communities are sometimes referred as the simplest in the world and

present few predators that all rely largely upon one main prey, lemmings (*Dycrostony spp.,*

Lemmus spp. ; Gilg et al. 2006, Reiter and Andersen 2011), African savannas present a high

diversity of prey and predators in which classical prey and predator couplings are not observed

(Sinclair et al. 2003). In arctic ecosystems, arctic fox only shifts to other prey when lemmings

become very scarce (Gilg et al. 2006), which makes direct and observable repercussion of

changes in their populations dynamics on that of lemmings more likely. In contrast, hyaena is

known to be a generalist predator with a very plastic feeding behaviour, selecting food sources

depending on their abundance (Kruuk 1972, Périquet et al. 2015). Consequently, if hyaena

predation starts to reduce the abundance of one prey, it is likely that the predator shifts easily

to another type of prey, allowing the other species to recover.

Furthermore, carrion can be considered as pulsed resources whose characteristics can

affect the consumer dynamics (Yang et al. 2008). Interestingly, the spatio-temporal

predictability of elephant carrion pulses differed between periods 1 and 3. During period 3, they

were predictable and limited in time and space, as elephants die naturally near waterholes at the

end of the dry season (Conybeare and Haynes 1984). From a hyaena perspective, a cursorial

hunter, the end of the dry season is the time of the year when prey are the weakest and hence

easier to catch (Cozzi et al. 2015). Consequently, mortality is likely to be compensatory,

(meaning that individuals killed by hyaena were weak and likely to have died from other causes,

such that population's total mortality remained unchanged) and resource pulses at the end of

the dry season may not matter that much. In contrast, during period 1, carcasses were highly

aggregated (because entire elephant families were removed through culling) and unpredictable

in space and time (Cumming 1981). They were more likely to influence hyaena fitness and

hence demography with later consequences on prey demography. However, the unpredictability

and short duration of carrion availability through these measures may explain the lack of strong

and observable impacts on population dynamics on the long term as we observed here.

The frequency and intensity of pulses of carrion are likely to be altered under global

changes and human activities. In large terrestrial mammals, peaks of herbivore carcasses are

mostly associated with extreme weather periods, such as droughts or severe winters, which

should become more frequent in the future (Intergovernmental Panel on Climate Change

(IPCC) 2023). Moreover, anthropogenic carcasses become increasingly present (Moreno-Opo

and Margalida 2019) through hunting activities (Mateo-Tomás et al. 2015), poaching

(Underwood et al. 2013) or increasing number of domestic herbivores (Arrondo et al. 2019).

Understanding how the characteristics of pulses of large herbivore carcasses may affect the

coupling between large carnivores and their live prey populations is key if we want to predict

the future of these populations.

Acknowledgment: We are indebted to the Wildlife Environment Zimbabwe and thank warmly

the Zimbabwe Parks and Wildlife Management Authority for their dedicated effort in

monitoring wildlife in Zimbabwe in general and in Hwange National Park in particular and for

providing this invaluable long-term dataset that spans over 50 years. We also thank Arnold

Tshipa, Makalolo Camp and Wilderness for providing their rainfall data for this study. This

manuscript benefited from comments from Eli Strauss and one anonymous reviewer, and a

preprint version of this article has been peer-reviewed and recommended by PCIEcology.

References
Abrams, P. A. 2000. The evolution of predator-prey interactions: Theory and evidence. - Annu. Rev.
Ecol. Syst. 31: 79–105.

Andrén, H. et al. 2011. Modelling the combined effect of an obligate predator and a facultative predator
on a common prey: lynx *Lynx lynx* and wolverine *Gulo gulo* predation on reindeer *Rangifer*
tarandus. - Wildl. Biol. 17: 33–43.

Arrondo, E. et al. 2019. Rewilding traditional grazing areas affects scavenger assemblages and carcass
consumption patterns. - Basic Appl. Ecol. 41: 56–66.

Bate, A. M. and Hilker, F. M. 2012. Rabbits protecting birds: Hypopredation and limitations of
hyperpredation. - J. Theor. Biol. 297: 103–115.

Burnham, K. P. et al. 2011. AIC model selection and multimodel inference in behavioral ecology: some
background, observations, and comparisons. - Behav. Ecol. Sociobiol. 65: 23–35.

Chamaillé-Jammes, S. et al. 2008. Resource variability, aggregation and direct density dependence in
an open context: the local regulation of an African elephant population. - J. Anim. Ecol. 77:
135–144.

Child, G. 2004. Elephant culling in Zimbabwe. - ZimConservation Opin.

Conybeare, A. and Haynes, G. 1984. Observations on elephant mortality and bones in water holes. -
Quat. Res. 22: 189–200.

Cortés-Avizanda, A. et al. 2009a. Carcasses increase the probability of predation of ground-nesting
birds: A caveat regarding the conservation value of vulture restaurants. - Anim. Conserv. 12:
85–88.

Cortés-Avizanda, A. et al.. 2009b. Effects of carrion resources on herbivore spatial distribution are
mediated by facultative scavengers. - Basic Appl. Ecol. 10: 265–272.

Courchamp, F. et al. 2000. Rabbits killing birds: modelling the hyperpredation process. - J. Anim. Ecol.
69: 154–164.

Cozzi, G. et al.. 2015. Effects of Trophy Hunting Leftovers on the Ranging Behaviour of Large
Carnivores: A Case Study on Spotted Hyenas. - PLOS ONE 10: e0121471.

Cumming, D. 1981. The management of elephant and other large mammals in Zimbabwe. - In: Problems
in management of locally abundant wild mammals. Academic Press, pp. 91–118.

DeVault, T. L. et al. 2003. Scavenging by vertebrates: behavioral, ecological, and evolutionary
perspectives on an important energy transfer pathway in terrestrial ecosystems. - Oikos 102:
225–234.

Dudley, J. P. et al. 2001. Drought mortality of bush elephants in Hwange National Park, Zimbabwe. -
Afr. J. Ecol. 39: 187–194.

Fallows, C. et al. 2013. White Sharks (*Carcharodon carcharias*) Scavenging on Whales and Its Potential
Role in Further Shaping the Ecology of an Apex Predator - PLoS ONE 8: e60797.

Ferreira, S. M. and Funston, P. J. 2016. Population estimates of spotted hyaenas in the Kruger National
Park, South Africa. - Afr. J. Wildl. Res. 46: 61–70.

Fuller, T. and Sievert, P. 2001. Carnivore demography and the consequences of changes in prey avail-
ability. - In: Carnivore Conservation. Cambridge University Press, pp. 163–178.
412

Gilg, O. et al. 2006. Functional and numerical responses of four lemming predators in high arctic
Greenland. - Oikos 113: 193–216.

Gomo, G. et al. 2017. Scavenging on a pulsed resource: quality matters for corvids but density for
mammals. - BMC Ecol. 17: 22.

Grange, S. et al. 2015. Demography of plains zebras (*Equus quagga*) under heavy predation. - Popul.
Ecol. 57: 201–214.

Grosbois, V. et al. 2008. Assessing the impact of climate variation on survival in vertebrate populations.
- Biol. Rev. 83: 357–399.

Guiden, P. W. et al. 2019. Predator-Prey Interactions in the Anthropocene: Reconciling Multiple Aspects
of Novelty. - Trends Ecol. Evol. 34: 616–627.

Hayward, M. W. et al. 2007. Carrying capacity of large African predators: Predictions and tests. - Biol.
Conserv. 139: 219–229.

Hebblewhite, M. et al. 2002. Elk population dynamics in areas with and without predation by
recolonizing wolves in Banff National Park, Alberta. - Can. J. Zool. 80: 789–799.

Holmes, E., E. et al. 2012. MARSS: Multivariate Autoregressive State-space Models for Analyzing
Time-series Data. - R J. 4: 11.

Holmes, E., E. et al. 2021. MARSS: Multivariate Autoregressive State-Space Modeling.

Intergovernmental Panel on Climate Change (IPCC) 2023. Climate Change 2021 – The Physical Science
Basis: Working Group I Contribution to the Sixth Assessment Report of the Intergovernmental
Panel on Climate Change. - Cambridge University Press.

Krebs, C. J. et al. 2001. What Drives the 10-year Cycle of Snowshoe Hares? - BioScience 51: 25.

Kruuk, H. 1972. The spotted hyena, a study of predation and social behavior. - Univ Chic. Press: 335.

Loveridge, A. J. et al. 2016. Conservation of large predator populations: Demographic and spatial
responses of African lions to the intensity of trophy hunting. - Biol. Conserv. 204: 247–254.

Luna, Á. et al. 2021. Predation and Scavenging in the City: A Review of Spatio-Temporal Trends in
Research. - Diversity 13: 46.

Mateo-Tomás, P. et al. 2015. From regional to global patterns in vertebrate scavenger communities
subsidized by big game hunting. - Divers. Distrib. 21: 913–924.

Mellard, J. P. et al. 2021. Effect of scavenging on predation in a food web. - Ecol. Evol. 11: 6742–6765.

Moleón, M. and Sánchez-Zapata, J. A. 2015. The Living Dead: Time to Integrate Scavenging into
Ecological Teaching. - BioScience 65: 1003–1010.

Moleón, M. et al. 2014. Inter-specific interactions linking predation and scavenging in terrestrial
vertebrate assemblages. - Biol. Rev. Camb. Philos. Soc. 89: 1042–1054.

Moreno-Opo, R. and Margalida, A. 2019. Human-Mediated Carrion: Effects on Ecological Processes. -
In: Olea, P. P. et al. (eds), Carrion Ecology and Management. Wildlife Research Monographs.
Springer International Publishing, pp. 183–211.

Okabe, A. et al. 2009. Spatial Tessellations: Concepts and Applications of Voronoi Diagrams. - John
Wiley & Sons.

Oro, D. et al. 2013. Ecological and evolutionary implications of food subsidies from humans. - Ecol.
Lett. 16: 1501–1514.

Pereira, L. M. et al. 2014. Facultative predation and scavenging by mammalian carnivores: seasonal,
regional and intra-guild comparisons: Predation vs. scavenging in carnivores. - Mammal Rev.
44: 44–55.

Périquet, S. et al. 2015. Spotted hyaenas switch their foraging strategy as a response to changes in
intraguild interactions with lions. - J. Zool. 297: 245–254.

Périquet, S. et al. 2021. Dynamic interactions between apex predators reveal contrasting seasonal
attraction patterns. - Oecologia 195: 51–63.

R Core Team 2021. R: A language and environment for statistical computing. - R Foundation for
Statistical Computing, Vienna, Austria.

Reiter, M. E. and Andersen, D. E. 2011. Arctic Foxes, Lemmings, and Canada Goose Nest Survival at
Cape Churchill, Manitoba. - Wilson J. Ornithol. 123: 266–276.

Rodewald, A. D. et al. 2011. Anthropogenic resource subsidies decouple predator—prey relationships.
- Ecol. Appl. 21: 936–943.

Roth, J. 2003. Variability in marine resources affects arctic fox population dynamics. - J. Anim. Ecol.
72: 668–676.

Sinclair, A. R. E. et al. 2003. Patterns of predation in a diverse predator–prey system. - Nature 425: 288–
290.

Slotow, R. et al. 2008. Lethal Management of Elephants. - In: Scholes, R. J. and Mennell, K. G. (eds),
Elephant Management: A Scientific Assessment for South Africa. Wits University Press.

Underwood, F. M. et al. 2013. Dissecting the Illegal Ivory Trade: An Analysis of Ivory Seizures Data.
- PLoS ONE 8: e76539.

Valeix, M. et al. 2008. Fluctuations in abundance of large herbivore populations: insights into the
influence of dry season rainfall and elephant numbers from long-term data. - Anim. Conserv.
11: 391–400.

Wang, H. et al. 2009. The roles of predator maturation delay and functional response in determining the
periodicity of predator-prey cycles. - Math. Biosci. 221: 1–10.

Wilson, E. E. and Wolkovich, E. M. 2011. Scavenging: how carnivores and carrion structure
communities. - Trends Ecol. Evol. 26: 129–135.

Yang, L. H. et al. 2008. What can we learn from resource pulses? - Ecology 89: 621–634.

ZPWMA 1998. Hwange National Park Management Plan 1999-2004. Report, Harare, Zimbabwe.

484