Prominin 1 and Tweety Homology 1 both induce extracellular vesicle formation

Tristan A. Bell[1,2]*, Bridget E. Luce[1], Pusparanee Hakim[1], Virly Y. Ananda[1], Hiba Dardari[1], Tran H. Nguyen[1],

Arezu Monshizadeh[1], and Luke H. Chao[1,2]*

[1]Department of Molecular Biology, Massachusetts General Hospital, Boston, MA, 02114

[2]Department of Genetics, Blavatnik Institute, Harvard Medical School, Boston, MA, 02115

*For correspondence: Tristan Bell, 185 Cambridge St., CPZN 7250, Boston, MA, 02114,

bell@molbio.mgh.harvard.edu; Luke Chao, 185 Cambridge St., CPZN 7250, Boston, MA, 02114,

chao@molbio.mgh.harvard.edu.

Running Title: Extracellular vesicle formation in Prominin-family proteins

Keywords: prominin-1, extracellular vesicles, membrane biophysics, protein conformation, cholesterol-

binding protein

Preprint Servers: Deposited with bioRxiv under a CC-BY 4.0 license (BIORXIV/2023/566258)

Abstract

Prominin-1 (Prom1) is a five-transmembrane-pass integral membrane protein that associates with curved

regions of the plasma membrane. Prom1 interacts with membrane cholesterol and actively remodels the

plasma membrane. Membrane bending activity is particularly evident in photoreceptors, where Prom1

loss-of-function mutations cause failure of outer segment homeostasis, leading to cone-rod retinal

dystrophy (CRRD). The Tweety Homology (Ttyh) protein family has been proposed to be homologous to

Prominin, but it is not known whether Ttyh proteins have an analogous membrane-bending function.

Here, we characterize the membrane-bending activity of human Prom1 and Ttyh1 in native bilayer

membranes. We find that Prom1 and Ttyh1 both induce formation of extracellular vesicles (EVs) in

cultured mammalian cells and that the EVs produced are biophysically similar. Ttyh1 is more abundant in

EV membranes than Prom1 and produces EVs with membranes that are more tubulated than Prom1 EVs.

We further show that Prom1 interacts more stably with membrane cholesterol than Ttyh1 and that this

may contribute to membrane bending inhibition in Prom1 EVs. Intriguingly, a loss-of-function mutation in

Prom1 associated with CRRD induces particularly stable cholesterol binding. These experiments provide

mechanistic insight into Prominin function in CRRD and suggest that Prom and Ttyh belong to a single

family of functionally related membrane-bending, EV-generating proteins.

Introduction

Mammalian cells interact with the extracellular environment through proteins, lipids, and glycans at the

plasma membrane. Organized protrusive structures of the membrane such as microvilli and cilia are

hotspots for nutrient absorption, cell cycle regulation, and extracellular signaling[1,2]. Membrane

protrusions are organized both by interactions with the cytoskeleton and by sorting of proteins and lipids

within the membrane bilayer[2–4]. Prominin-1 (Prom1) is a five-transmembrane pass integral membrane

protein that interacts with membrane cholesterol at sites of membrane protrusion[5,6]. Prom1 was first

characterized as the target of AC133-1, a monoclonal antibody raised against pluripotent human epithelial

stem cells[7,8]. Since then, Prom1 has been identified in the apical membranes of most epithelial and

neuroepithelial cell types, but is only natively recognized by AC133-1 in stem cells and photoreceptors[9–

11]. In photoreceptors, Prom1 associates with a retinal cadherin (Protocadherin-21, Pcdh21) to promote

normal outer segment membrane homeostasis[12–14], and several Prom1 mutations are linked to hereditary

cone-rod retinal dystrophy (CRRD)[12,15–18]. Most animal genomes also encode at least one paralogous

prominin gene (Prominin-2) that is expressed in non-retinal cell types and localizes to positively curved

membrane regions[19].

When overexpressed, Prom1 dramatically reorganizes the plasma membrane of cultured mammalian cells

into long protrusions[20,21]. Small-molecule inhibitors of actin and tubulin do not impair the tubulation

phenotype, indicating that cytoskeletal interactions are not strictly required for membrane bending[20,21].

However, cells treated with cholesterol biosynthesis inhibitors or depletion agents do not exhibit

membrane tubulation, suggesting that cholesterol plays a pivotal role in the regulation of Prom1

function[20,21].

Prom1 also induces release of extracellular vesicles (EVs) that bleb from the apical plasma membrane of

differentiating epithelial stem cells[22]. Large Prom1 EVs (500-700 nm in diameter) appear to form from bulk

release of membrane from the cell midbody, whereas small Prom1 EVs (< 250 nm in diameter) likely bleb

off the membrane at organized membrane protrusions such as microvilli and cilia[22–24]. Prom1 EVs can be

detected in saliva, urine, semen, neural tube fluid, and lacrimal fluid of healthy adults[23,25].

Recently, the Tweety homology (Ttyh) protein family was hypothesized to be a distant homolog of the

prominins[20]. Ttyh proteins share prominins' five-transmembrane topology but have a more minimal

extracellular domain[26]. Most animals have three paralogous Ttyh proteins (Ttyh1, Ttyh2, and Ttyh3)[27] that

all are predominantly expressed in neural tissues[28–33]. Notably, overexpression of Ttyh1 in cultured cells

induces plasma membrane tubulation that is strikingly similar to that observed with Prom1[27,34–36].

Defining mechanisms of membrane bending by Prominin-family proteins is foundational for

understanding retinal disease and stem cell development. Here, we reconstitute EV formation by Prom1

and Ttyh1 to characterize their functions in a native membrane bilayer system. To our knowledge, we

present the first evidence that Ttyh1 induces formation of EVs that are functionally and biophysically

analogous to Prom1 EVs. Ttyh1 EVs are more tubulated and protein-rich than Prom1 EVs. Prom1 forms a

more stable interaction with cholesterol than Ttyh1, and when depleted of cholesterol, Prom1 EVs mimic

the tubulation observed in Ttyh1 EVs. A CRRD-associated Prom1 mutant (W795R) binds cholesterol more

stably than wild-type protein, suggesting that a less dynamic interaction between Prom1 and cholesterol

inhibits membrane bending. These findings contribute to a path toward mechanistic understanding of

Prominin-family protein function in different tissues, including the role of Prom1 in retinal pathology.

Results

Reconstitution and purification of Prom1 EVs

To understand how Prom1 interacts with and reshapes native membranes, we sought a method to purify

Prom1 without disturbing protein-lipid interactions. Because Prom1 induces EV formation in epithelial

stem cells[23], we asked whether Prom1 can also form EVs in cultured cells. C-terminally Strep-tagged Prom1

was detected in immunoblots of the conditioned media from transfected cells, but not in cells treated

with a no-plasmid mock transfection (Fig 1A, 1B). To validate that the ~120 kDa Strep-reactive band we

observe is indeed Prom1-Strep, we immunoblotted with the Prom1-specific antibody AC133-1[37] and

observed robust staining (Fig 1C). To confirm that EV production is not an artifact of transfection, we

introduced Prom1-Strep into Expi293 cells by lentiviral transduction to generate a stable polyclonal

overexpression cell line. A Strep-reactive band consistent with mature Prom1 was detectable in the

conditioned media from these cells over several rounds of cell passaging (Fig S1A). We assessed the

protein composition of Prom1 EVs using SDS-PAGE and silver nitrate staining. Prom1 was a constituent of

our samples alongside several other proteins (Fig S1B). We cannot conclude whether these proteins

specifically interact with Prom1 in EVs, are hitchhikers enclosed within the EVs, or originate from

endogenous co-purifying EVs.

We next purified the Prom1 EVs using differential centrifugation, ultracentrifugal concentration, and size-

exclusion chromatography (SEC) based on isolation methods previously described for endogenous small

Prom1 EVs (Fig 1A)[23,38]. Using dynamic light scattering (DLS), we measured the solution sizes of the purified

Prom1 EVs. The sample was monodisperse with an average diameter of 164 ± 14 nm, considerably larger

than previously reported 50-80 nm EVs measured by negative-stain transmission electron microscopy (NS-

TEM) in samples purified from biological fluids[23] (Fig 1D). Upon treatment with PNGase F, an enzyme that

cleaves *N*-glycan groups from proteins, the ~120 kDa Prom1 gel band shifted down to its predicted

molecular weight of 102 kDa (Fig 1E).

To verify that the purified particles are truly EVs (secreted particles with intact bilayer membranes), we

vitrified purified samples and imaged them using transmission cryo-electron microscopy (cryo-TEM) (Fig

1F). We observed spherical EVs with bilayer membranes (Fig 1F, inset). The diameters of EVs measured

from cryo-TEM images was bimodal with an average diameter of 117 ± 58 nm, somewhat lower than our

measurements from DLS (Fig 1H). We next used NS-TEM to directly compare our reconstituted Prom1 EVs

with previously characterized endogenous Prom1 EVs[23] (Fig 1G). We observed a distribution of largely

circular EVs ranging in size from ~50 nm to ~250 nm in diameter, but more skewed toward smaller

diameters than observed in the cryo-TEM data (Fig 1H). In addition, EVs had rough edges and internal

depressions in NS-TEM, a characteristic feature of EV fixation and dehydration[23] (Fig 1H). Because solution

DLS measurements (164 ± 14 nm) suggest slightly larger EV diameters than our NS-TEM (123 ± 73 nm) or

cryo-TEM (117 ± 58 nm) measurements, we speculate that sample fixation/dehydration or vitrification

may induce deformation and potentially fission of the reconstituted EVs. This effect may have similarly

impacted previous characterization of endogenous Prom1 EVs[23]. We conclude that reconstituted Prom1

EVs have similar morphology to endogenous EVs but may be slightly larger in size.

We next asked whether Prom1 in purified EVs retains known functional behavior of Prom1 from

endogenous membranes. Prom1 stably binds the retinal cadherin Pcdh21 in photoreceptors[12–14]. To

determine if Prom1 can co-traffic with Pcdh21 in our EV samples, we co-transfected Expi293 cells with

Strep-tagged Prom1 and Flag-tagged Pcdh21 and looked for the presence of each component in cells,

conditioned media (CM), and clarified conditioned media (CCM). Pcdh21 was only observed in CCM

samples when co-expressed with Prom1, indicating that Prom1 is necessary to traffic Pcdh21 into this

class of EVs (Fig 1I). To establish a direct interaction between Prom1 and Pcdh21, we co-expressed Prom1-

Strep with mNeonGreen-tagged Pcdh21 and immunopurified Prom1 from EVs solubilized with 1% n-

dodecyl-ß-D-maltoside (DDM). Pcdh21-associated mNeonGreen reproducibly co-purified with detergent-

solubilized Prom1 compared to a control condition lacking Prom1, indicating that the two proteins

physically interact in purified EVs (Fig 1J).

Figure 1. (A) EV expression and purification graphic protocol. **(B)** Anti-Strep immunoblot of cell pellet,

conditioned media (CM), clarified conditioned media (CCM), or SEC-purified EVs from mock-transfected

or Prom1-Strep-transfected Expi293 cells. Arrowhead indicates the expected molecular weight of

Prom1. **(C)** AC133-1 immunoblot of SEC-purified Prom1 EVs. Arrowhead indicates the expected molecular weight of Prom1. **(D)** Dynamic light scattering (solution size) measurement of purified Prom1 EVs. **(E)** Anti-Strep immunoblot of Prom1 EVs treated with or without PNGase F to remove *N*-glycan moieties. **(F)** Cryo-TEM images of purified Prom1 EVs. Inset image is magnified to emphasize membrane bilayer density. Images are lowpass filtered to 5 Å to enhance contrast. **(G)** NS-TEM image of purified Prom1 EVs. **(H)** Measured diameters of Prom1 EVs from Cryo-TEM or NS-TEM images. (n = 322 and n = 176 for NS-TEM and cryo-TEM measurements, respectively.) **(I)** Anti-Strep (Prom1) and anti-Flag (Pcdh21) immunoblots of CM and CCM from cells transfected with Prom1-Strep, Pcdh21-Flag, or both. Note that Pcdh21 is only detected in CCM when co-expressed with Prom1. Filled and empty arrows indicate expected molecular weights of glycosylated Prom1-Strep and Pcdh21-Flag, respectively. **(J)** Comparative fluorescence measurements of Pcdh21-mNeonGreen co-immunoprecipitated with or without Prom1-Strep (n = 3, ** p < 0.01 by Student's two-tailed unpaired t test).

Mutations in the Prom1 transmembrane domain impair EV formation

Prom1 binds cholesterol in native membranes and the Prom1-cholesterol interaction is necessary for remodeling the plasma membrane[21], but the mechanism of cholesterol-dependent Prom1 function is unclear. We analyzed the sequence of human Prom1 to identify candidate cholesterol recognition amino acid consensus (CRAC, [L/V]-X_{1-5}-[Y/F]-X_{1-5}-[K/R]) or mirrored CRAC (CARC, [K/R]-X_{1-5}-[Y/F]-X_{1-5}-[L/V]) sequences in the transmembrane helices, as these motifs often predict cholesterol binding in membrane proteins[39]. Human Prom1 contained four CRAC and two CARC motifs, of which none were completely conserved among metazoans and only one (CRAC-3) was modestly evolutionarily conserved (Figs S2A, S2B). We therefore turned to a more comprehensive evolutionary analysis of prominin proteins to identify evolutionarily conserved Prom1 transmembrane residues.

We curated prominin sequences from across eukaryotes, considering a prominin to be a sequence with five predicted transmembrane helices, two large extracellular loops, two small intracellular loops, and homology to annotated metazoan prominin sequences. Putative prominin sequences were identified across metazoans as well as in fungi, excavates, SAR (stramenopiles, alveolates, and rhizarians), and green plants (Figs S3A, S3B). Strikingly, Trp-795 was nearly perfectly conserved across metazoa and eukaryotic outgroups, making it by far the most conserved residue (excluding Cys residues positioned to form disulfides in AlphaFold2 models) across the curated prominin sequences (Figs 2A, S3A). The conserved Trp-795 residue was of particular interest as a missense mutation at this site (W795R) is implicated in hereditary CRRD cases[40].

We generated Prom1-Strep variants with mutations to disrupt each CRAC or CARC motif, the disease-
associated W795R mutation, and point mutations directed against residues predicted to be in close
contact with Trp-795 in Alphafold2 models (Fig 2B, Table ST1)[41]. We then assessed EV formation by
quantifying the Prom1-Strep signal secreted in EVs or retained in the cellular membranes. We found that
several mutations to the Prom1 TM domains significantly disrupted the proportion of Prom1 secreted in
EVs, with the CARC-1 and W795R mutants demonstrating the most severe effects (Figs 2C, S4). In most
cases, this effect arose from both decreased expression of Prom1 and increased retention of protein in
cells. However, the CRAC-1 and CARC-2 mutants were more highly expressed than WT Prom1, with the
entire effect arising from cellular membrane retention. To better characterize the morphology of the
Prom1 variants, we expressed and purified a subset of mutant Prom1 EVs at larger scale. Each of the
mutants produced EVs that are 150-200 nm in diameter and monodisperse by DLS analysis, generally
similar to WT Prom1 EVs (Fig 2D). Thus, transmembrane domain mutations in Prom1 primarily alter the
quantity of EVs produced rather than EV size.

To better understand the mechanism of reduced Prom1 EV formation by the CRRD-associated W795R
mutant, we engineered HeLa cells that stably express C-terminally StayGold-tagged WT (Prom1-SG) or
W795R Prom1 (Prom1[W795R]-SG) under the EF-1α promoter. Prom1-SG signal colocalized with Wheat
Germ Agglutinin (WGA)-stained plasma membrane and intracellular vesicles, but Prom1[W795R]-SG
concentrated on WGA-positive intracellular vesicles (Fig 2E, Movie SM1, Movie SM2). Despite the
apparent low signal of plasma-membrane localized Prom1-SG, line scan analysis shows correlation
between Prom1-SG signal across the cell junctions (Fig S5 left). In contrast, there is a marked absence of
Prom1[W795R]-SG signal localized on the plasma membrane (Fig S5 right). We infer that fluorescent
W795R Prom1 is trafficked to the plasma membrane less efficiently than the wild-type protein.

Figure 2. (A) AlphaFold2 model of human Prom1[41] with residues color-coded by level of conservation across a multiple-sequence alignment of metazoan prominin sequences. **(B)** Possible network of interactions between Trp-795 and several adjacent aromatic residues in human Prom1. **(C)** Quantification of anti-Strep immunoblots of cellular and clarified conditioned media pools of Prom1-Strep mutants relative to wild-type Prom1-Strep (n = 3, [n.s.] p > 0.05, * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001 by Student's two-tailed unpaired t test). Raw blot images are included in Supplemental Figure S4. **(D)** Mean particle diameter of purified Prom1-Strep EVs measured by dynamic light scattering. Error bars indicate S.D. (n = 5, [n.s.] p > 0.05, * p < 0.05 by Student's two-tailed unpaired t test). **(E)** Confocal fluorescence microscopy images of HeLa cells stably expressing WT (top) or W795R (bottom) Prom1-StayGold (yellow), stained with wheat germ agglutinin (WGA) (blue). Scale bar is 10 μm. Line scan traces across cell junctions are included in Supplemental Figure S5.

Prominin homolog Ttyh1 produces EVs

Tweety homology (Ttyh) proteins are proposed prominin homologs that share the five-transmembrane topology of prominins but have shorter extracellular domains (~120 amino acids in Ttyh vs ~280 amino

acids in Prom)[20] (Fig 3A). Given the predicted sequence homology between Prom and Ttyh, we

hypothesized that Ttyh may also produce EVs. Evolutionary analysis of metazoan Ttyh proteins does not

indicate conserved CRAC or CARC sites, nor does it suggest any conserved transmembrane residue

analogous to Trp-795 in metazoan prominins (Fig 3B).

We expressed C-terminally Strep-tagged Ttyh1 in Expi293 cells and purified EVs using the same procedure

as for Prom1 EVs (Fig 1A). We detected Ttyh1-Strep in CM, CCM, and SEC-purified EV fractions (Fig 3C).

Dynamic light scattering indicated that purified Ttyh1 EVs are monodisperse with an average diameter of

180 ± 10 nm, similar to the average diameter of WT Prom1 EVs (164 ± 14 nm) (Fig 3D).

We next characterized the morphology of purified Ttyh1 EVs using NS-TEM. We found that Ttyh EVs

adopted striking long and bent tubular structures with much higher frequency than Prom1 EVs (6.4% of

Ttyh1 EVs versus 0.3% of Prom1 EVs, n = 1357 and n = 322 respectively) (Figs 3E, 3H). Ttyh1 EVs visualized

by NS-TEM were on average smaller than Prom1 EVs, and were similarly smaller than expected from

solution DLS measurement (Figs 3G, 3H). Furthermore, we observed that the smallest Ttyh1 EVs were

similar in diameter to the short-axis caliper diameter of the tubular EVs (42 ± 6 nm) and we observed cases

where tubular EVs appeared to be in the process of dividing into smaller EVs (Figs 3E white arrowheads,

3G). Although the NS-TEM conditions deviate from a solvated physiological state, they do suggest that

Ttyh1 supports comparatively greater membrane curvature than Prom1 in these EVs.

We further analyzed Ttyh1 EVs by cryo-TEM to verify that the purified sample indeed contained EVs with

intact bilayer membranes (Fig 3F). Like Prom1 EVs, Ttyh1 EVs have a bimodal size distribution with local

maxima around 60 nm and 140 nm (Fig 3G). We also observed a population of EVs exhibiting the tubular

phenotype seen in NS-TEM that again substantially exceeded that seen with Prom1 EVs (8.8% of Ttyh1

EVs versus 0.8% of Prom1 EVs, n = 2224 and n = 176 respectively) (Fig 3H). This supports the observation

that EV bending is more frequent in Ttyh1 EVs than Prom1 EVs.

Molecular crowding is known to contribute to membrane bending[42]. To test if Ttyh1 may promote EV

membrane bending through a crowding mechanism, we quantified relative protein-to-membrane ratios

for Prom1 and Ttyh1 EVs. From identical amounts of transfected cell media, purified EVs contained 8.8 ±

1.0-fold more Ttyh1 protein than Prom1 (Figs 3I, S6A). Both Prom1 and Ttyh1 EV membrane surface area

was significantly more abundant than the background level of EVs produced in a mock-transfected control

culture, but Ttyh1 produced 5.6 ± 0.8-fold more total EV membrane than Prom1 (Figs 3J, S6B). From these

measurements, we estimate that Ttyh1 is present at ~5-fold higher concentration in EV membranes (Fig

S6C). Thus, molecular crowding effects may be a factor contributing to the increased membrane bending

we observe in Ttyh1 EVs.

To further characterize purified Prom1 and Ttyh1 EVs, we subjected purified EVs to equilibrium sucrose

gradient sedimentation to resolve populations by density. We observed a single population of Prom1 EVs

(centered on fraction 7) but resolved two distinct populations of Ttyh1 EVs with densities lower (centered

on fraction 6) and higher (centered on fraction 9) than the Prom1 EVs (Fig 3K). When Prom1 and Ttyh1

were co-expressed, we observed that the resulting EVs contained more Prom1 than Ttyh1 but followed

the bimodal distribution of Ttyh1 EVs (Fig 3L). Immunoblots of the sucrose gradient fractions showed that

co-expressed Prom1 and Ttyh1 peak in the same sucrose gradient fractions, suggesting that Prom1 and

Ttyh1 co-elute in the same EV populations (Fig 3L, 3M). We conclude that Prom1 and Ttyh1 can be

secreted into the same EVs, and thus likely originate from the same plasma membrane domains.

Figure 3. (A) Comparison of a Ttyh1 subunit from cryo-TEM structure 7P5J[32] with an AlphaFold2-
predicted Prom1 monomer[41]. **(B)** Residue-level conservation among metazoan Ttyh proteins plotted
onto a subunit of human Ttyh1. No residue analogous to Prom1 Trp-795 is present in Ttyh. **(C)** Anti-strep
immunoblot comparing Prom1-Strep EVs and Ttyh1-Strep EVs throughout different stages of
purification. Filled and empty arrowheads indicate the expected positions of Prom1 and Ttyh1,
respectively. Doublet and higher bands in Ttyh1 lanes are products of on-gel disulfide crosslinking in

concentrated samples. **(D)** Dynamic light scattering measurement of EV diameter for purified Prom1-
Strep, Ttyh1-Strep, or Prom1-Strep + Ttyh1-Strep co-expression EVs. **(E)** Representative NS-TEM images
of Ttyh1-Strep EVs. White arrowheads indicate possible sites of EV fission. **(F)** Representative cryo-TEM
images of Ttyh1-Strep EVs. Magnified insets show bilayer density at highly curved membrane segments.
Images are lowpass filtered to 5 Å to enhance contrast. **(G)** Comparison of Prom1 or Ttyh1 EV diameter
in NS-TEM or cryo-TEM images. (n = 322, n = 1357, n = 176, and n = 2224 for Prom1 NS-TEM, Ttyh1 NS-
TEM, Prom1 cryo-TEM, and Ttyh1 cryo-TEM measurements, respectively.) **(H)** Comparison of Prom1 or
Ttyh1 EV roundness in NS-TEM or cryo-TEM images. Inset plots only include EVs with roundness ≤ 0.8. (n
= 322, n = 1357, n = 122, and n = 1546 for Prom1 NS-TEM, Ttyh1 NS-TEM, Prom1 cryo-TEM, and Ttyh1
cryo-TEM measurements, respectively.) **(I)** Quantification of Prom1 and Ttyh1 protein levels in EVs from
anti-Strep Western blots (n = 3). Immunoblots used for quantification are included in Supplementary
Figure S6A. **(J)** Quantification of total EV membrane area from NS-TEM micrographs (n = 236, *** p <
0.001 by unpaired Mann-Whitney test). Representative micrographs used for quantification are included
in Supplementary Figure S6B. **(K)** Anti-strep immunoblots of fractions from sucrose gradient equilibrium
sedimentation of Prom1-Strep EVs (top) or Ttyh1-Strep EVs (bottom). **(L)** Anti-strep immunoblots of
fractions from sucrose gradient equilibrium sedimentation of EVs from cells co-expressing Prom1-Strep
and Ttyh1-Strep. Filled and empty arrowheads indicate the expected positions of Prom1 and Ttyh1,
respectively. **(M)** Quantification of immunoblots in panels *K* (top) and *L* (bottom).

Ttyh1 binds cholesterol less stably than Prom1

Because cholesterol interaction is known to regulate membrane bending by Prom1, we asked whether

Prom1 and Ttyh1 bind cholesterol similarly[20,21]. To comparatively measure binding, we developed a

cholesterol co-immunopurification assay (hereafter referred to as chol-IP) to quantify interaction of

Prom1 with fluorophore-labeled cholesterol (Fig 4A). Prom1-Strep or Ttyh1-Strep were transfected into

Expi293 cells, and cells were labeled with a low concentration of fluorescent cholesterol. EVs were purified

from the conditioned media, solubilized with nonionic detergent (1% DDM), and immunoprecipitated with

Strep resin trace labeled with blue fluorescent protein (mTagBFP2-Strep). We then collected

epifluorescence micrographs of the resin particles, computationally segmented and filtered each image,

and quantified bound cholesterol using mTagBFP2 as a normalizing control (Fig 4A). This approach permits

sensitive quantification of bound lipid while efficiently excluding autofluorescent and refractive artifacts.

We performed chol-IP analysis on Prom1 or Ttyh1 EVs labeled with BODIPY-cholesterol. Surprisingly,

fluorescent cholesterol robustly copurified with Prom1 but not with Ttyh1 (Fig. 4B). To verify that this

difference is not a result of nonspecific fluorophore-protein interactions, we replicated the observation

using a different fluorescent cholesterol analog (AlexaFluor647-cholesterol) (Fig S7). We further verified

that the cholesterol is specifically bound to Prom1 by quantifying bound cholesterol at different

concentrations of solubilizing detergent (Fig 4C). Nonspecifically associated fluorescent cholesterol in

Ttyh1 samples is delipidated at 0.04-0.2% DDM, 5-25-fold below the assay working concentration,

suggesting that the assay informs on specific lipid binding that is resistant to delipidation. Although this

method cannot inform on transient or kinetically unstable protein-lipid interactions, we infer that Prom1

and Ttyh1 differ in how stably they interact with membrane cholesterol.

Given the striking difference in cholesterol binding stability between Prom1 and Ttyh1, we next asked

whether Prom1 mutants also disrupt cholesterol interaction stability. We purified BODIPY-cholesterol-

labeled EVs produced by WT and mutant forms of Prom1 and subjected equal input concentrations of

Prom1 to chol-IP analysis. To our surprise, none of the mutants substantially disrupted cholesterol

binding, but the W795R mutant and the two CARC domain mutants bound cholesterol significantly more

stably than WT Prom1 (Fig 4D). We chose to focus our efforts on the W795R mutant as it is a naturally

occurring single-residue mutation with exceptional evolutionary conservation and a clinically validated

disease phenotype. To verify that differences in cholesterol binding between wild-type and W795R Prom1

do not arise from a gross oligomerization defect, we purified EVs from cells co-transfected with Strep-

tagged and mScarlet-tagged Prom1 and measured co-purification of Prom1-mScarlet on Strep resin.

Prom1 W795R co-purified with ~60% as much mScarlet fluorescence as WT Prom1, indicating that W795R

indeed multimerizes, albeit with reduced efficiency (Fig 4E).

Figure 4. (A) Cholesterol co-immunoprecipitation (Chol-IP) assay graphic protocol. **(B)** Comparison of relative cholesterol bound by Prom1-Strep or Ttyh1-Strep (n = 3, ** p < 0.01 by Student's two-tailed unpaired t test). **(C)** Fluorescent cholesterol co-immunopurified by Prom1-Strep or Ttyh1-Strep at different concentrations of DDM detergent. Error bars indicate S.D. (n = 3). **(D)** BODIPY-cholesterol binding measurements for WT and mutant variants of Prom1. Error bars indicate S.D. (n = 3, $^{n.s.}$ p > 0.05, * p < 0.05, ** p < 0.01 by Student's two-tailed unpaired t test). **(E)** Red fluorescence signal from anti-Strep immunopurification of DDM-solubilized EVs from cells expressing Prom1-mScarlet ("mSc") or both Prom1-mScarlet and Prom1-Strep ("Strep + mSc"). Error bars indicate S.D. (n = 3).

Cholesterol binding stability inversely correlates with Prom1 EV membrane bending

Because Ttyh1 promotes greater membrane curvature than Prom1 but does not stably bind cholesterol, we hypothesized that cholesterol binding stability may contribute to negative regulation of membrane bending by prominin-family proteins. If this is the case, then W795R Prom1, which binds cholesterol more stably than WT protein, should not produce EVs that exhibit the tubular morphology observed in Ttyh1 EVs. After purifying W795R Prom1 EVs, we observed a similar solution size for WT (164 ± 14 nm) and W795R EVs (186 ± 13 nm) (Fig 5A). We measured the size and shape of W795R EVs by cryo-TEM and observed largely spherical vesicles with some local deformations, similar to what we observed with WT Prom1 EVs (Fig 5B). Generally, WT and W795R Prom1 EVs are both spherical, with W795R having no vesicles that fall into our tubular morphology classification (n = 1211) (Fig 5C). Though both large and small

diameter EV populations were observed by cryo-TEM, W795R Prom1 had a larger fraction of small EVs

than WT Prom1, indicating possible decreased stability or increased fissile propensity in the W795R Prom1

EVs (Fig 5D).

If stable cholesterol binding negatively regulates membrane bending, then depleting cholesterol away

from EV membranes could induce a Ttyh1-like morphology in Prom1 EVs. We tested this model directly

by purifying WT Prom1 or Ttyh1 EVs and treating them with methyl beta cyclodextrin (mBCD), a

compound that extracts cholesterol from membranes[21,43]. After re-purifying EV samples away from free

mBCD and mBCD-cholesterol complexes, Prom1 EVs treated with mBCD (2.5-10 mM) exhibited a decrease

in cholesterol content (Fig 5E). A control set of Ttyh1 EVs treated with 0 or 10 mM mBCD showed a similar

change in cholesterol content (Fig 5E). Treated and untreated EVs were then analyzed by NS-TEM to

compare EV size and morphology (Fig 5F). We observed a 5.4-fold increase in the fraction of Prom1 EVs

that deviated from spherical membrane topology after 10 mM mBCD treatment compared to untreated

EVs (3.0% versus 0.6% of EVs, n = 232 and n = 356 respectively) (Fig 5G). Importantly, identical treatment

of Ttyh1 EVs was used as a nonspecific control for cholesterol depletion from EVs. Ttyh1 EVs treated with

mBCD under identical conditions only showed a 1.4-fold increase in deformed EVs (2.1% at 10 mM mBCD

versus 1.5% in mock-treated EVs, n = 433 and n = 535 respectively) (Fig 5G). Thus, cholesterol-depleted

Prom1 EVs are more prone to membrane deformation than untreated EVs.

Figure 5. (A) Dynamic light scattering EV diameter measurements of purified WT or W795R Prom1-Strep EVs. **(B)** Representative cryo-TEM images of W795R Prom1-Strep EVs. Magnified insets show bilayer density. Images are lowpass filtered to 5 Å to enhance contrast. **(C)** Comparison of WT or W795R EV roundness in cryo-TEM images. Inset plots only include EVs with roundness ≤ 0.8. (n= 176 and n = 1211 for WT and W795R Prom1 cryo-TEM measurements, respectively.) **(D)** Comparison of WT or W795R Prom1 EV diameter in cryo-TEM images. (n= 122 and n = 821 for WT and W795R Prom1 cryo-TEM measurements, respectively.) **(E)** Cholesterol content of Prom1 or Ttyh1 purified EV samples after treatment with methyl-beta cyclodextrin (mBCD). Error bars indicate S.D. (n = 3). **(F)** Representative NS-

TEM images of purified mBCD-treated EVs. **(G)** Quantification of EV diameter (*top*) and roundness (*middle*) from NS-TEM images, as well as quantification of tubulated EVs (roundness < 0.6) (*bottom*).

Discussion

Prom1 and Ttyh1 bend membranes and form EVs

We report here that both Prom1 and Ttyh1 can induce EV formation in cultured cells upon overexpression (Figs 1B, 3C). Though we do not claim that Ttyh1 endogenously induces EV formation, we note that it is sufficient to do so in a recombinant system with similar or higher efficiency than Prom1. Given similar membrane remodeling and tubulation behavior of Prom1 and Ttyh1 in cell culture[20,35], Ttyh1 EVs may form by a similar mechanism to Prom1 EV formation. We find that Prom1 and Ttyh1 form EVs that are of similar size and that the two proteins exist in the same pool of EVs when co-expressed (Figs 3L, 3M). Because Prom1 EVs arise directly from blebbing of the plasma membrane[23], this finding suggests that Prom1 and Ttyh1 traffic to similar regions of the plasma membrane.

Membrane bending by Prom1 is critical for maintaining outer segment membrane homeostasis in photoreceptors. Intriguingly, photoreceptors with impaired expression of the disc rim stabilizing protein Peripherin accumulate EVs 200-280 nm in diameter at the outer segment base[44]. We observe formation of Prom1 EVs of relatively similar size in our reconstituted system (Fig 1D). This may be an example of endogenous regulation of Prom1-induced membrane curvature, with curvature-stabilizing proteins like Peripherin preventing the evaginating membrane from budding into EVs.

Ttyh1 forms EVs with membranes that are more tubulated than Prom1 EV membranes

Ttyh1 produces EVs similar in size and density to Prom1 EVs (Figs 3D, 3K, 3L, 3M). However, Ttyh1 EVs are dramatically more tubulated that Prom1 EVs (Figs 1F, 1G, 3E, 3F, 3H). Several factors could contribute to this increase in membrane bending. First, Ttyh1 is present in EVs at concentrations approximately 5-fold higher than Prom1 (Figs 3I, 3J). Membrane proteins can induce curvature through molecular crowding when enriched at sufficiently high local concentrations[42]. Second, Ttyh1 appears to bind cholesterol less stably than Prom1 does (Fig 4B). Membrane cholesterol is necessary for Prom1-mediated remodeling[20,21] and is present at similar levels in purified Prom1 and Ttyh1 EVs (Fig 5E), indicating that it is undoubtedly important for EV formation by both proteins. Our results do not indicate that Ttyh1 does not bind

cholesterol, merely that it does not form an interaction that is sufficiently kinetically stable to be co-
immunoprecipitated.

Ideally, molecular-crowding and cholesterol-binding effects could be differentiated by modulating
protein-to-lipid ratios in EV membranes. Unfortunately, our native purified EVs do not provide a tractable
path to perform these experiments. Future experiments reconstituting Prom1 and Ttyh1 into artificial
liposomes with variable cholesterol content will be necessary to fully elucidate the mechanistic
differences in membrane bending between Prominin and Ttyh proteins.

Cholesterol binding stability may contribute to membrane bending regulation by Prom1
We find here that Prom1 in EVs forms a stable interaction with cholesterol that is resistant to delipidation
by nonionic detergent (Figs 4B, 4C). A disease-associated mutation in a strikingly conserved residue in the
fifth transmembrane helix of Prom1, W795R, substantially stabilizes cholesterol binding (Figs 2A, 4D). In
AlphaFold2 models, Trp-795 forms a series of base-stacking interactions with Phe residues in neighboring
transmembrane helices (Fig 2B)[41]. We find that conservative mutations in several of these adjacent
aromatic residues impair EV formation by Prom1, but do not mimic the stable cholesterol binding of
W795R (Figs 2C, 4D). Cholesterol is asymmetrically distributed in eukaryotic membranes, with a bias
toward the outer leaflet[45]. Prom1 Trp-795 is predicted to be positioned toward the outer leaflet, possibly
potentiating interactions with cholesterol. We also observe increases in cholesterol binding stability for
mutations to two CARC domains in Prom1 (Fig 4D). This effect could arise because these residues interface
with cholesterol or because they destabilize the transmembrane domain in a manner that mimics the
effect of the W795R mutation. Given the poor evolutionary conservation of the two CARC motifs in human
Prom1, we favor the latter explanation.

We further show that the W795R mutation prevents efficient trafficking and plasma membrane insertion
of Prom1 from the endomembrane to the plasma membrane (Fig 2E). This observation explains the low
yield of W795R EVs produced in cell culture (Fig 2C). W795R Prom1 is expressed well but strongly localizes
to WGA⁺ intracellular vesicles. These vesicles are believed to traffic downstream of the trans-Golgi,
suggesting that the roadblock in W795R Prom1 endomembrane trafficking occurs between the trans-Golgi
and the plasma membrane[46]. Several studies have demonstrated that protein trafficking through the ER
and Golgi is highly sensitive to cholesterol levels[47,48]. Our findings suggest that the dynamic interaction of

Prom1 with cholesterol may promote efficient maturation and trafficking of Prom1 between the

endomembrane system and the plasma membrane.

Prominin and Tweety homology proteins are homologous and functionally analogous

Prominin and Ttyh proteins are both biologically implicated in membrane bending. Prominins localize to

cholesterol-rich domains of the plasma membrane and drive protrusion-shed EV formation in

differentiating stem cells[21–23]. Ttyh proteins are associated with dendritic spikes in healthy neurons and

with tumor microtube formation in aggressive astrocytoma[27,34–36]. Given their sequence homology and

transmembrane topology, and their similarities in membrane localization, membrane bending and EV

formation behavior, we suggest that Prom and Ttyh proteins be considered members of a broader

prominin family of membrane remodeling proteins. Further work will be required to fully characterize the

functional specialization of Prom and Ttyh proteins.

Materials and Methods

Expression Constructs

All Prom1 expression constructs were generated by site-directed mutagenesis from pCS2-Prom1-YFP (a

gift from N. Sasai, Nara Institute of Science and Technology). The base Prom1 construct used is the human

S1 isoform (NCBI accession NP_001139319.1)[49]. Prom1-Strep was subcloned into pLV-EF1a vector (a gift

from K. Hochedlinger, Massachusetts General Hospital) for lentiviral transduction. Human Pcdh21 isoform

1 (NCBI accession NP_149091.1) was synthesized (GenScript) and cloned into a pCDNA3.1 vector (Thermo

Fisher) for mammalian cell transfection. mTagBFP2 was subcloned from pBAD (Addgene #54572, a gift

from Michael Davidson) into a pET28a vector with a C-terminal Strep tag for bacterial overexpression.

Ttyh1 was expressed from a pLX304 vector after addition of C-terminal Strep and His tags to an existing

construct (Addgene #161676, a gift from Mike McManus). To stably express fluorescently tagged Prom1

WT and W795R variants in HeLa cells for live cell imaging, Prom1-StayGold and Prom1[W795R]-StayGold

sequences were synthesized (GenScript) and cloned into a P2555 vector (a gift from S. Jakobs, Max Planck

Institute for Biophysical Chemistry) to yield constructs pAH18 and pAH20, respectively.

Cell Line Construction

Cell lines were regularly tested for mycoplasma contamination. Lentiviral transduction of Prom1-Strep

into Expi293 cells was performed using a modification of published protocols[50,51]. Briefly, lentiviral

particles were produced in 293T cells (ATCC) by transient transfection of pLV-EF1a-Prom1-Strep with

VSVG and Delta 8.9 plasmids (a gift from K. Hochedlinger, Massachusetts General Hospital) using the

Lipofectamine 3000 system (Thermo Fisher) and incubated overnight at 37 °C, 5% CO_2. Expi293 cells

(Thermo Fisher) raised for several passages in suspension culture were seeded onto adherent tissue

culture plates in adherent culture media (DMEM (Gibco) supplemented with 10% Fetal Bovine Serum

(Gibco) and 1x Penicillin/Streptomycin (Gibco)) and incubated overnight at 37 °C, 5% CO_2 to form an

adherent monolayer. Transfected cells were exchanged into fresh adherent culture media. Culture media

was harvested after an additional 48 h, filtered through a 0.45 µm vacuum unit, and concentrated from

70 mL to 0.3 mL in PBS buffer by ultracentrifugation according to established protocols[51]. Adherent

Expi293 cells were infected with concentrated lentiviral particles at 75% confluency in adherent culture

media supplemented with 8 µg/mL polybrene (Sigma-Aldrich), then incubated for 48 h at 37 °C, 5% CO_2.

Cells were gently washed with PBS, exchanged into adherent culture media, and incubated for 24 h at

37 °C, 5% CO_2. Subsequently, the cells were subjected to antibiotic selection by exchange into adherent

culture media containing 2 µg/mL blasticidin (Gibco) for 10 days with regular exchange into fresh selective

media and passaging to prevent cells from achieving full confluency. Selection was considered complete

when the majority of cells died and antibiotic-resistant foci recolonized the culture plate. The cells were

then trypsinized (Gibco) and transitioned back to suspension culture in modified suspension culture media

(Expi293 media (Gibco) supplemented with 1% Fetal Bovine Serum) at a density of 1.0×10^6 live cells per

mL of culture and incubated for 48 h at 37 °C, 8% CO_2 with 125 rpm rotation. Once the suspension culture

reached a density of 3.0×10^6 live cells per mL of culture, cells were re-passaged in 1.0×10^6 live cells per

mL of culture in fresh modified suspension culture media supplemented with 1.5 µg/mL Blasticidin, and

subsequently re-passaged into this media condition every 2 days.

Stable cell lines were constructed using lentiviral and AAV transduction methods according to established

protocols[50,51]. For generation of stable HeLa cells expressing WT Prom1-StayGold and Prom1[W795R]-

StayGold off the AAVS1 locus, the donor plasmids pAH18 or pAH20 were co-transfected with the nuclease

plasmid PX458-AAVS1 (a gift from S. Jakobs, Max Planck Institute for Biophysical Chemistry) using

Lipofectamine 3000 (Thermo Fisher). Transfected cells were selected with 10 µg/mL blasticidin (Gibco)

starting 48 h post-transfection for 7 days. Stable clones were expanded for 10 days, and single-cell GRP-

positive clones were obtained using a FACS AriaII Cell Sorter (BD Biosciences). After clonal expansion,

positive clones were detected and verified by fluorescence imaging.

Prom1 and Ttyh1 EV Reconstitution

Prom1 and Ttyh1 EVs were reconstituted by expression in Expi293 suspension cells. Briefly, Expi293 cells

grown in serum-free Expi293 media at 37 °C, 8% CO_2 with 125 rpm rotation to a density of 3.0 x 10^6 live

cells per mL of culture were transiently transfected with an appropriate plasmid at 1 µg of DNA per 1 mL

of culture using the Expifectamine transfection kit (Thermo Fisher) according to the manufacturer

protocol. After 48 h, cultures were centrifuged for 5 min at 500 x g, the media discarded, and the cells

resuspended in the same volume of fresh Expi293 media and returned to incubate for an additional 48 h.

After this final incubation, cultures were centrifuged for 5 min at 1500 x g and the conditioned media

transferred to clean 50 mL conical tubes.

EVs labeled with fluorescent cholesterol analogs were generated as described above with the following

modifications. Two days after transfection, cells were transferred to 50 mL conical tubes, centrifuged for

5 min at 500 x g, then resuspended in an equal volume of Expi293 media with fluorescent cholesterol

added to a final concentration of 4 µM. Cells were transferred back to shaker flasks and incubated for 24

h before conditioned media was harvested.

EV Purification

Conditioned media was clarified immediately after harvest by centrifuging for 30 min, 10,000 x g at 4 °C;

then transferring the supernatant to clean tubes and centrifuging again for 1 h, 21,100 x g at 4 °C. The

supernatant was transferred to clean 50 mL conical tubes and stored at 4 °C until ready for further

purification. Clarified conditioned media was transferred to Seton 7030 tubes and each tube underlaid

with a 100 µL cushion of 50% sucrose. Tubes were centrifuged in an SW-41 Ti rotor (Beckman Coulter) for

1 h, 36,000 rpm at 4 °C, then ≥ 200 µL of volume was harvested from the bottom of each tube. In cases

where the total harvested volume exceeded 500 µL, the harvested volume was diluted to 11 mL with

sterile-filtered PBS buffer, transferred to a final Seton 7030 tube, underlaid with a 100 µL cushion of 50%

sucrose, re-centrifuged as described above, and 500 µL of volume harvested from the bottom of the tube.

Concentrated EVs were then purified by size exclusion chromatography into sterile-filtered PBS buffer

using qEV2-35 nm gravity columns (Izon) at ambient temperature (0.5 mL load volume, 2.5 mL void

volume, 1.2 mL harvest volume). Purified EVs were stored at 4 °C for up to 8 weeks, over which time no

evidence of sample degradation was observed.

Immunoblots

Protein samples were run on 4-20% or 7.5% Mini-PROTEAN TGX SDS-PAGE gels (BioRad), then transferred

to PVDF membranes using the TurboBlot semi-dry transfer system (BioRad). Blots were washed briefly 3

times with 10 mL of PBST buffer, then incubated for 1-2 h at room temperature in PBST with blocking

agent (5 mg/mL Bovine Serum Albumin (Sigma Aldrich) for anti-Strep blots, 5% nonfat dry milk for all other

blots). Blocking solution was removed and primary antibody solution in PBST with blocking agent (1:2000

rabbit anti-Strep (Abcam #76949), 1:4000 mouse anti-Flag (Millipore Sigma #F3165), 1:2500 mouse

AC133-1 (Miltenyi #130-111-756)) for 2 h at ambient temperature or for 12-72 h at 4 °C. The blots were

then washed 3 times for 5-10 min with 10 mL PBST. Secondary antibody solution (1:5000 ECL anti-Mouse

(Cytiva) or 1:10,000 ECL anti-Rabbit (Cytiva)) in PBST with appropriate blocking agent was then added to

the blots and incubated for 1 h at ambient temperature. Blots were incubated with 5 mL Western Lighting

ECL solution (PerkinElmer) for 1 min and imaged using the chemiluminescence setting on an Amersham

680 gel imager (GE Healthcare). Blots were adjusted for brightness and contrast using GIMP (GNU Project)

and annotated with Illustrator (Adobe). After blotting and imaging, images were adjusted for brightness

and contrast and subjected to digital densitometry with ImageJ[52]. Resulting measurements were reported

normalized to WT Prom1 on each blot to allow comparisons between blots.

Dynamic Light Scattering

Dynamic light scattering measurements were performed using an SZ-100 Nano Particle Analyzer (Horiba).

EVs diluted in PBS to a volume of 1 mL were transferred to a disposable plastic cuvette (Fisher) and

measurements were taken using settings for polydisperse liposomes in aqueous buffer. All measurements

were taken at 25 °C in multiple replicates to control for instrument sampling error.

Silver Stain

EV samples were run on 7.5% SDS-PAGE Tris-Glycine gels and stained with Pierce Silver Stain for Mass

Spectrometry (Thermo Scientific) according to the manufacturer's protocol.

Glycosylation Assays

PNGase F (New England Biolabs) was used to remove *N*-glycan moieties from proteins under denaturing

conditions according to the manufacturer's instructions.

Cryo-TEM Sample Preparation and Imaging

Prom1 and Ttyh1 EVs were vitrified on 300-mesh gold Quantifoil R 1.2/1.3 + 2 nm Carbon grids (Electron

Microscopy Sciences). Briefly, grids were glow discharged in an EasiGlow device (Pelco) set to 5 mA, 30 s,

0.39 mbar, with a 15 s vacuum hold time. The grids were then treated with 5 µL of purified EVs, incubated

for 60 s to allow EVs to adhere to the carbon layer, then blotted with a VitroBot Mark IV (Thermo Scientific)

set to 22 °C, 5 s blot time, +15 blot force, 100% humidity; and plunge frozen in liquid ethane. Vitrified

samples were imaged on a Titan Krios microscope (Thermo Scientific) with a Falcon 4 direct electron

detector (Thermo Scientific) using counted detection mode, 105,000 x nominal magnification, 0.83 Å pixel

size, with 49-frame fractionated collection, 49.8 e$^-$/Å^2 total dose, and defocus ranging from -0.8 to -2.0

µm in 0.1 µm increments. Images were processed and analyzed with CryoSparc v. 4.2.1 (Structura

Biotechnology). Vesicles were defined and diameter (all EVs) and roundness (only EVs completely visible

on one micrograph) were calculated using custom scripts that extend CryoSparc, made publicly available

at https://github.com/tribell4310/vesicle-quantification.

NS-TEM Sample Preparation and Imaging

Formvar carbon film 400-mesh copper grids (Electron Microscopy Sciences) were glow discharged in an

EasiGlow device (Pelco) set to 30 mA, 30 s, 0.39 mbar, with a 15 s vacuum hold time. 5 µL of EV sample

was applied to a glow-discharged grid and incubated for 60 s at room temperature. The grid was then

blotted manually with filter paper (Whatman #43), briefly washed 3 times with 20 µL of PBS buffer,

blotted, washed 2 times with deionized water, blotted, washed 1 time with 1.25% (w/v) uranyl formate

(Electron Microscopy Sciences), and blotted. The grid was then floated for 10 s on a 20 µL of drop of 1.25%

uranyl formate, blotted, and allowed to air dry. Imaging was performed on a Tecnai T12 transmission

electron microscope (FEI) equipped with an XR16 detector (AMT) operated at an accelerating voltage of

120 kV, 30,000 x nominal magnification, 4.32 Å pixel size, and -1.5 µm defocus. Pixel size was manually

calibrated prior to image acquisition using a dedicated calibration waffle grid (Ted Pella). Data collected

for EV membrane area quantification was automated using SerialEM[53]. Vesicles were defined and

analyzed in CryoSparc (Structura Biotechnology) as described above for cryo-TEM data.

Quantification of EV Membrane Area from NS-TEM Images

NS-TEM images were collected using an automated acquisition script. The images were processed to

resolve membrane features from background using a custom image processing script made publicly

available at https://github.com/tribell4310/vesicle-quant. EV membrane area was calculated for each

micrograph as the fraction of pixels in the imaging area that were occupied by membrane features.

Equilibrium Sucrose Gradient Sedimentation Analysis

Linear sucrose gradients were prepared in Seton 7030 tubes using sterile-filtered 5% and 30% (w/v)

sucrose dissolved in PBS buffer using a Gradient Station IP (BioComp). Freshly prepared gradients were

loaded into an SW-41 Ti swinging bucket rotor and 200 μL of the purified EVs were layered atop

gradients immediately prior to centrifugation. Samples were centrifuged for 5 h, 22,000 rpm, 4 °C, then

fractionated into 13 fractions of 930 μL using the Gradient Station IP. Gradient fractions were analyzed

by SDS-PAGE and immunoblotting, then quantified as described above.

Evolutionary Analysis of Prom and Ttyh proteins

We identified sequences homologous to human Prom1 using BLAST[54] and InterPro[55]. Putative prominin

sequences were curated to include only sequences with five predicted transmembrane helices by

TMHMM[56] in a 2+2+1 pattern, containing large extracellular loops (> 300 amino acids) and 2 small

intracellular loops (< 25 amino acids). This broad search revealed prominin sequences across the

eukaryotic tree ranging from metazoans to green plants. Multiple sequence alignment was performed

using MAFFT[57] and phylogenetic relationships inferred using IQ-TREE with MODELFIND for evolutionary

model selection[58,59]. Branch supports were calculated using the approximate likelihood ratio test[60]. A

smaller tree was also constructed spanning only metazoan sequences, with fungi included as an outgroup

for rooting. For direct comparison between Prom and Ttyh sequences, homologs of human Ttyh1 were

identified for all species included in the prominin metazoan tree and aligned as described above. Figures

showing multiple sequence alignments were generated using JalView[61]. Figures showing trees were

generated using IcyTree[62].

Conservation of each residue within the Prom and Ttyh metazoan trees was calculated using the

Livingstone and Barton algorithm implemented in JalView[63]. Conservation scores were then plotted onto

the AlphaFold2 structure model[41] of human Prom1 or a subunit of the solved cryo-EM structure of human

Ttyh1[32] using PyMol (Schrödinger).

Live Cell Fluorescence Microscopy and Analysis

Confluent monoclonal HeLa cells stably expressing WT Prom1-StayGold and Prom1[W795R]-StayGold

were harvested, seeded onto 35 mm glass-bottom dishes (MatTek Life Sciences) coated with poly-D-lysine

(0.1 mg/mL) and allowed to grow overnight at 37˚C under 5% CO_2. Cells were stained with wheat germ

agglutinin (WGA) conjugated with AlexaFluor 647 (W32466, Thermo Fisher Scientific) at 5 µg/mL for 10

min at 37 ˚C, then washed twice with 1x PBS. Cells were placed in Live Cell Imaging Solution (Invitrogen)

prior to imaging using a Nikon A1R HD25 point scanning confocal microscope with GaAsP and PMT

detectors, equipped with an Apo TIRF 60x/1.49 NA objective lens and Ti2 Z-drive. Temperature, humidity,

and CO_2 concentrations were controlled with a Live Cell environmental chamber (Oko-Lab). Image

acquisition was performed in NIS-Elements (Nikon Instruments Inc.) and line-scan analysis was performed

using Fiji[64].

Preparation of fluorescent cholesterol analogs

BODIPY-cholesterol was procured commercially (TopFluor cholesterol, Avanti Polar Lipids) and

resuspended at 1 mM in ethanol. AlexaFluor647-cholesterol was synthesized from Alkyne Cholesterol and

AZDye 647 Azide Plus (Click Chemistry Tools) by Copper(I)-catalyzed azide-alkyne cycloaddition. Briefly, 1

mM AZDye 647 Azide Plus and 2 mM alkyne cholesterol (each from a 10 mM stock prepared in anhydrous

DMSO) were combined with 2 mM tetrakis(acetonitrile)copper(I) tetrafluoroborate (from a 40 mM stock

prepared in ethanol) in a 350 µL reaction brought up to volume with ethanol. The reaction was incubated

at 42 °C for 30 min then at 70 °C for 2 h with the reaction vessel open to allow solvent to evaporate.

Cholesterol was extracted from the final mixture by diluting the solution to 650 µL with PBS buffer and

adding 1 mL methanol and 0.5 mL chloroform. The mixture was centrifuged for 2 min at 14,000 x *g* and

the supernatant transferred to a clean vessel, after which 1 mL of chloroform and 2 mL of glacial acetic

acid were added and mixed by vortexing. The solution was then concentrated by evaporation in a SPD1010

SpeedVac instrument (Savant) until dried, then resuspended in 100 µL ethanol. This preparation was

considered to be at 5 mM labeled cholesterol (assuming 100% yield) for downstream calculations.

Protein Expression and Purification

A plasmid encoding mTagBFP2-Strep was transformed into *E. coli* BL21 (DE3) pLysS, grown at 37 °C in LB

media to OD_{600} 0.5-0.7, induced with 0.5 mM IPTG (Gold Biotechnology), and harvested after 3 h of

expression. Cells were resuspended in 25 mL per liter of culture of Buffer A (25 mM HEPES-NaOH pH 7.5,

500 mM NaCl, 20 mM imidazole, 0.5 mM dithiothreitol (DTT)) supplemented with 10 µM leupeptin (Sigma

Aldrich), 1 µM pepstatin A (Sigma Aldrich), 1 mM phenylmethylsulfonyl fluoride (Sigma Aldrich), 1 mg/mL

chicken egg lysozyme (Fisher), and 250 U benzonase nuclease (Sigma Aldrich); and incubated with stirring

for 1 h at 4 °C. Cells were then sonicated for 3 min in an ice/water bath with 5 s on / 10 s off pulses. The

lysate was then clarified by centrifugation in a JA-25.5 rotor (Beckman Coulter) for 45 min, 15,000 rpm,

4 °C. The supernatant was then loaded onto two 5-mL HisTrap columns (Cytiva) plumbed in series

equilibrated in Buffer A using a peristaltic pump at 1.5 mL / min flow rate. The column was washed with

100 mL of Buffer A and eluted with 20 mL of Buffer B (25 mM HEPES-NaOH pH 7.5, 500 mM NaCl, 300 mM

Imidazole, 0.5 mM DTT). The protein was found to be ~95% pure by SDS-PAGE, and the concentration of

the eluted material was measured using absorbance signal at 280 nm. The sample was divided into small

aliquots, flash frozen in liquid nitrogen and stored at -80 °C.

Chol-IP Assays

Prior to running chol-IP assays, input EVs were quantified by SDS-PAGE and immunoblotting to ensure

equal inputs of Prom1-Strep and/or Ttyh1-Strep in each assay. EVs were mixed 4:1 with buffer CIA (25

mM HEPES-NaOH pH 7.8, 150 mM NaCl, 5 mM $CaCl_2$, 5% n-dodecyl-beta-maltoside (DDM) (Anatrace)) and

incubated for 1 h at 4C with end-over-end rotation, protected from ambient light. During incubation,

0.025 µL of StrepTactinXT 4Flow resin (IBA) per condition was equilibrated with buffer CIB (25 mM HEPES-

NaOH pH 7.8, 150 mM NaCl, 5 mM $CaCl_2$, 1% DDM) in a single pooled reaction. mTagBFP2-Strep was added

to the resin at a ratio of 2 fmol mTagBFP2-Strep per 1 µL of resin, and incubated for 15 min at 4 °C with

end-over-end rotation, protected from ambient light. After one additional wash with buffer CIB, the resin

was then divided equally across low-binding 1.5 mL tubes (USA Scientific) so that each condition being

tested plus one negative control condition had equal inputs of BFP-labeled resin. The DDM-treated EVs

were then added to the appropriate resin tube and incubated for 1 h at 4 °C with end-over-end rotation,

protected from ambient light. Each condition was then washed twice with buffer CIB for 5 min at 4 °C with

end-over-end rotation, resuspended in 45 µL of buffer CIB, and stored on ice, protected from light. Each

condition was sequentially pipetted with a cut pipette tip onto a glass microscope slide (Fisher) and gently

covered with an 18 mm x 18 mm no. 1 glass cover slip (Matsunami) layered on carefully to minimize

trapped air bubbles. Montaged images of the resin beads were collected using an Axio Observer TIRF

microscope (Zeiss) in epifluorescence mode with a Prime 95B camera (Photometrics) running SlideBook

v. 6.0.24 software (3i). Custom scripts were then used to identify resin beads in each image using DAPI

(BFP) signal to both identify beads and normalize fluorescence signal in the FITC (BODIPY-cholesterol) or

CY5 (AlexaFluor647-cholesterol) channels. Images of all resin particles were manually reviewed to ensure

the exclusion of air bubbles, or other non-resin fluorescent artifacts from downstream analysis. Analysis

scripts have been made publicly available at https://github.com/tribell4310/bead-assay.

Data Availability

Electron microscopy datasets and all biochemical data are deposited in a Zenodo repository at

doi:10.5281/zenodo.10034616.   Custom   software   packages   are   available   at

https://github.com/tribell4310.

Supporting Information

This article contains supporting information.

Acknowledgments

We thank D. Syau (Harvard Medical School) and members of the Chao lab for helpful feedback, and Z. Li,

S. Sterling, R. Walsh, M. Mayer, and S. Rawson from the Harvard Medical School Cryo-EM Center for

support with cryo-TEM data collection. We are grateful to Dr. Richard Bouley at the Microscopy Core of

the Program in Membrane Biology (PMB) at Massachusetts General Hospital for providing helpful advice

and equipment access for live-cell confocal microscopy. We thank the staff at the Center for Regenerative

Medicine (CRM) Flow Cytometry Core Facility for helpful support in cytometry sorting and analysis. This

work was supported by National Institutes of Health R35GM142553 (to LHC) and the Helen Hay Whitney

Foundation (to TAB). The content is solely the responsibility of the authors and does not necessarily

represent the official views of the National Institutes of Health.

References

1. Sharkova, M., Chow, E., Erickson, T., and Hocking, J.C. (2023). The morphological and functional
diversity of apical microvilli. J. Anat. *242*, 327–353. 10.1111/joa.13781.

2. Satir, P., and Christensen, S.T. (2007). Overview of structure and function of mammalian cilia. Annu.
Rev. Physiol. *69*, 377–400. 10.1146/annurev.physiol.69.040705.141236.

3. Brown, D.A., and London, E. (2000). Structure and Function of Sphingolipid- and Cholesterol-rich
Membrane Rafts*. J. Biol. Chem. *275*, 17221–17224. 10.1074/jbc.R000005200.

4. Corbeil, D., Marzesco, A.-M., Wilsch-Bräuninger, M., and Huttner, W.B. (2010). The intriguing links
between prominin-1 (CD133), cholesterol-based membrane microdomains, remodeling of apical
plasma membrane protrusions, extracellular membrane particles, and (neuro)epithelial cell
differentiation. FEBS Lett. *584*, 1659–1664. 10.1016/j.febslet.2010.01.050.

5. Corbeil, D., Karbanová, J., Fargeas, C.A., and Jászai, J. (2013). Prominin-1 (CD133): Molecular and
Cellular Features Across Species. In Prominin-1 (CD133): New Insights on Stem & Cancer Stem Cell
Biology Advances in Experimental Medicine and Biology., D. Corbeil, ed. (Springer), pp. 3–24.
10.1007/978-1-4614-5894-4_1.

6. Thamm, K., Šimaitė, D., Karbanová, J., Bermúdez, V., Reichert, D., Morgenstern, A., Bornhäuser, M.,
Huttner, W.B., Wilsch-Bräuninger, M., and Corbeil, D. (2019). Prominin-1 (CD133) modulates the
architecture and dynamics of microvilli. Traffic *20*, 39–60. 10.1111/tra.12618.

7. Yin, A.H., Miraglia, S., Zanjani, E.D., Almeida-Porada, G., Ogawa, M., Leary, A.G., Olweus, J., Kearney,
J., and Buck, D.W. (1997). AC133, a Novel Marker for Human Hematopoietic Stem and Progenitor
Cells. Blood *90*, 5002–5012. 10.1182/blood.V90.12.5002.

8. Miraglia, S., Godfrey, W., Yin, A.H., Atkins, K., Warnke, R., Holden, J.T., Bray, R.A., Waller, E.K., and
Buck, D.W. (1997). A Novel Five-Transmembrane Hematopoietic Stem Cell Antigen: Isolation,
Characterization, and Molecular Cloning. Blood *90*, 5013–5021. 10.1182/blood.V90.12.5013.

9. Corbeil, D., Röper, K., Weigmann, A., and Huttner, W.B. (1998). AC133 Hematopoietic Stem Cell
Antigen: Human Homologue of Mouse Kidney Prominin or Distinct Member of a Novel Protein
Family? Blood *91*, 2625–2626. 10.1182/blood.V91.7.2625.

10.   Kemper, K., Tol, M.J.P.M., and Medema, J.P. (2010). Mouse Tissues Express Multiple Splice
Variants of Prominin-1. PLOS ONE *5*, e12325. 10.1371/journal.pone.0012325.

11.   Mak, A.B., Nixon, A.M.L., Kittanakom, S., Stewart, J.M., Chen, G.I., Curak, J., Gingras, A.-C.,
Mazitschek, R., Neel, B.G., Stagljar, I., et al. (2012). Regulation of CD133 by HDAC6 Promotes β-

Catenin Signaling to Suppress Cancer Cell Differentiation. Cell Rep. *2*, 951–963.
10.1016/j.celrep.2012.09.016.

12.   Yang, Z., Chen, Y., Lillo, C., Chien, J., Yu, Z., Michaelides, M., Klein, M., Howes, K.A., Li, Y.,
Kaminoh, Y., et al. (2008). Mutant prominin 1 found in patients with macular degeneration disrupts
photoreceptor disk morphogenesis in mice. J. Clin. Invest. *118*, 2908–2916. 10.1172/JCI35891.

13.   Burgoyne, T., Meschede, I.P., Burden, J.J., Bailly, M., Seabra, M.C., and Futter, C.E. (2015). Rod
disc renewal occurs by evagination of the ciliary plasma membrane that makes cadherin-based
contacts with the inner segment. Proc. Natl. Acad. Sci. *112*, 15922–15927.
10.1073/pnas.1509285113.

14.   Rattner, A., Smallwood, P.M., Williams, J., Cooke, C., Savchenko, A., Lyubarsky, A., Pugh, E.N.,
and Nathans, J. (2001). A Photoreceptor-Specific Cadherin Is Essential for the Structural Integrity of
the Outer Segment and for Photoreceptor Survival. Neuron *32*, 775–786. 10.1016/S0896-
6273(01)00531-1.

15.   Zacchigna, S., Oh, H., Wilsch-Bräuninger, M., Missol-Kolka, E., Jászai, J., Jansen, S., Tanimoto, N.,
Tonagel, F., Seeliger, M., Huttner, W.B., et al. (2009). Loss of the Cholesterol-Binding Protein
Prominin-1/CD133 Causes Disk Dysmorphogenesis and Photoreceptor Degeneration. J. Neurosci. *29*,
2297–2308. 10.1523/JNEUROSCI.2034-08.2009.

16.   Eidinger, O., Leibu, R., Newman, H., Rizel, L., Perlman, I., and Ben-Yosef, T. (2015). An intronic
deletion in the PROM1 gene leads to autosomal recessive cone-rod dystrophy. Mol. Vis. *21*, 1295–
1306.

17.   Liang, J., She, X., Chen, J., Zhai, Y., Liu, Y., Zheng, K., Gong, Y., Zhu, H., Luo, X., and Sun, X. (2019).
Identification of novel PROM1 mutations responsible for autosomal recessive maculopathy with rod-
cone dystrophy. Graefes Arch. Clin. Exp. Ophthalmol. *257*, 619–628. 10.1007/s00417-018-04206-w.

18.   Maw, M.A., Corbeil, D., Koch, J., Hellwig, A., Wilson-Wheeler, J.C., Bridges, R.J.,
Kumaramanickavel, G., John, S., Nancarrow, D., Röper, K., et al. (2000). A frameshift mutation in
prominin (mouse)-like 1 causes human retinal degeneration. Hum. Mol. Genet. *9*, 27–34.
10.1093/hmg/9.1.27.

19.   Fargeas, C.A. (2013). Prominin-2 and Other Relatives of CD133. In Prominin-1 (CD133): New
Insights on Stem & Cancer Stem Cell Biology Advances in Experimental Medicine and Biology., D.
Corbeil, ed. (Springer), pp. 25–40. 10.1007/978-1-4614-5894-4_2.

20.   Hori, A., Nishide, K., Yasukuni, Y., Haga, K., Kakuta, W., Ishikawa, Y., Hayes, M.J., Ohnuma, S.,
Kiyonari, H., Kimura, K., et al. (2019). Prominin-1 Modulates Rho/ROCK-Mediated Membrane
Morphology and Calcium-Dependent Intracellular Chloride Flux. Sci. Rep. *9*, 15911. 10.1038/s41598-
019-52040-9.

21.   Röper, K., Corbeil, D., and Huttner, W.B. (2000). Retention of prominin in microvilli reveals
distinct cholesterol-based lipid micro-domains in the apical plasma membrane. Nat. Cell Biol. *2*, 582–
592. 10.1038/35023524.

22.   Dubreuil, V., Marzesco, A.-M., Corbeil, D., Huttner, W.B., and Wilsch-Bräuninger, M. (2007).
Midbody and primary cilium of neural progenitors release extracellular membrane particles enriched
in the stem cell marker prominin-1. J. Cell Biol. *176*, 483–495. 10.1083/jcb.200608137.

23.   Marzesco, A.-M., Janich, P., Wilsch-Bräuninger, M., Dubreuil, V., Langenfeld, K., Corbeil, D., and
Huttner, W.B. (2005). Release of extracellular membrane particles carrying the stem cell marker
prominin-1 (CD133) from neural progenitors and other epithelial cells. J. Cell Sci. *118*, 2849–2858.
10.1242/jcs.02439.

24.   Wood, C.R., and Rosenbaum, J.L. (2015). Ciliary ectosomes: transmissions from the cell's
antenna. Trends Cell Biol. *25*, 276–285. 10.1016/j.tcb.2014.12.008.

25.   Hurbain, I., Macé, A.-S., Romao, M., Prince, E., Sengmanivong, L., Ruel, L., Basto, R., Thérond,
P.P., Raposo, G., and D'Angelo, G. (2022). Microvilli-derived extracellular vesicles carry Hedgehog
morphogenic signals for Drosophila wing imaginal disc development. Curr. Biol. *32*, 361-373.e6.
10.1016/j.cub.2021.11.023.

26.   Campbell, H.D., Kamei, M., Claudianos, C., Woollatt, E., Sutherland, G.R., Suzuki, Y., Hida, M.,
Sugano, S., and Young, I.G. (2000). Human and Mouse Homologues of the Drosophila melanogaster
tweety (tty) Gene: A Novel Gene Family Encoding Predicted Transmembrane Proteins. Genomics *68*,
89–92. 10.1006/geno.2000.6259.

27.   Matthews, C.A., Shaw, J.E., Hooper, J.A., Young, I.G., Crouch, M.F., and Campbell, H.D. (2007).
Expression and evolution of the mammalian brain gene Ttyh1. J. Neurochem. *100*, 693–707.
10.1111/j.1471-4159.2006.04237.x.

28.   Halleran, A.D., Sehdev, M., Rabe, B.A., Huyck, R.W., Williams, C.C., and Saha, M.S. (2015).
Characterization of tweety gene (ttyh1-3) expression in Xenopus laevis during embryonic
development. Gene Expr. Patterns *17*, 38–44. 10.1016/j.gep.2014.12.002.

29.   Bae, Y., Kim, A., Cho, C.-H., Kim, D., Jung, H.-G., Kim, S.-S., Yoo, J., Park, J.-Y., and Hwang, E.M.
(2019). TTYH1 and TTYH2 Serve as LRRC8A-Independent Volume-Regulated Anion Channels in Cancer
Cells. Cells *8*, 562. 10.3390/cells8060562.

30.   Han, Y.-E., Kwon, J., Won, J., An, H., Jang, M.W., Woo, J., Lee, J.S., Park, M.G., Yoon, B.-E., Lee,
S.E., et al. (2019). Tweety-homolog (Ttyh) Family Encodes the Pore-forming Subunits of the Swelling-
dependent Volume-regulated Anion Channel (VRACswell) in the Brain. Exp. Neurobiol. *28*, 183–215.
10.5607/en.2019.28.2.183.

31.   Li, B., Hoel, C.M., and Brohawn, S.G. (2021). Structures of Tweety Homolog Proteins TTYH2 and
TTYH3 reveal a Ca2+-dependent switch from intra- to inter-membrane dimerization
10.1101/2021.08.15.456437.

32.   Sukalskaia, A., Straub, M.S., Deneka, D., Sawicka, M., and Dutzler, R. (2021). Cryo-EM structures
of the TTYH family reveal a novel architecture for lipid interactions. Nat. Commun. *12*, 4893.
10.1038/s41467-021-25106-4.

33.   Melvin, E., Shlush, E., Giladi, M., and Haitin, Y. (2021). TTYH family members form tetrameric
complexes within the cell membrane. Preprint at bioRxiv, 10.1101/2021.09.05.459021
10.1101/2021.09.05.459021.

34.   Jung, E., Osswald, M., Blaes, J., Wiestler, B., Sahm, F., Schmenger, T., Solecki, G., Deumelandt, K.,
Kurz, F.T., Xie, R., et al. (2017). Tweety-Homolog 1 Drives Brain Colonization of Gliomas. J. Neurosci.
37, 6837–6850. 10.1523/JNEUROSCI.3532-16.2017.

35.   Stefaniuk, M., Swiech, L., Dzwonek, J., and Lukasiuk, K. (2010). Expression of Ttyh1, a member of
the Tweety family in neurons in vitro and in vivo and its potential role in brain pathology. J.
Neurochem. *115*, 1183–1194. 10.1111/j.1471-4159.2010.07023.x.

36.   Wiernasz, E., Kaliszewska, A., Brutkowski, W., Bednarczyk, J., Gorniak, M., Kaza, B., and Lukasiuk,
K. (2014). Ttyh1 Protein is Expressed in Glia In Vitro and Shows Elevated Expression in Activated
Astrocytes Following Status Epilepticus. Neurochem. Res. *39*, 2516–2526. 10.1007/s11064-014-1455-
3.

37.   Kemper, K., Sprick, M.R., Bree, M. de, Scopelliti, A., Vermeulen, L., Hoek, M., Zeilstra, J., Pals,
S.T., Mehmet, H., Stassi, G., et al. (2010). The AC133 Epitope, but not the CD133 Protein, Is Lost upon
Cancer Stem Cell Differentiation. Cancer Res. *70*, 719–729. 10.1158/0008-5472.CAN-09-1820.

38.   Huttner, W.B., Schiebler, W., Greengard, P., and De Camilli, P. (1983). Synapsin I (protein I), a
nerve terminal-specific phosphoprotein. III. Its association with synaptic vesicles studied in a highly
purified synaptic vesicle preparation. J. Cell Biol. *96*, 1374–1388. 10.1083/jcb.96.5.1374.

39.   Fantini, J., and Barrantes, F.J. (2013). How cholesterol interacts with membrane proteins: an
exploration of cholesterol-binding sites including CRAC, CARC, and tilted domains. Front. Physiol. *4*.
10.3389/fphys.2013.00031.

40.   Boulanger-Scemama, E., El Shamieh, S., Démontant, V., Condroyer, C., Antonio, A., Michiels, C.,
Boyard, F., Saraiva, J.-P., Letexier, M., Souied, E., et al. (2015). Next-generation sequencing applied to
a large French cone and cone-rod dystrophy cohort: mutation spectrum and new genotype-
phenotype correlation. Orphanet J. Rare Dis. *10*, 85. 10.1186/s13023-015-0300-3.

41.   Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K.,
Bates, R., Žídek, A., Potapenko, A., et al. (2021). Highly accurate protein structure prediction with
AlphaFold. Nature *596*, 583–589. 10.1038/s41586-021-03819-2.

42.   Derganc, J., and Čopič, A. (2016). Membrane bending by protein crowding is affected by protein
lateral confinement. Biochim. Biophys. Acta BBA - Biomembr. *1858*, 1152–1159.
10.1016/j.bbamem.2016.03.009.

43.   Zidovetzki, R., and Levitan, I. (2007). Use of cyclodextrins to manipulate plasma membrane
cholesterol content: Evidence, misconceptions and control strategies. Biochim. Biophys. Acta BBA -
Biomembr. *1768*, 1311–1324. 10.1016/j.bbamem.2007.03.026.

44.   Salinas, R.Y., Pearring, J.N., Ding, J.-D., Spencer, W.J., Hao, Y., and Arshavsky, V.Y. (2017).
Photoreceptor discs form through peripherin-dependent suppression of ciliary ectosome release. J.
Cell Biol. *216*, 1489–1499. 10.1083/jcb.201608081.

45.   Lorent, J.H., Levental, K.R., Ganesan, L., Rivera-Longsworth, G., Sezgin, E., Doktorova, M., Lyman,
E., and Levental, I. (2020). Plasma membranes are asymmetric in lipid unsaturation, packing and
protein shape. Nat. Chem. Biol. *16*, 644–652. 10.1038/s41589-020-0529-6.

46.   Kanazawa, T., Takematsu, H., Yamamoto, A., Yamamoto, H., and Kozutsumi, Y. (2008). Wheat
germ agglutinin stains dispersed post-golgi vesicles after treatment with the cytokinesis inhibitor
psychosine. J. Cell. Physiol. *215*, 517–525. 10.1002/jcp.21328.

47.   Ridsdale, A., Denis, M., Gougeon, P.-Y., Ngsee, J.K., Presley, J.F., and Zha, X. (2006). Cholesterol
Is Required for Efficient Endoplasmic Reticulum-to-Golgi Transport of Secretory Membrane Proteins.
Mol. Biol. Cell *17*, 1593–1605. 10.1091/mbc.e05-02-0100.

48.   Stüven, E., Porat, A., Shimron, F., Fass, E., Kaloyanova, D., Brügger, B., Wieland, F.T., Elazar, Z.,
and Helms, J.B. (2003). Intra-Golgi Protein Transport Depends on a Cholesterol Balance in the Lipid
Membrane*. J. Biol. Chem. *278*, 53112–53122. 10.1074/jbc.M300402200.

49.   Fargeas, C.A., Huttner, W.B., and Corbeil, D. (2007). Nomenclature of prominin-1 (CD133) splice
variants – an update. Tissue Antigens *69*, 602–606. 10.1111/j.1399-0039.2007.00825.x.

50.   Elegheert, J., Behiels, E., Bishop, B., Scott, S., Woolley, R.E., Griffiths, S.C., Byrne, E.F.X., Chang,
V.T., Stuart, D.I., Jones, E.Y., et al. (2018). Lentiviral transduction of mammalian cells for fast, scalable
and high-level production of soluble and membrane proteins. Nat. Protoc. *13*, 2991–3017.
10.1038/s41596-018-0075-9.

51.   Kutner, R.H., Zhang, X.-Y., and Reiser, J. (2009). Production, concentration and titration of
pseudotyped HIV-1-based lentiviral vectors. Nat. Protoc. *4*, 495–505. 10.1038/nprot.2009.22.

52.   Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image
analysis. Nat. Methods *9*, 671–675. 10.1038/nmeth.2089.

53.   Mastronarde, D.N. (2005). Automated electron microscope tomography using robust prediction
of specimen movements. J. Struct. Biol. *152*, 36–51. 10.1016/j.jsb.2005.07.007.

54.   Sayers, E.W., Bolton, E.E., Brister, J.R., Canese, K., Chan, J., Comeau, D.C., Connor, R., Funk, K.,
Kelly, C., Kim, S., et al. (2022). Database resources of the national center for biotechnology
information. Nucleic Acids Res. *50*, D20–D26. 10.1093/nar/gkab1112.

55.   Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B.L., Salazar, G.A., Bileschi, M.L.,
Bork, P., Bridge, A., Colwell, L., et al. (2023). InterPro in 2022. Nucleic Acids Res. *51*, D418–D427.
10.1093/nar/gkac993.

56.   Sonnhammer, E.L., von Heijne, G., and Krogh, A. (1998). A hidden Markov model for predicting
transmembrane helices in protein sequences. Proc. Int. Conf. Intell. Syst. Mol. Biol. *6*, 175–182.

57.   Katoh, K., Misawa, K., Kuma, K., and Miyata, T. (2002). MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Res. *30*, 3059–3066.
10.1093/nar/gkf436.

58.   Minh, B.Q., Schmidt, H.A., Chernomor, O., Schrempf, D., Woodhams, M.D., von Haeseler, A., and
901   Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the
902   Genomic Era. Mol. Biol. Evol. *37*, 1530–1534. 10.1093/molbev/msaa015.

903   59.   Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., and Jermiin, L.S. (2017).
904   ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods *14*, 587–589.
905   10.1038/nmeth.4285.

906   60.   Anisimova, M., and Gascuel, O. (2006). Approximate Likelihood-Ratio Test for Branches: A Fast,
907   Accurate, and Powerful Alternative. Syst. Biol. *55*, 539–552. 10.1080/10635150600755453.

908   61.   Clamp, M., Cuff, J., Searle, S.M., and Barton, G.J. (2004). The Jalview Java alignment editor.
909   Bioinformatics *20*, 426–427. 10.1093/bioinformatics/btg430.

910   62.   Vaughan, T.G. (2017). IcyTree: rapid browser-based visualization for phylogenetic trees and
911   networks. Bioinformatics *33*, 2392–2394. 10.1093/bioinformatics/btx155.

912   63.   Livingstone, C.D., and Barton, G.J. (1993). Protein sequence alignments: a strategy for the
913   hierarchical analysis of residue conservation. Bioinformatics *9*, 745–756.
914   10.1093/bioinformatics/9.6.745.

915   64.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
916   Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image
917   analysis. Nat. Methods *9*, 676–682. 10.1038/nmeth.2019.