Interleukin-17 regulates epidermal remodeling via keratinocyte cytonemes

Yi Wang[1], Qingan He[1], Oliver Has[1], Kiarash Forouzesh[1], Dae Seok Eom[1,2*]

[1]Department of Developmental and Cell Biology, The University of California, Irvine, CA 92617, USA, [2]Center for Complex Biological Systems, University of California, Irvine, CA 92697, USA

*Corresponding author: dseom@uci.edu

Abstract:
The skin, the largest organ in our body, serves as our primary defense mechanism. Epidermal stem cells supply undifferentiated keratinocytes, which undergo differentiation as they migrate towards the outermost skin layer. This replenishment mechanism is conserved in vertebrates and various human skin diseases exhibit disruption of this process. However, its underlying cellular and molecular mechanisms remain elusive. Through high-resolution live imaging and *in vivo* manipulations, we unveil that cytoneme-mediated Notch signaling between keratinocytes is essential for their differentiation and proliferation. We show that IL-17, associated with human skin diseases like *psoriasis*, cell autonomously regulate keratinocyte cytonemes. Our findings highlight the significance of keratinocyte communication via cytonemes in epidermal remodeling and maintenance. Moreover, this could shed light on potential therapeutic targets for treating human skin diseases.

Introduction:
The vertebrate skin is a multilayered structure comprised of three main layers: the epidermis, the dermis, and the hypodermis (Akat et al., 2022). Each layer has distinct anatomical characteristics and serves specific functions. Of these layers, the epidermis, as the outermost layer, plays a pivotal role as the body's first line of defense against environmental threats such as pathogens. It also plays a crucial role in regulating body temperature and preventing water loss into the surrounding environment. The epidermis consists of various layers, including the stratum basale, which houses a single layer of stem cells, and several layers of differentiated cells, including the stratum spinosum, stratum granulosum, and stratum lucidum. The outermost layer, the stratum corneum, primarily comprises dead keratinocytes that are continuously shed and replenished by underlying cells differentiated from epidermal stem cells in mammals (Segre, 2006).
In zebrafish, the epidermis shares a similar composition to that of mammals. However, unlike mammals, aquatic animals lack a cornified layer and instead rely on mucous layer to protect the epidermis in the aquatic environment. The basal layer of zebrafish, equivalent to mammalian stratum basale, consists of stem cells that provide keratinocytes for replenishment during epidermal remodeling or wound healing. The periderm, comparable to the mammalian stratum granulosum and lucidum, is where fully differentiated keratinocytes are found. Similar to mammalian stratum granulosum, an intermediate layer is present between the basal layer and the periderm, housing undifferentiated keratinocytes (Chang and Hwang, 2011). This intermediate layer is

formed during metamorphosis and persists throughout adulthood in zebrafish (Lee et al., 2014).

The maintenance of a healthy epidermal barrier requires on the intricately controlled proliferation and differentiation of keratinocytes, which are supplied from underlying basal stem cells (Blanpain and Fuchs, 2009). As keratinocytes migrate apically from the stem cell layer, they undergo differentiation while concurrently exhibiting proliferation within the intermediate layer. Ultimately, fully differentiated keratinocytes replenish the periderm. This replenishment mechanism is well-conserved in both zebrafish and humans. Its mis-regulation constitutes a pivotal factor in the etiology of various human skin diseases, including *psoriasis*, *atomic dermatitis*, and others (Chang and Hwang, 2011; Lowes et al., 2014; Nowell and Radtke, 2013; Schempp et al., 2009; Zhou et al., 2022). Studies have also revealed that maintaining a healthy epidermis requires complex intercellular communications between keratinocytes and the immune system, adding complexity to our efforts to comprehensively understand the mechanism.

One of the major signaling pathways critical for keratinocyte differentiation and proliferation is the Notch pathway. Studies have demonstrated that Notch receptors and ligands exhibit spatially restricted expression patterns in layers in epidermis (Nowell and Radtke, 2013). Notch 1, 2, and 3 are notably prevalent in the undifferentiated keratinocytes in mice (Blanpain et al., 2006; Nickoloff et al., 2002; Thelu et al., 2002). Dysregulated Notch signaling has been linked to abnormal keratinocyte differentiation and hyperproliferation, which are one of the hallmarks of various skin diseases including *psoriasis* and *atopic dermatitis* (Gratton et al., 2020), underscoring the importance of proper Notch signaling in epidermal maintenance to retain the balance between keratinocyte differentiation and proliferation (Ota et al., 2014; Rangarajan et al., 2001). Nonetheless, the precise mechanisms underlying the initiation of Notch signaling in undifferentiated keratinocytes are not yet understood.

Furthermore, various combinations of cytokines produced by different immune cells under inflammatory conditions can significantly contribute to the onset or progression of these skin diseases. Among these, interleukin-17 (IL-17) is a well-characterized cytokine closely associated with the symptoms of these conditions. The overproduction of IL-17 by T helper (Th)17 cells and other immune cells is a primary driver of certain skin diseases, such as *psoriasis*. Biologic treatments that inhibit IL-17 have proven to effectively alleviating the symptoms of these conditions (Mosca et al., 2021). Unfortunately, discontinuing these treatments often lead to symptom recurrence. Also known side effects associated with these inhibitors include headaches, nasopharyngitis, and infections (Campa et al., 2016). Consequently, a definitive cure remains elusive currently (Fragoulis et al., 2016; Lowes et al., 2014).

Thus, IL-17 is one of the key molecules that contribute to the development of *psoriasis* and other skin diseases. The IL-17 pathway exerts its effects on keratinocytes, endothelial cells, and immune cells resulting in the production of additional cytokines that promote positive feedback loops, sustaining the inflammatory state (Mosca et al., 2021). However, the underlying cellular and molecular mechanisms of this process are not yet fully understood. Therefore, gaining a comprehensive understanding of the interplay between keratinocytes, the Notch pathway, IL-17, and other players is essential for advancing the treatment of these diseases.

Intercellular signaling plays a critical role in skin development and maintenance. In recent years, a growing body of evidence has demonstrated that cells can establish communication through long, thin cellular protrusions extended by cells involved in either sending or receiving signals. These protrusions can be categorized based on their cytoskeletal composition, mode of signal delivery, morphology, and other characteristics. Examples include cytonemes, airinemes, tunneling nanotubes, intercellular bridges, migrasomes, exophers, and more (Eom, 2020; Kornberg and Roy, 2014; Ma et al., 2015; Melentijevic et al., 2017; Zhang and Scholpp, 2019). These specialized cellular protrusions have been observed in diverse cell types and species, ranging from fruit flies and sea urchins to zebrafish and cultured mammalian cells, with their signaling roles experimentally confirmed *in vivo* (Daly et al., 2022; Eom, 2020; Kornberg and Roy, 2014; Zhang and Scholpp, 2019) .

These specialized protrusions are significantly longer than typical filopodia and establish direct contact with target cells and serve as highways for transporting signaling molecules for major signaling pathways such as Hedgehog, Wnt, TGFβ, FGF, Notch in many *in vivo* and *in vitro* contexts (Daly et al., 2022). These signaling cellular protrusions temporally exist and have thin actin or actin/tubulin-based filaments. Consequently, live-imaging is considered one of the most effective methods for observing these structure. For that reason, zebrafish has emerged as an excellent model system for studying these cellular protrusions. The transparent nature of their early developing embryos makes zebrafish particularly well-suited for this purpose. Moreover, zebrafish are vertebrates that share common mechanisms with mammals in terms of epidermal remodeling and maintenance (Eisenhoffer et al., 2017; Li et al., 2011; Martinez-Navarro et al., 2019).

In this study, we show evidence that cytonemes extended by fully differentiated keratinocytes play a crucial role in activating Notch signaling in undifferentiated keratinocytes, thus contributing to the maintenance of epidermal homeostasis and remodeling in zebrafish. Furthermore, we demonstrate that IL-17, a key player in the pathogenesis of human skin diseases, and *clint1*, a gene required for epidermal homeostasis, regulate keratinocyte cytonemes (Dodd et al., 2009; Sahlen et al., 2021). Collectively, our findings shed light on the mechanisms that govern the regulation of skin replenishment through keratinocyte cytonemes, providing a novel perspective on understanding how skin diseases can originate from keratinocytes autonomously through the mediation of cellular protrusions.

Differentiated keratinocytes extend cytoneme-like cellular protrusions
In previous studies, we have reported a unique type of signaling cellular protrusions known as 'airinemes' and have investigated their signaling roles in pigment cells of zebrafish (Bowman et al., 2023; Eom, 2020; Eom et al., 2015; Eom and Parichy, 2017; Park et al., 2022). To extend our understanding of airinemes in other cell types, we employed random cell labelling and identified several cell types that extend airineme-like protrusions featured by highly curved filaments and large vesicles at their tips (Eom, 2020; Eom et al., 2015; Lee et al., 2014). Fully differentiated keratinocytes, marked by *krt4*, were among the cell types displaying these airineme-like protrusions (Lee et al., 2014) (Fig. 1A, arrowhead). These keratinocytes intriguingly also extend cytoneme-like

protrusions, characterized by relatively straight filaments and a lack of vesicle at the tips (Eom, 2020; Kornberg and Roy, 2014; Zhang and Scholpp, 2019) (Figure 1A, B, arrowheads). We observed that while airineme-like protrusions are rare, cytoneme-like are more frequently observed in these keratinocytes (Fig. 1C). Consequently, we conducted a more in-depth characterization of these cytoneme-like protrusions. It is note that cytonemes primarily consist of actin filaments, whereas airinemes are composed of both actin and tubulin-based structures (Eom, 2020). To affirm their nature, we utilized an actin marker, *LifeAct-mRuby*, which clearly labeled these cytoneme-like protrusions while showing no co-localization with *tubulin-mCherry* (Fig. 1D, E). Additionally, cytoneme-like protrusions were significantly reduced with actin inhibitor treatment (fig. S1). Therefore, based on their cytoskeletal composition and morphology, we classified them as cytonemes.

Keratinocytes marked by *krt4* expression extend cytonemes with the highest frequency during post-embryonic stages, specifically between 6.5 to 7.5 SSL (Standardized Standard Length (Parichy et al., 2009)) (Fig. 1C). It is worth noting that embryonic keratinocytes are replaced by post-embryonic cells around these stages, suggesting the potential role of cytonemes in epidermal remodeling and maintenance (Lee et al., 2014; Parichy et al., 2009). These keratinocyte cytonemes exhibit an approximate speed of 2.98µm/min during extension and retract at 1.9µm/min, with an average length of 18.21µm (Fig. 1F, G).

Keratinocyte cytonemes establish physical contact with underlying undifferentiated keratinocytes

Similar to mammals, the zebrafish epidermis consists of three layers. We specifically visualized the periderm, which is the outermost layer and consists of *krt4+* fully differentiated keratinocytes. The intermediate layer, where undifferentiated keratinocytes reside, was marked with *krtt1c19e* promoter driving tdTomato expression, pseudo-colored in magenta (Chang and Hwang, 2011; Lee et al., 2014). The basal layer was also labelled with *krtt1c19e* but exhibited a typical polygonal morphology (Fig. 2A-A"). This allowed us to distinguish all three epidermal layers using the *Tg(krt4:palmEGFP; krtt1c19e:tdTomato)*.

In addition to *krt4+* keratinocytes, we explored whether undifferentiated keratinocytes (*krtt1c19e+*) also extend cytonemes or other types of cellular protrusions during metamorphic stages. However, we observed that cytoneme-like protrusions from undifferentiated keratinocytes were significantly fewer in number compared to those from fully differentiated keratinocytes, and we did not observe cytoneme-like protrusions from basal stem cells (Fig. 2B). In this study, our focus was primarily on cytonemes from *krt4+* differentiated keratinocytes.

To investigate the role of keratinocyte cytonemes, we first asked what the target cells of those cytonemes are. High resolution confocal imaging revealed that cytonemes originating from peridermal keratinocytes (*krt4+*) establish physical contact with underlying undifferentiated keratinocytes (*krtt1c19e+*) in the intermediate layer. Cross-sectional views clearly showed that the tips of these cytonemes halt at the intermediate layer, with no observed instances reaching the basal layer (Fig. 2C, red dashed circles).

Cytoneme-mediated signaling regulates keratinocyte differentiation and proliferation

The observations that cytonemes most often seen during epidermal remodeling and their target cells are undifferentiated keratinocytes led us to hypothesize that cytoneme-mediated signaling play a pivotal role in regulating keratinocyte differentiation and proliferation.

To test, we generated a transgenic line capable of temporally expressing a dominant negative form of cdc42 (cdc42DN) in peridermal keratinocytes, a strategy frequently used to inhibit cytoneme extension in various contexts (Daly et al., 2022; Zhang and Scholpp, 2019). We implemented a dually inducible TetON system in the transgene, denoted as *Tg(krt4:TetGBDTRE-v2a-cdc42DN),* and immersed the transgenic and control fish in water containing inducer drugs, doxycycline and dexamethasone, for temporal cdc42DN transgene induction (Eom et al., 2015; Knopf et al., 2010). To minimize potential off-target effects of cdc42 manipulation, we determined the drug concentrations at which normal filopodial or lamellipodial activities were maintained while inhibiting cytoneme extensions (Movies S1). We employed two control groups in this experiment: one consisting of a transgenic line with no drugs and the other comprising non-transgenic fish treated with the same drug concentrations as the experimental group. Under these conditions, we observed a significant reduction in the frequency of cytoneme extension in the cdc42DN-expressing transgenic line within the drug treated group (Fig. 3A).

In contrast to the two control groups, the peridermal layer in the cytoneme-inhibited group displayed severe disorganization, with areas devoid of *krt4+* keratinocytes (Fig. 3B", arrowhead). Furthermore, the expression of the undifferentiated keratinocyte marker, *krtt1c19e* (magenta), was dramatically increased in the periderm when cytoneme extension was inhibited (Fig. 3B-B", C). Additionally, we found a substantial increase in the number of keratinocytes within the peridermal layer in the cytoneme-inhibited group (Fig. 3D). However, it is important to note that cytoneme inhibition did not affect the cell death rate or the number of goblet cells (Fig. 3E, 2A', fig. S2A). These findings strongly suggest that cytoneme-mediated signaling controls keratinocyte differentiation and proliferation.

Cytonemes activate Notch in undifferentiated keratinocytes

Previous studies have well established the crucial role of Notch signaling in controlling keratinocyte differentiation and proliferation. However, the mechanisms triggering Notch signaling in keratinocytes have remained obscure (Moriyama et al., 2008; Nguyen et al., 2006; Rangarajan et al., 2001). To investigate whether the target cells of cytonemes, specifically undifferentiated keratinocytes in the intermediate layer, are responsive to Notch signaling, we crossed Notch signal reporter line, *Tg(Tp1:H2B-mCherry),* with *Tg(krt4:palmEGFP; krtt1c19e:lynTdtomato)* to label all three epidermal layers (Parsons et al., 2009). We observed that only undifferentiated keratinocytes in the intermediate layer exhibit Notch responsiveness, whereas keratinocytes in the periderm or basal layer do not (Fig. 4A, arrowheads, fig. S3).

Subsequently, we tested whether Notch activation in undifferentiated keratinocytes is dependent on cytonemes. First, we validated the *Tp1* Notch reporter line, *Tg(Tp1:EGFP)* by measuring cytoplasmic EGFP signal before and after Notch inhibitor,

LY411575 treatment, which demonstrated a significant decrease in Notch signaling, as expected (fig. S4A, B). Next, we measured the difference in *Tp1* intensity before and after cytoneme inhibition. Notably, we observed a significant reduction in Notch signal under conditions where cytonemes were compromised (Fig. 4B, fig. S4C). We also found more abundant Notch activated keratinocytes during metamorphic stages (fig. S4D, Fig. 1C). These suggest cytonemes are responsible for Notch activation in undifferentiated keratinocytes.

We then asked why Notch activation is dependent on cytonemes since peridermal keratinocytes and underlying intermediate keratinocytes are in constant contact with each other. We revealed that intermediate keratinocytes exclusively express *lfng* (lunatic fringe), which potentially act as a default inhibitor of Notch activation in these cells (Dale et al., 2003; Panin et al., 1997) (fig. S5A, S3). However, this needs further studies to understand how cytonemes overcome this Notch signaling barrier or potentiate the signal.

Next, we hypothesized that if Notch activation by cytonemes is indeed essential for keratinocyte differentiation and proliferation, then direct inhibition of Notch signal in undifferentiated keratinocytes should yield phenotypes akin to those observed when cytonemes are inhibited (Gratton et al., 2020). To investigate this, we generated a transgenic line capable of temporally expressing dominant negative form of Suppressor of Hairless (SuHDN) in the undifferentiated keratinocytes, *Tg(krtt1c19e:TetGBDTRE-v2a-SuHDN)* (Eom et al., 2015). We observed that Notch inhibition in the undifferentiated keratinocytes resulted in a disorganized periderm, abnormally increased expression of undifferentiated keratinocyte marker, and hyperproliferation of keratinocytes in the periderm (Fig. 4C-C", D, E, F, fig. S4C). These outcomes closely mirrored the phenotypes observed when we inhibited cytoneme extension in peridermal keratinocytes (Fig. 3B-B", C, D, fig. S4C). These observations suggest that Notch signaling, activated by cytonemes in undifferentiated keratinocytes, plays a critical role in epidermal remodeling and maintenance.

It has been documented that both hypo- or hyper-activation of Notch signaling can lead to the onset of skin diseases (Abdou et al., 2012; Gratton et al., 2020). Thus, we further investigated whether gain-of-function experiments for Notch also disrupt epidermal homeostasis. We overexpressed the Notch IntraCellular Domain (NICD) in undifferentiated keratinocytes, using the *Tg(krtt1c19e:nVenus-v2a-NICD)*. Consistent with previous studies in mammals and the cytoneme or Notch loss-of-function experiments, we observed disorganized periderm, abnormal keratinocyte differentiation and hyper-proliferation (fig. S5B-B', C, D). These results imply that uncontrolled Notch signaling in keratinocytes could lead to disruption of epidermal remodeling and maintenance in both zebrafish and mammals.

Interleukin-17 regulates keratinocyte cytonemes

The aberrant differentiation and hyperproliferation of keratinocytes observed when cytoneme extension was inhibited in peridermal keratinocytes (*krt4+*) or Notch signaling was inhibited in undifferentiated keratinocytes (*krtt1c19e+*) are hallmark features of many human skin diseases such as *psoriasis*, *atopic dermatitis* and more (Armstrong and Read, 2020; Gratton et al., 2020) (Fig. 3, 4). Also, it has been well established that impaired Notch signaling is associated with some of these human skin disorders

(Gratton et al., 2020; Nowell and Radtke, 2013). Thus, we asked which key molecules are critical for the onset or progression of these skin diseases and whether they have an impact on keratinocyte cytoneme signaling.

One of them is interleukin-17 (IL-17) and we hypothesized IL-17 signaling could influence cytoneme extension in peridermal keratinocytes. Given that IL-17 is a cytokine released into the extracellular space, and to test its role in peridermal keratinocytes, we opted to overexpress the IL-17 receptors. It is note that interleukin-17 receptors and ligands are evolutionarily conserved across chordates (Gonzalez-Fernandez et al., 2020; Wu et al., 2011).

To test, we generated transgenic lines that cell-autonomously overexpressed il17 receptor D (*il17rd*) or il17 receptor A1a (*il17ra1a*) in fully differentiated peridermal keratinocytes (*krt4+*), *Tg(krt4:nVenus-v2a-il17rd)* and *Tg(krt4:il17ra1a-v2a-mCherry)*. Both of these IL-17 receptor overexpressing transgenic lines exhibited a significant reduction in cytoneme extension, resulting in a decreased Notch activation in undifferentiated keratinocytes (Fig. 5A, B). Moreover, similar to the two previous manipulations, we observed a severely disorganized periderm, an upregulation of the undifferentiated keratinocytes marker expression, and hyperproliferation of keratinocytes within the periderm (Fig. 5C-C', D, E, fig. S3).

To assess whether IL-17 signaling regulates keratinocyte differentiation and proliferation through cytonemes, we crossed two transgenic lines: *Tg(krt4:TetGBDTRE-v2a-cdc42DN)* and *Tg(krt4:nVenus-v2a-il17rd)*. If cytoneme extension and IL-17 signaling are part of the same pathway, we would expect to observe a more severe inhibition of cytoneme extension when these manipulations are combined compared to individual manipulations. Indeed, when inhibited cytoneme extension with the cdc42DN along with *il17rd* overexpression, we observed a more significant inhibition of cytoneme extension than when manipulated individually (Fig. 5F).

Next, we sought to understand how the IL-17 signal impacts cytoneme extension. Studies have shown that IL-17 signaling can regulate actin cytoskeleton in keratinocytes (Borowczyk et al., 2020; Das et al., 2019). To evaluate whether *il17rd* overexpression alters the actin cytoskeleton, thus cytonemes, we compared keratinocyte microridges, which are actin cytoskeleton-rich structures, between control and il17rd-overexpressed peridermal keratinocytes (Pinto et al., 2019; van Loon et al., 2020). Our observation revealed severe disorganization of microridges in the *il17rd* overexpressed keratinocytes, coupled with a significant up- or down-regulation of *RhoA* and *Cdc42* GTPases expression in these cells (Fig. 5G, H). These data suggest that IL-17 signaling negatively regulates cytoneme extension by unbalancing small GTPases that are essential in control of actin cytoskeleton (Daly et al., 2022; Hall, 1998; Jaffe and Hall, 2005; Stanganello et al., 2015).

Our data collectively support the idea that cytoneme-mediated intercellular signaling between peridermal and underlying undifferentiated keratinocytes is essential for epidermal remodeling and maintenance. IL-17 functions as a regulator capable of affecting cytoneme extension by modulating the actin cytoskeleton in keratinocytes. Consequently, our data suggest that the overproduction of IL-17 ligands by immune cells in conditions such as psoriatic or other skin diseases leads to epidermal imbalance through altered cytoneme signaling in keratinocytes.

In addition, recent human Genome-Wide Association Studies (GWAS) have identified a close association between *clint1* and individuals with *psoriasis* and *atopic dermatitis* (Sahlen et al., 2021). It has also been reported that *clint1* is important for epidermal homeostasis in zebrafish (Dodd et al., 2009). Given these reported findings, we investigated to determine whether *clint1* plays a role in epidermal maintenance through its influence on keratinocyte cytonemes. Indeed, we found that *clint1* mutants exhibited a significant decrease in cytoneme extension in peridermal keratinocytes, accompanied by reduced Notch signaling in undifferentiated keratinocytes. As a result, these mutant fish displayed abnormal keratinocyte differentiation and hyperproliferation of keratinocytes (fig. S6). These findings collectively suggest that *clint1* is yet another regulator of cytoneme-mediated intercellular signaling in keratinocytes.

In conclusion, the intercellular signaling facilitated by cytonemes between fully differentiated and undifferentiated keratinocytes is crucial for the maintenance and remodeling of the epidermis. Our study implies that the patients with human skin diseases may have defective cytoneme signaling between keratinocytes. However, it is important to note that further verification of the existence of cytonemes in humans and their functional validation is needed in the future.

Discussion
It is intriguing to observe that keratinocytes have the ability to extend both cytoneme and airineme-like protrusions. While these protrusions differ in their extension frequency and developmental context, this observation implies that a single cell type can employ two distinct types of cellular protrusions for signaling, each capable of transferring unique signaling molecules to potentially different targets.
Previous research in *Drosophila* has demonstrated that the air sac primordium (ASP) can extend multiple cytonemes that appear morphologically identical but convey different signaling molecules (Roy et al., 2011). However, it remains unknown whether the same cell type can extend two morphologically (and functionally) distinct protrusions that might also serve distinct functions in separate contexts. Consequently, future studies aimed at unraveling the regulation of these two types of protrusions could provide insights into how cells choose their signaling modalities based on their specific signaling requirements and environmental cues.

We have asked why cytoneme-mediated signaling is necessary between fully differentiated- and undifferentiated keratinocytes, even though they are separated by opposing cell membranes. Notably, we have identified the Notch signaling pathway as being involved, which operates through membrane receptor and ligand interactions between adjacent cells. This led us to hypothesize that Notch signaling between cells in these layers is suppressed by Fringe proteins, which modulate Notch signaling. In our research, we have discovered that *lunatic fringe (lfng)* is exclusively expressed in the intermediate layer in zebrafish ((Panin et al., 1997), fig. S3). Early studies in mice have also reported that three fringe genes exhibit differential expression in different epidermal layers (Thelu et al., 1998). Hence, it appears that Notch signaling is tightly regulated or inhibited by default between epidermal layers in both zebrafish and mice. While it is well documented that Notch signaling is essential for keratinocyte differentiation and

proliferation, the mechanism by which Notch signaling is initiated remains unclear. Our study suggests that cytonemes from fully differentiated keratinocytes in the periderm can activate Notch signaling in undifferentiated keratinocytes in intermediate layer. However, further investigations are necessary to fully comprehend how cytoneme-mediated signaling coordinate with Lunatic fringe or other Fringes to trigger the Notch signal.

It is widely believed that the overproduction of IL-17 is a major contributing factor to *psoriasis*, and many studies have demonstrated that IL-17 signaling triggers inflammation (Liu et al., 2020). IL-17 affects various cellular targets, such as keratinocytes, neutrophils, endothelial cells, promoting tissue inflammation (Blauvelt and Chiricozzi, 2018). Thus, further studies are essential to determine whether uncontrolled cytoneme-mediated signaling in keratinocytes correlates with the onset or progression of psoriatic or other skin diseases. Nevertheless, our data strongly indicate that cell-autonomous overexpression of *il17rd* in peridermal keratinocytes (*krt4*+) can induce psoriasis-like phenotypes by inhibiting cytoneme extensions, which subsequently leads to a reduction in Notch signaling in undifferentiated keratinocytes. A diminished Notch signal has been observed in psoriatic lesions (Ota et al., 2014). Our study suggests that cytoneme-mediated intercellular communication between keratinocytes plays a crucial role in epidermal maintenance, and its dysregulation may contribute to the development of human skin diseases. This idea has yet to be fully appreciated, and it could open up new avenues for understanding the onset or progression of human skin diseases.

We have demonstrated that IL-17 can influence cytoneme extension by regulating RhoA or Cdc42 GTPases, ultimately affecting actin polymerization. Consequently, it would be intriguing to investigate whether genes associated with *psoriasis* or other human skin diseases also function as regulators of cytonemes. For instance, the epidermal-specific deletion of inhibitor of NF-kB kinase 2 (IKK2) or the double knock out of the c-Jun and JunB genes in mice resulted in psoriasis-like phenotypes in a T cell-independent manner (Pasparakis et al., 2002; Zenz et al., 2005). Among the various roles of IKK and JunB, they are also known to regulate the actin cytoskeleton (Dubin-Bar et al., 2008; Lennikov et al., 2018; Otani et al., 2016). Thus, it is conceivable that dysregulated cytoneme-mediated signaling may contribute to psoriatic conditions in these contexts. Similarly, we have shown that *clint1* mutants exhibited compromised cytoneme signaling and keratinocyte hyperproliferation (fig. S6) (Dodd et al., 2009). Clint1(Clathrin Interactor 1) plays an important role in vesicle trafficking, and it is well established that endocytic pathways are critical for multiple steps in cytoneme-mediated morphogen delivery. Therefore, it is interesting to investigate whether clint1 functions as a cytoneme regulator could provide valuable insights.

Another important question requires an answer: the proper remodeling and maintenance of the epidermis seemingly depend on a well-controlled supply of keratinocytes from basal stem cells. Our study demonstrated the relay of cytoneme-mediated signals from the periderm to the underlying undifferentiated keratinocytes, leading to their differentiation and subsequent replacement of peridermal keratinocytes. However, the mechanism through which basal stem cells detect the need for an

increased population of keratinocytes to maintain the intermediate layer remains unclear. This process doesn't necessarily involve cellular protrusion-mediated signaling. However, we discovered cytoneme-like protrusions originating from undifferentiated keratinocytes (Fig. 2B). Our preliminary data, not presented in this manuscript, suggests that these cellular protrusions project downward, indicating their interaction with basal stem cells. Although further investigation is needed, it is conceivable that a replenishment signal from external sources or the periderm can be transmitted to stem cells through a multi-step cytoneme-mediated intercellular signaling pathways.

Figure 1. Differentiated keratinocytes extend cytoneme-like cellular protrusions
(A) Airineme-like protrusion (arrowhead), and (B) cytoneme-like protrusions (arrowheads) labelled in *krt4:palmEGFP* injected keratinocytes. (C) Keratinocytes expressing *krt4* extend cytoneme-like protrusions most frequently during metamorphic stages [6.5-7.5 SSL (*Parichy et al., 2009*); N=457 cells, 19 larvae total]. A cytoneme-like protrusion (arrowhead) labeled for (D) F-actin, revealed by *LifeAct-mRuby* but did not colocalize (E) with tubulin, revealed by *Tuba-mCherry*. (F) Keratinocyte cytonemes extend faster than they retract (N=125 cells, 4 larvae total), with an average length (G) of 18.21μm (N=190 cells, 4 larvae total). Scale bars: 20μm (A, B, D, E). Error bars indicate mean ± SEM.

Figure 2. Keratinocyte cytonemes establish physical contact with underlying undifferentiated keratinocytes

(A-A") Postembryonic zebrafish epidermis of *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* comprises three distinct layers. (A)Expression of *lyn-EGFP* in *krt4+* cells in the periderm layer. (A'-A") Expression of *lyn-tdTomato* (pseudo-colored magenta) in *krtt1c19e+* cells in (A') the intermediate layer and (A") the basal layer. Note that the green circular labelling in (A', arrowheads) represents mucous-secreting goblet cells (Chang and Hwang, 2011). (B) Undifferentiated keratinocytes (KC) extend significantly fewer cytoneme-like protrusions (*P*<0.0001, N=49, 3 larvae; N=91 cells, 6 larvae total). (C) Cytonemes from fully differentiated keratinocytes make physical contact with undifferentiated keratinocytes in the intermediate layer (red dotted circles).

Scale bars: 20µm (A-A", C). Error bars indicate mean ± SEM.

Figure 3. Cytoneme-mediated signaling regulates keratinocyte differentiation and proliferation
(A) Expression of *cdc42DN* effectively inhibits cytoneme extension ($F_{2, 231}$ = 25.38, *P*<0.0001, N=234 cells, 10 larvae total). (B) Periderm layer of *Tg(krt4:TetGBDTRE-v2a-Cdc42DN;krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* without drug treatment. (B') Periderm layer of *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* with drug treatment. (B'') Periderm layer of *Tg(krt4:TetGBDTRE-v2a-cdc42DN;krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* with drug treatment exhibits disorganization and regions lacking *krt4+* keratinocytes (arrowhead). Cytoneme inhibition increases *krtt1c19e* expression in the periderm (C) ($F_{2, 387}$ = 226.7, *P*<0.0001, N=39 larvae total) and leads to an increased number of keratinocytes within the periderm (D) ($F_{2, 18}$ = 27.36, *P*<0.0001, N=21 larvae total). (E) The cell death rate in the periderm is not affected by cytoneme inhibition. Tested with acridine orange incorporation ($F_{2, 17}$ = 3.192, *P*=0.0666, N=20 larvae total). Statistical significances were assessed by One-way ANOVA followed by Tukey's HSD post hoc test. Scale bars: 20μm (B-B"). Error bars indicate mean ± SEM.

Figure 4. Cytonemes activate Notch in undifferentiated keratinocytes
(A) Postembryonic zebrafish epidermis of *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato;TP1:H2BmCherry)* shows Notch responsiveness in undifferentiated keratinocytes within the intermediate layer but not in the periderm or basal layer. (B) Inhibition of cytonemes reduces Notch responsiveness in undifferentiated keratinocytes. ($F_{2, 33}$ = 48.27, P < 0.0001, N=36 larvae total). (C-C") Notch inhibition by expressing SuHDN in undifferentiated keratinocytes results in the disorganization of periderm (C"). (C) Properly organized periderm layer of *Tg(krtt1c19e:tetGBDTRE-v2a-SuHDN;krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* without drug treatment and (C') *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* with drug treatment. (D-F) Notch inhibition (D) increases *krtt1c19e* signal in the periderm ($F_{2, 397}$ = 89.47, P < 0.0001, N=40 larvae total), (E) reduces Notch responsiveness in undifferentiated keratinocytes ($F_{2, 32}$ = 13.18, P < 0.0001, N=35 larvae total), and leads to (F) an increased number of keratinocytes in the periderm ($F_{2, 19}$ = 6.620, P = 0.0066, N=22 larvae total). Statistical significances were assessed by One-way ANOVA followed by Tukey's HSD post hoc test. Scale bars, 20μm (C-C"). Error bars indicate mean ± SEM.

Figure 5. Interleukin-17 regulates keratinocyte cytonemes
Overexpression of *il17rd* or *il17ra1a* in *krt4+* keratinocytes results in (A) a significant reduction in cytoneme extension ($F_{2, 325}$ = 38.48, *P* < 0.0001, N=328 cells, 10 larvae total), and (B) decreased Notch responsiveness ($F_{2, 34}$ = 23.16, *P* < 0.0001, N=37 larvae total). (C-C') The periderm is disrupted in (C') *Tg(krt4:nVenus-v2a-il17rd;krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* compared to the (C) properly arranged periderm of *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)*, resulting in (D) an increased *krtt1c19e* signal in the periderm (D) ($F_{2, 287}$ = 109.7, *P* < 0.0001, N=29 larvae total) and (E) an increased number of keratinocytes in the periderm ($F_{2, 15}$ = 27.79, *P* < 0.0001, N=18 larvae total). (F) Simultaneous overexpression of *cdc42DN* and *il17rd* in *krt4+* keratinocytes further reduces cytoneme extension further (*P* < 0.0001, N=164 cells, 7 larvae total). (G) *il17rd*-overexpressing *krt4+* keratinocytes exhibit disorganization of microridges (arrowhead). (H) Peridermal cells sorted by FACS show significantly elevated *RhoA* and decreased *Cdc42* expression. Statistical significances were assessed by One-way ANOVA followed by Tukey's HSD post hoc test and Student t test. Scale bars: 20µm (C, G). Error bars indicate mean ± SEM.

Acknowledgements

We thank for the invaluable discussions with Drs. Plikus, Andersen, Parsons and all the members of Eom lab in maintaining our fish lines.

Funding: This study was supported by the National Institutes of Health grant R35GM142791 (to D.S.E.) and National Institutes of Health T32 training grant AR080622 (to Y.W.).

Author Contributions:
Conceptualization: YW, DSE
Investigation: YW, QH, OH, KF, DSE
Writing – Original Drafting: YW, DSE
Writing – Review & editing: YW, DSE
Funding Acquisition: DSE, YW
Supervision: DSE

Competing interests: The authors declare no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials.

Materials and Methods

Zebrafish husbandry and maintenance
Adult zebrafish were maintained at a constant temperature of 28.5°C under a 16:8 (L:D) cycle. The zebrafish used in this study were wild-type AB^{wp} or its derivative WT(ABb). Additionally, transgenic lines used include *Tg(krt4:lyn-EGFP)^{sq18}, Tg(krtt1c19e:lyn-tdTomato)^{sq16}* provided by T. Carney; *Tg(Tp1:EGFP)^{um14}, Tg(Tp1:H2BmCherry)^{s939}* provided by M. Parsons*; clint1a$^{hi1520Tg/+}$(AB),* obtained from the Zebrafish International Resource Center (ZIRC). The sexes of individual fish could not be distinguished since experiments were performed before the development of secondary sexual characteristics. All zebrafish stocks used in this study were known to produce balanced sex ratios, ensuring that experiments sampled similar numbers of males and females. All animal work conducted in this study received approval from the University of California Irvine Institutional Animal Care and Use Committee (protocol #AUP-22-023) and adhere to institutional and federal guidelines for the ethical use of animals.

Transgenesis and transgenic line production
Transgenic fish lines with cell-type specific and temporally inducible expression of human *cdc42DN* and *SuHDN* were generated following the protocol described by Eom et al., (2015). These constructs were driven by a 2.2kb *krt4* promoter obtained from T. Carney for the expression in fully differentiated keratinocytes and a 3.9kb *krtt1c19e* promoter, also from T. Carney, for the expression in undifferentiated keratinocytes. The efficiency of gene expression in both *Tg(krt4:TetGBDTRE-v2a-cdc42DN)$^{ir.rt3}$* and *Tg(krtt1c19e:TetGBDTRE-v2a-SuHDN)$^{ir.rt18}$*, induced by doxycycline (Dox; Sigma-Aldrich) and dexamethasone (Dex; Sigma-Aldrich), was performed in both F0 mosaic fish and non-mosaic stable lines, with consistent results. The coding sequence of *il17rd* (ZDB-GENE-020320-5) and *il17ra1a* (ZDB-GENE-070705-242) were isolated from WT(ABb) cDNA and used to generate constructs driven by the *krt4* promoter, which were assembled using Gateway assembly into the pDestTol2CG2 destination vector to produce *Tg(krt4:nVenus-v2a-il17rd)$^{ir.rt16}$* and *Tg(krt4:il17ra1a-v2a-mCherry)$^{ir.rt17}$*. Constitutive Notch activation used the intracellular domain of notch1a (NICD) linked by 2a sequence to nVenus and driven by the *krtt1c19e* promoter for the production of *Tg(krtt1c19e:nVenus-v2a-NICD)$^{ir.rt19}$*. The lunatic fringe reporter line, *Tg(lfng:mCherry)$^{ir.rt20}$*, was generated by first amplifying the *lfng* promoter sequence (ZDB-GENE-980605-16) using WT gDNA and then used Gateway assembly into the pDestTol2-exorh:EGFP destination vector [Addgene #195983]. These constructs were then injected into WT(ABb) fish to generate mosaic F0 fish, which were then bred in pairs to find germline carriers to produce non-mosaic transgenic lines. Actin was examined with *LifeAct-mRuby* [Addgene #166977] (Barros-Becker et al., 2017) driven by *krt4* promoter. Microtubules in *krt4*+ keratinocytes and cytonemes were labelled using subclones from Tuba1-mCherry [(Addgene #49149] (Friedman et al., 2010).

Drug treatments
Zebrafish were kept in E3 medium with or without drugs (Dox/Dex) during both the light and the dark cycles for long-term drug treatment experiments. For overnight time-lapse images, fish subjected to drug-treatment were transferred to fresh E3 medium 30min

before explant preparation to minimize background fluorescence from the drugs. Zebrafish with *krt4:palmEGFP* injected TetGBD-containing transgenics and non-transgenic siblings were treated with 37.5µM Dox and 50µM Dex one day prior to time-lapse imaging, while fish for long-term drug treatment received 27.5 µM Dox and 50µM Dex starting from SSL7.0. Notch inhibitor, LY411575 (10mM; Thomas Scientific) was prepared in DMSO (Sigma-Aldrich), and fish were treated with drug [LY411575 (3µM)] or vehicle from SSL7.0. For acute drug administration, Cdc42 inhibitor ML141 (Sigma-Aldrich), prepared in DMSO, was added to fish medium [ML141 (2µM)] 30min prior to explant preparation for time-lapse imaging.

Acridine Orange Cell Death Assay
Acridine Orange (5mg/mL; Sigma-Aldrich) was prepared in Invitrogen UltraPure™ Distilled Water. Fish were washed in 1x E3 medium 30min before being transferred to 1x E3 medium containing AO (2µg/mL). The fish were incubated in AO for 30min and then rinsed in regular 1x E3 medium three times for 10min each to minimize background fluorescence during imaging. Dead cells, pseudo-colored in magenta, were counted for all groups.

Time-lapse and still imaging
Ex vivo time-lapse imaging of zebrafish epidermal cells within their native tissue environment was acquired at 3-minute intervals for 10 hours. This was achieved using a 40x water-immersion objective on a Leica TCS SP8 confocal microscope equipped with a resonant scanner and two HyD detectors. Larvae were imaged at SSL7.5, unless otherwise specified. For still images, the region directly underneath the dorsal fin of the zebrafish was captured, with a total of 10 tiles per fish.

RT-PCR, qRT-PCR and FACS
Zebrafish at SSL7.5 were skinned (N=15 per group), and the dissected tissues were placed in cold 1x PBS. Tissues were washed by pipetting up and down and then centrifuged to remove the supernatant. Next, 1mL Gibco Stem® Pro® Accutase Cell Dissociation Reagent was added to resuspend tissues, and the tube was incubated for at least 10 min at 37°C to dissociate the cells. After incubation, the cells were passed through 40µm cell strainers to remove large tissue chunks before subjecting them to FACS (Fluorescence-activated cell sorting). FACS enabled the separation and collection of desired cell populations based on different cell membrane markers. EGFP+ and tdtomato+ cells were collected for downstream non-quantitative RT-PCR and quantitative RT-PCR. For cDNA synthesis, the Invitrogen Cells Direct™ One-Step qRT-PCR Kit was employed. Quantitative PCR was conducted on an Applied Biosystems QuantStudio™ 3 Real-Time PCR Instrument using custom TaqMan probes for *rhoab* (Dr03439241_mH) and *cdc42* (Dr03139234_m1). Quantitative PCRs were run with at least three triplicates for each sample. Non-quantitative RT-PCR amplifications were performed with 40 cycles (*actb1, dlc, lfng, krt4, krt19, il17rd, notch2, notch3, il17ra1a*) using Takara PrimeStar GXL DNA Polymerase. *actb1*: 5'-CATCCGTAAGGACCTGTATGCCAAC- 3', 5'- AGGTTGGTCGTTCGTTTGAATCTC- 3' (Hu et al., 2016); *krt4*: 5'- TCAACCAGGTCTATCTCTTACTCC- 3', 5'-AACAGGCTCTGGTTGACAGTTAC- 3'; *krt19*: 5'- GCTACCACCTTCTCCAGCGGAAG-

3', 5'- TGCAGCACCAAATCCTCCACCAG- 3'; *lfng*: 5'-
CATATCGCGGAAAAGTCGTGGTC- 3', 5'- GCTTATCCGCTTCCCTCCTGCTG- 3'; *dlc*:
5'- CGGGAATCGTCTCTTTGATAAT- 3', 5'- CTCACCGATAGCGAGTCTTCTT- 3';
il17rd: 5'- CTGGGTCACGGAGACTTGCGC- 3', 5'- TGGCGTGTTTCCCCAGCGGAC-
3'; *il17ra1a*: 5'- TCACCTTTATACCTGTAGTGCTG- 3'; 5'-
CATGTTTTCTCTGACATCAAAAC-3'; *notch2*: 5'- GAGTGTGTGGACCCGTTAGTATG-
3', 5'- GCAGGCATCATCAATGTGACAC- 3'; *notch3*: 5'-
TCAGGATTGTTCTCTCGTTGATG- 3', 5'- GTGTTAAAGCATGTACCACCATTG- 3'.

Cell Counts
For all experiments, 6 out of 10 tiles of still images, which represent the central trunk
area directly beneath the dorsal fin, were selected for cell counts. To ensure
consistency across all samples and minimize discrepancies resulting from variations in
fish trunk widths, a region of interest (ROI), 290μm x 200μm per tile, was created using
the rectangle selection tool in ImageJ. Z-stack imaging was confined to the
superficial/periderm layer based on the cross-sectional view, and the final count
included both *krt4+* and *krtt1c19e+* cells at the periderm layer.

Intensity analyses
To analyze Notch expression level, tile images were merged separately for each fish.
Merged images were then sectioned in ImageJ using the polygon selection tool to
encompass only the fish trunk, and maximal projection was utilized to obtain the mean
gray value. For *krtt1c19e+* expression in the periderm layer, the z-stack was set as
previously described, and the mean gray value of *krtt1c19e+* expression was obtained.

Statistical analyses
GraphPad Prism software version 9.0.0 for Windows (GraphPad Software, San Diego,
CA, USA) was used to perform statistical analyses. Continuous data were evaluated by
Student t-test and Post hoc means were compared by Tukey-Kramer HSD.

Figure S1. Effect of actin inhibitor treatment on cytoneme extension
(A) Treatment with the actin inhibitor (ML141) significantly reduced cytoneme extension frequency
(P<0.0001, N=3 larvae per group). Statistical significance was assessed using Student t test. Error bars
indicate mean ± SEM.

Figure S2. Goblet cell count in the zebrafish epidermis
(A) Quantification of goblet cell numbers revealed no significant difference between two controls and
experimental *Tg(krt4:TetGBDTRE-v2a-cdc42DN)* zebrafish ($F_{2,20}$ = 2.763, P = 0.0872, N=23 larvae total).
(B) Quantification of goblet cell numbers showed no significant difference between two controls and
experimental *Tg(krtt1c19e:TetGBDTRE-v2a-SuHDN)* zebrafish ($F_{2, 30}$ = 3.040, P = 0.0628, N= 33 larvae
total). Statistical significances were assessed by One-way ANOVA followed by Tukey's HSD post hoc
test.

Figure S3. Gene expressions in keratinocytes
RT-PCR revealed the endogenous expression of *krt4, dlc, il17ra1a*, and *il17rd* in keratinocytes FACS-
sorted for EGFP+ cells from *Tg(krt4:lyn-EGFP)*. Additionally, *krtt1c19e, notch2, notch 3*, and *Ifng* were
detected in tdTomato+ cells sorted from *Tg(krtt1c19e:tdTomato).*

Figure S4. Notch reporter (*Tp1*) expression
(A) Images before and after treatment with DMSO and LY411575 (Notch inhibitor) in the *Tg(Tp1:EGFP)*.
(B) Treatment with LY411575 resulted in a significant overall reduction in Notch expression levels in the TP1 Notch reporter line compared to the DMSO control group ($P<0.001$, N=5 larvae per group). (C) Images before and after cytoneme inhibition through drug induction of cdc42DN-expression in *krt4+* keratinocytes and direct inhibition of Notch signaling via drug induction of SuHDN expression in *krtt1c19e+* keratinocytes, along with DMSO treatment. (D) The number of *Tp1+* keratinocytes peaked during epidermal remodeling (SSL6.5-7.5, N=58 larvae total) in zebrafish. Statistical significance was assessed by Student t test. Scale bars: 100μm (A, C). Error bars indicate mean ± SEM.

Figure S5. Notch signal modifiers in undifferentiated keratinocytes
(A-A') Images showing *lunatic fringe (lfng)* expression in the intermediate layer of epidermis. (B-B')
Overexpression of Notch IntraCellular Domain (NICD) in undifferentiated keratinocytes resulted in (C, B')
an increased krtt1c19e+ signal in the periderm (*P*<0.0001, N= 6 larvae per group) and (D) a greater
number of keratinocytes within the periderm (*P*<0.001, N=6 larvae per group). Statistical significance was
assessed using Student t test. Scale bars: 20µm (A-A', B-B'). Error bars indicate mean ± SEM.

Figure S6. *clint1* regulates epidermal maintenance via keratinocyte cytonemes
(A-A') *clint1* mutants exhibit a disorganized periderm. (B-E) *clint1* mutants show (B) a lower frequency of cytoneme extension as compared to controls (WT, *cx41.8*[-/-]) ($F_{2, 186}$ = 14.55, *P*<0.0001, N=10 larvae total), (C) reduced Notch responsiveness (*P*<0.0001, N=12 mutants, N=10 controls), (D) increased *krtt1c19e* expression in the periderm (*P*<0.0001, N=13 mutants, N=6 controls), and (E) a greater number of keratinocytes within the periderm (*P*<0.001, N=8 mutants, N=6 controls). Statistical significances were assessed by One-way ANOVA followed by Tukey's HSD post hoc test and Student t test. Scale bars, 20µm (A-A'). Error bars indicate mean ± SEM.

Movies S1. Cytoneme extension in keratinocytes
In cdc42DN-expressing *krt4+* (green) keratinocytes (right), cytoneme extension is significantly inhibited. It is worth nothing that normal filopodial and lamellipodial extension seem unaffected. In the control groups (left and middle), cytonemes actively extend. The movie was captured at 3-minute intervals, and the hours and minutes elapsed are displayed in the upper right corner.

References

Abdou, A.G., Maraee, A.H., Sharaf, A., and Elnaidany, N.F. (2012). Up-regulation of Notch-1 in psoriasis: an immunohistochemical study. Ann Diagn Pathol *16*, 177-184.

Akat, E., Yenmis, M., Pombal, M.A., Molist, P., Megias, M., Arman, S., Vesely, M., Anderson, R., and Ayaz, D. (2022). Comparison of vertebrate skin structure at class level: A review. Anat Rec (Hoboken) *305*, 3543-3608.

Armstrong, A.W., and Read, C. (2020). Pathophysiology, Clinical Presentation, and Treatment of Psoriasis: A Review. JAMA *323*, 1945-1960.

Barros-Becker, F., Lam, P.Y., Fisher, R., and Huttenlocher, A. (2017). Live imaging reveals distinct modes of neutrophil and macrophage migration within interstitial tissues. J Cell Sci *130*, 3801-3808.

Blanpain, C., and Fuchs, E. (2009). Epidermal homeostasis: a balancing act of stem cells in the skin. Nat Rev Mol Cell Biol *10*, 207-217.

Blanpain, C., Lowry, W.E., Pasolli, H.A., and Fuchs, E. (2006). Canonical notch signaling functions as a commitment switch in the epidermal lineage. Genes Dev *20*, 3022-3035.

Blauvelt, A., and Chiricozzi, A. (2018). The Immunologic Role of IL-17 in Psoriasis and Psoriatic Arthritis Pathogenesis. Clin Rev Allergy Immunol *55*, 379-390.

Borowczyk, J., Buerger, C., Tadjrischi, N., Drukala, J., Wolnicki, M., Wnuk, D., Modarressi, A., Boehncke, W.H., and Brembilla, N.C. (2020). IL-17E (IL-25) and IL-17A Differentially Affect the Functions of Human Keratinocytes. J Invest Dermatol *140*, 1379-1389 e1372.

Bowman, R.L., Wang, D., and Eom, D.S. (2023). A macrophage subpopulation promotes airineme-mediated intercellular communication in a matrix metalloproteinase-9 dependent manner. Cell Rep *42*, 112818.

Campa, M., Mansouri, B., Warren, R., and Menter, A. (2016). A Review of Biologic Therapies Targeting IL-23 and IL-17 for Use in Moderate-to-Severe Plaque Psoriasis. Dermatol Ther (Heidelb) *6*, 1-12.

Chang, W.J., and Hwang, P.P. (2011). Development of zebrafish epidermis. Birth Defects Res C Embryo Today *93*, 205-214.

Dale, J.K., Maroto, M., Dequeant, M.L., Malapert, P., McGrew, M., and Pourquie, O. (2003). Periodic notch inhibition by lunatic fringe underlies the chick segmentation clock. Nature *421*, 275-278.

Daly, C.A., Hall, E.T., and Ogden, S.K. (2022). Regulatory mechanisms of cytoneme-based morphogen transport. Cell Mol Life Sci *79*, 119.

Das, S., Srinivasan, S., Srivastava, A., Kumar, S., Das, G., Das, S., Dwivedi, A., Karulkar, A., Makkad, K., Bilala, R.*, et al.* (2019). Differential Influence of IL-9 and IL-17 on Actin Cytoskeleton Regulates the Migration Potential of Human Keratinocytes. J Immunol *202*, 1949-1961.

Dodd, M.E., Hatzold, J., Mathias, J.R., Walters, K.B., Bennin, D.A., Rhodes, J., Kanki, J.P., Look, A.T., Hammerschmidt, M., and Huttenlocher, A. (2009). The ENTH domain protein Clint1 is required for epidermal homeostasis in zebrafish. Development *136*, 2591-2600.

Dubin-Bar, D., Bitan, A., Bakhrat, A., Kaiden-Hasson, R., Etzion, S., Shaanan, B., and Abdu, U. (2008). The Drosophila IKK-related kinase (Ik2) and Spindle-F proteins are part of a complex that regulates cytoskeleton organization during oogenesis. BMC Cell Biol *9*, 51.

Eisenhoffer, G.T., Slattum, G., Ruiz, O.E., Otsuna, H., Bryan, C.D., Lopez, J., Wagner, D.S., Bonkowsky, J.L., Chien, C.B., Dorsky, R.I.*, et al.* (2017). A toolbox to study epidermal cell types in zebrafish. J Cell Sci *130*, 269-277.

Eom, D.S. (2020). Airinemes: thin cellular protrusions mediate long-distance signalling guided by macrophages. Open Biol *10*, 200039.

Eom, D.S., Bain, E.J., Patterson, L.B., Grout, M.E., and Parichy, D.M. (2015). Long-distance communication by specialized cellular projections during pigment pattern development and evolution. Elife *4*.

Eom, D.S., and Parichy, D.M. (2017). A macrophage relay for long-distance signaling during postembryonic tissue remodeling. Science *355*, 1317-1320.

Fragoulis, G.E., Siebert, S., and McInnes, I.B. (2016). Therapeutic Targeting of IL-17 and IL-23 Cytokines in Immune-Mediated Diseases. Annu Rev Med *67*, 337-353.

Friedman, J.R., Webster, B.M., Mastronarde, D.N., Verhey, K.J., and Voeltz, G.K. (2010). ER sliding dynamics and ER-mitochondrial contacts occur on acetylated microtubules. J Cell Biol *190*, 363-375.

Gonzalez-Fernandez, C., Chaves-Pozo, E., and Cuesta, A. (2020). Identification and Regulation of Interleukin-17 (IL-17) Family Ligands in the Teleost Fish European Sea Bass. Int J Mol Sci *21*.

Gratton, R., Tricarico, P.M., Moltrasio, C., Lima Estevao de Oliveira, A.S., Brandao, L., Marzano, A.V., Zupin, L., and Crovella, S. (2020). Pleiotropic Role of Notch Signaling in Human Skin Diseases. Int J Mol Sci *21*.

Hall, A. (1998). Rho GTPases and the actin cytoskeleton. Science *279*, 509-514.

Hu, Y., Xie, S., and Yao, J. (2016). Identification of Novel Reference Genes Suitable for qRT-PCR Normalization with Respect to the Zebrafish Developmental Stage. PLoS One *11*, e0149277.

Jaffe, A.B., and Hall, A. (2005). Rho GTPases: biochemistry and biology. Annu Rev Cell Dev Biol *21*, 247-269.

Knopf, F., Schnabel, K., Haase, C., Pfeifer, K., Anastassiadis, K., and Weidinger, G. (2010). Dually inducible TetON systems for tissue-specific conditional gene expression in zebrafish. Proc Natl Acad Sci U S A *107*, 19933-19938.

Kornberg, T.B., and Roy, S. (2014). Cytonemes as specialized signaling filopodia. Development *141*, 729-736.

Lee, R.T., Asharani, P.V., and Carney, T.J. (2014). Basal keratinocytes contribute to all strata of the adult zebrafish epidermis. PLoS One *9*, e84858.

Lennikov, A., Mirabelli, P., Mukwaya, A., Schaupper, M., Thangavelu, M., Lachota, M., Ali, Z., Jensen, L., and Lagali, N. (2018). Selective IKK2 inhibitor IMD0354 disrupts NF-kappaB signaling to suppress corneal inflammation and angiogenesis. Angiogenesis *21*, 267-285.

Li, Q., Frank, M., Thisse, C.I., Thisse, B.V., and Uitto, J. (2011). Zebrafish: a model system to study heritable skin diseases. J Invest Dermatol *131*, 565-571.

Liu, T., Li, S., Ying, S., Tang, S., Ding, Y., Li, Y., Qiao, J., and Fang, H. (2020). The IL-23/IL-17 Pathway in Inflammatory Skin Diseases: From Bench to Bedside. Front Immunol *11*, 594735.

Lowes, M.A., Suarez-Farinas, M., and Krueger, J.G. (2014). Immunology of psoriasis. Annu Rev Immunol *32*, 227-255.

Ma, L., Li, Y., Peng, J., Wu, D., Zhao, X., Cui, Y., Chen, L., Yan, X., Du, Y., and Yu, L. (2015). Discovery of the migrasome, an organelle mediating release of cytoplasmic contents during cell migration. Cell Res *25*, 24-38.

Martinez-Navarro, F.J., Martinez-Menchon, T., Mulero, V., and Galindo-Villegas, J. (2019). Models of human psoriasis: Zebrafish the newly appointed player. Dev Comp Immunol *97*, 76-87.

Melentijevic, I., Toth, M.L., Arnold, M.L., Guasp, R.J., Harinath, G., Nguyen, K.C., Taub, D., Parker, J.A., Neri, C., Gabel, C.V.*, et al.* (2017). C. elegans neurons jettison protein aggregates and mitochondria under neurotoxic stress. Nature *542*, 367-371.

Moriyama, M., Durham, A.D., Moriyama, H., Hasegawa, K., Nishikawa, S., Radtke, F., and Osawa, M. (2008). Multiple roles of Notch signaling in the regulation of epidermal development. Dev Cell *14*, 594-604.

Mosca, M., Hong, J., Hadeler, E., Hakimi, M., Liao, W., and Bhutani, T. (2021). The Role of IL-17 Cytokines in Psoriasis. Immunotargets Ther *10*, 409-418.

Nguyen, B.C., Lefort, K., Mandinova, A., Antonini, D., Devgan, V., Della Gatta, G., Koster, M.I., Zhang, Z., Wang, J., Tommasi di Vignano, A.*, et al.* (2006). Cross-regulation between Notch and p63 in keratinocyte commitment to differentiation. Genes Dev *20*, 1028-1042.

Nickoloff, B.J., Qin, J.Z., Chaturvedi, V., Denning, M.F., Bonish, B., and Miele, L. (2002). Jagged-1 mediated activation of notch signaling induces complete maturation of human keratinocytes through NF-kappaB and PPARgamma. Cell Death Differ *9*, 842-855.

Nowell, C., and Radtke, F. (2013). Cutaneous Notch signaling in health and disease. Cold Spring Harb Perspect Med *3*, a017772.

Ota, T., Takekoshi, S., Takagi, T., Kitatani, K., Toriumi, K., Kojima, T., Kato, M., Ikoma, N., Mabuchi, T., and Ozawa, A. (2014). Notch signaling may be involved in the abnormal differentiation of epidermal keratinocytes in psoriasis. Acta Histochem Cytochem *47*, 175-183.

Otani, T., Ogura, Y., Misaki, K., Maeda, T., Kimpara, A., Yonemura, S., and Hayashi, S. (2016). IKKepsilon inhibits PKC to promote Fascin-dependent actin bundling. Development *143*, 3806-3816.

Panin, V.M., Papayannopoulos, V., Wilson, R., and Irvine, K.D. (1997). Fringe modulates Notch-ligand interactions. Nature *387*, 908-912.

Parichy, D.M., Elizondo, M.R., Mills, M.G., Gordon, T.N., and Engeszer, R.E. (2009). Normal table of postembryonic zebrafish development: staging by externally visible anatomy of the living fish. Dev Dyn *238*, 2975-3015.

Park, S., Kim, H., Wang, Y., Eom, D.S., and Allard, J. (2022). Zebrafish airinemes optimize their shape between ballistic and diffusive search. Elife *11*.

Parsons, M.J., Pisharath, H., Yusuff, S., Moore, J.C., Siekmann, A.F., Lawson, N., and Leach, S.D. (2009). Notch-responsive cells initiate the secondary transition in larval zebrafish pancreas. Mech Dev *126*, 898-912.

Pasparakis, M., Courtois, G., Hafner, M., Schmidt-Supprian, M., Nenci, A., Toksoy, A., Krampert, M., Goebeler, M., Gillitzer, R., Israel, A.*, et al.* (2002). TNF-mediated inflammatory skin disease in mice with epidermis-specific deletion of IKK2. Nature *417*, 861-866.

Pinto, C.S., Khandekar, A., Bhavna, R., Kiesel, P., Pigino, G., and Sonawane, M. (2019). Microridges are apical epithelial projections formed of F-actin networks that organize the glycan layer. Sci Rep *9*, 12191.

Rangarajan, A., Talora, C., Okuyama, R., Nicolas, M., Mammucari, C., Oh, H., Aster, J.C., Krishna, S., Metzger, D., Chambon, P.*, et al.* (2001). Notch signaling is a direct determinant of keratinocyte growth arrest and entry into differentiation. EMBO J *20*, 3427-3436.

Roy, S., Hsiung, F., and Kornberg, T.B. (2011). Specificity of Drosophila cytonemes for distinct signaling pathways. Science *332*, 354-358.

Sahlen, P., Spalinskas, R., Asad, S., Mahapatra, K.D., Hojer, P., Anil, A., Eisfeldt, J., Srivastava, A., Nikamo, P., Mukherjee, A.*, et al.* (2021). Chromatin interactions in differentiating keratinocytes reveal novel atopic dermatitis- and psoriasis-associated genes. J Allergy Clin Immunol *147*, 1742-1752.

Schempp, C., Emde, M., and Wolfle, U. (2009). Dermatology in the Darwin anniversary. Part 1: Evolution of the integument. J Dtsch Dermatol Ges *7*, 750-757.

Segre, J.A. (2006). Epidermal barrier formation and recovery in skin disorders. J Clin Invest *116*, 1150-1158.

Stanganello, E., Hagemann, A.I., Mattes, B., Sinner, C., Meyen, D., Weber, S., Schug, A., Raz, E., and Scholpp, S. (2015). Filopodia-based Wnt transport during vertebrate tissue patterning. Nat Commun *6*, 5846.

Thelu, J., Rossio, P., and Favier, B. (2002). Notch signalling is linked to epidermal cell differentiation level in basal cell carcinoma, psoriasis and wound healing. BMC Dermatol *2*, 7.

Thelu, J., Viallet, J.P., and Dhouailly, D. (1998). Differential expression pattern of the three Fringe genes is associated with epidermal differentiation. J Invest Dermatol *111*, 903-906.

van Loon, A.P., Erofeev, I.S., Maryshev, I.V., Goryachev, A.B., and Sagasti, A. (2020). Cortical contraction drives the 3D patterning of epithelial cell surfaces. J Cell Biol *219*.

Wu, B., Jin, M., Zhang, Y., Wei, T., and Bai, Z. (2011). Evolution of the IL17 receptor family in chordates: a new subfamily IL17REL. Immunogenetics *63*, 835-845.

Zenz, R., Eferl, R., Kenner, L., Florin, L., Hummerich, L., Mehic, D., Scheuch, H., Angel, P., Tschachler, E., and Wagner, E.F. (2005). Psoriasis-like skin disease and arthritis caused by inducible epidermal deletion of Jun proteins. Nature *437*, 369-375.

Zhang, C., and Scholpp, S. (2019). Cytonemes in development. Curr Opin Genet Dev *57*, 25-30.

Zhou, X., Chen, Y., Cui, L., Shi, Y., and Guo, C. (2022). Advances in the pathogenesis of psoriasis: from keratinocyte perspective. Cell Death Dis *13*, 81.