# Plasticity-led and mutation-led evolutions are discrete modes of the same developmental gene regulatory network.

Eden Tian Hwa Ng and Akira R. Kinjo

Department of Mathematics, Faculty of Science,

Universiti Brunei Darussalam,

Jalan Tungku Link, BE1410, Brunei Darussalam

November 9, 2023

**Abstract**

Plasticity-led evolution (PLE), where evolution is initiated by the environmental induction of novel adaptive phenotypes, which is followed by genetic accommodation, has been proposed to remedy the difficulties of mutation-led evolution (MLE) implied by the standard evolutionary theory, according to which evolution is initiated by mutations causing phenotypic variation to be naturally selected. Under PLE, developmental systems should be able to respond to environmental changes adaptively. However, developmental systems are known to be robust against environmental or mutational perturbations. Therefore, we expect a transition from a robust state to a plastic one. Here, we simulated the evolution of a gene regulatory network (GRN) model that incorporates hierarchical developmental regulation and environmental cues over different magnitudes of environmental changes. We show that this GRN model exhibits PLE under large environmental changes and MLE under small environmental changes. Furthermore, this GRN model is susceptible to environmental or mutational fluctuations under large environmental changes but is robust under small environmental changes, indicating a breakdown of robustness

due to large environmental changes. It is also observed that developmental bias emerges and is aligned to the environmental changes before the breakdown of robustness, as though the population is readying itself for rapid adaptive evolution. These observations suggest that the evolutionary transition from mutation-led to plasticity-led evolution is underlain by a developmental transition from robust to susceptible regimes over increasing magnitudes of environmental change. Thus, the GRN model can unify these conflicting theories of evolution.

**Keywords:** phenotypic plasticity, environmental change, developmental bias, cryptic mutations, computer simulation.

**Significance statement:** The standard evolutionary theory posits that evolution is initiated by mutations causing heritable variation to be naturally selected. This mutation-led evolution scheme has been challenged by an alternative theory called plasticity-led evolution, whereby environmentally induced adaptive phenotypes are genetically accommodated. By performing evolutionary simulations using a gene regulatory network model incorporating hierarchical developmental regulation and environmental cues, we show that this model transitions between mutation-led and plasticity-led evolutionary schemes depending on the magnitude of environmental changes. Furthermore, we found that this evolutionary transition is underlain by a developmental transition from a state robust against environmental and mutational perturbations to a susceptible one, highlighting the critical role of the developmental process, primarily neglected in the standard evolutionary theory.

# Introduction

Plasticity-led evolution (PLE) is a process in which novel adaptive phenotypes initially induced by a new environment are genetically accommodated over generations [1, 2, 3]. The ability to respond to environmental cues by changing the phenotype without changing the genotype is called phenotypic plasticity [1, 4, 5]. PLE has been proposed to remedy the problem of gradualism implied by the Modern Evolutionary Synthesis [6, 7, 8, 9, 4].

In contrast, mutation-led evolution (MLE) refers to evolution initiated by a mutation that induces a novel adaptive phenotype [5]. Since mutations are random, an adaptive mutation is expected to appear only after a long time, assuming the population has not yet gone extinct. While the Modern Evolutionary Synthesis is necessarily a form of MLE, MLE itself encompasses more general concepts, particularly hierarchical developmental process [5, 10]. Our previous work shows that, even without environmental cues, the developmental process is still essential for adaptive evolution in a plausible time scale [11]. In comparison, it has been observed that quantitative genetics models, not incorporating developmental processes, evolve far more slowly even with environmental cues [12, 13]. In this work, we consider a model for MLE that incorporates hierarchical developmental processes.

In our previous work [11], we proposed a gene regulatory network (GRN) model that exhibits behaviors compatible with PLE upon large environmental changes [2]. However, this model is highly robust against mutations and environmental noises in already adapted environments [11, 14, 15, 16]. How can the same system be robust in one environment yet exhibit a plastic response to another? This observation suggests there should be a transition from a robust to a plastic regime, depending on the magnitude of environmental change. In the following, we show that this transition manifests as a breakdown of robustness. Furthermore, we demonstrate that the system undergoes MLE under the robust regime and PLE under the plastic regime, thereby unifying two conflicting evolutionary schemes as two modes of the same developmental system.

# Modeling

Here, we briefly describe our evolutionary models based on gene regulatory networks. For the details, see Methods and our previous paper [11].

## Development

In the following, we compare two variants [11] of the gene regulatory network (GRN) model introduced by A. Wagner [14]. In these models, the phenotype is a vector, and the genome is a set of interaction matrices. The first model, the *Full* model,

incorporating environmental cues, developmental processes, and hierarchical regulation, exhibits PLE under large environmental changes. The environmental cue serves as the environmental input to the network. The developmental process is a recursive process that updates the system's internal state, integrating environmental and genetic information into the phenotype. The hierarchical regulation reflects the multilayered nature of biological regulation. The other model, the *NoCue* model, is identical to the Full model, except it does not take in environmental cues. As a result, it does not exhibit phenotypic plasticity and, therefore, serves as a model for MLE. See Methods for the details on the model architectures and how the developmental process determines phenotypes. We note that these models are mathematically equivalent to artificial recurrent neural networks so that they have generic learning capabilities. In this context, adaptive evolution can be viewed as a process of learning the environment by changing the genome (network weights) [17]. Notably, the Full model can adaptively respond to environmental cues through development.

## Environment

In plasticity-led evolution, the environment plays two roles: selector and inducer of phenotypes [1]. We assume that these two roles of the environment are highly correlated [11].

We model the environment-as-selector as a 200-dimensional vector $\mathbf{e}$ of $\pm 1$. This vector is also the optimal phenotype used to calculate the fitness of each individual's phenotype (also modeled as a 200-dimensional vector; see Methods). For the Full model, we model the environment-as-inducer as each individual's environmental cue $e$. This vector is obtained by randomly flipping 5% of the elements of $\mathbf{e}$. The environmental cue is absent in the NoCue model.

## Evolution

We simulate evolution by applying a genetic algorithm [18] to a population of 1000 individuals [11]. We let the population adapt to a constant environment $\mathbf{e}$ for 200 generations. This duration is called an *epoch*. Before simulating for another epoch, we change the environment $\mathbf{e}$ by flipping a specified proportion of the elements of

this vector. We iterate this procedure for many epochs. Before the production runs given in the Results section, we ran 40 epochs of preparatory runs under 40 completely uncorrelated environments. We did this to equilibrate the randomly initialized population and to let the system learn how to respond to environmental changes [11]. In the following, the novel and ancestral environments refer to current and previously adapted environments.

# Results

In the following, we study evolution over various magnitudes of environmental change. We show that the Full model exhibits characteristics consistent with plasticity- or mutation-led evolution when the environmental change is large or small, respectively. We then demonstrate that a breakdown in robustness due to large environmental changes induces adaptive plastic responses and uncovers cryptic mutations in the Full model. We show that developmental bias, the skew in phenotype distribution induced by environmental changes, precedes the onset of PLE and is already aligned with environmental changes.

## The Full model exhibits plasticity-led evolution under large environmental changes

In this subsection, we briefly summarize our previous results [11] to introduce fundamental notions describing PLE and MLE. We simulated evolution under large environmental changes for ten epochs, flipping 50% of the elements of the environment vector at the end of each epoch. We visualized evolutionary trajectories using the genotype-phenotype plots (Fig. 1 **a,b**) [11]. Phenotypic values of 0 and 1 correspond to phenotypes perfectly adapted to the ancestral and novel environments, respectively. Similarly, genotypic values of 0 and 1 correspond to the average genotypes adapted to the ancestral and novel environments. Generally, the evolution proceeds from the lower left corner to the upper right corner of the genotype-phenotype plot.

For the Full model (Fig. 1 **a**), phenotypic values are greater in the novel environment than in the ancestral environment on average, indicating an adaptive plastic

Figure 1: **Evolution under large environmental changes.** A large environmental change is modeled by flipping 50% of the ancestral environment vector **e**. The genotype-phenotype plot of an arbitrary epoch of the (**a**) Full and (**b**) NoCue model. Trajectories of (**c**) Mismatch and (**d**) Genetic Variance for ten different consecutive epochs. These results were reproduced from our previous work [11].

response. The standard deviation in the phenotypic values in the ancestral environment is minimal, indicating the robustness of adapted phenotypes. The standard deviation in the phenotypic values in the novel environment is much greater than that in the ancestral environment in the first generation, indicating the uncovering of cryptic mutations. The phenotypic and genotypic values in the novel environment increase somewhat quickly, indicating rapid genetic accommodation. These features of the genotype-phenotype plot assert that the Full model exhibits plasticity-led evolution under large environmental changes.

The NoCue model (Fig. 1 **b**) exhibits neither adaptive plastic response nor uncovering of cryptic mutations (by design). The cluttered distribution of points in the lower left corner demonstrates that the early adaptation stage of these trajectories is prolonged. The sparse distribution of points around the center of the plot indicates a rapid change in genotype once adaptive mutations are found.

We next tracked the trajectories of mismatch (Fig. 1**c**). Here, the mismatch

measures adaptedness using the distance between the phenotype and environment (Recall that the environment-as-selector is the optimal phenotype in our models). The mismatch of the Full model decreases immediately and rapidly. In contrast, the NoCue model exhibits a delay of around 20–30 generations before the mismatch decreases rapidly, a manifestation of gradualism.

Finally, we checked the trajectories of genetic variance (Fig. 1**d**). Here, the genetic variance is computed as the sum of the variance in each element of the genome matrices. For the Full model, the genetic variance decreases immediately and rapidly to a minimum before gradually increasing again. The initial rapid decrease corresponds to a purifying selection of uncovered cryptic variations, whereas the later increase indicates the accumulation of cryptic mutations. This behavior contrasts the NoCue model, for which the genetic variance gradually increases at the start of each epoch before decreasing to a minimum. The initial increase in the genetic variance coincides with the initial delay of mismatch (Fig. 1**c**), which is attributed to the time taken to find adaptive mutations in the novel environment. We identify these initial behaviors of the NoCue model as a key signature of MLE.

## The Full model exhibits mutation-led evolution under small environmental changes

We simulated evolution under small environmental changes for ten epochs, flipping 5% of the elements of the environment vector at the end of each epoch. The evolutionary trajectories of the Full and NoCue models (Figs. 2 **a,b**) are nearly identical under small environmental changes. That is, the Full model (Fig. 2 **a**) exhibits neither adaptive plastic response nor uncovering of cryptic mutations. The NoCue model exhibits similar evolutionary trajectories under small (Fig. 2 **b**) and large (Fig. 1 **b**) environmental changes. Therefore, under small environmental changes, both the Full and NoCue models exhibit the characteristics of MLE. This observation is consolidated by the trajectories of mismatch and genetic variance (Fig. 2 **c,d**).

Figure 2: **Evolution under small environmental change.** A small environmental change is modeled by flipping 5% of the ancestral environment vector **e**. Genotype-Phenotype plots in an arbitrary epoch of the **(a)** Full and **(b)** NoCue model. Trajectories of **(c)** Mismatch and **(d)** Genetic Variance for ten different consecutive epochs.

## The Full model undergoes a phase transition from MLE to PLE

We have seen that the Full model exhibits PLE or MLE under large or small environmental changes. This observation suggests a transition from MLE to PLE as environmental change increases. To see how this transition occurs, we examined the mean projected phenotype (Fig. **3a**) in the first generation after environmental changes of different magnitudes. The mean projected phenotypic value suddenly increases (indicating adaptive plastic response) at 30% environmental change. The variance in projected phenotype (Fig. **3b**) also suddenly increases (indicating the uncovering of cryptic mutations) at 30% environmental change.

To further analyze the phenotypic variation under various environmental changes, we examined the cross-covariance matrix between phenotype and environmental cue (Pheno-Cue cross-covariance) or mutations (Pheno-Geno cross-covariance). These cross-covariance matrices quantify how phenotypic variation is affected by environ-

Figure 3: **Transition from MLE to PLE**. **(a)** Projected phenotype in the first generation. **(b)** Variance in projected phenotype in the first generation. **(c)** First singular value of phenotype-environmental cue (Pheno-Cue) cross-covariance matrix. **(d)** First singular value of phenotype-genome (Pheno-Geno) cross-covariance matrix.

mental noise or mutations. In particular, we only consider the first singular value of the Pheno-Cue (Fig. 3**c**) and Pheno-Geno (Fig. 3**d**) cross-covariance matrices as it explains most phenotypic variation. We found that the first singular values also suddenly increase at 30% environmental change. This observation indicates that the phenotypes become susceptible to environmental noise or mutations at this transition point, suggesting a breakdown of environmental and mutational robustness.

## Developmental bias precedes PLE

The cross-covariance matrices introduced in the previous subsection contain various information about phenotype distribution, particularly the extent of phenotypic diversity and the skew in phenotype distribution [19]. There, we studied the extent of phenotypic diversity measured by the first singular values. We now study the skew in phenotype distribution, also called the developmental bias [20]. In the previ-

Figure 4: **Developemental bias under various environmental changes.** Percentage contribution of the first singular value of the Pheno-Cue **(a)** and Pheno-Geno **(b)** cross-covariance matrices. Alignment (normalized dot product) between environmental change and the first phenotype singular vector of the Pheno-Cue **(c)** and Pheno-Geno **(d)** cross-covariance matrices.

ous work, we observed large developmental biases aligned with large environmental changes, whereas such biases were absent in adapted environments [11].

To study the transition in developmental bias, we measured the proportion of the first singular value of the Pheno-Cue (Fig. 4**a**) and Pheno-Geno (Fig. 4**b**) cross-covariance matrices over environmental changes of different magnitudes. There is an increase in both proportions at around 25% environmental change. Compared to the transition from MLE to PLE (Fig 3), this increase in bias is less abrupt and occurs at smaller environmental changes. Thus, the developmental bias is more susceptible to environmental changes than the extent of phenotypic diversity. Still, it does not manifest as adaptive plastic responses or increased phenotypic variance due to the system's robustness when the environmental changes are small. When the bias becomes sufficiently large, the system's robustness breaks down, as indicated by an abrupt increase in the cross-covariances (Fig 3**c,d**).

We next examined the orientation of the developmental bias. Specifically, we

measured the alignment between the environmental change and the first pheno-
type singular vector of the Pheno-Cue (Fig. 4**c**) or Pheno-Geno (Fig. 4**d**) cross-
covariance matrices over different magnitudes of environmental changes. The align-
ment is small for less than 25% environmental changes. Above 25% environmental
change, developmental bias is well-aligned with environmental change. Thus, the
emergent developmental bias facilitates selection by magnifying the difference be-
tween the most and least fit individuals. Note that the transition in alignment
also happens before the increase in phenotypic variance (see Fig 3**c,d**). Thus, the
population seems "ready" for rapid adaptive evolution before selection, although
plastic responses are invisible.

# Discussion

From an evolutionary perspective, the Full model exhibits PLE under large en-
vironmental changes and MLE under small environmental changes. From a de-
velopmental perspective, the Full model is susceptible to environmental noise and
mutations under large environmental changes but is robust under small environmen-
tal changes. When the Full model is susceptible to perturbations, adaptive plastic
phenotypes are induced, cryptic mutations are uncovered, and developmental bias
is aligned with environmental change, enabling PLE. When it is robust against
environmental and genetic fluctuations, phenotypic variation is suppressed, cryptic
mutations accumulate, and developmental bias is virtually absent, necessitating an
extensive search for favorable mutations to adapt to small environmental changes,
which leads to MLE. Thus, the developmental transition from a robust regime to
a susceptible one underlies the evolutionary transition from MLE to PLE.

The developmental transition from robust to susceptible regimes may be consid-
ered as the switching of an evolutionary capacitor from the "charging" (accumulat-
ing cryptic mutations) to "discharging" (purifying uncovered variations) states [21].
Indeed, evolutionary capacitance has been suggested to be a generic feature of the
GRNs, which are non-linear multi-body feedback systems with gene-environment
interactions [22]. The present work suggests that the switching of the evolutionary
capacitor also switches the evolutionary regimes from MLE to PLE.

The transition from robust to susceptible regimes over environmental changes

implies the existence of a force opposing robustness. According to the Theory of Facilitated Variation [23], robustness and *adaptability* are (apparently) competing properties of the developmental process. Here, adaptability means the ability of the developmental process to respond to environmental changes adaptively. In our results, adaptability manifests in two ways: (1) adaptive plastic response (Fig. 3) and (2) developmental bias favorably aligned with environmental change (Fig. 4). In the robust regime, the adaptive plastic response is not visible (Fig. 3), but the developmental bias is already aligned with environmental change (Fig. 4 **c,d**). As the magnitude of environmental change increases, the induced adaptability overwhelms the system's robustness, manifests as adaptive plastic responses, and triggers PLE (Fig. 3).

The adaptability of the Full model is a consequence of its learning capability, as it is mathematically equivalent to an artificial recurrent neural network [16, 17, 24, 25, 11]. Many studies attempted to explain adaptive plastic response in terms of this learning capability of GRNs [16, 17, 24, 26, 25]. However, these works assume that the novel environment should be sufficiently similar to one of the ancestral environments to exhibit adaptive plastic response ("developmental memory"). This type of plasticity is genetically encoded because it has been selected through evolution [10], and we still face the problem of gradualism if the novel environment is entirely different from any of the ancestral environments. Furthermore, with genetically encoded plasticity, the plastic responses are expected to be uniform across the population. This implies that there will not be a wide range of variations for selection to act upon [27], hence limiting the efficiency of genetic accommodation. In contrast, our results show that the GRNs (the Full model) can exhibit adaptive plastic responses even in entirely new environments (Fig. 3). Since the novel environments in our simulations have no correlations to any of the ancestral environments, such adaptive plastic responses cannot be (entirely) genetically encoded but must be an emergent property of the GRNs [11]. This type of plasticity may be called developmental plasticity instead of genetically encoded plasticity. Furthermore, the adaptive plastic response caused by developmental plasticity compromises the robustness of the developmental system to expose a wide range of variations under large environmental changes for selection to act upon, hence accelerating

genetic accommodation. In summary, the adaptability of GRNs can manifest not only as genetically encoded plasticity but also as developmental plasticity, but it is the latter plasticity that is the key to rapid adaptation to large environmental changes.

# Methods

Here, we summarize the developmental models and data analysis methods. For details, see our previous paper [11].

## Development

In our developmental GRN models, let $p(s)$ be the phenotype expressed at the $s$-th stage of development and $\tilde{p}(s)$ (and $v(s)$) be the exponential moving average (and variance) of the phenotype. To reflect the hierarchical regulation of GRN elements (such as epigenetic marks, RNAs, and proteins), we introduce layers of 200-dimensional vectors $f(s), g(s)$, and $h(s)$ to represent epigenetic marks, gene expression and higher-order complexes (such as proteins, supramolecular complexes, etc.), respectively. Thus, we assume the following mutually recursive equations:

$$
\begin{aligned}
f_i(s) &= \sigma_f \left( \sum_{j=1}^{200} G_{ij} g_j(s-1) + \sum_{j=1}^{200} E_{ij}(e_j - \tilde{p}_j(s-1)) \right), \\
g_i(s) &= \sigma_g \left( \sum_{j=1}^{200} F_{ij} f_j(s) \right), \\
h_i(s) &= \sigma_h \left( \sum_{j=1}^{200} H_{ij} g_j(s) + \sum_{j=1}^{200} J_{ij} h_j(s-1) \right), \\
p_i(s) &= \sigma_p \left( \sum_{j=1}^{200} P_{ij} h_j(s) \right), \\
\tilde{p}_i(s) &= \alpha p_i(s) + (1-\alpha)\tilde{p}_i(s-1), \\
v_i(s) &= (1-\alpha)\{v(s-1) + \alpha[\tilde{p}_i(s-1) - p_i(s)]^2.\}
\end{aligned}
\tag{1}
$$

where $E$ is a matrix that represents environmental regulation of epigenetic marks, $F$ is a matrix representing epigenetic regulation of gene expression levels, $G$ is a matrix representing genetic regulation of epigenetic marks, $H$ is a matrix representing

genetic regulation of higher-order complexes, $J$ is a matrix representing interactions among higher-order complexes, $P$ is a matrix representing regulation of the phenotype. The matrix ensemble $E, F, G, H, J, P$ represents the individual's genome. All matrices are sparse with density 0.02. $\sigma_f, \sigma_g, \sigma_h, \sigma_p$ are activation functions [11]. The initial conditions are $f(0) = \mathbf{0}$, $g(0) = \mathbf{1}$, $h(0) = \mathbf{0}$ and $p(0) = \mathbf{0}$ where $\mathbf{0}$ is the zero vector and $\mathbf{1}$ is a vector where all elements are equal to 1. We recursively compute the state vectors $f(s), g(s), h(s), p(s)$ for $s = 1, 2, \cdots$ until the total phenotypic exponential moving variance $\sum_i v_i(s) < 10^{-5}$ for some $s \leq 200$ (i.e., the phenotype has converged) and take the adult phenotype as $\tilde{p}_i(s)$. Otherwise, we say that the phenotype does not converge.

The NoCue model is identical to the Full model, except it uses the following for $f_i(s)$ instead:

$$f_i(s) = \sigma_f \left( \sum_{j=1}^{200} G_{ij} g_j(s-1) + \sum_{j=1}^{200} E_{ij} \tilde{p}_j(s-1) \right). \tag{2}$$

# Singular Value Decomposition of Cross-Covariance Matrices

We use cross-covariance matrices to study the correlation between selected phenotypes and environmental noise or mutations. We define the Pheno-Cue or Pheno-Geno cross-covariance matrices as

$$C_{ij} = \frac{1}{N} \sum_{k=1}^{N} (p_{ik} - \overline{p}_i)(x_{jk} - \overline{x}_j) \tag{3}$$

where $p_{ik}$ is the $i$-th trait of the phenotype vector of the $k$-th individual, $\overline{p}_i$ is the population average of the $i$-th trait, $x_{jk}$ is the $j$-th environmental factor (for Pheno-Cue) or the $j$-th element of the vectorized genome (for Pheno-Geno) of the $k$-th individual and $\overline{x}_j$ is the corresponding population average value.

We apply singular value decomposition [28] to the cross-covariance matrix $C$ to obtain orthonormal components as follows [29]:

$$C = \sum_i \sigma_i u_i v_i^\top \tag{4}$$

where the superscript $\top$ indicates transpose. In Eq. (4), $u_i$ and $v_i$ are the $i$-th left and right singular vectors, interpreted as the $i$-th principal axis of phenotypic variation (the left singular vectors) in response to the corresponding principal axis (the right singular vectors) of environmental noises (or mutations); $\sigma_i$ is the $i$-th singular value, arranged in decreasing order, which represents the cross-covariance between the left and right principal components.

To quantify developmental bias, we used the proportion of the (squared) first singular value to the total cross-covariance:

$$\frac{\sigma_1^2}{\sum_i \sigma_i^2}.$$ (5)

The alignment between the first principal axis of phenotypic variation and environmental change is measured by the magnitude of the normalized dot product:

$$\frac{|u_1 \cdot (\mathbf{e}_n - \mathbf{e}_a)|}{\|\mathbf{e}_n - \mathbf{e}_a\|}$$ (6)

where $\mathbf{e}_n$ is the novel environment and $\mathbf{e}_a$ is the ancestral environment.

# Acknowledgments

The authors thank David Marshall and Daphne Teck Ching Lai for their helpful comments and Haziq Jamil for stimulating discussions. E.T.H.N. was supported by the UBD Bursary Award.

## Author contributions

A.R.K. conceived the models and ran preliminary simulations. E.T.H.N. ran large-scale simulations and collected the data. E.T.H.N. and A.R.K. designed the work plan, wrote the computer programs, analyzed the data, and wrote the manuscript.

## Competing interests

The authors declare no competing interests.

# References

[1] West-Eberhard MJ (2003) *Developmental plasticity and evolution.* (Oxford University Press, New York).

[2] Levis NA, Pfennig DW (2016) Evaluating 'plasticity-first'evolution in nature: key criteria and empirical approaches. *Trends Ecol. Evol.* 31(7):563–574.

[3] Levis NA, Pfennig DW (2021) Innovation and diversification via plasticity-led evolution in *Phenotypic Plasticity & Evolution*, ed. Pfennig DW. (CRC Press, Boca Raton, FL), pp. 211–240.

[4] Pfennig DW (2021) *Phenotypic plasticity & evolution: causes, consequences, controversies.* (Taylor & Francis).

[5] Gilbert SF, Epel D (2009) *Ecological Developmental Biology: Integrating Epigenetics, Medicine, And Evolution.* (Sinauer Associates, Sunderland, MA).

[6] Laland K, et al. (2014) Does evolutionary theory need a rethink? *Nature* 514(7521):161–164.

[7] Ehrenreich IM, Pfennig DW (2016) Genetic assimilation: a review of its potential proximate causes and evolutionary consequences. *Ann. Bot.* 117(5):769–779.

[8] Nishikawa K, Kinjo AR (2014) Cooperation between phenotypic plasticity and genetic mutations can account for the cumulative selection in evolution. *Biophysics* 10:99–108.

[9] Nishikawa K, Kinjo AR (2018) Mechanism of evolution by genetic assimilation: Equivalence and independence of genetic mutation and epigenetic modulation in phenotypic expression. *Biophys. Rev.* 10(2):667–676.

[10] Parsons KJ, McWhinnie K, Pilakouta N, Walker L (2020) Does phenotypic plasticity initiate developmental bias? *Evol. Dev* 22(1-2):56–70.

[11] Ng ETH, Kinjo AR (2023) Plasticity-led evolution as an intrinsic property of developmental gene regulatory networks (bioRxiv doi: 10.1101/2023.05.25.542372).

[12] Lande R (2009) Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. *J. Evol. Biol.* 22(7):1435–1446.

[13] Scheiner SM, Levis NA (2021) The loss of phenotypic plasticity via natural selection: genetic assimilation in *Phenotypic Plasticity & Evolution*, ed. Pfennig DW. (CRC Press, Boca Raton, FL), pp. 161–181.

[14] Wagner A (1996) Does evolutionary plasticity evolve? *Evolution* 50(3):1008–1023.

[15] Espinosa-Soto C, Martin OC, Wagner A (2011) Phenotypic robustness can increase phenotypic variability after nongenetic perturbations in gene regulatory circuits. *J. Evol. Biol.* 24(6):1284–1297.

[16] Watson RA, Wagner GP, Pavlicev M, Weinreich DM, Mills R (2014) The evolution of phenotypic correlations and "developmental memory". *Evolution* 68(4):1124–1138.

[17] Watson RA, Szathmáry E (2016) How can evolution learn? *Trends Ecol. Evol.* 31(2):147–157.

[18] Mitchell M (1998) *An introduction to genetic algorithms.* (MIT press, Cambridge, Massachusetts).

[19] Noble DW, Radersma R, Uller T (2019) Plastic responses to novel environments are biased towards phenotype dimensions with high additive genetic variation. *Proc. Natl. Acad. Sci. USA* 116(27):13452–13461.

[20] Uller T, Moczek AP, Watson RA, Brakefield PM, Laland KN (2018) Developmental bias and evolution: A regulatory network perspective. *Genetics* 209(4):949–966.

[21] Masel J (2013) Q&A: evolutionary capacitance. *BMC Biol.* 11(1):1–4.

[22] Bergman A, Siegal ML (2003) Evolutionary capacitance as a general feature of complex gene networks. *Nature* 424(6948):549–552.

[23] Gerhart J, Kirschner M (2007) The theory of facilitated variation. *Proc. Natl. Acad. Sci. USA* 104(suppl 1):8582–8589.

[24] Kouvaris K, Clune J, Kounios L, Brede M, Watson RA (2017) How evolution learns to generalise: Using the principles of learning theory to understand the evolution of developmental organisation. *PLoS Comput. Biol.* 13(4):e1005358.

[25] Ng ETH, Kinjo AR (2022) Computational modelling of plasticity-led evolution. *Biophys. Rev.* 14:1359–1367.

[26] Szilágyi A, Szabó P, Santos M, Szathmáry E (2020) Phenotypes to remember: Evolutionary developmental memory capacity and robustness. *PLoS Comput. Biol.* 16(11):e1008425.

[27] Riley CL, Oostra V, Plaistow SJ (2023) Does the definition of a novel environment affect the ability to detect cryptic genetic variation? *J Evol. Biol.*

[28] Yanai H, Takeuchi K, Takane Y (2011) *Projection Matrices, Generalized Inverse Matrices, and Singular Value Decomposition.* (Springer, New York, NY).

[29] Navarra A, Simoncini V (2010) *A Guide to Empirical Orthogonal Functions for Climate Data Analysis.* (Springer Netherlands, Dordrecht), pp. 97–106.