1
2   **Distinct functions of the kinesin-9 family in mammalian motile cilia**
3

Chuyu Fang[1,2,†], Xinwen Pan[1,3,†], Di Li[4,†], Yawen Chen[1,2], Luan Li[1,2], Qi Gao[1,2], Dong Li[4,*],

Xueliang Zhu[1,2,3,5,*] and Xiumin Yan[6,*]

[1] State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology,

Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, 320 Yueyang

Road, Shanghai 200031, China

[2] University of Chinese Academy of Sciences, Beijing 100049, China

[3] School of Life Science and Technology, ShanghaiTech University, China

[4] National Laboratory of Biomacromolecules, CAS Center for Excellence in

Biomacromolecules, Institute of Biophysics, Chinese Academy of Sciences, Beijing, China

[5] School of Life Science, Hangzhou Institute for Advanced Study, University of Chinese

Academy of Sciences, Hangzhou 310024, China

[6] Ministry of Education-Shanghai Key Laboratory of Children's Environmental Health,

Xinhua Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200092,

China

[†] Co-first author

* Corresponding author (e-mail: lidong@ibp.ac.cn, xlzhu@sibcb.ac.cn or

yanx@shsmu.edu.cn)

Abstract

Motile cilia, with 9+2 axonemes, propel directional fluid flows by coordinated beating

for various physiological functions. The kinesin-9 family comprises Kif6 and Kif9, known for

their involvement in regulating flagella motility across species. However, the precise roles of

mammalian kinesin-9 family remain unclear. In this study, we reveal the distinct localizations

and functions of Kif6 and Kif9 in mammalian motile cilia. Kif6 associates with intraflagellar

transport (IFT) particles and exhibits bidirectional movements within motile cilia. Knockout

of *Kif6* results in severe hydrocephalus due to impaired planar polarity and directional fluid

flow in the ependyma. Conversely, Kif9 localizes to the axonemal central lumen, and its

depletion causes brain ventricular expansion and defective sperm motility. These findings

shed light on the diverse functions of the kinesin-9 family in motile cilia and flagella,

providing valuable insights into the complex regulatory mechanisms underlying ciliary

trafficking and motility.

Keywords

Kinesin; intraflagellar transport; ependymal cells; ciliary motility; planar cell polarity; Kif6;

Kif9; hydrocephalus; basal body

Introduction

Cilia are evolutionarily conserved, microtubule (MT)-based organelles that protrude from the cell surface and are generally categorized into primary and motile cilia (Klena and Pigino, 2022). Primary cilia, with a 9+0 MT arrangement, are present in most mammalian cell types and serve as an 'antenna' to sense environmental cues such as mechanical, chemical, and light signals (Goetz and Anderson, 2010; Nachury and Mick, 2019). On the other hand, motile cilia/flagella typically possess a 9+2 MT axoneme with a central pair (CP) of MT singlets (C1 and C2) surrounded by peripheral nine doublet MTs (DMTs) (Klena and Pigino, 2022). The CP MTs are decorated with proteinaceous projections and the peripheral DMTs are equipped with dynein arms and radial spokes. The signal from CP apparatus transmits through radial spokes to govern the sliding of axonemal dyneins along adjacent MTs, generating cilia/flagella beating(Hoek et al., 2022; Lechtreck et al., 2009). With the exception of sperm flagella, motile cilia are mainly present as multicilia with dozens to hundreds of cilia within a single cell in mammals. Multicilia are found in epithelia of various organs such as the brain ventricles, the airway and the oviduct, playing critical roles in cerebrospinal fluid (CSF) circulation, mucus clearance, and ovum transport (Marshall and Kintner, 2008; Ohata and Alvarez-Buylla, 2016).

To generate directional fluid flow, proper polarities are needed to establish at both cellular and tissue levels. During the differentiation of multiciliated cells, basal bodies (BBs) initially amplify and dock randomly to the cell membrane, as indicated by the orientiation of their basal feet (BFs), which align with the direction of effective strokes (Gibbons, 1961; Guirao et al., 2010; Ohata and Alvarez-Buylla, 2016). As maturation progress, the orientation of BBs within individual cells gradually aligns in a coordinated manner along the direction of fluid flow, establishing rotational polarity (Boutin et al., 2014; Guirao et al., 2010; Werner et al., 2011). In ependymal cells, in addition to rotational polarity, BBs exhibit a specific clustering into a patch on one side of the cell surface, establishing translational polarity that distinguishes them from other multiciliated cells such as tracheal epithelial cells (Mirzadeh et

66 al., 2010). These processes ultimately establish the planar polarity of multiciliated tissues. The

67 establishment of planar polarity involves planar cell polarity (PCP) pathway and

68 hydrodynamic force flow, guiding the process over several days (Boutin et al., 2014; Guirao

69 et al., 2010; Ohata and Alvarez-Buylla, 2016).

70  Kinesins are motor proteins that play essential roles in directional transport of cargos

71 along MTs and regulation of MT dynamics in various cellular processes such as mitosis,

72 ciliogenesis and intracellular transport across eukaryotes (Cason and Holzbaur, 2022;

73 Hirokawa et al., 2009). They are classified into N-kinesins, M-kinesins, and C-kinesins based

74 on the position of motor domain. Two kinesin families, kinesin-2 and 9, have been found to

75 function in cilia or flagella. Among them, the kinesin-2 family motors (Kif3a/b/c) are well-

76 known for their involvement in IFT, a specialized bidirectional transportation machinery

77 (Bhogaraju et al., 2013; Scholey, 2013; Taschner et al., 2016; Zhao et al., 2012). IFT relies on

78 two multi-subunit complexes, IFT-A and IFT-B, which are powered by kinesin-2 or

79 cytoplasmic dynein-2 respectively for anterograde or retrograde trafficking along the

80 axonemal MTs. This process enables the transport of ciliary proteins in and out of cilia,

81 critical for cilia assembly, maintenance, and function (Jordan et al., 2018; Klena and Pigino,

82 2022; Meleppattu et al., 2022; Pazour et al., 1998; Pigino, 2021).

83  The kinesin-9 family consists of two members, Kif6 and Kif9, which share a high

84 sequence identity at their motor domains. They belong to the N-kinesin group, typically

85 involved in plus-end-directed transports of cargos. Unlike the kinesin-2 family, which plays a

86 critical role in both primary and motile cilia, the kinesin-9 family is predominantly associated

87 with flagellated species (Demonchy et al., 2009), suggesting distinct or redundant roles in

88 regulating flagella motility. In *Chlamydomonas*, only KLP1, the homolog of Kif9, has been

89 studied. It specifically localizes to the CP MTs and is proposed as an active motor to regulate

90 flagellar motility (Bernstein et al., 1994; Han et al., 2022; Yokoyama et al., 2004).

91 Interestingly, in *Trypanosoma*, the kinesin-9 family has been demonstrated distinct functions

92 in flagella motility (Demonchy et al., 2009). Depletion of *KIF9A*, the *Trypanosoma* homolog

93 of *Kif9*, reduces flagellum beating and cell movement, while knockdown of *KIF9B*, the

homolog of *Kif6*, results in cell paralysis and defective assembly of the paraflagellar rod

(PFR). The PFR, a unique large structure found exclusively in euglenoids and kinetoplastids,

is asymmetrically attached to the axoneme at DMTs 4-7, critical for flagella beating and cell

locomotion (Demonchy et al., 2009; Portman and Gull, 2010; Zhang et al., 2021). Although

there is no observed movement of KIF9B within flagella, it is speculated that KIF9B is

involved in PFR transport or the recognition of axoneme asymmetry for PFR assembly in

Trypanosoma (Demonchy et al., 2009). This raises the question of whether Kif6, in other

species, also contributes to ciliary trafficking of essential materials involved in cilia beating

and recognition or establishment of axonemal polarity. In mammals, mutations in *Kif9* have

been associated with defective sperm motility and impaired fertility (Meng et al., 2022;

Miyata et al., 2020). A frameshift mutation of *KIF6* has been found in a patient with

intellectual disability and megalocephaly (Buchan et al., 2014; Konjikusic et al., 2018). Mice

carrying a *Kif6* C-terminal truncation mutant (*Kif6^{p.G555fs}*) develop severe hydrocephalus

possibly due to defective ciliogenesis in the ependyma (Buchan et al., 2014; Konjikusic et al.,

2018). Despite these achievements, a comprehensive understanding of the roles of Kif6 and

Kif9 in mammalian ciliary trafficking and cilia motility is still lacking.

In this study, we investigate the functions of Kif6 and Kif9 in mice. Kif6 is a motile-cilia

specific kinesin and plays a critical role in ciliary transportation, essential for the

establishment of planar polarization in the ependyma, while Kif9 likely functions within the

CP projection for the regulation of motile cilia/flagella motility.

Results

Mammalian Kif6 and Kif9 are motile cilia-specific kinesins with distinct localizations

Murine full-length Kif6 (NP_796026.2) and Kif9 (NP_001157041.1) are highly

conserved proteins composed of 802 and 810 amino acid residues, respectively. Structurally,

both proteins have a motor domain located at the N-terminus, followed by two coiled-coil

regions and a tail at the C-terminus (Fig. 1a) (Hirokawa et al., 2009; Miki et al., 2005). The

motor domains of the two proteins share ~43% sequence identity, while their C-terminus are

less conserved (Fig.1a and Supplementary Fig. 1a, b) (Miki et al., 2005). Phylogenetic

122 analysis showed Kif6 and Kif9 co-existed in eukaryotes with motile cilia/flagella, dating back

123 to protists like *Chlamydomonas* (Fig. 1b and Supplementary Fig. 1b) (Bernstein et al., 1994;

124 Vincensini et al., 2011; Yokoyama et al., 2004). In constrast, Kif6 and Kif9 homologues are

125 absent in *Caenorhabditis elegans* which only contain immotile sensory cilia (Fig. 1b) (Inglis

126 et al., 2007; Vincensini et al., 2011). These data suggest that Kif6 and Kif9 may have

127 important roles not only in flagella but also in motile cilia.

128   Analysis on our previous microarray results (Xu et al., 2015) demonstrated that *Kif6* and

129 *Kif9* were highly upregulated during the multiciliation of mouse tracheal epithelial cells

130 (mTECs) (Fig. 1c). Immunoblots confirmed a substantial increase in the protein levels of Kif6

131 and Kif9 during multiciliation of mTECs (Fig. 1d). Furthermore, Kif6 and Kif9 were

132 specifically expressed in tissues and cells abundant in motile cilia, such as the ependyma, the

133 trachea, the lung, the testis, and primary cultured mouse ependymal cells (mEPCs), but were

134 hardly detected in primary ciliated tissues and cells (Fig. 1e, f) (Duan et al., 2021; Fliegauf et

135 al., 2007). To visualize their subcellular localizations, we immunostained mEPCs and IMCD3

136 cells with antibodies against Kif6 or Kif9 (Fig. 1g-i). Confocal analysis revealed that both

137 Kif6 and Kif9 localized to the axoneme of motile cilia in mEPCs (Fig. 1h), but not to the

138 axoneme of primary cilia in IMCD3 cells (Fig. 1i). To validate their ciliary localization, we

139 infected mEPCs with adenovirus to express Kif6-GFP or Kif9-GFP (Fig. 1g, j). Consistently,

140 Kif6-GFP and Kif9-GFP were clearly observed in ciliary axonemes of mEPCs (Fig. 1j).

141   Super-resolution microscopy analysis revealed that ciliary Kif9 was distributed at the CP

142 region, when Hydin, a C2 projection protein (Lechtreck et al., 2008), and acetylated tubulin

143 (Ac-tub) were respecisvely used as CP and axoneme markers (Fig. 1k), suggesting a

144 conserved localization in the C2 projection as *Chlamydomonas* KLP1 (Han et al., 2022). In

145 sharp contrast, Kif6 localized along the axonemes as puncta (Fig. 1l). Similar ciliary

146 localization patterns were observed when GFP-tagged Kif6 and Kif9 were expressed in

147 mEPCs and co-immunostained with Spef1, another CP protein (Supplementary Fig. 2a, b)

148 (Zheng et al., 2019).

Collectively, our results demonstrate the specific localization of Kif6 and Kif9 within

motile cilia. Kif6 localizes as punta along the peripheral DMTs of axonemes, whereas Kif9 is

specifically enriched in the CP lumen. Their distinct localization patterns imply potential

unique roles for the two kinesins in mammalian motile cilia.

Kif6, but not Kif9, moves along ciliary axoneme and colocalizes with IFT particles

As kinesins with a motor domain at their N-terminus typically function as plus-end-

directed motor proteins (Cason and Holzbaur, 2022; Hirokawa et al., 2009), we examined

whether Kif6 and Kif9 could move along DMTs and central MTs, respectively. We employed

the high spatiotemporal resolution, grazing incidence structured illumination microscopy (GI-

SIM) (Guo et al., 2018; Qiao et al., 2022) to record the movements of GFP-tagged Kif6 and

Kif9 in mEPCs (Fig. 2a-e and Movies 1-3). In these experiments, we used one IFT-B

component Ift81-GFP as a positive control (Kanie et al., 2017). Immunoblotting demonstrated

a comparable expression level of Kif6-GFP and Ift81-GFP with their endogenous proteins,

while Kif9-GFP showed higher expression than endogenous Kif9 (Fig. 2b). As expected,

Ift81-GFP particles displayed clear bidirectional movements along the axonemes (Fig. 2c and

Supplementary Fig. 3a, top; Movie 1). Strikingly, Kif6-GFP particles also underwent robust

and processive movements in both the anterograde and retrograde directions along the

axonemes, resembling the movement of Ift81-GFP particles (Fig. 2d and Supplementary Fig.

3a, middle; Movie 2). In contrast, we did not detect any processive movement of Kif9-GFP

along the entire axonemes (150 cilia from 10 cells) (Fig. 2e and Supplementary Fig. 3a,

bottom; Movie 3).

Qunatification results revealed that Ift81-GFP moved at velocities of 0.73 ± 0.40 μm/s

for anterograde and 0.67 ± 0.36 μm/s for retrograde (Fig. 2f). The velocities were slightly

faster than the IFT in primary ciliated IMCD3 cells and the olfactory sensory neurons but

slower than those in *C. elegans* and *Chlamydomonas* (Williams et al., 2014; Wingfield et al.,

2017). Kif6-GFP labeled puncta moved at similar velocities to Ift81-GFP, with the mean

velocities of 0.71 ± 0.34 μm/s for anterograde and 0.64 ± 0.34 μm/s for retrograde (Fig. 2f)

transports. The frequency of anterograde movement for Kif6-GFP particles (13.3 ± 11.3

counts/min) was similar to that of Ift81-GFP particles (14.0 ± 10.0 counts/min). However, the

retrograde frequency of Kif6-GFP puncta appeared slightly decreased compared to Ift81-GFP

particles (11.0 ± 9.0 vs 13.2 ± 9.9 counts/min) (Fig. 2g), suggesting that a portion of Kif6 may

diffuse back to the ciliary base like kinesin-2 after it reached to ciliary tip (Chien et al., 2017).

Since Kif6 exhibited comparable velocity and frequency characteristics to those of IFT

particles, we expressed Kif6-GFP in mEPCs and co-immunostained with antibodies against

two components of IFT-B complexes (Ift56 and Ift88) to further explore their relationship

(Kanie et al., 2017; Taschner et al., 2014). Confocal imaging unveiled a notable colocalization

between Ift56 and Kif6-GFP at the axoneme (Fig. 2h). Three-dimensional structured

illumination microscopy (3D-SIM) images further showed their punctate colocalization along

the axonemal DMTs (Fig. 2i). Similar results were observed when Kif6-GFP was co-

immunostained with Ift88 (Supplementary Fig. 3b, c).

Taken together, our findings demonstrate the bidirectional movements of Kif6 within

motile cilia, which is not observed for Kif9. The similarity in movement behaviors,

velocities and frequencies between Kif6 and IFT particles strongly suggest a potential

involvement of Kif6 in IFT trafficking.

Kif6-GFP puncta exhibit comovement with IFT particles along axonemal MTs

To gain further insights into the relationship between Kif6 and IFT particles, we co-

expressed Kif6-GFP and Ift27-Halo, another IFT-B component (Kanie et al., 2017), in

mEPCs by adenoviral infection and performed dual-color time-lapse GI-SIM imaging (Fig.

3). Before imaging, a calibration of the microscope was performed routinely with fluorescent

beads, ensuring that different fluorescent channels were acculartely aligned in GI-SIM

(Supplementary Fig. 3d). Interestingly, we observed three distinct bidirectional moving

particles within cilia: particles labeled with both Ift27-Halo and Kif6-GFP (Ift27[+]Kif6[+]

particles), those labeled only with Kif6-GFP (Kif6[+] particles), and those labeled only with

Ift27-Halo (Ift27[+] particles) (Fig. 3a-c; Movies 4-5). The corresponding komographs

confirmed the presence of these three types of particles (Fig. 3a-c and Movies 4-5). The ratio

of these three trafficking types were approximately 1:1.5:1 (Ift27$^+$Kif6$^+$:Kif6$^+$:Ift27$^+$) in both

anterograde and retrograde directions (45 cilia in total from 4 cells) (Fig. 3d). These particle

types exhibited similar velocities for both anterograde and retrograde transports (Fig. 3e and

Movies 4-5). Specifically, Ift27$^+$ particles showed velocities of 0.70 ± 0.31 μm/s for

anterograde and 0.62 ± 0.28 μm/s for retrograde transport. Kif6$^+$ particles moved at velocities

of 0.74 ± 0.32 μm/s and 0.65 ± 0.34 μm/s, respectively. Ift27$^+$Kif6$^+$ particles moved at

velocities of 0.75 ± 0.32 μm/s and 0.66 ± 0.28 μm/s, respectively (Fig. 3e).

Taken together, the distribution of Kif6 observed in fixed samples and time-lapse

experiments indicate that Kif6, unlike Kif9, is not a structural component of motile cilia,

functions as an active motor in ciliary trafficking. The presence of IFT-free Kif6 particles

suggests independent movements of Kif6 along the cilia.

Mice lacking *Kif6* or *Kif9* develop hydrocephalus and exhibit male infertility

Function of Kif6 has been explored using *Kif6$^{p.G555fs}$* mice which develop severe

hydrocephalus (Konjikusic et al., 2018). Since the mutant mice still express the motor domain

of Kif6, the observed hydrocephalus phenotype could be caused by the dominant negative

effect of the remaining motor domain. Additionally, depletion of *Kif9* results in defective

sperm motility in mice, with its function in motile cilia remaining unexplored (Miyata et al.,

2020). To uncover their physiological function *in vivo* and the mechanism involved, we

generated *Kif6* and *Kif9* knockout mice (Fig. 4a, b) using the CRISPR/Cas9 system (Joung et

al., 2017). Immunoblotting confirmed complete depletion of Kif6 or Kif9 (Fig. 4c). Both *Kif6^{-}*

$^{/-}$ and *Kif9$^{-/-}$* mice were born at the expected Mendelian ratio of genotypes. The majority of

Kif6$^{-/-}$ mice experienced growth failure and often exhibited a dome-shaped skull compared to

wild-type littermates after P21, suggestive of severe hydrocephalus (Fig. 4d, indicated by

arrows). Some of these mice displayed more pronounced cranial deformities and skull

expansion, leading to a forward rounding of the back or kyphosis with a smaller stature

compared to their littermates (Fig. 4d, indicated by arrowhead). Careful examination of brain

dissections or coronal sections revealed that every analyzed *Kif6$^{-/-}$* mouse (100%; total n = 60)

displayed lateral ventricle expansion when compared to the wild-type controls (Fig. 4f, h).

This ventricle enlargement was evident even in mice that displayed a normal skull appearance

and longer lifespan (Fig. 4f, bottom). In sharp contrast to *Kif6*[-/-] mice, *Kif9*[-/-] mice appeared

normal similar to their wild-type littermates based on macroscopic examination (Fig. 4e).

However, coronal brain sections revealed enlarged ventricles in over half of *Kif9*[-/-] mice (Fig.

4g, h), indicative of hydrocephalus. Survival curve analysis revealed that approximately 77%

of *Kif6*[-/-] mice died before reaching P90, while *Kif9*[-/-] mice exhibited a normal lifespan similar

to their wild-type littermates during a 12-month observation period (Fig. 4i).

We did not observed noticeable defects in organs where primary cilia are critical, such as

polydactyly or polycystic kidney (Duan et al., 2021; Fliegauf et al., 2007), suggesting that

primary cilia might not be affected. Consistent with the previous study, although *Kif9*[-/-] male

mice displayed a normal sperm count. However, there was a decline in sperm motility and

poor male fertility (Movie 6), similar to previous observation (Miyata et al., 2020). Contrarily,

Kif6[-/-] mice that reached reproductive age (~ 2 months) showed a significant reduction in

sperm count. Notably, while the sperm of *Kif6*-deficiency mice had normal morphology, they

exhibited almost no motility (Fig. 4j, k and Movie 7). These observations emphasize the

critical but distinct roles of Kif6 and Kif9 in motile cilia and sperm flagella.

***Kif6* but not *Kif9* deficiency disrupts the planar polarity of ependymal motile cilia**

Ependymal multiciliated cells are responsible for generating unidirectional CSF flow

through planar-polarized cilia beating in the brain ventricle (Marshall and Kintner, 2008;

Ohata and Alvarez-Buylla, 2016). Abnormalities in cilia formation and motility of ependymal

cells cause hydrocephalus (Matsuo et al., 2013; Mirzadeh et al., 2010; Tissir et al., 2010).

Given that the knockout of *Kif6* and *Kif9* resulted in hydrocephalus, we thus assessed the

impact of Kif6 and Kif9 on ciliogenesis and cilia motility (Fig. 5a). As Kif6 was found to be

associated with IFT in ependymal cilia, we firstly employed scanning electron microscopy

(SEM) to determine if Kif6 is required for ciliogenesis. We examined whole-mounts of

ependyma from *Kif6*[-/-] mice and their wild-type littermates at P4, the time of cilia formation

(Spassky et al., 2005). Our SEM analysis revealed that both the number and morphology of

cilia in ependymal cells remained unaffected by *Kif6* deficiency (Fig. 5b, top). Simliar

observations were obtained when using mice at P30 (Fig. 5b, bottom), implying that depletion

of *Kif6* does not affect ciliogenesis, different from the defective ciliogenesis observed in

Kif6$^{p.G555fs}$ mice (Konjikusic et al., 2018). Moreover, transmission electron microscopy (TEM)

exhibited no discernible ultrastructure abnormalities in the ciliary axoneme of *Kif6*-deficient

mice (Supplementary Fig. 4a).

As ependymal cells display unidirectional beating at tissue level after P21 (Guirao et al.,

2010; Ohata and Alvarez-Buylla, 2016), to assess the effect of *Kif6* or *Kif9* knockout on

ciliary motility, we labeled the lateral ventricles derived from mice after P21 with SiR-

tubulin, a live-cell microtubule fluorescent probe (Lukinavicius et al., 2014) and conducted

live-cell imaging using high-speed spinning disk confocal microscopy (Fig. 5a). A ciliary beat

cycle typically consists of a quick effective stroke and a slower recovery stroke (Lechtreck et

al., 2009). To clarify the ciliary beat pattern, we traced and analyzed the movement of cilia

from both side and top views. Strikingly, we observed that cilia in individual *Kif6$^{-/-}$*

ependymal cells, regardless of the severity of hydrocephalus, exhibited metachronal

synchrony, beating in a back-and-forth asymmetric manner, resembling the behavior seen in

wild-type littermates (Fig. 5c and Supplementary Fig. 4b). Notably, the ciliary beat

frequencies of *Kif6$^{-/-}$* cilia (23.4 ± 6.3 Hz) were similar to those of wild-type counterparts

(23.9 ± 6.4 Hz) (Fig. 5d). In contrast, at the tissue level, wild-type ependymal cells exhibited

metachronal coordination, predominantly beating in one direction (Fig. 6e and Supplementary

Fig. 4c; Movie 8). However, the coordination of ciliary beat directions within *Kif6$^{-/-}$*

ependyma was significantly disrupted (Fig. 5e and Supplementary Fig. 4c; Movie 9),

indicating impaired PCP (Mirzadeh et al., 2010; Ohata and Alvarez-Buylla, 2016).

Similar to the loss of *Kif6*, depletion of *Kif9* did not significantly alter the number and

length of cilia. Additionally, individual ependymal cells from *Kif9$^{-/-}$* littermates exhibited

well-coordinated, asymmetric beating patterns (Fig. 5f and Supplementary Fig. 4d). In

contrast to *Kif6$^{-/-}$* ependyma (Fig. 5e and Supplementary Fig. 4c; Movie 9), multicilia in *Kif9$^{-/-}$*

ependyma displayed metachronal coordination at the tissue level, resembling the wild-type

controls (Fig. 5g and Supplementary Fig. 4e; Movies 10-11). Moveover, cilia beat frequencies

in $Kif9^{-/-}$ ependymal cilia remained unchanged compared to wild-type controls

(Supplementary Fig. 4f).

Given that the knockout of $Kif6$ resulted in uncoordinated ciliary beating across the

ependyma (Fig. 5h, i), we performed a beads assay to investigate whether disrupted cilia

beating in $Kif6^{-/-}$ mice leads to impaired fluid flow, ultimately contributing to the development

of hydrocephalus. Latex fluorescent microbeads were added to freshly prepared brain slices

and their movements were tracked to quantitatively assess dynamic fluid flow. In wide-type

samples, the beads displayed a relatively uniform drift in one direction (Fig. 5h; Movie 12).

Conversely, in $Kif6^{-/-}$ samples, the uncoordinated cilia beats mainly drove the beads to

oscillate or whirl regionally (Fig. 5h; Movie 13). The velocities of beads in $Kif6^{-/-}$ mice were

46.3 ± 15.4 μm/s slower than those in wild-type mice (63.6 ± 27.1 μm/s) (Fig. 5i).

Taken together, our results suggest that the loss of $Kif6$ and $Kif9$ does not affect

ciliogenesis. However, the absence of $Kif6$, but not $Kif9$, does impair the coodinated ciliary

beats at tissue-level which may contribute to the development of hydrocephalus.

Kif6 is crucial for the establishment of rotational polarity of BBs in ependymal tissues

Since the coordinated cilia beating closely relates with the orientation of BBs (Mirzadeh et

al., 2010; Ohata and Alvarez-Buylla, 2016), we examined the BB polarity in both $Kif6^{-/-}$ and

$Kif9^{-/-}$ ependyma. We immunostained whole-mounts of ependyma from wild type, $Kif6^{-/-}$ and

$Kif9^{-/-}$ mice using Cep164, a transition fiber component (Graser et al., 2007; Siller et al.,

2017), to mark BBs, and γ-tubulin or Centriolin to label BFs (Fig. 6a-e and Supplementary

Fig. 5) (Nguyen et al., 2020; Ohata et al., 2014). 3D-SIM images revealed a uniform

rotational polarity of BB patches and consistent tissue-level polarity in both wild type and

$Kif9^{-/-}$ wholemounts (Fig. 6b, c and Supplementary Fig. 5a, b). In contrast, both the rotational

and tissue polarity of BBs were markedly impaired in $Kif6$-deficient ependyma, regardless of

the severity of hydrocephalus (Fig. 6d, e and Supplemenatry Fig. 5c, d). To assess the extent

of the rotational polarity in individual cells, we measured the mean vector length by

considering each arrow marking the BF orientation as a unit vector (Bustamante-Marin et al.,

2019; Werner et al., 2011). As expected, our analysis revealed that the mean vector length of BBs in wild-type and $Kif9^{-/-}$ ependymal cells was close to 1 (0.95 ± 0.03 for wild type; 0.95 ± 0.04 for $Kif9^{-/-}$) (Fig. 6f). In contrast, the mean vector length of BBs was significantly reduced in $Kif6^{-/-}$ samples (0.32 ± 0.15 for mice with mild hydrocephalus; 0.28 ± 0.14 for mice with severe hydrocephalus) (Fig. 6f), suggesting an impaired rotational polarity of BBs. Similar quantitative results were obtained when used Centriolin as the BF marker (Supplementary Fig. 5e). TEM images further confirmed the proper alignment of BFs in wild-type ependyma (mean vector length = 0.97 ± 0.07) and the impaired rotational polarity of BBs in $Kif6^{-/-}$ ependyma, in both mild (0.35 ± 0.17) and severe (0.31 ± 0.18) hydrocephalus mice (Fig. 6g, h). Our results highlight the crucial role of Kif6, rather than Kif9, in the establishment of rotational polarity of BBs in ependyma, essential for coordinated cilia beating at the tissue level (Fig. 6i).

In contrast to the rotational polarity, BB patches displayed conspicuous asymmetric distributions within the cells of both mild and severe hydrocephalus $Kif6^{-/-}$ mice (Supplementary Fig. 6a-c). The area of BB patches also remained unaffected in $Kif6^{-/-}$ mice with a mild hydrocephalus, however, we did observe a slight increase in the BB patches area in samples with severe hydrocephalus (Supplementary Fig. 6d), potentially influenced by the pathological effects of hydrocephalus, suggesting that Kif6 may be dispensable for translational polarity (Ohata and Alvarez-Buylla, 2016). Notably, the depletion of $Kif6$ and $Kif9$ did not affect the number of BBs (Fig. 6j), consistent with unaffected ciliogenesis (Fig. 5b, e, g and Supplementary Fig. 4c, e).

Discussion

In this study, we show that Kif6 and Kif9, two kinesin-9 family members, have unique ciliary localizations and functions in mammalian motile cilia and flagella (Fig. 1b and 6k). Kif9 is mainly found in the CP lumen in motile cilia, while Kif6 localizes along the DMTs and can move bidirectionally along the ciliary axoneme (Fig. 1-3 and Supplementary Fig. 3). In knockout mice, neither Kif6 nor Kif9 are necessary for ciliogenesis (Fig. 5 and

Supplementary Fig. 4). Specifically, *Kif6* knockout leads to reduced sperm motility (Fig. 4k

and Movie 7) and disrupted cilia motility coordination (Fig. 5e and Supplementary Fig. 4c;

Movie 9), causing hydrocephalus (Fig. 3d, f, h) and male infertility. Interestingly, *Kif6*

depletion causes flagella paralysis in *Trypanosoma* (Demonchy et al., 2009) and impairs

sperm flagella motility in mice (Fig. 4k and Movie 7), but it does not impact the beating

frequency of motile cilia (Fig. 5d). *Kif9* knockout affects flagella motility with minimal

impact on motile cilia (Fig. 5 and Supplementary Fig. 4; Movie 6). These differences suggest

distinct regulatory mechanisms, potentially linked to different beating patterns like the back-

and-forth pattern of ependymal cilia, and the non-planar waveform pattern of sperm flagella

(Gadadhar et al., 2021; Lechtreck et al., 2009; Miyata et al., 2020).

The kinesin-9 family belong to N-kinesins, usually facilitate plus-end-directed cargo

transports (Cason and Holzbaur, 2022; Demonchy et al., 2009). Using GI-SIM, we find that

Kif6 but not Kif9 undergoes processive movement in ependymal cilia (Figs. 2-3 and

Supplementary Fig. 3; Movies 2-4). The result of mammalian Kif9 is consistent with the

observation in *Xenopus* epidermal multiciliated cells (Konjikusic et al., 2023), suggesting that

Kif9 is not involved in the ciliay transport along the cilia. Considering the conserved CP

localization of Kif9 and brain ventricle expansion phenotype observed in over half of *Kif9*[-/-]

mice (Fig. 4g, h), Kif9 may regulate ependymal cilia beating, potentially mediating

conformation changes of CP MTs during beating. Future studies such as cryo-electron

tomography technology (Cryo-ET) studies will be needed to investigate this hypothesis. Most

strikingly, we observe the bidirectional movement of Kif6 along the ciliary axoneme and

association of IFT particles (Figs. 2, 3 and and Supplementary Fig. 3). Although Kif6 is not

essential for ciliogenesis (Fig. 4 and Supplementary Fig. 4), it is the only known kinesin

specifically involved in the IFT process within motile cilia. Its co-movement with IFT

particles suggests Kif6 may serve as a cargo adaptor or as an active motor contributing to the

delivery of proteins critical for establishing planar polarity (Fig. 6k). Further investigations

are required to confirm this.

Kif6 deficiency does not affect translational polarity of BBs, but it does disrupt the

rotational polarity of BBs and planar polarity of cilia in ependyma (Figs. 5-7 and

Supplementary Figs. 4-5; Movie 9). Since PCP pathway is key for translational polarity

(Mirzadeh et al., 2010; Ohata and Alvarez-Buylla, 2016), Kif6 likely does not function

upstream of the PCP pathway. Instead, it may transport specific cargos like signalling

molecules within motile cilia that respond to hydrodynamic forces or downstream of PCP

signaling (Fig. 6k). This idea is supported by the misalignment of BBs in $Kif6^{-/-}$ mice, similar

to obervations in cultured mEPCs lacking external flow force (Guirao et al., 2010).

Furthermore, this speculation is also supported by $Kif6^{-/-}$ mice not showing obvious airway

defects, where the rotational polarity of BBs could align properly without flow forces *in vitro*

(Herawati et al., 2016). In *Trypanosoma*, the absence of *KIF9B*, a *Kif6* homolog, disrupts the

assembly of the PFR, a structure asymmetrically assembled at DMTs 4-7 along the flagellar

axoneme, required for flagella motility (Demonchy et al., 2009; Zhang et al., 2021).

Interestingly, in mammals, "9+2"-type motile cilia also exhibit an asymmetric axonemal

orientation. For instance, the CP plane is perpendicular to the direction of ciliary beat, and the

direction of BFs aligns with the effective stroke of cilia (Butler and Wallingford, 2017; Clare

et al., 2014; Lindemann and Lesich, 2021; Ohata and Alvarez-Buylla, 2016). The disturbance

of the rotational polarity of BBs in $Kif6^{-/-}$ ependymal cells suggests a role of Kif6 to help

ensure proper axonemal asymmetry and BB polarity. Notably, $Kif6^{-/-}$ sperm flagella are nearly

immobile, resembling the *KIF9B*-defective *Trypanosoma* flagella. As sperm flagella have

multiple extra-axonemal structures, further research into function of Kif6 in these structures

could provide interesting insights.

In summary, our findings from *Kif6* and *Kif9* knockout mice and GI-SIM experiments

have elucidated the crucial role of Kif6 in BB polarization and its necessity for coordinated

cilia beating, potentially through its involvement in motile cilia-specific transport.

Furthermore, bidirectional trafficking of Kif6 suggests that it may deliver materials critical for

coordinated cilia beating across ependymal tissues. Some Kif6 puncta appear independent of

IFT particles, hinting at separate protein transport from IFT, though further investigation is

needed. Identifying molecules or substructures associated with Kif6 could provide more

insight into the regulatory mechanism. Although *Kif9* depletion does not significantly impact

ciliary motility, its role in sperm motility and brain ventricle expansion suggests a critical

function in flagella/motile cilia motility.

Methods

Plasmids

The cDNAs encoding full-length *Kif6* (GenBank: NM_177052.3), *Kif9* (GenBank:

NM_001163569.1), *Ift81* (GenBank: NM_001358917.1), and *Ift27* (GenBank:

NM_025931.3) were amplified using PCR with cDNAs extracted from cultured mEPCs (Day

10) as the template. To generate adenoviral expression constructs, the cDNAs fragments were

PCR-amplified and subsequently cloned into the entry vector pYr-1.1-GFP (YRBio, China) or

pYr-1.1-Halo to express GFP-tagged or Halo-tagged proteins, respectively. The LR

recombination reactions between the entry constructs and the destination vector

pAd/BLOCK-iT-DEST (YRBio, China) were carried out using LR Clonase II enzyme mix

(Thermo Fisher, 11791020), as described previously (Zhao et al., 2021).

To generate polyclonal antibodies of Kif9, the cDNA fragments encoding the indicated

amino acids (348-810 aa) were amplified by PCR and subcloned into pET-28a or pGEX-4T-1,

to express His-tagged or GST-tagged Kif9, respectively.

All constructs were validated by sequencing. The primers used for PCR amplification are

listed in Supplementary Table 1.

Mice

All animal experiments were performed following guidelines approved by the Institutional

Animal Care and Use Committee of Institute of Biochemistry and Cell Biology. The mice

used in this study had a genetic background of C57BL/6J. $Kif6^{+/-}$ and $Kif9^{+/-}$ mice were

obtained from GemPharmatech, China. Ages of mice used in this study were indicated in the

figures or the legends. gRNA sequences used for targeting *Kif6* or *Kif9* and primers used for

genotyping are listed in Supplementary Table 1.

Cell culture and transfection

HEK293T and HEK293A cells were grown in Dulbecco's modified Eagle's medium

(DMEM) (Thermo Fisher, 12430-054) supplemented with 10% fetal bovine serum (FBS)

(Ausbian, VS500T), 0.3 mg/ml glutamine (Sigma, G8540), 100 U/ml penicillin (Solarbio,

P8420) and 100 U/ml streptomycin (Solarbio, S8290). NIH3T3 cells were cultured in DMEM

medium with 10% normal goat serum (NGS), 0.3 mg/ml glutamine, 100 U/ml penicillin, and

100 U/ml streptomycin. hTERT-RPE1 and IMCD3 cells were cultured in DMEM/F12

medium (GE Healthcare) supplemented with 10% FBS, 0.3 mg/ml glutamine, 100 U/ml

penicillin, and 100 U/ml streptomycin. Additionally, RPE1 culture medium was supplemented

with 10 μg/ml hygromycin B (Thermo Fisher). All cell lines were passaged at a 1:4 ratio once

they reached 80%-90% confluency. HEK293A cells were used for packaging adenovirus

expressing GFP-tagged proteins following the guidelines (YRBio, China). All the cells were

routinely tested for mycoplasma contamination.

Primary culture of mouse ependymal cells (mEPCs) was prepared and cultured as

previously described (Delgehyr et al., 2015; Zhao et al., 2021). Newborn C57BL/6J mice

were anesthetized on ice, and the cerebellum, olfactory bulbs, meninges, and hippocampus

were carefully removed using sharp tweezers (Dumont, 1214Y84) in cold dissection solution

(5 mM Hepes, 161 mM NaCl, 5 mM KCl, 1 mM $MgSO_4$, 3.7 mM $CaCl_2$, and 5.5 mM

Glucose, pH 7.4) under a stereo microscope. The remaining telencephala were then digested

in a dissection solution containing 10 U/ml papain (Worthington-Biochem, LS003126),

0.2 mg/ml L-cysteine, 1.5 mM NaOH, 1 mM $CaCl_2$, 0.5 mM EDTA, and 0.15% DNase I

(Sigma-Aldrich, D5025) for 30 min at 37°C. After gentle pipetting and centrifugation at 1,500

rpm for 5 min at room temperature (r.t.), the cells were resuspended in DMEM medium

supplemented with 10% FBS and 1% penicillin/streptomycin and seeded into flasks (Sigma-

Aldrich, L2020) pre-coated with 10 μg/ml Human Plasma Fibronectin (Sigma-Aldrich,

FC010). Neurons were mechanically shaken off and removed by patting the flask

approximately 2 days after inoculation. When reaching ~90% confluence, cells were digested

with trypsin and re-plated into fibronectin-coated 29-mm glass-bottomed dishes (Cellvis,

D29-14-1.5-N) for immunofluorescence staining. Once cells were cultured to ~90%

confluency, FBS was removed from the medium to induce differentiation.

For the primary culture of mTECs, we followed the method described (Zhao et al.,

2021). mTECs were used to investigate the expression profile of *Kif6* and *Kif9* during the

differentiation of mTECs, because of their high induced differentiation efficiency.

For transfection of plasmids, cells were transfected at ~70% confluency with

Lipofectamine 2,000 (Thermo Fisher).

459   Preparation and culture of whole-mounts of lateral ventricle walls

Whole-mounts of the lateral ventricles were prepared as described (Ohata et al., 2014)

with minor modifications. Mice were euthanized with CO_2, and the lateral ventricles were

carefully dissected at a thickness of 500-1000 μm using Vannas Scissors (66VT, 54140B) in

pre-warmed (37°C) dissection solution (25 mM Hepes,117.2 mM NaCl, 5.3 mM KCl,

0.81 mM $MgSO_4$, 1.8 mM $CaCl_2$, 26.1 mM $NaHCO_3$, 1 mM $NaH_2PO_4 \cdot 2H_2O$, and 5.6 mM

Glucose, pH 7.4). The samples were subsequently cultured in DMEM supplemented with 0.3

mg/ml glutamine, 100 U/ml penicillin, and 100 U/ml streptomycin.

467   Viral production and transfection

Adenovirus particles were utilized to infect mEPCs or IMCD3 cells and produced in

HEK293A cells as previously described (Zhao et al., 2021). In brief, the adenovirus

expression constructs were linearized using *P*acI restriction enzyme (NEB) and transfected

into HEK293A cells cultured in 6-well plates using Lipofectamine 2,000 to produce the initial

adenovirus. When approximately 80% of the cells exhibited a cytopathic effect (CPE), the

culture medium was collected and infected HEK293A cells in a 10-cm dish. Once about 80%

of the cells showed a CPE, the culture medium was collected to infect fresh HEK293A cells

for viral amplification. When ~80% of the cells exhibited a CPE, both cells and the culture

medium were collected. To ensure complete release of the viral particles, three freeze-thaw

cycles (-80°C and 37°C) were performed, followed by centrifugation at 1,500 rpm for 5 min

to remove cell debris. The harvested adenoviral particles were used at a dilution of 1:1000 or

1:200.

480   Immunofluorescence staining and imaging

IMCD3 cells were fixed with 4% paraformaldehyde (PFA) in PBS, whereas mEPCs

grown on 29 mm glass-bottom dishes and freshly dissected whole-mounts were pre-

permeabilized with 0.5% Triton X-100 in PBS for 30 sec before fixation to remove soluble

proteins. After fixation, the cells were extracted with 0.5% Triton X-100 in PBS for 15 min

and blocked in 4% BSA in TBST at r.t. for 1 h. Next, the cells were incubated with primary

antibodies overnight at 4°C. After three washes with the blocking solution for 5 min each, the

cells were incubated with secondary antibodies at r.t. for 1h, and mounted in Dako (Dako,

S3023) or ProLong™ (Thermo Fisher, 2273640). All the antibodies used are listed in

Supplementary Table 2.

Fluorescent images were captured using a Leica TCS SP8 system with a HC PL APO

CS2 63 × /1.40 OIL objective or an Olympus Xplore SpinSR10 microscope with UPLAPO

OHR 60 ×/1.50 Objective Lens, and images were processed with maximum intensity

projections. 3D-SIM super-resolution images were taken with a DeltaVision OMX SR system

(GE Healthcare) and the z-axis scanning step was 0.125 μm. The original images were

processed for OMX SI Reconstruction, OMX Image Registration, and maximum intensity

projection with SoftWoRx software.

To perform GI-SIM imaging for mEPCs, the cells were seeded onto fibronectin-coated

12-mm coverslips and were infected with adenovirus one day prior to serum starvation. To

fluorescently label their multicilia, mEPCs were incubated with 100 nM SiR-tubulin

(Spirochrome, SC002) for 30 min or 200 nM Janelia Fluor® 549 HaloTag ligand (Promega,

GA1110) for 20 min. After labeling, mEPCs were gently laid flat against the bottom coverslip

and GI-SIM imaging was performed as described (Guo et al., 2018; Qiao et al., 2022). For

dual-color imaging, instrument calibration was conducted routinely with fluorescent beads.

The images and movies were processed using Fiji and Huygens softwares.

To record ciliary beating, the whole mounts of lateral ventricles were dissected and cut at

a thickness of 500-1000 μm as previously described (Ohata et al., 2014; Zhao et al., 2021). To

label the cilia, 100 nM SiR-tubulin was added to the culture medium and incubated for 1 h.

High-speed live imaging was conducted at 37°C using an Olympus Xplore SpinSR 10

microscope equipped with UPLAPO OHR 60 × /1.50 Objective Lens, Hamamatsu ORCA-

Fusion camera, 4,000 rpm CSU Disk Speed, and OBIS Laser. Ciliary beat was captured at

100 frames per second (fps) by a time-lapse collection of single optical sections with 80%

laser power (640 nm) to achieve an exposure time of 5 ms. Kymographs were generated for

the first 500 ms using Fiji.

The ependymal flow assay was performed as described (Liu et al., 2021; Mirzadeh et al.,

2010) with minor modifications. Briefly, the freshly dissected ependyma was sliced using a

vibratome (Leica VT1000S) to obtain slices with a thickness of 250 μm. Before imaging,

multicilia were labeled with 100 nM SiR-tubulin for 1 h and small fluorescent latex beads

with a diameter of 250 nm at 1:500 dilution (Invitrogen, F8809) were added to the culture

medium. Time-lapse microscopy was performed at 37 ℃ using using an Olympus Xplore

SpinSR10 microscope equipped with UPLAPO OHR 60 × /1.50 Objective Lens. The laser

power for the beads channel (561 nm) was adjusted based on the fluorescent intensity, and

exposure time was set to 100 ms. The laser power for the SiR-tubulin channel (640 nm) was

set to 80% to achieve an exposure time of 5 ms.

During recording, fluorescent beads suspended in the culture medium containing the

living ependymal tissues were imaged at a fixed z-plane to record their motility.

Subsequently, multicilia in the same areas were visualized by adjusting the focal plane, and

live imaging of a single optical section was conducted to record ciliary motility. All frames of

beads in the same area were projected together from Movies 11-12 and superimposed with

trajectories of traceable, rapidly-moving bead aggregates in the first 500 ms to show flow

directions and tracks. The velocity of beads was meausured using Fiji.

To record sperm motility, sperm were released from cauda epididymidis into buffer

(Njabsw, M1130), and followed by live imaging with a spinning disk confocal microscope at

a fixed z-plane and at 25-ms intervals. Two pair of P88 wild-type mice and $Kif6^{-/-}$ litermates

were performed.

Immunoblotting

Immunoblotting experiments were performed as described (Zhao et al., 2021). To

prepare tissue lysates, freshly dissected tissues from mice were placed in ice-cold RIPA buffer

(50 mM Tris-HCl, 150 mM NaCl, 1% NP-40, 0.5% Sodium Deoxycholate, 0.1% SDS, and 1

mM EDTA, pH7.5) supplemented with protein inhibitors cocktail. The tissues were

homogenized using the homogenizer Precellys 24 system (Bertin, France). The lysates were

cleared by centrifugation at 13,200 rpm at 4°C for 15 min, and mixed with an equal volume of

2 × SDS-PAGE loading buffer (100 mM Tris-HCl, 4% SDS, 20% glycerol, 2% 2-

mercaptoethanol, 0.1% bromophenol blue, pH 6.8). For cultured cells, cells were washed with

pre-warmed PBS and lysed in 2 × SDS-PAGE loading buffer. The lysates were boiled at

100°C for 10 min and then subjected to immunoblotting.

Electron microscopy

For scanning electron microscopy (SEM), mice were deeply anesthetized with 360 mg/kg body weight of avertin (Sigma-Aldrich, T48402) by intraperitoneal injections. The tissues were dissected after transcardial perfusion with 4% PFA in PBS using an injection pump (Smiths medical, WZS-50F6, 250 ml/h). The dissected tissues were post-fixed with 2.5% glutaraldehyde at 4°C overnight. Subsequently, the tissues were treated with 1% OsO_4 at 4°C for 1 h and dehydrated with a graded series of ethanol. Critical point drying (Leica EM CPD300) was performed before metal shadowing (40 mA, 60 s). Images were captured using a FEI Quanta 250 scanning electron microscope.

For transmission electron microscoy (TEM), the tissues were fixed in 2.5% glutaraldehyde/PBS at 4°C overnight, washed with PBS, and treated with 1% OsO_4 for 1.5 h at r.t.. The tissues were dehydrated with a graded series of ethanol and further dehydrated with acetone. Subsequently, the samples were embedded in Epon 812 resin at 60°C for 48 h. Thin sections of 70 nm thickness were obtained using an ultramicrotome (Leica EM PACT2) and stained with 2% uranyl acetate for 15 min, followed by 1% lead citrate for 5 min. The images were captured using a FEI Tecnai G2 Spirit Twin transmission electron microscope and analyzed using Fiji software.

Coronal brain section

Coronal brain slices with a thickness of 250 μm were achieved using a vibratome sectioning technique, as previously described (Chen et al., 2020). Mice were deeply anesthetized with 360 mg/kg body weight of avertin administered via intraperitoneal injections. Transcardial perfusion was performed using PBS followed by 4% PFA in PBS, utilizing an injection pump (Smiths medical, WZS-50F6, 250 ml/h). Whole brains were dissected from the mice and cut using a vibratome to obtain coronal slices with a thickness of 250 μm. These slices were placed into 35-mm glass-bottom dishes (Cellvis, D35-20-1.5-N) and were imaged with an Olympus SZX16 Stereo Microscope.

Reproductive performance test

Mating test was perform as previously described (Miyata et al., 2020). 2-3 month-old $Kif6^{-/-}$ and $Kif9^{-/-}$ male mice with normal appearance were individually caged with two age-

576 appropriate heterozygous female mice for four months and two *Kif6*[-/-] and *Kif9*[-/-] female mice

577 were separately caged with one heterozygous male mice, and then were checked regularly

578 every several days. The number of pups was counted during the period. Two independent

579 experiments were carried out.

Phylogenomic analysis and multiple alignment

The phylogenetic tree and taxonomic groups were constructed based on literatures and

Taxonomy Browser of National Center for Biotechnology Information (NCBI) (Cetkovic et

al., 2018; Mukherjee and Brocchieri, 2013). The protein database of NCBI were searched to

identify genes with orthologous proteins of Kif6 and Kif9 across a wide range of species

(such as mammals, birds, amphibians, fishes, jawless vertebrates, cephalochordates,

urochordates, echinoderms, nematodes, mollusks, flatworms, cnidarians, placozoans and

protozoans) using the full-length murine protein sequences of Kif6 and Kif9 as queries.

The selected sequences were aligned using COBALT searches with default settings.

Genebank accession numbers of Kif6 and Kif9 orthologues were provided in Supplemenatry

Figure 1a. Only alignment columns without gaps were colored, with highly conserved

positions shown in red and less conserved positions shown in blue.

Analysis of BB rotational polarity

BB rotational polarity was analyzed as previously described (Guirao et al., 2010;

Mirzadeh et al., 2010; Ryu et al., 2021). In TEM images, the BB rotational polarity was

determined by drawing a unit vector from the center of a BB to the vertex of its BF. The mean

vector length within a given field in each full-size electron micrograph (2,048 pixels × 2,048

pixels) was calculated using Microsoft Excel software (Microsoft). In 3D-SIM images, the

BB rotational polarity was determined by drawing a unit vector from the geographic center of

Cep164 ring (BB distal end marker) to the center of either Centriolin or γ–tubulin signals (BF

marker). The mean vector length within a cell was calculated using Excel software (Microsoft)

to demostrate the extent of the rotational polarity.

Quantification and statistical analysis

BB numbers were assessed without knowledge of the genotypes and were measured

from 3D-SIM images using Cep164-positive ring as a BB marker. The BB center and cell

border center were determined using the default setting of the "ROI Manager" function in

Fiji.

All experiments, unless otherwise stated, were independently repeated at least twice for

both microscopic or biochemical results. Statistical results are presented as mean ± SD.

Unpaired two-tailed student's t-test was performed using GraphPad Prism 9.0 software, and

differences were considered significant at $P < 0.05$. Data analyses were performed in a

blinded manner, with the researchers performing the quantification being unaware of the

genotype.

Acknowledgements.

The authors thank Profs. Lan Bao and Dr. Lei Diao (Center for Excellence in Molecular

Cell Science, CAS) for the reagents and technical assistance, Profs. Xin Liang (Tsinghua

University) and Profs. Wei Feng (Institute of Biophysics, CAS) for technical help and

valuable discussions, and the staff members of the Integrated Laser Microscopy System at the

National Facility for Protein Science in Shanghai (NFPS) (Shanghai Advanced Research

Institute, CAS) and core facilities of the Center for Excellence in Molecular Cell Science

(CEMCS) for instrumental and technical supports. This work was supported by National

Natural Science Foundation of China (31991192, 32230027 and 32270725).

Declaration of interests

The authors declare no competing financial interests.

Author contributions

X.Y. and X.Z. conceived and supervised the project; C.F., X.P. and Di Li performed the

research; Dong Li contributed to GI-SIM imaging; Y.C., L.L. and Q.G. assisted with

experiments; X.Y., X.Z., C.F. and X.P. collectively analyzed data and wrote the manuscript,

with input from all authors.

Figure legends

Figure 1. Molecular characterization and subcellular localization of mammalian *Kif6*

and *Kif9*. (a) Schematics of full-length murine Kif6 and Kif9. The motor and coiled-coil

domains were identified using Conserved Domain Search Service of NCBI and SMART

(Simple Modular Architecture Research Tool) (http://smart.embl.de). **(b)** Conservation of *Kif6*

and *Kif9* across species with motile cilia and/or flagella throughout the evolution process. The

phylogenetic tree and taxonomic groups were referrenced from literature (Cetkovic et al.,

2018; Mukherjee and Brocchieri, 2013). **(c)** mRNA expression profile of *Kif6* and *Kif9* during

multiciliation. The gene expression profiles were obtained from our previous cDNA

microarray analysis of mTECs cultured at an air-liquid interface (ALI) for the indicated days

(Xu et al., 2015). Expression patterns of genes crucial for cilia formation (*Ift88* and *Ift122*)

and BB amplification (*Cep152*), were listed for comparison. **(d)** Expression patterns of Kif6

and Kif9 during multiciliation of mTECs. Multicilia formation is indicated by increased levels

of Ift81, Bbs3 and acetylated tubulin (Ac-tub). Gapdh served as loading control. **(e, f)**

Abundance of Kif6 and Kif9 in mouse tissues (**e**) and cultured cells (**f**) containing motile cilia.

Tissues from a 2-month-old male mouse were dissected and lysed. mEPCs were harvested on

day 8 post serum starvation. RPE1, NIH3T3 and IMCD3 cells were serum-starved for 48 h to

induce ciliogenesis. Gapdh or β-actin served as loading control. **(g)** Experimental scheme for

(**h, j-l and Supplementrary Fig. 2**). Precursors of mEPCs were isolated from P0 mouse brain

and serum-starved on day 0 to induce to differentiate into multiciliated cells. Cells were fixed

on day 8 for immunofluorescence (IF) staining and imaged by confocal microscopy or 3D-

SIM. To express GFP-tagged proteins, cells were infected with adenovirus on day-1. **(h)**

Endogenous Kif6 and Kif9 localized in multicilia. Ac-tub labeled ciliary axonemes

(arrowheads). **(i)** Kif6 and Kif9 did not localize in primary cilia. IMCD3 cells were harvested

at day 2 post serum starvation, and fixed for immunostaining, followed by confocal

microscopy. DAPI marked nucleus. Cilia pointed by arrowheads were zoomed in 2× to show

details. **(j)** Exogenous Kif6-GFP and Kif9-GFP localized at the axonemes of multicilia. **(k, l)**

Typical 3D-SIM images of endogenous Kif9 and Kif6. Kif9 was enriched in the central lumen

(**k**), while Kif6 localized as punta along peripheral DMTs (**l**). Ac-tub and Hydin served as

axoneme and CP markers, respectively. Cilia pointed were magnified by 200% for details.

Corresponding line scans at the indicated region of the magnified insets showed

colocalization of Kif9 with Hydin (**k**), and Kif6 distributed along the axoneme with Ac-tub

(**l**). Illustrations are provided to aid comprehensions.

Figure 2. Kif6, differed from Kif9, moves along axonemal MTs and colocalizes with IFT

particles. Quantification results are presented as mean ± SD and subjected to unpaired two-

tailed student's t-test: $*P < 0.05$. **(a)** Experimental scheme for ciliary trafficking recorded by

GI-SIM. mEPCs were infected with adenovirus at day-1 and imaged during day 5-8. **(b)**

Expression levels of GFP-tagged proteins in day 7 mEPCs. Immunoblotting was performed

using GFP, Kif6, Kif9 and Ift81 antibodies, respectively. A star indicates a non-specific band

closed to Ift81-GFP. **(c-e)** Ciliary movement of Ift81-GFP (**c**), Kif6-GFP (**d**) and Kif9-GFP

(**e**). Both Ift81-GFP (**c**) and Kif6-GFP (**d**) were able to move along the axonemal outer DMTs

while the motility of Kif9-GFP (**e**) was not detectable. Images were recorded at 25 frames per

sec (fps). The trafficking trajectories of GFP-tagged puncta through a time projection of the

first 200 frames from Movies 1-3 were showed. Typical cilia framed were magnified by 300%

to show detailed bidirectional transports of GFP-fusion puncta. The clear and traceable GFP-

tagged particles were marked and corresponding kymographs were presented. In (**c**), two

GFP-tagged particles (labeled by 1, 3) and two (labeled by 2, 4) were showed for anterograde

and retrograde transports, respectively. In (**d**), two GFP-tagged particles (labeled by 1, 2) and

one (labeled by 3) were showed for anterograde and retrograde transports, respectively. **(f, g)**

Quantification results of velocities and frequencies from the indicated numbers of particles (**f**)

and cilia (**g**). Antero: Anterograde transport; Retro: Retrograde transport. **(h, i)** Colocalization

of Kif6-GFP and IFT protein (Ift56) at axonemes by confocal (**h**) and 3D-SIM (**i**). Ac-tub

marked ciliary axoneme. A typical cilium was magnified by 400% to show details. Line scans

at the indicated regions (I and II) exihibited the colocalization of Kif6-GFP and Ift56 along

the axoneme. Co-localized punta were pointed by yellow arrowheads.

Figure 3. Kif6-GFP punta move together with IFT particles along the axoneme. (a-c)

Ciliary trafficking of fluorescent particles. mEPCs were infected with adenovirus expressing

of Kif6-GFP and Ift27-Halo and dual-color time-lapse live imaging was taken at 14 fps. Three

distinct bidirectional moving particles within cilia were observed: particles labeled with both

Ift27-Halo and Kif6-GFP (Ift27[+]Kif6[+] particles) (**a**, **b**), those labeled only with Kif6-GFP

(Kif6$^+$ particles), and those labeled only with Ift27-Halo (Ift27$^+$ particles). Ciliary trafficking

trajectories of GFP/Halo-tagged puncta were projected by the first 200 frames from Movies 4-

5. Typical cilia pointed were magnified by 300% to display traceable anterograde and

retrograde transports of Ift27$^+$Kif6$^+$ particles (**a, b**), Kif6$^+$ particles (**c**) and Ift27$^+$ particles (**c**)

from representative frames, respectively. The clear and traceable particles are indicated by

colored arrows or arrowheads in magnified cilia. Corresponding kymographs were presented.

(**d, e**) Quantification and velocities of three types particles of anterograde and retrograde

transports. 45 clear cilia from 4 cells co-expressed Kif6-GFP with Ift27-Halo were scored.

The corresponding particles number (n) was presented.

Figure 4. Knockout of *Kif6* or *Kif9* leads to hydrocephalus and male infertility. (a)

Schematic for generating *Kif6* (top) and *Kif9* (bottom) knockout mice using CRISPR/Cas9

technique. The targeting of sgRNA (scissors) and genotyping PCR primers were indicated for

each knockout model. (**b**) Representative genotyping PCR results. Genomic DNAs were

extracted from P21 mice tails and amplified by PCR using the indicated primers in (**a**). (**c**)

Complete depletion of Kif6 and Kif9 proteins in the indicated tissues. Tissue lysates were

prepared from 2-month-old mice of the indicated genotypes. Gapdh served as a loading

control. (**d, e**) Morphologies of *Kif6* and Kif9 deficient mice. *Kif6$^{-/-}$* mice exhibited a dome-

shaped skull (arrow) and kyphosis (arrowhead) with skinny bodies (**d**). In contrast, *Kif9$^{-/-}$*

mice showed a normal appearance (**e**). (**f, g**) Representative images of coronal brain sections

of *Kif6$^{-/-}$* (**f**) or *Kif9$^{-/-}$* (**g**) and their corresponding wide-type littermates. M: month. (**h**)

Quantification of mice with enlarged ventricles (n = 60 for *Kif6$^{-/-}$*; n = 11 for *Kif9$^{-/-}$* mice). (**i**)

Survival curves of mice in indicated genotypes during a 12-month observation period.

Numbers of mice used for each genotype were indicated. (**j, k**) Morphology and motility of

sperm from wide-type and *Kif6$^{-/-}$* littermates. Sperm motility trajectories were color-coded

through projection of the first 50 frames (**k**). Two pairs of P88 *Kif6$^{-/-}$* mice and wild-type

liitermates were used. Refer to Movie 7.

Figure 5. Knockout of *Kif6* impairs planar polarity of cilia across the ependyma.

Quantification results are pooled together (mean ± SD plus sample dots) and subjected to

unpaired two-tailed student's *t*-test: *$P < 0.05$. M: mild hydrocephalus mice with normal

appearance but larged ventricles; S: severe hydrocephalus mice with the observable dome-

shaped skull like Fig. 1d. **(a)** Schematic of experimental design. Whole-mounts of ventricle

walls were freshly dissected and subjected to electron microscopy (EM) or live cell imaging.

SiR-tubulin, a fluorescent dye for MTs, was used to label cilia. Ciliary motility was live-cell

imaged at a single optical section with 10-ms intervals. An illustration of the beat pattern of

an ependymal cilium, depicted as a side view, is provided for better comprehension. **(b)**

Representative SEM images of ependymal multicilia from wild-type and *Kif6*[-/-] littermates at

P4 (top) and P30 (bottom). **(c)** Representative side-view cilia beat images. Cilia beat patterns

in individual cells were unaffected by *Kif6*-deficiency. Two consecutive cycles of cilia beat

patterns were showed. The beat direction was indicated by color-coded arrows, followed by

pseudo-coloring and merging. Representative top-view images from another pair of

littermates were presented in Supplementary Fig. 4b. **(d)** Quantification of ciliary beating

frequencies. Results were from 5 pairs of littermates with 60 cells per mouse. **(e)**

Unidirectional multicilia beat across ependyma was disrupted by the loss of Kif6. The first

three consecutive frames cropped from Movies 8-9, were pseudo-colored and overlaid to

show ciliary motilities. Beating directions of multicilia are indicated by arrows.

Representative images of another two pairs of littermates were presented in Supplementary

Fig. 4c. **(f)** Representative top-view images of cilia beat patterns in *Kif9*-deficient mice. Two

consecutive cycles of cilia beat patterns are presented. Representative side-view fields from

another pair of littermates were presented in Supplementary Fig. 4d. **(g)** Tissue-wide polarity

of cilia beating was not altered in *Kif9*-deficient ependyma. The first three consecutive frames

cropped from Movies 9-10, were pseudo-colored and merged to show ciliary motilities.

Additional representative images were presented in Supplementary Fig. 4e. **(h)**

Representative projections of bead movements from Movies 12-13. Trajectories of traceable,

rapidly-moving beads in the first 500 ms were superimposed to show flow directions. **(i)**

Quantification of bead velocities. Results from two pairs of P36 and one pair of P48

littermates, with 60 traceable beads for each.

Figure 6. *Kif6* deficiency impairs rotational polarity of BBs in ependymal tissues.

Quantification results are pooled together (mean ± SD plus sample dots) and subjected to

unpaired two-tailed student's *t*-test: *$P < 0.05$. **(a)** Structural features of a motile cilium. Top

and bottom panels illustrate the side view of the cilium and typical transverse view of the BB,

respectively. BB polarity marked by its conical BF that extends laterally to the MT triplet. **(b-**

f) The loss of Kif6, but not Kif9, impaired rotational polarity. Ependymal tissues from the

indicated mice were immunostained for Cep164 to visualize BBs and γ-tubulin to visualize

BFs. Framed regions **(c, e)** were magnified to show the rotational polarity of BBs in

individual cells **(b, c)**. The BF orientations are indicated by arrows **(b, d)**. Note the

misorangized rotational polarity in both cellular and tissue-level in *Kif6*[-/-] ependyma. Mean

vector lengths **(f)** were scored from 20 cells each mouse. Mice used were two 6-month-old

(6M) and one 7-month-old (7M) mice for *Kif9*[-/-] and three of their wild-type littermates; two

P34, one P43 mice with severe hydrocephalus (Se) and one P62 mouse with mild

hydrocephalus (Mi) for *Kif6*[-/-] and their corresponding wild-type littermates. **(g, h)**

Representative TEM images **(g)** and quantification analysis on the extent of rotational polarity

(h). BF orientations are indicated by arrows. The mean vector length **(h)** was calculated from

BBs with clear BFs in at least ten full-size electron micrographs and over 130 BFs were

scored for each genotype. **(i)** Illustrations are provided to aid comprehensions on the functions

of Kif9 (top) and Kif6 (bottom) in the rotational polarity of BBs. **(j)** The number of BBs in

ependymal cells was not altered by *Kif6* or *Kif9*-deficiency. 50 cells were measured per

mouse. **(k)**. A summarizing model showing the distinct functions of *Kif6* and *Kif9* in motile

cilia. Kif6 may be involved in material transport associated with a specific IFT machinery, or

transport cargo on its own, thus thereby ensuring BB polarity and the coordinated ciliary beat.

Kif9 may regulate ciliary beat by inducing changes on subtle structure related to

conformation. BB, basal body; BF, basal foot; CP, central pair; DMT, doublet microtubule;

TF, transition fiber; TZ, transition zone.

Legends for Supplementary Figures

Supplementary Figure 1. Comparison of murine Kif6 and of Kif9 (related to Fig. 1). (a)

Sequence alignment of Kif6 (left) and Kif9 (right) orthologues as indicated Fig. 1b. Highly

conserved positions were shown in red and less conserved positions shown in blue. The

sequences of Kif6 orthologues used were: mouse (NP_796026.2), human (NP_659464.3),

chicken (XP_003640993.5), frog (NP_001106482.1), zebrafish (NP_001070899.1), lancelet

(XP_035685117.1), ascidian (XP_002124970.1), sea urchin (XP_030854411.1), mussel

(CAG2221785.1), flatworm (PAA61917.1), sea anemone (XP_032228212.2), *Trichoplax*

(RDD42533.1), *Tetrahymena* (XP_001024804.2), *Trypanosoma* (XP_846346.1) and green

algae (XP_042928646.1). The sequences of Kif9 orthologues used were: mouse

(NP_001157041.1), human (NP_001400904.1), chicken (XP_040522129.1), frog

(XP_012821147.1), zebrafish (XP_001922460.1), lancelet (XP_035663761.1), ascidian

(XP_018668035.1), sea urchin (XP_030830023.1), mussel (CAG2242108.1), flatworm

(PAA61899.1), sea anemone (XP_001636554.1), *Trichoplax* (RDD45058.1), *Tetrahymena*

(XP_001022313.1), *Trypanosoma* (RHW71316.1) and green algae (P46870.1). Multiple

sequence alignments were obtained using COBALT (NCBI). **(b)** Protein sequence alignment

of motor domains (Kif6-MD, 1-353 aa; Kif9-MD, 1-348 aa). The alignment was generated by

the COBALT sequence alignment program using default parameters. Genebank accession

numbers of proteins were as follows: Kif6 (NP_796026.2) and Kif9 (NP_001157041.1). The

motors were identified using SMART.

Supplementary Figure 2. The localization of exogenous Kif6 and Kif9 within motile cilia

(related to Fig. 1). mEPCs were harvested at day 8 and fixed for immunostaining, followed

by 3D-SIM. **(a, b)** Localization of Kif6-GFP (**a**) and Kif9-GFP (**b**) in the axonemes of

multicilia. Kif6-GFP exhibited punctate localization along the axonemal DMTs whereas Kif9-

GFP was enriched in CP lumen. Spef1 served as the CP marker. Typical cilia were magnified

by 200% to highlight details. Line scans were generated at the indicated positions by white

lines in merged images.

Supplementary Figure 3. Kif6 moves along the axoneme and colocalizes with IFT

particles (related to Fig. 2 and 3). (a) Kif6-GFP, but not Kif9-GFP, resembling Ift81-GFP,

moved along the axonemal DMTs. The trafficking trajectories of GFP-tagged puncta were

visualized through projection of the first 200 frames. Typical cilia were magnified by 300% to

show the detailed anterograde transport of GFP-fusion puncta. The clear and traceable GFP-

tagged particles were marked and corresponding kymographs were presented. **(b, c)** Kif6-

GFP co-localized with IFT protein (Ift88) at axonemes from confocal (**b**) and 3D-SIM images

(**c**). Ac-tub marked ciliary axoneme. A typical cilium was zoomed in 4× for details. Line scan

at the indicated region exihibited the colocalization of Kif6-GFP and Ift88 along the axoneme

(**c**). (**d**) Calibration of the microscope. Instrument calibration was conducted routinely with

fluorescent beads before dual-color GI-SIM imaging experiments. A line scan was performed

along the white line to determine the alignment.

Supplementary Figure 4. *Kif6* deficiency disrupts PCP of cilia beat across ependyma

(related to Fig. 6). Quantification results are pooled together (mean ± SD plus sample dots)

and subjected to unpaired two-tailed student's *t*-test. (**a**) Representative TEM images of

ependymal cilia from wild-type and *Kif6*$^{-/-}$ mice. CP MTs (arrow); inner and outer dynein

arms (arrowheads); transition zone (TZ, indicated by yellow lines); basal body (BB, indicated

by yellow lines) (n =2 biological replicates). (**b**) Representative top-view images of Cilia beat

pattern from *Kif6*$^{-/-}$ mice and wild-type littermates. At least two beat cycles from consecutive

images were exhibited. The beat directions of the initial cycle are indicated by arrows,

followed by pseudo-coloring and merging. (**c**) Cilia beat directions across ependyma by *Kif6*-

deficency. The first three consecutive frames were pseudo-colored and overlaid to show

ciliary beat directions. (**d**) Representative side-view cilia beat images. At least two

consecutive cycles of cilia beat were presented. The beat direction is indicated by color-coded

arrows, followed by pseudo-coloring and merging. (**e**) Cilia beat directions across ependyma

by *Kif9*-deficiency. Beat directions of multicilia (white arrows) in representative microscopic

fields were presented. (**f**) Ciliary beat frequencies of *Kif9*$^{-/-}$ ependymal cilia. Quantification

results were from 3 pairs of littermates with 50 cells per mouse. S: Severe hydrocephalus

mice with obvious dome-shaped skull. M: mild hydrocephalus mice with normal

morphological appearance but enlarged ventricles.

Supplemental Figure 5. *Kif6* deficiency disrupts the rotational polarity of BBs in

ependyma (related to Fig. 7). (a-d) BBs in *Kif6*$^{-/-}$ (**c, d**) but not in *Kif9*$^{-/-}$ (**a, b**) ependymal

cells displayed impaired rotational polarity. Ependymal tissues from the indicated littermates

were co-immunostained for Cep164 to visualize BBs and Centriolin to visualize BFs. Framed

regions in (**b, d**) were maginified to show BF orientations (arrows) in (**a, c**). Arrows in (**b, d**)

indicate rotational polarities of the wild-type, $Kif9^{-/-}$ and $Kif6^{-/-}$ BB patches, which broadly

point to one direction in the microscopic field. **(e)** Quantification of the rotational polarity.

Results from three mice per genotype were presented as mean ± SD plus sample dots and

subjected to unpaired two-tailed student's t-test: *$P < 0.05$. Mice used were: two 6-month-old

(6M) and one 7-month-old (7M) mice for $Kif9^{-/-}$ and three of their wild-type littermates; three

P35 mice with severe (S) hydrocephalus for $Kif6^{-/-}$ and three of their wild-type littermates. 50

ependymal cells were scored for each mouse.

Supplementary Figure 6. The translational polarity was not affected by *Kif6* deficiency

(related to Fig. 7). Pooled results from three mice per genotype were presented as mean ± SD

plus sample dots and subjected to unpaired two-tailed student's t-test: *$P < 0.05$. **(a-d)** *Kif6*

deficiency did not impair the transitional polarity of BBs. Ependymal tissues from two P62

and one P68 $Kif6^{-/-}$ mice of mild (M) hydrocephalus, three severe (S) hydrocephalus (P34,

P38 and P43) and six of their wild-type littermates were immunostained for ZO1 and Cep164

to visualize cell borders and BBs, respectively **(a)**. BB patch displacement relative to the

cellular geographic center **(c)** and relative BB patch area **(d)** were quantified as illustrated in

(b). 50 ependyma cells were scored for each mouse.

Legends for supplementary movies

Movie 1. Bidirectional motility of ciliary Ift81-GFP puncta (related to Fig. 2c). mEPCs

were infected with adenovirus at day-1 and imaged during day 5-8 by GI-SIM. Images were

recorded at 25 frames per sec (fps) and the playback speed is 20 fps. Typical cilia framed

were magnified by 300% to show detailed bidirectional transports of Ift81-GFP puncta.Two

particles (labeled by 1, 3) and two (labeled by 2, 4) were showed for anterograde and

retrograde transports, respectively.

Movie 2. Bidirectional motility of ciliary Kif6-GFP puncta (related to Fig. 2d). mEPCs

were treated, imaged, and presented as described in Movie 1. Typical cilia framed were

magnified by 300% to show detailed bidirectional transports of Kif6-GFP puncta. Two

particles (labeled by 1, 2) and one (labeled by 3) were showed for anterograde and retrograde

transports, respectively.

Movie 3. Kif9-GFP does not exhibit progressive movement along the cilia (related to Fig. 2e). mEPCs were treated, imaged, and presented as described in Movie 1. Typical cilia framed were magnified by 300% to show details.

Movie 4. Ciliary Kif6-GFP punta move together with IFT particles (related to Fig. 3a, b). mEPCs were infected with adenovirus expressing Kif6-GFP and Ift27-Halo, and dual-color time-lapse live imaging was taken by GI-SIM. Images were recorded at 14 fps and the playback speed is 20 fps. Typical cilia pointed were magnified by 300% to display traceable anterograde (Fig. 3a) and retrograde (Fig. 3b) transports of Ift27$^+$Kif6$^+$ particles from representative frames, respectively. Two particles (labeled by 1, 2) and one (labeled by 3) were showed for anterograde and retrograde transports, respectively.

Movie 5. Ciliary Kif6-GFP punta and IFT particles move separately (related to Fig. 3c). mEPCs were treated, imaged, and presented as described in Movie 4. Typical cilia pointed were magnified by 300% to display traceable anterograde and retrograde transports of Kif6$^+$ particles or Ift27$^+$ particles (related to Fig. 3c) from representative frames, respectively. Two particles (labeled by 5, 6) and two (labeled by 4, 7) were showed for anterograde and retrograde transports, respectively.

Movie 6. A representative sperm motility from *Kif9$^{+/+}$* and *Kif9$^{-/-}$* male mice. Sperm were released from cauda epididymidis, followed by live imaging with a spinning disk confocal microscope at a fixed z-plane and at 25-ms intervals. The playback speed is 20 fps. One pair of P80 wild-type (top) mice and *Kif9$^{-/-}$* (bottom) litermates were presented.

Movie 7. A representative sperm motility from *Kif6$^{+/+}$* and *Kif6$^{-/-}$* male mice (related to Fig. 5k). Sperm were released, imaged, and presented as described in Movie 6. One pair of P88 wild-type (top) mice and *Kif6$^{-/-}$* (bottom) litermates were presented.

Movie 8. Representative ciliary beat patterns in *Kif6$^{+/+}$* ependyma (related to Fig. 6e). P31 WT ependyma was freshly dissected and labeled with SiR-tubulin, followed by live imaging with a spinning disk confocal microscope at a fixed z-plane and at 10-ms intervals. The playback speed is 5 fps. Images of the first three frames are presented in Figure 6e.

Movie 9. Representative ciliary beat patterns in *Kif6$^{-/-}$* ependyma (related to Fig. 6e). P31

Kif6$^{-/-}$ ependyma was treated, imaged, and presented as described in Movie 8. The mouse used

was a littermate of the one in Movie 9, and exhibited severe hydrocephalus. Images of the

first three frames are presented in Figure 6e.

Movie 10. Representative ciliary beat patterns in *Kif9$^{+/+}$* ependyma (related to Fig. 6g).

7M WT ependyma was treated, imaged, and presented as described in Movie 9. Images of the

first three frames are presented in Figure 6g.

Movie 11. Representative ciliary beat patterns in *Kif9$^{-/-}$* ependyma (related to Fig. 6g).

7M *Kif9$^{-/-}$* ependyma was treated, imaged, and presented as described in Movie 8. The mouse

was a littermate of the one in Movie 10. Images of the first three frames are presented in

Figure 6g.

Movie 12. Representative liquid flows driven by ciliary beat in *Kif6$^{+/+}$* ependyma (related

to Fig. 6h). Fresh ependyma from a P36 wild-type mouse were incubated with SiR-tubulin to

label cilia and fluorescent latex beads as flow indicators. Beads motilities were imaged with a

spinning disk confocal microscope at 100-ms intervals and at a fixed z-plane. The playback

speed is 5 fps. Trajectories of 8 traceable, rapidly-moving aggregated beads in the first 500

ms were shown.

Movie 13. Representative liquid flows driven by ciliary beat in *Kif6$^{-/-}$* ependyma (related

to Fig. 6h). Fresh ependymal tissues from a P36 *Kif6$^{-/-}$* mouse with severe hydrocephalus

were treated, imaged, and presented as described in Movie 12. Trajectories of 13 traceable,

rapidly-moving aggregated beads in the first 500 ms were shown.

Legends for Supplementary Tables

Supplementary Table 1 Sequences of primers and gRNAs.

Supplementary Table 2 List of antibodies.

References

1. Bernstein, M., Beech, P.L., Katz, S.G., and Rosenbaum, J., L. (1994). A New Kinesin-like Protein
(Kip1) Localized to a Single Microtubule of the Chlamydomonas Flagellum. J Cell Biol *125*,
1313-1326.

2.   Bhogaraju, S., Cajanek, L., Fort, C., Blisnick, T., Weber, K., Taschner, M., Mizuno, N., Lamla, S.,
917   Bastin, P., Nigg, E.A.*, et al.* (2013). Molecular basis of tubulin transport within the cilium by
918   IFT74 and IFT81. Science *341*, 1009-1012.

919   3.   Boutin, C., Labedan, P., Dimidschstein, J., Richard, F., Cremer, H., Andre, P., Yang, Y.,
920   Montcouquiol, M., Goffinet, A.M., and Tissir, F. (2014). A dual role for planar cell polarity genes
921   in ciliated cells. Proc Natl Acad Sci U S A *111*, E3129-3138.

922   4.   Buchan, J.G., Gray, R.S., Gansner, J.M., Alvarado, D.M., Burgert, L., Gitlin, J.D., Gurnett, C.A.,
923   and Goldsmith, M.I. (2014). Kinesin Family Member 6 (kif6) Is Necessary for Spine
924   Development in Zebrafish.

925   5.   Bustamante-Marin, X.M., Yin, W.N., Sears, P.R., Werner, M.E., Brotslaw, E.J., Mitchell, B.J.,
926   Jania, C.M., Zeman, K.L., Rogers, T.D., Herring, L.E.*, et al.* (2019). Lack of GAS2L2 Causes
927   PCD by Impairing Cilia Orientation and Mucociliary Clearance. Am J Hum Genet *104*, 229-245.

928   6.   Butler, M.T., and Wallingford, J.B. (2017). Planar cell polarity in development and disease. Nat
929   Rev Mol Cell Biol *18*, 375-388.

930   7.   Cason, S.E., and Holzbaur, E.L.F. (2022). Selective motor activation in organelle transport along
931   axons. Nat Rev Mol Cell Biol *23*, 699-714.

932   8.   Cetkovic, H., Harcet, M., Roller, M., and Bosnar, M.H. (2018). A survey of metastasis suppressors
933   in Metazoa. Lab Invest *98*, 554-570.

934   9.   Chen, Y., Zheng, J., Li, X., Zhu, L., Shao, Z., Yan, X., and Zhu, X. (2020). Wdr47 Controls
935   Neuronal Polarization through the Camsap Family Microtubule Minus-End-Binding Proteins. Cell
936   Rep *31*, 107526.

937   10.   Chien, A., Shih, S.M., Bower, R., Tritschler, D., Porter, M.E., and Yildiz, A. (2017). Dynamics of
938   the IFT machinery at the ciliary tip. eLife *6*.

939   11.   Clare, D.K., Magescas, J., Piolot, T., Dumoux, M., Vesque, C., Pichard, E., Dang, T., Duvauchelle,
940   B., Poirier, F., and Delacour, D. (2014). Basal foot MTOC organizes pillar MTs required for
941   coordination of beating cilia. Nat Commun *5*.

942   12.   Delgehyr, N., Meunier, A., Faucourt, M., Bosch Grau, M., Strehl, L., Janke, C., and Spassky, N.
943   (2015). Ependymal cell differentiation, from monociliated to multiciliated cells. Methods Cell
944   Biol *127*, 19-35.

945   13.   Demonchy, R., Blisnick, T., Deprez, C., Toutirais, G., Loussert, C., Marande, W., Grellier, P.,
946   Bastin, P., and Kohl, L. (2009). Kinesin 9 family members perform separate functions in the
947   trypanosome flagellum. J Cell Biol *187*, 615-622.

948   14.   Duan, S., Li, H., Zhang, Y., Yang, S., Chen, Y., Qiu, B., Huang, C., Wang, J., Li, J., Zhu, X.*, et al.*
949   (2021). Rabl2 GTP hydrolysis licenses BBSome-mediated export to fine-tune ciliary signaling.
950   EMBO J *40*, e105499.

951   15.   Fliegauf, M., Benzing, T., and Omran, H. (2007). When cilia go bad: cilia defects and ciliopathies.
952   Nat Rev Mol Cell Biol *8*, 880-893.

953   16.   Gadadhar, S., Alvarez Viar, G., Hansen, J.N., Gong, A., Kostarev, A., Ialy-Radio, C., Leboucher,
954   S., Whitfield, M., Ziyyat, A., Touré, A.*, et al.* (2021). Tubulin glycylation controls axonemal
955   dynein activity, flagellar beat, and male fertility. Science *371*.

956   17.   Gibbons, I.R. (1961). The relationship between the fine structure and direction of beat in gill cilia
957   of a lamellibranch mollusc. J Biophys Biochem Cytol *11*, 179–205.

958   18.   Goetz, S.C., and Anderson, K.V. (2010). The primary cilium: a signalling centre during vertebrate
959   development. Nat Rev Genet *11*, 331-344.

19.   Graser, S., Stierhof, Y.-D., Lavoie, S.b.B., Gassner, O.S., Lamla, S., Le Clech, M., and Nigg, E.A.
(2007). Cep164, a novel centriole appendage protein required for primary cilium formation. The
Journal of Cell Biology *179*, 321-330.

20.   Guirao, B., Meunier, A., Mortaud, S., Aguilar, A., Corsi, J.M., Strehl, L., Hirota, Y., Desoeuvre,
A., Boutin, C., Han, Y.G.*, et al.* (2010). Coupling between hydrodynamic forces and planar cell
polarity orients mammalian motile cilia. Nat Cell Biol *12*, 341-350.

21.   Guo, Y., Li, D., Zhang, S., Yang, Y., Liu, J.J., Wang, X., Liu, C., Milkie, D.E., Moore, R.P., Tulu,
U.S.*, et al.* (2018). Visualizing Intracellular Organelle and Cytoskeletal Interactions at Nanoscale
Resolution on Millisecond Timescales. Cell *175*, 1430-1442 e1417.

22.   Han, L., Rao, Q., Yang, R., Wang, Y., Chai, P., Xiong, Y., and Zhang, K. (2022). Cryo-EM
structure of an active central apparatus. Nat Struct Mol Biol *29*, 472-482.

23.   Herawati, E., Taniguchi, D., Kanoh, H., Tateishi, K., Ishihara, S., and Tsukita, S. (2016).
Multiciliated cell basal bodies align in stereotypical patterns coordinated by the apical
cytoskeleton. J Cell Biol *214*, 571-586.

24.   Hirokawa, N., Noda, Y., Tanaka, Y., and Niwa, S. (2009). Kinesin superfamily motor proteins and
intracellular transport. Nat Rev Mol Cell Biol *10*, 682-696.

25.   Hoek, H., Klena, N., Jordan, M., Viar, G., Righetto, R., Schaffer, M., Erdmann, P., Wan, W.,
Geimer, S., Plitzko, J.*, et al.* (2022). In situ architecture of the ciliary base reveals the stepwise
assembly of intraflagellar transport trains. Science *377*, 543–548

26.   Inglis, P.N., Ou, G., Leroux, M.R., and Scholey, J.M. (2007). The sensory cilia of Caenorhabditis
elegans. WormBook, 1-22.

27.   Jordan, M.A., Diener, D.R., Stepanek, L., and Pigino, G. (2018). The cryo-EM structure of
intraflagellar transport trains reveals how dynein is inactivated to ensure unidirectional
anterograde movement in cilia. Nat Cell Biol *20*, 1250-1255.

28.   Joung, J., Konermann, S., Gootenberg, J.S., Abudayyeh, O.O., Platt, R.J., Brigham, M.D.,
Sanjana, N.E., and Zhang, F. (2017). Genome-scale CRISPR-Cas9 knockout and transcriptional
activation screening. Nat Protoc *12*, 828-863.

29.   Kanie, T., Abbott, K.L., Mooney, N.A., Plowey, E.D., Demeter, J., and Jackson, P.K. (2017). The
CEP19-RABL2 GTPase Complex Binds IFT-B to Initiate Intraflagellar Transport at the Ciliary
Base. Dev Cell *42*, 22-36 e12.

30.   Klena, N., and Pigino, G. (2022). Structural Biology of Cilia and Intraflagellar Transport. Annu
Rev Cell Dev Biol *38*, 103-123.

31.   Konjikusic, M.J., Lee, C., Yue, Y., Shrestha, B.D., Nguimtsop, A.M., Horani, A., Brody, S.,
Prakash, V.N., Gray, R.S., Verhey, K.J.*, et al.* (2023). Kif9 is an active kinesin motor required for
ciliary beating and proximodistal patterning of motile axonemes. J Cell Sci *136*.

32.   Konjikusic, M.J., Yeetong, P., Boswell, C.W., Lee, C., Roberson, E.C., Ittiwut, R.,
Suphapeetiporn, K., Ciruna, B., Gurnett, C.A., Wallingford, J.B.*, et al.* (2018). Mutations in
Kinesin family member 6 reveal specific role in ependymal cell ciliogenesis and human
neurological development. PLoS Genet *14*, e1007817.

33.   Lechtreck, K.F., Delmotte, P., Robinson, M.L., Sanderson, M.J., and Witman, G.B. (2008).
Mutations in Hydin impair ciliary motility in mice. J Cell Biol *180*, 633-643.

34.   Lechtreck, K.F., Sanderson, M.J., and Witman, G.B. (2009). High-speed digital imaging of
ependymal cilia in the murine brain. Methods Cell Biol *91*, 255-264.

35.   Lindemann, C.B., and Lesich, K.A. (2021). The many modes of flagellar and ciliary beating:

Insights from a physical analysis. Cytoskeleton (Hoboken) *78*, 36-51.

36. Liu, H., Zheng, J., Zhu, L., Xie, L., Chen, Y., Zhang, Y., Zhang, W., Yin, Y., Peng, C., Zhou, J.*, et*
al. (2021). Wdr47, Camsaps, and Katanin cooperate to generate ciliary central microtubules. Nat
Commun *12*, 5796.

37. Lukinavicius, G., Reymond, L., D'Este, E., Masharina, A., Gottfert, F., Ta, H., Guther, A.,
Fournier, M., Rizzo, S., Waldmann, H.*, et al.* (2014). Fluorogenic probes for live-cell imaging of
the cytoskeleton. Nat Methods *11*, 731-733.

38. Marshall, W.F., and Kintner, C. (2008). Cilia orientation and the fluid mechanics of development.
Curr Opin Cell Biol *20*, 48-52.

39. Matsuo, M., Shimada, A., Koshida, S., Saga, Y., and Takeda, H. (2013). The establishment of
rotational polarity in the airway and ependymal cilia: analysis with a novel cilium motility mutant
mouse. Am J Physiol Lung Cell Mol Physiol *304*, L736-745.

40. Meleppattu, S., Zhou, H., Dai, J., Gui, M., and Brown, A. (2022). Mechanism of IFT-A
polymerization into trains for ciliary transport. Cell *185*, 4986-4998 e4912.

41. Meng, Z., Meng, Q., Gao, T., Zhou, H., Xue, J., Li, H., Wu, Y., and Lv, J. (2022). Identification of
bi-allelic KIF9 loss-of-function variants contributing to asthenospermia and male infertility in two
Chinese families. Front Endocrinol (Lausanne) *13*, 1091107.

42. Miki, H., Okada, Y., and Hirokawa, N. (2005). Analysis of the kinesin superfamily: insights into
structure and function. Trends Cell Biol *15*, 467-476.

43. Mirzadeh, Z., Han, Y.G., Soriano-Navarro, M., Garcia-Verdugo, J.M., and Alvarez-Buylla, A.
(2010). Cilia organize ependymal planar polarity. J Neurosci *30*, 2600-2610.

44. Miyata, H., Shimada, K., Morohoshi, A., Oura, S., Matsumura, T., Xu, Z., Oyama, Y., and Ikawa,
M. (2020). Testis-enriched kinesin KIF9 is important for progressive motility in mouse
spermatozoa. FASEB J *34*, 5389-5400.

45. Mukherjee, K., and Brocchieri, L. (2013). Ancient Origin of Chaperonin Gene Paralogs Involved
in Ciliopathies. J Phylogenetics Evol Biol *1*.

46. Nachury, M.V., and Mick, D.U. (2019). Establishing and regulating the composition of cilia for
signal transduction. Nat Rev Mol Cell Biol *20*, 389-405.

47. Nguyen, Q.P.H., Liu, Z., Albulescu, A., Ouyang, H., Zlock, L., Coyaud, E., Laurent, E.,
Finkbeiner, W., Moraes, T.J., Raught, B.*, et al.* (2020). Comparative Super-Resolution Mapping of
Basal Feet Reveals a Modular but Distinct Architecture in Primary and Motile Cilia. Dev Cell *55*,
209-223 e207.

48. Ohata, S., and Alvarez-Buylla, A. (2016). Planar Organization of Multiciliated Ependymal (E1)
Cells in the Brain Ventricular Epithelium. Trends Neurosci *39*, 543-551.

49. Ohata, S., Nakatani, J., Herranz-Perez, V., Cheng, J., Belinson, H., Inubushi, T., Snider, W.D.,
Garcia-Verdugo, J.M., Wynshaw-Boris, A., and Alvarez-Buylla, A. (2014). Loss of Dishevelleds
disrupts planar polarity in ependymal motile cilia and results in hydrocephalus. Neuron *83*, 558-
571.

50. Pazour, G.J., Wilkerson, C.G., and Witman, G.B. (1998). A Dynein Light Chain Is Essential for
the Retrograde Particle Movement of Intraflagellar Transport (IFT). J Cell Biol *141*, 979-992.

51. Pigino, G. (2021). Intraflagellar transport. Curr Biol *31*, R530-R536.

52. Portman, N., and Gull, K. (2010). The paraflagellar rod of kinetoplastid parasites: from structure
to components and function. Int J Parasitol *40*, 135-148.

53. Qiao, C., Li, D., Liu, Y., Zhang, S., Liu, K., Liu, C., Guo, Y., Jiang, T., Fang, C., Li, N.*, et al.*

(2022). Rationalized deep learning super-resolution microscopy for sustained live imaging of
rapid subcellular processes. Nat Biotechnol.

54.   Ryu, H., Lee, H., Lee, J., Noh, H., Shin, M., Kumar, V., Hong, S., Kim, J., and Park, S. (2021).
The molecular dynamics of subdistal appendages in multi-ciliated cells. Nat Commun *12*, 612.

55.   Scholey, J.M. (2013). Kinesin-2: a family of heterotrimeric and homodimeric motors with diverse
intracellular transport functions. Annu Rev Cell Dev Biol *29*, 443-469.

56.   Siller, S.S., Sharma, H., Li, S., Yang, J., Zhang, Y., Holtzman, M.J., Winuthayanon, W.,
Colognato, H., Holdener, B.C., Li, F.Q*., et al.* (2017). Conditional knockout mice for the distal
appendage protein CEP164 reveal its essential roles in airway multiciliated cell differentiation.
PLoS Genet *13*, e1007128.

57.   Spassky, N., Merkle, F.T., Flames, N., Tramontin, A.D., Garcia-Verdugo, J.M., and Alvarez-
Buylla, A. (2005). Adult ependymal cells are postmitotic and are derived from radial glial cells
during embryogenesis. J Neurosci *25*, 10-18.

58.   Taschner, M., Kotsis, F., Braeuer, P., Kuehn, E.W., and Lorentzen, E. (2014). Crystal structures of
IFT70/52 and IFT52/46 provide insight into intraflagellar transport B core complex assembly. J
Cell Biol *207*, 269-282.

59.   Taschner, M., Weber, K., Mourao, A., Vetter, M., Awasthi, M., Stiegler, M., Bhogaraju, S., and
Lorentzen, E. (2016). Intraflagellar transport proteins 172, 80, 57, 54, 38, and 20 form a stable
tubulin-binding IFT-B2 complex. EMBO J *35*, 773-790.

60.   Tissir, F., Qu, Y., Montcouquiol, M., Zhou, L., Komatsu, K., Shi, D., Fujimori, T., Labeau, J.,
Tyteca, D., Courtoy, P*., et al.* (2010). Lack of cadherins Celsr2 and Celsr3 impairs ependymal
ciliogenesis, leading to fatal hydrocephalus. Nat Neurosci *13*, 700-707.

61.   Vincensini, L., Blisnick, T., and Bastin, P. (2011). 1001 model organisms to study cilia and
flagella. Biol Cell *103*, 109-130.

62.   Werner, M.E., Hwang, P., Huisman, F., Taborek, P., Yu, C.C., and Mitchell, B.J. (2011). Actin and
microtubules drive differential aspects of planar cell polarity in multiciliated cells. J Cell Biol *195*,
19-26.

63.   Williams, C.L., McIntyre, J.C., Norris, S.R., Jenkins, P.M., Zhang, L., Pei, Q., Verhey, K., and
Martens, J.R. (2014). Direct evidence for BBSome-associated intraflagellar transport reveals
distinct properties of native mammalian cilia. Nat Commun *5*, 5813.

64.   Wingfield, J.L., Mengoni, I., Bomberger, H., Jiang, Y.-Y., Walsh, J.D., Brown, J.M., Picariello, T.,
Cochran, D.A., Zhu, B., Pan, J*., et al.* (2017). IFT trains in different stages of assembly queue at
the ciliary base for consecutive release into the cilium. eLife *6*.

65.   Xu, Y., Cao, J., Huang, S., Feng, D., Zhang, W., Zhu, X., and Yan, X. (2015). Characterization of
tetratricopeptide repeat-containing proteins critical for cilia formation and function. PLoS One *10*,
e0124378.

66.   Yokoyama, R., O'Toole, E., Ghosh, S., and Mitchell, D.R. (2004). Regulation of flagellar dynein
activity by a central pair kinesin. Proc Natl Acad Sci U S A *101*.

67.   Zhang, J., Wang, H., Imhof, S., Zhou, X., Liao, S., Atanasov, I., Hui, W.H., Hill, K.L., and Zhou,
Z.H. (2021). Structure of the trypanosome paraflagellar rod and insights into non-planar motility
of eukaryotic cells. Cell Discovery *7*.

68.   Zhao, C., Omori, Y., Brodowska, K., Kovach, P., and Malicki, J. (2012). Kinesin-2 family in
vertebrate ciliogenesis. Proceedings of the National Academy of Sciences *109*, 2388-2393.

69.   Zhao, H., Chen, Q., Li, F., Cui, L., Xie, L., Huang, Q., Liang, X., Zhou, J., Yan, X., and Zhu, X.

(2021). Fibrogranular materials function as organizers to ensure the fidelity of multiciliary
assembly. Nat Commun *12*, 1273.

70. Zheng, J., Liu, H., Zhu, L., Chen, Y., Zhao, H., Zhang, W., Li, F., Xie, L., Yan, X., and Zhu, X.
(2019). Microtubule-bundling protein Spef1 enables mammalian ciliary central apparatus
formation. J Mol Cell Biol *11*, 67-77.

1097

Fig.1

Fig. 2

Fig. 3

a Ift27-Halo Kif6-GFP

(I) Antero- particles (Ift27⁺Kif6⁺)

b (II) Retro- particles (Ift27⁺Kif6⁺)

c Ift27-Halo Kif6-GFP

Antero- particles (Kif6⁺ or Ift27⁺)

Retro- particles (Kif6⁺ or Ift27⁺)

d
Anterograde
- Ift27⁺Kif6⁺ 24.4%
- Kif6⁺ 45.8%
- Ift27⁺ 29.8%
- n=131

Retrograde
- Ift27⁺Kif6⁺ 28.3%
- Kif6⁺ 40.7%
- Ift27⁺ 31.0%
- n=113

e
n=32 60 39   41 46 36

Velocity of particles (μm/s)

P=0.51   P=0.50
P=0.54   P=0.64
P=0.89   P=0.87

Ift27⁺Kif6⁺ | Kif6⁺ | Ift27⁺ | Ift27⁺Kif6⁺ | Kif6⁺ | Ift27⁺

Anterograde   Retrograde

Fig. 4

a Kif6 / Kif9

Exon #: 1 2 13 14 23
WT allele ATG F1 R1 TGA
~139.0 kb

Exon #: 1 14 23
KO allele F2 R2

Uncoding region
Coding region

Kif9
Exon #: 1 2 3 17 18 21
WT allele ATG F3 R3 TAA
~34.0 kb

Exon #: 1 2 18 21
KO allele F4 R4

b Kif6: +/+ +/- +/+   Kif9: +/+ +/- -/-

c Ependyma / Trachea
Kif6: +/+ +/- -/-   +/+ +/- -/-   kDa
Kif6 — 100
Gapdh — 35

Testis / Lung
Kif9: +/+ +/- -/-   +/+ +/- -/-   kDa
Kif9 — 100
Gapdh — 35

d Kif6⁺/⁺ (P45)   Kif6⁻/⁻ (P45)
Top view / Side view   2cm

e Kif9⁺/⁺ (P130)   Kif9⁻/⁻ (P130)
Top view / Side view   2cm

f Kif6 Anterior → Posterior
+/+ (2M) / -/- (2M) / +/+ (4M) / -/- (4M)   5mm

g Kif9 Anterior → Posterior
+/+ (5M) / -/- (5M) / +/+ (6M) / -/- (6M)   5mm

h

Genotype		Normal Mice (n)	Enlarged Mice (n)	Total
Kif6	+/+	60	0	60
	-/-	0	60	60
Kif9	+/+	9	0	9
	-/-	5	6	11

i Viable mice (%)
Kif6⁺/⁺ (n=56)
Kif9⁺/⁺ (n=20)
Kif9⁻/⁻ (n=20)
Kif6⁻/⁻ (n=56)
0 60 120 180 240 300 360 (day)

k Sperm trajectories Frames: 1-50
Kif6⁺/⁺ (P88)
Kif6⁻/⁻ (P88)   10μm

j Sperm
Kif6⁺/⁺ (P88)   10μm
Kif6⁻/⁻ (P88)

Fig. 5

a Brain → Ependyma / Hippocampus → EM → Live imaging (SiR-tubulin)

Effective stroke / Recovery stroke

b *Kif6*: +/+   -/-
P4   P4
P30   P30, S

c Side view   Effective stroke ←→ Recovery stroke
+/+ (P30)   0 (ms) 10 20 30 40   50 60 70 80   0-30ms
Kif6: -/- (P30,S)   0 (ms) 10 20 30 40   50 60 70 80   0-30ms

d Mice=5   5   P=0.35   Ciliary beat frequency (Hz)   *Kif6^{+/+}*   *Kif6^{-/-}*

e SiR-tubulin   *Kif6^{+/+}* (P31)   Frame:1 2 3   *Kif6^{-/-}* (P31, S)   *Kif6^{-/-}* (P57, M)

f Top view   Effective stroke ←→ Recovery stroke
+/+ (7M)   0 (ms) 10 20 30 40   50 60 70 80
Kif9: -/- (7M)   0 (ms) 10 20 30 40   50 60 70 80   0-40ms

g SiR-tubulin   Frame:1 2 3   *Kif9^{+/+}* (7M)   *Kif9^{-/-}* (7M)

h Bead tracks   *Kif6^{+/+}* (P36)   *Kif6^{-/-}* (P36, S)   *Kif6^{-/-}* (P48, M)

i Mice=3   3   P<0.0001   Beads velocity (μm/s)   *Kif6^{+/+}*   *Kif6^{-/-}*

Fig. 6

Fig.S1

a

b

```
                                                        Motor domain                                        Amino acid
Kif6  KQTIQIFARVKPTVRKQQQGIYSIDEDEKLTHSLEIVLPRDLADGFVNNKRESYKFKFQRIFDQEAKQEIIFEIIAKPVA     80
Kif9  RRKVQAFVRVRPT---DDFAHEMIKYGED-NKSIDIHLKKDTRRGVVNNQQTDWSFKLDGVLHN-ASQDLVYETVAKDAV      75

      ESTLAGYNGTIFAYGQTGSGKTFTITGGAERYSDRGIIPRTLSYIFEQLQKDSSKIYTTHISYLEIYNECGYDLLDPRHE    160
      SQALDGYNGTIMCYGQTGAGKTYTMTGATENYKHRGILPRALQQVFRMIEERPTHAITVRVSYLEIYNENLFDLLSTLPY    155

      ASKLEDLPKVTILEDPDQNIHLKNLSLHQATTEEEALNLLFLGDTNRMIAETPMNQASTRSHCIFTVHLS--SKEPGSAT    238
      VGP--SVTPMTIVENP-QGIFIKGLSVHLTSQEEDAFSLLFEGETNRIIASHTMNKNSSRSHCIFTIYMEAHSRTLSDEK    232

      VRHAKLHLVDLAGSERVSKTGVGGLLLTEAKYINLSLHYLEQVIIALSEKHRTHIPYRNSMMTSVLRDSLGGNCMTTMIA    318
      YITSKINLVDLAGSERLSKTGSEGRVLKEATYINKSLSFLEQAIIALGDQNRDHVPFRQSKLTHALKDSLGGNCNMVLVT    312

      TLSLEKRNIDESISTCRFAQRVALIKNEAILNE    351
      NIYGEAAQLDETLSSLRFASRMKLVTTEPAINE    345
```

Fig.S2

Fig. S3

Fig. S4

a

Cross section | Longitudinal section

+/+ (P47) (9+2)

Kif6: -/- (P32, S) (9+2)

CP
BB   TZ
200nm

b Top view Effective stroke ← → Recovery stroke

+/+ (P62) 27 Hz
0 (ms) 10 20 30 10μm
40 50 60 70
80 90 100 0-40ms

Kif6: -/- (P62,M) 28 Hz
0 (ms) 10 20 30
40 50 60 70
80 90 100

c SiR-tubulin *Kif6:* +/+   -/-

P30 | P30, S Frame:1 2 3
P62 | P62, M 10μm

d Side view Effective stroke ← → Recovery stroke

+/+ (6M) 28 Hz
0 (ms) 10 20 30 10μm
40 50 60 70
80 90 100 0-40ms

Kif9: -/- (6M) 28 Hz
0 (ms) 10 20 30
40 50 60 70
80 90 100

e SiR-tubulin *Kif9:* +/+   -/-

8M | 8M Frame:1 2 3
10μm

f

Mice=3   3
P=0.35

Cilary beat frequency (Hz)

Kif9$^{+/+}$ *Kif9*$^{-/-}$

Fig. S5

a Cep164/Centriolin

+/+ (6M)

Kif9: -/- (6M)

2μm   1μm

b Cep164/Centriolin   *Kif9:* +/+ (6M)   -/- (6M)

5μm

c Cep164/Centriolin

+/+ (P35)

Kif6: -/- (P35, S)

2μm   1μm

d Cep164/Centriolin   *Kif6:* +/+ (P35)   -/- (P35, S)

5μm

e

Mice=3   3   3   3

P=0.90   P≤0.0001

Mean vector length per cell

+/+   -/-   +/+   -/-(S)

Kif9   *Kif6*

Fig. S6

a *Kif6*$^{+/+}$ (P62)   *Kif6*$^{-/-}$ (P62, M)   *Kif6*$^{-/-}$ (P43, S)

ZO1/Cep164

10 µm

b

S_{patch}
S_{cell}

a
b

• Center of apical surface
• Center of BB patch

BB patch displacement: a/b
BB patch distribution: S_{patch}/S_{cell}

c

Mice= 3   3   3

P=0.15

P=0.94

BB patch displacement (a / b)

0.8
0.6
0.4
0.2
0.0

Kif6: +/+   -/- (M)   -/- (S)

d

Mice= 3   3   3

P<0.0001

P=0.75

Relative BB patch area (%) (S_{patch} / S_{cell})

Kif6: +/+   -/- (M)   -/- (S)