Evaluating diversionary feeding as a method to resolve conservation conflicts in a recovering ecosystem.

List of Authors: Jack A. Bamber 0000-0002-1699-0083 (1), Kenny Kortland (2,3), Chris Sutherland 0000-0003-2073-1751 (4), Ana Payo-Payo 0000-0001-5482-242X (1, 5), Xavier Lambin 0000-0003-4643-2653 (1*).

1 School of Biological Sciences, University of Aberdeen, AB24 2TZ Aberdeen, Scotland, UK.

2 Forestry and Land Scotland, Smithton, Inverness, Scotland, UK

3 Cairngorms Connect, Achantoul, Aviemore, Scotland, UK.

4 Centre for Research into Ecological & Environmental Modelling, University of St Andrews, St Andrews, Scotland, UK.

5 Departamento de Biodiversidad, Ecología y Evolución, Universidad Complutense, 28040 Madrid, Spain.

Keywords: Predator management, Predator control, Conservation intervention, Western capercaillie, Depredation, Pine Marten, Non-lethal control, conservation conflict.

Abstract

1.   The recovery of mammalian predators of conservation concern in Europe is a success story,

but their impact on some prey species of conservation concern may cause conservation

dilemmas. This calls for effective intervention strategies that mitigate predator impacts

without compromising their recovery.

2.   We evaluated diversionary feeding as a management intervention tool to reduce depredation

on nests of rapidly declining Western capercaillies in Scotland. We studied the influence of

diversionary feeding provision on the fates of artificial nests deployed using a replicated and

representative randomised landscape-scale experiment. This comprised paired control (no

diversionary feeding) and treatment (diversionary feeding applied) sites, each with six artificial

nests distributed across 600 km^2. The experiment was conducted over two years, and in the

second year, the control-treatment pairs were swapped.

3.   Diversionary feeding substantially reduced depredation of artificial nests, translating into an

83% increase in predicted nest survival over 28 days of incubation. The increase in survival

was mostly accounted for by a reduction in the probability that a pine marten, the main nest

predator, consumed or cached eggs. Diversionary food also significantly reduced nest

predation by badgers, although the magnitude of this effect varied by year.

4.   Diversionary feeding is an easily employable method shown in this study to reduce predator

impact (functional) without lethal (numerical) intervention. Managers should proceed with its

application for conserving capercaillie in Scotland without delay.

Introduction

Many conservation interventions lack sufficient evidence of effectiveness before being implemented, despite multiple pressing issues requiring effective intervention owing to limited resources and vast demands. This often leads to inefficient and ineffective management that fails to achieve, or worse, are detrimental to, broader conservation goals. The result is often conflict among managers, stakeholders, scientists, and the public.

With legal protection and reduced lethal predator control, the recovery of many mammalian predators in Europe is a success story for conservation. However, many of these predators exploit a range of prey, some of which are also of conservation concern or financial interest, leading to conservation conflict (Redpath et al., 2013). Lethal control of generalist predators is widely used as a management intervention, aiming to maximise a harvestable surplus or to improve the conservation status of declining species (Gibson, 2006). Lethal control is widely accepted when it eradicates or controls some damaging non-native predators (Zavaleta et al., 2001). In contrast, lethal control of native predators to protect native species is ethically debatable and has often been shown to be ineffective, except at small spatial and short temporal scales.

Many generalist predators readily compensate for losses through increased immigration and reproduction, leading to only short-term reductions (months) in predator density unless control is applied continually, with culling foxes during breeding periods showing recovery by the following February (Lieury et al., 2015)). Accordingly, many programmes cannot achieve the standards required to have a substantial impact (Kämmerle and Storch, 2019). Additionally, protected predators cannot easily be lethally controlled, requiring relevant authorities to change conservation goals, policy, and licensing (Sainsbury et al., 2019). This means lethal control may only extend to a portion of the predator guild, and the most impactful species may not be influenced. Culling also disrupts the predator guild, with consequent changes in the behaviour and density of non-target species (Rees et al., 2023) while disrupting regulatory ecosystem services that some predators provide (Sheehy et al.,

2018; Williams et al., 2018). Moreover, even for species that are not presently protected, ethical and

public perception issues are associated with the acceptability of lethal control of predators (Santiago-

Ávila et al., 2018). Therefore, interventions to alter the impact of predation, as opposed to

interventions that seek to reduce the abundance of predators, may be more effective for

conservation.

The diet of mammalian generalist meso-predators is often dominated by the most abundant prey

within the system, typically small mammals (e.g., voles), but in the face of episodic scarcity of (cyclic)

primary prey species, prey switching results in increased predation on alternative preys, such as

ground-nesting birds and their nests (Kjellander and Nordström, 2003). Increased nest loss to

generalist predators has been implicated as contributing to the decline of multiple ground-nesting

birds, including forest grouse and waders (Ewing et al., 2022; Ibáñez-álamo et al., 2015). Roos et al.

(2018) systematic review found compelling evidence that ground-nesting seabirds, waders, and

gamebird populations can be suppressed by predation. Specifically, populations of long-lived species

with high adult survival and late onset of breeding are more likely to be impacted by predation.

Many ground-nesting birds have adaptations to reduce the impact of nest predation, such as laying

large clutches and re-laying, which reduces variation in population growth rate (Lima, 2009, 1987). Via

these adaptations, native predators and prey have historically coexisted. However, the level of nest

loss a population can withstand depends on its demographic and ecological context (Banks, 1999). For

example, if nest depredation rates are increased under human influence, whether owing to the

elevated density of generalist predators (e.g. via food subsidy (Pringle et al., 2019) or lack of

competitors (Petty et al., 2003)) or more vulnerable nests (e.g. a reduction in safe nesting habitat

(Kaasiku et al., 2022)) and compounded by deleterious climate and habitat influences (Ibáñez-Álamo

et al., 2015) losses may become additive. Therefore, there is a threshold below which decline will

eventually lead to deterministic extinction and where management intervention may be warranted

or, in some cases, is the only option.

One promising non-lethal management option is diversionary feeding: the deliberate provisioning of

food to change the behaviour of target species and reduce unwanted behaviour (Kubasiewicz et al.,

2016) by exploiting the propensity of foraging individuals to exploit the most easily accessed resources

(Pyke, 1984). It has been used to reduce the predation impact of single predator species, such as red

kites (*Milvus milvus*) preying on lapwings (*Vanellus vanellus)* (Mason et al., 2021) or kestrels (*Falco*

tinnunculus) exploiting little terns (*Sternula albifrons*) (Smart and Amar, 2018).

A notable 22-year-long diversionary feeding management trial, supplemented with a short-term

experiment, in a boreal forest landscape (Norway) resulted in increases in black grouse (*Tetrao tetrix)*

and Western capercaillie (monitored via brood counts with pointer dogs), attributed to a reduction in

fox predation (Finne et al., 2019). A similar experiment that provisioned foxes with dog food

(Lindström et al., 1987) found decreases in nest predation during cyclical vole crashes. These studies

are valuable starting points for understanding how diversionary feeding may influence nest predators

specifically. However, as noted by Kubasiewicz et.al (2016), the success of diversionary feeding is often

species and context-specific. Therefore, assessing species not evaluated in previous studies is crucial

for understanding the effectiveness of diversionary feeding for multiple species. Landscape-scale

experimental evidence is limited yet vital in establishing how diversionary feeding can function as a

widely applied conservation intervention to alleviate nest predation pressure.

Forest grouse species' (Tetraonidae) are the focus of much interest from a game and conservation

management perspective. Culling nest predators is often promoted as a key intervention for grouse

population maintenance (Fletcher et al., 2010). One species with a significant conservation focus

across Europe is the Western capercaillie *(Tetrao urogallus)*. In Scotland, several well-funded

conservation initiatives have failed to durably halt the pronounced decline of capercaillie since the

1970s, with evidence of a further 50% reduction from 2016 to 2020 and extinction is predicted within

the next 50 years (Baines and Aebischer, 2023). Climate change is the likely ultimate driver of decline

through reduced food sources for chicks and hens (Wegge et al., 2022). However, multiple proximate

factors are also implicated in the decline (i.e. fence collisions (Baines and Summers, 1997)), including

a significant impact of predation of nests and chicks (Summers et al., 2004).

Lethal control of foxes and crows is common practice across many shooting estates in Scotland; even

with this intervention, capercaillies have disappeared from all but a few shooting estates, with the

core remnant populations now found in regions where predator control is not carried out (Baines and

Aebischer, 2023). In contrast, two other potential grouse predators, badger (*Meles meles)* and pine

marten (*Martes martes,* marten hereafter), are UK-protected species and cannot be routinely

controlled in the same manner (MacPherson and Wright, 2021). A correlative study in Scotland

implicated martens in the decline of capercaillie (Baines et al., 2016). As a result, lethal control of

martens has now been suggested as a possible capercaillie conservation option. However, tension

exists because this would risk undoing conservation gains (marten recovery) whilst also requiring

significant scale, effort, and cost to overcome compensation through immigration. Given these

legislative restrictions, practical difficulties, a lack of scientific consensus on efficacy, and the intraguild

complexities resulting from the disruption of predator communities, there is an urgent need for

alternatives, such as diversionary feeding, to be evaluated. Particularly given its potential to influence

the behaviour of multiple members of the predator guild simultaneously.

Considering that evidence on the effectiveness and practicalities of diversionary feeding has been

mixed, we respond to a need to evaluate the extent to which diversionary feeding could decrease nest

predation by a guild of predators and to assess the practicalities and feasibility of diversionary feeding

as a management tool. We do this through large-scale, experimental deployment of diversionary

feeding paired with control sites with no feeding. Specifically, we focused on the protected marten as

an important nest predator and the critically endangered capercaillie. Our experiment compared

artificial nest survival in a control and treatment design to evaluate how diversionary feeding

influences the rate at which nests are depredated. We also aimed to establish how the distance of

nests from feeding sites may influence the effectiveness of diversionary feeding. Our experimental

approach provides a robust, accurate, and comparable index of predation change, with nest failure

purely being related to predation pressure and not alternative factors, such as nest abandonment due

to adverse weather, as may be true with actual nests.

Materials and Methods

2.1 Study Area

This study was conducted in the Cairngorms Connect landscape (FLS, Wildland Ltd, RSPB, Naturescot),

a 600km^2 ecological restoration project on the western side of the Cairngorms National Park, Scotland

(57°09'47.5"N 3°42'47.0" W, Figure 1). The landscape consists of remnant Caledonian and plantation

pine forests (mainly *Pinus sylvestris*), with a mixture of bogs, heaths, and some deciduous woodlands.

Management includes intense culling of *Cervidae* (red and roe deer) to allow forest regeneration, and,

unlike in more traditional neighbouring estates, there is no control of predators. The area

encompasses the core of the remaining population of Scottish capercaillie (Baines and Aebischer,

2023). The predator community includes badger, fox, marten, carrion crow (*Corvus corone)*, common

buzzard (*Buteo buteo)*, and ten scarcer raptor species.

2.2 Experimental Design

We performed a randomised landscape-scale experiment with paired control and treatment sites

swapped between years. The sampling units were 60 paired, 1km^2 square grid cells restricted to

forested areas (National Forest Index, min of 1.7 hectares of forest cover) and the size of the grids was

chosen to encompass the typical daily home range of a marten (49 ha in females and 54 ha in males

(Zalewski et al., 2004)). Diversionary feeding treatment was assigned to all but 8 cells randomly, and

the paired control cell was selected to be approximately>1km^2 from its matching diversionary feeding

cell to maximise treatment independence, using empty cells as spacing (Figure 1). The centre of

treatment cells had a feeding station (see below), and each pair of treatment and control cells

contained six artificial nests as response variable (see artificial nests below). Due to constraints

resulting from the convoluted shape of the study area, 8 cells with edges closer than 1 km were given

the same treatment to maintain the independence of treatment. The response variable to treatment

was the fate of six artificial nests. The experiment was conducted over nine weeks between 24[th] April

and 1[st] July, coinciding with tetraonids' nesting, re-nesting, and early brooding periods (Summers et

al., 2004) in 2021 and 2022. Diversionary feeding and control treatments were swapped within pairs

between years.

2.3 Diversionary Feeding Stations

To maximise the applied relevance of the experiment, we provided supplementary by-products from

ongoing deer culling. This is a resource known to be consumed by predators when left in situ as

gralloch (organs left after culling) and carcasses, which was more cost-effective than provisioning

other food sources. The nine-week timeframe selected for our feeding provisioning was deliberately

short to avoid any numerical predator response, a potential risk with diversionary feeding

(Kubasiewicz et al., 2016). The feeding period also coincides with the time of more abundant food

(Spring), meaning that diversionary food is more likely to provide an alternative, not a supplement.

By-products from deer culling presented in times of food scarcity (winter) are more likely to be

supplements (Whitney et al., 2018)

Inception of diversionary feeding was timed close to egg laying to avoid any increase in predator

abundance within the vicinity of feeding stations after territories and breeding decisions of predators

were likely fixed. Feeding stations were deployed within ~100m of the centre point of grid cells and

were replenished every two weeks with ~10kg of deer carrion (for 8 weeks), and the remaining food

weights were recorded with a spring scale to monitor depletion. Replenishment ensured food was

always available, even if predators were to have found decaying meat unappealing (Moleón and

Sánchez-Zapata, 2021). When possible, food movement and the distance from central feeding stations

were also recorded.

All feeding stations were monitored using remote camera traps (Browning Recon Force Advantage

model: BTC-7A), set to record three-shot bursts with a five-second interval between captures to

establish uptake of diversionary feeding treatment by target and non-target species. These settings

were selected to allow detection of predators quickly moving through sites and removing food. We

considered an independent detection as an image taken within >30 minutes of each other.

2.4 Artificial Nests

We used the fate of artificial nests rather than nests of wild capercaillie and black grouse as the

response variable owing to the scarcity of these focal prey and to minimise disturbance.

Approximately 7 days after establishing the feeding sites, we constructed 3 artificial nests, containing

7 eggs, in each control and treatment grid (N=180). The three nests were placed 100m, 300m and

500m from the centre of the cell. A second deployment of nests, mimicking re-laying, occurred at the

predation date or hatch date of the first deployment at the same distance interval from the cell centre

but 50-100m away from the previous nest to avoid predator bias. Secondary nests contained only 3

hen eggs plus one wax egg, mimicking the lower clutch size seen in relaying (Storaas et al., 2000) and

were not replaced once depredated or if they survived 28 days. Meaning 360 nests were deployed

each year. All nests were checked every 14 days, with two visits spanning the 28-day capercaillie

incubation. Checks were conducted visually from 3-5 m away. The location of the artificial nests at

each distance was determined using a randomly generated compass bearing from the centre of the

grid at the specified distance.

The nests were made to resemble actual capercaillie nests: a shallow depression at the base of a tree

filled with plant material, covered with dwarf bushes twigs to mimic the visual camouflage an

incubating hen provides. We ensured that at least 1.5 eggs were visible to an observer standing 3-5 m

away from the nest to allow for nest detection by a visual (avian) predator. Each artificial nest had six

small domestic hen (*Gallus gallus domesticus)* eggs, resembling capercaillie eggs in size and colour

(Mortola and Al Awam, 2010; Rosenberger et al., 2017). A 7[th] egg was drained and filled with "Parasoy"

soy wax blend to aid in identifying predators through tooth and bill marks. Wax eggs had a 50cm clear

fishing line attached to a 3mm nail inside the egg, with the other end tethered to the ground with a

30cm tent peg. To reduce human scent that may affect discovery rates (Weldon, 2021), eggs were

stored on pheasant feathers for seven days before deployment, rubber gloves were worn when

handling eggs, and rubber boots and field clothes worn during deployment were stored outdoors.

If a nest site was disturbed or the 1.5 visible eggs could not be seen during nest checks, the nest was

inspected in more detail. Depredation was deemed to have occurred if any hen egg was damaged or

removed. Field signs, including marks on wax eggs and patterns of nest disturbance, were recorded to

ascertain which predator was likely responsible (Summers et al., 2004). The assumed predator was

assigned for each depredation event before checking camera trap data to allow unbiased validation.

In addition to the 14-day visual checks, camera traps were set up at one-third (60, per treatment, per

year) of nests (Browning Recon Force Advantage model: BTC-7A or Browning Recon Force Elite Model:

BTC-7E-HP4). This allowed the specific identification of nest predators and was used to validate field

interpretation of wax egg markings and nest signs. Cameras trained on artificial nests were set to

record 10-second-long videos and distributed equally, but randomly, between treatments and

distances. Thus, 10 nests per distance from the cell centre had a camera for each treatment.

Figure 1. Illustration of the experimental design. The main map shows the forested areas of the Cairngorms connect landscapes, which was our scope of inference. The highlight region zooms into the sampling grid cell design (squares), showing the division of the sampling area into 1km^2 suitable sample grids (based on being within the CC area and containing the forest layer). The primary control and test deployments are colour coded, with feeding sites (2021) in red and control sites (2021) in blue. The bottom left 1km2 area shows an example of a sample grid showing the internal structure of a diversionary feeding site, including the general nest structure and a central feeding station. Control sites mimicked the structure without the feeding station. An exemplary artificial nest with heather cover is illustrated.

Data extraction

Camera trap photos taken at feeding stations were identified at the species level using the metadata

tagging software DigiKam 7.3.0, following the 'camtrapR' workflow (Niedballa et al., 2016). Videos

taken at artificial nests were viewed, and species-specific detection histories were generated manually

by recording the time and date of a depredation event. The assumed responsible predator of

depredated nests for nests without cameras was inferred from field signs. Video footage from camera

traps allowed validation of assumed predators from signs with confirmed predators on camera traps,

showing a 99% (N= 109) success rate in correctly identifying nest predators when an assumed predator

was assigned (See Appendix 1). We excluded 15 nests, from further analysis, known to have been

depredated by a fox (n=2), corvids (n=7), or rodent (n=6) owing to low sample size and 28 nests

depredated by non-identified species as they likely formed a heterogeneous group (12 Test: 31

Control).

<u>Statistical analysis</u>

We modelled the three fates of artificial nests using multinomial logistic regression: survived 28 days,

depredated by marten, and depredated by badger. We analysed the multinomial responses using

covariates that reflected the experimental design: diversionary feeding treatment to quantify the

effect of diversionary feeding, distance from the grid centre to quantify any spatial decay of the

feeding effect, and interaction between distance and treatment. We also included year as a fixed

effect to account for annual variation, e.g., in field vole prey abundance. We added grid cell identity

(n=60) as a random effect to account for any local influences on nest predation, such as local predator

abundance and habitat, across the two years of study.

Multinomial models were implemented using a generalised additive model (GAM) in package 'mgcv'

(Wood, 2017). We used the 'mn' response model as it allows the inclusion of random effects. We used

1000 simulations from the model using the function "Predict" to produce population-level (marginal)

estimated fate probabilities and confidence intervals. All statistical analysis was performed using R

(version 4.1.3).

Results

<u>Uptake and usage of feeding stations</u>

On average, 57.5kg (range: 38 - 81 kg) of deer meat was deployed per feeding station each year, with

an average of 10.5kg (range: 6 - 14 kg) added every two weeks according to the weighed assessment

of depletion at restocking visits.

Over 340,000 photographs were collected from the 60 feeding stations across 3,912 camera trap days.

They include 142,179 images of potential egg predators recorded in 3,726 independent visits. Martens

accounted for 13.0% of these visits (19374 images, 486 independent detections), badgers for 22.5%

(54024, 839), and foxes for 10.3% (12946, 383). We also detected avian nest predators: 31837 (1306,

35.0% detections) of the visits were by raptor sp. (buzzard, golden eagle, red kite) and 20410 by corvids

(639, jay, crow). Fifty-four out of 60 feeding stations were visited by predators (86.6% in 2021, 93.3%

in 2022). Badgers, foxes, and buzzards visited similar proportions of feeding stations (58%, 57%, and

55%, respectively), followed by martens (43.33 %) (See Appendix 2). As predators had moved deer

carrion away from the camera fields of view at 132 restocking visits (n= 240), it was not possible to

estimate the amount of diversionary food consumed by each species.

<u>Nest fate with diversionary feeding</u>

Forty-nine percent of artificial nests (353/720) survived the full 28 days, meaning that 51% of nests

experienced depredation. Recorded nest predators were martens (38%, 268), badgers (6.9%, 50), and

other species (unknown, fox, corvid, and rodent (6.8%, 49)). However, fewer nests survived in control

sites (128, 35.5%) than in diversionary feeding sites (228, 63.3%), reflecting fewer nests depredated

by martens (Control: 170, Test: 98) and badgers (Control: 31, Test 19; (See Appendix 3).

Multinomial logistic regression revealed that nest fates were associated with the three experimental

variables (treatment, distance, and year) but in different ways, see Table 1. The probability of marten

depredation was substantially reduced when that nest was at a treatment site relative to a control

site (-1.494, +/-0.309, p<0.00), and this did not vary between years (-0.007, +/-0.172, P= 0.969). There

was an increase in the probability of depredation by marten with increasing distance between the

nests and the feeding site. However, this effect was small and was only evident at the furthest distance

of 500m (0.698 +/-0.424, P= 0.099). The probability of badger depredation was also substantially and

significantly reduced by diversionary feeding (-1.723 +/-0.622, p=0.006), with a significant additive

influence of year reflecting higher badger depredation in 2022 (0.694 +/-0.325, P=0.033). Distance,

with an interaction between treatments, did not show any significant effect. Predictions were based

on the provision of diversionary food; we can see an increase in the mean predicted probability of

nests surviving from 0.406 (CI 0.303- 0.523) up to 0.744 (CI 0.645- 0.828), an increase of 83%. This

change occurs mainly due to the change in the predicted probability of marten depredation, reducing

in value from 0.52 (CI 0.40 0.64) in control to 0.22 (Ci 0.151- 0.318) with diversionary food provision.

There is also a change in predicted predation by badgers, with diversionary food treatment reducing

predictions from 0.085 (CI 0.0187-0.227) control to 0.03 (0.005- 0.098) in 2021 and an increase in

overall predicted badger predation in 2022 to 0.15 (CI 0.04-0.366) control and with 0.058 (CI 0.011-

0.175) however, confidence intervals of badger overlap. See Figure 2.

When assessing the fate of all nests (N=720), including those not included in the multinomial model

due to uncertain or sparsely represented fates, Kaplan-Meier analysis showed an increase in survival

with treatment. Estimated 28 days survival of control nests was 0.345 (CI 0.281- 0.422), and that of

nests in diversionary feeding treatment was 0.650 (CI 0.585-0.723), an 88% increase in line predictions

from the multinomial model (See Appendix 4).

Table 1. Coefficients of multinomial logistic regression showing changes to log probability of fate of artificial nests. The reference fate for the model is that the nest survived for 28 days. Significant results are highlighted in bold.

	Dependant Variable (Fate)							
	Pine Marten (B)				Badger (C)			
Independent Variables	Coefficient	Std. Err.	z	P>|z|	Coefficient	Std. Err.	z	P>|z|
Intercept	0.278	0.235	1.184	0.237	**-1.794**	**0.409**	**-4.390**	**1.14e-05**
Treatment	**-1.494**	**0.309**	**-4.832**	**1.35E-06**	**-1.723**	**0.622**	**-2.771**	**0.006**
Distance 300m	0.005	0.285	0.019	0.984	-0.547	0.533	-1.027	0.304
Distance 500m	0.206	0.297	0.691	0.489	0.319	0.473	0.675	0.499
Year (2022)	-0.007	0.172	-0.038	0.969	**0.694**	**0.325**	**2.137**	**0.033**
Treatment*500m	0.155	0.425	0.366	0.715	1.170	0.839	1.394	0.163
Treatment*300m	0.698	0.424	1.647	0.099	0.736	0.795	0.926	0.355

Fate A (Survived) Calculated as Difference in Interecept (100m, Control, 100*Control, 2021)
Significant Results in Bold

Figure 2. Multinomial logistic regression results show the predicted probability of nest fate: depredated by badger, depredated by marten, and survived. Predictions were created for nests at 100m from grid centres as the baseline for predictions. The year was not shown to influence the fates pine marten or survived with predictions based on 2021. We present predictions for Badger for both 2021 and 2022, as it was highlighted to have a significant difference within multinomial modelling (2021 circle, 2022 triangle). Treatment is shown with control in black and test in red 97.5% confidence intervals are shown via error bars.

Discussion

We evaluated how diversionary feeding alters the rates of nest depredation by meso-carnivores in a

boreal forest landscape. We found that diversionary feeding almost halved depredation rates of pine

martens and badgers on artificial nests over the 28-day incubation period of tetraonid grouse.

Predicted nest survival probability increased from 0.406 to 0.733 with the provision of diversionary

feeding. Using a fully randomised, well-replicated, landscape-scale experiment allows us to infer

causality between short-term provisioning of diversionary feeding and reduced depredation of

artificial nest predation. Should they extrapolate to real capercaillie nests, our results present

diversionary feeding as a viable non-lethal option for reducing nest predation on ground-nesting birds

of conservation concern across the boreal zone, providing a form of control of the impact of predator

presence without sacrificing ecosystem benefits, or garnering negative public interaction. The

presence of a validated alternative to lethal intervention raises questions as to whether practitioners

have the social licence to cull one protected species for the protection of another.

<u>Intervention for Protected Predators</u>

Diversionary feeding almost halved both marten and badger depredation rates but from different

baselines. Because of these two species' reduction in nest depredation, artificial nest survival

increased by 83% (Figure 2). The risk of depredation by marten was five times higher than badgers.

Hence, the reduction of pine marten impact is proportionally greater for nest survival. This is despite

martens having confirmed access at only 43% of feeding stations from cameras. This could be due to

the localised redistribution of diversionary food into the area surrounding feeding stations by other

predators, with large pieces of carrion found up to 50m away from feeding stations, causing

imperfect detection. Similar food redistribution around carcasses by badgers and foxes has been

seen at a maximum of 103m (Young et al., 2015).

Both badger and marten numbers have been suppressed historically due to persecution. Marten was

driven to localised extinction but has recovered since the first re-sightings in the study area in 1994. It

has been suggested that this recovery and the perceived high density of pine martens may be the

reason for unsustainably high nest predation rates. Surveys in 2012 estimated 0.07 - 0.38 individuals

per km^2 using spatially explicit capture-recapture of non-invasively collected hair in sites within our

research area (Kubasiewicz et al., 2017). A 2020 survey using similar field and analytical methods

indicates no rise in density Hobson et al. (2023). These values are within density estimates for pine

marten elsewhere, such as Bialoweiza Forest, ranging from 0.363-0.757 individuals per km^2 (Zalewski

and Jedrzejewski, 2006). This indicates that modifying marten feeding behaviour through diversionary

feeding is a realistic, sustainable, and evidence-based alternative to lethal control, especially if marten

populations are not above normal densities.

Land manager perceptions (before this experiment) are that badger numbers have increased; while

no formal estimates have been presented, we found that badgers were widely distributed, accessing

58% of feeding stations and depredating artificial nests in a pine forest, where historically, they were

assumed mostly absent. This could reflect a recovery by badgers and another reinstated source of

predation pressure that could also be halved by diversionary feeding.

<u>Interpreting artificial nest data</u>

There are known caveats to interpreting absolute predation rate on artificial nests and necessary

conditions for interpreting differences in predation rate in an experimental context such as ours. To

avoid overemphasising some predators' impact and underestimating others' impact, the predators

responsible for actual nest predation must match those of actual nests (Pärt and Wretenberg, 2002).

For marten, the main predator in this study, we can be confident that changes seen in nest predation

may translate to actual nests, as they have been seen to predate actual nests at a similarly high rate

in other studies. Summers et al. (2009) sampled actual capercaillie nests in the same region with

camera traps (N=22); in this instance, all confirmed losses were due to marten predation.

Consideration of other studies on predation rates on the capercaillie nests reveals that the fate of

artificial nests in this study aligns with what is seen elsewhere in their range. Predation rates of

capercaillie nests in a stable population in southern Norway ranged from 48 to 90% according to the

stage of the vole cycles (Wegge and Storaas, 1990). In Scotland, observed predation rates revealed by

camera traps deployed on actual nests ranged between 42 and 68% (Summers et al., 2009). Our

estimates of 65% predation rates on our artificial nests in control sites are within these ranges, with

the observed 37% in the presence of diversionary feeding as low as the lowest value seen in Norway

in a peak vole year. Thus, present nest predation rates in Scotland are not abnormally high, especially

considering our study took place during low vole years when predation rates were expected to be at

the high end of observed rates.

In this study, fox and corvid depredation of artificial nests was lower than on actual nests studied

by(Baines et al., 2004) in the same region. The presence of camera traps at some, but not all, false

nests may have deterred foxes, given the prevailing persecution (Zalewska et al., 2021). Conversely,

our efforts to reduce human activity at our artificial nests may have precluded the inflated corvid

predation that often occurs where humans interact with nests (such as trails and markers (Picozzi,

1975)). While some raptor species influence red grouse recruitment, the impact occurs through chicks,

not nest predation (Thirgood et al., 2000), so our results are not unexpected.

Consideration of risk factors:

The duration of our experiment, from late April to early July, was chosen to reduce the risk of a

numerical response through aggregation by predators potentially subsidised by deer carrion. Predator

territories and the number of embryos were long established before the deployment of diversionary

feeding. Thus, we infer diversionary feeding changed predator foraging (functional response), not

numbers (numerical response). However, deer culling activities mainly occur in winter in Scotland. It

is not unlikely that predators' numbers are elevated by the overwinter provision of gralloch (when it

is not removed) across the landscape and before our experiments. This is most likely in years of low

vole abundance, such as in 2021 and 2022. Anecdotally, case partners in Glenfeshie have been leaving

gralloch on the landscape across all seasons for over 10 years and have not seen any negative

influences of carrion.

Increased nest survival may not directly translate to more chicks reaching adulthood, hence

productivity (Saniga, 2002). If decreased nest predation and increased chicks make a breeding area

more attractive to predators aggregating in areas with diversionary feeding, this may elevate the

predation rate (Pakanen et al., 2022). Capercaillie chick biomass is tiny relative to all other prey

exploited by martens, badgers and other mesopredators. We view this as unlikely but worthy of

further study.

Management Implications:

Using one fieldworker, this experiment covered most of Scotland's core residual range of capercaillie.

Deployment of 30 feeding stations across five land ownership areas took five days. The focus on good

experimental design for robust inference meant that deployment was labour-intensive; practical

deployment would likely be easier, with no need for strict separation and designation of "control" and

"test" sites. Using by-products from existing deer culling efforts means the cost of providing food was

low and may even have reduced the disposal cost. Based on the total food deployed at each feeding

station across our experiment, at maximum, a feeding station would require approximately 80kg of

carrion, which is the equivalent of an adult male red deer (Reby and McComb, 2003). With large

numbers of deer being culled, there is no limitation to the amount of deer viscera that could be made

available by real-time supply or freezing byproducts during peak cull periods. Our design with nests at

three distances from the food dump found no clear evidence that the depredation rate changed

significantly with increasing distance from the feeding station up to 500m. A minimum effective

density of one station per km^2 of suitable habitat was shown in this study. However, with no significant

reduction in efficacy within that distance, the spacing could be more comprehensive for practical

deployment. Logically, the influence of feeding on nest predation rate must reduce eventually, as seen

with the 1.5 km^2 separation of feeding stations to control site nests. Thus, there is little doubt that

diversionary feeding could be rolled out across the remaining range at little cost with potentially

substantial benefits to a ground-nesting species in decline. To establish if a reduction in nest predation

alone can lead to capercaillie recovery in the face of climate change, monitoring the influence of

diversionary feeding should be performed, emphasising evaluating the full impact on productivity.

Another research priority is establishing if and when supplementary and diversionary food might

inflate mesopredator abundance.

The observed substantial reduction in artificial nest depredation demonstrates the potential of

diversionary feeding as an effective non-lethal intervention for conserving ground-nesting birds. It can

be used with protected and recovering predators of conservation concern, hence alleviating

conservation conflicts. Particularly, given pine marten recolonisation following legal protection,

implementing non-lethal management action that mitigates the impact of predation is feasible and

supported by the evidence presented here. No major obstacles should exist to implementing

diversionary feeding whilst further monitoring impacts throughout the historical range of capercaillie

now shared with the native pine marten.

Author contribution:

Conceptualisation: JB, CS, XL, KK, APP; **Fieldwork**: JB; **Statistical analyses,** JB, CS; **Data curation**: JB; **Writing—original draft preparation**: JB; **Writing—review and editing**: JB, CS, KK, XL, APP; **Funding acquisition**: XL, CS, KK. All authors have read and agreed to the published version of the manuscript.

Conflict of Interest:

None.

Acknowledgements:

JB was funded under the NERC Scottish Universities Partnership for Environmental Research (SUPER) Doctoral Training Partnership (DTP) (Grant reference number NE/S007342/1, https://superdtp.st-andrews.ac.uk/). Additional funding has been provided by the School of Biological Sciences (University of Aberdeen) and Forestry and Land Scotland. We also thank Thomas MacDonell and Wildland Ltd. for their in-kind contributions and continued support. We thank all Cairngorms Connect partner organisations and the staff who supported our work; Forestry and Land Scotland (Bobby Innes, Cieran Watson, Tom Cameron), Wildland Ltd (Davie McGibbon, Donnie Ross, Ronan Dugan, Grant Ashley, Marc Willis), the Royal Society for the Protection of Birds (Richard Mason, Michael Butler, Cameron Waite, Fraser Cormack, Jack Ward) and Nature Scot, for providing site access, assistance, and equipment. We also thank Elsa Fleurial for camera trap image processing and Titouan Sanglard for field assistance.

References:

Baines, D., & Aebischer, N. J. (2023). Estimating capercaillie Tetrao urogallus population size in

Scotland from annual leks and counts of broods over the period 2010–2020. Wildlife Biology.

https://doi.org/10.1002/wlb3.01104

Baines, D., Moss, R., & Dugan, D. (2004). Capercaillie breeding success in relation to forest habitat and

predator abundance. Journal of Applied Ecology, 41(1). https://doi.org/10.1111/j.1365-

2664.2004.00875.x

Baines, D., & Summers, R. W. (1997). Assessment of Bird Collisions with Deer Fences in Scottish

Forests. The Journal of Applied Ecology, 34(4). https://doi.org/10.2307/2405284

Banks, P. B. (1999). Predation by introduced foxes on native bush rats in Australia: Do foxes take the

doomed surplus? Journal of Applied Ecology, 36(6). https://doi.org/10.1046/j.1365-

2664.1999.00463.x

Doherty, J. F., & Ruehle, B. (2020). An Integrated Landscape of Fear and Disgust: The Evolution of

Avoidance Behaviors Amidst a Myriad of Natural Enemies. Frontiers in Ecology and Evolution, 8.

https://doi.org/10.3389/fevo.2020.564343

Ewing, H., Franks, S., Smart, J., Burton, N., & Gill, J. A. (2022). Nest survival of threatened Eurasian

Curlew (Numenius arquata) breeding at low densities across a human-modified landscape. Ibis.

https://doi.org/10.1111/ibi.13180

Finne, M. H., Kristiansen, P., Rolstad, J., & Wegge, P. (2019). Diversionary feeding of red fox in spring

increased productivity of forest grouse in southeast Norway. Wildlife Biology, 2019(1).

https://doi.org/10.2981/wlb.00492

Fletcher, K., Aebischer, N. J., Baines, D., Foster, R., & Hoodless, A. N. (2010). Changes in breeding

success and abundance of ground-nesting moorland birds in relation to the experimental deployment

of legal predator control. Journal of Applied Ecology, 47(2). https://doi.org/10.1111/j.1365-

2664.2010.01793.x

Gibson, L. (2006). The Role of Lethal Control in Managing the Effects of Apparent Competition on

Endangered Prey Species. Wildlife Society Bulletin, 34(4). https://doi.org/10.2193/0091-

7648(2006)34[1220:trolci]2.0.co;2

Hobson, K., Sutherland, C., Lambin, X., 2023, The influence of forest, and land management practices

on spatial variation in density of recovering pine marten populations, PhD Thesis, University of

Aberdeen, 101

Ibáñez-álamo, J. D., Magrath, R. D., Oteyza, J. C., Chalfoun, A. D., Haff, T. M., Schmidt, K. A., Thomson,

R. L., & Martin, T. E. (2015). Nest predation research: Recent findings and future perspectives. In

Journal of Ornithology (Vol. 156). https://doi.org/10.1007/s10336-015-1207-4

Kaasiku, T., Rannap, R., & Männil, P. (2022). Predation-mediated edge effects reduce survival of wader

nests at a wet grassland-forest edge. Animal Conservation, 25(5). https://doi.org/10.1111/acv.12774

Kämmerle, J. L., & Storch, I. (2019). Predation, predator control and grouse populations: a review.

Wildlife Biology, 2019(1). https://doi.org/10.2981/wlb.00464

Kjellander, P., & Nordström, J. (2003). Cyclic voles, prey switching in red fox, and roe deer dynamics -

A test of the alternative prey hypothesis. Oikos, 101(2). https://doi.org/10.1034/j.1600-

0706.2003.11986.x

Kubasiewicz, L. M., Bunnefeld, N., Tulloch, A. I. T., Quine, C. P., & Park, K. J. (2016). Diversionary

feeding: an effective management strategy for conservation conflict? In Biodiversity and Conservation

(Vol. 25, Issue 1). https://doi.org/10.1007/s10531-015-1026-1

Kubasiewicz, L. M., Quine, C. P., Summers, R. W., Coope, R., Cottrell, J. E., A'Hara, S. W., & Park, K. J.

(2017). Non-invasive genotyping and spatial mark-recapture methods to estimate European pine

marten density in forested landscapes. Hystrix, 28(2). https://doi.org/10.4404/hystrix-00007-2017

Lieury, N., Ruette, S., Devillard, S., Albaret, M., Drouyer, F., Baudoux, B., & Millon, A. (2015).

Compensatory immigration challenges predator control: An experimental evidence-based approach

improves management. Journal of Wildlife Management, 79(3). https://doi.org/10.1002/jwmg.850

Lima, S. L. (1987). Clutch size in birds: a predation perspective. Ecology, 68(4).

https://doi.org/10.2307/1938378

Lima, S. L. (2009). Predators and the breeding bird: Behavioral and reproductive flexibility under the

risk of predation. In Biological Reviews (Vol. 84, Issue 3). https://doi.org/10.1111/j.1469-

185X.2009.00085.x

Lindström, E., Angelstam, P., Widén, P., Andrén, H., Lindstrom, E., Widen, P., & Andren, H. (1987). Do

Predators Synchronize Vole and Grouse Fluctuations?: An Experiment. Oikos, 48(2).

https://doi.org/10.2307/3565847

Macpherson, J., & Wright, P. (2021). Long-term strategic recovery plan for pine martens in Britain.

Vincent Wildlife Trust, 72.

Mason, L. R., Green, R. E., Hirons, G. J. M., Skinner, A. M. J., Peault, S. C., Upcott, E. v., Wells, E., Wilding,

D. J., & Smart, J. (2021). Experimental diversionary feeding of red kites Milvus milvus reduces chick

predation and enhances breeding productivity of northern lapwings Vanellus vanellus. Journal for

Nature Conservation, 64. https://doi.org/10.1016/j.jnc.2021.126051

Moleón, M., & Sánchez-Zapata, J. A. (2021). The role of carrion in the landscapes of fear and disgust:

A review and prospects. In Diversity (Vol. 13, Issue 1). https://doi.org/10.3390/d13010028

Mortola, J. P., & Al Awam, K. (2010). Growth of the chicken embryo: Implications of egg size.

Comparative Biochemistry and Physiology - A Molecular and Integrative Physiology, 156(4).

https://doi.org/10.1016/j.cbpa.2010.03.011

Niedballa, J., Sollmann, R., Courtiol, A., & Wilting, A. (2016). camtrapR: an R package for efficient

camera trap data management. Methods in Ecology and Evolution, 7(12).

https://doi.org/10.1111/2041-210X.12600

Pakanen, V. M., Tornberg, R., Airaksinen, E., Rönkä, N., & Koivula, K. (2022). The abundance of small

mammals is positively linked to survival from nest depredation but negatively linked to local

recruitment of a ground nesting precocial bird. Ecology and Evolution, 12(9).

https://doi.org/10.1002/ece3.9292

Pärt, T., & Wretenberg, J. (2002). Do artificial nests reveal relative nest predation risk for real nests?

Journal of Avian Biology, 33(1). https://doi.org/10.1034/j.1600-048X.2002.330107.x

Petty, S. J., Anderson, D. I. K., Davison, M., Little, B., Sherratt, T. N., Thomas, C. J., & Lambin, X. (2003).

The decline of Common Kestrels Falco tinnunculus in a forested area of northern England: The role of

predation by Northern Goshawks Accipiter gentilis. Ibis, 145(3). https://doi.org/10.1046/j.1474-

919X.2003.00191.x

Picozzi, N. (1975). Crow Predation on Marked Nests. The Journal of Wildlife Management, 39(1).

https://doi.org/10.2307/3800478

Pringle, H., Wilson, M., Calladine, J., & Siriwardena, G. (2019). Associations between gamebird releases

and generalist predators. Journal of Applied Ecology, 56(8). https://doi.org/10.1111/1365-2664.13451

Pyke, G. H. (1984). Optimal foraging theory: a critical review. Annual Review of Ecology and

Systematics. Vol. 15. https://doi.org/10.1146/annurev.ecolsys.15.1.523

Reby, D., & McComb, K. (2003). Anatomical constraints generate honesty: Acoustic cues to age and

weight   in   the   roars   of   red   deer   stags.   Animal   Behaviour,   65(3).

https://doi.org/10.1006/anbe.2003.2078

Redpath, S. M., Young, J., Evely, A., Adams, W. M., Sutherland, W. J., Whitehouse, A., Amar, A.,

Lambert, R. A., Linnell, J. D. C., Watt, A., & Gutiérrez, R. J. (2013). Understanding and managing

conservation   conflicts.   In   Trends   in   Ecology   and   Evolution   (Vol.   28,   Issue   2).

https://doi.org/10.1016/j.tree.2012.08.021

Rees, M. W., Pascoe, J. H., Le Pla, M., Robley, A., Birnbaum, E. K., Wintle, B. A., & Hradsky, B. A. (2023).

Mesopredator release among invasive predators: Controlling red foxes can increase feral cat density

and alter their behaviour. Journal of Applied Ecology. https://doi.org/10.1111/1365-2664.14402

Rosenberger, J., Deeming, D. C., Lukaszewicz, E., & Kowalczyk, A. (2017). Changes of eggshell

characteristics during embryo development in Western Capercaillie (Tetrao urogallus). Avian Biology

Research, 10(2). https://doi.org/10.3184/175815617X14878495604607

Sainsbury, K. A., Shore, R. F., Schofield, H., Croose, E., Campbell, R. D., & Mcdonald, R. A. (2019). Recent

history, current status, conservation and management of native mammalian carnivore species in Great

Britain. In Mammal Review (Vol. 49, Issue 2). https://doi.org/10.1111/mam.12150

Saniga, M. (2002). Nest loss and chick mortality in capercaillie (Tetrao urogallus) and hazel grouse

(Bonasa bonasia) in West Carpathians. Folia Zoologica, 51(3).

Santiago-Ávila, F. J., Lynn, W. S., & Treves, A. (2018). Inappropriate consideration of animal interests

in predator management: Towards a comprehensive moral code. In Large Carnivore Conservation and

Management: Human Dimensions. https://doi.org/10.4324/9781315175454

Sheehy, E., Sutherland, C., O'Reilly, C., & Lambin, X. (2018). The enemy of my enemy is my friend:

Native pine marten recovery reverses the decline of the red squirrel by suppressing grey squirrel

populations. Proceedings of the Royal Society B: Biological Sciences, 285(1874).

https://doi.org/10.1098/rspb.2017.2603

Smart, J., & Amar, A. (2018). Diversionary feeding as a means of reducing raptor predation at seabird

breeding colonies. Journal for Nature Conservation, 46. https://doi.org/10.1016/j.jnc.2018.09.003

Storaas, T., Wegge, P., & Kastdalen, L. (2000). Weight-related renesting in capercaillie Tetrao urogallus.

Wildlife Biology, 6(4). https://doi.org/10.2981/wlb.2000.030

Summers, R. W., Green, R. E., Proctor, R., Dugan, D., Lambie, D., Moncrieff, R., Moss, R., & Baines, D.

(2004). An experimental study of the effects of predation on the breeding productivity of capercaillie

and black grouse. Journal of Applied Ecology, 41(3). https://doi.org/10.1111/j.0021-

8901.2004.00891.x

Summers, R. W., Willi, J., & Selvidge, J. (2009). Capercaillie Tetrao urogallus nest loss and attendance

at Abernethy Forest, Scotland. Wildlife Biology, 15(3). https://doi.org/10.2981/08-036

Thirgood, S., Redpath, S., Newton, I., & Hudson, P. (2000). Raptors and red grouse: Conservation

conflicts and management solutions. Conservation Biology, 14(1). https://doi.org/10.1046/j.1523-

1739.2000.99013.x

Tobajas, J., Oliva-Vidal, P., Piqué, J., Afonso-Jordana, I., García-Ferré, D., Moreno-Opo, R., & Margalida,

A. (2021). Scavenging patterns of generalist predators in forested areas: The potential implications of

increase in carrion availability on a threatened capercaillie population. Animal Conservation.

Wegge, P., Moss, R., & Rolstad, J. (2022). Annual variation in breeding success in boreal forest grouse:

Four decades of monitoring reveals bottom-up drivers to be more important than predation. Ecology

and Evolution, 12(10). https://doi.org/10.1002/ece3.9327

Wegge, P., & Storaas, T. (1990). Nest loss in capercaillie and black grouse in relation to the small rodent

cycle in southeast Norway. Oecologia, 82(4). https://doi.org/10.1007/BF00319796

Weldon, P. J. (2021). Does human scent bias the survival of bird nests? In Ibis.

https://doi.org/10.1111/ibi.12966

Whitney, T. D., Sitvarin, M. I., Roualdes, E. A., Bonner, S. J., & Harwood, J. D. (2018). Selectivity

underlies the dissociation between seasonal prey availability and prey consumption in a generalist

predator. Molecular Ecology, 27(7). https://doi.org/10.1111/mec.14554

Williams, S. T., Maree, N., Taylor, P., Belmain, S. R., Keith, M., & Swanepoel, L. H. (2018). Predation by

small mammalian carnivores in rural agro-ecosystems: An undervalued ecosystem service? Ecosystem

Services, 30. https://doi.org/10.1016/j.ecoser.2017.12.006

Wood, S. N. (2017). Generalized additive models: An introduction with R, second edition. In

Generalized Additive Models: An Introduction with R, Second Edition. CRC Press.

https://doi.org/10.1201/9781315370279

Young, A., Márquez-Grant, N., Stillman, R., Smith, M. J., & Korstjens, A. H. (2015). An Investigation of

Red Fox (Vulpes vulpes) and Eurasian Badger (Meles meles) Scavenging, Scattering, and Removal of

Deer Remains: Forensic Implications and Applications. Journal of Forensic Sciences, 60(s1).

https://doi.org/10.1111/1556-4029.12554

Zalewska, K., Waggershauser, C. N., Kortland, K., & Lambin, X. (2021). The best defence is not being

there: avoidance of larger carnivores is not driven by risk intensity. Journal of Zoology.

https://doi.org/10.1111/jzo.12910

Zalewski, A., Jedrzejewski, W., & Jedrzejewska, B. (2004). Mobility and home range use by pine

martens (Martes martes) in a Polish primeval forest. Ecoscience, 11(1).

https://doi.org/10.1080/11956860.2004.11682815

Zalewski, A., & Jedrzejewski ,W. (2006). Spatial organisation and dynamics of the pine marten Martes

martes population in Białowieza Forest (E Poland) compared with other European woodlands. In

Ecography (Vol. 29, Issue 1). https://doi.org/10.1111/j.2005.0906-7590.04313.x

Zavaleta, E. S., Hobbs, R. J., & Mooney, H. A. (2001). Viewing invasive species removal in a whole-

ecosystem context. In Trends in Ecology and Evolution (Vol. 16, Issue 8).

https://doi.org/10.1016/S0169-5347(01)02194-2

<u>Appendix Document:</u>

<u>Appendix 1: Nest Camera Trapping Assumed and Confirmed:</u>

Comparison of assumed predators from signs with confirmed predators on camera traps showed a
high success rate in correctly identifying nest predators when an assumption was made. In many
cases where an observer was incorrect in assigning an assumed predator (pine marten only), the
pine marten was the primary nest predator. Still, the nest site was visited afterwards by a secondary
predator. When assigning an unknown mammal, in some instances, it was due to multiple predators
interacting with the nest site before the observer could look at the predation signs, changing the
indications, or, in the case of the fox, having so few instances of occurrence, signs were not clear.
Out of 115 nest sites predated by predators with a camera trap deployed on them, 106 were able to
detect the true nest predator successfully, and six failed to detect any interaction with the nest due
to camera trap failure. Overall, observers were correct over 99% of the time when assigning an
assumed predator. This gave us confidence to use assumed and confirmed predators in further fate
analysis.

<u>Table 1 Fates of nests of camera-trapped sites, with confirmed predation. The total number of</u>
<u>assumed assignments is compared with the number of confirmed predators and if the assignment</u>
<u>was correct. In the cases of incorrect assignment, the true predator was highlighted in "Incorrect</u>

classification". In the case of unknown mammals, the incorrect classification highlights who the truly
responsible predator was

Comparison of Assumed Accuracy (Nests with Camera Traps)					
Predator	Assumed Assignments (Total)	Number Confirmed	Correct Assignments	Accuracy (%)	Incorrect Classification
Pine Marten	72	72	70	97.2	1 Marten & Fox, 1 Marten & Badger
Badger	18	18	18	100	NA
Fox	0	2	0	NA	NA
Rodent	4	4	4	100	NA
Avian	0	0		NA	NA
Unknown_Mammal	13	NA	NA	NA	1 Badger, 8 Pine Marten, 2 Fox, 1 Dog & Pine Marten, 1 Badger & Pine Marten.
					N=109 (6 Camera traps failed to detect a predator)

Appendix 2: Feeding Station Access:

Feeding stations were monitored via camera traps across the entire sampling regime; camera images
were tagged using the methodology described in the main text across the 9 weeks of camera trap
monitoring of diversionary feeding. We highlight below the minimum number of feeding sites with
at least 1 detection of a feeding predator. The minimum is due to a large amount of food dispersal
from the central monitored point preventing perfect detection. Raptor species were grouped into
one category to include rare detections, such as White-tailed eagles, but mostly were detections of
buzzards. Domestic dogs were accidentally detected at 25% of feeding sites, with them not being
kept on leads as is advised during the breeding season. Further breakdown of key predators can be
found in the main text.

Uptake of diversionary feeding

Figure 2. Shows the proportion of feeding stations with confirmed feeding by predatory species via camera trapping. The y-axis shows the percentage of all 60 sites that had a detection of each of the species listed on the X-axis.

Appendix 3; Nest Fates:

Raw nest data was explored to evaluate the primary visual effect of treatment on nest fate. Before robust monitoring, simply from base values alone, there is a noticeable difference between the number of nest predation events alongside the treatment variable. Overall, regardless of treatment, the two fates with the highest occurrences survived and were depredated by pine marten. Forty-nine per cent of artificial nests (353/720) survived the full 28 days, meaning that 51% of nests experienced depredation. Recorded nest predators were martens (38%, 268), badgers (6.9%, 50), and other species (unknown, fox, corvid, and rodent (6.8%, 49)). However, fewer nests survived in control sites (128, 35.5%) than in diversionary feeding sites (228, 63.3%), reflecting fewer nests depredated by martens (Control: 170, Test: 98) and badgers (Control: 31, Test 19).

Nest Predators

N=720

Figure 2. Proportional nest loss by all nests (confirmed and assumed combined), by treatment and species. The figure is split by treatment type, with control (un-fed) and test sites (feeding). Sites are ordered in order of overall prevalence from bottom to top.

Appendix 4; Survival without Fate Kaplan Meir:

The daily status of each artificial nest over 28 days was coded as a string of 1 (intact) and 0 (depredated) using the known predation date for nests with cameras and the mid-point between visual checks for depredated nests without cameras. We estimated the influence of treatment by fitting the Kaplan-Meier survival probability curve using the 'survival' package (Therneau *et al.,* 2023). This allowed us to assess baseline fate (survived or predated) for all 720 nests over the 28-day sampling period. Large numbers of deaths occur on day 7 and day 21 due to the assumption of a central death date of nests between 14 and 28-day nest checks. These large dips indicate the trapping regime, not actual deaths. Links with broader multinomial analysis are made in the main text.

636 <u>Figure 3. Kaplan Meir survival curve visualising artificial nest survival over 28 days for control and</u>

637 <u>diversionary feeding (legend) diversionary feeding (N=720). Survival across our research period is</u>

638 <u>visualised using Kaplan Meier across 28 days of life for each nest; check days at 14 and 28 days are</u>

639 <u>shown with central death dates at days 7 and 21.</u>

Table 1. The coefficient table of multinomial logistic regression shows the changes to the log probability of the fate of an artificial nest. The base reference fate for model (A) is that a nest survived for 28 days. Significant factors are highlighted in bold.

Independent Variables	Dependant Variable (Fate)							
	Pine Marten (B)				Badger (C)			
	Coefficient	Std. Err.	z	P>|z|	Coefficient	Std. Err.	z	P>|z|
Intercept	0.278	0.235	1.184	0.237	**-1.794**	**0.409**	**-4.390**	**1.14e-05**
Treatment	**-1.494**	**0.309**	**-4.832**	**1.35E-06**	**-1.723**	**0.622**	**-2.771**	**0.006**
Distance 300m	0.005	0.285	0.019	0.984	-0.547	0.533	-1.027	0.304
Distance 500m	0.206	0.297	0.691	0.489	0.319	0.473	0.675	0.499
Year (2022)	-0.007	0.172	-0.038	0.969	**0.694**	**0.325**	**2.137**	**0.033**
Treatment*500m	0.155	0.425	0.366	0.715	1.170	0.839	1.394	0.163
Treatment*300m	0.698	0.424	1.647	0.099	0.736	0.795	0.926	0.355

Fate A (Survived) Calculated as Difference in Interecept (100m, Control, 100*Control, 2021)
Significant Results in Bold

641