# Glial ferritin maintains neural stem cells via transporting iron required for self-renewal in *Drosophila*

**Zhixin Ma,[1] Wenshu Wang,[1] Xiaojing Yang,[1] Menglong Rui,[1] Su Wang [1,2,3,*]**

[1]School of Life Science and Technology, The Key Laboratory of Developmental Genes and Human Disease,

Ministry of Education, Southeast University, 2 Sipailou Road, Nanjing 210096, China

[2]Co-innovation Center of Neuroregeneration, Nantong University, Nantong, China

[3]Lead Contact

[*]Correspondence: wangsu@seu.edu.cn

**Abstract**

Stem cell niche is critical for regulating the behavior of stem cells. *Drosophila* neural stem cells (Neuroblasts, NBs) are encased by glial niche cells closely, but it still remains unclear whether glial niche cells can regulate the self-renewal and differentiation of NBs. Here we show that ferritin produced by glia, cooperates with Zip13 to transport iron into NBs for the energy production, which is essential to the self-renewal and proliferation of NBs. The knockdown of glial ferritin encoding genes causes energy shortage in NBs, which leads to the low proliferation and premature differentiation of NBs. Moreover, the level of glial ferritin production is affected by the status of NBs, establishing a bicellular iron homeostasis. In this study, we demonstrate that glial cells are indispensable to maintain the self-renewal of NBs, unveiling a novel role of the NB glial niche during brain development.

**Keywords:** glia, ferritin, neural stem cells, iron homeostasis, self-renewal

## Introduction

Neural stem cells (NSCs) possess the remarkable ability to self-renew for maintaining their identity and differentiate into diverse neural cell types required for neurogenesis during brain development. The self-renewal, differentiation, and proliferation of NSCs are regulated by both intrinsic programs inside stem cells and extrinsic cues from the surrounding niches. The mammalian NSC niche found in the neurogenic regions is composed of diverse cell types including choroid plexus, vascular endothelia cells, pericytes, microglia, and meninges, which regulate NSC behaviors such as proliferation, self-renewal, and differentiation (Bjornsson, Apostolopoulou, Tian, & Temple, 2015). The diversity and complexity of these cell populations confer niche the ability of integrating local and systemic signals, but make it complicated to study the interaction between niches and NSCs in mammals, which is a critical issue to understand neurogenesis and improve clinical applications of NSCs.

*Drosophila* larval central nervous system (CNS), composed of central brain (CB), optic lobe (OL) and ventral nerve cord (VNC), is a simple but sophisticated model to decipher the regulatory mechanisms underlying NSC behaviors mediated by niche signals. In this system, *Drosophila* NSCs, called neuroblasts (NBs), and their progenies are encased by a variety of glial cells including perineurial glia, subperineurial glia, cortex glia, astrocyte-like glia and ensheathing glia (Hartenstein, 2011; Ito, Urban, & Technau, 1995; Pereanu, Shy, & Hartenstein, 2005). These glial cells not only

form the physical support structure for NBs, but also provide extrinsic cues that regulate NB reactivation, survival, and proliferation (Read, 2018; Speder & Brand, 2014). However, it remains elusive whether the glial-cell-forming niche is involved in the regulation of the prominent characteristics of NBs, self-renewal and differentiation.

Iron is a vital micronutrient for almost all living organisms since it participates in many crucial biological processes, including mitochondrial electron transport and cellular respiration that both support ATP production used for most cellular activities. High levels of iron are found in many organs including the brain, where iron is required for myelination and formation of neuronal dendritic trees (Lozoff et al., 2006; Moller et al., 2019; Rice & Barone, 2000). Iron deficiency in infants and early childhood has been well documented to be associated with long-term cognitive, social, and emotional problems (Fretham, Carlson, & Georgieff, 2011). Thus, it is intriguing to investigate whether iron is also required in NSCs during neurogenesis.

Ferritin, a ubiquitous protein present in most organisms, is composed of 24 subunits that assemble into a hollow-sphere complex capable of storing up to 4,500 iron atoms in a bioavailable form (Andrews, 2005; Knovich, Storey, Coffman, Torti, & Torti, 2009). Ferritin in both mammal and insect is a heteropolymer, consisting of heavy and light chain homologue subunits. In *Drosophila*, these subunits are encoded by *ferritin 1 heavy chain homologue* (*Fer1HCH*) and *ferritin 2 light chain homologue* (*Fer2LCH*) respectively (Arosio, Ingrassia, & Cavadini, 2009; Pham & Winzerling, 2010), which both contain secretion signal peptides that direct them to the ER during translation (Lind, Ekengren, Melefors, & Soderhall, 1998). Consequently, *Drosophila* ferritin is confined to the secretory pathway, abundant in the hemolymph, and absent from the cytosol (Nichol, Law, & Winzerling, 2002). Previous researches have shown that *Drosophila* ferritin not only directly participates in dietary iron absorption by removing iron from enterocytes across the basolateral membrane, but competes with transferrin to deliver iron between the gut and the fat body (Tang & Zhou, 2013; Xiao, Liu, Zhao, Wang, & Zhou, 2019). These findings underscore the pivotal role of ferritin in iron transport. Interestingly, iron-loaded ferritin, serving as an essential mitogen, promotes proliferation of cultured *Drosophila* cells (Li, 2010). Therefore, it is tempting to investigate whether ferritin is required for self-renewal and proliferation of NSCs during brain development.

Here we show that glial ferritin supplies iron into NBs to form iron-sulfur (Fe-S) clusters, which is necessary for the production of ATP, providing the energy required for self-renewal and proliferation of NBs, highlighting ferritin as a potential target for suppressing tumor.

**Results**

**Glial ferritin is required for NB maintenance and proliferation**

To identify new niche factors involved in the balance between NB self-renewal and differentiation, we used *repo-GAL4* to perform an RNAi mediated screen in glial cells. *Fer1HCH* and *Fer2LCH*, two ferritin subunit encoding genes, were identified to be essential for NB maintenance and brain development. Specifically, knocking down *Fer1HCH* or *Fer2LCH* in glial cells significantly reduced the number of NBs in the CB and VNC at the third-instar larval stage, which resulted in decreased size of the CNS (**Figure 1A and B and Figure 1-figure supplement 1A**). To further assess the impact on NB proliferation, we employed the mitotic marker phosphohistone H3 (PH3) and the thymidine analogue 5-ethynyl-2'-deoxyuridine (EdU), revealing a substantial decrease in the proliferation rate of NBs upon glial ferritin knockdown (**Figure 1A and C and Figure 1-figure**

**supplement 1B and C**). These results indicated that glial ferritin is required for maintaining NBs
and their proliferation.
To identify the glial subpopulation in which ferritin functions, we employed subtype-restricted
GAL4 to perform ferritin knockdowns (**Table 1**). We observed that knocking down ferritin genes in
cortex glia led to a similar yet milder phenotype compared to that observed in the pan-glia
knockdown (**Figure 1-figure supplement 1D**). This milder effect may be attributed to potential
compensatory mechanisms from other glial subpopulations. Interestingly, knocking down ferritin in
other glial subtypes did not yield any discernible phenotype. These findings suggest that ferritin
functions collectively across all glial populations to regulate the activity of NBs.
We confirmed the phenotype with multiple independent RNAi lines for *Fer1HCH* or *Fer2LCH*,
and observed similar abnormalities of larval NBs and brain development (**Figure 1-figure
supplement 2A**). Hereafter, *UAS-Fer1HCH-RNAi* (THU5585) and *UAS-Fer2LCH-RNAi*
(TH01861.N) were used for most studies, and we only quantified NB number and proliferation rate
in the CB for convenience in this research.
To evaluate the efficiency of RNAi for *Fer1HCH* or *Fer2LCH*, RT-PCR was performed with total
RNA isolated from whole brain dissected from third-instar larvae. The abundance of *Fer1HCH* or
*Fer2LCH* mRNA was significantly reduced in glial *Fer1HCH* or *Fer2LCH* knockdown respectively
(**Figure 1D and E**). Western blot analysis of larval brain protein extract using polyclonal antibodies
against Fer1HCH or Fer2LCH further confirmed that both subunits were down-regulated in glial
knockdown of either subunit (**Figure 1F-I**), consistent with previous research in *Drosophila*
(Missirlis et al., 2007). To exclude the possibility of RNAi off-target effects, we used *repo-GAL4* to
overexpress RNAi-resistant *Fer1HCH* (or *Fer2LCH*) containing silent mutations in the region
targeted by shRNA, which resulted in a significant increase in the number and proliferation rate of
NBs compared to glial ferritin knockdown alone (**Figure 1J-L and Figure 1-figure supplement
2B, C**). Additionally, similar but weaker phenotypes were observed when Fer1HCH$^{DN}$, a
ferroxidase-inactive form of Fer1HCH (Missirlis et al., 2007), was overexpressed in glial cells
(**Figure 1-figure supplement 2D**). These results support the conclusion that ferritin in glial cells is
essential for the maintenance and proliferation of NBs, suggesting that iron homeostasis is involved
in this process.
116
**Ferritin is produced mainly in glial cells and secreted into NBs through a vesicle-dependent
pathway**
To investigate the role of iron homeostasis in *Drosophila* CNS, we tried to knock down genes
involved in iron transport, storage, and regulation in NBs, neurons, and glial cells, respectively.
Interestingly, only glial knockdown of ferritin-related genes led to NB loss and defective brain
development. It appears that ferritin generated in glial cells is the only crucial protein for regulating
the behavior of NBs and, ultimately, the larval brain development. To recapitulate the endogenous
ferritin expression pattern, we took advantage of a Fer1HCH/Fer2LCH enhancer trapped GAL4 line
to drive a variant of DsRed protein with a nuclear localization signal (DsRed Stinger). Surprisingly,
our immunostaining experiments revealed that most, if not all, DsRed was co-localized with the glia
nuclear protein Repo, instead of the NB nuclear protein Dpn (**Figure 2A**). These results further
confirm that glial cells are the main producers of ferritin in the larval brain.
*Fer1HCH$^{G188}$* encodes a GFP-tagged Fer1HCH subunit (Missirlis et al., 2007), which labels the
distribution of Fer1HCH, representing the pattern of ferritin. As shown in the **Figure 2B**, ferritin

was detected in NBs besides glial cells, despite that ferritin is specifically produced in glial cells.
To explore the possibility of ferritin transfer from glial cells to NBs, we observed the GFP signal of
*Fer1HCH^{G188}* with glial ferritin knockdown, and found that GFP in the brain was almost completely
depleted (**Figure 2B**). Furthermore, we used *repo-GAL4* to drive the expression of GFP-tagged
Fer1HCH and mCherry-tagged Fer2LCH in glial cells. We observed that both Fer1HCH::GFP and
Fer2LCH::mCherry were present in NBs, while the control GFP and mCherry were restricted to
glial cells (**Figure 2C**). Altogether, we conclude that ferritin is generated in glial cells and then
transferred to NBs in the larval brain.

In *Drosophila*, both ferritin subunits contain secretion signal peptides that direct protein to the
endoplasmic reticulum during translation, which allows the ferritin to be transported out of the cell.
To investigate whether the ferritin was transported from glia to NBs via the vesicular transport, we
expressed a temperature-sensitive dominant-negative Shibire dynamin protein (shi^{ts}) to block
vesicle trafficking from glial cells, and found that Fer1HCH::GFP accumulated in glia,
accompanying with significantly diminished Fer1HCH signal in NBs (**Figure 2D and E**),
suggesting glial ferritin secretes into NBs through vesicle system.
146
**Glial ferritin defects lead to iron deficiency in NBs**
Given that ferritin is required for iron storage as well as delivery, glial ferritin knockdown might
increase free iron level, leading to glial iron overload. Alternatively, it will block the iron transport
from glia to NBs, resulting in NB iron deficiency. To determine the underlying cause of the brain
development defects observed upon glial ferritin knockdown, we used iron chelator or ferric
ammonium citrate (FAC) to manipulate iron intake and detect how it will influence NBs. We
observed that depriving iron intake by adding iron chelator bathophenanthrolinedisulfonic acid
disodium (BPS) to the food led to a dramatic decrease in the number and proliferation rate of NBs,
compared to control larvae on normal food (**Figure 3A**). In line with this result, blocking iron
absorption by knocking down intestinal ferritin (Tang & Zhou, 2013) resulted in the similar defects
of brain (**Figure 3-figure supplement 1A-C**). However, supplementing the diet with FAC did not
induce any observable defects (**Figure 3A**). We also tested another iron chelator deferiprone (DFP),
and found that DFP alone did not cause any brain defects in wild-type (WT) larvae, but it
significantly exacerbated the effect of glial ferritin knockdown on defective NB maintenance
(**Figure 3B and C**). Furthermore, we attempted to rescue the brain defects observed in glial ferritin
knockdown by providing dietary FAC, but the effect was negligible (**Figure 3-figure supplement
1D and E**). This may be due to the possibility that the knockdown of glial ferritin largely, if not
completely, prevented the transport of iron from glia to NBs, which hindered the additional iron
from entering NBs. These results indicate that the NB loss phenotype induced by glial ferritin
knockdown is primarily due to iron deficiency in NBs.
Zip13, generally known as a zinc transporter, functions as an iron transporter, cooperating with
ferritin on iron transport in *Drosophila* intestine (Xiao, Wan, Fan, Tang, & Zhou, 2014). Specifically,
knockdown of *Zip13* leads to failure of iron loading in secretory vesicles and consequently defective
iron efflux (Xiao et al., 2014). To investigate whether the NB loss phenotype of glial ferritin
knockdown is caused by defective iron transport, we next sought to identify the role of Zip13 in the
larval brain. Firstly, we examined the expression pattern of Zip13 and found that, similar to ferritin,
the DsRed stinger driven by *Zip13-GAL4* was co-localized with Repo in glial cells rather than Dpn
in NBs (**Figure 3D**). Subsequently, we knocked down *Zip13* in glial cells and observed a similar

but weaker phenotype with the slightly reduced NB number and proliferation rate (**Figure 3E-G**). Consistently, the defective brain development was confirmed in *Zip13* mutant (**Figure 3-figure supplement 1F-H**). Moreover, the effect of double knockdown of both *Zip13* and *Fer2LCH* was much more severe than that of either single knockdown, with a significant reduction in NB number and proliferation rate (**Figure 3E-G**). Furthermore, we found that the reduced proliferation rate caused by *Zip13* knockdown could be partially rescued by iron supplementation in the food (**Figure 3H and I**). Overall, these results suggest that glial ferritin collaborates with Zip13 to deliver iron from glia to NBs.

Since previous studies have demonstrated the essential role of glial cells in NB survival and proliferation (Bailey et al., 2015; Read, 2018), it is possible that iron overload in glial cells up-regulated the level of reactive oxygen species (ROS), leading to apoptosis of glial cells, which ultimately resulted in the loss of NBs. To rule out this possibility, firstly, we examined the ROS level of glial ferritin knockdown via a transgenic reporter GstD-GFP for oxidative stress signaling (Sykiotis & Bohmann, 2008), and found an obvious increase as expected (**Figure 3-figure supplement 2A and B**). However, overexpressing anti-oxidant genes in glial cells or NBs did not restore the NB number and proliferation (**Figure 3-figure supplement 2C and D**). ROS accumulation leads to lipid peroxidation, which in turn results in ferroptosis (Hirschhorn & Stockwell, 2019). Mitochondria in cells undergoing ferroptosis typically exhibit a characteristic shrinkage and damage, while the nuclei remain structurally intact (Mumbauer, Pascual, Kolotuev, & Hamaratoglu, 2019). In order to investigate whether the NB loss is due to ferroptosis, we examined the mitochondrial morphology using Transmission Electron Microscopy (TEM). However, our observation revealed no discernible defects in the mitochondria of NBs after glial ferritin knockdown (**Figure 3-figure supplement 2E and F**). Taken with the findings from iron chelator feeding together, we conclude that NB loss resulting from glial ferritin knockdown is not a consequence of ferroptosis. Secondly, we examined the apoptosis in glial cells via Caspase-3 or TUNEL staining, and found the apoptotic signal remained unchanged after glial ferritin knockdown (**Figure 3-figure supplement 3A-D**). Furthermore, we overexpressed the baculovirus protein P35 to inhibit apoptosis in glial cells, but failed to restore the lost NBs (**Figure 3-figure supplement 3E**). Given that glial morphology is important for NB lineages (Speder & Brand, 2018), we used GFP driven by *repo-GAL4* to denote glia cells and found that glial chamber was almost intact after glial ferritin knockdown (**Figure 3-figure supplement 3F**). Together, we conclude that it is iron deficiency in NBs, rather than iron overload in glial cells, that leads to NB loss induced by ferritin defects.

**Glial ferritin defects result in impaired Fe-S cluster activity and ATP production**

To further prove that defective iron transport leads to the iron deficiency in NBs, we examined the iron availability by measuring aconitase activity in the brain. The activity of aconitase is sensitive to the level of Fe-S synthesis, so it can serve as an indicator for the availability of iron in the cell (Haile et al., 1992; Tong & Rouault, 2006). As shown in the **Figure 4A**, aconitase activity of ferritin knockdown diminished significantly, indicating that the availability of iron in the brain was reduced. This result also implies Fe-S clusters in NBs are probably deficient when knocking down ferritin in glia. Fe-S clusters are essential to activate the aconitase that converts citrate into isocitrate in the tricarboxylic acid cycle（TCA cycle) in mitochondria, and are one type of critical electron carrier for electron transport chain (ETC), especially for the Complex I, II and III function. Considering

that TCA cycle and ETC are critical biological processes for ATP production, we proposed that knockdown of glial ferritin might lead to ATP deficiency due to the block of TCA cycle and ETC, which resulted in the failure of maintaining NBs and their proliferation. To verify this hypothesis, we first assessed the function of Fe-S clusters in NBs by knocking down the components of the cytosolic iron-sulfur protein assembly (CIA) complex. After knocking down the CIA component *CG17683*, *galla2*, and *ciao1* in NBs respectively, the proliferation level of NBs decreased significantly and brain size became smaller (**Figure 4B and C**), which phenocopied the knockdown of *Fer1HCH* or *Fer2LCH* in glia. This result implies that Fe-S clusters are indispensable for NBs to maintain their proliferation.

To identify the biological processes related to Fe-S clusters in NB lineages, we performed transcriptome analysis of sorted NB lineages from glial ferritin knockdown as well as the control groups. We used NB driver *Insc-GAL4* to express *UAS-DsRed*, labeling NB lineages, and then sorted these cells via fluorescence-activated cell sorting (FACS) (**Figure 4D**) (Harzer, Berger, Conder, Schmauss, & Knoblich, 2013). We found that thousands of genes were significantly altered in the *Fer1HCH* and *Fer2LCH* knockdown groups compared to the control group. Specifically, 880 genes were simultaneously downregulated among the significantly changed genes in both the *Fer1HCH* and *Fer2LCH* knockdown groups contrasted to the control group (**Figure 4E**). Given that iron deficiency in NBs probably leads to the defects of energy production and biosynthesis, we performed gene enrichment analysis in these genes based on Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG). GO term analysis showed that mitochondrial organization, translation, transport, oxidative phosphorylation (OxPhos), and ETC were enriched in the Biological Processes (**Figure 4F**), indicating NB mitochondrial functions including OxPhos and ETC were abnormal after glial ferritin knockdown. Moreover, NADH dehydrogenase and ribosome related terms were also enriched whether in the Cellular Component or Molecular Function (**Figure 4F**), suggesting that NADH dehydrogenase and biosynthesis were defective. Further, we validated the enriched term associated with mitochondrial function by qRT-PCR (**Figure 4-figure supplement 1A**). And the enriched KEGG pathways included OxPhos and ribosome (**Figure 4G**), aligning with the GO enrichment results and supporting the hypothesis that glial ferritin knockdown disrupted mitochondrial function including OxPhos and ETC as well as further biosynthesis.

To confirm whether the function of NADH dehydrogenase was disrupted, we used the SoNar that could bind to $NAD^+$ with specific conformation and excitatory wavelength to monitor the $NAD^+$ level (Bonnay et al., 2020). SoNar was driven by the *Insc-GAL4* to indicate the $NAD^+$ level in NBs. Considering the protein accumulation in NBs due to the lower proliferation rate in glial ferritin knockdown group, the signal of SoNar was normalized to DsRed Stinger driven by *Insc-GAL4*. We observed significantly declined SoNar/DsRed after knocking down *Fer2LCH* (**Figure 4H and I**), suggesting the normalized level of $NAD^+$ in NBs was decreased, which supports the conclusion that the activity of NADH dehydrogenase was declined. To further verify the change of energy production, we utilized the ATP Determination Kit which offers a convenient bioluminescence assay for quantitative determination of ATP with recombinant firefly luciferase and its substrate D-luciferin. We found that ATP level decreased significantly after glial ferritin knockdown when compared to the control (**Figure 4J**). Therefore, we tried to supply the energy for NBs by restoring bioenergy-dependent $NAD^+$ metabolism. As expected, the NB number could be partially rescued by overexpressing proton-pump-independent yeast mitochondrial NADH dehydrogenase Ndi1 in NBs under the background of glial ferritin knockdown (**Figure 4K and L**) (Bonnay et al., 2020). In

summary, the iron deficiency leads to NB loss due to the insufficient energy required for essential biological processes.

**Glial ferritin maintains NBs via preventing premature differentiation**

To investigate how glial ferritin knockdown leads to NB loss, we firstly examined whether NB origin was affected via altering the timing of ferritin knockdown. We used the temperature-sensitive GAL4 inhibitor GAL80$^{ts}$ to suppress ferritin RNAi at a permissive temperature (18°C) during embryonic stage, and then shifted to a restrictive temperature (29°C), or vice versa until we dissected them at the third-instar larval stage. Ferritin knockdown during the larval stage led to severe brain development defects (**Figure 5A and B**), while the knockdown only during the embryonic stage showed no apparent abnormality (**Figure 5-figure supplement 1A and B**). Together, these results indicate that glial ferritin is required for NB maintenance at the larval stage. Consistent with this result, we found no obvious defects of brain at the first-instar larval stage (0-4 hours after larval hatching) when knocking down glial ferritin (**Figure 5-figure supplement 1C**). Since newborn NBs are delaminated from the neuroectoderm during the embryonic stage (stage 9 to 11), and ferritin RNAi only during the embryonic stage did not show any phenotype, suggesting that NB loss after glial ferritin knockdown is not ascribed to defective NB origin.

As mentioned before, we did not detect any apoptotic signal in the larval brain after glial ferritin knockdown, suggesting that there is no association between apoptosis and NB loss. The double binary systems, lexA/lexAop and GAL4/UAS, were employed to further exclude the possibility of apoptosis. We overexpressed P35 in NBs by *Insc-GAL4* and knocked down ferritin by *repo-lexA*, which failed to restore the decreased number of NBs (**Figure 5-figure supplement 1D**). These results reinforce the notion that NB loss occurs independent of the apoptotic pathway.

Next, we hypothesized that the NB loss might be attributed to nuclear-Prospero (Pros)-dependent premature differentiation of NBs (Cabernard & Doe, 2009; Choksi et al., 2006). Pros in NB nuclear was hardly detected in third-instar WT larvae, but was increased significantly with glial ferritin knockdown (**Figure 5C and D**). To further verify that the NB loss caused by glial ferritin knockdown was due to nuclear-Pros-dependent premature differentiation, we used the *Insc-GAL4* to drive *UAS-pros-RNAi* in order to prevent the NB loss in glial ferritin knockdown background. We found that the NB number was substantially restored (**Figure 5E and F**). Interestingly, despite of dramatic NB number increase, the decreased proliferation rate was not rescued. We proposed that *pros* knockdown in NBs blocked premature terminal differentiation, but it could not replenish the reduced energy caused by glial ferritin knockdown. As NB self-renewal and proliferation are both energy-consuming processes, the self-renewal was rescued via inhibiting differentiation by *Insc-GAL4* driving *UAS-pros-RNAi*, but at the expense of energy consumption for proliferation, leaving the proliferation rate unchanged, or even lower. Together, we conclude that glial ferritin maintains the number of NBs by preventing the nuclear-Pros-dependent premature differentiation.

Given that reduced proliferation is a distinctive characteristic of quiescent NBs, we tried to determine whether NBs were in a quiescent state after glial ferritin knockdown by examining the other two key features associated with quiescent NBs: decreased cell size and the formation of the extended cellular protrusions (Chell & Brand, 2010; Ly et al., 2019). However, we found that the NB size remained at the same level as control, and the extended cellular protrusion was not observed after glial ferritin knockdown (**Figure 2B and Figure 5-figure supplement 2A**). Furthermore, since insulin signaling is necessary and sufficient to reactivate NBs from quiescence (Sousa-Nunes, Yee,

307 & Gould, 2011), we tried to overexpress a constitutively active form of InR (InR$^{act}$) in NBs to
308 promote reactivation at the background of glial ferritin knockdown. However, InR$^{act}$ overexpression
309 failed to ameliorate the NB abnormalities including the decreased proliferation (**Figure 5-figure**
310 **supplement 2B and C**). In conclusion, NBs exhibited a low proliferation rate after glial ferritin
311 knockdown, but were not in a quiescent state.
312
**Ferritin functions as a potential target for tumor suppression**
We have demonstrated that glial ferritin is essential for NB self-renewal and proliferation by
providing glia-to-NB iron transport to support energy production in NBs. We next tried to determine
whether manipulating this biological function could suppress tumor development. We induced
tumor development by driving *UAS-brat-RNAi* or *UAS-numb-RNAi* under the control of *Insc-GAL4*,
and knocked down ferritin genes under the control of *repo-lexA*. We observed that the size of tumors
was strikingly decreased in the larval brains lacking glial ferritin (**Figure 6A and B**). This result
suggests that glial ferritin is required for tumor development, highlighting a potential target for
tumor treatment.
To verify whether iron is involved in the tumor suppression induced by blocking glial ferritin, we
tried to decrease iron level by adding the iron chelator BPS in the *Drosophila* food. Tumor formation
in the brain at the third instar larval stage was induced with the same method as mentioned above.
The addition of BPS into dietary food significantly reduced the tumor size (**Figure 6C and D**),
suggesting that iron is required for tumor development in *Drosophila* larval brain. Considering the
conservation of iron's fundamental function as an essential nutrient for cell growth and proliferation,
we thus tried to suppress brain tumor in mice via iron deficiency. Initially, we constructed orthotopic
glioma model in C57BL/6J mice. One week after inoculation of glioma cells (GL261-luc), the iron
chelator DFP was intraperitoneally injected every two days based on the murine weight until being
dissected on day 7, 14, and 21, respectively. Histological examination with HE staining revealed
that the size of glioma in mice, injected with DFP, decreased significantly on day 14 and 21 (**Figure**
**6-figure supplement 1A-C**). Since glioma cells GL261-luc carry the luciferase gene, we injected
the substrate luciferin potassium salt into the tumor-bearing mice and detected the bioluminescence
signal of mouse gliomas using IVIS Spectrum. This result showed that the bioluminescence signal
of glioma in mice with the injection of 10 mg/ml DFP was significantly reduced (**Figure 6E and**
**F**). The above results suggested that iron deficiency induced by DFP could suppress glioma in mice
effectively. Furthermore, we assessed the survival time of the tumor-bearing mice with DFP or
saline injection and found that 10 mg/ml DFP injection could significantly prolong the survival time
of tumor-bearing mice (**Figure 6G**). Altogether, blocking iron into tumor is a potential therapy for
tumor suppression.
342
**Glial ferritin level is regulated by NBs**
The aforementioned results have highlighted the significant role of glial ferritin in controlling NB
number and proliferation. We next wonder whether NBs could provide feedback to regulate glial
ferritin level. To answer this question, we analyzed ferritin level after altering NB proliferation.
Previous studies have corroborated that NB proliferation was diminished by blocking Tor signaling
pathway or enhanced by knocking down tumor suppressor genes, such as *brat* or *numb* (Bowman
et al., 2008; Reichardt et al., 2018; Sousa-Nunes et al., 2011). When NBs were in low proliferative
status, Fer2LCH subunits in the brain were significantly decreased, while Fer1HCH subunits

declined slightly with no significance (**Figure 7A-D**). Conversely, highly proliferative NBs, able to
form tumors, induced significantly increased level of the ferritin (**Figure 7A-D**). In summary, the
production of ferritin in glial cells depends on the level of NB proliferation.
354

**Discussion**

Here we demonstrate that glia regulates iron metabolism in the brain to sustain NBs and their
proliferation. Glial ferritin coordinates Zip13 to provide iron for NBs. Once ferritin in glia is
defective, NBs proliferate much slower and differentiate prematurely, implying that iron is a
necessary metallic element for NBs. Iron participates in the synthesis of Fe-S clusters which activate
aconitase required for TCA and facilitate electron-transfer processes involved in OxPhos. Since
TCA and OxPhos are essential steps of ATP production, glial ferritin supplies iron into NBs to
ensure energy production for maintaining NBs and their proliferation (**Figure 7E**).
363

**NSCs and glial niche**

Previous literatures have reported how glial niche regulates NB reactivation, proliferation, and
survival. Blood-brain barrier (BBB) glia secretes dILP2 and dILP6 that bind on InR of NBs, which
switches on Insulin signaling to reactivate NBs (Chell & Brand, 2010; Speder & Brand, 2014). In
addition, Pvr (PDGFR ortholog) signaling in cortex glia sustains NB survival and proliferation in a
PI3K-DE-cadherin-dependent manner (Read, 2018). However, it is unknown whether glial niche is
required for maintaining the self-renewal of NBs. In this study, we demonstrate that glial niche
maintains the self-renewal and proliferation of NBs through supplying the necessary microelement
iron into NBs for enough energy production, which implies that mammalian glia likewise provides
critical microelement for maintaining NSC function. Oligodendrocytes have been reported to secret
ferritin to protect neuron (Mukherjee et al., 2020), however, it remains unknown whether the ferritin
secretion is employed by oligodendrocytes to regulate NSCs. Astrocytes regulate NSC proliferation
via secreting growth factors (Shetty, Hattiangady, & Shetty, 2005), but it is unclear whether ferritin
from astrocytes regulates NSC proliferation and other NSC behaviors, such as self-renewal and
differentiation. Based on our findings, it is feasible to regulate the self-renewal and differentiation
of NSCs indirectly through niche cells, which would help to reduce the risk of directly manipulating
NSCs to treat neural diseases.
381

**Ferritin and iron**

Our results show that ferritin generated outside the brain was not detected in the CNS and vice versa
(data not shown), suggesting ferritin can hardly cross the BBB glia, possibly because ferritin is too
large to permeate. In this way, BBB glia separates the labile systemic iron from local iron inside the
brain to guarantee the normal brain development, as the neural cells, including NBs, require a stable
iron level for continuous energy generation during neurogenesis.
Although the dominant function of ferritin is iron storage in mammal, increasing evidence
suggests that ferritin also can transport iron. In mice, ferritin heavy chain is secreted by
oligodendrocytes via extracellular vesicles (Mukherjee et al., 2020), which possibly endows ferritin
with the function of iron transport. Consistently, previous results showed that ferritin can deliver
iron across the cultured BBB (Chiou et al., 2019; Fisher et al., 2007). Our results emphasize the
critical role of ferritin in iron transport in the CNS. Future studies would be interesting to investigate
whether ferritin is able to transport iron in the mammalian brain.

Ferritin specifically targets on the tumor cells, which has been applied to tumor visualization and targeted drug delivery (Fan et al., 2012; Fan et al., 2018). Our study shows that the block of ferritin or iron can inhibit tumor effectively, which offers an explanation that ferritin targets on tumor is to transport iron for maintaining the cell proliferation, highlighting a potential target for tumor suppression.

Iron is an essential component for Fe-S clusters which participate in TCA and OxPhos, providing energy required for self-renewal and proliferation of NBs. Previous study claimed that OxPhos is required for NB proliferation (van den Ameele & Brand, 2019). However, another research observed that OxPhos is dispensable in type II NBs (Bonnay et al., 2020). Our data imply that energy deficiency in NBs caused by iron shortage leads to NB loss and low proliferation possibly through simultaneous disruption of TCA and OxPhos. Consistently, iron-loaded ferritin nanoparticles can be used to improve the self-renewal ability and differentiation potential of human NSCs in vitro (Lee, Yang, Cho, & Cho, 2018). Additionally, previous study has shown that iron-loaded ferritin can promote the proliferation of cultured cell from *Drosophila* (Li, 2010). Furthermore, iron depletion was reported to suppress human cancer cell growth at G1/S by inducing cyclin D1 proteolysis (Nurtjahja-Tjendraputra, Fu, Phang, & Richardson, 2007). These studies suggest that iron, as a necessary microelement for energy production, is essential to NSC maintenance and proliferation.

**Bicellular iron homeostasis**

Iron homeostasis can be regulated systemically by the liver-derived hormone hepcidin and the iron exporter ferroportin, and it is also regulated cellularly by the IRP/IRE system (Hentze, Muckenthaler, Galy, & Camaschella, 2010). Here we propose a bicellular iron homeostasis between NB and glial niche cell, which is distinct from classic cellular and systemic iron homeostasis. The bicellular iron homeostasis requires an elaborate mechanism to guarantee normal neurogenesis.

In a proliferative state, NBs receive iron from glial cells via ferritin to fulfill the energy requirements for proliferation and self-renewal. Therefore, the iron required by NBs is mainly stored in the ferritin of glial cells and allocated as needed. The bicellular iron homeostasis mechanism makes glial cells the main pool for ferritin production, relieving NBs from the need to produce ferritin and transferring the potential risk of iron overload-induced oxidative stress to glial cells. Furthermore, glial cells can subsequently clear oxidative stress through lipid droplets (Bailey et al., 2015), ensuring normal NB function during neural development. It is intriguing to explore whether bicellular iron homeostasis exists in other stem cells and their niches.

When normal NBs are converted into immortal tumor NBs, becoming highly proliferative, they need more nutrients including iron (Bonnay et al., 2020; van den Ameele & Brand, 2019). In response, more ferritin is generated to transport iron from the glial cells to NBs. However, it is unclear how NBs communicate with glial cells to determine the level of ferritin production. It's interesting to identify the specific cues from NBs that regulate the level of ferritin in future studies. In addition, iron distribution in *Drosophila* brain is not observed due to low iron concentration and technical limitations in iron detection, which results in a lack of direct evidence for iron trafficking via ferritin.

**Figure legends**

**Figure 1. Ferritin in glia is required for NB maintenance and proliferation**

(A) NB number and proliferation rate after *Fer1HCH* or *Fer2LCH* RNAi in glia.

(B) Quantification of NB number in CB (A).

(C) Quantification of NB proliferation rate in CB (A).

(D and E) RT-PCR analysis of Fer1HCH and Fer2LCH mRNA levels (D) and normalized quantification (E) after glial ferritin knockdown.

(Fand H) Western blot of Fer1HCH (F) and Fer2LCH (H) after ferritin knockdown in glia. Tubulin was used as a loading control.

(G and I) Quantification of normalized Fer1HCH intensity in (G) and Fer2LCH intensity in (I).

(J) Rescue of NB number and proliferation when simultaneously overexpressing RNAi-resistant *Fer1HCH* and *Fer1HCH RNAi* (or *Fer2LCH* and *Fer2LCH RNAi*) in glia.

(K and L) Quantification of NB number and proliferation rate in (J).

(B, n=11; C, n=10; E, G and I, n=3; K, n=8,8,9,7,9; L, n=9,8,8,7,12; ns, not significant; *: $p<0.05$; **: $p<0.01$; ***: $p<0.001$; ****: $p<0.0001$; Dpn, NB nuclei, red; PH3, green; DAPI, nuclei, blue).

**Figure 1-figure supplement 1. Glial ferritin knockdown leads to low NB proliferation and number**

(A) Quantification of NB number in VNC.

(B) Quantification of NB proliferation rate in VNC.

(C) EdU incorporation in CNS after knocking down glial ferritin.

(D) Knocking down *Fer2LCH* by *nrv2-GAL4* led to similar but weaker phenotype in brain in comparison with pan-glial ferritin knockdown.

(A, n=11; B, n=10; ****: $p<0.0001$)

**Figure 1-figure supplement 2. Verify the phenotype induced by glial ferritin knockdown using different manipulations**

(A) NB number and proliferation in other RNAi lines targeting on ferritin.

(B) Rescue of NB number in VNC when simultaneously overexpressed RNAi-resistant Fer1HCH and Fer1HCH RNAi (or Fer2LCH and Fer2LCH RNAi) in glia.

(C) Rescue of NB proliferation rate in VNC when simultaneously overexpressed RNAi-resistant *Fer1HCH* and *Fer1HCH RNAi* (or *Fer2LCH* and *Fer2LCH RNAi*) in glia.

(D) NB number and proliferation in *Fer1HCH^{DN}*.

(B, n=8,7,9,7,8; C, n=9,8,8,7,12; ****: $p<0.0001$)

**Figure 2. Ferritin is produced mainly in glial cells in the *Drosophila* CNS and secreted into NBs through a vesicle dependent pathway**

(A) The pattern of Dsred Stinger driven by *Fer1HCH/Fer2LCH-GAL4* in CNS.

(B) The distribution of ferritin labeled by *Fer1HCH^{G188}* in CNS of control and glial ferritin knockdown.

(C) Overexpressed ferritin tagged with GFP and mCherry in glia secreted into NBs.

(D) The distribution of *Fer1HCH^{G188}* when blocking glial vesicular trafficking via overexpressing dominant-negative dynamin.

(E) Quantification of (D), GFP signal in NBs when overexpressing *shi^{ts}* in glia.

(E, n=50, 48; **: $p<0.01$)

**Figure 3. Glial ferritin defects lead to iron deficiency in NBs**

(A) The conditions of brain development in *Drosophila* with feeding iron chelator BPS or supplementary iron FAC.

(B) Iron depletion by adding DFP to food exacerbated the phenotype of NB loss induced by glial ferritin knockdown.

(C) Quantification of NB number in (B).

(D) Zip13 in CNS is produced mainly by glial cells, but not NBs.

(E) Brain defects in *Zip13* knockdown or double knockdown of *Zip13* and *Fer2LCH*.

(F and G) Quantification of NB number and proliferation rate in (E).

(H) Iron supplement can rescue the NB proliferation in *Zip13 RNAi* driven by *repo-GAL4*.

(I) Quantification of NB proliferation in (H).

(C, n=6; F, n=6,10,11,8; G, n=13,10,10,5; I, n=8; ns, not significant; *: p<0.05; **: p<0.01; ****: p<0.0001)

**Figure 3-figure supplement 1. Brain defects induced by glial ferritin knockdown were caused by iron deficiency**

(A) Iron deficiency by blocking iron absorption in gut also induced similar brain defects caused by BPS addition.

(B) Quantification of NB number in (A).

(C) Quantification of NB proliferation in (A).

(D) Brain development of glial ferritin knockdown after feeding FAC.

(E) The ratio of brain with NB loss after adding FAC.

(F) NB number and proliferation of *Zip13* mutant.

(G and H) Quantification of NB number and proliferation rate in (F).

(B, n=10; C, n=5; E, n=3; G, n=10,9; H, n=9,8; *: p<0.05; **: p<0.01; ***: p<0.001; ****: p<0.0001)

**Figure 3-figure supplement 2. ROS accumulates in CNS after glial ferritin knockdown, but inhibiting ROS cannot restore NB number and proliferation**

(A) ROS level marked by Gstd-GFP after knocking down ferritin in glia.

(B) Quantification of ROS level in (A).

(C) Overexpress antioxidant sod1 or cat in glial cells.

(D) Overexpress antioxidant sod1 or cat in NBs.

(E) Mitochondrial morphology by TEM upon glial ferritin knockdown.

(F) Quantification of normal mitochondria in (E). Intact mitochondria with their cristae visible were considered normal.

(B, n=5,4,5; F, n=23,27,28; ****: p<0.0001; ns, not significant)

**Figure 3-figure supplement 3. Glial ferritin knockdown does not induce apoptosis in the CNS**

(A) Apoptotic signal labeled by Caspase-3 of glial ferritin knockdown.

(B) Quantification of Caspase-3 negative NB ratio.

(C) TUNEL staining of *Fer1HCH RNAi* driven by *repo-GAL4*.

(D) Quantification of TUNEL positive cell per $1 \times 10^5 \ \mu m^2$.

(E) P35 overexpression in glia could not rescue brain defects of ferritin knockdown.

(F) Glial morphology was intact when knocking down glial ferritin.

(B, n=5,4,4; D, n=3; ns, not significant)

**Figure 4. Glial ferritin defects result in impaired Fe-S cluster activity and ATP production**

(A) The determination of cytosolic aconitase activity.

(B) Knockdown of CIA complex in NBs.

(C) Quantification of NB proliferation in (B).

(D) The schematic diagram of NB sorting procedure.

(E) Venn diagram of downregulated genes after glial ferritin knockdown.

(F) GO enrichment of down-regulated genes in glial ferritin knockdown comparing with control.

(G) KEGG pathway enrichment of down-regulated genes after glial ferritin knockdown.

(H) $NAD^+$ level was indicated by SoNar.

(I) Quantification of normalized SoNar signal in (H).

(J) The determination of ATP level in CNS.

(K) NB number in glial ferritin knockdown was rescued by Ndi1 overexpression in NBs.

(L) Quantification of NB number in (K).

(A and J, n=3; C, n=5,6,6,9; I, n=37,42; L, n=7,7,6,6,7; *: p<0.05; **: p<0.01; ***: p<0.001;

****: p<0.0001)

**Figure 4-figure supplement 1. Validation of RNA-seq data by qRT-PCR using sorted NB**

**lineages**

(A) qRT-PCR analysis of *ND-15* and *CG15715* to validate the enrichment result.

(n=3; *: p<0.05; **: p<0.01)

**Figure 5. Glial ferritin defects lead to the premature differentiation of NBs**

(A) Temporal window of glial ferritin maintaining NBs. The temperature shift of flies with glial

ferritin knockdown from the permissive temperature (18°C) to the restrictive temperature (29°C).

(B) Quantification of NB number and proliferation in (A).

(C) Pros staining of knocking down glial ferritin.

(D) Quantification of ratio of $Pros^+Dpn^+$ NBs to $Dpn^+$ NBs.

(E) Knockdown of *pros* in NBs rescues the NB number.

(F) The ratio of brain with NB loss in (E).

(B, n=6, 5, 6; n=6, 5, 4; D, n=5; F, n=3; *: p<0.05; ***: p<0.001; ****: p<0.0001)

**Figure 5-figure supplement 1. NB loss upon glial ferritin knockdown is not due to NB origin**

**and apoptosis**

(A) Temporal window of glial ferritin maintaining NBs. The temperature shift of flies with glial

ferritin knockdown from the restrictive temperature (29°C) to the permissive temperature (18°C).

(B) Quantification of NB proliferation in (A).

(C) The CNS of the first-instar larvae with glial ferritin knockdown.

(D) P35 overexpression in NBs could not rescue the brain defects induced by glial ferritin

knockdown.

(B, n=5; ns, not significant)

**Figure 5-figure supplement 2. Glial ferritin defects do not affect NB reactivation**

(A) NB size upon glial ferritin knockdown was not changed.

(B) Activation of Insulin signaling could not restore NB number and proliferation.

(C) Quantification of NB proliferation in (B).

(A, n=100; C, n=8, 5; ns, not significant)

**Figure 6. Ferritin functions as a potential target for tumor suppression**

(A) Knockdown of ferritin in glia could inhibit the tumor induced by *brat RNAi* or *numb RNAi*.

(B) Quantification of tumor size in (A).

(C) Iron chelator BPS supplemented in the food suppressed brain tumor growth.

(D) Quantification of tumor size in (C).

(E) Bioluminescence signal in mice with glioma on day 14.

(F) Quantification of bioluminescence signal in (E).

(G) Kaplan-Meier survival curve of mice treated with DFP.

(B, n=10,7,7,5; D, n=8,6,6,6; F, n=6,7; G, n=7,6; *: p<0.05; ***: p<0.001; ****: p<0.0001)

**Figure 6-figure supplement 1. Iron chelator DFP inhibits glioma progression in mice**

(A) The appearance and HE staining of brains in mice with DFP injection on day 7, 14 and 21.

Scar bar: 1000 μm.

(B and C) Quantification of HE staining in murine brains on day 14 (B) and 21 (C).

(B, n=6; C, n=9,8; ***: p<0.001; ****: p<0.0001.)

**Figure 7. The level of ferritin is controlled by NB proliferation**

(A) Fer1HCH subunit level in *Drosophila* CNS after manipulating proliferation.

(B) Fer2LCH subunit level in *Drosophila* CNS after manipulating proliferation.

(C) Quantification of Fer1HCH level in (A).

(D) Quantification of Fer2LCH level in (B).

(E) The model of glial ferritin regulating NBs.

(C and D, n=3; ns, not significant; *: p<0.05; **: p<0.01; ***: p<0.001)

**Materials and methods**

**Fly Genetics**

Fly strains were bred on yeast-containing medium at constant temperature 25°C in 12h light/dark

cycle unless otherwise stated. UAS-Fer1HCH-RNAi (THU5585) and UAS-Fer2LCH-RNAi

(TH01861.N) were used for most experiments except Figure1-figure supplement 2A (BDSC: 60000,

VDRC: 14491 and VDRC: 12925) and Figure3-figure supplement 3E (VDRC: 14491). *UAS-*

*Fer1HCH*, *UAS-Fer2LCH*, *lexAop-Fer1HCH-RNAi* and *lexAop-Fer2LCH-RNAi* were generated in

this paper. *GstD1-GFP was* from Fisun Hamaratoglu. *repo-lexA* was from Margaret S Ho. *UAS-*

*Ndi1 and UAS-SoNar* were from Juergen A. Knoblich. *Insc-GAL4* was from Yan Song. *UAS-*

*Fer1HCH::GFP;UAS-Fer2LCH::mCherry* was from Fanis Missirlis. *Fer1HCH^{G188}* was from Bing

Zhou. Other fly lines used in this study were bought from Bloomington Drosophila Stock Center,

Tsinghua Fly Center, Vienna Drosophila Resource Center, and Drosophila Genetic Resource Center

(Table 2 and 3).

**Mouse husbandry**

All mouse experiments were performed under the guidelines of the Institutional Animal Care and

Use Committee at Southeast University. Mice were maintained in a barrier facility, at 25°C, on a

regular 12h light and 12h dark cycle.

**DNA and Plasmids**

For generating RNAi-resistant UAS-Fer1HCH and UAS-Fer2LCH constructs used for transgenic

fly lines, Fer1HCH cDNA (PA type) and Fer2LCH cDNA (PA type) with silent mutations in the

region targeted by siRNA were inserted into pUAST plasmids, respectively. The cDNA with silent

mutations was synthesized by Sangon Biotech (Shanghai) Co., Ltd. To generate transgenic fly lines

of lexAop-Fer1HCH-RNAi and lexAop-Fer2LCH-RNAi, the shRNA sequences targeted on

Fer1HCH and Fer2LCH were generated by PCR amplification of UAS-Fer1HCH-RNAi

(TH01861.N) and UAS-Fer2LCH-RNAi (THU5585) DNA using primers (5'-ACGGAGCGACAATTCAATTCA-3'/5'-TGATGCCTACCTGATGCCAA-3'), respectively. The fragments were digested by EcoRI and XbaI, then ligated into lexAop vectors. The constructs were verified by sequencing.

**Immunostaining and Antibodies**

Larval brains were dissected in PBS (1×), fixed in 4% paraformaldehyde/PBS for 20 min and washed 3 times with 0.3% PBST (Triton X-100). Then the brains blocked in 1% BSA for 1h at room temperature. Finally, these brains were incubated with antibodies overnight at 4°C. Primary antibodies: Guinea pig anti-Dpn (1:1000, gift from Juergen A. Knoblich), rabbit anti-PH3 (1/100, CST, 9701), rat anti-Mira (1/1000, abcam, ab197788), mouse anti-Repo (1/100, DSHB, 8D12), rabbit anti-GFP (1/1000, Torrey Pines Biologies, TP401), rabbit anti-Caspase3 (1/100, CST, 9661), mouse anti-Pros (1/20, DSHB), H2v (1/100, ROCKLAND, 600401914).

To detect primary antibodies, the following Alexa-Flour conjugated second antibodies from Invitrogen were used: goat anti-rat Alexa 555 (1/100, A21434), goat anti-guinea pig Alexa 555 (1/100, A21435), donkey anti-rabbit Alexa 488 (1/100, A21206), donkey anti-rat Alexa 488 (1/100, A21208), donkey anti-mouse Alexa 488 (1/100, A21202), goat anti-rat Alexa 633 (1/100, A21094), donkey anti-mouse Alexa 647 (1/100, A31571). All immunostaining images were captured using an LSM700 or an LSM900 (Zeiss) confocal microscope.

**EdU and TUNEL**

Dissected freshly brains were incubated in EdU solution for about 30 min to incorporate EdU, then fixed and permeabilized referring to the steps from immunostaining. EdU was detected by Click-iT EdU Alexa Fluor 555 Imaging Kit according to the instructions. TUNEL detection was determined using kit based on the manufacturer's instructions.

**RNA Isolation, Semiquantitative RT-PCR, and Quantitative Real-time PCR**

Total RNA was extracted from larval brains by TRIzol reagent. cDNA was reverse-transcribed from total RNA using kit. Semiquantitative RT-PCR was performed using the specific primers to amplify the targeted gene segment. Quantitative Real-time PCR was performed on LightCycler 96 (Bio-Rad) using qPCR SuperMix (TransGen Biotech). Following primers were used:

rp49 F: GCACCAAGCACTTCATCC

rp49 R: CGATCTCGCCGCAGTAAA

Fer1HCH F: ATGGTGAAACTAATTGCTAGC

Fer1HCH R: TCAGATCGCTGACTCCCTC

Fer2LCH F: GCATGCTCTACGTCAGCCT

Fer2LCH R: TTACTGCTTCTGCAGATACTC

ND15 F: ATGTCGCTTACCCCCTTTCTACG

ND15 R: CTAGTAGGCATCAACGCGTGG

CG15715 F: ATGGCACGTGGACACCAG

CG15715 R: TCAGACCTCCTTCAGCTCCTCG

**Western blot and Antibodies**

Brains were dissected in PBS, then homogenized in the RIPA buffer with SDS. Homogenate was

centrifuged at 12000 rpm for 5 min at 4℃ three times and boiled for 10 min. The protein was separated on 15% SDS-PAGE and transferred to nitrocellulose membranes. The membranes were blocked in 5% milk and incubated overnight at 4℃ with antibodies. The blots were detected using Tanon 5200. Primary antibodies used: rabbit anti-Fer1HCH (1/500, this study), rabbit anti-Fer2LCH (1/500, this study), mouse anti-tubulin (1/1000, abcam, ab7291), mouse anti-Repo (1/200, DSHB, 8D12). Rabbit polyclonal antibodies were generated against recombinant proteins His-Fer1HCH or His-Fer2LCH. In brief, the aa 23–200 of Fer1HCH (PA type) or aa 41–227 of Fer2LCH (PA type), contained by different Fer1HCH or Fer2LCH cDNA isoforms, was synthesized and cloned into pET28a vector respectively. The recombinant proteins were expressed in *E. coli*, extracted, and injected into rabbit for generating the antibodies (Abclonal Biotech.). Then the antibodies were affinity-purified and applied in western blot.

**Temperature Shift**

GAL80$^{ts}$ was introduced into this system to control the function of GAL4 according to the temperature for determining the temporal window during which ferritin functions to maintain NBs. The eggs of *tubulin-GAL80$^{ts}$; repo-gal4* crossing with *UAS-Fer1HCH RNAi* or *UAS-Fer2LCH RNAi* were collected, and cultured at 18℃ (permissive temperature for GAL80$^{ts}$) before hatching, then transferred to 29℃ (restrictive temperature for GAL80$^{ts}$) after hatching, or vice versa until third-instar larval stage. Then brains were observed and analyzed.

**Culture Medium**

Supplemented iron salt and iron chelators (1×) in *Drosophila* food were used as follows: 5 mM ferric ammonium citrate (FAC) (sigma), 0.1 mM bathophenanthrolinedisulfonic acid disodium (BPS) (sigma), 2.5 mM deferiprone (DFP) (sigma).

**Cell Dissociation and FACS**

NB lineages were labeled with DsRed drived by *Insc-GAL4*. These cells were dissociated and sorted according to previous protocol (Harzer et al., 2013). Briefly, brains were dissected in Schneider's medium and transferred into Rinaldini's solution. Then Rinaldini's solution was removed, and dissociation solution was added to the brains to incubate at 30℃ for 1h. After washing these brains twice in Rinaldini's solution, the brains were disrupted in PBS. Finally, NB lineages were sorted by flow cytometer (Beckman) based on the red signal.

**RNA Sequencing and Data Processing**

Smart-Seq2 libraries preparation
The sorted NBs were collected into microtubes with ribonuclease inhibitor and lysis component. An Oligo-dT primer was added to the reverse transcription reaction for first-strand cDNA synthesis, followed by PCR amplification to enrich the cDNA and magbeads purification to clean up the production. Then qualified cDNA was sheared randomly by ultrasonic waves for Illumina library preparation protocol including DNA fragmentation, end repair, 3' ends A-tailing, adapter ligation, PCR amplification, and library validation. After library preparation, PerkinElmer LabChip®GX Touch and Step OnePlus™ Real-Time PCR System were introduced for library quality inspection. Qualified libraries were then loaded on Illumina Hiseq platform for PE150 sequencing (Annoroad

Gene Technology Co., Ltd).

Quantification and Differential Expression Analysis

The reads containing adapter sequences, low-quality reads or undetermined bases were removed

from raw data using fastp program. The clean reads were aligned to the reference genome of

*Drosophila melanogaster* using HISAT2 (v2.1.0). The Ensembl database was chosen as the

annotation reference. The gene expression was calculated as Fragments Per Kilobase Million

mapped reads (FPKM). The differential expression of genes was analyzed with DEGSeq package.

The criterion of |log2 (fold-change)| $\geq 1$ and p-value $< 0.05$ was used to identify differentially

expressed genes.

GO and KEGG Pathway Enrichment

Gene Ontology (GO) enrichment analysis and Kyoto Encyclopedia of Genes and Genomes (KEGG)

pathway analysis were performed to investigate the function of differentially expressed genes. GO

analysis was used to explore the function of genes, such as biological process, cellular component,

and molecular function. KEGG analysis focused on finding and analyzing interactions in biological

systems, such as signal transduction and disease pathways. Only the gene sets with p-value $< 0.05$

were significantly enriched.

**Aconitase Activity Assay**

Protein was extracted with assay buffer from Aconitase Activity Assay Kit (sigma, MAK051). The

concentration of protein was determined by Pierce™ Rapid Gold BCA Protein Assay Kit (Thermo

Fisher, A53225). Cytosolic aconitase was measured according to the Aconitase Activity Assay Kit.

**TEM observation and analysis**

TEM samples were prepared by standard procedures (Gan, Geng, Zhang, Mou, & Xie, 2020). In

brief, brains from third instar larvae upon WT and glial ferritin knockdown were dissected in fresh

PBS and fixed at 4°C overnight in a mixture of 2% glutaraldehyde and 2% formaldehyde in 0.1 M

sodium cacodylate buffer (pH 7.4), followed by several rinses with cacodylate buffer. The samples

were then postfixed for 2 h with 1% $OsO_4$ in cacodylate buffer and rinsed twice with distilled water.

The preparations were stained for 2 h with 2% saturated uranyl acetate in distilled water and rinsed

twice with distilled water. The specimens were dehydrated in an ethanol series, passed through

propylene oxide two times, and embedded into a sheet in Epon812 (SPI Science). Each slice was

90 nm thick, 30–40 slices were gathered into a group and attached to a grid. Samples were analyzed

with Hitachi H-7650 electron microscope operated at 80kV.

**Orthotopic Glioma Model and DFP Injection**

To establish an orthotopic glioma model (B. Wang et al., 2022), C57BL/6 mice (4-6 weeks) were

anesthetized using 2% isoflurane and positioned in a stereotactic instrument. GL261-luc cell

suspension ($1 \times 10^5$ cells in 3 μL PBS) was injected into the striatum. Specifically, the injection site

is at 0.5 mm anterior, 2 mm lateral to the bregma and 3.5 mm below the skull. The injection was

done slowly about 10 min.

The mice with glioma were injected intraperitoneally with DFP every 48 h (Eybl, Kotyzova,

Kolek, Koutensky, & Nielsen, 2002). DFP was dissolved in saline (0.9% NaCl) to 10 mg/ml. The

doses of injection in mice were 150 mg/kg using 10 mg/ml DFP. The injection volume was determined based on murine weight.

**Bioluminescence Imaging**

For tumor imaging, mice were injected with luciferin (150 μg/g) in PBS (S. L. Wang et al., 2019). After 10 min, the mice were imaged using the IVIS Spectrum (PerkinElmer).

**HE Staining**

At time of sacrifice, mouse brains were removed and fixed in 4 % paraformaldehyde for 24 h. After fixation, brains were immersed in 30 % sucrose and embedded in OCT for cryosectioning. Sections were stained by Hematoxylin and Eosin (HE). Tumor volumes were estimated using Image J.

**Quantifications and Statistical Analysis**

For quantification of NSCs, Dpn- or Mira-positive NSC of CB or the thoracic VNC at the indicated stage were counted. Mitotic index is the number of PH3-positive cells among Dpn-positive cells. For tumor size in *Drosophila*, the area of CNS was measured by NIH ImageJ. For quantifications of NB size, fluorescence or western blot intensity, NIH ImageJ was used to measure.

The figure legends showed the number of samples, statistical parameters, and significance. GraphPad Prism 6 was used to perform statistical analyses. For comparisons between two groups, unpaired two-sided Student's t test was employed to assess statistical significance. For comparisons between three of more groups, statistical significance was determined using one-way ANOVA with a Bonferroni test. The log-rank test was used to compare the survival curves of mice. The sample distributions were assumed to be normal and no particular statistical tests were used to check for normality within the samples. Statistical results were presented as means ± SEM and p values of less than 0.05 were considered statistically significant. Asterisks indicate critical levels of significance (*: $p<0.05$; **: $p<0.01$; ***: $p<0.001$; ****: $p<0.0001$).

**Data availability**

RNA-seq data are available at the GEO database, accession number GEO GSE237124 and reviewer token: cnqxqoqonzapjst

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

**Acknowledgments**

We are grateful to Drs. Bing Zhou, Margaret S Ho, Juergen A. Knoblich, Fanis Missirlis, Fisun Hamaratoglu, Yan Song, Andrea H. Brand; Bloomington Drosophila Stock Center (Indiana University, Bloomington, Indiana), the Drosophila Genetic Resource Center (Kyoto Institute of Technology, Kyoto, Japan), the Vienna Drosophila RNAi Center (Vienna, Austria), and Tsinghua Fly Center (Tsinghua University) for fly stocks and reagents. We thank Dr. Bing Zhou for the useful advice on this study. We thank Drs. Junhua Geng, and Mingdao Mu for comments on the manuscript. This study was supported by the National Natural Science Foundation of China (31970675).

**Funding**

National Natural Science Foundation of China (grant no. 31970675; grant recipient, Su Wang).

**Author contributions**

Zhixin Ma, Resources, Data curation, Formal analysis, Validation, Investigation, Methodology, Writing - original draft; Wenshu Wang, Resources, Formal analysis, Investigation, Methodology; Xiaojing Yang, Formal analysis, Methodology; Menglong Rui, Writing - review and editing; Su Wang, Conceptualization, Resources, Supervision, Funding acquisition, Investigation, Project administration, Writing - review and editing.

**Competing interests**

The authors declare no competing interests.

**References**

Andrews, N. C. (2005). Molecular control of iron metabolism. Best Pract Res Clin Haematol, 18(2), 159-169. doi:10.1016/j.beha.2004.10.004

Arosio, P., Ingrassia, R., & Cavadini, P. (2009). Ferritins: A family of molecules for iron storage, antioxidation and more. Biochimica Et Biophysica Acta-General Subjects, 1790(7), 589-599. doi:10.1016/j.bbagen.2008.09.004

Bailey, A. P., Koster, G., Guillermier, C., Hirst, E. M., MacRae, J. I., Lechene, C. P., . . . Gould, A. P. (2015). Antioxidant Role for Lipid Droplets in a Stem Cell Niche of Drosophila. Cell, 163(2), 340-353. doi:10.1016/j.cell.2015.09.020

Bjornsson, C. S., Apostolopoulou, M., Tian, Y., & Temple, S. (2015). It takes a village: constructing the neurogenic niche. Dev Cell, 32(4), 435-446. doi:10.1016/j.devcel.2015.01.010

Bonnay, F., Veloso, A., Steinmann, V., Kocher, T., Abdusselamoglu, M. D., Bajaj, S., . . . Knoblich, J. A. (2020). Oxidative Metabolism Drives Immortalization of Neural Stem Cells during Tumorigenesis. Cell, 182(6), 1490-1507 e1419. doi:10.1016/j.cell.2020.07.039

Bowman, S. K., Rolland, V., Betschinger, J., Kinsey, K. A., Emery, G., & Knoblich, J. A. (2008). The tumor suppressors brat and numb regulate transit-amplifying neuroblast lineages in Drosophila. Developmental Cell, 14(4), 535-546. doi:10.1016/j.devcel.2008.03.004

Cabernard, C., & Doe, C. Q. (2009). Apical/Basal Spindle Orientation Is Required for Neuroblast Homeostasis and Neuronal Differentiation in Drosophila. Developmental Cell, 17(1), 134-141. doi:10.1016/j.devcel.2009.06.009

Chell, J. M., & Brand, A. H. (2010). Nutrition-responsive glia control exit of neural stem cells from quiescence. Cell, 143(7), 1161-1173. doi:10.1016/j.cell.2010.12.007

Chiou, B., Neal, E. H., Bowman, A. B., Lippmann, E. S., Simpson, I. A., & Connor, J. R. (2019). Endothelial cells are critical regulators of iron transport in a model of the human blood-brain barrier. J Cereb Blood Flow Metab, 39(11), 2117-2131. doi:10.1177/0271678X18783372

Choksi, S. P., Southall, T. D., Bossing, T., Edoff, K., de Wit, E., Fischer, B. E., . . . Brand, A. H. (2006). Prospero acts as a binary switch between self-renewal and differentiation in Drosophila neural stem cells. Developmental Cell, 11(6), 775-789. doi:10.1016/j.devcel.2006.09.015

Eybl, V., Kotyzova, D., Kolek, M., Koutensky, J., & Nielsen, P. (2002). The influence of deferiprone (L1) and deferoxamine on iron and essential element tissue level and parameters of oxidative status in dietary iron-loaded mice. Toxicology Letters, 128(1-3), 169-175. doi:Pii

S0378-4274(01)00541-0

Doi 10.1016/S0378-4274(01)00541-0

Fan, K., Cao, C., Pan, Y., Lu, D., Yang, D., Feng, J., . . . Yan, X. (2012). Magnetoferritin
nanoparticles for targeting and visualizing tumour tissues. Nat Nanotechnol, 7(7), 459-464.
doi:10.1038/nnano.2012.90

Fan, K., Jia, X., Zhou, M., Wang, K., Conde, J., He, J., . . . Yan, X. (2018). Ferritin Nanocarrier
Traverses the Blood Brain Barrier and Kills Glioma. ACS Nano, 12(5), 4105-4115.
doi:10.1021/acsnano.7b06969

Fisher, J., Devraj, K., Ingram, J., Slagle-Webb, B., Madhankumar, A. B., Liu, X., . . . Connor, J. R.
(2007). Ferritin: a novel mechanism for delivery of iron to the brain and other organs. Am
J Physiol Cell Physiol, 293(2), C641-649. doi:10.1152/ajpcell.00599.2006

Fretham, S. J., Carlson, E. S., & Georgieff, M. K. (2011). The role of iron in learning and memory.
Adv Nutr, 2(2), 112-121. doi:10.3945/an.110.000190

Gan, G. M., Geng, J. H., Zhang, C. C., Mou, Y., & Xie, W. (2020). Neurexin and Neuroligins
Maintain the Balance of Ghost and Satellite Boutons at the Drosophila Neuromuscular
Junction. Frontiers in Neuroanatomy, 14. doi:ARTN 19

10.3389/fnana.2020.00019

Haile, D. J., Rouault, T. A., Harford, J. B., Kennedy, M. C., Blondin, G. A., Beinert, H., & Klausner,
R. D. (1992). Cellular regulation of the iron-responsive element binding protein:
disassembly of the cubane iron-sulfur cluster results in high-affinity RNA binding. Proc
Natl Acad Sci U S A, 89(24), 11735-11739. doi:10.1073/pnas.89.24.11735

Hartenstein, V. (2011). Morphological Diversity and Development of Glia in Drosophila. Glia, 59(9),
1237-1252. doi:10.1002/glia.21162

Harzer, H., Berger, C., Conder, R., Schmauss, G., & Knoblich, J. A. (2013). FACS purification of
Drosophila larval neuroblasts for next-generation sequencing. Nat Protoc, 8(6), 1088-1099.
doi:10.1038/nprot.2013.062

Hentze, M. W., Muckenthaler, M. U., Galy, B., & Camaschella, C. (2010). Two to Tango: Regulation
of Mammalian Iron Metabolism. Cell, 142(1), 24-38. doi:10.1016/j.cell.2010.06.028

Hirschhorn, T., & Stockwell, B. R. (2019). The development of the concept of ferroptosis. Free
Radical Biology and Medicine, 133, 130-143. doi:10.1016/j.freeradbiomed.2018.09.043

Ito, K., Urban, J., & Technau, G. M. (1995). Distribution, Classification, and Development of
Drosophila Glial-Cells in the Late Embryonic and Early Larval Ventral Nerve Cord. Rouxs
Archives of Developmental Biology, 204(5), 284-307. doi:Doi 10.1007/Bf02179499

Knovich, M. A., Storey, J. A., Coffman, L. G., Torti, S. V., & Torti, F. M. (2009). Ferritin for the
clinician. Blood Rev, 23(3), 95-104. doi:10.1016/j.blre.2008.08.001

Lee, J. S., Yang, K., Cho, A. N., & Cho, S. W. (2018). Ferritin nanoparticles for improved self-
renewal and differentiation of human neural stem cells. Biomater Res, 22, 5.
doi:10.1186/s40824-018-0117-y

Li, S. (2010). Identification of iron-loaded ferritin as an essential mitogen for cell proliferation and
postembryonic development in Drosophila. Cell Res, 20(10), 1148-1157.
doi:10.1038/cr.2010.102

Lind, M. I., Ekengren, S., Melefors, O., & Soderhall, K. (1998). Drosophila ferritin mRNA:
alternative RNA splicing regulates the presence of the iron-responsive element. FEBS Lett,
436(3), 476-482. doi:10.1016/s0014-5793(98)01186-7

Lozoff, B., Beard, J., Connor, J., Barbara, F., Georgieff, M., & Schallert, T. (2006). Long-lasting
neural and behavioral effects of iron deficiency in infancy. Nutr Rev, 64(5 Pt 2), S34-43;
discussion S72-91. doi:10.1301/nr.2006.may.s34-s43

Ly, P. T., Tan, Y. S., Koe, C. T., Zhang, Y., Xie, G., Endow, S., . . . Wang, H. (2019). CRL4Mahj E3
ubiquitin ligase promotes neural stem cell reactivation. PLoS Biol, 17(6), e3000276.
doi:10.1371/journal.pbio.3000276

Missirlis, F., Kosmidis, S., Brody, T., Mavrakis, M., Holmberg, S., Odenwald, W. F., . . . Rouault, T.
A. (2007). Homeostatic mechanisms for iron storage revealed by genetic manipulations and
live imaging of Drosophila ferritin. Genetics, 177(1), 89-100.
doi:10.1534/genetics.107.075150

Moller, H. E., Bossoni, L., Connor, J. R., Crichton, R. R., Does, M. D., Ward, R. J., . . . Ronen, I.
(2019). Iron, Myelin, and the Brain: Neuroimaging Meets Neurobiology. Trends Neurosci,
42(6), 384-401. doi:10.1016/j.tins.2019.03.009

Mukherjee, C., Kling, T., Russo, B., Miebach, K., Kess, E., Schifferer, M., . . . Simons, M. (2020).
Oligodendrocytes Provide Antioxidant Defense Function for Neurons by Secreting Ferritin
Heavy Chain. Cell Metab, 32(2), 259-272 e210. doi:10.1016/j.cmet.2020.05.019

Mumbauer, S., Pascual, J., Kolotuev, I., & Hamaratoglu, F. (2019). Ferritin heavy chain protects the
developing wing from reactive oxygen species and ferroptosis. PLoS Genet, 15(9),
e1008396. doi:10.1371/journal.pgen.1008396

Nichol, H., Law, J. H., & Winzerling, J. J. (2002). Iron metabolism in insects. Annual Review of
Entomology, 47, 535-559. doi:DOI 10.1146/annurev.ento.47.091201.145237

Nurtjahja-Tjendraputra, E., Fu, D., Phang, J. M., & Richardson, D. R. (2007). Iron chelation
regulates cyclin D1 expression via the proteasome: a link to iron deficiency-mediated
growth suppression. Blood, 109(9), 4045-4054. doi:10.1182/blood-2006-10-047753

Pereanu, W., Shy, D., & Hartenstein, V. (2005). Morphogenesis and proliferation of the larval brain
glia in Drosophila. Developmental Biology, 283(1), 191-203.
doi:10.1016/j.ydbio.2005.04.024

Pham, D. Q., & Winzerling, J. J. (2010). Insect ferritins: Typical or atypical? Biochim Biophys Acta,
1800(8), 824-833. doi:10.1016/j.bbagen.2010.03.004

Read, R. D. (2018). Pvr receptor tyrosine kinase signaling promotes post-embryonic morphogenesis,
and survival of glia and neural progenitor cells in Drosophila. Development, 145(23).
doi:10.1242/dev.164285

Reichardt, I., Bonnay, F., Steinmann, V., Loedige, I., Burkard, T. R., Meister, G., & Knoblich, J. A.
(2018). The tumor suppressor Brat controls neuronal stem cell lineages by inhibiting
Deadpan and Zelda. EMBO Rep, 19(1), 102-117. doi:10.15252/embr.201744188

Rice, D., & Barone, S., Jr. (2000). Critical periods of vulnerability for the developing nervous
system: evidence from humans and animal models. Environ Health Perspect, 108 Suppl 3,
511-533. doi:10.1289/ehp.00108s3511

Shetty, A. K., Hattiangady, B., & Shetty, G. A. (2005). Stem/progenitor cell proliferation factors
FGF-2, IGF-1, and VEGF exhibit early decline during the course of aging in the
hippocampus: role of astrocytes. Glia, 51(3), 173-186. doi:10.1002/glia.20187

Sousa-Nunes, R., Yee, L. L., & Gould, A. P. (2011). Fat cells reactivate quiescent neuroblasts via
TOR and glial insulin relays in Drosophila. Nature, 471(7339), 508-512.
doi:10.1038/nature09867

Speder, P., & Brand, A. H. (2014). Gap junction proteins in the blood-brain barrier control nutrient-
dependent reactivation of Drosophila neural stem cells. Dev Cell, 30(3), 309-321.
doi:10.1016/j.devcel.2014.05.021
Speder, P., & Brand, A. H. (2018). Systemic and local cues drive neural stem cell niche remodelling
during neurogenesis in Drosophila. Elife, 7. doi:10.7554/eLife.30413
Sykiotis, G. P., & Bohmann, D. (2008). Keap1/Nrf2 signaling regulates oxidative stress tolerance
and lifespan in Drosophila. Dev Cell, 14(1), 76-85. doi:10.1016/j.devcel.2007.12.002
Tang, X., & Zhou, B. (2013). Ferritin is the key to dietary iron absorption and tissue iron
detoxification in Drosophila melanogaster. FASEB J, 27(1), 288-298. doi:10.1096/fj.12-
213595
Tong, W. H., & Rouault, T. A. (2006). Functions of mitochondrial ISCU and cytosolic ISCU in
mammalian iron-sulfur cluster biogenesis and iron homeostasis. Cell Metab, 3(3), 199-210.
doi:10.1016/j.cmet.2006.02.003
van den Ameele, J., & Brand, A. H. (2019). Neural stem cell temporal patterning and brain tumour
growth rely on oxidative phosphorylation. Elife, 8. doi:10.7554/eLife.47887
Wang, B., Tang, M. P., Yuan, Z. W., Li, Z. Y., Hu, B., Bai, X., . . . Zhang, X. Q. (2022). Targeted
delivery of a STING agonist to brain tumors using bioengineered protein nanoparticles for
enhanced       immunotherapy.       Bioactive       Materials,       16,       232-248.
doi:10.1016/j.bioactmat.2022.02.026
Wang, S. L., Yao, F. L., Lu, X. H., Li, Q., Su, Z. P., Lee, J. H., . . . Du, L. Y. (2019). Temozolomide
promotes immune escape of GBM cells via upregulating PD-L1. American Journal of
Cancer Research, 9(6), 1161-+.
Xiao, G., Liu, Z. H., Zhao, M., Wang, H. L., & Zhou, B. (2019). Transferrin 1 Functions in Iron
Trafficking and Genetically Interacts with Ferritin in Drosophila melanogaster. Cell Rep,
26(3), 748-758 e745. doi:10.1016/j.celrep.2018.12.053
Xiao, G., Wan, Z., Fan, Q., Tang, X., & Zhou, B. (2014). The metal transporter ZIP13 supplies iron
into   the   secretory   pathway   in   Drosophila   melanogaster.   Elife,   3,   e03191.
doi:10.7554/eLife.03191

| AL4 driver | Stock center | Stock number | Expression pattern | Fer1HCH RNAi | Fer2LCH RNAi |
|------------|--------------|--------------|--------------------|--------------|--------------|
| *repo-GAL4* | BDSC | 7415 | all glia | +++ | +++ |
| *nrv2-GAL4* | BDSC | 6799 | cortex glia | + | + |
| *mdr65-GAL4* | BDSC | 50472 | subperineurial glia | - | - |
| *moody-GAL4* | BDSC | 90883 | subperineurial glia | - | - |
| *NP6293-GAL4* | DGRC | 105188 | perineurial glia | - | - |
| *alrm-GAL4* | BDSC | 67032 | astrocyte-like glia | - | - |
| *NP6520-GAL4* | DGRC | 105240 | ensheathing glia | - | - |

952  **Table 1. The phenotype of ferritin knockdown in different glial subpopulations.** BDSC:
953  Bloomington Drosophila Stock Center; DGRC: Drosophila Genetic Resource Center; +++: strong
954  phenotype; ++: weak phenotype; -: no phenotype.

# Figure 1

# Figure 1—figure supplement 1

**Figure 1—figure supplement 2**

**A**

*repo>*

*mCherry RNAi*　　*Fer1HCH RNAi* (BL60000)　　*Fer1HCH RNAi* (V12925)　　*Fer2LCH RNAi* (V14491)

Dpn
PH3
50 µm

**B**

NB number

****
****　****　****

GFP; mCherry RNAi
GFP; Fer1HCH RNAi
Fer1HCH; Fer1HCH RNAi
GFP; Fer2LCH RNAi
Fer2LCH; Fer2LCH RNAi

*repo>*

**C**

Dpn$^+$PH3$^+$ NB/ Dpn$^+$ NB

0.4
0.3
0.2
0.1
0.0

****
****　****　****

GFP; mCherry RNAi
GFP; Fer1HCH RNAi
Fer1HCH; Fer1HCH RNAi
GFP; Fer2LCH RNAi
Fer2LCH; Fer2LCH RNAi

*repo>*

**D**

*repo>*

*GFP*　　*Fer1HCH$^{DN}$*

Dpn
PH3
50 µm

**Figure 2**

**A**

*Fer1HCH/Fer2LCH-GAL4>Dsred Stinger*

Dpn
DsRed
50 µm

Repo
DsRed

**B**

*Fer1HCH$^{G188}$, repo-GAL4 >*

*mCherry RNAi*  |  *Fer1HCH RNAi*  |  *Fer2LCH RNAi*

GFP
Mira
50 µm

**C**

*repo>*

*Fer1HCH:GFP, Fer2LCH:mCherry*  |  *GFP, mCherry*

Mira
2 µm
GFP

mCherry
Merge

Mira
2 µm
GFP

mCherry
Merge

**D**

*Fer1HCH$^{G188}$, repo-GAL4>*

*LacZ*  |  *shi$^{ts}$*

GFP
Dpn
50 µm

**E**

GFP intensity per NB

**

*LacZ*  *shi$^{ts}$*

*Fer1HCH$^{G188}$, repo-GAL4>*

**Figure 3**

**Figure 3—figure supplement 1**

# Figure 3—figure supplement 2

**A**

*repo>*

mCherry RNAi      Fer1HCH RNAi      Fer2LCH RNAi

GstD1-GFP

50 µm

**B**

Mean intensity of GstD1-GFP

****
****

mCherry RNAi   Fer1HCH RNAi   Fer2LCH RNAi

*repo>*

**C**

*repo>*

Fer1HCH RNAi     Fer1HCH RNAi, cat     Fer2LCH RNAi, sod1

Dpn
PH3

50 µm

**D**

*repo-lexA>Fer2LCH RNAi*

Insc-GAL4>UAS-sod1     Insc-GAL4>UAS-cat

Dpn
PH3

50 µm

**E**

*repo>*

mCherry RNAi      Fer1HCH RNAi      Fer2LCH RNAi

**F**

Normal Mitochondria (%)

ns
ns

mCherry RNAi   Fer1HCH RNAi   Fer2LCH RNAi

*repo>*

# Figure 4

**Figure 4—figure supplement 1**

A

# Figure 5

# Figure 5—figure supplement 1

**A**

29℃ — Embryo — RNAi activated — Dissection

RNAi activated / RNAi inactivated — 18℃

(29℃ to 18℃) *Tubulin-GAL80 ts; repo-GAL4>*

*Fer1HCH RNAi*        *Fer2LCH RNAi*

Dpn
PH3
50 μm

**B**

Dpn⁺PH3⁺ NB/ Dpn⁺ NB

ns
ns

*repo>mCherry RNAi*    *Fer1HCH RNAi*    *Fer2LCH RNAi*

*Tubulin-GAL80 ts; repo-GAL4>*

**C**

*repo>mCherry RNAi*

repo    Dpn    PH3    DAPI    Merge
20 μm

*repo>Fer1HCH RNAi*

*repo>Fer2LCH RNAi*

**D**

*repo-lexA*        *repo-lexA>Fer2LCH RNAi*

*Insc-GAL4>UAS-p35*

Dpn
PH3
50 μm

**A**

**B**

*repo-lexA>Fer2LCH RNAi*

*Insc-GAL4>UAS-InR $^{act}$*

Dpn
PH3

50 μm

**C**

ns

*Insc-GAL4>*
*UAS-InR $^{act}$*

*repo-lexA>Fer2LCH RNAi*

# Figure 6

**Figure 6—figure supplement 1**

# Figure 7