1   **Structural and biochemical requirements for secretory component interactions with**

2   **dimeric Immunoglobulin A**

Sonya Kumar Bharathkar and Beth M. Stadtmueller[1,2,3*]

[1]Department of Biochemistry, University of Illinois Urbana-Champaign, Urbana, Illinois 61801 USA

[2]Department of Biomedical and Translational Sciences, Carle Illinois College of Medicine, University of

Illinois Urbana-Champaign, Urbana, Illinois 61801 USA

[3]Carle R. Woese Institute of Genomic Biology

*Address correspondence to bethms@illinois.edu–

ABSTRACT

Secretory (S) Immunoglobulin (Ig) A is the predominant mucosal antibody that protects host epithelial

barriers and promotes microbial homeostasis. SIgA production occurs when plasma cells assemble two

copies of monomeric IgA and one joining chain (JC) to form dimeric (d) IgA, which is bound by the

polymeric Ig receptor (pIgR) on the basolateral surface of epithelial cells and transcytosed to the apical

surface. There, pIgR is proteolytically cleaved, releasing SIgA, a complex of the dIgA and the pIgR

ectodomain, called secretory component (SC). The pIgR has five Ig-like domains (D1-D5) that undergo a

conformational change upon binding dIgA, ultimately contacting four IgA heavy chains and the JC in SIgA.

Here we report structure-based mutational analysis combined with surface plasmon resonance binding

assays that identify key residues in mouse SC D1 and D3 that mediate SC binding to dIgA. Residues in D1

CDR3 are likely to initiate binding whereas residues that stabilize the D1-D3 interface are likely to promote

the conformation change and stabilize the final SIgA structure. Additionally, we find that the three C-

terminal residues of JC play a limited role in dIgA assembly but a significant role in pIgR/SC binding to

dIgA. Together results inform new models for the intricate mechanisms underlying IgA transport across

epithelia and functions in the mucosa.

INTRODUCTION

Secretory Immunoglobin A (SIgA) is a crucial antibody in the mucosal immune system that helps maintain the balance of microorganisms and protect the mucosal barrier (1),(2). The assembly of SIgA begins in the lamina propria, where plasma cells link one Joining chain (JC) and between two to five monomeric (m) IgA to form or polymeric (p) IgA; in humans the predominant form is dimeric (d) IgA(3),(4),(5). Subsequently, pIgA binds to polymeric Ig Receptor (pIgR) on the basolateral surface of mucosal epithelia and is then transcytosed to the apical side(6),(7). In the mucosa, proteases cleave the pIgR ectodomain from the cell surface, releasing a complex containing pIgR's five Ig-like domains (D1-D5), collectively referred to as Secretory Component (SC), bound to the pIgA(8),(9). The SC-pIgA complex, which typically includes dIgA, is referred to as SIgA(10).

Each mIgA consists of two heavy chains (HC) and two light chains (LC) that together form two antigen binding fragments (Fabs) and one constant fragment (Fcα), and thus SIgA contains at least four Fabs and two Fcs (Fig.1). The cryo-EM structures of SIgA(11),(12),(13) and dIgA(11) revealed a remarkably asymmetric molecule with individual mIgAs bent and tilted with respect to each other. This arrangement results in each HC (HC-A-D) adopting a unique structure and unique contacts with other chains, despite sharing the same sequence. The greatest structural variability is found among copies of the HC's 18-22 residue C-terminal extension, called the tailpiece (Tp) (11),(12),(13). The four Tps (TpA-D) co-fold with JC to form a β-sandwich, in which the Tps closest to JC, TpA and TpC, are disulfide bonded to the JC with their penultimate Cys residue(11),(12),(13),(14). The other two Tps, TpB and TpD, are partially disordered in dIgA, but are ordered in SIgA, indicating a possible role in pIgR binding and successful SIgA formation(11). In addition to contacting the Tps, the JC also contacts HC residues in the Fcα regions, through several β-hairpins, termed *wings*. JC_{wing-1} and JC_{wing-2} contact HC-C and HC-A respectively, and appear to contribute to the bent, asymmetric conformation observed for both dIgA and SIgA. Additionally JC also contacts parts of SC with its C-terminal residues interfacing with SC-D1, and JC_{wing-1} interfacing with SC-D4-D5 (11),(12),(13).

Figure-1

Figure-1: The structure of liganded and unliganded SC. (Top) Cartoon representation of a mouse pIgR ectodomain homology model based on the human SC crystal structure (PDB:5D4K). The pIgR is depicted bound to an epithelial cell membrane with its five constitutive immunoglobin-like domains (D1-D5) adopting a closed, unliganded conformation. Upon encountering dIgA, D1-D5 undergoes conformational change (indicated by red arrows) to form SIgA (PDB:7JG2), which is shown as a cartoon representation in two orientations (Fabs are disordered and represented with dotted schematics in grey). The dimeric SIgA contains four heavy chains (HC-A to D), J-chain (JC), and one Secretory Component (SC). All chains are colored according to the key. The Fc formed by HC-A and HC-B is indicated as Fc_{AB} (green) and the Fc formed by HC-C and HC-D is indicated as Fc_{CD} (of blue). The JC occupies the center of the SIgA complex with Fc_{AB} and Fc_{CD} on opposite sides of. The SC has five immunoglobin domains (D1-D5), and each domain (D1-D5) are colored with different shades of cyan. The yellow asterisk marks the location of D1 CDR loops in unliganded and liganded structures and the orange asterisk indicated the location of the D1-D3 interface in the liganded (SIgA) structure. Color scheme and labeling approach are adapted from S. Kumar Bharathkar *et al.*, "The structures of secretory and dimeric immunoglobulin A," *eLife*, vol. 9, p. e56098, Oct. 2020 [11].

Given critical roles in SIgA transport and function, the mechanisms of pIgR binding to pIgA have long been of interest[8],[15],[16],[17],[18]. Earlier studies demonstrated that pIgR (and SC) D1 is both necessary and sufficient for binding to dIgA, but that D2-D5 further stabilize the SIgA complex[15],[16],[18],[19], and structural and biophysical studies have demonstrated that pIgR/SC domains undergo a massive conformational change upon dIgA binding (Fig.1) [8]. The SIgA structures further revealed contacts between D1 and Fc_{AB} (the $Fc\alpha$ formed by HC-A and HC-B) and TpC and TpD, as well as D4-D5 contacts with Fc_{CD} (the $Fc\alpha$ formed by HC-C and HC-D)[11],[12],[13]. Notably these interactions were accompanied by new inter-domain contacts between D1-D3 and D3-D4 not observed in the unliganded SC structure (Fig.S1). Furthermore, pIgR (and SC) binding to pIgA has long been appreciated to require the JC[4],[20],[21]. SIgA structures suggest that the JC directly provides contacts for pIgR (or SC) D1 binding and indirectly provides the bent asymmetric conformation needed for other pIgR (or SC) domains to bind[11],[12],[13].

Taken together, pIgR binding to pIgA can be described as a multi-step process involving multiple interactions between pIgA and multiple SC domains (D1-D5), that undergo conformational changes to bind four IgA HCs and the JC, while also forming new inter-domain contacts within SC (Fig.1, Fig.S1); yet, aspects of this remarkably intricate mechanism remain unclear, such as the importance of inter-domain contacts within SC (D1-D5), and the role of JC and Tps in pIgR/SC binding. Building upon recent structural findings, we hypothesized that the conformational change-induced formation of pIgR/SC inter-domain contacts, together with binding to the C terminus of JC is a critical step in SIgA assembly. To test this hypothesis, we carried out structure-based mutational analysis combined with Surface Plasmon Resonance (SPR) binding assays, that test the roles of pIgR/SC and JC residues in SIgA complex formation. Results signify that formation of the D1-D3 interface and D1 binding to the C-terminal residues of JC and Tps are pivotal steps necessary to achieve stable SIgA formation and subsequent effector functions.

RESULTS

To investigate the molecular mechanisms underlying SIgA assembly we analyzed existing SC(8), dIgA(11) and SIgA(11),(12),(13) structures and identified SC and JC residues with potential to mediate the pIgR binding mechanism, for example, residues that may promote SC conformational changes and/or stabilize SIgA interfaces. Subsequently, we mutated target residues in recombinantly expressed full-length SC and D1-only variants and tested binding to immobilized recombinant dIgA lacking the Fabs (dFcα) using SPR binding assays; we also tested unmutated SC and D1-only binding to dFcα containing JC mutant variants. SC and D1 analytes provide an established, soluble model for pIgR binding to dIgA(8). Consistent with the ligand-binding-induced conformational change observed for full-length SC, published SC-dIgA binding studies do not fit single state kinetic models, and thus similar to prior studies we followed the approach of qualitatively monitoring differences in the SPR binding profiles to identify differences among protein variants(8),(22), while also calculating equilibrium binding affinities for variants containing only D1. All experiments utilized mouse SC, Fcα and JC sequences.

D1-CDR3 plays a crucial role in the initiation of SC binding.

SC D1-D5 Ig-like domains each contain up to three loops (1-3) that are structurally similar to antibody complementarity determining regions (CDRs)(8). SIgA structures revealed residues in SC D1, including those in CDRs 1-3, mediating contacts with all four HCs and JC (Fig.S1C and Fig.S1F). However, the unliganded SC crystal structure revealed that a subset of these residues, including those in D1-CDR1 and CDR2, interact with D4 and D5 leaving them sterically occluded and likely unavailable to initiate binding between SC-D1 and dIgA, while D1-CDR3 is solvent exposed and does not participate in any interdomain interactions (Fig.S1C and Fig.S1F). This observation led us to hypothesize that D1-CDR3, which is solvent exposed and sterically accessible in unliganded SC, initiates pIgR binding to dIgA.

Figure-2

Figure-2: Contribution of mSC-D1-CDR3 towards binding dIgA. (A) Detailed view of mSC-D1-CDR3 and dIgA interactions found in mouse SIgA (PDB: 7JG2) (Fig.1, yellow asterisk); the structure is shown as a cartoon representation with interacting residues shown as sticks and labeled. D1-CDR3 residues that were mutated are underlined, and the unmutated and mutated sequences are shown on the right. (B) SPR sensorgrams showing the response (top) and normalized response (bottom) of 0.5uM mSC-wt (black) and mSC-D1-CDR3mut (red) binding to dFcα-wt. (C) SPR sensorgrams showing the response of 0.5uM of mD1-wt (black) and mD1-CDR3mut (red) binding to dFcα-wt (top) and the concentration dependent binding and steady state fit (bottom). The calculated K_D values for each analyte are indicated to the right of the name.

In SIgA structures, SC-D1-CDR3 residues contact HC-B, TpC and TpD though hydrogen bonds and ionic interactions, and additionally, lie within 3.5-6Å of JC, where they may mediate Van-der Waals interactions (Fig.2A). To explore the hypothesis that SC-D1-CDR3 is crucial for SC and dIgA binding, we generated SC expression constructs, in which the SC-D1-CDR3 residues were mutated to Glycine and Alanine (CDR3-wt: GTSNRGL to CDR3-mut: GAGGAGA), limiting the possible side chain interactions with dIgA, while preserving the length of the loop (Fig.2A). The first construct incorporated this mutation into full-length SC (mSC-D1-CDR3mut) and the second incorporated it into an expression construct encoding only SC-D1 (mD1-CDR3mut). We expressed and purified variants along with unmutated controls, mSC-wt and mD1-wt, and tested their binding to dFcα-wt (Fig.2 and Fig.S2).

The mSC-D1-CDR3mut SPR sensorgrams revealed markedly lower response units (RU) compared to concentration-matched mSC-wt responses, with an overall maximal response (R_{max}) of mSC-D1-CDR3mut that was 9.87% of the mSC-wt R_{max} for the 0.5μM analyte sample (Fig.2B and Fig.S2A). Furthermore, overlay of normalized responses revealed more rapid association and dissociation for mSC-D1-CDR3mut compared to mSC-wt (Fig.2B). Similarly, the mD1-CDR3mut SPR sensograms revealed markedly lower response, ~9.6% of the concentration-matched mD1-wt response (Fig.2C, Fig.S2B) and an overall reduction in affinity from K_D=0.374μM for mD1-wt to K_D=1.199μM for mD1-CDR3mut (Fig.2C). In contrast to comparisons between mSC-D1-CDR3mut and concentration matched mSC-wt, the responses revealed similar association and dissociation phases for mD1-CDR3mut and mD1-wt. Taken together, these data indicate that mutations in D1-CDR3 reduce D1 binding affinity in the context of SC and isolated D1, but alter the binding kinetics only in the context of SC. These results point toward a central role for D1-CDR3 in promoting D1-dIgA interactions, but also a role in promoting changes in other SC domains upon dIgA binding, as evidenced by differences in binding kinetics for mSC-wt and mSC-D1-CDR3mut. These differences may result from conformational changes that expose more of D1 for binding and/or the binding of other SC domains.

The formation of the SC D1-D3 clamp is crucial for stabilization of SC-binding.

In unliganded human SC, D1 forms extensive interfaces with SC domains D2, D4 and D5, yet shares minimal contacts with D3 (no contacts in mouse SC, one contact in human SC)(8). However, in the SIgA structure, D1 contacts only D3 through an extensive hydrophobic interface described as a "D1-D3 clamp" (Fig.3B)(11),(12),(13). The hydrophobic interface involves D1 residues F5, P7, Y23, P25 and D3 residues L226, Y228, F326. Notably, D3 L226 and Y228 contact SC D4 in the unliganded SC structure, implying that D3 swaps interfaces between D4 (unliganded) and D1 (liganded) during SC binding in what we hypothesize is a dominant step in SC's conformational change associated with SIgA formation (Fig.3A and Fig.S1D, S1E).

To investigate the significance of the D1-D3 clamp, we generated three mutated SC variants to minimize the hydrophobic interface: (1) mSC-D1mut, which includes D1 F5A, Y23A, and P25A mutations, (2) mSC-D3mut, which includes D3 L226A, Y228A, F326A mutations, and (3) mSC-D1-D3mut, which includes D1 F5A, Y23A, P25A and D3 L226A, Y228A, F326A mutations (Fig.3B) and tested their binding to mouse dFcα-wt. Compared to mSC-wt, SPR sensorgrams revealed lower RUs and lower R_{max} for the binding of concentration-matched mSC-D1mut (15.7% mSC-wt R_{max}), mSC-D3mut (25.3% mSC-wt R_{max}), and mSC-D1-D3mut (20.7% mSC-wt R_{max}) (Fig.3C and Fig.S3), indicative of weaker binding. Qualitatively, the reduction in RU appeared similar for all three mutant variants, with R_{max} values being between 15-26% of mSC-wt for 0.5μM analytes; the combination of mutations in both D1 and D3 did not appear to produce an additive effect, indicating that disruption of interface residues in either D1 or D3 was sufficient to disrupt the interface.

Overlay of concentration matched, normalized responses for mSC-wt and mSC-mutant variants also revealed differences in the binding kinetics, with mSC-D1mut exhibiting faster association and faster dissociation phases compared to mSC-wt (Fig.3C). Responses of mSC-D1mut were similar to those of mD1-wt (Fig.3C, Fig.3D), which may indicate that only D1 (and not D4 and D5) is binding. However, mSC-D3mut and mSC-D1-D3mut exhibited association phases similar to mSC-wt, but faster dissociation phases compared to mSC-wt. Compared to mSC-D1mut, mSC-D3mut and mSC-D1-D3mut exhibited

157   moderate differences in the dissociation phase compared to mSC-D1 mut, suggesting that mutation of the

158   D3 interface may impact binding differently than mutation of the D1 interface (Fig.3C and Fig.S3).

Figure-3.

Figure-3: Role of SC-D1-D3 Clamp for the stabilization of SC and dIgA binding. (A) Detailed view of the unliganded D3-D4 interface from the homology model of mSC and the D1-D3 interface from the SIgA cryo-EM structure (Fig.1, orange asterisk) (PDB code 7JG2). In (A) and (B) structures are shown as a cartoon representation with interacting residues, shown as sticks and labeled. SC D1 and D3 residues that were mutated are underlined, and the unmutated and mutated sequences are shown on the right. (C) SPR sensorgrams showing the response (top) and normalized response (bottom) of 0.5uM mSC-wt (black) mSC-D1mut (red), mSC-D3mut (blue) and mSC-D1-D3mut (green) binding to dFcα-wt. (D) SPR sensorgrams showing the response of 0.5uM of mD1-wt (black) and mD1-mut binding to dFcα-wt (top) and the concentration dependent binding and steady state fit (bottom). The calculated K_D values for each analyte are indicated to the right of the name.

To explore the possibility that mutation of D1 F5A, Y23A, P25A influenced D1 binding

independent of the other SC domains we generated mD1-mut, which contained D1 F5A, Y23A and P25A

mutations, and compared its binding affinity to mD1-wt (Fig.3D and Fig.S3). The mD1-mut binding

affinity, K_D=0.445μM, was comparable to the mD1-wt binding affinity, K_D=0.374μM, indicating that D1

interactions with dIgA were not disrupted. Thus, in the context of full-length SC, the mutations F5A, Y23A

and P25A appear to reduce binding to dIgA indirectly, presumably through disruption of the D1-D3 clamp.

Taken together, results imply that D1-D3 interface formation contributes to stable SIgA assembly but is not

necessary for SC (or D1) binding. As noted, the kinetics of the three D1-D3 interface mutants closely

resemble that of isolated SC D1 (Fig.3 and Fig.S3) and thus it is plausible that disruption of the D1-D3

clamp prevents subsequent stages of SC binding (e.g., part of the conformation change and D4-D5 binding

to dIgA) resulting in only D1-dIgA contacts. Alternatively, D1-D3 clamp disruption may markedly decrease

the efficiency of SC D4-D5 binding and/or the stability of the resulting SIgA.

JC C-terminal residues mediate pIgR binding to dIgA.

The structures of human and mouse SIgA reveal similar conformations for SC, JC and similar

contacts between chains. Among these interactions, SC-D1 contacts with JC are conserved, and contribute

to 63% of the total SC- JC interface. Of the three SC-D1 CDRs, JC interfaces primarily with SC-D1-CDR1,

forming hydrogen bonds and ionic interactions between JC residues R106, D107, D132, Y135 and D137

and SC-D1-CDR1 residues S28, D30, R31, H32 and T33 (Fig.4A). JC Y135 and D137 are part of the

conserved C-terminal *YPD* motif that follows JC wing-2; JC-YPD are disordered in the dIgA structure(11)

and therefore we hypothesized that these residues play a dominant role in pIgR binding, rather than dIgA

assembly. Furthermore, in unliganded SC, D1-CDR1 R31 and H32 form contacts with D4-D5(Fig.S1C)(8)

and thus, JC YPD may function to displace the D1-D4D5 interface and/or promote D1-JC binding while

freeing D4-D5 to bind the Fc_{CD}. To investigate these possibilities, we created ten JC variants with mutations

or truncations in YPD.

Figure-4.

Figure-4: Role of the JC YPD motif in binding mSC-wt or mD1-wt. (A) The interaction of JC with SC-D1, and specifically D1-CDR1 shown as red loop with larger radius. The interacting residues are shown as sticks at this interface. Y135, P136 and D137 are underlined in JC, indicating that these residues were subjected to mutations, such as alanine substitutions or truncations. (B) The SEC chromatograms of the indicated dFcα-wt and dFcα mutant variants resulting from 25ml co-transfections and following Nickel-NTA purification; fractions corresponding to dimeric forms were used for SPR experiments. (C) SPR sensorgrams of 0.5uM mD1-wt binding to the indicated dFcαs with JC- YPD alanine mutations or truncations. (D) The steady state affinity analysis of mD1-wt binding to the indicated wt or mutant dFcα variants. The respective calculated K_D values are listed in the adjacent table. (E) SPR sensorgrams of 0.5uM mSC-wt binding to different dFcα with JC- YPD alanine mutations or truncations. (F) SPR sensorgrams showing the normalized response of 0.5uM mSC-wt analytes binding to the indicated dFcα variants.

Initially we tested the effect of YPD mutations or truncations on dFcα assembly by co-expressing

each JC variant with Fcα and found that all could be incorporated into dFcα, indicating that mutations did

not inhibit dFcα assembly. In most cases, the relative percentage of dFcα and monomeric Fcα, which is

found in lower abundance in all dFcα preparations, was similar for preparations containing unmutated or

mutated JC. However, truncation of YPD resulted in higher percentages of monomeric Fcα and lower dFcα

yield, suggesting that the efficiency of JC incorporation, and therefore dIgA assembly, may rely in-part on

the presence of JC C-terminal residues (Fig.4B). Accordingly, we used mutated dFcα variants as

immobilized ligands in mD1-wt and mSC-wt SPR binding assays.

Initially, we tested binding of mD1-wt to dFcα-wt and dFcα-mutant variants having single (dFcα-

APD, dFcα-YAD, dFcα-YPA), double (dFcα-AAD, dFcα-ADA, dFcα-YAA) or triple (dFcα-AAA) alanine

substitutions in JC YPD and calculated the binding affinity for each variant. Results revealed binding

affinities of K_D=0.374μM, K_D = 1.104uM, K_D = 0.329uM and K_D = 0.810uM for dFcα-wt and single alanine

variants dFcα-APD, dFcα-YAD, and dFcα-YPA, respectively (Fig.4C, Fig.4D). Binding affinities for

double mutants dFcα-AAD, dFcα-APA, and dFcα-YAA were K_D =1.728uM, K_D =3.692uM and K_D =1.191

uM, respectively (Fig.4C, Fig.4D). Collectively, these results demonstrate a marked reduction in binding

affinity when either Y135 or D137 is mutated and a further reduction in binding affinity when both Y135

and D137 are mutated to alanine. The P136A mutation resulted in a comparatively minor changes in binding

on its own; however, did further reduce binding when combined with Y135A or D137A mutations. The

binding affinity for the triple mutant dFcα-AAA was K_D = 4.246uM, indicating the substitution of all YPD

residues has the most pronounced effect on binding (Fig.4C, Fig.4D).

We also tested the mSC-wt binding to all dFcα YPD-alanine substitutions. Although resulting SPR

responses do not fit a single-state kinetic model or reach equilibrium during the experiment, differences in

dissociation phases for concentration matched, normalized responses were apparent when comparing mSC-

wt binding to dFcα-wt and dFcα YPD-alanine substitutions (Fig.4E, Fig.4F). Specifically, we observed that

all mutations except P136A altered dissociation. Consistent with experiments that tested D1 binding, single

mutation Y135A exhibited the most pronounced effect, followed closely by D137A; the combination of

Y135A and D137A further altered the dissociation phase (more rapid dissociation) and the triple mutant

containing Y135A, P136A, and D137A exhibited the most significant change in dissociation compared to

unmutated control, dFcα-wt. Qualitatively, we also observed subtle differences in association and overall

reduction in R_{max} values for mSC-wt binding to the majority of dFcα YPD-alanine substitutions (Fig.4E and

Fig.S4).

Taken together YPD-alanine substitutions reveal a central role for JC-YPD in mediating D1 binding

to dIgA. Y135 and D137 appear to play prominent and unique roles since individual mutations in these

residues reduced binding affinity more than mutation of P136. Both Y135 and D137 appear to contribute

uniquely since dFcα YPD-alanine substitutions that combined them exhibited further reductions in binding.

While a more minor role for P136 was observed, its mutation to alanine did reduce binding affinity, most

notably when combined with the Y135A and/or D137A mutations. Notably, none of the dFcα YPD-alanine

substitutions completely abrogated mD1-Wt or mSC-wt binding indicating that either YPD main-chain

interactions and/ or other JC-SC interfaces are sufficient for SIgA formation.

To investigate the role of JC length and YPD main chain atoms on interactions with SC, we tested

SC binding to JC deletion mutants, in which D137 (dFcα-YPX), P136 and D137 (dFcα-YXX), or Y135,

P136 and D137(dFcα-XXX) were deleted. Initially, we tested mD1-wt binding to dFcα-wt and dFcα YPD

deletions, dFcα-YPX, dFcα-YXX and dFcα-XXX and calculated associated binding affinities. Results

revealed binding affinities of K_D= 0.737uM, K_D= 1.434uM and K_D= 0.430uM for dFcα-YPX, dFcα-YXX

and dFcα-XXX, respectively (Fig.4C, Fig.4D), indicating a reduction in binding upon the deletion of one

or more JC C-terminal residues compared to dFcα-wt K_D=0.374μM. Reduction in binding affinity was not

correlated with the number of deleted residues indicating that a change in length has some unknown impact

on D1 binding. We also tested the mSC-wt binding to all dFcα YPD-deletion variants and similar to dFcα

YPD alanine substitutions, observed differences in the dissociation phases for concentration matched,

normalized responses when comparing mSC-wt binding to dFcα-wt, dFcα-YPX, dFcα-YXX and dFcα-

XXX (Fig.4E, Fig.4F). While we observed more rapid dissociation for mSC-wt binding to dFcα-YPX and

dFcα-YXX (compared to dFcα-wt), binding to dFcα-XXX exhibited dissociation more similar to dFcα-wt,

albeit still altered. Taken together, these results further support a role for JC YPD in SIgA formation and

suggest that both sidechain identity and the presence of mainchain atoms support the interaction.

DISCUSSION

The pIgR binding to dIgA is a critical step in SIgA formation, its transcytosis to mucosal secretions,

and downstream effector functions(23),(24). Prior work has unveiled an intricate, multi-step process, in

which domains D1-D5 undergo conformational rearrangements that uncover D1 residues, ultimately

allowing them to bind all four HCs and the JC, while also repositioning D4 and D5 to bind a single HC(8),

(11),(12),(13),(16),(17),(18). Here we characterized the contributions of specific residues in the D1 CDR3,

the D1-D3 clamp and the JC C-terminal YPD residues, which provide new details on how SIgA assembles.

In SIgA, D1-CDR3 residues account for just 34.3% of D1's total buried surface, yet in our studies

mutation of these residues reduced D1 binding affinity by ~3 times and altered SC association and

dissociation phases. Prior studies demonstrated that D1 is necessary and sufficient for dIgA

binding(16),(19), and that substituting the highly conserved D1 CDR loops with alternative sequences (e.g.

from other SC domains) resulted in limited dIgA binding (18). D1 CDR1, which contains a conserved

helical turn has long been thought to play a pivotal role in the process(15),(18); however, our data and

analysis also point toward a central role for D1-CDR3 in SIgA assembly. As noted, D1-CDR3 residues,

unlike D1-CDR1 and D1-CDR2 residues, are fully exposed in the unliganded SC structure(8) and thus, our

results are consistent with the model in which exposed D1-CDR3 residues are able to initiate binding by

contacting Tps and JC residues, promoting D1-CDR1 and D1-CDR2 contact with dIgA, and likely initiating

the conformational change. CDR3 residues also appear to contribute considerable stability to SC

interactions with pIgA in the fully assembled structure and interface with at least four different chains. In

the mouse SIgA structure(11), SC-D1-CDR3 residues participate in a network of hydrogen bonds and ionic

interactions with HC-B, TpC, and TpD as well as putative Van-der Waals interactions with the JC.

Furthermore, these Tps (specifically TpD) and JC residues appear to be flexible (disordered) in dIgA and

we speculate that binding to CDR3 helps stabilize their positions and following the conformational change

and promote additional interactions with D1 (e.g. CDR1 and CDR2).

D1-CDR3 may also promote binding to higher order pIgA and pIgM. Our own inspection of the

human dimeric (PDB: 6UE7 and 6LX3)(12),(13), tetrameric (PDB: 6UE8 and 6UE9), and pentameric

(PDB:6UEA) SIgA core structures(12), reveals D1-CDR3 contacting one Tp (the others are not resolved)

in dimeric SIgA and contacting two Tps in tetrameric and pentameric SIgA; in pentameric SIgA, a third Tp

contacts D1 outside the CDR regions (11),(12),(13). These observations are consistent with earlier reports

that SC bound tetrameric IgA2m2 with better affinity than dimeric IgA1 and dimeric IgA2m1(5) and

suggest D1-CDR3 may support high affinity binding to higher order pIgA. Our inspection of the related

SIgM core structure (PDB: 6KXS and 7K0C)(25),(26) indicates that SC D1 forms additional contacts with

the Tps in pentameric SIgM (similar to pentameric SIgA and in contrast to dimeric SIgA) and thus evolution

of D1 may have been driven by interactions with higher order polymers of IgM and IgA and their respective

Tps.

The structures of SIgA (and SIgM) revealed that the pIgR's ligand-induced conformational change

is associated with the formation of the D1-D3 clamp and data presented here revealed that disruption of the

D1-D3 clamp markedly reduces dIgA binding. Structurally, the unliganded SC D3-D4 interface must be

disrupted for D1-D3 interface to form, which implies that either the D1-D3 interface formation frees D4-

D5 for binding to dIgA, or that the binding of D4-D5 to dIgA frees D3 to bind D1. Indeed, SC variants with

mutated clamp residues exhibited SPR binding profiles that are similarly to that of D1 only, which may

indicate that in the absence of the clamp, D4 and D5 are not able to bind efficiently. While this could result

from failure of the clamp to form and/or from the failure of the unliganded SC D3-D4 interface to break; it

is notable that among D1-D3 interface mutants, mSC-D1mut is the weakest binder. The mSC-D1mut lacks

the L226A and Y228A mutations found in mSC-D3mut and mSC-D1-D3mut, which are expected to disrupt

the D3-D4 interface in unliganded SC. Thus, it appears plausible that D1-D3 clamp formation frees D4 and

D5 and positions them optimally to bind dIgA. Regardless, it is clear is that both disruption of unliganded

SC D3-D4 contacts and formation of the liganded D1-D3 clamp and are crucial for the formation of a stable

SIgA complex.

The JC has long been known to be necessary for dIgA assembly in mammals and for pIgR binding

to dIgA; yet gaps in our understanding of how it mediates both processes remain (27). The mouse dIgA

structure revealed numerous contacts between JC and two copies of mIgA; however, the conserved C-

terminal JC YPD motif was disordered (11), hinting at a limited role in dIgA assembly. Our data

demonstrate that dIgA assembly can tolerate mutations in JC YPD, and thus, its identity is not necessary

for dIgA assembly. A limited role for the JC C-terminus in dIgA is consistent with earlier studies, in which

deletion of the 25 or more residues from the C-terminus did not completely abrogate polymeric IgA

assembly (21). However, our data indicates the presence of three C-terminal residues in the YPD location

may contribute significantly to the dIgA assembly process since deletion of all three residues resulted in

markedly lower levels of dIgA expression. Additionally, the mouse SIgA structure revealed an ordered YPD

motif contacting SC-D1 (11) and we find that mutations of these residues, particularly Y and D, markedly

impact SC binding. Yet, mutation or deletion of JC YPD doesn't completely inhibit SC binding *in vitro*.

Thus, other contacts between the JC and SC and/or dIgA appear to be sufficient for binding in our

experiments. While these contacts may include other JC-SC interfaces, it may also be that SC-SC and/or

SC-HC contacts help compensate for JC mutations and that those contacts are dependent on the overall bent

dIgA conformation, conferred by largely by the JC (but not directly by the YPD). Regardless, our data

support the notion that direct contacts with the JC C-terminal YPD motif are an inherent part of the pIgR-

dIgA binding mechanism.

Notably, none of our SC or JC mutant variants completely abrogated SC-dIgA binding. This is

valuable for determining the relatively contributions of mutated residues; however, we also speculate that

the reductions in binding affinity and apparent changes in kinetics that we observe could translate to severe

disruption of pIgR-dependent transcytosis and or reduced SIgA stability in the mucosa. For example,

mutations in the D1-D3 clamps and/or mutations disrupting interfaces may increase the exposure and/or

flexibility of SC and/or dIgA motifs making them more susceptible to proteolysis. They may also limit

subsequent stages of pIgR binding to dIgA. For example, mD5 C470 (C468 in human) can form a disulfide

bond with HC C306 (C311 in human IgA) that stabilizes SIgA after SC D5 has bound dIgA

(11),(12),(13),(28),(29). As noted in our results, mutations in SC D1 and/or D3 resulted in kinetic profiles

similar to D1 only variants, and thus we speculate that these mutations may limit D4/D5 binding and

subsequent disulfide bond formation; however, in our SPR experiments (and previously reported binding

studies (8),(22)) SC C470 and C504 were mutated to alanine to prevent irreversible binding to dIgA

surfaces.

MATERIALS AND METHODS

Construct design and cloning

To make the mouse Fcα expression construct, the gene segment encoding the *mus musculus* IgA

HC constant region (Uniprot P01878) CH2 and CH3 domains was codon optimized, fused to an N-terminal

TPA signal sequence (residues MDAMKRGLCCVLLLCGAVFVSPAGA) and a C-terminal hexahistidine

affinity tag and cloned into expression vector pD2610v1 (ATUM) using Electra cloning (ATUM),

analogous to previously reported full-length IgA HC expression constructs(11). Constructs encoding the

mus musculus JC (Uniprot P01592; native signal peptide) and *mus musculus* pIgR ectodomain residues 1-

567 (Uniprot P01592; native signal peptide), were used as previously reported and served as templates for

cloning truncated and/or mutant variants(11). SC and JC mutations were introduced into coding sequence

fragments using overlap-extension PCR(30) with primers encoding the desired mutations (Integrated DNA

Technologies; IDT) and Electra cloning sites; resulting products were cloned into expression vector

pD2610v1 (ATUM) using Electra cloning (ATUM). SC-D5 cysteine mutations (C470A and C504A) were

introduced into all SC expression constructs to prevent the formation of disulfide bonds with dIgA, as

previously described(8). Constructs encoding the SC mD1-wt (residues 1-117) and mD1-mutant variants

were generated by PCR and overlap extension PCR as described for other SC variants; coding sequence

retained the native signal peptide and a C-terminal hexahistidine tag and were cloned into pD2610v1 using

Electra cloning (ATUM).

Protein expression and purification

All constructs were transiently transfected (SC and D1 variants) or co-transfected (Fcα and JC) into Expi293F (Gibco: A14527) cells using the ExpiFectamine 293 Transfection kit (Gibco: A14525) and following previously described methods (11). Four to five days after transfection, supernatants were harvested and filtered through 0.22μm PES bottletop filters (Millipore Sigma). Proteins and complexes were purified from the cell supernatants using Nickel-NTA affinity resin (Qiagen: 30210), followed by Superose 6 (Cytiva) or Superdex 200 (Cytiva) size exclusion chromatography (SEC) on an AKTA FPLC (Cytiva). During and after SEC, purified molecules were maintained in TRIS-buffered saline (20mM Tris+ 150mM NaCl, pH=7.4). The SEC-eluted protein peaks corresponding to the correct retention volume based on the molecular weight standards (BioRad: 1511901) were collected, concentrated using Amicon Ultra centrifugal filters (Millipore Sigma) to stock concentrations between 2-4mg/ml for SC, 1.5-2mg/ml for D1, and 0.4-0.7mg/ml for dFcα constructs, which were then filtered using 0.22μm spin filters (Millipore Sigma) and utilized for SPR experiments.

SPR experiments and data analysis.

SPR experiments were carried out using a Sierra SPR-32 Pro (Bruker) operating at room temperature. Solutions containing 0.25uM dFcα-wt or dFcα mutant variant ligands in sodium acetate buffer (pH=4), were immobilized on lanes B1-8 and D1-8 of a High-Capacity Amine (HCA) 32-spot sensor (Bruker) using a standard amine coupling kit (Bruker). Lanes A and C were mock coupled using sodium acetate buffer pH=4 and served as references. The binding of soluble analytes, including mSC-wt, mSC-mutants, mD1-wt and mD1-mutants was tested using a two-fold (mSC and variants) or four-fold (mD1 and variants) dilution series starting at 2uM in HBS-EP+ running buffer (0.01 M HEPES, 0.15 M NaCl, 3 mM EDTA, 0.005% v/v Surfactant P20, pH 7.4). Analyte responses were measured on all spots and responses on lanes A and C (internal references) were subtracted to account for any non-specific binding. Kinetic experiments were carried out using a 25ul/min flow rate, 120s contact time, and 300s dissociation time.

After every cycle the surface was regenerated using 2.5M MgCl2 three times, followed by a wash step with

the running buffer (HBS-EP+) to achieve the best baseline for the next cycle.

Resulting SPR data were analyzed using the Bruker Sierra Analyzer software version R4. Briefly,

all data were reference subtracted and buffer-subtracted and K_D values were calculated for the D1-only

constructs using "steady state affinity" analysis program and 1:1 Langmuir + offset fitting model. Raw

sensor-grams (Response Units versus time) for all data and the steady state affinity data for mD1 variants

(Response Units vs concentration), were exported and re-plotted using OriginPro 2020 graphing and

analysis software (OriginLab: https://www.originlab.com/). Normalization of the respective sensorgrams

for concentration matched samples were carried out in OriginPro 2020, by dividing RU by R_{max} of the

respective analyte.   Figures were assembled and in Photoshop (Adobe) or Inkscape (https://inkscape.org/).

AUTHOR CONTRIBUTIONS

The study was conceived by B.M.S and S.K.B.; experiments were conducted by S.K.B; Authors contributed

to data analysis and manuscript writing.

ACKNOWLEDGEMENTS

We thank Paul Ritter and Hniang Khamh for help with SPR training and troubleshooting, Sarah Leonard

for technical assistance, and members of the Stadtmueller Lab for insightful discussions regarding this

work. This work was supported by NIH grant R01AI165570 to B.M.S., University of Illinois start-up

funding to B.M.S., and the Lowell P. Hager Fellowship to S.K.B.

REFERENCES

1. Corthesy, B. 2013. Multi-Faceted Functions of Secretory IgA at Mucosal Surfaces. *Frontiers in Immunology* 4.

2. Mantis, N. J., N. Rol, and B. Corthésy. 2011. Secretory IgA's complex roles in immunity and mucosal homeostasis in the gut. *Mucosal Immunol* 4: 603–611.

3. de Sousa-Pereira, P., and J. M. Woof. 2019. IgA: Structure, Function, and Developability. *Antibodies* 8: 57.

4. Zikan, J., J. Mestecky, R. Kulhavy, and J. C. Bennett. 1986. The stoichiometry of J chain in human secretory dimeric IgA. *Molecular Immunology* 23: 541–544.

5. Lombana, T. N., S. Rajan, J. A. Zorn, D. Mandikian, E. C. Chen, A. Estevez, V. Yip, D. D. Bravo, W. Phung, F. Farahi, S. Viajar, S. Lee, A. Gill, W. Sandoval, J. Wang, C. Ciferri, C. A. Boswell, M. L. Matsumoto, and C. Spiess. 2019. Production, characterization, and in vivo half-life extension of polymeric IgA molecules in mice. *mAbs* 11: 1122–1138.

6. Kaetzel, C. S., J. K. Robinson, K. R. Chintalacharuvu, J. P. Vaerman, and M. E. Lamm. 1991. The polymeric immunoglobulin receptor (secretory component) mediates transport of immune complexes across epithelial cells: a local defense function for IgA. *Proc Natl Acad Sci U S A* 88: 8796–8800.

7. Brandtzaeg, P. 2013. Secretory IgA: Designed for Anti-Microbial Defense. *Frontiers in Immunology* 4.

8. Stadtmueller, B. M., K. E. Huey-Tubman, C. J. López, Z. Yang, W. L. Hubbell, and P. J. Bjorkman. 2016. The structure and dynamics of secretory component and its interactions with polymeric immunoglobulins. *eLife* 5: e10640.

9. Asano, M., N. Takenouchi-Ohkubo, N. Matsumoto, Y. Ogura, H. Nomura, H. Suguro, and I. Moro. 2004. Multiple cleavage sites for polymeric immunoglobulin receptor. *Immunology* 112: 583–589.

10. Woof, J. M., and M. W. Russell. 2011. Structure and function relationships in IgA. *Mucosal Immunol* 4: 590–597.

11. Kumar Bharathkar, S., B. W. Parker, A. G. Malyutin, N. Haloi, K. E. Huey-Tubman, E. Tajkhorshid, and B. M. Stadtmueller. 2020. The structures of secretory and dimeric immunoglobulin A. *eLife* 9: e56098.

12. Kumar, N., C. P. Arthur, C. Ciferri, and M. L. Matsumoto. 2020. Structure of the secretory immunoglobulin A core. *Science* 367: 1008–1014.

13. Wang, Y., G. Wang, Y. Li, Q. Zhu, H. Shen, N. Gao, and J. Xiao. 2020. Structural insights into secretory immunoglobulin A and its interaction with a pneumococcal adhesin. *Cell Res* 30: 602–609.

14. Brunke, C., S. Lohse, S. Derer, M. Peipp, P. Boross, C. Kellner, T. Beyer, M. Dechant, L. Royle, L. P. Liew, J. H. Leusen, and T. Valerius. 2013. Effect of a tail piece cysteine deletion on biochemical and functional properties of an epidermal growth factor receptor-directed IgA2 m(1) antibody. *mAbs* 5: 936–945.

15. Hamburger, A. E., A. P. West, and P. J. Bjorkman. 2004. Crystal Structure of a Polymeric Immunoglobulin Binding Fragment of the Human Polymeric Immunoglobulin Receptor. *Structure* 12: 1925–1935.

16. Frutiger, S., G. J. Hughes, W. C. Hanly, M. Kingzette, and J. C. Jaton. 1986. The amino-terminal domain of rabbit secretory component is responsible for noncovalent binding to immunoglobulin A dimers. *Journal of Biological Chemistry* 261: 16673–16681.

17. Braathen, R., V. S. Hohman, P. Brandtzaeg, and F.-E. Johansen. 2007. Secretory Antibody Formation: Conserved Binding Interactions between J Chain and Polymeric Ig Receptor from Humans and Amphibians1. *The Journal of Immunology* 178: 1589–1597.

18. Coyne, R. S., M. Siebrecht, M. C. Peitsch, and J. E. Casanova. 1994. Mutational analysis of polymeric immunoglobulin receptor/ligand interactions. Evidence for the involvement of multiple complementarity determining region (CDR)-like loops in receptor domain I. *Journal of Biological Chemistry* 269: 31620–31625.

19. Norderhaug, I. N., F.-E. Johansen, P. Krajči, and P. Brandtzaeg. 1999. Domain deletions in the human
polymeric Ig receptor disclose differences between its dimeric IgA and pentameric IgM interaction.
European Journal of Immunology 29: 3401–3409.
20. Vaerman, J.-P., A. E. Langendries, D. A. Giffroy, C. S. Kaetzel, C. M. Tamer Fiani, I. Moro, P.
Brandtzaeg, and K. Kobayashi. 1998. Antibody against the human J chain inhibits polymeric Ig receptor-
mediated biliary and epithelial transport of human polymeric IgA. *European Journal of Immunology* 28:
171–182.
21. Johansen, F.-E., R. Braathen, and P. Brandtzaeg. 2001. The J Chain Is Essential for Polymeric Ig
Receptor-Mediated Epithelial Transport of IgA1. *The Journal of Immunology* 167: 5185–5192.
22. Stadtmueller, B. M., Z. Yang, K. E. Huey-Tubman, H. Roberts-Mataric, W. L. Hubbell, and P. J.
Bjorkman. 2016. Biophysical and biochemical characterization of avian secretory component provides
structural insights into the evolution of the polymeric Ig receptor. *J Immunol* 197: 1408–1414.
23. Johansen, F.-E., and C. S. Kaetzel. 2011. Regulation of the polymeric immunoglobulin receptor and
IgA transport: new advances in environmental factors that stimulate pIgR expression and its role in
mucosal immunity. *Mucosal Immunology* 4: 598–602.
24. Wei, H., and J.-Y. Wang. 2021. Role of Polymeric Immunoglobulin Receptor in IgA and IgM
Transcytosis. *Int J Mol Sci* 22: 2284.
25. Kumar, N., C. P. Arthur, C. Ciferri, and M. L. Matsumoto. 2021. Structure of the human secretory
immunoglobulin M core. *Structure* 29: 564-571.e3.
26. Li, Y., G. Wang, N. Li, Y. Wang, Q. Zhu, H. Chu, W. Wu, Y. Tan, F. Yu, X.-D. Su, N. Gao, and J. Xiao.
2020. Structural insights into immunoglobulin M. *Science* 367: 1014–1017.
27. Hockenberry, A., E. Slack, and B. M. Stadtmueller. 2023. License to Clump: Secretory IgA Structure–
Function Relationships Across Scales. *Annual Review of Microbiology* 77: 645–668.
28. Chintalacharuvu, K. R., A. S. Tavill, L. N. Louis, J. P. Vaerman, M. E. Lamm, and C. S. Kaetzel.
1994. Disulfide bond formation between dimeric immunoglobulin A and the polymeric immunoglobulin
receptor during hepatic transcytosis. *Hepatology* 19: 162–173.
29. Underdown, B. J., J. De Rose, and A. Plaut. 1977. Disulfide bonding of secretory component to a
single monomer subunit in human secretory IgA. *J Immunol* 118: 1816–1821.
30. Ho, S. N., H. D. Hunt, R. M. Horton, J. K. Pullen, and L. R. Pease. 1989. Site-directed mutagenesis
by overlap extension using the polymerase chain reaction. *Gene* 77: 51–59.
463

Supplementary Information for:

Structural and biochemical requirements for secretory component interactions with

dimeric Immunoglobulin A

Sonya Kumar Bharathkar and Beth M. Stadtmueller[1,2,3*]

[1]Department of Biochemistry, University of Illinois Urbana-Champaign, Urbana, Illinois 61801 USA

[2]Department of Biomedical and Translational Sciences, Carle Illinois College of Medicine, University of Illinois Urbana-Champaign, Urbana, Illinois 61801 USA

[3]Carle R. Woese Institute of Genomic Biology

*Address correspondence to bethms@illinois.edu–

Supplemental Figure-1: Different interfaces of SC-domains in unliganded and liganded structures. (A) Homology model of unliganded mouse SC based on the unliganded human SC structure (PDB: 5D4K) and the mouse SIgA Cryo-EM structure (PDB: 7JG2). The SC is shown as a cartoon, with its five domains (D1-D5) colored different shades of cyan. In SIgA, dIgA components are shown as a molecular surface and colored as Fig. 1. The boxed regions are enlarged in B-G and detail residues engaged in molecular interactions for the (B) liganded SC D1-D3 clamp in SIgA, (C) unliganded SC D1 and D4-D5 interface, (D) unliganded D3 and D4 interface, (E) liganded D3 and D4 interface in SIgA, (F) liganded D1 and dIgA interface (G) the liganded D4-D5 and dIgA interface. In all panels, interfacing residues are shown as sticks. The D1-CDRs are shown as thicker loops denoted as 1, 2 and 3 corresponding to CDR1, CDR2 and CDR3 respectively, and they are colored brick red.

Supplemental Figure-2: mSC-wt and mSC-D1-CDR3mut binding to dFcα-wt at various concentrations (A) SPR sensorgrams showing the responses for the entire dilution series of mSC-wt and mSC-D1-CDR3mut binding to dFcα-wt, shown with the same scale. (B) SPR sensorgrams showing the responses for the entire dilution series of mD1-wt and mD1-CDR3mut binding to dFcα-wt. The responses for each analyte dilution are colored according to the keys (right), shown with the same scale.

Supplemental Figure-3: Role of D1-D3 in binding stabilization and conformation change. (A) SPR sensorgrams showing the responses for the entire dilution series of mSC-wt, mSC-D1mut, mSC-D3mut and mSC-D1-D3mut, binding to dFcα-wt (2-fold dilution series starting from 2uM). All shown in same scale for comparison. (B) SPR sensorgrams showing the responses for the entire dilution series of mD1-wt, and mD1mut, binding to dFcα-wt (4-fold dilution series from 2uM). The responses for each analyte dilution are colored according to the keys (right).

Supplemental Figure-4.

Supplemental Figure-4: (A) SPR sensorgrams showing the responses for the entire dilution series of mSC-wt binding to the indicated dFcα variants (shown on the same scales). (B) SPR sensorgrams showing the responses for the entire dilution series of mD1-wt binding to the indicated dFcα variants (shown on the same scales). The responses for each analyte dilution are colored according to the keys (right).