1

Examining models of modern human origins through the analysis of 43 fully sequenced human Y chromosomes

Shi Huang*

Center for Medical Genetics, Hunan Key Laboratory of Medical Genetics, School of Life Sciences, Central South University, 110 Xiangya Road, Changsha, Hunan 410078, P.R. China

*Corresponding author: Shi Huang (shihuang1@gmail.com)

Short title: Testing human origin models with fully sequenced Y chromosomes
Keywords: Y chromosomes, maximum genetic diversity theory, neutral theory, infinite site model, Recent Out of East Asia model, Recent Out of Africa model,

Abstract

Molecular studies have yielded two primary models for understanding the uniparental DNA phylogenetic trees of modern humans: the Recent Out of Africa (ROA) and the Recent Out of East Asia (ROE) models. These models differ in their underlying assumptions, particularly in relation to early stem haplotypes, even though they share many haplotype relationships. Leveraging the wealth of new genetic variants within the male-specific region unveiled through the comprehensive sequencing of 43 diverse human Y chromosomes, we here investigated the presence of shared variants among different haplotypes to determine which model better aligns with the genetic data. We were able to corroborate the existence of stem haplotypes specific to the ROE model, but not those exclusive to the ROA model. We found that A0b and A1a shared the most variants with each other, aligning with the A00A1a clade of the ROE model. Also, stem haplotypes specific to the ROE model showed the expected relationships, with A00A1a closest to B, AB closest to E, E closest to B, A, and C, ABDE closest to C. Our findings also revealed extensive variant sharing independent of common ancestry, consistent with the maximum genetic diversity theory underlying the ROE model but challenging the neutral theory behind the ROA model. These new genetic data lend robust support to the ROE model as the more accurate representation of modern human origins.

Introduction:

40
The origins of modern humans remain a subject of ongoing and intensive research.
Molecular analyses of contemporary human populations have generated two distinct
models to explain the evolutionary history of modern humans: the "Recent Out of Africa"
(ROA) model and the "Recent Out of East Asia" (ROE) model. These models propose
contrasting geographical locations as the point of origin for modern humans, with ROA
suggesting Africa and ROE suggesting East Asia. The ROA model gained prominence in
the 1980s, primarily based on mitochondrial DNA (mtDNA) studies (1, 2). On the other
hand, the ROE model, also initially suggested by mtDNA studies in the 1980s (1), had
been relatively overlooked until its recent independent rediscovery (3-6). Moreover, it
has been revealed that all non-African Y chromosome haplotypes can be traced back to
their origin in the southern part of East Asia (7).
The foundation of rooting uniparental DNAs in Africa rests on two fundamental
assumptions: the molecular clock and the neutral theory (1, 2, 8, 9). Within this neutral
framework, the "infinite site assumption" holds particular significance. This assumption
posits that mutations occur only once in the course of evolutionary history, thereby
excluding the possibility of back mutations and convergent mutations. It serves as a
critical factor in deducing the derived alleles that are pivotal in shaping the phylogenetic
tree topology of the ROA model (8, 10). However, the widely recognized downfall of the
molecular clock hypothesis (11)—initially proposed as an *ad hoc* explanation for the
genetic equidistance phenomenon—has effectively undermined the neutral theory,
which was inspired by the molecular clock and supposed to explain or predict it (12).
Quantifying constraint in the human mitochondrial genome has shown that most variants
in mtDNA, including those in the non-coding region, are subject to selection (13). Direct
experimental tests of the predictions of the neutral theory have disproven it (14). Strong
evidence suggests that mutation saturation and natural selection are more prevalent
than initially believed (3, 15-20). In the Y chromosome phylogenetic tree of the ROA
model, numerous incompatible alleles have emerged, showcasing recurrent mutations
(21). Ancient DNA has revealed common occurrence of convergent mutations in the
human Y chromosome (22).
Genetic diversity patterns have continued to elude full explanation under the neutral
theory (23), prompting the emergence of a novel framework in molecular evolution
known as the "maximum genetic diversity theory" (17, 24-27). Central to this new theory
is the intuitive notion, initially suggested by Fisher nearly a century ago, that mutations of
a given size are less likely to be favorable in complex than simple organisms (28, 29).
The most recent supporting evidence for this concept lies in the observation that lower
species tend to harbor a greater number of common variants, which, however, are often
detrimental in humans (30). In accordance with this new theory, and bolstered by a
wealth of strong evidence, genetic diversities or distances are primarily observed to be
at optimal saturation levels, rather than continually scaling with time, as assumed by the
molecular clock and neutral theory. Upon mutation saturation, convergent evolution is
very common where independent lineages share variants due to independent mutations
and common selection.
The Y chromosome phylogenetic tree within the ROE model is constructed based
on this novel theoretical framework, setting it apart from the ROA model in three
significant ways (Fig. 1). Firstly, in ROE, haplotypes are defined by haplotype-specific
alleles, irrespective of whether they are derived or result from back mutations. Secondly,
the rooting in the ROE model is grounded in a straightforward premise: the original
haplotype should be the one most commonly shared among individuals, as mutations
leading to alternative haplotypes should be relatively rare occurrences (1). Finally, the
haplotype representing the ancestral lineage is expected to exhibit a closer genetic

relationship to ancient DNA samples from the time when modern humans initially
appeared, in contrast to other contemporary haplotypes (3-5).
Both models, ROA and ROE, rely on assumptions that carry inherent uncertainties,
necessitating independent testing to establish their validity. Human Y chromosomes
have until recently only been partially sequenced (53.8%) and the phylogenetic tree of Y
chromosome is built by using single nucleotide variants (SNVs, which are equivalent to
SNP, with SNP often referring to common variants with frequency >0.01 whereas rare
variants are sometimes referred as SNVs) located in the non-recombining Y region
(NRY) or the male specific Y region (MSY). These variants are located within a 10 MB
region amendable to short read sequencing (21). The human Y chromosome is 57 MB in
length with 95% being MSY and 5% being the pseudoautosomal regions (PARs). The
MSY region, unlike the PAR region, does not undergo recombination and is therefore
informative to genealogy. A groundbreaking development is the recent achievement of
complete sequencing for 43 human Y chromosomes, representing a wide array of major
human groups or haplotypes (31). This comprehensive sequencing has unveiled a
plethora of new SNVs and insertions or deletions (indels) located in the MSY region,
which now offer a valuable resource for an independent examination of the human origin
models of the paternal lineages.
In our study, we scrutinized these variants to ascertain which model finds stronger
support in the context of this new variant data. Our results indicate that numerous
variants are shared across different haplotypes, providing an informative pattern that
sheds light on the relationships among these haplotypes. Importantly, this analysis
validates the Y chromosome phylogenetic tree of the ROE model, offering compelling
evidence in its favor.
115
Methods and Materials:
Datasets
The data utilized in this study were sourced from Supplementary Tables S22, S24,
and S25, as published in the paper by Hallast et al. in 2023 (31). These tables provide
comprehensive listings of large and small size indels (Table S22 and S24) and SNVs
(Table S25), along with the occurrences of these variants across the 43 Y
chromosomes/samples. In our analysis, we combined the information from Table S22
and S24, which includes a total of 24,325 indels and 53,744 SNVs for our study. From
these, we picked out variants from the MSY region for our studies based on their
locations and Y-class designations as found in the source tables, including 16,679 indels
and 34,764 SNVs.
127
Calculation of the shared variant fraction between two haplotypes
For each haplotype/sample, we first compiled a list of SNVs or indels that are
present within that specific haplotype/sample. We then employed custom scripts
designed for this purpose to tally the count of shared genetic variants between the given
haplotype/sample and all the other haplotypes/samples among the 43 Y chromosomes
analyzed in our study. To quantify the extent of shared variants between two haplotypes,
we computed the fraction of shared variants. For example, when the haplotype A0b
(sample HG01890) was analyzed for the degree of sharing with each of the other 43
samples, the shared fraction was calculated by dividing the number of shared variants
by the total number of variants in the A0b sample (6,793 SNVs or 3521 indels). These
fraction values are listed in the A0b column of Table 2 and in Supplementary Tables S1
and S2. This approach allowed us to systematically assess the genetic relationships
between the various haplotypes within our dataset.
141

Calculation of the shared variant fraction between a stem haplotype and other
haplotypes
A stem haplotype is ancestral and not directly observed in the current populations
but is inferred to have existed in the past and is represented as the common ancestor of
several modern haplotypes. The shared variant fraction between a stem haplotype and
other haplotypes was calculated as follows, using the O stem haplotype as an example
to illustrate our method. 1) Identification of SNVs or indels defining the O stem haplotype:
SNVs or indels defining a stem haplotype, such as O, should be present in all terminal
branch haplotypes. In the case of the O stem haplotype, there were one O1b sample
and four O2a samples in the 43 Y chromosomes dataset. All SNVs or indels that were
present in all five O samples were retrieved and designated as the O stem set. 2)
Counting shared SNVs or indels: For a specific haplotype/sample, the number of shared
SNVs or indels between that haplotype and the O stem set was counted. 3) Calculation
of the shared fraction: The shared fraction for the O stem set was obtained by dividing
the number of shared variants by the total number of variants in the O stem set. 4)
Handling multiple samples: For haplotypes with multiple samples or terminal haplotypes
represented in the 43 Y chromosomes dataset, such as haplotype E, the shared fraction
between each individual E haplotype/sample and the O stem haplotype was
independently calculated. These shared fractions for all E haplotypes/samples were then
aggregated to calculate an average value. This average value was used to represent the
shared fraction between the E haplotype as a whole and the O stem haplotype. This
method allowed us to systematically assess and compare the shared genetic variants
between the O stem haplotype and other haplotypes.
165
166 **Statistical methods**
167　　Standard methods including student's t test and chi squared tests were used.
168
Results:
We used variants from the MSY region for our studies, including 16,679 indels and
34,764 SNVs located mostly in the ~22 MB non-PAR euchromatin region (31). Within
this dataset, we identified numerous variants that serve to uniquely characterize the non-
controversial haplotypes shared by both competing models. As an example, the NO
stem haplotype is defined by 75 SNVs and 9 indels, while the O stem haplotype is
delineated by 106 SNVs and 15 indels (Table 1). Similarly, the RQ stem haplotype is
defined by 6 SNVs. Furthermore, we uncovered a range of variants that define the stem
haplotypes exclusive to the ROE model, including A00A1a, AB, ABDE, and ABDEC
(Table 1 and Fig. 1A). An illustration of this is that A0b and A1a share 718 SNVs and
143 indels that are absent in other haplotypes/samples, supporting the A00A1a clade in
the ROE model. In contrast, no variants were found that would support or uniquely
define the ROA specific stem haplotype BT or CT (Table 1). The stem haplotype here
refers to a haplotype that is found on an internal branch or node of a phylogenetic tree,
representing a common ancestor from which several descendant haplotypes branch off.
The terminal haplotype refers to a haplotype that is found at the tips (or leaves) of the
phylogenetic tree. The results therefore support the ROE model and raise questions
regarding the validity of the ROA model.
Notably, relatively to the number of variants found previously that define the ROE
stem haplotypes, the full sequencing of 43 Y chromosomes has uncovered many new
variants (Table 1). There are now 718 SNVs that define A00A1a, compared to previous
39 SNVs, 324 SNVs that define AB, compared to previous 177 SNVs, for example (22).
Additionally, there are now a large number of indel variants that define these stem
haplotypes (Table 1) but few such indels have previously been found. The results thus

show that the full sequencing work has indeed as expected produced substantially more
variants that are informative to phylogenetic analyses.
We also observed that many variants are present in two or more haplotypes that do
not share a most recent common ancestor (MRCA), indicating convergent mutations.
For example, H1a and G2a share 3 SNVs and 4 indels that are absent in other
haplotypes/samples (H1a,G2a in Table 1). If these mutations had arisen in the MRCA of
G and H, they would have been present in all H, I, J, and K haplotypes, as these 4
haplotypes share a more recent MRCA than with G. Other representative examples of
sharing between two distantly related haplotypes include A1a,B2b; G2a,J2; G2a,J1;
B2b,G2a; B2b,H1a; C1a,G2a; C1a,H1a;N1a,H1a (Table 1). These cases indicate the
common occurrence of convergent mutations.
We systematically analyzed the pattern of shared variants among the 43 samples
(Table 2, see Supplementary Table S1 for an expanded version of Table 2 containing all
43 samples and Supplementary Table S2 for the same analysis of indels). For example,
among 6973 SNVs present in A0b, only 3275 are unique to A0b and the rest are shared
in various degrees with each of the other haplotypes (8.3-39.6% shared) represented by
the 43 samples (see A0b column in Table 2 and Supplementary Table S1). The results
again indicate that convergent evolution may be common.
Haplotypes that are closely related, sharing a MRCA, are expected to exhibit a
higher degree of variant sharing among them. This shared genetic material may arise
from common ancestry and convergent mutations. Convergent mutations, for instance,
may result from shared adaptations and two closely related groups with different Y
haplotypes are expected to share more adaptations. To assess the phylogenetic
relationships of different Y haplotypes, one can examine the extent of shared variants
among them. If, among the 43 haplotypes or samples, haplotype 1 shares the highest
fraction of variants with haplotype 2, and vice versa, these two haplotypes are
considered the closest to each other or are likely to belong to the same clade. For
example, sample NA19317 shared 95.4% of its variants with sample NA19347, a higher
percentage than that observed between NA19317 and any other sample listed in the
NA19317 column in Table 2. Similarly, NA19347 shared 91.1% of its variants with
NA19317, which is greater than the percentage shared with any other sample listed in
the NA19347 column in Table 2. These results suggest that these two samples are more
closely related to each other than to any of the other samples studied here. Indeed, the
two samples belong to the same clade, E1b1a1a1a1c1a1a3a1.
We determined the closest pairs among the 43 samples expected from the ROE
model or the ROA model based on the known haplotype identity of each sample (Table
3). There are 12 pairs according the ROE model and 11 according to the ROA model.
The two models differ in the expected pairing only with regard to the A0b-A1a pair, which
are most related to each other in the ROE model belong to the A00A1a clade (Fig. 1a).
In contrast, according to the ROA model, A1a, together with A1b, belongs to the A1
clade and A1a should be more closely related to haplotypes within the A1b clade, such
as B, C, E, or F, among others (see Fig. 1B).
Our approach identified 6 of 12 pairs in the ROE model that emerged as each
other's closest relatives based on the shared SNV fractions among the 43 samples
(Table 2 and Supplementary Table S1). Remarkably, the A0b and A1a pair was among
the 6 identified.
One possibility might explain that not all expected pairs could be identified by our
approach here. Two haplotypes may have diverged for very long time such as J1 and J2
and may each accumulate many independent mutations. Convergent evolution could
result in each being more related to a haplotype from a different clade. This could also
similarly account for recently diverged haplotypes not sharing the most with each other;

they may have encountered strong adaptive evolution with a haplotype from a distantly
related clade.
While not all closely related pairs of haplotypes could be identified as such by our
approach, it is important to note that there were not any pair of unrelated or distantly
related haplotypes that were mis-identified as the most closely related by our method.
For instance, G2a/HG03579 shared the most SNVs (57.7%) with the distantly
J1/HG01258, while J1/HG01258 shared the most (51.8%) with the distantly related
E1b1a1b1E/HG03065, rather than with G2a/HG03579 (Supplementary Table S1). Given
this observation, it is highly unlikely for the A0b-A1a pair, otherwise distantly related
according to the ROA model, to represent an exception to the pattern that only closely
related haplotypes could be identified as such by our approach.
We also performed the same analysis by using indel variants and obtained
consistent results (Supplemental Table S2). Together, these findings strongly support
the A00A1a clade within the ROE model, which encompasses A0b and A1a as terminal
branches. The data therefore once again support the ROE model and challenge the
ROA model.
According to the ROA model, the expected topological relationship between A1a
and A0b is akin to that between B and A1b1 (or A1a) or between E and B (as depicted in
Fig. 1B). Thus, if the ROA model predicts that A1a should exhibit the highest degree of
variant sharing with A0b, it would logically follow that E should share the most variants
with B. To put this prediction to the test, we conducted a thorough analysis of the sharing
patterns among all individual haplotypes and samples, as presented in Fig. 2. We used
the average value of the shared fractions of all individual E haplotypes or samples to
represent the value of E. The results revealed a significant disparity from the expected
outcomes under the ROA model. E was found to share more variants with C1a than with
B (Fig. 2A). Similar analysis using indel variants revealed consistent results (Fig. 2B).
Therefore, the close relationship observed between A0b and A1a contradicts the ROA
model.
The approach of inferring close relationships based on the level of shared variants
offers a fresh perspective for evaluating the two competing models of human origins. We
next focused on variants that define a stem haplotype. These variants are present in all
individual terminal haplotypes belonging to the respective stem haplotype. Such stem-
defining variants are more informative to the relationship between a stem haplotype and
another haplotype compared to the variants found in a terminal haplotype. We examined
whether the stem haplotypes exclusive to the ROE model exhibited the anticipated
relationships. As depicted in Fig. 3, the stem haplotype A00A1a of the ROE model
shared the most variants with B2b, a finding that aligns with the presence of the AB stem
haplotype. The AB stem haplotype in the ROE model shared the most variants with E,
while the stem haplotype E shared the most with A and B (except in the case of indels
where E shared more with C than A00A1a, Fig. 3B), all consistent with the existence of
the ABDE stem haplotype. Notably, the ABDE stem haplotype shared the most variants
with C, reinforcing the presence of the ABDEC stem haplotype. Additionally, F was
found to share variants nearly equally with A, B, E, and C, which is in line with
expectations derived from the ROE model. However, F should be closer to C than to A,
B, and E if the CF clade in the ROA model is real. These results lend substantial support
to the actual existence of all the stem haplotypes unique to the ROE model. By contrast,
the predictions derived from the ROA model do not align with the observed relationships
in Fig. 3. Therefore, these findings further challenge the ROA model.
All the stem haplotypes within the ROE model that are common in African
populations, including A00A1a, AB, ABDE, and ABDEC, are anticipated to exhibit equal
genetic relatedness to all F subtypes such as G, H, J, N, O, Q, and R. The observations

presented in Fig. 3 is largely in line with this expectation where G, H, and J shared
similar number of variants with these African stem haplotypes. However, haplotype
within the K2 clade (R, Q, O, and N) shared lower number of variants with these African
stem haplotypes compared to G, H, or J. This can likely be attributed to the fact that the
reference genome used for variant calling (GRCh38 Y) corresponds to the R1b
haplotype, which would inherently result in a reduced number of variants called for R
and its closely related Q, O, or N haplotype, as demonstrated in Table 2 and
Supplementary Table S1-S2 where the total number of variants called for each
haplotype or sample are listed. Some of the variants that were not called could include
some shared variants with the African stem haplotypes.
We extended our analysis to assess the stem haplotypes exclusive to the ROA
model, including BT, CT, and CF. According to the ROA model's predictions, BT should
exhibit the closest relationship to A1a, CT should be most closely related to B, and CF
should have the strongest affinity with E. However, our findings deviate from these
expectations. Our results indicated that BT is similarly related to A0b and A1a, or shares
slightly fewer variants with E1a than with A0b. CT is similarly related to B2b, A1a, and
A0b, while CF is similarly related to E, B2b, A1a, and A0b (Fig. 4). These results raise
questions about the validity of the stem haplotypes unique to the ROA model,
suggesting that they may not correspond to actual genetic relationships.
Finally, we studied the proportion of variant sharing between two haplotypes that is
due to convergent evolution. As shown in Fig. 4, for the O1b (NA18534) and O2a
(HG00673) pair, O1b (NA18534) has 3279 SNVs and shared 1651 with O2a (HG00673)
and 1254 with all four O2a samples in the dataset. So, the number of shared variants
between the O1b and O2a samples that are not due to common ancestry is 397 (1651-
1254). The proportion of shared variants due to convergent evolution is thus 0.24
(397/3279). Similarly, the proportion of convergent evolution for the E1a (NA19239) and
E1b (HG03471) pair is calculated to be 0.31, and for the R1a (HG03492) and R1b
(HG00731) pair is calculated to be 0.48 (Fig. 5). Analyses using indel variants showed
similar results but the proportion of convergent evolution is higher compared to SNVs
(Fig. 5). These data indicate that a substantial fraction of shared variants between any
two haplotypes is due to convergent evolution.
326
Discussions:
Within the ROE model, a well-defined hierarchy is observed, where A0b and A1a fall
within the A00A1a clade, A00A1a resides within the AB clade, AB is within ABDE clade,
E is encompassed within the ABDE, ABDE is nested within the ABDEC clade, and C
finds its place within the ABDEC clade. Remarkably, the anticipated variant sharing
patterns are in good alignment with the actual observations we have presented in our
study. These findings offer compelling support for the representation of the genetic
relationships among these haplotypes as offered by the ROE model.
Our study presents three observations that pose significant challenges to the ROA
model. First, in contrast to the predictions by the ROA model, we found that A1a
exhibited the closest relationship with A0b. However, this observation was not paralleled
by the expected proximity of E to B as prescribed by the ROA model, if one were to
consider the possibility of this observation being compatible with the ROA model.
Second, according to the ROA model, the E stem haplotype should be more closely
related to C than to B, and the F stem haplotype should have a closer genetic affinity
with C than with E. Our study did not align with these expectations. Lastly, the stem
haplotypes unique to the ROA model, namely BT, CT, and CF, all failed to exhibit the
anticipated sharing patterns. These observations collectively cast doubt on the

explanatory power of the ROA model to account for the genetic relationships we
observed in our study.
Influenced by the neutral theory, sharing of variants between any two haplotypes of
Y chromosome is previously thought to be all due to common ancestry. Our findings
here however showed that a substantial fraction of shared variants may be due to
convergent evolution (Fig. 5). Previous studies only used variants located in certain Y
chromosomal regions that can be easily identified by short read sequencing and did not
observe significant convergent evolution (21).
We have also employed ancient DNA analysis to provide independent validation of
the ROE model, as detailed in our previous work (22). The ancient DNA study also
revealed the common occurrence of convergent evolution. Ancient samples only show
mutations in a portion of the loci defining a haplotype, whether it is a basal haplotype or
a terminal haplotype. This implies that the diversification of terminal haplotypes had
actually occurred before the basal haplotype had completely mutated all relevant loci,
contrary to what the neutral theory predicts. This also suggests that the process of the
basal haplotype transitioning from partial to complete mutation of all loci occurs
simultaneously with the full formation of the terminal haplotype, necessitating the
involvement of convergent mutations. This is akin to the growth of a living tree, where
the development of leaves or branches occurs in sync with the growth of the trunk.
The ROE model is also strongly supported by the recent discoveries in human
fossils. The Yunxian crania from Hubei China, dated to 1 million years ago, are more
modern than *H. erectus* (reduced alveolar prognathism and no sagittal crest) and are
likely on the lineage leading to modern humans (32-36). The Hualongdong skull HLD6
from Anhui China at 300,000 years ago also has features of modern humans such as a
weak chin, high eye socket, flat face, a pronounced mental trigone, and a weak alveolar
prognathism (37, 38). The Dingcun teeth from Shanxi China are ~270,000 years old and
show modern features that are not found in Southern Africa fossils older than 100,000
years (39). It should be noted that different skull traits are not equally informative when
identifying modern human forms. Alveolar prognathism appears to be a highly
informative trait, yet it has long been somewhat overlooked. Most human fossils from
Africa, regardless of age, including Jebel Irhoud at 300,000 year ago (40), as well as a
high fraction of present-day sub-Sahara Africans, have alveolar prognathism. If humans
with some modern features had been living in East Asia at 1 million or 300,000 years
ago, it would be expected—barring any unforeseen random events—that modern
humans with the most complete modern traits would also have lived in East Asia during
the period when fully modern humans first emerged, i.e., between 100,000 and 50,000
years ago. Thus, the ROE model is well aligned with the fossil record.
The ROE model is more consistent with first principle logics. The group with the
leading cognitive abilities and other phenotypic traits should be the first to cross the
threshold into modern human status and maintain that status, barring any unexpected
events. Compared to other ancient human groups, the first group to become modern
humans should have a genome most distant from chimpanzees and maintain this
distinction to the present. The first population to become modern humans should have
lower maximum genetic diversity and maintain this low diversity unless disrupted by
unexpected events.
The regions suitable for the evolution of *Homo erectus* into modern humans should
also be suitable for the continued evolution of modern humans. Conversely, regions
where current populations are less advanced should be less likely to have been the
cradle of early modern humans. The ROA hypothesis contains an inherent contradiction:
it assumes that, while Africa was more conducive to human progress between 2 million
and 50,000 years ago, Eurasia became more suitable only in the last 50,000 years. As a

result, modern humans in Eurasia are genetically further removed from the great apes.
(41).
The complex and diverse environment of East Asia provides moderate challenges
suitable for progressive human evolution (42). Tropical regions have more pathogens,
which require populations living there to have stronger immunity. This increased
immunity, however, may not be conducive to reducing genetic diversity. As Gage noted
about the higher genetic diversity of chimpanzees compared to humans, "Humans might
rely less on biological variation than cultural and cognitive evolution—in the form of
medicine and engineering, for example—to adapt to changing environments." (43). By
this sound reasoning, human populations with relatively lower genetic diversity may
depend more on cultural and cognitive evolution than on biological evolution. Existing
research on culture and cognitive abilities appears to support this intuitive perspective.
A model should be self-consistent. Yet, the ROA model has a major self-
contradiction. For the first seven major haplotype splits in the Y chromosome tree
(Figure 1B, from A00 to F), the population associated with each newly derived haplotype
is consistently much larger than the ancestral population. For example, the A0-T
population is orders of magnitude greater than the ancestral A00 population, the BT
population is orders of magnitude greater than the ancestral A population, and so on,
with the F population being much larger than the C population at the seventh split,
implying a fitness advantage for each of the newly formed haplotypes, therefore
contradicting the neutral assumption required for building the tree in the first place. The
same issue also exists for the mtDNA tree of the ROA model.
If all haplotypes are equally fit, as assumed by the neutral theory, the original
ancestral haplotype of Y chromosome or mtDNA should be the most abundant and
widespread. This is because mutations leading to other haplotypes are rare events,
occurring only in a fraction of the entire population. Yet, the ancestral haplotype in the
ROA model, A00 for Y chromosome and L0 for mtDNA, is the rarest. In contrast, the
ancestral type in the ROE model, F for Y chromosome and R for mtDNA, is the most
abundant (Figure 1A).
The ROE theory does not necessitate concealing any truths, whereas the ROA
theory often overlooks certain scientific realities, such as the differences in genetic and
phenotypic distances between various human populations and great apes (41).
Furthermore, the ROA advocates tend to downplay the foundational role of the neutral
theory in the ROA model and the considerable uncertainty or controversy surrounding
that theory.
The ROA theory and its associated neutral theory provide only a limited
understanding of human uniqueness in terms of DNA, lacking insights into the
differences between humans and lower animals, as well as the distinctions between
modern and archaic humans. If one lacks a correct understanding of the specific
genomic features of humans, it would be difficult to infer human evolutionary processes
based on DNA data.
The results presented in our study offer yet another strong support for the ROE
model. Future molecular tests to further assess these Y chromosome models may
concentrate on obtaining more comprehensive sequencing data from additional ancient
and contemporary human Y chromosomes, with particular emphasis on populations from
Africa and East Asia. Expanding the scope of investigations to encompass a wider array
of genetic data will contribute to a more comprehensive and refined understanding of
human origins.
444
445 **Acknowledgements:**
446 This research was conducted without the direct support of any specific research grants.

447
Competing interests:
The author declares no competing interests.

References:

1.   Johnson MJ, Wallace DC, Ferris SD, Rattazzi MC, Cavalli-Sforza LL. Radiation of human mitochondria DNA types analyzed by restriction endonuclease cleavage patterns. J Mol Evol. 1983;19(3-4):255-71.
2.   Cann RL, Stoneking AC, Wilson AC. Mitochondrial DNA and human evolution. Nature. 1987;325:31-6.
3.   Yuan D, Lei X, Gui Y, Wang M, Zhang Y, Zhu Z, et al. Modern human origins: multiregional evolution of autosomes and East Asia origin of Y and mtDNA. bioRxiv. 2017:doi.org/10.1101/101410
4.   Zhang Y, Huang S. The Out of East Asia theory of modern human origins supported by recent ancient mtDNA findings. Acta Anthropologica Sinica. 2019;38(4):491-8.
5.   Zhang Y, Huang S. The Out of East Asia model versus the African Eve model of modern human origins in light of ancient mtDNA findings. bioRxiv. 2019:doi/10.1101/546234.
6.   Huang S. Molecular models of modern human origins. Prehistoric Archaeology 史前考古 (in Chinese). 2024:DOI : https://doi.org/10.3724/2097-3063.
7.   Hallast P, Agdzhoyan A, Balanovsky O, Xue Y, Tyler-Smith C. A Southeast Asian origin for present-day non-African human Y chromosomes. Human genetics. 2021;140(2):299-307.
8.   Underhill PA, Shen P, Lin AA, Jin L, Passarino G, Yang WH, et al. Y chromosome sequence variation and the history of human populations. Nat Genet. 2000;26(3):358-61.
9.   Ke Y, Su B, Song X, Lu D, Chen L, Li H, et al. African origin of modern humans in East Asia: a tale of 12,000 Y chromosomes. Science. 2001;292(5519):1151-3.
10.   Ingman M, Kaessmann H, Paabo S, Gyllensten U. Mitochondrial genome variation and the origin of modern humans. Nature. 2000;408:708-13.
11.   Mello B, Schrago CG. Modeling Substitution Rate Evolution across Lineages and Relaxing the Molecular Clock. Genome Biol Evol. 2024;16(9).
12.   Hu T, Long M, Yuan D, Zhu Z, Huang Y, Huang S. The genetic equidistance result, misreading by the molecular clock and neutral theory and reinterpretation nearly half of a century later. Sci China Life Sci. 2013;56:254-61.
13.   Lake NJ, Ma K, Liu W, Battle SL, Laricchia KM, Tiao G, et al. Quantifying constraint in the human mitochondrial genome. Nature. 2024.
14.   Lynch M, Wei W, Ye Z, Pfrender M. The genome-wide signature of short-term temporal selection. Proc Natl Acad Sci U S A. 2024;121(28):e2307107121.
15.   Kreitman M. The neutral theory is dead. Long live the neutral theory. Bioessays. 1996;18(8):678-83; discussion 83.
16.   Kern AD, Hahn MW. The Neutral Theory in Light of Natural Selection. Mol Biol Evol. 2018;35(6):1366-71.
17.   Huang S. New thoughts on an old riddle: What determines genetic diversity within and between species? Genomics. 2016;108(1):3-10.

18.   Teitz LS, Pyntikova T, Skaletsky H, Page DC. Selection Has Countered High Mutability to Preserve the Ancestral Copy Number of Y Chromosome Amplicons in Diverse Human Lineages. Am J Hum Genet. 2018;103(2):261-75.

19.   Zhu Z, Yuan D, Luo D, Lu X, Huang S. Enrichment of Minor Alleles of Common SNPs and Improved Risk Prediction for Parkinson's Disease. PLoS ONE. 2015;10(7):e0133421.

20.   Wang M, Huang S. The collective effects of genetic variants and complex traits. J Hum Genet. 2023;68:255-62.

21.   Poznik GD, Xue Y, Mendez FL, Willems TF, Massaia A, Wilson Sayres MA, et al. Punctuated bursts in human male demography inferred from 1,244 worldwide Y-chromosome sequences. Nat Genet. 2016;48(6):593-9.

22.   Xia Z, Chen H, Zhang Y, Huang S. Ancient uniparental DNAs in distinguishing the competing theories of molecular evolution and modern human origins. Research Square. 2021:DOI:10.21203/rs.3.rs-1087730/v1.

23.   Leffler EM, Bullaughey K, Matute DR, Meyer WK, Segurel L, Venkat A, et al. Revisiting an old riddle: what determines genetic diversity levels within species? PLoS Biol. 2012;10(9):e1001388.

24.   Bickel D. Testing Hypotheses of Molecular Evolution. In: Phylogenetic Trees and Molecular Evolution.: Springer, Cham.; 2022.

25.   黄石, 朱作斌. 系统生物学与进化理论. 朱作斌, 张潇, 王亮, editors. 南京: 东南大学出版社; 2022.

26.   Zhu Z, Han C, Huang S. New insights shed light on the enigma of genetic diversity and species complexity. Science China Life Sciences. 2024:https://doi.org/10.1007/s11427-023-2610-2.

27.   Huang S. The maximum genetic diversity theory of molecular evolution. Communications in Information and Systems. 2023;23:359-92.

28.   Fisher RA. The genetical theory of natural selection. Oxford, U.K.: Oxford University Press; 1930.

29.   Orr HA. Adaptation and the cost of complexity. Evolution. 2000;54(1):13-20.

30.   Gao H, Hamp T, Ede J, Schraiber JG, McRae J, Singer-Berk M, et al. The landscape of tolerated genetic variation in humans and primates. Science. 2023;380(6648):eabn8153.

31.   Hallast P, Ebert P, Loftus M, Yilmaz F, Audano PA, Logsdon GA, et al. Assembly of 43 human Y chromosomes reveals extensive complexity and variation. Nature. 2023;621(7978):355-64.

32.   Feng X, Lu D, Gao F, Fang Q, Feng Y, Huang X, et al. The phylogenetic position of the Yunxian cranium elucidates the origin of Dragon Man and the Denisovans. bioRxiv. 2024: https://doi.org/10.1101/2024.05.16.594603.

33.   Ji Q, Ji Y, Chen S, Liu A, Han K. The first descendants of Homo erectus recognized from China. J Geology. 2024(https://link.cnki.net/urlid/32.1796.p.20241031.1105.002).

34.   Li T, Etler DA. New Middle Pleistocene hominid crania from Yunxian in China. Nature. 1992;357(6377):404-7.

35.   Zhang Y. Fossil human crania from Yunxian: morphological comparison with Homo erectus crania from Zhoukoudian. Acta Anthropologica Sinica. 1995;14:1-7.

36.   Li T, Wang Z, Li W, Feng X, Wu X. Morphological features of human skulls from Quyuan River mouth, Yunzian, Hubei and their place in human evolution. Acta Anthropologica Sinica. 1994;13:104-16.

37.   Wu X, Pei S, Cai Y, Tong H, Zhang Z, Yan Y, et al. Morphological and
morphometric analyses of a late Middle Pleistocene hominin mandible from
Hualongdong, China. J Hum Evol. 2023;182:103411.
38.   Wu XJ, Pei SW, Cai YJ, Tong HW, Li Q, Dong Z, et al. Archaic human remains
from Hualongdong, China, and Middle Pleistocene human continuity and variation. Proc
Natl Acad Sci U S A. 2019;116(20):9820-4.
39.   Hu Y, Fan A, Shao Q-F, Li S-H, Hou Y-M, Zhao L-X, et al. New age of the
Dingcun 54:100 hominin site in northern China. J Archaeological Science: Reports.
2024;55:https://doi.org/10.1016/j.jasrep.2024.104502.
40.   Hublin JJ, Ben-Ncer A, Bailey SE, Freidline SE, Neubauer S, Skinner MM, et al.
New fossils from Jebel Irhoud, Morocco and the pan-African origin of Homo sapiens.
Nature. 2017;546(7657):289-92.
41.   Mallick S, Li H, Lipson M, Mathieson I, Gymrek M, Racimo F, et al. The Simons
Genome Diversity Project: 300 genomes from 142 diverse populations. Nature.
2016;538(7624):201-6.
42.   Zeller E, Timmermann A, Yun KS, Raia P, Stein K, Ruan J. Human adaptation to
diverse biomes over the past 3 million years. Science. 2023;380(6645):604-8.
43.   Gage F, Nair P. QnAs with Fred Gage. Proc Natl Acad Sci U S A.
2014;111(36):12958-9.
561
562
563

564   **Tables:**
565
566   **Table 1. Number of variants uniquely found for each haplotype or combination of**
567   **haplotypes.**
568   The listed haplotypes represent only a fraction of the complete set.
569

Haplotype	# SNVs	# indels	Previous #SNVs
A00A1a	718	143	39
AB	324	60	177
ABDE	5	1	4
ABDEC	192	41	155
CT	0	0	0
BT	0	0	0
RQ	6	0	n.d.
NO	75	9	40
O	106	15	n.d.
A1a,B2b	3	27	n.d.
H1a, G2a	3	4	n.d.
G2a,J2	1	2	n.d.
G2a,J1	3	1	n.d.
B2b,G2a	1	6	n.d.
B2b,H1a	13	8	n.d.
C1a,G2a	3	4	n.d.
C1a,H1a	6	6	n.d.
N1a,H1a	0	2	n.d.

Group	ID	Haplotype	# Total	# Unique	# Shared	ACB HG01890 A0b	GWD HG02666 A1a	LWK NA19384 B2b	YRI NA19239 E1a2a1a1a	GWD HG03248 E1a2b1a1a1	GWD HG02572 E1a2b1a2	MSL HG03471 E1b1a1a1a	ACB HG02486 E1b1a1a1a1c1a1a	LWK NA19317 E1b1a1a1c	LWK NA19347 E1b1a1a1a1c
ACB	HG01890	A0b	6973	3275	3698		**0.465**	0.470	0.454	0.522	0.468	0.490	0.484	0.484	0.511
GWD	HG02666	A1a	5677	2486	3191	**0.390**		0.481	0.415	0.534	0.383	0.441	0.432	0.428	0.419
LWK	NA19384	B2b	4126	1649	2477	0.278	0.391		0.421	0.550	0.404	0.462	0.459	0.449	0.447
YRI	NA19239	E1a2a1a1a	3645	256	3389	0.237	0.295	0.372		0.892	0.739	0.594	0.558	0.542	0.522
GWD	HG03248	E1a2b1a1a1	2753	151	2602	0.206	0.317	0.367	0.674		0.600	0.502	0.489	0.478	0.459
GWD	HG02572	E1a2b1a2	4076	255	3821	0.274	0.301	0.399	0.827	0.888		0.611	0.588	0.580	0.577
MSL	HG03471	E1b1a1a1a1c	3587	255	3332	0.252	0.268	0.402	0.584	0.654	0.538		0.845	0.808	0.782
ACB	HG02486	E1b1a1a1a1c1a1a	3654	101	3553	0.254	0.319	0.407	0.559	0.649	0.527	0.861		0.890	0.863
LWK	NA19317	E1b1a1a1a1c1a1a3a1	3714	92	3622	0.258	0.336	0.404	0.552	0.645	0.528	0.837	0.905		**0.911**
LWK	NA19347	E1b1a1a1a1c1a1a3a1	3889	12	3877	0.285	0.342	0.421	0.557	0.649	0.551	0.848	0.919	**0.954**	
ESN	HG03371	E1b1a1a1a1c1a1a3c1b	3819	187	3632	0.253	0.324	0.413	0.567	0.646	0.534	0.876	0.930	0.903	0.881
ACB	HG02011	E1b1a1a1a1c1b	3607	126	3481	0.252	0.299	0.410	0.578	0.644	0.544	0.882	0.868	0.872	0.833
GWD	HG02717	E1b1a1a1a1c2c3a2a	3967	371	3596	0.265	0.293	0.407	0.580	0.657	0.535	0.865	0.872	0.848	0.816
ACB	HG02554	E1b1a1a1a2a1	3734	157	3577	0.260	0.313	0.403	0.570	0.642	0.520	0.847	0.857	0.844	0.806
ESN	HG02953	E1b1a1a1a2a1	3800	160	3640	0.260	0.325	0.423	0.579	0.660	0.551	0.883	0.905	0.864	0.842
ASW	NA19705	E1b1a1a1a2a1a3b1a1	3744	171	3573	0.256	0.325	0.400	0.558	0.633	0.534	0.853	0.871	0.866	0.823
MSL	HG03065	E1b1a1b1	4095	348	3747	0.244	0.303	0.398	0.610	0.638	0.582	0.785	0.786	0.780	0.755
PUR	HG01109	E1b1a1b1a	3579	443	3136	0.223	0.301	0.408	0.540	0.632	0.525	0.572	0.570	0.555	0.545
PUR	HG01106	E1b1a1b1a1	3837	522	3315	0.256	0.276	0.373	0.607	0.640	0.570	0.616	0.610	0.591	0.597
CLM	HG01457	E1b1b1b2a1a4	3816	496	3320	0.244	0.285	0.400	0.603	0.617	0.562	0.634	0.626	0.623	0.607
LWK	NA19331	E1b1b1b2b2a1	3959	523	3436	0.251	0.304	0.362	0.617	0.655	0.576	0.601	0.569	0.556	0.578
JPT	NA18989	C1a	5466	2522	2944	0.257	0.322	0.397	0.525	0.522	0.493	0.515	0.511	0.498	0.482
MSL	HG03579	G2a	3318	966	2352	0.211	0.260	0.339	0.459	0.442	0.448	0.451	0.426	0.425	0.411
BEB	HG03009	H1a	3161	1039	2122	0.220	0.233	0.343	0.431	0.468	0.401	0.471	0.444	0.430	0.423
CLM	HG01258	J1	3986	791	3195	0.218	0.264	0.358	0.487	0.456	0.486	0.470	0.452	0.447	0.444
PJL	HG02492	J2	2950	674	2276	0.189	0.296	0.345	0.358	0.463	0.333	0.387	0.380	0.361	0.354
FIN	HG00358	N1a	2447	698	1749	0.185	0.237	0.310	0.321	0.426	0.306	0.345	0.337	0.328	0.324
CHB	NA18534	O1b	3279	748	2531	0.203	0.246	0.310	0.453	0.424	0.447	0.424	0.402	0.394	0.384
CHS	HG00673	O2a2a1a1a	2771	446	2325	0.197	0.228	0.325	0.396	0.433	0.371	0.367	0.359	0.349	0.373
CHS	HG00621	O2a2b1a1a1	2761	280	2481	0.212	0.243	0.304	0.412	0.441	0.381	0.430	0.402	0.397	0.406
CHS	HG00512	O2a2b1a2a1a	3878	561	3317	0.231	0.243	0.348	0.483	0.438	0.478	0.451	0.429	0.432	0.458
KHV	HG01596	O2a1b1a1a1a1c	3077	553	2524	0.217	0.259	0.323	0.408	0.442	0.373	0.419	0.413	0.413	0.400
PEL	HG01928	Q1b1a1a	2481	207	2274	0.170	0.198	0.276	0.359	0.355	0.366	0.372	0.357	0.358	0.346
PEL	HG01952	Q1b1a1a	2566	283	2283	0.162	0.217	0.256	0.378	0.339	0.357	0.355	0.326	0.324	0.314
PJL	HG03492	R1a1a1b2a1a2c	2160	556	1604	0.139	0.168	0.206	0.349	0.292	0.292	0.309	0.281	0.265	0.257
PUR	HG00731	R1b1a1b1a1a2a5	1105	66	1039	0.091	0.138	0.136	0.216	0.186	0.186	0.215	0.192	0.184	0.184
PUR	HG01243	R1b1a1b1a1a2a5	1028	97	931	0.084	0.164	0.119	0.192	0.179	0.166	0.190	0.175	0.163	0.155
MXL	NA19650	R1b1a1b1a1a2b1a1	1135	124	1011	0.083	0.140	0.121	0.205	0.179	0.185	0.192	0.177	0.164	0.157
TSI	NA20509	R1b1a1b1a1a2b1a1	1314	91	1223	0.088	0.140	0.127	0.260	0.181	0.223	0.198	0.183	0.170	0.162
CLM	HG01358	R1b1a1b1a1a2b1a1	1319	121	1198	0.113	0.151	0.124	0.183	0.164	0.198	0.194	0.179	0.169	0.204
GBR	HG00096	R1b1a1b1a1a2c1a1f1a2	1216	189	1027	0.089	0.180	0.129	0.195	0.185	0.177	0.201	0.177	0.177	0.169
IBS	HG01505	R1b1a1b1a1a2c1a4b2d1	1322	132	1190	0.086	0.141	0.129	0.253	0.185	0.211	0.200	0.185	0.172	0.165
ITU	HG03732	R2	2019	597	1422	0.143	0.237	0.249	0.271	0.312	0.272	0.307	0.315	0.309	0.309

Table 3. Expected and observed pair of closely related samples within the 43 samples dataset.

Group	ID	Haplotype	Expected closest relative		Observed	
			OOEA	OOA	SNVs	Indels
ACB	HG01890	A0b				
GWD	HG02666	A1a	HG01890		yes	yes
LWK	NA19384	B2b				
YRI	NA19239	E1a2a1a1a				
GWD	HG03248	E1a2b1a1a1	HG02572	same	no	no
GWD	HG02572	E1a2b1a2				
MSL	HG03471	E1b1a1a1a1c				
ACB	HG02486	E1b1a1a1a1c1a1a				
LWK	NA19317	E1b1a1a1a1c1a1a3a1	NA19347	same	yes	yes
LWK	NA19347	E1b1a1a1a1c1a1a3a1				
ESN	HG03371	E1b1a1a1a1c1a1a3c1b				
ACB	HG02011	E1b1a1a1a1c1b				
GWD	HG02717	E1b1a1a1a1c2c3a2a	HG02011 or HG02554/3	same	yes	no
ACB	HG02554	E1b1a1a1a2a1				
ESN	HG02953	E1b1a1a1a2a1				
ASW	NA19705	E1b1a1a1a2a1a3b1a1	HG02953 or HG02554	same	yes	no
MSL	HG03065	E1b1a1b1				
PUR	HG01109	E1b1b1a1b1a				
PUR	HG01106	E1b1b1b1a1				
CLM	HG01457	E1b1b1b2a1a4	NA19331	same	no	no
LWK	NA19331	E1b1b1b2b2a1				
JPT	NA18989	C1a				
MSL	HG03579	G2a				
BEB	HG03009	H1a				
CLM	HG01258	J1	HG02492	same	no	no
PJL	HG02492	J2				
FIN	HG00358	N1a				
CHB	NA18534	O1b				
CHS	HG00673	O2a2a1a1a				
CHS	HG00621	O2a2b1a1a1				
CHS	HG00512	O2a2b1a2a1a	HG00621 or HG00596	same	yes	yes
KHV	HG01596	O2a1b1a1a1a1c				
PEL	HG01928	Q1b1a1a	HG01952	same	yes	yes
PEL	HG01952	Q1b1a1a				
PJL	HG03492	R1a1a1b2a1a2c				
PUR	HG00731	R1b1a1b1a1a2a5	HG01243	same	no	no
PUR	HG01243	R1b1a1b1a1a2a5				
MXL	NA19650	R1b1a1b1a1a2b1a1				
TSI	NA20509	R1b1a1b1a1a2b1a1	NA19650 or HG01358	same	no	no
CLM	HG01358	R1b1a1b1a1a2b1a1				
GBR	HG00096	R1b1a1b1a1a2c1a1f1a2	HG01505	same	no	yes
IBS	HG01505	R1b1a1b1a1a2c1a4b2d1				
ITU	HG03732	R2				

588
589 **Figure legends:**
590
Fig. 1. Y chromosome phylogenetic trees of modern humans. A. The Out of East
Asia model. B. The Out of Africa model. Only major branches and representative SNVs
are shown with branch lengths not to scale. The haplotype topologies for F subtypes
under the ROA model, including G, H, I, J, and K, are consistent and identical between
the ROA model and the ROE model. These haplotypes share the same genetic
relationships and branching patterns in both models.
597

598
599
600

Fig. 2. Fraction of shared variants among individual terminal haplotypes. A.
Fraction of shared SNVs. B. Fraction of shared indels. In the case of haplotypes like E, J,
O, Q, and R that have more than one individual terminal haplotype/sample represented
in the dataset of 43 chromosomes, the average value of all samples under a haplotype is
shown with error bars representing the standard deviation of the data. The information
used to generate the figures were extracted from Supplementary Table S1 and S2.
607

Fig. 3. Fraction of shared variants between a stem haplotype in the ROE model and other individual terminal haplotypes in the 43 human Y chromosomes. A. Fraction of shared SNVs. B. Fraction of shared indels. For haplotypes such as A, E, J, O, Q, and R that have more than one individual terminal haplotype/sample represented in the 43 chromosomes, the average value of all samples under a haplotype is shown with error bars representing the standard deviation.

Fig. 4. Fraction of shared variants between a stem haplotype in the ROA model and other individual terminal haplotypes in the 43 human Y chromosomes. A. Fraction of shared SNVs. B. Fraction of shared indels. For haplotypes such as E that have more than one individual terminal haplotype/sample represented in the 43 chromosomes, the average value of all samples under a stem haplotype is shown with error bars representing the standard deviation.

632 **Fig. 5. Number and fraction of shared variants due to convergent evolution in**
633 **selected haplotypes.** A. Number of variants. B. Fraction of shared variants due to
634 convergent evolution.

636