Wintering, rather than breeding, oceanic conditions may modulate declining survival in a long-distance migratory seabird

Running page head: oceanic conditions influence survival

Katherine R. S. Snell[1,2,3] [§][*], Inês Alexandre Machado dos Santos[1] [§], Rob SA van Bemmelen[4], Børge Moe[5] & Kasper Thorup[1,6]

[1] Center for Macroecology, Evolution and Climate, Globe Institute, University of Copenhagen, Universitetsparken 15, DK-2100 Copenhagen, Denmark [2] MPI-AB, Am Obstberg 1, Radolfzell, 78315, Germany [3] Centre for the Advanced Study of Collective Behaviour, University of Konstanz, Konstanz, 78464, Germany [4] Waardenburg Ecology, Culemborg, 4101 CK, the Netherlands and Wageningen Marine Research, IJmuiden, 1976 CP, the Netherlands [5] NINA, Trondheim, 7485, Norway [6] Copenhagen Bird Ringing Centre, Natural History Museum of Denmark, University of Copenhagen, Øster Voldgade 5-7, DK-1350 Copenhagen, Denmark

[§] These authors contributed equally to this work [*] corresponding author

ksnell@ab.mpg.de

ABSTRACT

Steep declines in Arctic skua populations in the southern extent of their breeding range have been reported during the last half of the 20[th] century. We used 24 years of ringing and re-encounter data from the Faroe Islands, North Atlantic, to investigate if patterns in survival probabilities can be explained by large scale climatic events. Having first determined the migratory phenology and wintering regions, we estimated the effects on survival of the North Atlantic Oscillation (NAO) index during breeding and Oceanic Niño index (ONI) during the non-breeding period within a capture-mark-recapture framework. Temporal trends, and direct and time-lagged effects of the environment on annual survival were modelled. We found support for a substantial decrease in adult annual survival, from ca. 0.93 in 1985 to ca. 0.77 in 2008, and weak support for a decrease in young (1[st] year) survival over the duration of the study period. Furthermore, models indicated increased young survival following an El Niño winter. We suggest this time-lagged effect reflects downstream propagation of environmental conditions, particularly food availability, or a potential carry-over effect of El Niño conditions positively impacting the performance of the parents in the subsequent breeding season, leading to improved young survival prospects. While adult mortality cannot be attributed to the oceanic climate oscillations tested here; the negative trend in survival may account for the substantial population declines observed over the last decades.

Keywords: mortality, carry-over effects, CMR modelling, *Stercorarius parasiticus*

1. INTRODUCTION

Seabirds occupy the upper trophic level and are important in island and marine ecosystem processes, function and resilience (Paleczny et al. 2015). Their distribution and population dynamics are strongly connected with their feeding ecology and patterns of food availability which are dependent on climatic and oceanographic factors, such as water temperature and currents (Meier et al. 2017). Therefore, changes in any such environmental variables can substantially influence seabird survival and productivity (Irons et al. 2008, Waugh et al. 2015, Champagnon et al. 2018). Steep declines in seabird numbers have been reported during the last half of the 20th century. Palenczny et al. (2015) assessed the population trend of the world's monitored seabirds from a global dataset representing ca. 19% of the global seabird population and found an overall decline of 69.7% in between 1950-2010: a pattern reflected in the population trends of seabirds in the North Atlantic (Irons et al., 2008). Moreover, population declines are more apparent in migratory species (Kubelka et al. 2021).

Given that a broad range of factors may affect demographic parameters, identifying the factors responsible for population dynamics is often difficult. Unravelling effects of different factors is especially challenging for migratory species, as they depend on multiple, widely separated areas that may contrast in their (rate of change in) conditions. As migratory species spend the majority of their annual cycle on migration or overwintering, the habitat conditions experienced during these periods are likely to influence survival of all age classes post-fledging. Likewise, conditions on breeding grounds influence reproductive success, with decreases in productivity in years where conditions are poorer (Fife et al. 2018). In addition to these within-season effects, conditions experienced in one season can affect the condition and performance of

individuals in later seasons (Harrison et al. 2011). The effects of climate on species

survival have received increased interest during the last few decades due to accelerated

climate change (Jóhannesson et al. 1995, Ballerini et al. 2015, Salas et al. 2017). As

shown in several studies, climate indices can influence survival and breeding success of

both migrant and non-migrant birds (England 2000, Irons et al. 2008, Boano et al. 2010,

Genovart et al. 2013, Waugh et al. 2015). For example, blue-footed booby (*Sula*

nebouxii) in the Gálapagos suffered increased mortality during an El Niño year, when

sea surface temperature (SST) was elevated; similarly, chick growth rates of dark-

rumped petrel (*Pterodroma phaeopygia*) were retarded during the 1983 nesting season,

one of the strongest El Niño events since records began (England 2000). Formal

analysis of long-term data revealed a complex interaction of stochastic oceanic

processes on chick quality of sooty shearwaters (*Puffinus griseus*), including a time-

lagged effect of El Niño oscillations (Humphries & Möller 2017). In adult blue petrels

(*Halobaeria caerula*), Barbraud and Weimerskirch (2003) found that increased

mortality occurred when the El Niño Southern Oscillation phase was associated with

warmer water temperatures. The influence of climate indices on seabird survival rates

are likely explained by the important impact of stochastic ocean scale environmental

processes on ecosystem cascades affecting nutrients, temperature and primary

productivity (Polis et al. 1997). Formally linking survival and reproductive rates to

climate indices, such as the El Niño Southern Oscillation (ENSO) and North Atlantic

Oscillation (NAO), can be a first step towards understanding the ultimate or proximate

drivers of population dynamics (Descamps et al. 2010).

The influence of the ENSO in the Pacific Ocean triggers a cascade of events globally

across equatorial regions and the Southern Ocean, with delayed propagation of

conditions across the Atlantic, albeit less robust (Kim et al. 2023). During El Niño

periods, warm Pacific waters are associated with increased atmospheric stability and

suppressed tropical cyclone genesis over the Atlantic; while during La Niña years, cold

equatorial Pacific SSTs are associated with decreased stability and increased tropical

cyclones in the Atlantic between *ca.* 0-30°N (Kim et al. 2023). In the year following an

El Niño event, warm water in the Atlantic may contribute to increased hurricane

formation from Cape Verde towards the Americas (Kim et al. 2023). Overall higher

wind speeds and lower SSTs around Atlantic upwellings are associated with significant

increases in primary productivity (chlorophyll *a*) (Oviatt et al. 2015), which likely

propagates into higher trophic levels. In contrast, periods of above average SSTs leads

to reduced egg and fish abundance (Franco et al. 2020). The NAO index is simply the

pressure differential between Iceland and the Azores. In the North Atlantic region the

positive phase is related to warmer/wetter ambient conditions and more severe storms

(Hurrell et al. 2001, Báez et al. 2021). Milder weather conditions lead to an advanced

reproductive phenology (Báez et al. 2021), while increased precipitation and extreme

weather events may have a direct impact on higher mortality (Stenseth et al. 2002,

Frederiksen et al. 2008), and an increased energetic cost of migration (Nourani et al.

2023). Negative NAO phases are linked to relatively warmer water, cooler air

temperature but dry/stable summers. NAO influences zooplankton abundance; and

reflects fish both abundance and body condition, with fish larvae particularly sensitive

to variation in water conditions (Stenseth et al. 2002, Drinkwater et al. 2003). In the

waters east of the Faroes, NAO has a significant positive relationship with lower trophic

levels: phytoplankton and zooplankton (Drinkwater et al. 2003). Around the Faroes,

overall krill abundance was attributed to temperature and chlorophyll *a* (Silva et al.

2013); and the positive phase of the NAO is favourable for cod stocks, however the

opposite is found in the North Sea, although the effect size is small (Drinkwater et al.

2003, Overland et al. 2010). Due to intense fishing practices there are difficulties in

attributing prey demographics to oceanic processes in many regions (Oviatt et al. 2015).

Cory's shearwaters (*Calonectris borealis*) increased their foraging effort in negative

phase NAO years, which was attributed to lower marine productivity in the eastern mid-

North Atlantic (Paiva et al. 2017). In Scotland, NAO phase is a predictor of breeding

phenology in black-legged kittiwakes (*Rissa tridactyla*) and common guillemots (*Uria*

aalge), and breeding success of northern fulmars (*Fulmarus glacialis*) and subsequent

population trajectories (Thompson & Ollason 2001, Frederiksen et al. 2004). While

NAO clearly can be an important factor in seabird ecology, there are difficulties in

making specific predictions for a given population.

Indices of large-scale ocean processes reflect multiple variations in the marine

environment, including water and air temperatures, winds, salinity, run-off, and

precipitation, and capture more of the environmental stochasticity than individual

metrics. In general the influence of the oceanic phase is likely to be cumulative and

dynamic and lags can be expected for some effects (Stenseth et al. 2002). While the

underlying bottom-up mechanisms are complex and geographically explicit, they

provide a valuable tool to elicit variation in biological processes.

Like many populations of vulnerable seabird species in Northwest Europe, (Dias et al.

2019), Arctic skua (*Stercorarius parasiticus*) populations are declining rapidly (Perkins

et al. 2018). The Arctic skua is a migratory seabird with a circumpolar breeding range in

high temperate to high Arctic zones. It is long-lived, with immature birds recruiting to

the breeding population from an age of 4 years (O'Donald 1983, Furness 1987). They

are kleptoparasitic and, thus, highly dependent on other seabirds (Furness 1987), most of which have also experienced sharp population declines (BirdLife International 2016). Although the Arctic skua is listed as 'least concern' globally, with a world population estimated at 4-600 000, it is considered 'endangered' within the European Union (Burfield & van Bommel 2004). In the United Kingdom the species has declined by *ca.* 66% during the period 2015-2021 (Burnell et al. 2023), and in the Faroe Islands the population declined from 1200-1500 pairs in 1981 (Bengtson & Bloch 2003) to *ca.* 207 pairs in 2017 (Santos 2018). The proposed drivers of these declines are geographically limited but, in the United Kingdom and Norway, associations were found with food shortages and predation of eggs and young (Perkins et al. 2018, van Bemmelen et al. 2021).

The decline of European Arctic skua populations has hitherto been mainly attributed to factors during the breeding periods on reproductive success, and adult survival was considered to be sufficiently high to maintain a stable population. However, a full understanding of the population declines requires quantification of potential effects from both the breeding and non-breeding periods, on both adult and immature mortality (Flack et al. 2022). We hypothesise that for long-lived, long-distance migratory seabirds non-breeding environmental conditions, rather than breeding conditions, influence survival rates and thereby population dynamics. Population rates for these life history traits are expected to be most sensitive to adult survival (Morris et al. 2011), as such are generally buffered against variations in environmental conditions experienced (Hilde et al. 2020). However, extreme weather events influence survival (Frederiksen et al. 2008) and are likely to be encountered more often during the non-breeding period, when conditions are less predictable (Lisovski et al. 2017). Here, we investigated annual

survival of adult and young Arctic skuas breeding in the Faroe Islands. Within a formal

capture-mark-recapture framework we explored annual survival, trends in annual

survival and the influence of oceanic scale climatic events, precisely the El Niño

Southern Oscillation (ENSO) and North Atlantic Oscillation (NAO), using a ringing

dataset spanning over two decades. We predict that non-breeding conditions, the largest

component of the annual cycle (van Bemmelen et al. 2024), will influence annual

survival of both young and adults, with El Niño years having an overall negative effect

due to increased water temperatures leading to reduced marine productivity.

Furthermore, in long-lived species, we expect adult survival to be largely buffered

against environmental variability; as such, mortality will firstly be greater in young than

in adults, and secondly, instead of only affecting survival, poor feeding conditions may

be expected to carry over to the breeding performance due to the adults' poorer

condition upon arrival and a subsequent increased mortality of offspring.

2. MATERIALS & METHODS

2.1 Study Species and sites

The Faroe Islands (61°53'N, 6°58'W) are located between the UK and Iceland, in the middle of the North Atlantic current which creates stable temperatures and a maritime climate (Fosaa et al. 2004). The Faroese archipelago consists of 18 islands and is an important breeding site for seabirds (Bakken et al. 2006). Ringing of Arctic skuas has been conducted since 1924 across the archipelago but was not systematic (Hammer et al. 2014).

As part of a tracking programme covering the Atlantic AMAs (Arctic Marine Areas) (Kuletz et al. 2017), tracking data was obtained from 20 birds in the Fugloy colony from 2016-7 (Fig. 1). Full details of the analysis and interpretation of light level data are detailed in van Bemmelen et al. (2024). The Faroese Arctic skua population winter on the Patagonian Shelf, Discovery Seamounts, coastally from the Iberian plain to Cape Verde Islands, the Gulf of Guinea, and off the Namibian coast and the Cape of Good Hope (Fig.1). Spatio-temporal data was used to select appropriate environmental indices. Mean arrival and departure dates were 27 April and 8 August for breeding and 9 September and 21 March for over-wintering sites, respectively.

2.2 Environmental Variables

Standardised large-scale oceanic indices were selected appropriate for a spatio-temporal schedule of the long-distance migrant. The Oceanic Niño index (ONI) quantifies the phase and strength of El Niño Southern Oscillation (ENSO) events, where negative values correspond to La Niña years and positive values to El Niño years (NOAA

2018a). The North Atlantic Oscillation index (NAO) is the air pressure differential

across the North Atlantic and represents the phases of the North Atlantic Oscillation

(NOAA 2018b). Both ONI and NAO data are presented as aggregated 3-month

overlapping temporal periods with values ranging from -1.64 to 2.24 and -1.4 to 1.8,

respectively (Fig. 2). Wintering and breeding periods were derived from the GLS data.

We calculated mean ONI during the stationary non-breeding season (September to

March) and mean NAO during the breeding season (May to July; Fig. 2). Values were

used as covariates in the survival model both as a direct effect (same year) and as a

time-lagged effect from the previous year.

2.3 Survival Analysis

We used a subset (1985 to 2008) of the ringing and recovery data which had been

digitised and had the greatest continuity of data (prior to 1985 gaps of >2 years in

ringing effort occurred). Where discrepancies made the record unreliable, we excluded

these individuals from our dataset (n=2). The dataset totalled 1060 individuals, of which

530 were ringed as chicks and 530 ringed as adults; 85 individuals were subsequently

re-encountered (alive and/or dead) during the 24 years (suppl. info. Fig. S1). Live re-

encounters consisted of ring reading during breeding season fieldwork; dead recoveries

included both individuals recovered from the study colony and rings reported to the

ringing scheme. The reporting probability structure is similar to the ringing effort for

the period included in this study. A year was defined as starting on the 1[st] May and

ending on the 30[th] April of the following calendar year, to coincide with the birds'

breeding cycle. For young birds the initial year is the period from ringing (typically

from 10 days after hatching) until the following breeding season – and therefore

includes late chick stage, fledging, and dependant period.

Annual survival was estimated from a formal capture-mark-recapture framework with the Mark software, version 10.1 (White & Burnham 1999). Firstly, we use the Burnham model for joint live and dead encounters (Burnham 1993). This method calculates parameter estimates of apparent survival (S), live re-encounter probability (p), dead recovery probability (r) and fidelity (F). An encounter history was created for each individual in the classic 2 age-class structure: for young (1[st] year) and for adults and included both live and dead encounters. We modelled fidelity as constant due to known high philopatry and no reported post-natal dispersal between the Faroe Islands and the nearest colonies in the UK (Hammer et al. 2014): F was found to be high, F=0.94 +/- 0.065 SE. As Arctic skuas generally delay breeding until their 4[th] season (O'Donald 1983, Furness 1987), re-encounter (p) and recovery probability (r) were modelled as a function of 3 age-classes: young, immature (Imm) and adults. During the immature years (age 1 to 3 inclusive), birds are thought to be entirely marine, not returning to land once they leave the colony (Furness 1987); as such p(Young), p(Imm) and r(Imm) were fixed to 0. As ringed young birds could be recovered dead in the colony after the breeding season had finished r(Young) was estimable (there were three occurrences in the data where birds ringed as chicks were recovered dead within the first year). Survival probability (S) was modelled as a function of 2 age-class (Young and Adults) with the assumption that once birds are full-grown immatures, after 1 year, they will have the same mortality as breeding adults. As the joint model includes more parameters than a live-only re-encounter model we also repeated the analysis by modelling apparent survival with a Cormack-Jolly-Seber (CJS) model (Lebreton et al. 1992). This method calculates parameter estimates of apparent survival (Φ) and live re-encounter probability (p). The encounter history for each individual was created in the

same 2 age-class structure: for young (1[st] year) and for adults using only live encounter

data. The structure of the model followed the Burnham model, and p(Young) and

p(Imm) were fixed to 0.

We derived a basic model (i.e., a single survival probability over all years) and

modelled year-dependent survival for each age-class. We then compared the following *a*

priori candidate models: linear temporal trends of survival rates for both age-classes

combined and for each age-class independently, and the effects of the two

environmental covariates (detailed above) on survival probabilities. We estimated a

dispersion coefficient with the bootstrap goodness-of-fit function in MARK to the

general model and some over-dispersion was indicated. Residuals were visually

inspected and minor departures from goodness-of-fit did not indicate a structural issue.

Therefore, we calculated the variance inflation factor (ĉ) via the median c-hat procedure

in MARK, and then used this to adjust AICc values (Akaike 1973). We tested the

overall CJS model encounter histories using the overall_CJS() function in the R

package R2ucare (Gimenez et al. 2018; a formal test for the Burnham model is not

available). This tests the basic assumptions of that model and indicated the CJS models

were appropriate for the data and any overdispersion in the data was not significant (χ^2

= 16.181, p = 0.973, df = 29). No formal statistical method for testing goodness-of-fit

(GOF) for models with linear covariates (here, trends over time or environmental

covariates) is available. Model selection was based on the ranking of Akaike's

information criteria corrected for small sample size adjusted for median-ĉ (AICc).

Survival estimates ± SE are given as back transformed from the logit link function Beta

parameter estimates. Linear trends are visualised with 95% confidence intervals.

3. RESULTS

The basic joint Burnham model (model B, Table 1) estimated the overall survival of young as 0.43±0.14 and adults 0.92±0.07. Year dependant annual survival was estimable in 10 of the 24 years for adults, ranging from 0.16±0.15 to 0.87±0.51, and in 8 years for young, ranging from 0.13±0.12 to 0.66±0.67 (the time-dependant model is not included in Table 1 due to the high number of non-estimable parameters).

Three models were well supported (ΔAICc < 2, Table 1; model C1, T1 & T2). The highest ranked model (model C1), indicated a combined negative trend for adult annual survival and the influence of a time-lagged effect of ONI (Oceanic Niño Index) on young survival (Table 1; suppl. info. Table S1 for parameter estimates); it should be noted that this model is nested. Evidence for a negative trend in adult annual survival is further supported by model T1 (ΔAICc = 2.86 from model B, the basic model (Table 1; suppl. info. Table S2 for parameter estimates)). We found some support for a decreasing trend for young birds also (Table 1; suppl. info. Table S3 for parameter estimates). Based on model T2, the adults' survival showed a decline of 17% in 24 years, with 0.93±0.05 probability of survival in 1985 and 0.77±0.09 in 2008, whereas young survival decreased 57%, from 0.60±0.19 in 1985 to 0.26±0.14 on 2008 (Fig. 3a,b). Adult survival estimates from model C1 and T1 were comparable: within the error range of model T2 estimates.

Model support indicated that young survival could be influenced in part by ONI from the preceding year. El Niño events were related to an increase in apparent survival while La Niña events a reduction (Fig. 3c & Table 1). When testing for the effect of

time-lagged ONI on young in isolation (model E1; suppl. info. Table S4 for parameter

estimates) the model was only weakly supported, suggesting a coupled interaction with

a temporal trend. The model which included temporal trends for both age-classes and

ONI carry-over effect on young survival included several non-estimable parameters and

was not supported. No models including the NAO (North Atlantic Oscillation) during

breeding season as a covariate were supported (Table 1; Group E).

The live-only CJS model corroborated results from the joint Burnham model, firstly

with a similar pattern in apparent survival in the basic model. The highest ranked model

was the same: a trend in adult survival and time-lagged ONI effect on young (suppl.

info. Table S5 & S6). Model ranking was equivalent when using a ΔAIC of <2 to

indicate equal support (suppl. info. Table S5). Because of the limited number of live

young in the CJS model, we focus on results from the Burnham model.

4. DISCUSSION

Survival rates and productivity are important parameters in population dynamics and, in

long-lived species, survival probability is the greatest contributing factor (Lebreton &

Clobert 1991). The numerous ecological factors that are acting upon survival rates, and

the tendency for survival to be buffered against environmental stochasticity in long-

lived species, complicate determining the driving factors. However, survival rates of

seabirds are known to be affected by climatic events, and increased variation in

environmental conditions leads to increased variation in vital rates (Morris et al. 2011).

In our study of Arctic skuas breeding on the Faroe Islands, a declining trend in survival

of adult birds with a possible decline in survival of young birds (1^{st} year) coupled with

a time-lagged/carry-over effect of ENSO best explained the variation in the data. These

findings may in part explain the notable decline of the Arctic skua population in the

Faroe Islands in recent times, and should be considered for other seabird species

experiencing similar declines.

As K-selected species, many seabirds share the life history trait of high annual survival

for adults (Szostek & Becker 2015). Overall high survival probabilities of this age-class

are commensurate with reported values and are expected for Arctic skuas (O'Donald

1983, Phillips 2001, Davis et al. 2005, van Bemmelen et al. 2021). We found strong

support for a negative trend over the 24-year study period. Because annual survival of

adults is important for maintaining a stable population, a decrease of 17% in survival

probability in this timeframe is certainly biologically significant to population

persistence (van Bemmelen et al. 2021). Due to their demographic strategy, such a

significant decline in adult survival is both unusual and alarming. In general, in long-

lived vertebrates, the rate of population growth is most sensitive to adult survival. The

number of Arctic skuas breeding in the Faroe Islands has decreased substantially during

our study period (Santos 2018) and the negative trend in adult survival observed on our

study might explain this decline. While the cause of increased mortality in this

population has yet to be determined, the potential impact on the viability of this

population is concerning. In adults, we found no support for the influence of large-scale

oceanic indices on their survival, either during the breeding season or non-breeding

period. This may be due to the masking of any effect by the strong downward trend, or

be due to an environmental variable trending over the duration of the study period (such

as air/sea surface temperature) as it would be difficult to resolve drivers of the decline

over collinearity (Grosbois et al. 2008), or limitations in the data. As such, exploration

of this pattern demands further investigation, aided in part by our improved

understanding of the spatial ecology of this population and species. The closest

neighbouring population, in the UK, is also suffering heavy population declines (Jones

et al. 2008, Perkins et al. 2018). Here, studies aimed at understanding these population

declines have focussed mainly on explaining population dynamics by the variation in

reproductive success (Perkins et al. 2018), but have left open the possibility that

population declines were (also) driven by reduced survival rates due to factors outside

the breeding period. Furthermore, Perkins et al. (2018) demonstrated that low breeding

success affects the stability of the population. The breeding success of Arctic skuas in

the Faroe Islands has received no formal analysis to date, but sharp declines of the

population are expected during our study period if productivity declined here also,

together with the decreases in adult survival observed in our study.

ENSO has been linked with survival in various seabird species (Boersma 1998, England

2000, Champagnon et al. 2018). These examples, however, only assessed chick survival

from birth to fledging. In the Kiritimi Island Black noddy (*Anous minutos*) populations,

no nestlings survived during an El Niño year, when heavy rains flooded the nests, and in

the Galapagos archipelago reduced chick size and survival was reported in three species

during an El Niño year (Jaksic & Fariña 2010). While we found no direct effect of

ENSO on adult survival, we did find some support for an influence of ENSO conditions

(ONI) from the previous year on first year survival (specifically in the period from

ringing until the following breeding season). One explanation is a possible carry-over

effect observed on young birds that might be a consequence of the quality of their

parents' non-breeding environment. Alternatively, a time-lagged response may be

attributed to downstream ecological cascades. A time-lagged effect is commonly

reported in relation to life cycles of prey species as the influence of the environment is

propagated up the food chain (Stenseth et al. 2002). As such, the effect of conditions

during the previous year may indicate that survival of young skuas is governed by food

availability. Szostek and Becker (2015) found such a correlation between primary

productivity during winter with recruitment in the following breeding season in

common terns (*Sterna hirundo*). Primary productivity can be used as a proxy for food

availability (Lindeman 1942) and this suggests that food availability during the non-

breeding period is limiting survival and recruitment success in common terns. As ENSO

has effects on primary productivity, ENSO would be expected to correlate with species

vital rates, as indicated in our study. Potentially, in Arctic skuas, as in the case of

common terns, the quality of the environment during the non-breeding period for adult

birds, expressed by food availability, will be reflected on the quality of their young and

their ability to complete their first year. There is likely to be a disparity between the

impact of south Atlantic conditions on locally breeding populations and migratory birds

from thousands of kilometres away. Alternatively, the time-lagged influence of ONI

may result from the direct interaction of a delayed effect of environmental processes in

the Atlantic from ENSO phenomena originating in the Pacific; specifically, increased

wind associated with El Niño events in the previous year around important upwellings

where skuas are known to winter (Iberian and Benguela currents), and increased

primary production, may lead to more profitable foraging (Oviatt et al. 2015, Kim et al.

2023). Although our model support for an effect of the wintering environment was not

strong, we did find some support for a downward trend in survival probability in young

over time (the model including constant survival probability was equally supported with

the temporal trend). The coupled interaction of the stochastic ENSO events masking the

trend over time may explain the limited model support. However, the magnitude of the

decline is substantial (57%), as such the potential biological significance of lowered

recruitment cannot be ignored over and above a potential interaction with ONI and/or

the temporal occurrence of ringing data (see below). In terms of quantifying higher-

level processes, we acknowledge that while the trend is indicated, the magnitude of the

effect is uncertain (i.e., large confidence intervals around the estimate). For such

vulnerable populations, employing alternative analysis, such as integrated population

models, may improve the robustness of estimates (Schaub & Abadi 2011).

We found no evidence of the effects of NAO on survival of either adults or young. This

suggests the lack of a strong influence of oceanic conditions on the survival of both

young and adults during the breeding season. One explanation is that all skuas,

regardless of age spend the majority of their annual cycle in the wintering grounds

rather than the breeding region in the north Atlantic (Hurrell et al. 2001, Stenseth et al.

2003), and so have more limited exposure to the effects of NAO. Furthermore, the

summer season is characterised by clement weather and abundant resources which is

likely to buffer the influence of oceanic processes; in addition, the influence of NAO

effects are regionally specific and in our area the magnitude and direction of the effect

on fish abundance, and hence prey availability is variable (Drinkwater et al. 2003,

Overland et al. 2010, Báez et al. 2021). It is likely that the environmental conditions

during the breeding season (and the influence of NAO) have greater effects on breeding

success than on survival (Thompson & Ollason 2001, Frederiksen et al. 2004,

Frederiksen et al. 2007).

While we made attempts to use the best available dataset to cover the range of

stochastic environmental conditions (in particular intense El Niño and La Niña events)

and this modelling approach takes into account a lack of data in the estimates, the

structure of the ringing and re-encounter data precluded a fully time-dependent model

(year by year probability estimates) for all years of the study period. This limits our

ability to detect an effect of the linear covariates and influences how the results are

interpreted. We conservatively state that the absence of an effect of the environment,

consistent with the demographic buffering hypothesis (Hilde et al. 2020) cannot be

ruled out. While we found a signal of an effect of time-lagged ONI on survival of young

birds, AIC support was not definitive; furthermore, this alone does not indicate the

environmental or ecological mechanisms potentially involved in driving this

relationship.

Not all variation in survival could be attributed to the variables tested in this study. This

may be explained by the complex interactions and regionally specific effects of

stochastic oceanic events, or the inability of these fundamental models to fully capture

nuanced ecological processes (including variation in other sources of mortality such as

predation, hunting etc.). A better understanding of direct, carry-over, time-lagged

effects of the climatic event or time-lagged trophic propagation is needed to describe the

mechanistic processes at play. An exciting knowledge gap, yet to be explored due to the

current lack of technology, is an understanding of the movements of young and

immature birds in comparison to non-breeding movements of adults. Furthermore,

population changes can be investigated with an approach from breeding ecology, which

is outwith the scope of this survival study. Future studies should include the analysis of

the contribution of other demographic parameters as a function of climate variables, to

also help understand how climate influences survival from hatching until the chicks

fledge.

5. CONCLUSIONS

From a conservation perspective, studying population dynamics and understanding the

underlying process is important for targeting effective conservation measures. The

declining trend on annual survival of adult birds observed in our study is an important

component in understanding population declines in the Faroe Islands. Additionally, the

time-lagged effect of ENSO on survival of young birds can inform predictions of

population trends, especially those due to climate change effects and the prediction of

an increase of ENSO events, despite our incomplete understanding of the complex

interactions of ENSO environmental and ecological processes on survival rates. We

highlight the importance of continuing ringing efforts and longitudinal monitoring

programmes, particularly in light of acute conservation threats such as hunting

(BirdLife International 2017) and highly pathogenic strains of avian influenza (Banyard

et al. 2022).

FUNDING

KS, KT and IAMS were supported by the Danish Council for Independent Research

which funded the MATCH project (1323-00048B) and Danish National Research

Foundation supported Center for Macroecology, Evolution and Climate (DNRF96). KS

was supported by Marie Skłodowska-Curie Fellowship (892006 TesSEH) and the

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under

Germany's Excellence Strategy – EXC 2117 – 422037984. RvB was funded by the

Netherlands Organisation for Scientific Research (project number 866.13.005).

ACKNOWLEDGEMENTS

Ringing data is courtesy of the Danish Bird Ringing Centre at the Natural History

Museum of Denmark, University of Copenhagen. We are indebted to Willy Mardel for

his long-term ringing efforts and his help resolving any data uncertainties in archival

records. We gratefully acknowledge the support of Faroes Museum of Natural History

and Leivu Janus Hansen, Sjúrður Hammer, Jón Aldará and Anthony Wetherhill for their

substantial involvement in the GLS tracking project fieldwork. We kindly acknowledge

the landowners on Fugloy for permission to work on their property. Catching and

deploying geolocators was approved in the Faroe Islands by the Copenhagen Bird

Ringing Centre and The National Museum of the Faroe Islands. A pre-print of this work

is available at https:// https://www.biorxiv.org/content/10.1101/2023.11.10.566398v1.

We kindly thank the anonymous reviewers for their comments which greatly improved

the manuscript.

REFERENCES

Akaike H (1973) Information theory and an extension of the maximum likelihood

505   principle Proceedings of the 2nd International Symposium on Informationc,

506   Czaki, Akademiai Kiado, Budapest

507   Báez JC, Gimeno L, Real R (2021) North Atlantic Oscillation and fisheries

508   management during global climate change. Rev Fish Biol Fish 31:319-336

509   Bakken V, Boertmann D, Mosbech A, Olsen B, Petersen A, Strøm H, Goodwin H

510   (2006) Nordic seabird colony databases: results of a Nordic project on seabird

511   breeding colonies in Faroes, Greenland, Iceland, Jan Mayen and Svalbard,

512   Nordic Council of Ministers, Copenhagen

513   Ballerini T, Tavecchia G, Pezzo F, Jenouvrier S, Olmastroni S (2015) Predicting

514   responses of the Adélie penguin population of Edmonson Point to future sea ice

515   changes in the Ross Sea. Frontiers in Ecology and Evolution 3:8

516   Banyard AC, Lean FZ, Robinson C, Howie F and others (2022) Detection of highly

517   pathogenic avian influenza virus H5N1 clade 2.3. 4.4 b in great Skuas: a species

518   of conservation concern in Great Britain. Viruses 14:212

519   Barbraud C, Weimerskirch H (2003) Climate and density shape population dynamics of

520   a marine top predator. Proc Biol Sci 270:2111-2116

521   Bengtson S-A, Bloch D (2003) The Arctic Skua *Stercorarius parasiticus* on the Faeroe

522   Islands: abundance and plumage polymorphism. Dansk Ornitologisk Forenings

523   Tidskrift 97:210-220

524   BirdLife International (2016) IUCN Red List for birds. http://www.birdlife.org

525   (accessed 11 April 2016)

BirdLife International (2017) Summary of National Hunting Regulations: Faroe Islands

http://www.birdlife.org (accessed 31 July 2022)

BirdLife International and Handbook of the Birds of the World (2016) Bird species

distribution maps of the world. Version 6.0. Available at

http://datazone.birdlife.org/species/requestdis

Boano G, Brichetti P, Foschi U (2010) 'La Niña'-driven Atlantic storms affect winter

survival of Mediterranean Cory's Shearwaters. Italian Journal of Zoology

77:460-468

Boersma PD (1998) Population trends of the Galápagos penguin: impacts of El Niño

and La Niña. Condor 100:245-253

Burfield I, van Bommel F (2004) Birds in Europe: population estimates, trends and

conservation status, Birdlife International

Burnell D, Perkins AJ, Newton SF, Bolton M, Tierney TD, Dunn TE (2023) Seabirds

count: A census of breeding seabirds in Britain and Ireland (2015–2021), Lynx

Nature Books, Barcalona

Burnham KP (1993) A theory for combined analysis of ring recovery and recapture

data. Marked individuals in the study of bird population:199-213

Champagnon J, Lebreton JD, Drummond H, Anderson DJ (2018) Pacific Decadal and

El Niño oscillations shape survival of a seabird. Ecology 99:1063-1072

Davis SE, Nager RG, Furness RW (2005) Food availability affects adult survival as

well as breeding success of parasitic jaegers. Ecology 86:1047-1056

Descamps S, Yoccoz NG, Gaillard JM, Gilchrist HG and others (2010) Detecting

population heterogeneity in effects of North Atlantic Oscillations on seabird

body condition: get into the rhythm. Oikos 119:1526-1536

Dias MP, Martin R, Pearmain EJ, Burfield IJ and others (2019) Threats to seabirds: a

global assessment. Biol Conserv 237:525-537

Drinkwater KF, Belgrano A, Borja A, Conversi A and others (2003) The response of

marine ecosystems to climate variability associated with the North Atlantic

Oscillation. Geophysical Monograph-American Geophysical Union 134:211-

234

England MC (2000) A review of bird responses to El Niño-Southern Oscillation

conditions in the Neotropics. Cotinga 13:83-88

Fife DT, Davis SE, Robertson GJ, Gilchrist HG, Stenhouse IJ, Shutler D, Mallory ML

(2018) Correlating tropical climate with survival of an Arctic-breeding, trans-

equatorial migrant seabird. Arctic Science:1-13

Flack A, Aikens EO, Kolzsch A, Nourani E and others (2022) New frontiers in bird

migration research. Curr Biol 32:R1187-R1199

Fosaa AM, Sykes MT, Lawesson JE, Gaard M (2004) Potential effects of climate

change on plant species in the Faroe Islands. Glob Ecol Biogeogr 13:427-437

Franco BC, Defeo O, Piola AR, Barreiro M and others (2020) Climate change impacts

on the atmospheric circulation, ocean, and fisheries in the southwest South

Atlantic Ocean: a review. Climatic Change 162:2359-2377

Frederiksen M, Harris MP, Daunt F, Rothery P, Wanless S (2004) Scale☐dependent

climate signals drive breeding phenology of three seabird species. Glob Change

Biol 10:1214-1221

Frederiksen M, Furness RW, Wanless S (2007) Regional variation in the role of bottom-

up and top-down processes in controlling sandeel abundance in the North Sea.

Marine Ecology Progress Series 337:279-286

Frederiksen M, Daunt F, Harris MP, Wanless S (2008) The demographic impact of

extreme events: stochastic weather drives survival and population dynamics in a

long-lived seabird. J Anim Ecol 77:1020-1029

Furness RW (1987) The Skuas, T & A.D. Poyser, U.K.

Genovart M, Sanz-Aguilar A, Fernandez-Chacon A, Igual JM, Pradel R, Forero MG,

Oro D (2013) Contrasting effects of climatic variability on the demography of a

trans-equatorial migratory seabird. J Anim Ecol 82:121-130

Grosbois V, Gimenez O, Gaillard JM, Pradel R and others (2008) Assessing the impact

of climate variation on survival in vertebrate populations. Biol Rev Camb Philos

Soc 83:357-399

Hammer S, Madsen JJ, Jensen J-K, Pedersen KT, Bloch D, Thorup K (2014) Færøsk

Trækfugleatlas: the Faroese bird migration atlas, Fróðskapur spf.

Harrison XA, Blount JD, Inger R, Norris DR, Bearhop S (2011) Carry□over effects as

drivers of fitness differences in animals. J Anim Ecol 80:4-18

Hilde CH, Gamelon M, Saether BE, Gaillard JM, Yoccoz NG, Pelabon C (2020) The

Demographic Buffering Hypothesis: Evidence and Challenges. Trends Ecol

Evol 35:523-538

Humphries GRW, Möller H (2017) Fortune telling seabirds: sooty shearwaters

(Puffinus griseus) predict shifts in Pacific climate. Mar Biol 164

Hurrell JW, Kushnir Y, Visbeck M (2001) The north Atlantic oscillation. Science

291:603-605

Irons DB, Anker-Nilssen T, Gaston AJ, Byrd GV and others (2008) Fluctuations in

circumpolar seabird populations linked to climate oscillations. Glob Change

Biol 14:1455-1463

Jaksic F, Fariña J (2010) El Niño and the birds: a resource-based interpretation of

climatic forcing in the southeastern pacific. Anales del Instituto de la Patagonia

38:121-140

Jóhannesson T, Jónsson T, Källén E, Kaas E (1995) Climate change scenarios for the

Nordic countries. Clim Res 5:181-195

Jones T, Smith C, Williams E, Ramsay A (2008) Breeding performance and diet of

Great Skuas Stercorarius skua and Parasitic Jaegers (Arctic Skuas) S. parasiticus

on the west coast of Scotland. Bird Study 55:257-266

Kim D, Lee SK, Lopez H, Foltz GR, Wen C, West R, Dunion J (2023) Increase in Cape

Verde hurricanes during Atlantic Nino. Nat Commun 14:3704

Kubelka V, Sandercock BK, Székely T, Freckleton RP (2021) Animal migration to

northern latitudes: environmental changes and increasing threats. Trends Ecol

Evol

Kuletz K, Mallory M, Gilchrist G, Robertson G and others (2017) Seabirds. In:

Secretariat AC (ed) State of the Arctic Marine Biodiversity Report.

Conservation of Arctic Flora and Fauna

Lebreton J, Clobert J (1991) Bird population dynamics, management, and conservation:

the role of mathematical modelling. Bird population studies:105-125

Lebreton J-D, Burnham KP, Clobert J, Anderson DR (1992) Modeling survival and

testing biological hypotheses using marked animals: a unified approach with

case studies. Ecol Monogr 62:67-118

Lindeman RL (1942) The Trophic-Dynamic Aspect of Ecology. Ecology 23:399-417

Lisovski S, Ramenofsky M, Wingfield JC (2017) Defining the Degree of Seasonality

and its Significance for Future Research. Integr Comp Biol 57:934-942

Meier RE, Votier SC, Wynn RB, Guilford T and others (2017) Tracking, feather moult

and stable isotopes reveal foraging behaviour of a critically endangered seabird

during the non☐breeding season. Diversity and Distributions 23:130-145

Morris WF, Altmann J, Brockman DK, Cords M and others (2011) Low demographic

variability in wild primate populations: fitness impacts of variation, covariation,

and serial correlation in vital rates. Am Nat 177:E14-28

NOAA (2018a).

https://origin.cpc.ncep.noaa.gov/products/analysis_monitoring/ensostuff/ONI_v

5.php. (accessed 1 Sept 2018)

NOAA (2018b). https://www.ncei.noaa.gov/access/monitoring/nao/ (accessed 1 June

2018)

Nourani E, Safi K, de Grissac S, Anderson DJ and others (2023) Seabird morphology

determines operational wind speeds, tolerable maxima, and responses to

extremes. Curr Biol 33:1179-1184. e1173

O'Donald P (1983) The Arctic skua: A study of the ecology and evolution of a seabird.

Cambridge University Press, Cambridge (UK)

Overland JE, Alheit J, Bakun A, Hurrell JW, Mackas DL, Miller AJ (2010) Climate

controls on marine ecosystems and fish populations. J Mar Syst 79:305-315

Oviatt C, Smith L, McManus M, Hyde K (2015) Decadal Patterns of Westerly Winds,

Temperatures, Ocean Gyre Circulations and Fish Abundance: A Review.

Climate 3:833-857

Paiva VH, Pereira J, Ceia FR, Ramos JA (2017) Environmentally driven sexual

segregation in a marine top predator. Sci Rep 7:2590

Paleczny M, Hammill E, Karpouzi V, Pauly D (2015) Population trend of the world's

monitored seabirds, 1950-2010. PloS one 10:e0129342

Perkins A, Ratcliffe N, Suddaby D, Ribbands B and others (2018) Combined bottom-up

and top-down pressures drive catastrophic population declines of Arctic skuas in

Scotland. J Anim Ecol 87:1573-1586

Phillips RA (2001) *Stercorarius parasiticus* Arctic Skua. BWP Update 3:25-41

Polis GA, Hurd SD, Jackson CT, Piñero FS (1997) El Niño effects on the dynamics and

control of anisland ecosystem in the Gulf of California. Ecology 78:1884-1897

Salas EAL, Seamster VA, Boykin KG, Harings NM, Dixon KW (2017) Modeling the

impacts of climate change on Species of Concern (birds) in South Central US

based on bioclimatic variables. AIMS Environmental Science 4:358-385

Santos IAMd (2018) Survival and breeding success of the declining Arctic skua

population of the Faroe Islands., University of Copenhagen

Schaub M, Abadi F (2011) Integrated population models: a novel analysis framework

for deeper insights into population dynamics. Journal of Ornithology 152:227-

237

Silva T, Gislason A, Licandro P, Marteinsdottir G, Astthorsson OS (2013) Environment

induced changes in krill abundance in the North Atlantic Ocean ICES CM

2013/B:82

Stenseth NC, Mysterud A, Ottersen G, Hurrell JW, Chan K-S, Lima M (2002)

Ecological effects of climate fluctuations. Science 297:1292-1296

Stenseth NC, Ottersen G, Hurrell JW, Mysterud A and others (2003) Studying climate

effects on ecology through the use of climate indices: the North Atlantic

Oscillation, El Nino Southern Oscillation and beyond. Proc Biol Sci 270:2087-

2096

Szostek KL, Becker PH (2015) Survival and local recruitment are driven by

environmental carry-over effects from the wintering area in a migratory seabird.

Oecologia 178:643-657

Thompson PM, Ollason JC (2001) Lagged effects of ocean climate change on fulmar

population dynamics. Nature 413:417-420

van Bemmelen RS, Schekkerman H, Hin V, Pot MT and others (2021) Heavy decline of

the largest European Arctic Skua Stercorarius parasiticus colony: interacting

effects of food shortage and predation. Bird Study 68:88-99

van Bemmelen RSA, Moe B, Schekkerman H, Hansen SA and others (2024)

Synchronous timing of return to breeding sites in a long-distance migratory

seabird with ocean-scale variation in migration schedules. Mov Ecol 12:22

Waugh SM, Barbraud C, Adams L, Freeman AND and others (2015) Modeling the

demography and population dynamics of a subtropical seabird, and the influence

of environmental factors. Condor 117:147-164

White GC, Burnham KP (1999) Program MARK: survival estimation from populations

of marked animals. Bird Study 46:S120-S139

FIGURES

Fig. 1. Non-breeding areas of Arctic skuas breeding in Fugloy, Faroe Islands, based on

positional data from 20 GLS tracked birds 2016-7 (van Bemmelen et al., 2024): blue

transparent circles are daily wintering GLS positions and blue circles with black

outlines indicate the centroid of each bird. Birds largely over-winter on the Patagonian

Shelf, Discovery Seamounts, coastally from the Iberian Peninsula to the Cape Verde

Islands, the Gulf of Guinea, the Angola Current and from the Benguela Current

westwards to Tristan da Cunha. The breeding range is given in green and the Faroe

Islands study site is indicated by the dark green circle. Breeding range data from

BirdLife International and Handbook of the Birds of the World (2016); a Robinson

projection was used to create the map.

Fig. 2 a. The Oceanic Niño index (ONI) and b. North Atlantic Oscillation index (NAO).

Lines represents 3-month rolling mean per calendar month. Bars shows the mean

aggregate for period of interest: a. non-breeding period (September to March) and b.

breeding period (May to July), of which values were used as covariates in the survival

models.

Fig. 3. Best supported models describing variation in survival probabilities for (a) adult

and (b-c) young. Plots represent the back transformed (logit link function) derived linear

trend with (a-b) year (Table 1 [model T2]; suppl. info. Table S3) and (c) the time-lagged

oceanic Niño index, ONI(t-1) (Table 1 [model E1]; suppl. info. Table S4), with 95%

confidence intervals. Solid lines indicate strongly supported models, dashed lines are

weakly supported (i.e. there is equal support for no trend: Table 1 [model T1]).

TABLE

Table 1. Annual survival model results for Arctic Skuas, relative to the highest ranked model

[C1] and grouped into temporal trend models [T1-3] and environmental variable models [E1-6].

The basic model is given by [B]. Akaike's information criterion values adjusted for median-$\hat{c}$;

(AICc), differences in AICc (ΔAICc), within-group ΔAICc, weights, model likelihood, number

of estimable parameters and deviance are given. 2Age represents 2 age-classes. We present the

best supported environmental indices: ONI(t-1) is the time lagged (-1 year) Oceanic Niño index,

NAO is the North Atlantic Oscillation index. The three equally supported best models are

highlighted by bold ΔAICc. Live re-encounter probability (p), dead recovery probability (r) and

fidelity (F) remain constant across models: $p(3Age_{Young=Imm=0})$, $r(3Age_{Imm=0})$, $F(.)$.

	Model	AICc	ΔAICc	Within-group ΔAICc	AICc Weights	Model Likelihood	no. Parameters	Deviance
C1	$2Age + Trend_{Adult} + ONI(t-1)_{Young}$	2383.7	**0.00**	0.00	0.245	1.00	8	1772.72
B	2Age	2387.2	3.56	3.56	0.041	0.17	6	1780.34
	Temporal Trend Models							
T1	$2Age + Trend_{Adult}$	2384.4	**0.70**	0.00	0.173	0.71	7	1775.45
T2	$2Age + Trend_{2Age}$	2385.1	**1.38**	0.68	0.123	0.50	7	1776.13
B	2Age	2387.2	3.56	2.86	0.041	0.17	6	1780.34
T3	$2Age + Trend_{Young}$	2389.0	5.32	4.62	0.017	0.07	7	1780.07
	Environmental Variable Models							
E1	$2Age + ONI(t-1)_{Young}$	2385.9	2.19	0.00	0.082	0.33	7	1776.94
B	2Age	2387.2	3.56	1.37	0.041	0.17	6	1780.34
E2	$2Age + NAO_{Adult}$	2388.3	4.58	2.39	0.025	0.10	7	1779.34
E3	$2Age + ONI(t-1)_{Adult}$	2388.5	4.84	2.65	0.022	0.09	7	1779.59
E4	$2Age + NAO_{Young}$	2389.3	5.59	3.40	0.015	0.06	7	1780.34
E5	$2Age + ONI(t-1)$	2393.8	10.1	7.91	0.001	0.01	6	1786.92
E6	$2Age + NAO$	2397.9	14.2	12.01	0.001	0.01	6	1791.02