Paroxetine-induced transient apoptosis with delayed neurogenesis induces brain remodeling in developing zebrafish

Tomomi Sato[1,2]*, Kaito Saito[3], Tsubasa Oyu[3], Kazuna Yamaguchi[3], Maiko Osawa[4], Yuki Shiko[4], Yohei Kawasaki[4], Tomohiro Kurisaki[1], Sachiko Tsuda[3], Takeshi Kajihara[2], Masabumi Nagashima[1]

[1]Department of Anatomy, School of Medicine, Saitama Medical University, Morohongo 38, Moroyama-machi, Saitama 350-0495, Japan

[2]Department of Obstetrics and Gynecology, School of Medicine, Saitama Medical University, Morohongo 38, Moroyama-machi, Saitama 350-0495, Japan

[3]Division of Life Science, Graduate School of Science and Engineering, Saitama University, 255 Shimo-Okubo, Sakura-ku, Saitama City, Saitama 338-8570, Japan

[4]Department of Biostatics, School of Medicine, Saitama Medical University,

Morohongo 38, Moroyama-machi, Saitama 350-0495, Japan

*Corresponding author

[1]Department of Anatomy, School of Medicine, Saitama Medical University, Morohongo

38, Moroyama-machi, Saitama 350-0495, Japan

Tel: +81-49-276-1148

Fax: +81-49-276-1148

Email: tosato@saitama-med.ac.jp (TS)

Abstract

Autism spectrum disorder (ASD) is a neurodevelopmental condition caused by various genetic and environmental factors. This disorder has the cardinal symptoms including impaired social behavior involving the amygdala. Ingesting antidepressants such as paroxetine in early pregnancy increases the risk of ASD in offspring. However, a comprehensive picture of the underlying pathogenic mechanisms remains elusive. Here we demonstrate that early exposure of zebrafish embryos to paroxetine retards neurogenesis in the optic tectum and the telencephalon that includes a region corresponding to the mammalian amygdala. Paroxetine-treated embryos exhibit impaired growth, with small heads and short body lengths resulting from transient apoptosis. This is reminiscent of the early-onset fetal growth restriction (FGR) associated with ASD. Interestingly, the delayed neurogenesis in the small heads was found to be restored after the cessation of paroxetine. This restoration was accompanied by extended retinotectal projections, suggesting brain-preferential remodeling. Finally, social behavior of the paroxetine-treated fish was examined. Our findings imply a biological mechanism underlying early exposure to paroxetine, a risk factor associated with the pathogenesis of ASD.

Introduction

Autism spectrum disorder (ASD) is a neurodevelopmental disorder diagnosed in

childhood [1]. The cardinal symptoms include impaired social behavior, communication

disorder, and stereotyped repetitive behaviors [2]. To date, a large number of genes have

been identified as related to ASD. These are involved in synaptic functions,

transcriptional regulation, and brain development [1,3]. Furthermore, de novo mutations,

paternal age, maternal infection and inflammation during pregnancy have also been

implicated as causative factors [3-5]. Thus, various genetic and environmental factors

contribute to the impaired brain development and functions observed in ASD [4-6].

Maternal depression and anxiety during pregnancy are also associated with an increased

risk of ASD in offspring [7]. Selective-serotonin reuptake inhibitors (SSRIs), one of the

conventional types of antidepressants, are prescribed to pregnant women based on

risk-benefit considerations [8]. The drugs act on the maternal brain to exert their

therapeutic effects but also pass through the placenta to the fetus [9,10]. Several studies

suggest that maternal use of SSRIs during the first trimester of pregnancy increases the

risk of ASD [11-14]. However, a comprehensive picture of the underlying pathogenic

mechanisms remains elusive.

Fetal growth restriction (FGR) is a common complication during pregnancy and

possibly caused by various maternal, placental, and fetal factors [15]. Early-onset FGR

manifests as a morphologically symmetrical fetus with a small head and body due to

fetal growth failure [16]. The causative factors include genetic abnormalities, congenital

malformation, intrauterine infection, and teratogenic effects [15]. This type of FGR is

also associated with ASD [17-19]. However, the pathogenesis of ASD and the

pathophysiological role of SSRIs in the early-onset FGR is poorly understood.

Serotonin (5-hydroxytryptamine [5-HT]) is a monoamine neurotransmitter involved

in emotional disorders such as depression and anxiety by modulating synaptic

transmission in the brain [20,21]. SSRIs exert their therapeutic effects by inhibiting the

reuptake of 5-HT via serotonin transporter (SERT), to sustain the concentration of

extracellular 5-HT [22,23]. 5-HT also plays an important role in adult neurogenesis by

promoting the self-renewal of neural stem cells in the hippocampus [24]. Moreover,

maternal 5-HT synthesized in the placenta of pregnant mice has been found to

transiently localize in the forebrain of early-stage embryos [25], raising the possibility

that placenta-derived 5-HT regulates early brain development [26-28]. However, the

effects of early-stage SSRI and 5-HT on embryonic brain development and postnatal

behaviors are unclear. Elucidation of the primary causes is essential for early prediction

and prevention of the risk for ASD. While SSRI during early pregnancy increases the

risk of ASD, the primary cause remains ambiguous [11,13,14]. The involvement of

various genetic and environmental factors makes it difficult to comprehensively

understand the causes and consequences for the pathogenesis of ASD [1-4]. Thus, there

is a need to elucidate the primary actions of SSRI on early brain development and the

subsequent effects on social behavior. To address this question, we use zebrafish as a

model of human embryos, because easy access to early embryos in a living and intact

condition enables us to investigate the direct actions on those embryos. In zebrafish,

various models for neurodevelopmental disorders have already been established,

suggesting evolutionary conserved pathophysiological mechanisms between humans

and zebrafish in the disorders [29-31]. We focus on the SSRI, paroxetine, because of its

broad applicability across affective disorders, including major depressive and anxiety

disorders [32]. In this study, we assess the effects of paroxetine and compare those with

the control and 5-HT, on morphology, neurogenesis, projection in the brain during

embryonic, larval stages, and finally on social behavior in juvenile stages.

Results

Amino acids interacting with paroxetine are completely conserved between human and zebrafish SERT

To investigate the effects of paroxetine on brain development in zebrafish embryos, we first examined whether the amino acid sequence of the serotonin transporter (SERT) targeted by paroxetine, is conserved between humans and zebrafish (Fig 1). Three-dimensional structures of paroxetine-binding sites and interacting amino acids in human SERT have already been well-defined by X-ray crystallography (Fig 1A) [33,34]. The entire amino acid sequence was 70% identical between the two species (Fig 1B). All the amino acids that interact with paroxetine at the central site of human SERT were found to be identical between the two species (Fig 1A, B). We then examined the expression of zebrafish SERT proteins by immunohistochemistry (Fig 1C and S1 Fig). SERT proteins were weakly but broadly expressed in the brains at 30 hours post-fertilization (hpf), at the time when pet1-positive serotonergic neurons develop in the raphe nuclei (S1 Fig) [35]. We found that SERT proteins were localized to the apical surfaces of radial glial cells in the ventricular zone at 24 hpf, implying that SERT play some roles in neural stem/progenitor cells.

Fig 1. Amino acids interacting with paroxetine are conserved between human and zebrafish SERT. (A) X-ray structure of the ts3 human SERT complexed with paroxetine at the central site (516X, RCSB Protein Data Bank). Amino acids interacting with paroxetine (light blue), indicated in pink. An amino acid forming hydrophobic cavity, indicated in green. (B) Alignments of amino acid sequences of human and zebrafish SERT (P31645 and NP001035061). Amino acids interacting with paroxetine in the central site are completely identical between human and zebrafish. (C) SLC6A4 (SERT) proteins localized in the apical surface of the ventricular zone of the optic tectum (green, open arrowheads). A scheme of the embryonic brain in a dorsal view (Left). The corresponding region shown in the right at high magnification is indicated (Bold rectangle). A scheme of an RFP-expressing radial glial cell and the SLC6A4 localization is shown (Right).

Early transient paroxetine generates embryos with a short body and a small head which is restored at later larval stages

To clarify the effects of paroxetine on embryonic development, we treated embryos with paroxetine and the effector 5-HT between 26–50 hpf, which roughly corresponds

to the first trimester of human embryos (Fig 2A) [36]. To determine subthreshold

concentrations appropriate for treatment, we treated embryos and calculated survival

rates at different concentrations of paroxetine and 5-HT. Both compounds had no effect

on the survival rate at less than 20 μM (Fig 2B). Then, we measured the head-tail body

length and head area of paroxetine-treated embryos to parallel the indication of human

fetal development estimated by crown-lump length (CRL) between weeks 8–11 of the

first trimester in human pregnancy [37]. At first, we monitored the typical development

of wild-type embryos. In these experiments, body lengths and head areas were gradually

but significantly greater than those embryos at 50 hpf according to developmental

stages (S2 Fig). We found that embryos treated with 10 μM paroxetine exhibited

restricted growth with a slight but significant decrease in body length and head area at

50 hpf, whereas no significant decreases were detected in embryos treated with 1 μM

paroxetine (Fig 2C-E). This closely resembled early-onset FGR of human embryos in

the first trimester [16]. In contrast, 5-HT did not affect both the body length and head

area of the embryos at the concentrations of 10 μM and 1 μM, indicating that the

observed embryonic growth restriction is a paroxetine-selective action (Fig 2C-E).

Embryos treated with 1 μM paroxetine for two days exhibited significantly shorter body

lengths and smaller head areas at 73 hpf (S3 Fig). In addition, treatment with 10 μM

paroxetine for one day had a similar effect on larval morphology at 3 dpf, suggesting

that transient early-stage treatment is sufficient for paroxetine to exert its

growth-inhibiting effects (S3 Fig). Therefore, we investigated the effects of early

transient paroxetine on later development. 5-dpf larvae treated with 10 µM paroxetine

during 26–50 hpf showed significant decreases in body lengths and head areas

compared to the controls (S4 Fig). Interestingly, the paroxetine-treated larvae exhibited

restoration of their head areas compared to the controls at 7 dpf despite persistent

decreases in their body lengths (Fig 2F-I). This observation implied that

brain-preferential remodeling occurrs between 5–7 dpf.

Fig 2. Early paroxetine induces embryonic growth restriction followed by preferential restoration in the head at a larval stage. (A) A scheme for the time course of treatment. (B) Dose-responses of survival rates for paroxetine- (blue) and 5-HT- (magenta) treated embryos. (C) Lateral views of 50-hpf zebrafish embryos treated with parox, 5-HT, or the control DMSO. (D, E) Quantification of the body lengths (D) and head areas (E) of the treated embryos. A representative image of an embryo indicating the quantified body length (black line, D) and head area (dotted line, E) are shown in the bottom. (F) Scheme for the treatment schedule and major events in

zebrafish development. (G) Lateral views of 7-dpf larvae treated with parox, 5-HT, or

DMSO. (H, I) Quantification of the body lengths (H) and head areas (I) of the treated

embryos. A representative image of a larva indicating the quantified body length (black

line, H) and head area (dotted line, I) in the bottom.

Paroxetine but not 5-HT transiently induces local apoptosis in the head and tail tip

To elucidate the mechanisms underlying the different effects of paroxetine and 5-HT

on embryonic growth and development, we examined the induction of apoptosis at 31

hpf, 5 hours after treatment with paroxetine or 5-HT by immunostaining with an

antibody against active Caspase-3 (Fig 3A, B). We found a significant increase in the

immunofluorescent signals of active Caspase-3 in the whole brain of paroxetine-treated

embryos compared to the control dimethyl sulfoxide (DMSO)-treated embryos (Fig 3B,

D, parox). The apoptosis induced by paroxetine was increased in a dose-dependent

manner (Fig 3E). In contrast, no such increase was detected in 5-HT-treated embryos

(Fig 3B, D, 5-HT). This is consistent with our observations that 5-HT had no effects on

the head area (Fig 2C, E). Co-treatment of paroxetine with 5-HT also increased

apoptosis in the brain, suggesting that the apoptosis is directly induced by paroxetine

blocking the incorporation of 5-HT via SERT, but not through extracellular 5-HT (Fig

3F). Moreover, in addition to the apoptosis in the whole brain, apoptosis was also

induced locally in the tail tip of paroxetine-treated embryos in an almost all-or-nothing

manner (Fig 3C, G). Consistent with paroxetine-treated embryos, *slc6a4a*

(*serta*)-knockdown embryos also exhibited growth restriction in the body length and

increased apoptosis in the brain were observed in (S1 Fig). These results indicate that

the increased apoptosis was caused by reduced 5-HT transportation due to the inhibition

or knockdown of SERT. Thus. the local apoptosis in the brain and the tail tip

corresponded to the growth restriction of the paroxetine-treated embryos with decreased

head area and body length (Fig 2C-E). Next, we examined whether apoptosis

persistently occurred in the paroxetine-treated brain at 48 hpf. A brain-wide decrease of

Caspase-3-positive signals were detected in the paroxetine-treated embryos. Indeed, as

representative brain regions, the signals were significantly decreased in the optic tecta

(OT) and the telencephalons (T) of paroxetine-treated embryos, indicating that

apoptosis acutely induced after the treatment was suppressed in the brains at later stages

(Fig 3H-J).

Fig 3. Paroxetine transiently induces local apoptosis in the brain and tail tip of

treated embryos. (A) A scheme for the time course of treatment. (B) Lateral views of

the heads of 31-hpf embryos treated with DMSO, parox, or 5-HT and immunostained

for active Caspase-3 (white puncta). Images binarized above a threshold. Dotted line:

Morphological outlines. (C) Lateral views of the tail tips of 33-hpf embryos treated with

DMSO, parox, or 5-HT and immunostained for active Caspase-3. Anterior, left;

posterior, right. (D-G) Quantification of the mean intensities of active Caspase-3 signals

in the heads (D-F) and tail tips (G). (H) Lateral views of DMSO- or parox-treated

embryos immunostained for active Caspase-3 (white puncta) at 48 hpf. Images

binarized above a threshold. Dotted line: Morphological outlines. (I, J) Quantification of

the mean intensities of active Caspase-3 in the optic tecta (I) and telencephalons (J) of

the treated embryos. Scale bars: 100 μm (B, C, H). Data are presented as mean ± SEM.

*$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$; ns, not significant. Kruskal-Wallis with Dunn's

multiple comparisons tests; parox, paroxetine; 5-HT, serotonin; DMSO,

dimethylsulfoxide; hpf, hours post-fertilization; parox, paroxetine; SEM, standard error

of the mean. OT, optic tectum; T, telencephalon.

Paroxetine decreases post-mitotic neurons, whilst increasing

proliferation of neural progenitor cells in the optic tectum

We then explored the effect of paroxetine on neurogenesis in the optic tectum, since

it has been well defined as a visuomotor center in the zebrafish brain in terms of the

development and function of the neural circuits [31, 38-42]. For this purpose, we used

transgenic lines TgBAC(*neurod1:EGFP*), expressing enhanced green fluorescent

protein (EGFP) in differentiated neurons [43], and Tg(*elavl3:kaede*), expressing Kaede

under a pan-neuronal promoter *elavl3* (*huC*) [44] (Fig 4A and S4 Fig). To evaluate the

effect on neurogenesis by treatment with paroxetine and other SSRIs using intact whole

embryos, we examined fluorescence-positive area and the fluorescence intensity to

establish simple indicators of the number of post-mitotic neurons expressing a

fluorescent protein such as EGFP or Kaede. Previously we have revealed that GFP

intensity in post-mitotic neurons increases with developmental stages, because it is

proportional to the number and the time after differentiation of those neurons [31].

Imaging and relative quantification processes are shown in S4 Fig and Materials and

Methods. We found that the EGFP and Kaede fluorescence in post-mitotic neurons were

significantly decreased in the optic tectum and the telencephalon by the treatment with

paroxetine during 26–50 hpf compared to the control (Fig 4A-C and S4 Fig). Indeed,

the number of EGFP-positive neurons (with intensity above 70) in the optic tectum was

significantly decreased in paroxetine-treated embryos at 50 hpf (Fig 4D), suggesting

that the number of post-mitotic neurons was significantly decreased due to increased

apoptosis by paroxetine (Fig 3B, D-F). Treatment with other SSRIs, fluoxetine,

fluvoxamine and sertraline also showed decreased post-mitotic neurons in

TgBAC(*neurod1:EGFP*) similar to that with paroxetine, indicating that the effects on

post-mitotic neurons were shared with other SSRIs (Fig 4G and data not shown).

Immunohistochemistry for CyclinD1 showed a significant decrease in the protein

expression in the optic tectum of the SSRI-treated embryos (Fig 4H and S4 Fig). We

then investigated the effect of upregulation of extracellular 5-HT by treating embryos

with 5-HT during the same developmental stages. Similar to the paroxetine-treated

embryos, EGFP fluorescence was significantly decreased in the optic tectum of

5-HT-treated embryos compared to the controls (Fig 4E, F), indicating that the effect of

paroxetine on the post-mitotic neurons is recapitulated by 5-HT even in the absence of

induction of apoptosis (Fig 3B, D, F).

In adult neurogenesis of both rodents and humans, SSRIs are known to promote the

proliferation of neural stem cells [24,45]. Then, we examined the proliferation of neural

stem/progenitor cells in the optic tectum by immunostaining with anti-phospho-histone

H3 (pH3) antibody to label mitotic cells [31]. We found that paroxetine-treated embryos

showed an increase in the number of pH3-positive mitotic cells at 50 hpf (Fig 4I, J). The

increase was selectively observed in the subventricular zone (SVZ) where cell divisions

of intermediate progenitor cells occur for neurogenesis and proliferation (Fig 4I, K, L)

[46]. Expression of SOX2, a high-mobility group (HMG)-box transcription factor to

suppress cell differentiation and promote self-renewal in proliferating neural

stem/progenitor cells, was enhanced in paroxetine-treated embryonic brains, with a

concomitant decrease in Kaede-expressing neurons at 48 hpf, although an increase in

sox2 mRNA expression was not detected by RT-qPCR with whole embryos (S4 Fig)

[47]. These results suggest that paroxetine promotes the proliferation of neural

progenitor cells in the optic tectum.

Fig 4. Paroxetine decreases post-mitotic neurons and promotes proliferation of

neural progenitor cells in the optic tectum. (A) Representative 50-hpf

TgBAC(*neurod1:EGFP*) embryos treated with parox and the control DMSO shown in

lateral views. EGFP-positive region in the optic tectum (dotted ellipse) for

quantification is indicated (arrowhead). (B) Intensity frequency of the EGFP-positive

region in the optic tectum of DMSO- (top) and paroxetine- (bottom) treated embryos

(DMSO, 24.38 $\pm$ 1.359 pixels; Parox, 21.30 $\pm$ 0.882 pixels; each n = 5). (C)

Quantification of the mean EGFP intensity in the optic tectum (dotted ellipse with

arrowheads in A). Unpaired t test. (D) Quantification of the number of EGFP-positive

neurons with intensity above 70 in the optic tectum. (E) Lateral views of 50-hpf

TgBAC(*neurod1:EGFP*) embryos treated with parox, 5-HT and the control DMSO. (F)

Quantification of the mean EGFP intensity in the optic tecta (arrowheads in E) of the

embryos. Kruskal-Wallis with Dunn's multiple comparisons tests. (G) Lateral views of

50-hpf TgBAC(*neurod1:EGFP*) embryos treated with SSRIs (fluoxetine, fluvoxamine)

and the control. (H) Quantification of CyclinD1 protein expression in the optic tectum

treated with paroxetine, fluoxetine, fluvoxamine and the control. (I) High-magnification

images of the optic tecta of zebrafish embryos treated with DMSO or parox and

immunostained for pH3 (red), with DAPI (blue) co-staining for the nuclei. Open

arrowheads, pH3-positive cells in the SVZ. (J-L) Quantification of the number of

pH3-positive cells in the optic tecta (J), the ventricular zones (K), and the subventricular

zones (L). Scale bars: 100 µm (A, E, G); 25 µm (I). Data are presented as mean ± SEM.

p < 0.05, **p* < 0.01, ***p* < 0.001; ****p* < 0.0001; ns, not significant.

Mann-Whitney U test; 5-HT, serotonin; DAPI, 4',6-diamidino-2-phenylindole

fluorescent stain; DMSO, dimethylsulfoxide; e, eye; EGFP, enhanced green fluorescent

protein; H, hindbrain; hpf, hours post-fertilization; OT, optic tectum; parox, paroxetine;

fluox, fluoxetine; fluvox, fluvoxamine; pH3, phospho-histone H3; SEM, standard error

of the mean; SVZ, subventricular zone; T, telencephalon; VZ, ventricular zone.

The development of retinotectal projections is impaired by

early transient paroxetine and later restored with extended

retinal axons

To investigate whether the decreased neurons in the brain of paroxetine-treated

embryos impacts later neural circuit formation, we examined the retinotectal projections.

We used transgenic line Tg(*pou4f1-hsp70l:GFP*) expressing GFP in excitatory tectal

neurons and retinal ganglion cells (RGCs), and line Tg(*pou4f3:GAL4;UAS:mCherry*)

expressing mCherry in RGCs [43,44]. In wild-type larvae, GFP- and

mCherry-expressing areas gradually and significantly increased with developmental

stages in the optic tecta compared with those areas at 72 hpf (S2 Fig). In contrast, the

fluorescence of both the GFP and mCherry-expressing areas was significantly decreased

between 5 and 7 days post fertilization (dpf) during which activity-dependent

competition followed by dynamic synapse formation or elimination occurs in retinal

axons and tectal dendrites, respectively (S2 Fig) [38,39]. Treatment with 1 µM

paroxetine for two days caused significant decreases in both the area and the

fluorescence of the region expressing GFP and mCherry in the optic tectum at 73 hpf

(Fig 5A-E). One-day treatment with 10 µM paroxetine also showed a similar effect on

the GFP-expression region (S3 Fig). Thus, transient early-stage treatment of paroxetine

is responsible for suppressing the development of the retinotectal projections, consistent

with the growth-inhibiting effects on larval morphology (S3 Fig). Therefore, we

examined the effects of the early transient paroxetine on later development of the

retinotectal projections. In accordance with the restoration of the head area at 7 dpf (Fig

2G, I), paroxetine-treated 7-dpf larvae exhibited restoration of both the area and the

fluorescence of the GFP-expressing region and the fluorescence of the

mCherry-expressing region in the optic tectum by reverting the values to those in the

controls (Fig 5F-H, J). Furthermore, the areas of mCherry-expressing regions were

significantly expanded in paroxetine-treated larvae compared to the controls (Fig 5I,

parox). This expansion was recapitulated in 5-HT-treated larvae (Fig 5I, 5-HT),

suggesting that early transient paroxetine drives the extension of retinal axons through

5-HT at later neurodevelopmental stages.

Fig 5. Early transient paroxetine inhibits the development of the retinotectal

projection leading to brain-preferential remodeling. (A) Lateral views of the heads

of 73-hpf Tg(*pou4fl-hsp70l:GFP; pou4f3:GAL4;UAS:mCherry*) embryos treated with

DMSO or parox at 1 µM for 2 days from 26 hpf. Tectal neurons and retinal axons

express GFP (green) and retinal axons express mCherry (magenta). (B) Quantification

of the GFP-positive areas. (C) Mean GFP intensities in the optic tecta of the transgenic

embryos. (D) Quantification of the mCherry-positive areas. (E) Mean mCherry

intensities in the optic tecta of the transgenic embryos. A representative image

indicating the quantified area (dotted line) in the optic tectum are shown (B, D, top

insets). Mann-Whitney U test. (F) Lateral views of the heads of 7-dpf

Tg(*pou4fl-hsp70l:GFP;pou4f3:GAL4;UAS:mCherry*) larvae treated with parox, 5-HT,

or the control DMSO. Optic tecta (filled arrowheads) and retinal axons (open

arrowheads). (G-J) Quantification of the GFP-positive areas (G), the mean GFP

intensities (H), the mCherry-positive areas (I), and the mean mCherry intensities (J) in

the optic tecta of the transgenic larvae. Kruskal-Wallis with Dunn's multiple

comparisons tests; Scale bar, 100 µm. Data are presented as mean ± SEM. *$p < 0.05$,

**$p < 0.01$; ns, not significant. 5-HT, serotonin; DMSO, dimethylsulfoxide; GFP, green

fluorescent protein; hpf, hours post-fertilization; parox, paroxetine; SEM, standard error

of the mean.

Early paroxetine negatively impacting neurogenesis in the

telencephalon

Next, we investigated whether the early paroxetine treatment affects the

development of the telencephalon that includes a region corresponding to the

mammalian amygdala [29]. The amygdala plays a critical role in social behavior.

Amygdala dysfunction causes impaired social behavior characteristic of ASD [2]. To

assess the effects of paroxetine, SAGFF120A gene-trap line was used, because the line

locally expresses Gal4FF in neurons of the telencephalon which are essential to active

avoidance fear conditioning [49]. Fluorescence of the GFP-expressing areas was

significantly decreased in the telencephalons of paroxetine- and 5-HT-treated embryos

compared to the control at 76 hpf (parox, $p = 0.0279$; 5-HT, $p = 0.0250$, Fig 6B-E) as

observed in the optic tecta, although the decrease was significant for 5-HT but not for

paroxetine at 48 hpf (parox, $p = 0.3344$; 5-HT, $p = 0.0006$, S6 Fig). Consistently, the

GFP fluorescence in the telencephalon was restored at 7 dpf following washout of

paroxetine and 5-HT at 2 dpf (Parox, $p = 0.1419$; 5-HT, $p = 0.6623$; Fig 6C-F). This

restoration concomitantly occurred with that in the head area, corresponding to the

results observed in the larval optic tecta (Fig 2I, Fig 5F, H, J, S6 Fig).

Finally, we explored whether these neurodevelopmental effects of early transient

paroxetine have an impact on juvenile social behavior. This is because no obvious

differences in food intake and feeding behavior were observed between paroxetine-,

5-HT-treated and the control fish. Indeed, the body length of paroxetine-treated fish was

not significantly different from that of the control fish (data not shown). To assess social

behavior of a young fish, we developed a behavioral apparatus using a test arena

separated into three chambers: a test area, a sibling area, and a non-sibling area (Fig 7A;

see Methods). The swimming trajectories of the fish in the test area were recorded

following habituation in the absence (non-sibling trial) and presence (sibling trial) of

siblings in the sibling area (Fig 7B). We examined which area the fish was located in the

sibling (blue) or non-sibling area (red) per frame to determine whether the fish show a

social preference for the sibling frame. This was measured by calculating the social

preference index (SPI, -1 < SPI < 1) (Fig 7C; S1-S6 Videos; see Methods) [50]. Total

swimming distances showed no significant differences between paroxetine-,

5-HT-treated and the control DMSO-treated fishes both in non-sibling and sibling trials

(Fig 7D). The mean SPI difference between the sibling and non-sibling trials tended to

be more negative than the control for both paroxetine and 5-HT, although comparisons

of paroxetine and 5-HT with the control were not significantly different (paroxetine, $p =$

0.103, 5-HT, $p = 0.083$, permutation test) (Fig 7E). The behavioral analysis showed that

the incidence of positive SPI (sibling trial > non-sibling trial, white) was lower in the

fish treated with paroxetine and 5-HT than in the control fish, whereas that of negative

SPI (sibling trial < non-sibling trial, black) was increased in the paroxetine- and

5-HT-treated fish compared to the control (Fig 7F; DMSO, n = 13; parox, n = 7; 5-HT, n

= 6). This was also not significantly different in comparison with the control (paroxetine,

p = 0.195, 5-HT, p = 0.096, permutation test). However, these results raised the

possibility of a tendency for negative social preference in those fishes. On the other

hand, we found a difference in the effect on total swimming distance between

paroxetine and 5-HT. The total swimming distances were significantly decreased in

sibling trials of the control and paroxetine-treated fish, but not in 5-HT-treated fish (Fig

7G). Thus, the total swimming distances of the control exhibited negative values for the

differences between sibling and non-sibling trials (Fig 7H). This was observed in

paroxetine-treated fish, whereas the decrease in the distance was abolished and the

difference was significantly increased in the 5-HT-treated fish compared to the control

and paroxetine (control, p = 0.0427; parox, p = 0.0082; Fig 7G, H). These results

indicated a difference in the effects on behavior by early transient exposure between

paroxetine and 5-HT.

Fig 6. Early transient paroxetine exerts suppressive effects on neurogenesis

followed by restoration in the telencephalon. (**A**) A schematic diagram of treatment

with 1 μM paroxetine. (**B**, **C**) Quantification of GFP intensity in the telencephalon of

the paroxetine-treated (**B**) and 5-HT-treated larvae (**C**). (**D**, **E**) Dorsal views of 76-hpf

SAGFF120A;UAS:GFP larvae treated with paroxetine (**D**), 5-HT (**E**), and the control

DMSO. (**F**, **G**) Quantification of the mean GFP intensities in the telencephalons of the

paroxetine-treated (**F**) and 5-HT-treated larvae (**G**). $p = 0.1419$ (**F**), $p = 0.6623$ (**G**),

Mann-Whitney U test. (**H**, **I**) Dorsal views of 7-dpf SAGFF120A;UAS:GFP larvae

treated with paroxetine (**H**), 5-HT (**I**), and the control DMSO. Scale bar, 100 μm. Data

are presented as mean ± SEM. *$p < 0.05$, ***$p < 0.001$; ns, not significant. 5-HT,

serotonin; DMSO, dimethylsulfoxide; GFP, green fluorescent protein; hpf, hours

post-fertilization; parox, paroxetine; SEM, standard error of the mean.

Fig 7. Social behaviors of early transient paroxetine- and 5-HT-treated zebrafish.

(A) (top) A schematic diagram of the social behavioral test. (bottom) A test arena used

in this study. The sibling frames and non-sibling frames are represented in blue and red,

respectively. (B) Representative swimming trajectories of the fish treated with DMSO

(control), parox, or 5-HT in the non-sibling (top) and sibling trials (bottom). The SPIs

indicated below. (C) A total of 3,000 frames recorded over 5 min for the trials shown in

(B) with color-coded according to the frames in (A). (D) Total swimming distances of

DMSO-, Parox- and 5-HT-treated fish in non-sibling (left) and sibling (right) trials. (E)

Differences in SPIs between the sibling and non-sibling trials. Data are presented as

mean ± SEM, Permutation test. (F) Incidences of positive SPI (sibling > non-sibling,

white) and negative SPI (sibling < non-sibling, black) in the fish treated with DMSO,

parox, and 5-HT (n = 13, 7, 6, respectively). (G) Changes in the total swimming

distances between non-sibling and sibling trials for DMSO-, Parox- and 5-HT-treatd fish.

(H) Differences in the total swimming distances of the fish between the sibling and

non-sibling trials. Data are presented as mean ± SEM, $*p < 0.05$, $**p < 0.01$; ns, not

significant. Brown-Forsythe with Dunnett's multiple comparisons tests. 5-HT,

serotonin; ANOVA, analysis of variance; DMSO, dimethylsulfoxide; GFP, green

fluorescent protein; hpf, hours post-fertilization; parox, paroxetine; SEM, standard error

of the mean; SPI, social preference index.

442   Discussion

In this study, we show that early exposure to paroxetine restricts embryonic growth,

resulting in short body lengths and small heads in zebrafish. In humans, early-onset

FGR manifests as symmetrical morphology with a small head and body, and the

causative factors include teratogenic effects [15,16]. Thus, the paroxetine-treated

zebrafish embryos are considered as a model of early-onset FGR in humans. The growth

restriction is induced by paroxetine but not by 5-HT which is incorporated into cells

through SERT, the target protein blocked by paroxetine. Consistently, apoptosis is

locally induced by paroxetine in the brain and the tail tip, but not by 5-HT, suggesting

that apoptosis is the underlying mechanism for the paroxetine-induced embryonic

growth restriction. However, these results imply disparities between the effects of

paroxetine and 5-HT on embryogenesis. SSRIs including paroxetine have already been

reported to induce apoptosis in various malignant cell types, including gliomas,

neuroblastomas, leukemia, lymphomas, and colon tumors, in addition to normal cells

such as myeloid cells, osteoblasts, and osteoclasts [51-54]. These lines of evidence

suggest a shared mechanism of action of SSRIs for the apoptosis induction in

proliferating cells, potentially attributable to the etiology of adverse effects of SSRIs on

embryogenesis. In addition, another SSRI, fluvoxamine, has recently been identified

that have therapeutic potential as an immunomodulator due to its ability to suppress

immune responses which is likely to reduce the risk of deterioration in COVID-19

[55,56]. These pleiotropic effects of SSRIs are reminiscent of those of thalidomide.

Thalidomide also has proapoptotic properties to exert the teratogenic effects on

embryogenesis, and the exposure during early pregnancy has been associated with ASD

in the offspring [57,58]. Furthermore, the thalidomide derivatives are therapeutically

used as immunomodulatory drugs [59,60]. Taken together with those studies, our

findings imply a shared common mechanism underlying teratogenic and therapeutic

effects of SSRI which are related to the pathogenesis of ASD.

Here, we reveal that paroxetine- and 5-HT-treated embryos exhibit decreased

post-mitotic neurons in the optic tectum. The effect of paroxetine is reproduced by the

treatment with other SSRIs, suggesting that those effects are attributable to a shared

property of antidepressants, serotonin re-uptake inhibition. The SSRI-treated embryos

also show a significant decrease in CyclinD1 protein expression. These results imply

that neurogenesis is suppressed by SSRIs, because Cyclin D1 play a role in promoting

neurogenesis as a cell cycle-independent function [71]. Despite those effects on

post-mitotic neurons, the number of pH3-positive mitotic cells are selectively increased

in the subventricular zone. These observations raise the possibility that the proliferation

of neural stem/progenitor cells is promoted by paroxetine and increased 5-HT at the

expense of the generation of neurons from neural stem/progenitor cells during brain

development [24,31,61]. This is supported by increased expression of SOX2, a

transcription factor regulated by the PI3K/AKT signaling pathway that is not only

inevitable for the induction and maintenance of stem cells, but also a marker of

proliferating neural stem/progenitor cells in the brain [62,72]. Our observations are

consistent with the effect of SSRI that increases the number of neural stem/progenitor

cells in the dentate gyrus of the adult mammalian hippocampus through the increased

concentration of 5-HT [24,45,63]. The effect of SSRI is thought to represent the

underlying mechanism of the therapeutic effect on the maternal brain. On the other hand,

our results suggest a biological mechanism that underlie an adverse effect on the

embryonic brain which may be caused by SSRI ingestion during pregnancy [11-14] (S7

Fig). Indeed, we show that apoptosis is transiently induced and thereafter decreased in

the brain, which generates growth restriction of paroxetine-treated embryos. These

observations are consistent with a previous report showing that SERT-knockout mice

exhibit reductions in programmed cell death in the brain [64]. As for the impact of the

adverse effect, this apoptosis in the embryonic brain *per se* may elicit maternal

inflammation through damage-associated molecular patterns (DAMPs) [73]. However,

at present, it remains unclear how paroxetine induces apoptosis. Our results showing a

similar effect of SSRIs on the post-mitotic neurons, that is decreased neurogenesis,

suggest that incorporation of serotonin through SERT will be critical for survival of

neural stem/progenitor cells leading to generation of neurons (S7 Fig).

Although there is a difference in apoptosis induction at early stages between

paroxetine and 5-HT, both of which cause delayed neurogenesis. This implies that the

delayed neurogenesis is a leading cause of the later effects on the retinotectal projection

and social behavior, because both the treatments show a similar tendency for significant

extension and negative SPI, respectively. Intriguingly, we find that the head size is

restored at a later larval stage in early paroxetine-treated embryos, whereas body length

is not, implying brain-preferential remodeling. Likewise, our data support that the

delayed neurogenesis in the optic tectum decreases the areas and the fluorescence in the

retinotectal projection of the paroxetine-treated embryos, which are later restored after

the removal of paroxetine. These observations raise the possibility that remodeling

occurs in neural networks in the larval brains during the restoration processes, because

the remodeling period corresponds to the activity-dependent refinement stages in the

retinotectal projection [38,39]. Indeed, we reveal extended projections of retinal axons

in the restored brain. This phenotype in axonal projections is similar to that observed in

SERT-knockout mice which exhibit impaired segregation of retinal axons both in the

superior colliculus and the dorsal lateral geniculate nucleus, leading to the disruption of

sensory map formation [65,66]. Those extended axons are also observed in the optic

tecum treated with 5-HT. Similarly, the decreased neurogenesis is detected in both the

optic tecta treated with paroxetine and 5-HT. Thus, our findings suggest that decreased

neurogenesis is a cause of extended axonal projections, because those projections are

observed regardless of whether growth restriction and local apoptosis are induced by the

treatments. This provides insight into a biological basis for the neurodevelopmental

disorders associated with early-onset FGR. The decreased neurogenesis followed by the

restoration is also observed in the telencephalon treated with paroxetine and 5-HT using

the gene-trap line SAGFF120A. The extent of the decrease in the fluorescent protein

expression is different in the telencephalon of SAGFF120A from that of

Tg(*elavl3:Kaede*) at 48 hpf. This is likely due to the difference in the population of

neurons labeled by the fluorescent proteins. In the gene-trap line, neurons in the

telencephalon that includes a region corresponding to the mammalian amygdala are

visualized by GFP. These neurons have been identified that play a critical role in active

avoidance fear conditioning [29,49]. Therefore, the restoration in the telencephalon

implies that the treatments cause remodeling similarly to the optic tectum, which affects

the neural circuit function involved in fear conditioning. Indeed, in individuals with

ASD, the number of neurons in the amygdala has been found not to undergo

neurotypical alterations during their development [67].

Finally, we investigate the impacts of delayed neurogenesis on the development of

social behavior. In the control fish, the SPI differences between sibling and non-sibling

trials vary among individuals, and thereby these are no significant difference in

paroxetine- and 5-HT-treated fish as compared with the control. However, the incidence

of negative SPI (sibling > non-sibling trials) is markedly increased in both the treated

fish. No significant differences in the total swimming distances between paroxetine-,

5-HT-treated and the control fishes would exclude the possibility that the changes in

social behavior are due to those in locomotor activity. A significant decrease in sibling

trials of the control and paroxetine-treated fish can be explained by habituation, but not

explain a similar tendency for increased negative SPI in Paroxetine- and 5-HT-treated

fish. This will exclude the possibility of habituation. In addition, our observations of no

obvious defects of the treated fish in food intake and feeding behavior which mainly

depend on the visual system mostly rule out the possibility of visual impairment [74].

Therefore, our observations raise the possibility of a tendency to negatively impact the

development of social behavior by early exposure to paroxetine and 5-HT. However,

these early environmental conditions per se are less likely to be sufficient to have a

significant effect on social behavior, because the effects on SPIs do not show significant

differences. Nevertheless, transient apoptosis with delayed neurogenesis can be at risk

for neurodevelopmental disorders, even when that is induced by other genetic or

environmental factors. Of note, the effects on total swimming distance are significantly

different between paroxetine and 5-HT, implying a difference in the effects on neural

circuit functions. These findings may offer a biological mechanism underlying early risk

factors associated with the pathogenesis of ASD. However, a more detailed analysis is a

prerequisite for fully understanding the difference in the effects of paroxetine and 5-HT

on the structure and function of the relevant neural circuits. In conclusion, our study

provides evidence for the effects of early transient paroxetine and 5-HT on embryonic

brain development and behaviors. Furthermore, this will represent a potential simple

screening system for evaluating risk factors for ASD and neurodevelopmental disorders

affecting social behavior using zebrafish.

Methods
566

Zebrafish maintenance and strains
568

Wild-type and transgenic zebrafish were kept at 28ºC under 14 hours:10 hours

light/dark cycles as described previously [68], in an IWAKI Small fish breeding unit

LAbREED ITS. To obtain transgenic embryos, a wild-type strain Riken WT (RW) was

crossed with transgenic lines as follows: Tg(*elavl3:Kaede*) to visualize post-mitotic

neurons [44]; TgBAC(*neurod1:EGFP*) to visualize differentiated neurons[43];

Tg(*pau4f1-hsp70l:gfp*) to visualize development of post-mitotic excitatory neurons and

retinal axons in the optic tectum [40]; Tg(*pau4f3:GAL4;UAS:mCherry*) to visualize

retinal axons in the optic tectum [48]; and SAGFF120A;UAS:GFP to visualize neurons

in the telencephalon [49]. All embryos and larvae were maintained at 28ºC in E3

embryo medium (5 mM NaCl, 0.17 mM KCl, 0.33 mM MgSO4, 0.33 mM CaCl2), then

E3 medium containing 0.003% 1-phenyl 2-thiourea (PTU) from approximately 8 hpf,

and those media were refreshed daily. Embryos and larvae were removed from their

chorions and used at the stage noted in each experiment. Fish were treated with 0.04%

Tricine (MS-222, Ethyl m-Aminobenzoate Methanesulfonate, 14805-82, Nacalai

Tesque) in the medium for anesthesia, chilled on ice to alleviate suffering, and then

sacrificed by fixation in 4% Paraformaldehyde Phosphate Buffer Solution (09154-14,

Nacalai Tesque) at 4 ºC. All animal experiments for this study were approved by the

Animal Experimentation Committee of Saitama Medical University (Approval No.

4250) and performed in accordance with the Regulation on Animal Experimentation at

Saitama Medical University.

Microinjection

Preparation and injection of antisense morpholino oligonucleotides (MOs) were carried

out as previously described [31]. Briefly, MOs were diluted in $1\times$ Danieau buffer (58

mM NaCl, 0.7 mM KCl, 0.4 mM MgSO4, 0.6 mM Ca(NO3)2, 5.0 mM HEPES, pH 7.6)

and injected into 1 to 2-cell-stage embryos by using a microinjector FemtoJet 5247

(Eppendorf). The sequences and the concentrations of MOs (Gene Tools, OR, USA)

used in this study are as follows: slc6a4a-AMO (splice-blocking antisense MO,

MO1-slc6a4a, ZDB-MRPHLNO-141031-1, ZIRC) [69], 5'-

ACGCACTTACATGCACTTACACATA -3', at 4.178 ng/nl; slc6a4a-5mis (5mis;

5-bp-mismatched control), 5'- ACCCACTTAGATCCAGTTAGACATA -3', at 4.198

ng/nl.

Drug treatments

Preparation of paroxetine and 5-HT solution were carried out as described previously

[47]. Briefly, paroxetine (Mw 365.83, P1977, Tokyo Chemical Industry (TCI), Tokyo,

Japan), fluoxetine (Mw 345.79, F0750, TCI), fluvoxamine (Mw 434.41, F0858, TCI),

sertraline (Mw 342.69, 193-16191, Fujifilm Wako Chemicals, Osaka, Japan) and 5-HT

(Mw 212.68, S0370, TCI) were dissolved in dimethyl sulfoxide (DMSO) (Nacalai

Tesque, 13407-45, Kyoto, Japan) at 10 mM for stock solutions and diluted with E3

medium to adjust the final concentration from 1–20 μM in 1% DMSO. Paroxetine

pharmacokinetics is reported that the maximal blood concentration (C_{max}) and the

maximal time point (T_{max}) are 1.93 ± 1.38 ng/mL and 4.61 ± 1.04 h, respectively, in

adult healthy humans at a single dose of 10 mg paroxetine orally (Paxil Tablet, GSK).

Our treatment with 1–20 μM paroxetine by bath application in E3 medium corresponds

to 0.3748 and 7.4966 μg/mL *in vitro*, and thus approximately 200–3900 times higher

than the C_{max} in the blood. Zebrafish embryos were dechorionated with forceps and

immersed in the solution from 26-27 hpf, and then incubated at 28.5ºC. Most of

embryos treated under these conditions were survived (see Fig 2B), and those embryos

were euthanized for further analyses or after social behavioral analysis.

Immunohistochemistry

621 Whole-mount immunohistochemistry was carried out as previously described [31].

622 Briefly, embryos were fixed with 4% paraformaldehyde/PBS and incubated with

623 primary antibodies at 4ºC overnight, followed by incubation with secondary antibodies

624 against rabbit or goat IgG conjugated with Alexa Flour 488 or 555 (Abcam, Cambridge,

UK) together with DAPI (Thermo Fisher Scientific, MA, USA). The following primary

antibodies were used in this study: anti-active Caspase-3 (rabbit, 1:500 dilution, 559565,

BD Pharmingen, NJ, USA); anti-phospho histone H3 (pH3; 1:500 dilution, 9701, Cell

Signaling, MA, USA); Sox2 antibody (rabbit, 1:500 dilution, GTX124477, GeneTex,

Hsinchu, Taiwan); anti-Serotonin (goat, 1:500 dilution, ab66047, Abcam);

SLC6A4/5-HTT (rabbit, 1:100 dilution, PA5-49572, Thermo Fisher Scientific); Cyclin

D1 antibody [N1C3] (rabbit, 1:500 dilution, GTX108624, GeneTex). Following

immunostaining, phosphate-buffered saline (PBS), the wash solution of embryos was

gradually substituted by 30%, 50% and finally 70% glycerol/PBS for imaging under a

microscope.

RNA extraction and RT-qPCR

Total RNAs were extracted from whole embryos injected with MOs or treated with

drugs using miRNeasy mini prep kit according to the manufacture's protocols (Qiagen,

Hilden, Germany). The total RNAs were used for cDNA synthesis using SuperScript IV

reverse transcriptase (Thermo Fisher Scientific, MA, USA) with oligo dT primers, and

obtained cDNAs were used for qPCR using Power SYBR Green Master Mix (Thermo

Fisher Scientific, MA, USA) with gene specific primers. Primer sequences used in this

study   are: 5'-ATCTCTCCCGCTGAAGAGTTTTATG-3'   (slc6a4a-forward),

5'-GCCAGAGGTCTTGACCCCCTTC-3'   (slc6a4a-reverse),

5'-CGAGTCTAGTTCGAGTCCGC-3'   (sox2-forward),

5'-CGGGCAGGTACATACTGATC-3'   (sox2-rerverse),

5'-TCTGGAGGACTGTAAGAGGTATGC-3'   (rpl13α-forward)   [68],

5'-AGACGCACAATCTTGAGAGCA-3' (rpl13α-reverse) [68]. qPCR analysis was

performed using PikoReal 96 Real-Time PCR system (Thermo Scientific, MA, USA).

The expression levels of slc6a4a and sox2 relative to rpl13α were calculated by the

$2^{-\Delta\Delta ct}$ method.

Microscopy and imaging

Most of individuals were observed in lateral views under a microscopy unless otherwise

noted. Each individual was mounted on a slide glass and adjusted for the left-light body

axis to be parallel to the optical axis under a stereoscopic microscope as shown in S4

Fig. Whole body images were obtained using a stereoscopic microscope Leica MZ10F

equipped with MC170 HD camera. Multicolor fluorescence images of the head and the

tail were acquired using a confocal laser scanning microscope Zeiss LSM700 equipped

with EC Plan-NEOFLUAR 20×/NA 0.5 and LD Plan-NEOFLUAR 40×/NA 0.6

objectives. The acquisition and the analysis of images were performed after validating

our methods using different transgenic lines. To avoid the variance of fluorescence

intensity by image-acquiring conditions, the conditions were initially determined using

the control individuals per experiment, and all images were acquired under the same

conditions. Each confocal image was obtained by compiling approximately 30–40

optical sections at 2.0 µm intervals per hemisphere using a sequential-scanning mode

per each frame with 512×512 frame size and 8-bit depth. As a control, the mode of the

fluorescence intensity frequency distribution within the ROI set in the hindbrain (see S4

Fig, B, C), which was 20, was regarded as the background level and removed from the

compiled images. The same image processing was applied to all the images obtained by

each experiment.

Social behavioral analysis

A test arena (68 mm length $\times$ 39 mm width $\times$ 15 mm height) divided into three

compartments (sibling, no sibling areas, 10-mm length on both sides; test area, 38-mm

length in the center) was used to observe social behaviors of young fish (35–37 dpf).

Swimming behaviors in the test arena were recorded by capturing movies using a cMOS

camera (Blackfly S BFS-U3-32S4M, Point Grey) for 5 min at 50 frames per sec (fps).

Each experiment was performed in order of five steps as follows: (1) A fish was

transferred to the test area and habituated for 5 min. (2) Without siblings, swimming

behavior was recorded for 5 min (no sibling trial). (3) Siblings were transferred to the

sibling area and habituated for 5min. (4) With siblings, swimming behavior was

recorded for 5 min (sibling trial). (5) Captured movies were converted to avi format

using ImageJ software, and tracking analysis of the swimming fish was performed using

UMA Tracker software (Release 15). Social behaviors were analyzed by comparing (2)

with (4) for Social Preference Index (SPI) calculated as follows: SPI = (Sibling frames –

No sibling frames) / Total frames [50]. Sibling frames (blue) were defined as fish

located in a half region on the side of sibling area, while no sibling frames (red) were

defined as fish located in the other region in the test area. A value of SPI varied between

–1 and 1 [50]. SPI difference is positive when SPI in the sibling trial is larger than the

no sibling trial, and negative when SPI in the sibling trial is smaller than the no sibling

trial. Difference in total distance was defined as the value in sibling trial subtracting that

in no sibling trial.

Quantification and statistical analysis

Body length, head area and Kaede, GFP or mCherry-positive areas and intensities were

measured using Adobe Photoshop (2023, Ver. 24.4.1) and ImageJ (Ver. 1.53). Confocal

stacks were converted into 2D images using sum intensity projection. To eliminate

ectopic signals derived from noise, regions exceeding a defined threshold and

comprising contiguous pixels with an area of 3 pixels or more were considered as

foreground. The regions were defined as the fluorescence-positive area, and the area and

fluorescence mean intensity were measured. An average of control values was

calculated, and then each value of individuals was represented as a relative value to the

average. Thus, the average of controls was denoted as 1.0 for every experiment. For

statistical analysis of two or more samples, two-tailed, unpaired nonparametric

Mann-Whitney U test or Kruskal-Wallis with Dunn's multiple comparison tests were

performed, unless otherwise noted, using GraphPad Prism 9 (Ver. 9.5.1). Permutation

test was used to assess differences in SPI between sibling and non-sibling trials. For

each comparison, individual SPI differences were randomly permuted between groups,

and the test statistic was recalculated for each permutation. A total of 1000 permutations

were performed to generate a null distribution. All results were calculated using SAS

version 9.4 for Windows (SAS Institute, Cary, NC, USA).

Acknowledgements

We are grateful to Hiroshi Handa, Keisuke Miyazawa, Masaki Hiramoto (Tokyo Medical University) for valuable discussions and generous support. We thank members of Anatomy and Experimental Animals (Saitama Medical University) and Hiromi Kazama (Tokyo Medical University) for help and technical support. Tg(*elavl3:Kaede*), Tg(*pou4fl-hsp70l:GFP*), Tg(*gfap:Gal4FF;UAS:GFP*) and SAGFF120A were provided from Zebrafish National BioResource Project. Tg(*pou4f3:GAL4;UAS:mCherry*) was kindly provided by Herwig Baier (Max-Planck Institute).

Author contributions

T.S. conceived and designed the experiments. T.S., T.Ku. performed the anatomical and pharmacological experiments, and K.S., T.O. K.Y. performed behavioral experiments. T.S., K.S., S.T. analyzed the data. M.O., Y.S., Y.K. performed permutation test. T.S., S.T., T.Ka., and M.N. wrote and edited the manuscript.

References

1. Sestan N, State MW. Lost in Translation: Traversing the Complex Path from

Genomics to Therapeutics in Autism Spectrum Disorder. Neuron.

2018;100(2):406-23. doi: 10.1016/j.neuron.2018.10.015.

2. Amaral DG, Schumann CM, Nordahl CW. Neuroanatomy of autism. Trends

Neurosci. 2008;31(3):137-45. doi: 10.1016/j.tins.2007.12.005.

3. Iakoucheva LM, Muotri AR, Sebat J. Getting to the Cores of Autism. Cell.

2019;178(6):1287-98. doi: 10.1016/j.cell.2019.07.037.

4. Willsey HR, Willsey AJ, Wang B, State MW. Genomics, convergent neuroscience

and progress in understanding autism spectrum disorder. Nat Rev Neurosci.

2022;23(6):323-41. doi: 10.1038/s41583-022-00576-7.

5. Al-Haddad BJS, Oler E, Armistead B, Elsayed NA, Weinberger DR, Bernier R, et

al. The fetal origins of mental illness. Am J Obstet Gynecol. 2019;221(6):549-62.

doi: 10.1016/j.ajog.2019.06.013.

6. Amgalan A, Andescavage N, Limperopoulos C. Prenatal origins of

neuropsychiatric diseases. Acta Paediatr. 2021;110(6):1741-9. doi:

10.1111/apa.15766.

7. Wiggins LD, Rubenstein E, Daniels J, DiGuiseppi C, Yeargin-Allsopp M, Schieve

LA, et al. A Phenotype of Childhood Autism Is Associated with Preexisting

Maternal Anxiety and Depression. J Abnorm Child Psychol. 2019;47(4):731-40.

doi: 10.1007/s10802-018-0469-8.

8. Howard LM, Molyneaux E, Dennis CL, Rochat T, Stein A, Milgrom J.

Non-psychotic mental disorders in the perinatal period. Lancet.

2014;384(9956):1775-88. doi: 10.1016/S0140-6736(14)61276-9.

9. Hendrick V, Stowe ZN, Altshuler LL, Hwang S, Lee E, Haynes D. Placental

passage of antidepressant medications. Am J Psychiatry. 2003;160(5):993-6. doi:

10.1176/appi.ajp.160.5.993.

10. Ewing G, Tatarchuk Y, Appleby D, Schwartz N, Kim D. Placental transfer of

antidepressant medications: implications for postnatal adaptation syndrome. Clin

Pharmacokinet. 2015;54(4):359-70. doi: 10.1007/s40262-014-0233-3.

11. Croen LA, Grether JK, Yoshida CK, Odouli R, Hendrick V. Antidepressant use

during pregnancy and childhood autism spectrum disorders. Arch Gen Psychiatry.

2011;68(11):1104-12. doi: 10.1001/archgenpsychiatry.2011.73.

12. Velasquez JC, Goeden N, Bonnin A. Placental serotonin: implications for the

developmental effects of SSRIs and maternal depression. Front Cell Neurosci.

2013;7:47. doi: 10.3389/fncel.2013.00047.

13. Millard SJ, Weston-Green K, Newell KA. The effects of maternal antidepressant

use on offspring behaviour and brain development: Implications for risk of

neurodevelopmental disorders. Neurosci Biobehav Rev. 2017;80:743-65. doi:

10.1016/j.neubiorev.2017.06.008.

14. Andalib S, Emamhadi MR, Yousefzadeh-Chabok S, Shakouri SK, Hoilund-Carlsen

PF, Vafaee MS, et al. Maternal SSRI exposure increases the risk of autistic

offspring: A meta-analysis and systematic review. Eur Psychiatry. 2017;45:161-6.

doi: 10.1016/j.eurpsy.2017.06.001.

15. Nardozza LM, Caetano AC, Zamarian AC, Mazzola JB, Silva CP, Marcal VM, et al.

Fetal growth restriction: current knowledge. Arch Gynecol Obstet.

2017;295(5):1061-77. doi: 10.1007/s00404-017-4341-9.

16. Dall'Asta A, Brunelli V, Prefumo F, Frusca T, Lees CC. Early onset fetal growth

restriction. Matern Health Neonatol Perinatol. 2017;3:2. doi:

10.1186/s40748-016-0041-x.

17. Pels A, Knaven OC, Wijnberg-Williams BJ, Eijsermans MJC, Mulder-de Tollenaer

SM, Aarnoudse-Moens CSH, et al. Neurodevelopmental outcomes at five years

after early-onset fetal growth restriction: Analyses in a Dutch subgroup

participating in a European management trial. Eur J Obstet Gynecol Reprod Biol.

2019;234:63-70. doi: 10.1016/j.ejogrb.2018.12.041.

18. Ranzil S, Ellery S, Walker DW, Vaillancourt C, Alfaidy N, Bonnin A, et al.

Disrupted placental serotonin synthetic pathway and increased placental serotonin:

Potential implications in the pathogenesis of human fetal growth restriction.

Placenta. 2019;84:74-83. doi: 10.1016/j.placenta.2019.05.012.

19. Joseph RM, Korzeniewski SJ, Allred EN, O'Shea TM, Heeren T, Frazier JA, et al.

Extremely low gestational age and very low birthweight for gestational age are risk

factors for autism spectrum disorder in a large cohort study of 10-year-old children

born at 23-27 weeks' gestation. Am J Obstet Gynecol. 2017;216(3):304 e1- e16.

doi: 10.1016/j.ajog.2016.11.1009.

20. Gaspar P, Cases O, Maroteaux L. The developmental role of serotonin: news from

mouse molecular genetics. Nat Rev Neurosci. 2003;4(12):1002-12. doi:

10.1038/nrn1256.

21. Pourhamzeh M, Moravej FG, Arabi M, Shahriari E, Mehrabi S, Ward R, et al. The

Roles of Serotonin in Neuropsychiatric Disorders. Cell Mol Neurobiol.

2022;42(6):1671-92. doi: 10.1007/s10571-021-01064-9.

22. Otte C, Gold SM, Penninx BW, Pariante CM, Etkin A, Fava M, et al. Major

depressive   disorder.   Nat   Rev   Dis   Primers.   2016;2:16065.   doi:

10.1038/nrdp.2016.65.

23. Gross C, Zhuang X, Stark K, Ramboz S, Oosting R, Kirby L, et al. Serotonin1A

receptor acts during development to establish normal anxiety-like behaviour in the

adult. Nature. 2002;416(6879):396-400. doi: 10.1038/416396a.

24. Aimone JB, Li Y, Lee SW, Clemenson GD, Deng W, Gage FH. Regulation and

function of adult neurogenesis: from genes to cognition. Physiol Rev.

2014;94(4):991-1026. doi: 10.1152/physrev.00004.2014.

25. Bonnin A, Goeden N, Chen K, Wilson ML, King J, Shih JC, et al. A transient

placental source of serotonin for the fetal forebrain. Nature. 2011;472(7343):347-50.

doi: 10.1038/nature09972.

26. Rosenfeld CS. The placenta-brain-axis. J Neurosci Res. 2021;99(1):271-83. doi:

10.1002/jnr.24603.

27. St-Pierre J, Laurent L, King S, Vaillancourt C. Effects of prenatal maternal stress on

serotonin and fetal development. Placenta. 2016;48 Suppl 1:S66-S71. doi:

10.1016/j.placenta.2015.11.013.

28. Trowbridge S, Narboux-Neme N, Gaspar P. Genetic models of serotonin (5-HT)

depletion: what do they tell us about the developmental role of 5-HT? Anat Rec

(Hoboken). 2011;294(10):1615-23. doi: 10.1002/ar.21248.

29. Rea V, Van Raay TJ. Using Zebrafish to Model Autism Spectrum Disorder: A

Comparison of ASD Risk Genes Between Zebrafish and Their Mammalian

Counterparts. Front Mol Neurosci. 2020;13:575575. doi:

10.3389/fnmol.2020.575575.

30. Sakai C, Ijaz S, Hoffman EJ. Zebrafish Models of Neurodevelopmental Disorders:

Past, Present, and Future. Front Mol Neurosci. 2018;11:294. doi:

10.3389/fnmol.2018.00294.

31. Sato T, Sato F, Kamezaki A, Sakaguchi K, Tanigome R, Kawakami K, et al.

Neuregulin 1 Type II-ErbB Signaling Promotes Cell Divisions Generating Neurons

from Neural Progenitor Cells in the Developing Zebrafish Brain. PLoS One.

2015;10(5):e0127360. doi: 10.1371/journal.pone.0127360.

32. Bourin M, Chue P, Guillon Y. Paroxetine: a review. CNS Drug Rev.

2001;7(1):25-47. doi: 10.1111/j.1527-3458.2001.tb00189.x.

33. Coleman JA, Green EM, Gouaux E. X-ray structures and mechanism of the human

serotonin transporter. Nature. 2016;532(7599):334-9. doi: 10.1038/nature17629.

34. Coleman JA, Navratna V, Antermite D, Yang D, Bull JA, Gouaux E. Chemical and

structural investigation of the paroxetine-human serotonin transporter complex.

Elife. 2020;9. doi: 10.7554/eLife.56427.

35. Lillesaar C, Stigloher C, Tannhauser B, Wullimann MF, Bally-Cuif L. Axonal

projections originating from raphe serotonergic neurons in the developing and adult

zebrafish, Danio rerio, using transgenics to visualize raphe-specific pet1 expression.

J Comp Neurol. 2009;512(2):158-82. doi: 10.1002/cne.21887.

36. Ito T, Handa H. Deciphering the mystery of thalidomide teratogenicity. Congenit

Anom (Kyoto). 2012;52(1):1-7. doi: 10.1111/j.1741-4520.2011.00351.x.

37. Chen X, Liu H, Li Y, Zhang W, Zhou A, Xia W, et al. First-trimester fetal size,

accelerated growth in utero, and child neurodevelopment in a cohort study. BMC

Med. 2024;22(1):181. doi: 10.1186/s12916-024-03390-3.

38. Niell CM, Meyer MP, Smith SJ. In vivo imaging of synapse formation on a

growing dendritic arbor. Nat Neurosci. 2004;7(3):254-60. doi: 10.1038/nn1191.

39. Hua JY, Smear MC, Baier H, Smith SJ. Regulation of axon growth in vivo by

activity-based   competition.   Nature.   2005;434(7036):1022-6.   doi:

10.1038/nature03409.

40. Sato T, Hamaoka T, Aizawa H, Hosoya T, Okamoto H. Genetic single-cell mosaic

analysis implicates ephrinB2 reverse signaling in projections from the posterior

tectum to the hindbrain in zebrafish. J Neurosci. 2007;27(20):5271-9. doi:

10.1523/JNEUROSCI.0883-07.2007.

41. Isa T, Marquez-Legorreta E, Grillner S, Scott EK. The tectum/superior colliculus as

the vertebrate solution for spatial sensory integration and action. Curr Biol.

2021;31(11):R741-R62. Epub 2021/06/09. doi: 10.1016/j.cub.2021.04.001.

42. Xiao T, Baier H. Lamina-specific axonal projections in the zebrafish tectum require

the type IV collagen Dragnet. Nat Neurosci. 2007;10(12):1529-37. Epub

2007/11/06. doi: 10.1038/nn2002.

43. Obholzer N, Wolfson S, Trapani JG, Mo W, Nechiporuk A, Busch-Nentwich E, et al.

Vesicular glutamate transporter 3 is required for synaptic transmission in zebrafish

hair cells. J Neurosci. 2008;28(9):2110-8. doi:

10.1523/JNEUROSCI.5230-07.2008.

44. Sato T, Takahoko M, Okamoto H. HuC:Kaede, a useful tool to label neural

morphologies in networks in vivo. Genesis. 2006;44(3):136-42. doi:

10.1002/gene.20196.

45. Boldrini M, Hen R, Underwood MD, Rosoklija GB, Dwork AJ, Mann JJ, et al.

Hippocampal angiogenesis and progenitor cell proliferation are increased with

antidepressant use in major depression. Biol Psychiatry. 2012;72(7):562-71. doi:

10.1016/j.biopsych.2012.04.024.

46. Greig LC, Woodworth MB, Galazo MJ, Padmanabhan H, Macklis JD. Molecular

logic of neocortical projection neuron specification, development and diversity. Nat

Rev Neurosci. 2013;14(11):755-69. doi: 10.1038/nrn3586.

47. Ando H, Sato T, Ito T, Yamamoto J, Sakamoto S, Nitta N, et al. Cereblon Control of

Zebrafish Brain Size by Regulation of Neural Stem Cell Proliferation. iScience.

2019;15:95-108. doi: 10.1016/j.isci.2019.04.007.

48. Scott EK, Mason L, Arrenberg AB, Ziv L, Gosse NJ, Xiao T, et al. Targeting neural

circuitry in zebrafish using GAL4 enhancer trapping. Nat Methods.

2007;4(4):323-6. doi: 10.1038/nmeth1033.

49. Lal P, Tanabe H, Suster ML, Ailani D, Kotani Y, Muto A, et al. Identification of a

neuronal population in the telencephalon essential for fear conditioning in zebrafish.

BMC Biol. 2018;16(1):45. doi: 10.1186/s12915-018-0502-y.

50. Dreosti E, Lopes G, Kampff AR, Wilson SW. Development of social behavior in

young zebrafish. Front Neural Circuits. 2015;9:39. doi: 10.3389/fncir.2015.00039.

51. Levkovitz Y, Gil-Ad I, Zeldich E, Dayag M, Weizman A. Differential induction of

apoptosis by antidepressants in glioma and neuroblastoma cell lines: evidence for

p-c-Jun, cytochrome c, and caspase-3 involvement. J Mol Neurosci.

2005;27(1):29-42. doi: 10.1385/JMN:27:1:029.

52. Schuster C, Fernbach N, Rix U, Superti-Furga G, Holy M, Freissmuth M, et al.

Selective serotonin reuptake inhibitors--a new modality for the treatment of

lymphoma/leukaemia? Biochem Pharmacol. 2007;74(9):1424-35. doi:

10.1016/j.bcp.2007.07.017.

53. Wei P, Jia H, Li R, Zhang C, Guo S, Wei S, et al. Fluvoxamine prompts the

antitumor immune effect via inhibiting the PD-L1 expression on mice-burdened

colon tumor. Cell Biol Int. 2023;47(2):439-50. doi: 10.1002/cbin.11936.

54. Hodge JM, Wang Y, Berk M, Collier FM, Fernandes TJ, Constable MJ, et al.

Selective serotonin reuptake inhibitors inhibit human osteoclast and osteoblast

formation and function. Biol Psychiatry. 2013;74(1):32-9. doi:

10.1016/j.biopsych.2012.11.003.

55. Reis G, Dos Santos Moreira-Silva EA, Silva DCM, Thabane L, Milagres AC,

Ferreira TS, et al. Effect of early treatment with fluvoxamine on risk of emergency

care and hospitalisation among patients with COVID-19: the TOGETHER

randomised, platform clinical trial. Lancet Glob Health. 2022;10(1):e42-e51. doi:

10.1016/S2214-109X(21)00448-4.

56. Hashimoto Y, Suzuki T, Hashimoto K. Mechanisms of action of fluvoxamine for

COVID-19: a historical review. Mol Psychiatry. 2022;27(4):1898-907. doi:

10.1038/s41380-021-01432-3.

57. Asatsuma-Okumura T, Ando H, De Simone M, Yamamoto J, Sato T, Shimizu N, et

al. p63 is a cereblon substrate involved in thalidomide teratogenicity. Nat Chem

Biol. 2019;15(11):1077-84. doi: 10.1038/s41589-019-0366-7.

58. Miller MT, Stromland K, Ventura L, Johansson M, Bandim JM, Gillberg C. Autism

with ophthalmologic malformations: the plot thickens. Trans Am Ophthalmol Soc.

2004;102:107-20; discussion 20-1.

59. Sato T, Ito T, Handa H. Cereblon-Based Small-Molecule Compounds to Control

Neural Stem Cell Proliferation in Regenerative Medicine. Front Cell Dev Biol.

2021;9:629326. doi: 10.3389/fcell.2021.629326.

60. Ito T, Yamaguchi Y, Handa H. Exploiting ubiquitin ligase cereblon as a target for

small-molecule compounds in medicine and chemical biology. Cell Chem Biol.

2021;28(7):987-99. doi: 10.1016/j.chembiol.2021.04.012.

61. Kriegstein A, Alvarez-Buylla A. The glial nature of embryonic and adult neural

stem   cells.   Annu   Rev   Neurosci.   2009;32:149-84.   doi:

10.1146/annurev.neuro.051508.135600.

62. Suh H, Consiglio A, Ray J, Sawai T, D'Amour KA, Gage FH. In vivo fate analysis

reveals the multipotent and self-renewal capacities of Sox2+ neural stem cells in

the adult hippocampus. Cell Stem Cell. 2007;1(5):515-28. doi:

10.1016/j.stem.2007.09.002.

63. Sahay A, Hen R. Adult hippocampal neurogenesis in depression. Nat Neurosci.

2007;10(9):1110-5. doi: 10.1038/nn1969.

64. Persico AM, Baldi A, Dell'Acqua ML, Moessner R, Murphy DL, Lesch KP, et al.

Reduced programmed cell death in brains of serotonin transporter knockout mice.

Neuroreport. 2003;14(3):341-4. doi: 10.1097/00001756-200303030-00009.

65. Salichon N, Gaspar P, Upton AL, Picaud S, Hanoun N, Hamon M, et al. Excessive

activation of serotonin (5-HT) 1B receptors disrupts the formation of sensory maps

in monoamine oxidase a and 5-ht transporter knock-out mice. J Neurosci.

2001;21(3):884-96. doi: 10.1523/JNEUROSCI.21-03-00884.2001.

66. Upton AL, Ravary A, Salichon N, Moessner R, Lesch KP, Hen R, et al. Lack of

5-HT(1B) receptor and of serotonin transporter have different effects on the

segregation of retinal axons in the lateral geniculate nucleus compared to the

superior colliculus. Neuroscience. 2002;111(3):597-610. doi:

10.1016/s0306-4522(01)00602-9.

67. Avino TA, Barger N, Vargas MV, Carlson EL, Amaral DG, Bauman MD, et al.

Neuron numbers increase in the human amygdala from birth to adulthood, but not

in autism. Proc Natl Acad Sci U S A. 2018;115(14):3710-5. doi:

10.1073/pnas.1801912115.

68. Westerfield M. THE ZEBRAFISH BOOK, 5th ed. A guide for the laboratory use of

zebrafish (Danio rerio). Eugene: University of Oregon Press; 2007.

69. Waugh TA, Horstick E, Hur J, Jackson SW, Davidson AE, Li X, et al. Fluoxetine

prevents dystrophic changes in a zebrafish model of Duchenne muscular dystrophy.

Hum Mol Genet. 2014;23(17):4651-62. doi: 10.1093/hmg/ddu185.

70. Rassier GT, Silveira TLR, Remiao MH, Daneluz LO, Martins AWS, Dellagostin

EN, et al. Evaluation of qPCR reference genes in GH-overexpressing transgenic

zebrafish (Danio rerio). Sci Rep. 2020;10(1):12692. doi:

10.1038/s41598-020-69423-y.

71. Lukaszewicz AI, Anderson DJ. Cyclin D1 promotes neurogenesis in the developing

spinal cord in a cell cycle-independent manner. Proc Natl Acad Sci U S A.

2011;108(28):11632-7. doi: 10.1073/pnas.1106230108.

72. Schaefer T, Steiner R, Lengerke C. SOX2 and p53 Expression Control Converges

in PI3K/AKT Signaling with Versatile Implications for Stemness and Cancer. Int J

Mol Sci. 2020;21(14). doi: 10.3390/ijms21144902.

73. Chen R, Zou J, Liu J, Kang R, Tang D. DAMPs in the immunogenicity of cell death.

Mol Cell. 2025;85(20):3874-89. doi: 10.1016/j.molcel.2025.09.007.

74. Gahtan E, Tanger P, Baier H. Visual prey capture in larval zebrafish is controlled by

identified reticulospinal neurons downstream of the tectum. J Neurosci.

2005;25(40):9294-303. doi: 10.1523/JNEUROSCI.2678-05.2005.

Supporting information

S1 Fig. Distribution and knockdown of zebrafish SLC6A4a. (**A**) Immunostaining

with anti-SERT antibody for 30-hpf SLC6A4a-knockdown (KD) embryos by injection

of slc6a4a antisense MO (AMO) and the control (5mis). (**B**) Quantification of

immunofluorescent signals in SLC6A4a-KD embryos. (**C**) Expression of *slc6a4a*

mRNAs in SLC6A4a-KD embryos by RT-qPCR. (**D**) Lateral views of 32-hpf embryos

injected with antisense slc6a4a-AMO and the control slc6a4a-5mis. White line,

head-tail body length. (**E, F**) Quantification of the body length (**E**) and the head area (**F**)

of the injected embryos. (**G**) Lateral views of 32-hpf Tg(*elavl3:Kaede*) embryos

injected with MOs (top) and those embryos immunostained for active Caspase-3

(bottom). (**H**) Quantification of the intensity of active Caspase-3 signals in the brain of

the injected embryos. Scale bars: 200 µm (**D**); 100 µm (**A**, **G**). Data are presented as

mean ± SEM. *$p < 0.05$, ***$p < 0.001$, Mann-Whitney U test.

S2 Fig. Development of morphology and the retinotectal projection in wild-type

embryos. (**A**) Lateral views of wild-type embryos at 50 hpf, 72 hpf, 5 dpf, and 7 dpf.

White line, head-tail body length. (**B**, **C**) Quantification of the body length (**B**) and the

head area (**C**) of wild-type embryos. (**D**) Lateral views of wild-type

Tg(*pou4f1-hsp70l:GFP;pau4f3:GAL4;UAS:mCherry*) embryos at 72 hpf, 5 dpf, 7 dpf.

GFP (green) expressed in tectal neurons and retinal axons. mCherry (red) expressed in

retinal axons. (**E**, **F**) Quantification of GFP-positive (**E**) and mCherry-positive (**F**) areas

in the optic tectum. (**G**, **H**) Quantification of the intensity of GFP (**G**) and mCherry (**H**)

in the optic tectum. Scale bars: 200 µm (**A**); 100 µm (**D**). Data are presented as mean ±

SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$, ns, not significant.

Kruskal-Wallis with Dunn's multiple comparisons tests.

1000

S3 Fig. Effects of early paroxetine on embryogenesis and neurogenesis at 3 dpf. (**A**)

A schematic diagram of treatment with 1 µM paroxetine. (**B**) Lateral views of 73-hpf

embryos treated with paroxetine and the control DMSO. (**C**, **D**) Quantification of the

body length (**C**) and the head area (**D**) of the treated embryos. (**E**) A schematic diagram

of treatment with 10 µM paroxetine. (**F**) Lateral views of 72-hpf embryos treated with

paroxetine and the control DMSO. (**G**, **H**) Quantification of the body length (**G**) and the

head area (**H**) of the treated embryos. (**I**) Lateral views of 72-hpf

Tg(*pou4f1-hsp70l:GFP*) embryos treated with 10 µM paroxetine and the control DMSO.

(**J**) Quantification of GFP intensity in the optic tectum of the treated embryos. White

lines, head-tail body length. Parox, paroxetine. Scale bars: 200 µm (**B**, **F**); 100 µm (**I**).

Data are presented as mean ± SEM. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p <$

0.0001, Mann-Whitney U test.

1013

S4 Fig. Effects of early paroxetine on neurogenesis. (**A**) Lateral views of a 50-hpf

Tg(*neurod:EGFP*) embryo. Regions and lines for quantification of the intensity are

indicated in the bottom image. (**B**) Intensity profiles of the yellow arrows in (**A**, bottom)

in the optic tectum (green dots) and the hindbrain (gray dots). (**C**) Intensity frequencies

of the regions (white squares in **A**, bottom) in the optic tectum (green dots) and the

hindbrain (gray dots). (**D**) A schematic for the time course of treatment with paroxetine.

(**E**) Lateral views of 50-hpf Tg(*elavl3:Kaede*) embryos treated with 1 µM, 10 µM

paroxetine and the control DMSO. (**F**-**I**) Quantification of Kaede-positive area (**F**, **G**)

and the intensity (**G**, **I**) in the optic tectum (**F**, **G**) and the telencephalon (**H**, **I**) of 10 μM

paroxetine-treated embryos. (**J**) Expression of CyclinD1 proteins in the optic tectum.

Images immunestained for CyclinD1 (red) merged with nuclear staining with DAPI

(top) and binarized for CyclinD1 signals (bottom). (**K**) Lateral views of DMSO- and

paroxetine-treated embryos immunostained for SOX2 (magenta). (**L**, **M**) Quantification

of the expression of SOX2 proteins by immunohistochemistry (**L**), and *sox2* mRNA by

RT-qPCR (**M**). Parox, paroxetine. Cb, cerebellum; e, eye; H, hindbrain; OT, optic

tectum; T, telencephlon. Filled arrowhead, optic tectum; open arrowhead, telencephalon.

Scale bars: 100 μm (**A**, **E**, **K**); 10 μm (**J**). Data are presented as mean ± SEM. *$p < 0.05$,

**$p < 0.01$, Mann-Whitney U test.

S5 Fig. Growth restriction and decreased neurogenesis by early transient

paroxetine at 5 dpf. (**A**) A schematic diagram of treatment with 10 μM paroxetine and

5-HT. (**B**) Lateral views of 5-dpf larvae treated with paroxetine, 5-HT and the control

DMSO. White line, head-tail body length. (**C**, **D**) Quantification of the body length (**C**)

and the head area (**D**) of the treated larvae. (**E**) Lateral views of 5-dpf

Tg(*pou4f1-hsp70l:GFP*) larvae treated with 10 μM paroxetine and the control DMSO.

1039   (**F**, **G**) Quantification of GFP-positive area (**E**) and the intensity (**F**) in the optic tectum

1040   of the treated larvae. Parox, paroxetine. Scale bars: 200 μm (**B**); 100 μm (**E**). Data are

1041   mean ± SEM. *$p < 0.05$, ***$p < 0.001$, ns, not significant. Kruskal-Wallis with Dunn's

1042   multiple comparisons tests (**C**, **D**), Mann-Whitney U test (**F**, **G**).

1043

S6 Fig. Decreased neurogenesis in the telencephalon and restored head

morphology in paroxetine and 5-HT-treated 7-dpf larvae. (**A**) A schematic diagram

of treatment with 10 μM paroxetine and 5-HT. (**B**) Dorsal views of 48-hpf

SAGFF120A;UAS:GFP embryos treated with paroxetine, 5-HT, and the control DMSO.

(**C**) Quantification of the mean GFP intensities in the telencephalons of

SAGFF120A;UAS:GFP embryos. (**D**) A schematic diagram of treatment with 1 μM

paroxetine and 5-HT. (**E**) Lateral views of 7-dpf larvae treated with paroxetine, 5-HT

and the control DMSO. White line, head-tail body length. (**F**, **G**) Quantification of the

body length (**F**) and the head area (**G**) of the treated larvae. Parox, paroxetine. Scale

bars: 100 μm (**B**); 200 μm (**E**). Data are mean ± SEM. **$p < 0.01$, ***$p < 0.001$, ns, not

significant. Kruskal-Wallis with Dunn's multiple comparisons tests.

1055

1056   **S7 Fig. A summary of the results in this study.** The results obtained by treatment with

1057 paroxetine, other SSRIs and knockdown of SERT are shown in light blue, those by

1058 treatment with 5-HT are shown in magenta, and those with both SSRIs and 5-HT are

1059 shown in black on the background in pale blue.

1060

1061 **S1 Video. DMSO-treated fish in a no sibling trial.**

1062

1063 **S2 Video. DMSO-treated fish in a sibling trial.**

1064

1065 **S3 Video. Paroxetine-treated fish in a no sibling trial.**

1066

1067 **S4 Video. Paroxetine-treated fish in a sibling trial.**

1068

1069 **S5 Video. 5-HT-treated fish in a no sibling trial.**

1070

1071 **S6 Video. 5-HT-treated fish in a sibling trial.**

1072

1073

1074

A GFP/DAPI

DMSO

Paroxetine

TgBAC(*neurod1:EGFP*)

B DMSO

Paroxetine

C Relative EGFP mean intensity (a.u.)

D Relative No. of EGFP+ neurons (a.u.)

E DMSO | Paroxetine | 5-HT

EGFP/BF

TgBAC(*neurod1:EGFP*)

F Relative EGFP mean intensity (a.u.)

G DMSO | Fluoxetine | Fluvoxamine

EGFP/BF

TgBAC(*neurod1:EGFP*)

H Relative CyclinD1 expression in the OT (a.u.)

H DMSO | Parox

pH3/DAPI

basal
SVZ
VZ
apical

I No. of pH3-positive cells

J VZ
No. of pH3-positive cells

K SVZ
No. of pH3-positive cells

A GFP/mCherry/BF

DMSO

73 hpf

Paroxetine

Tg(*pou4f1-hsp70l:GFP; pou4f3:GAL4;UAS:mCherry*)

B Optic tectum

GFP

GFP-positive area (a.u.)

DMSO   Parox

*

C

GFP mean intensity (a.u.)

DMSO   Parox

**

D Retinal axons

mCherry

mCherry-positive area (a.u.)

DMSO   Parox

**

E

mCherry mean intensity (a.u.)

DMSO   Parox

**

F

| | DMSO | Paroxetine | 5-HT |

GFP/BF

7 dpf

mCherry/BF

Tg(*pou4f1-hsp70l:GFP;pou4f3:GAL4;UAS:mCherry*)

G

GFP-positive area (a.u.)

ns
ns

DMSO   Parox   5-HT

H

GFP mean intensity (a.u.)

ns
ns

DMSO   Parox   5-HT

I

mCherry-positive area (a.u.)

**
*

DMSO   Parox   5-HT

J

mCherry mean intensity (a.u.)

ns
ns

DMSO   Parox   5-HT