Dynamics of odor-source localization: Insights from real-time odor plume recordings and

head-motion tracking in freely moving mice

Mohammad F. Tariq[1,2], Scott C. Sterrett[1,2], Sidney Moore[2], Lane[2,4], David J. Perkel[3], David H.

Gire[2]

[1]Graduate Program in Neuroscience, [2]Department of Psychology, [3]Departments of Biology &

Otolaryngology, University of Washington, [4]Department of Psychology, Seattle University

Abstract

Animals navigating turbulent odor plumes exhibit a rich variety of behaviors, and employ

efficient strategies to locate odor sources. A growing body of literature has started to probe this

complex task of localizing airborne odor sources in walking mammals to further our understanding

of neural encoding and decoding of naturalistic sensory stimuli. However, correlating the

intermittent olfactory information with behavior has remained a long-standing challenge due to

the stochastic nature of the odor stimulus. We recently reported a method to record real-time

olfactory information available to freely moving mice during odor-guided navigation, hence

overcoming that challenge. Here we combine our odor-recording method with head-motion

tracking to establish correlations between plume encounters and head movements. We show that

mice exhibit robust head-pitch motions in the 5-14Hz range during an odor-guided navigation task,

and that these head motions are modulated by plume encounters. Furthermore, mice orient towards

the odor source upon plume contact. Head motions may thus be an important part of the

sensorimotor behavioral repertoire during naturalistic odor-source localization.

Short Title: Dynamics of odor-source localization in freely moving mice

Introduction

Animals exhibit rich sensorimotor transformations as part of their natural behavior. Studying naturalistic behaviors in rodents [1–3] offers the opportunity to understand the neural circuits involved in processing sensory information for goal-directed behaviors. For example, odor plume-guided navigation [4,5] is a tractable system in this pursuit, but has historically been under-studied due to the inability to experimentally monitor the complex odor plumes [6,7] that rodents encounter during free behavior. Lewis et al. [8] and Ackels et al. [9] have shown that the olfactory bulb of mice can encode the fast dynamics present in a naturalistic plume, but how this neural activity shapes behavior in response to intermittent plume encounters during naturalistic behaviors has not been clearly established.

Vickers and Baker [10] were the first to precisely correlate plume contacts to behavior in freely flying moths. Their seminal work, along with subsequent studies [11–16] revealed robust behavioral sequences exhibited by invertebrates during complex odor-source localization tasks. Additionally, walking flies navigating a turbulent odor plume also exhibit behavioral changes in response to plume encounters [17].

Mammalian responses to plume encounters during odor tracking using airborne plumes have not been characterized. Findley et al. [18] have recently shown head motions to be a key behavioral feature during an odor gradient-dependent search. However, the plume encounter-dependent changes in rodent head motions were not established in their study. In addition, Liao and Kleinfeld [19] have shown alterations in behavioral states of freely moving rats engaged in an exploratory behavior. They, along with Findley et al. [18], have also established important correlations between rodents' sniffing and their head motions. We have previously used metal oxide sensors to record real-time olfactory information during airborne odor-tracking in mice [20].

Here we combine this method with head-mounted accelerometers to study how head motions

change around plume encounters.

Like Findley et al. [18], and Liao and Kleinfeld [19] we establish head motions as a critical

component of the mouse natural behavioral repertoire. We extend upon these findings by

demonstrating that plume encounters in a turbulent setting modulate these head motions.

Specifically, we show that preceding plume contacts mice increase the amplitude of their head-

pitch motions in the active search band (5-14 Hz), with an immediate reduction after contact. In

addition, mice also lower their heads and switch to foraging after plume contacts, with a

corresponding increase in the frequency of the head-pitch motions and decrease in the body angle

with respect to the source. Our results thus suggest that mice are able to gather information about

the relative orientation of the source with each plume contact.

Similar head motions are also involved in visually-guided tasks [21–23], highlighting the

importance of head motions for sensory acquisition and motor guidance [3] during free behavior.

Hence, our work combined with that of Findley et al. [18] establishes the groundwork to study the

neural mechanisms underlying sensorimotor transformations during the naturalistic behavior of

odor-plume tracking in mammals.

Materials and Methods

Animal Housing and Protocol

All animal procedures were approved by the University of Washington Institutional

Animal Care and Use Committee (IACUC protocol #4356-01). Wild type mice of both genders

(n=7; 3 females/4 males), obtained from The Jackson Laboratory (Strain #000664) were used as

test subjects. Mice were maintained on a reverse 12 hrs light schedule (7AM lights off; 7PM lights

on). After surgery, described below, each mouse was housed individually.

Surgery

All the mice were between 19-24 weeks of age at implantation. Each mouse underwent

aseptic surgery for implantation of a 3D printed part, housing the female end of PCB connectors

(Mill-Max 833-93-100-10-001000; 2 rows of 3 pins). The 3D part was affixed to the skull with

cyanoacrylate. A 1-mm screw was also implanted over the caudal skull on top of cerebellum to

anchor the part. The glue was then built around the part and the screw to ensure long-term stability

of the part. The exposed area of the skull was covered with dental cement. After each surgery, the

mouse was monitored for 45 minutes in a bedding and chips free cage until the mouse was mobile

again. The mouse was then housed individually and monitored for 10 days after surgery for post-

surgery complications. Analgesics, Carprofen and topically applied Lidocaine, were administered

before the surgery. Post-surgery carprofen was administered if needed.

Behavioral training and task

After a >2weeks post-surgery period, the mice were water-restricted and trained to

associate the smell of ethanol with water. This was achieved by placing weigh boats containing

water and ethanol-soaked pads taped underneath in the home cage. Each mouse was given

approximately 15 minutes to consume the water from the weigh boat. The behavioral apparatus

and the task design for odor-source localization were as previously described [20,24]. Briefly, the

odor port in the custom-designed arena was placed randomly in an x-y location and the goal for

the mouse was to locate the source of the odor port to receive reward (water droplet). A 1-s tone

of 1 kHz signified the end of the trial. After each trial was completed, the mouse was coaxed back

into the housing cage located outside the arena, and the odor port was repositioned for the next

trial. The order in which the mice were tested was randomized for each day.

The arena was illuminated solely with infrared lights so that the mouse could not see the

port but an overhead camera (Basler acA640-90uc) could record the animal's position. In addition,

white noise was continuously played in the room to mask any acoustic cues. To minimize strain

by the cables, a linear actuator (Progressive Automation PA-15), with a slip-ring commutator

mounted on a small plexiglass sheet, to minimize electrical noise, was installed in the arena. The

actuator motion was manually controlled by an experimenter with a switch, located outside the

arena, based on the location of the mouse within the arena. Each mouse was weighed before water

restriction to establish the baseline weight and was then maintained at a weight >0.85 times the

baseline weight. In addition to the behavioral task, the mice were supplemented with water

following the same protocol as the ethanol-water association, described above, to maintain their

body weight.

Head-mounted odor sensor and accelerometer

A Figaro TGS 2620 sensor, as described previously [20], mounted on a custom-designed

Printed Circuit Board (PCB) was used to record the real-time olfactory information. Also mounted

on the PCB was a 3-axis accelerometer (Analog Devices ADXL 335). The orientation of the

accelerometer with respect to the ground can be seen in Figure 1A. The accelerometer and the

ethanol sensor were respectively powered by 3.3V and 5V outputs from an Arduino (Adafruit),

and shared a common ground. The PCBs were screwed onto a 3D printed part using miniature

screws (McMaster-Carr#96817A704) and the part also contained male PCB pins to connect with

the 3D printed part glued onto the mouse skull. A 30 Gauge, 8 conductor cable (McMaster-

Carr#3890N21) was soldered directly onto the PCB for signal conduction. The outer insulation

jacket was removed from the cable to make it more flexible to facilitate mouse locomotion. The

data were digitized using a National Instruments NI6212 DAQ, and acquired and stored using

LabView scripts. To acclimate the mice with the 3D printed part, a bare part without the PCBs

mounted was connected to the head piece after the odor-localization behavioral task each day. In

addition, before the odor-localization task in the arena, a smaller version of the apparatus was used

to acclimatize the mice with the head-mounted sensors in a bedding-free mouse cage.

Signal Analysis

All the time-series analysis was done prior to analyzing the position data to minimize the

location bias. Furthermore, the training for DeepLabCut (DLC) [25,26], described below, was

conducted by a researcher not involved in the data collection.

Head-mounted Sensors

All analysis code was written in MATLAB R2019a (MathWorks). The ethanol sensor

signal was low-pass filtered (5Hz cutoff) and then deconvolved using a difference-of-two-

exponentials kernel with τ_{Rise} and τ_{Decay} values set at 0.02 and 10s, respectively [20]. For plume-

event detection, the slope of the cumulative sum of the signal around all time points was calculated

for 1-s segments before and after that point. The onset of an event was defined as the time point

when the ratio of the forward and reverse slopes crossed a threshold. The threshold was determined

by pooling data from all of the odor-localization task sessions, and sessions involving the

presentation of a plume without an animal running in the arena. This allowed us to minimize

spurious detections that might result from the wind as temperature changes affect the signal over

longer time scales. Events that were separated by greater than 5s were pooled for the group data.

The accelerometer readings were calibrated before and after running animals by placing

the head piece in various orientations. The voltage recordings from the accelerometer were

converted to units of g by zeroing and scaling appropriately. The recordings were then low pass

filtered (20Hz cutoff), and the derivative of the signals was taken to convert the acceleration to

jerk [27]. The resulting jerk signal was then used for time-frequency analysis via the *cwt*

(Continuous Wavelet Transform) function. For determining when the head was raised, a 1-s

moving median of the Z- and X- axes was calculated. The median signal for the X-axis was then

subtracted from the Z-axis. The data from all the animal trials were then pooled and a three

Gaussian model was then fit to the distribution as shown in Figure 3A using a Gaussian Mixture

Model algorithm (MathWorks File Exchange; Matthew Roughan Commit Date: 2007). This

analysis was based on the work of Liao and Kleinfeld [19]. To determine the dominant frequency

of head-pitch motions, the band with the highest power within the active search (5-14 Hz) was

used. The real trajectories were defined as the time-window around the onset of events (described

above), while random trajectories were obtained by randomly selecting equal time-sized windows

throughout the trial, except for 1-s windows on either side of real events.

Position Data

We used DLC [25,26] to train a deep neural network for mouse position tracking. Trial

videos were cropped to the size of the arena. The neural network was trained with videos from

recorded trials without wire occlusion to reduce external artifacts. Our trained neural network

exhibited a test error of 2.33 pixels, satisfying the reported standard for human level accuracy of

2.7 pixels [25]. We then developed a MATLAB (2022b) package to aggregate and analyze the

position data alongside the existing accelerometer and ethanol sensor datasets. DLC output was

used for angle calculations.

To calculate the body angle, the mid-point between the neck and body coordinates was

determined. Then the angle between the vector connecting the mid-point with the body, and the

vector connecting the mid-point to the port was defined as the body angle with respect to the port.

The direction of the angle was taken as positive if the body to mid-point vector was clockwise to

the vector connecting the mid-point to port, and negative when it was counter-clockwise. The head

angle was defined as the angle between the vector connecting nose with neck, and the vector

obtained by a 180° rotation of the vector connecting neck with body. Hence, an egocentric right

head turn was thus considered negative while a left head turn was considered positive.

To minimize noise due to erroneous DLC labeling, any of the labels with a likelihood less

than 0.9 was set to NaN, removing those points from the calculation. The angles were analyzed

using the MATLAB circular statistics toolbox [28]. In addition to the time criteria (described

above), only events greater than 15 pixels from the source, approximately the length of an adult

mouse, at the time of encounters were pooled together for the group data (206 plume encounters

across 111 trials for the accelerometer data). For the directional analyses, 281 plume encounters

across 137 trials were grouped (additional trials with non-functional accelerometer were included

into this data set).

Results

To understand the plume-encounter dependent changes in mouse head motions during

airborne-odor source localization, we combined our previously published method of recording

real-time plume information [20] with head-motion monitoring using a 3-axis accelerometer.

Concurrent head-motion tracking with real-time plume information during odor-guided

navigation task

Figure 1A shows the 3D part designed to carry the Printed Circuit Board (PCB) with the

ethanol sensor and accelerometer. This device allowed us to record the real-time plume

information experienced by, and head motions of a mouse engaged in odor-guided navigation.

Figure 1. Concurrent head-motion tracking with real-time odor plume monitoring. A) The designed 3D-printed part with the custom designed PCB to monitor the plume information along with an accelerometer. The arrows correspond to the axis of orientation of the accelerometer and the color correspond with panel **C. B)** An individual trajectory of a mouse engaged in plume-tracking to locate the odor source for a water reward. Overlaid on the trajectory is the colormap of the real-time plume signal experienced by the mouse. The black asterisks represent the locations of plume encounters. The red "x" marks the location of the odor source. **C)** (Top) The read-out over time from the Z- and X- axis channels of the accelerometer (red & blue, respectively) from the trajectory shown in **B**. (Middle) The speed of the animal over time during the trial shown in **B**. (Bottom) The plume signal experienced by the mouse during its search. The color-scale goes from dark (low signal) to lighter (higher signal) and corresponds with the colors shown in **B**. Black dashed lines represent the times of plume encounters. **D1)** A 3-s (before and after) window showing the position (left) and time (right) evolution of the signals around the encounter labeled as **D1** in **B** and **C**. **D2)** Same as **D1** but for the encounter labeled as **D2** in panels **B** and **C**.

An example trial showing the spatial trajectory of the mouse is presented in Figure 1B,

while the accelerometer reading, speed, and olfactory information experienced by the mouse

during that trial are presented in Figure 1C. To better visualize the evolution of the signals over

time and space, zoomed-in views around the encounters labeled in Figures 1B and C are presented

in Figure 1D. Hence, this method allowed us to examine how the head motions of mice engaged

in the odor-localization task change with each odor-plume encounter.

Plume encounter-dependent changes in the amplitude of the mouse head-pitch motions

To better characterize how plume encounters affect the head-pitch motions, we calculated

the derivative of the X-axis signal from the accelerometer to convert the acceleration to jerk [27].

Since the X-axis is aligned with the direction of gravity, this channel was the most useful for

studying the up and down (pitch) motions of the mouse head. It was readily apparent that the pitch

motion had many frequency components embedded in the signal, as previously reported

[18,19,27]. We therefore aligned the jerk signals with respect to the onset of plume encounters

shown in Figure 2C, and conducted time-frequency analysis using the CWT of the jerk signal as

shown in Figure 2D. The dominant frequency band was found to be in the 5-14 Hz range, the

frequency band correlated with active exploration in freely moving rodents [19,29]. Interestingly,

this frequency band was also prominent during the non-plume encounter portions as seen in the

randomly selected trajectories (middle panels; Figures 2C-E).

To better characterize how the amplitude of head-pitch motions evolve over time, and in

particular around plume encounters, we examined the power within the active search band (5-14

Hz) over time presented in Figure 2E. Additionally, the power from the example trial presented in

Figures 1B-D is presented in Figure 2B. As seen in Figure 2E, there is a rapid increase in the

amplitude of pitch motions immediately before plume encounters, which decreases rapidly soon

after plume encounters. Thus, mice move their heads up and down more vigorously before plume

encounters, and dampen these motions following an encounter.

Figure 2. Plume encounters are preceded by increased amplitude of head-pitch motion in the active search (5-14 Hz) band, and immediately followed by decreased head-pitch motion. A) Same trajectory with the plume signal overlaid over space (left), and over time (right) as depicted in Figures **1B-D**. **B)** Amplitude of the mouse head-pitch motions in the 5-14 Hz band (see panel **D**) during the search shown in **A** over location (left) and time (right). The black asterisks, and dashed lines respectively represent the locations and times when plume encounters occurred. **C)** Group data showing the heat map of the plume signal in a 5-s window before and after a plume encounter for real (left) and randomly selected portions (middle), while the mean±SEM of all the encounters' and random trajectories' plume signals (Real: blue; Random: red) over time is presented (right). The signals in the left and middle heatmaps are sorted by the distance of the animal from the source when the encounter happened (Top: closest). **D)** A 5-s window (before and after plume encounter) CWT of the derivative of the accelerometer pitch motion showing 5-14 Hz predominant band (demarcated by the white dashed lines), referred to as the active search band, during real (left), and randomly selected trajectories (middle). The color scales are the same for the spectrograms for real and randomly selected data. In addition, mean power over the frequency bands are presented (right). Black dashed lines again demarcate the 5-14Hz band. **E)** The time-evolution of the power in the active search band during real (left) and randomly selected (middle) trajectories. The sorting of the signal is the same as in **C**. Furthermore, the color scale is the same for both heatmaps. Mean±SEM (right) of the power in the active search band over time is also presented.

Figure 3. Plume encounter results in a lowering of the head with a corresponding increase in the frequency of head-pitch motion. A) The difference in the median Z- and X-axes read out from the accelerometer pooled over all trials and all animals. The resulting distribution was then fit with the sum of three Gaussians (black dashed curve). The mean-S.D. (magenta dashed line) of the third Gaussian (magenta curve) was then set as a threshold to classify the head as either raised (values greater than the dashed magenta line) or lowered. **B)** Cumulative histograms of the dominant frequency of head-pitch motion in the active search band during the head-raised (magenta) and head-lowered (cyan) epochs. A leftward shift in the pitch-motion frequency is apparent during head-raised epochs. **C)** Mean ± SEM of the difference of median of Z- and X-axes readings for real (blue) and randomly selected (red) trajectories (same plume encounter events as those presented in Figures **2C-E**). **D)** Mean ± SEM of the dominant frequency of the head-pitch motions around plume encounters for real and randomly selected trajectories.

Mice lower their heads following plume encounters

Liao and Kleinfeld [19] have shown that rats engaged in free exploration switch from rearing to foraging, and that the sniff frequency during the foraging and rearing epochs are different. To test if such switching also occurred during our task, we examined the difference

between the moving medians of the Z- and X-axes. This is an indirect method to look at the relative

angle of the head with respect to the ground.

The distribution of this difference across all animals is presented in Figure 3A. This

distribution was then fit to a sum of three Gaussians (black dashed curve) and the third Gaussian

(magenta) was then used to establish the threshold for raising of the head [19]. To determine the

number of Gaussians, we applied the Bayesian Information Criteria (BIC). The BIC decreased

substantially with three Gaussians, and marginally decreased further with higher numbers so we

chose the three Gaussian model. We next looked at the dominant frequency of head-pitch motions

during epochs when the head was either raised (magenta) or lowered (cyan) as shown in Figure

3B. The leftward shift of the dominant frequency during head-raising is consistent with that

observed by Liao and Kleinfeld [19].

Furthermore, to study how plume encounters affected the position of the head we looked

at this difference measure over time for the trajectories presented in Figures 2C-E. As shown in

Figure 3C, there is an increase followed by a decrease right around plume encounters, suggesting

that the mice stop scanning the air upon plume contacts. This is also accompanied with a

corresponding increase in the dominant frequency of the head pitch motion as shown in Figure

3D. While we cannot correlate these head pitch-motion changes with changes of sniffing frequency

in our paradigm, many groups [18,19,27] have established a strong correlation between sniffing

and rodent head pitch motions in this frequency band.

Mice reduce their body angle with respect to the source upon plume encounters

We next asked if mice can determine the relative location of the source with each plume

encounter. To answer this question, we calculated the body angle with respect to the source (see

Methods). Figure 4A shows the body angles of the mouse over time for the trajectories presented

in Figure 1D. It was apparent that each mouse reduced the variability of its angle with respect to

the source after plume encounter. To further quantify this effect, we calculated the mean angle of

each trajectory 3s before and after the encounter, and then calculated the difference between the

instantaneous angles and the mean angles for that time period. Figure 4B shows the probability

heatmap over time for this angular difference for the real (blue; top) and random (red; bottom)

trajectories. The distribution of the difference angles for the real trajectories becomes narrower

(blue) after plume encounters than before encounters (gray), as can be seen in Figure 4C. No such

change is observed for the randomly selected trajectories (Figure 4C bottom; $p = 0.0002$ real & p

$= 0.6383$ randomly selected trajectories; Paired Wilcoxon signed rank test).

Furthermore, the mean angle over time reduces after plume encounter, as can be seen in

Figure 4D with a corresponding increase in the strength of the resultant vector for real (blue)

trajectories. The mean angle for the randomly selected trajectories (red; Figure 4D top), on the

other hand, hovers around 10° (close to no particular direction preference with respect to the

source). Furthermore, the high vector strength for the random trajectories (red; Figure 4D bottom)

signifies increased proportions of angles randomly distributed within a range of ±90° around the

mean angle.

Figure 4. Mice reduce their body angle with respect to the odor source upon plume encounter. A1) The body angle (see Methods for description) of the mouse with respect to the odor source during the trajectory **D1** shown in Figure **1**. **A2)** Same as **A1** but for the trajectory **D2** in Figure **1**. **B)** The probability heatmap of the angle difference between instantaneous angles and the mean angle 3s before (gray) and after for each real (blue; top) and randomly selected (red; bottom) trajectory over time. **C)** The distribution of the difference in instantaneous angles and the mean angles before (gray) and after plume encounter for real (blue; top) and randomly selected (red; bottom) trajectories. Notice no difference between the before and after distribution for randomly selected trajectories (red; bottom), while the distribution after encounter for real trajectories (blue; top) becomes narrower than the distribution before (gray; top) plume encounter. **D)** The mean angle (top) over time for real (blue) and randomly selected (red) trajectories with the corresponding strength (bottom) of the resultant vector.

Plume encounters reduce mouse head yaw angle variability

Casting crosswind of an odor source has been established as an efficient strategy to locate the plume in the horizontal plane for flying insects [10,12,13]. We therefore looked at the head yaw angle as a correlate for an active sensing strategy for walking mice to locate the plume in the horizontal plane. Figure 5A (top) shows the mean angle over time for the real (blue) and randomly selected (red) trajectories. While the mean angle for the real and randomly selected trajectories do not differ after plume encounters, there is a slight increase in the strength of the resultant vector (bottom) for real trajectories as compared to the random. This increase in strength for the real trajectories suggests that variability around the mean angle decreases peri-plume encounters.

Figure 5. Mice reduce the variability of the yaw angle of head motions after plume encounter during odor-guided search. A) The mean angle (top) over time for real (blue) and randomly selected (red) trajectories with the corresponding strength of the resultant vector (bottom). Notice the increase in the strength peri-plume encounter for the real trajectories. **B)** The probability heatmap over time for left (+ve) and right (-ve) head motions before (grey) and after (blue) plume encounter. **C)** The head angle distribution before (grey) and after (blue) plume encounters. **D)** Raster plots for 0° crossing events for real (blue; left) and randomly selected (red; right) trajectories. **E)** Mean ± SEM of the z-scored 0° crossing rolling rates for the real (blue) and random (red) trajectories.

In addition, the probability heatmap over time before (grey) and after (blue) plume

encounters is presented in Figure 5B, with the distribution of the yaw angles presented in Figure

5C. To further test if this reduction in the yaw angle variability was due to increased rate of yaw

turns, we looked at the number of 0° crossings before and after plume encounters. This measure is

presented as raster plots for the real (blue) and random (red) trajectories in Figure 5D, where each

tick represents a 0° crossing event. Figure 5E shows the mean ± SEM of the z-scored rolling rate

of the 0° crossings for the real (blue) and random (red) trajectories. There is a transient increase in

the rate for the real trajectories right after plume encounter while a decrease in the rate later after

the encounter.

Discussion

Correlating sporadic plume encounters to behavioral changes, and neural recordings during

odor-guided navigation has remained an elusive goal [6,7] due to the constraints in real-time

olfactory monitoring. We previously reported a method to record the odor information available

to a freely moving mouse using head-mounted sensors [20] that has been adopted by other groups

to fit their research goals [30,31]. Here we combined this method with head-motion tracking using

accelerometers to study how plume encounters shape head motions during an odor-localization

task.

We show that mice actively search for olfactory plume encounters in our odor-source

localization task by exhibiting robust head-pitch motions in the 5-14Hz range. These head motions

are increased in their amplitude just before the plume encounter, and robustly decrease right after

the encounter. This, in conjunction with our previously published result of a decrease in speed

after plume encounters [20] suggests that the decrease in the head-pitch motion might be a strategy

to localize the odor source. The vigorous head motions before encounter could be made to catch a

whiff of the plume, while the slowing of the head-pitch motions could be useful to more precisely

decipher the location of the source.

Consistent with this two-stage model for source localization, we find that following plume

encounters mice reduce their angle of travel with respect to the source. This suggests that mice can

extract source orientation information from each plume encounter. We also show that mice lower

their heads after plume encounters, with an increase in the frequency of their head motions. This

lowering of the head is consistent with normative theoretical strategies of switching to sampling

the air vs. sampling closer to the ground for a computational searcher [32].

In addition, we show that the yaw head angles are reduced in their variability after plume

encounters, with a subtle increase in the rate of yaw turns again supporting the idea that mice

change sampling strategy after a plume encounter [33]. While many mammals, including humans,

have been shown to adapt their sensing strategies while following odor trails [18,34–36], here we

show that mice adapt their strategies locked to plume encounters during odor-guided navigation.

Furthermore, the evolution of the change in the different behavioral parameters suggest switching

of behavioral motifs [37].

Together, our observations support a model in which mice engage in search behavior prior

to plume contact, then adjust this behavior upon a plume encounter to robustly extract information

from the plume, enabling mice to estimate source location from a single plume encounter. While

previous models of plume-guided search behavior emphasize the need for repeated plume

sampling over space and time to iteratively estimate source position, the results we present instead

suggest that mice have evolved a robust behavioral strategy, which combined with morphological

and neural adaptations, enables them to extract useful directional cues from even single plume

encounters. Our results suggest that mice, and perhaps all macrosmatic mammals, receive and

process vastly more directional information from odor plumes than previously thought.

Stats Table

Comparison		Figure			
Variable	Time	2E	3C	3D	5E
Real vs. Random	-5 to -4	0.6550	0.4722	0.3144	0.4801
	-4 to -3	0.8454	0.1310	0.2369	0.3794
	-3 to -2	0.8290	0.5793	0.4396	0.3002
	-2 to -1	0.1635	0.1362	0.2357	0.1867
	-1 to 0	0.0104	0.0007	0.1190	0.1154
	0 to 1	0.0235	0.00002	0.0569	0.0292
	1 to 2	0.0657	0.0456	0.0014	0.5007
	2 to 3	0.0660	0.2673	0.1764	0.0890
	3 to 4	0.0088	0.5906	0.0158	0.0091
	4 to 5	0.0155	0.4567	0.0892	0.1866

Acknowledgements

We like to acknowledge the members of the Perkel, and the Gire labs for helpful
comments and discussions on this project.

Funding

This work was supported by the National Institutes of Health Grant R01DC018789 to
D.H.G.; M.F.T. was supported by UW Auditory Neuroscience Training Grant (T32DC005361),
and an NIH NRSA Fellowship (F31DC018442).

Author Contributions

Conceptualization: M.F.T., D.J.P., D.H.G.

Data Curation: M.F.T., L.

Formal analysis: M.F.T., S.C.S., L.

Data Collection: M.F.T., S.M.

Writing – original draft: M.F.T.

Writing – review & editing: M.F.T., S.C.S., L., D.J.P., D.H.G.

References

1. Dennis EJ, Hady AE, Michaiel A, Clemens A, Tervo DRG, Voigts J, et al. Systems Neuroscience of Natural Behaviors in Rodents. J Neurosci. 2021;41: 911–919. doi:10.1523/JNEUROSCI.1877-20.2020

2. Miller CT, Gire D, Hoke K, Huk AC, Kelley D, Leopold DA, et al. Natural behavior is the language of the brain. Current Biology. 2022;32: R482–R493. doi:10.1016/j.cub.2022.03.031

3. Parker PRL, Brown MA, Smear MC, Niell CM. Movement-Related Signals in Sensory Areas: Roles in Natural Behavior. Trends in Neurosciences. 2020;43: 581–595. doi:10.1016/j.tins.2020.05.005

4. Baker KL, Dickinson M, Findley TM, Gire DH, Louis M, Suver MP, et al. Algorithms for Olfactory Search across Species. J Neurosci. 2018;38: 9383. doi:10.1523/JNEUROSCI.1668-18.2018

5. Gaeta G, Sullivan RM, Wilson DA. Neural Mechanisms for Odor-Guided Behavior. Oxford Research Encyclopedia of Neuroscience. Oxford University Press; 2023. doi:10.1093/acrefore/9780190264086.013.359

6. Marin AC, Schaefer AT, Ackels T. Spatial information from the odour environment in mammalian olfaction. Cell Tissue Res. 2021;383: 473–483. doi:10.1007/s00441-020-03395-3

7. Ackels T. Information about space from time: how mammals navigate the odour landscape. Neuroforum. 2022;28: 159–168. doi:10.1515/nf-2022-0006

8. Lewis SM, Xu L, Rigolli N, Tariq MF, Stern M, Seminara A, et al. Plume dynamics structure the spatiotemporal activity of glomerular networks in the mouse olfactory bulb. Front Cell Neurosci. 2021;15. doi:10.3389/fncel.2021.633757

9. Ackels T, Erskine A, Dasgupta D, Marin AC, Warner TPA, Tootoonian S, et al. Fast odour dynamics are encoded in the olfactory system and guide behaviour. Nature. 2021;593: 558–563. doi:10.1038/s41586-021-03514-2

10. Vickers NJ, Baker TC. Reiterative responses to single strands of odor promote sustained upwind flight and odor source location by moths. Proc Natl Acad Sci USA. 1994;91: 5756. doi:10.1073/pnas.91.13.5756

11. Cardé RT, Willis MA. Navigational Strategies Used by Insects to Find Distant, Wind-Borne Sources of Odor. Journal of Chemical Ecology. 2008;34: 854–866. doi:10.1007/s10886-008-9484-5

12. Mafra-Neto A, Cardé RT. Rate of realized interception of pheromone pulses in different wind speeds modulates almond moth orientation. Journal of Comparative Physiology A. 1998;182: 563–572. doi:10.1007/s003590050203

13. Mafra-Neto A, Cardé RT. Fine-scale structure of pheromone plumes modulates upwind
orientation of flying moths. Nature. 1994;369: 142.

14. van Breugel F, Dickinson MH. Plume-Tracking Behavior of Flying Drosophila Emerges
from a Set of Distinct Sensory-Motor Reflexes. Current Biology. 2014;24: 274–286.
doi:10.1016/j.cub.2013.12.023

15. Vickers N. Mechanisms of animal navigation in odor plumes. The Biological Bulletin.
2000;198: 203–212. doi:10.2307/1542524

16. Vickers NJ, Christensen TA, Baker TC, Hildebrand JG. Odour-plume dynamics influence
the brain's olfactory code. Nature. 2001;410: 466.

17. Demir M, Kadakia N, Anderson HD, Clark DA, Emonet T. Walking Drosophila navigate
complex plumes using stochastic decisions biased by the timing of odor encounters. eLife.
2020;9: e57524. doi:10.7554/eLife.57524

18. Findley TM, Wyrick DG, Cramer JL, Brown MA, Holcomb B, Attey R, et al. Sniff-
synchronized, gradient-guided olfactory search by freely moving mice. eLife. 2021;10:
e58523. doi:10.7554/eLife.58523

19. Liao S-M, Kleinfeld D. A change in behavioral state switches the pattern of motor output
that underlies rhythmic head and orofacial movements. Current Biology. 2023;
S0960982223004566. doi:10.1016/j.cub.2023.04.008

20. Tariq MF, Lewis SM, Lowell A, Moore S, Miles JT, Perkel DJ, et al. Using Head-Mounted
Ethanol Sensors to Monitor Olfactory Information and Determine Behavioral Changes
Associated with Ethanol-Plume Contact during Mouse Odor-Guided Navigation. eNeuro.
2021;8. doi:10.1523/ENEURO.0285-20.2020

21. Michaiel AM, Abe ET, Niell CM. Dynamics of gaze control during prey capture in freely
moving mice. eLife. 2020;9: e57458. doi:10.7554/eLife.57458

22. Bouvier G, Senzai Y, Scanziani M. Head Movements Control the Activity of Primary
Visual Cortex in a Luminance-Dependent Manner. Neuron. 2020;108: 500-511.e5.
doi:10.1016/j.neuron.2020.07.004

23. Meyer AF, O'Keefe J, Poort J. Two Distinct Types of Eye-Head Coupling in Freely
Moving Mice. Current Biology. 2020;30: 2116-2130.e6. doi:10.1016/j.cub.2020.04.042

24. Jackson BJ, Fatima GL, Oh S, Gire DH. Many Paths to the Same Goal: Balancing
Exploration and Exploitation during Probabilistic Route Planning. eNeuro. 2020;7:
ENEURO.0536-19.2020. doi:10.1523/ENEURO.0536-19.2020

25. Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW. Using DeepLabCut for 3D
markerless pose estimation across species and behaviors. Nat Protoc. 2019;14: 2152–2176.
doi:10.1038/s41596-019-0176-0

26.   Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, et al. DeepLabCut:
markerless pose estimation of user-defined body parts with deep learning. Nature
Neuroscience. 2018;21: 1281–1289. doi:10.1038/s41593-018-0209-y

27.   Karalis N, Sirota A. Breathing coordinates cortico-hippocampal dynamics in mice during
offline states. Nat Commun. 2022;13: 467. doi:10.1038/s41467-022-28090-5

28.   Berens P. **CircStat** : A *MATLAB* Toolbox for Circular Statistics. J Stat Soft. 2009;31.
doi:10.18637/jss.v031.i10

29.   Rojas-Lãbano D, Frederick DE, Egaña JI, Kay LM. The olfactory bulb theta rhythm
follows all frequencies of diaphragmatic respiration in the freely behaving rat. Front Behav
Neurosci. 2014;8. doi:10.3389/fnbeh.2014.00214

30.   Chen KS, Wu R, Gershow MH, Leifer AM. Continuous odor profile monitoring to study
olfactory navigation in small animals. eLife. 2023;12: e85910. doi:10.7554/eLife.85910

31.   Li Y, Swerdloff M, She T, Rahman A, Sharma N, Shah R, et al. Robust odor identification
in novel olfactory environments in mice. Nat Commun. 2023;14: 673. doi:10.1038/s41467-
023-36346-x

32.   Rigolli N, Reddy G, Seminara A, Vergassola M. Alternation emerges as a multi-modal
strategy for turbulent odor navigation. eLife. 2022;11: e76989. doi:10.7554/eLife.76989

33.   Reddy G, Shraiman BI, Vergassola M. Sector search strategies for odor trail tracking. Proc
Natl Acad Sci USA. 2022;119: e2107431118. doi:10.1073/pnas.2107431118

34.   Khan AG, Sarangi M, Bhalla US. Rats track odour trails accurately using a multi-layered
strategy with near-optimal sampling. Nature Communications. 2012;3: 703.

35.   Catania KC. Stereo and serial sniffing guide navigation to an odour source in a mammal.
Nature Communications. 2013;4: 1441.

36.   Porter J, Craven B, Khan RM, Chang S-J, Kang I, Judkewitz B, et al. Mechanisms of scent-
tracking in humans. Nat Neurosci. 2007;10: 27–29. doi:10.1038/nn1819

37.   Wiltschko AB, Johnson MJ, Iurilli G, Peterson RE, Katon JM, Pashkovski SL, et al.
Mapping Sub-Second Structure in Mouse Behavior. Neuron. 2015;88: 1121–1135.
doi:10.1016/j.neuron.2015.11.031

A

ETH

B

Power (5-14 Hz)

0.5au

0.1m

1au

2s

C

Real   Random

ETH

Real

Random

2dx/x

D

Frequency (Hz)

20
14.14

1.14

E

Power (5-14 Hz)

1au

1au

Plume Encounter   2s   Plume Encounter   2s

A

Probability

Head Raised

Difference

B

Cumulative Probability

Frequency (Hz)

C

Difference

Real

Random

-0.125g

Plume Encounter

0.05g

2s

D

Frequency (Hz)

9Hz

Plume Encounter

0.2Hz

2s

A1

A2

B Before   After/After   **C**

D

Mean Angle

Vector Strength

Plume Encounter   1s

A

Mean Angle

-2°

Random

1°

Vector Strength

Real

0.96

0.01

Plume Encounter 1s

B Before

40°

20°

0°

-20°

-40°

0   0.06   0.12

After

1s

C

0   0.04   0.08

D Real   Random

0° Crossings

Plume Encounter 1s

25 Encounters

E

0.1 z

1s