Local field potential sharp waves with diversified impact on cortical neuronal encoding of haptic input

Sofie S. Kristensen & Henrik Jörntell*

Neural Basis of Sensorimotor Control, Department of Experimental Medical Science, Lund University, Lund, Sweden

*corresponding author: Henrik.jorntell@med.lu.se

Acknowledgments

We thank Kaan Kesgin for inventing the dot product approach to identify the LFP sharp waves. We thank Dr. Jonas M.D. Enander for allowing us to use some of his data from the publication doi: 10.1016/j.celrep.2019.02.099. This research was funded by the EU Horizon 2020 research and innovation program under grant agreement No 861166 (INTUITIVE).

Abstract

In cortical sensory processing, internal activity substantially impact the cortical response to any given input. But controlling for that internal activity is difficult since the thalamocortical network is high-dimensional, perpetually active and its state can change at a high pace. Here we report on a type of spontaneous local field potential sharp wave (LFP-SPW) in the somatosensory cortex (S1) and explore its impact on spiking responses evoked in S1 neurons by tactile stimulation. LFP-SPWs that preceded the tactile input could have major impacts, depressing the tactile responses in some neurons and enhancing them in others. In spontaneous data, in contrast, LFP-SPW events triggered short-lasting but massive neuronal activation in all neurons recorded with a subset of neurons initiating their activation up to 20 ms before the LFP-SPW onset. LFP-SPWs often coincided with ECoG-SPWs recorded at the cortical surface 2 mm away from the patch electrode, suggesting that the LFP-SPWs could be part of a more global cortical signal with a variable extent. When LFP-SPWs and ECoG-SPWs coincided, the impact of the LFP-SPW on the neuronal tactile response could change substantially, including inverting the impact to its opposite. Such diversified impacts of the SPWs on different S1 neurons are well in line with previous observations of cortical neurons displaying diverse, complementary, response types to given sensory inputs. These cortical SPWs had similar overall activity patterns as reported for hippocampal SPWs and may be a marker for a particular type of state change that possibly involves both hippocampus and neocortex.

Introduction

How neocortical information processing is configured remains open to various alternative interpretations. In recent years, multiple observations suggest that representations of information are widespread across neuron populations (Allen et al., 2019; Enander and Jorntell, 2019; Enander et al., 2019) and that individual neurons within certain constraints carry complementary information (Luczak et al., 2009; Oddo et al., 2017; Stringer et al., 2019), even when they are located literally right next to each other and are provided with the exact same input (Kristensen and Jörntell, 2023; Oddo et al., 2017). However, also the time-continuous variations in internal brain state impact how the individual neurons represent the information. We have previously reported that the exact same tactile input pattern can trigger a wide variety of responses in individual primary somatosensory (S1) cortical neurons (Norrlid et al., 2021). The response variations are due to internal state variations in the neocortical neuronal network globally, which can be demonstrated by weakly perturbing either a remote area of the cortex (Etemadi et al., 2022; Wahlbom et al., 2019) or the output of the hippocampus (Etemadi et al., 2023). These response variations in individual neurons can be explained as a consequence of subtle changes in the state of the neocortex globally, which in turn can alter the number of "open" network pathways supplying the recorded neuron with tactile information. Hence, when interpreting the response of a neuron to any given input, it is important to also control for the current internal cortical state when the input is provided.

But controlling for that internal activity is difficult since the thalamocortical network is high-dimensional, perpetually active and its state can change at a high pace. There is therefore a value in identifying potential markers for such internally generated state changes. Here we apply a template-based signal extraction method to identify a wide category of events that in many respects resembles the hippocampal sharp wave, but which is recorded as spontaneous local field potential sharp waves (LFP-SPWs) at about mid-depth in S1.

The hippocampal sharp wave (Hipp-SPW) is a field potential signal that has been associated with many high-level brain functions, such as replay and recall (Buzsáki, 2015; Kanamori, 1985; Leonard and Hoffman, 2017; Liu et al., 2022). The Hipp-SPWs have a relatively distinct neurophysiological signature and are often superimposed by higher frequency components, ripples. The Hipp-SPW has been linked to activity changes in the neocortex in different ways (Logothetis et al., 2012; Mölle et al., 2006; Siapas and Wilson, 1998; Sirota et al., 2003). For example, Hipp-SPWs can trigger corresponding EEG-SPWs as demonstrated for the prefrontal cortical region (Mölle et al., 2006).

We found our LFP-SPWs to have similar overall activity patterns as reported for Hipp-SPWs. LFP-SPWs also strongly correlated with similar sharp waves recorded at the cortical surface using ECoG electrodes ('ECoG-SPWs'). As the ECoG-SPWs were recorded some distance away from the LFP-SPW recording site, this suggests that these SPWs represent some type of non-local cortical signal. In the spontaneous activity, both types of SPWs triggered a massive drive on the neuronal spiking activity. We found that preceding LFP-SPWs greatly impacted the S1 neuronal tactile responses, in some of the neurons by reducing and in other neurons instead by enhancing the evoked response. The ECoG SPWs, which occurred somewhat more rarely than the LFP-SPWs (but always coincided with them), could in turn substantially modify the impact that the LFP-SPW had on the tactile response.

Methods

Surgical procedure

Adult Sprague-Dawley rats (N=16, male sex, weight 306-420) were maintained under general anesthesia through a continuous infusion of ketamine and xylazine (ratio 20:1) mixed with Ringer acetate and glucose. The anesthetic mixture was administered as a continuous infusion through an intravenous catheter in the right femoral vein (approximately ~5 mg/kg per hour). The catheter was inserted by making an incision in the inguinal area of the hindlimb. Before inserting the catheter, animals were sedated with isoflurane (3% mixed with air for 60–120 s) and given an intraperitoneal injection of ketamine/xylazine (ratio 15:1) to induce anesthesia. While anesthetized, a small part of the skull (4x4 mm) was removed to expose the primary somatosensory (S1) cortex. The craniectomy extended from the coordinates 1 mm rostral to -3 mm caudally and 2-4 mm laterally relative to bregma (**Figure 1A**). An ECoG-electrode was placed on the surface of the cortex at the caudal end of the craniectomy. To prevent dehydration of the brain, the exposed brain area was covered in a thin layer of agarose (0.03 g/ml dissolved in physiological saline). Apart from recording SPWs, the ECoG signal was used to monitor the level of anesthesia by characterizing occurrences sleep spindles mixed with epochs of more desynchronized activity, an indication of sleep (Niedermeyer and da Silva, 2005). Note that ketamine/xylazine anesthesia has previously been shown to not affect the order of neuronal recruitment of a sheet of layer V neurons in spontaneous brain activity fluctuations and evoked responses as compared to the awake condition, suggesting that the neocortical network otherwise may work close to normal (Luczak and Barthó, 2012). Adequate anesthesia was further ensured by noxious pinching of the hind paw to confirm absence of withdrawal reflexes. Animals were sacrificed with pentobarbital (140 mg/kg IV) by the end of the experiment. All procedures related to animal experiments were approved by the Local Animal Ethics Committee of Lund, Sweden in advance (permit M13193-2017).

Recordings

Electrocorticographic recordings (ECoG) were made from the surface of the cortex at the rostral end of the craniectomy, approximately 4 mm laterally and 3 mm caudally to the bregma using a silver ball electrode (Ø 250 um at the cortex contact surface; no filters were applied, amplifier cut-off 50 kHz). Extracellular recordings of individual neurons were made with patch clamp pipettes in the loose-patch current clamp recording mode from the forepaw region of the S1, at the coordinates -1.0 – 1.0 mm relative to bregma. The location of the S1 was also estimated by evoked LFPs in response to stimulation of the second digit of the forepaw. Borosilicate glass capillaries were used to pull the patch pipettes to 10–30 MΩ using a Sutter Instruments (Novato, CA, USA) P-97 horizontal puller. Pipettes were filled with an electrolyte solution composited of (in mM) potassium-gluconate (135), HEPES (10), KCl (6.0), Mg-ATP (2), EGTA (10) titrated to 7.35–7.40 pH using 1 M KOH. The pipette was inserted into the cortex with an electrical stepping motor at a rate of 0.002 mm/s and neurons were recorded between cortical layers III and V. Unfiltered patch electrode recording data was digitized at 100 KHz using CED 1401 mk2 hardware and Spike2 software (CambridgeElectronic Design, CED, Cambridge, UK). The ECoG recording was digitized at 1 kHz using the same system.

Stimulation

We wanted to analyze the effect of LFP-SPWs on the evoked spike responses to haptic stimulation. We have previously shown that most S1 neurons have unique responses to

electrotactile spatiotemporal activation patterns mimicking mechanical skin-object interactions (Enander and Jorntell, 2019; Enander et al., 2019; Oddo et al., 2017).The patterns consisted of 5-33 pulses with intensities of 0.5 mA and durations of 0.14 ms (DS3 Isolated Stimulator, Digitimer, UK) delivered through four pairs of needle electrodes. Needle pairs were inserted into the volar side of the second digit of the forepaw with 2-3 mm distance between each pair. Patterns were delivered in pre-defined random order and lasted between 200 ms and 340 ms and were separated by about 1.8 s (random intervals). Each pattern was repeated 50 times in one experiment. These were thus the exact same set of spatiotemporal activation patterns we have used in earlier publications, but since we in the current paper only cared about the earliest part of the response evoked by the first stimulation pulses for each pattern, we simply considered pooled responses from all stimulation patterns and refer the interested reader to the original papers in order to view the full patterns.

Responsiveness to stimulation

Before pooling the responses, we calculated the responsiveness to each of the eight stimulation patterns for individual neurons. We constructed peristimulus time histograms (PSTHs) of the spike responses across all 100 repetitions of a stimulation pattern (in a few cases only 50 repetitions). The spike responses in the 200 ms time window preceding the onset of the stimulation were binned at 2 ms time bins and used to calculate the mean spike frequency and standard deviation (SD) of the baseline activity. If the average poststimulus activity (30 ms) exceeded the baseline activity by more than two SDs, then it was counted as being responsive. Stimulation patterns that did not evoke a response according to this definition, were excluded from the pooled responses of the neuron used for further analysis. Only neurons with responses to at least 3/8 of the patterns (i.e. at least 150 repetitions of tactile inputs) were included in the analysis.

Extracting neuron spikes, LFP-SPWs and ECoG-SPW s

Neuron spike events were extracted and time-stamped using an in-house software employing a template based spike identification (Mogensen et al., 2019). Local field potential sharp waves (LFP-SPW) and ECoG sharp waves (ECoG-SPW) were identified using a waveshape transform approach, which is related to continuous wavelet transform but uses a fixed waveshape rather than a wavelet. An LFP-SPW waveshape was based on visually identified LFP-SPWs, which had a latency to peak of 10-15 ms and a duration of 20-30 ms. The LFP waveshape vector was multiplied with the vector of the time continuous patch recording data, with a stride of 500 data sample points (5 ms), across the entire recording (typically 30-60 mins). At each stride, we used the dot product of the transform to define if an SPW was present by setting a threshold. We manually set the threshold of the dot product for each neuron recording individually, so that as many visually identified events as possible could be detected without including other, less structured variations in baseline activity. We removed detected events if they were detected less than 10 ms apart to avoid the same event being detected twice. The same method was used to detect ECoG-SPWs in the ECoG recording, except that here we used two waveshape templates; one with a positive voltage deflection and one with a negative voltage deflection. Here, the waveshape was multiplied with a stride of one data sample point (1 ms).

Spike responses to tactile stimulations, LFP-SPWs or ECoG-SPWs

To visualize the relationships between the spike activity and the tactile stimulations, the LFP-SPWs and the ECoG-SPWs, we constructed Kernel Density Estimations (KDEs) of the spike

173 responses. Each triggered trace consisted of a time series of spikes. Each individual spike
174 response, occurring at a specific time point relative to the trigger (which was a tactile
175 stimulation or a SPW), was convolved with a 2 ms Gaussian kernel. Then all the convolved
176 responses of that category were summed to obtain a KDE of the response around the trigger
177 point.

STATISTICAL ANALYSIS

Classifying the spike responses triggered by spontaneous SPWs

For the spike responses to the SPWs, the intensity and latency time of the spike activity that
preceded the onset of the LFP-SPWs were used to define three distinct groups; neurons with
no preceding activity, neurons with short preceding activity, and neurons with longer
preceding activity. To sort neurons into these groups, we calculated the normalized baseline
KDE activity and its SD in a time window starting 100 ms to 20 ms in the pre LFP-SPW
interval separately for each neuron. The time window was chosen because the earliest
preceding activity was observed to precede the LFP-SPW onset by 20 ms. If a neuron did not
have any spike activity exceeding its baseline by two SDs in the 20 ms time window
preceding LFP-SPW onset, the neuron would fall into the category "no preceding activity". If
a neuron exceeded baseline activity by two SDs within 7 ms preceding LFP-SPW onset, it
would be categorized as "short preceding activity". If a neuron exceeded baseline activity by
two SDs at more than 7 ms preceding LFP-SPW onset, it would be categorized as "longer
preceding activity". The same procedure was repeated using ECoG-SPWs as the trigger.

To statistically analyze if ECoG-SPWs evoked more spikes than LFP-SPWs, the amplitudes
of KDEs were measured for each individual neuron in each group and compared with
Welch's t-test. Before running the test, the distributions of the two groups were checked for
normality by plotting histograms. *Levene*'s test for homogeneity of variance failed which is
why Welch's t-test was chosen instead of a paired student's t-test.

LFP-SPW impacts on spike responses evoked by tactile stimulation

To analyze the effect of LFP-SPWs on the evoked spike responses for individual neurons, all
recording traces were divided into two groups for each neuron individually. The first group of
recording traces were the ones in which stimulations were not preceded by an LFP-SPW in
the 100 ms time window preceding stimulation onset. The second group consisted of traces
where stimulations were preceded by an LFP-SPW in this time window. To compare the
evoked responses to the stimulations for these two groups of traces, the number of spikes
occurring in the 30 ms post stimulation time window was counted for each trace in each
group. This resulted in two distributions of ordinal data. The Mann-Whitney U test was used
to test for a statistical difference between the two groups of traces (traces preceded by and
traces not preceded by an LFP-SPW) in terms of the number of spikes in each trace. If the test
result was significant, and the difference between the normalized sum of the evoked spikes
was negative, we classified the neuron as having a response that was "depressed" by the SPW.
If the difference between the normalized sum of the evoked spikes instead was positive, we
classified the neuron as being "excited" by the SPW. If the test result was not significant, we
classified the neuron as having "no effect" from the preceding SPW. Effect size was
calculated with: $r = \frac{z}{\sqrt{n}}$

ECoG-SPW effects on the LFP-SPW impact on the evoked responses

The same method as described above was used to compare evoked responses preceded by
LFP-SPWs in the 100 ms pre stimulus time window but with or without a coinciding ECoG-
SPW. Also here, neurons were classified as "depressed", "excited", or with "no effect"
depending on the significance of the test. In this case, depressed would indicate that the
coinciding ECoG-SPW reduced the evoked response, and vice versa for excitation. Only
neurons that had 15 or more traces per category were included in the corresponding analysis.

Results

222
Our aim was to investigate the effect of LFP-SPWs and ECoG-SPWs, which both stood out as
salient signals in our recordings (**Figure 1A,B**), on tactile-evoked neuronal spiking responses.
LFP-SPWs were recorded with the same patch electrode that was also used to record unitary
neuronal spikes, and the ECoG signal was obtained from the surface of the cortex at least 2
mm away from the patch electrode. Neuron recordings were made across layers III-V (depth
0.432-1.094 mm, except one neuron at depth 0.330 mm), which corresponded to the depths
where the LFP-SPWs were found to be the most prominent. A total number of 65 neurons
were recorded from 16 rats with neuronal responses evoked by haptic stimulation delivered to
the second digit of the forepaw. 48 neurons passed the inclusion criterion for analysis
(Methods).

Figure 1. Properties of LFP and ECoG sharp waves. (A) LFP and ECoG recording sites inside and outside the forelimb S1 region, respectively. (B) Raw recording data from a sample experiment. The asterisks indicate LFP sharp waves (LFP-SPWs) and ECoG sharp waves (ECoG-SPWs), respectively. Grey shading indicates periods of skin stimulation on the second digit. (C) Top: LFP-SPW template. Bottom: ECoG-SPW templates. (D) Zoom-in on a segment of the raw traces in A. Below, in green, are the equivalent traces of the dot product, obtained by multiplying the templates with the raw recordings across each time step. The SPW-detection threshold is indicated for each dot product trace as black dashes lines. (E). 25 superimposed LFP-SPWs and coinciding ECoG-SPWs to illustrate their variability in

appearance. (F) Occurrences of LFP-SPW triggered by occurrences of ECoG-SPW. (G) Inter-event intervals of LFP-SPWs. (H) Inter-event intervals of ECoG-SPWs. (I, J) Interspike intervals (ISIs) for two example neurons, for comparison with the Inter-event intervals in G, H. Note the much shorter time scales of the ISIs.

LFP-SPW	3.3 Hz (1.8 Hz)
EcoG-SPW	1.4 Hz (0.8 Hz)
LFP-SPW with ECoG-SPW	33.8 % (16.7 %)
LFP-SPW with spike	36.9 % (20.2 %)
LFP-SPW amplitude	-1.3 mV (0.7 mV)
LFP-SPW latency to peak	11.6 ms (5 ms)

Table 1: Frequencies of occurrence of LFP-SPWs and ECoG-SPWs across all 48 neurons. Coincidence frequency between LFP-SPWs and ECoG SPWs, and between LFP-SPWs and spikes. Quantification of the variety of LFP-SPW configurations.

Shape and frequency analysis of SPWs

Figure 1B illustrates a four-second-long segment of a parallel patch and ECoG recording from one example neuron. Light blue asterisks indicate LFP-SPWs in the patch recording and dark red asterisks indicate ECoG-SPWs in the ECoG recording. **Figure 1C** shows the wavelet templates used to detect the presence of LFP-SPWs and ECoG-SPWs, in the patch recording and the ECoG recording, respectively. **Figure 1D** shows a zoom in on 0.5 seconds of the segment together with the dot product traces that resulted from multiplying the waveshape templates with the raw data. **Figure 1E** illustrates that both LFP-SPWs and ECoG-SPWs were highly variable in their appearance (latency-to-peak and peak amplitude for example, see **Table 1**), which naturally made it impossible to capture all the occurrences of either type of event. According to our detection method (Methods), LFP-SPWs occurred more than 2 times more frequently than ECoG-SPWs (**Table 1**). ECoG-SPWs occurred almost exclusively in coincidence with LFP-SPWs, which is illustrated in **Figure 1F** where the occurrence of LFP-SPWs was triggered on ECoG-SPWs. **Figure 1G,H** illustrate the inter-event-intervals (IEIs) for LFP-SPWs (light blue) and ECoG-SPWs (dark red) respectively, from the same example neuron recording. IEIs of SPWs were not dissimilar from inter-spike-intervals (ISIs) (**Figure 1I,J**) in shape, i.e. ISIs and IEIs both had a tendency towards a sharp peak followed by a log-normal style of distribution, as previously described for a wide set of neurons (Nilsson and Jörntell, 2021; Spanne et al., 2014), but the SPWs occurred at a much slower overall rate. What can also be observed in **Figure 1B,D** is that spikes sometimes coincided with the LFP-SPWs (**Table 1**), but spikes could also occur outside these events.

The effect of LFP-SPWs on evoked spike responses

The first part of the analysis was centered in our main question, namely, if LFP-SPWs had an impact on the tactile-evoked spike responses. To analyze this, we divided all tactile-evoked recording traces of a neuron into two classes, depending on whether they were preceded or

not preceded by an LFP-SPW in the 100 ms prestimulus time window. **Figure 2A** illustrates example raw traces and KDEs of all responses for a sample neuron where the presence of a preceding LFP-SPW (in the grey box time window) resulted in a depression of the tactile-evoked response (measured in the red time zone). **Figure 2B** instead illustrates a neuron where the preceding LFP-SPW resulted in an excitation of the tactile-evoked response.

To statistically analyze the differences between the two classes of responses for every neuron individually, we used Mann-Whitney U as a test. The test showed a significant difference in the evoked responses between the populations of traces with a preceding LFP-SPW and the population of traces without a preceding LFP-SPW for 30 out 48 neurons (see supplementary data for details from individual tests). Of these 30 neurons, 21 showed a decrease in evoked spikes when preceded by an LFP-SPW, indicating that the presence of LFP-SPWs within 100 ms pre stimulus onset were depressing the evoked spike response. Nine of the neurons with a significant difference, showed an increase in evoked spikes when preceded by an LFP-SPW, indicating excitation of evoked responses. In 17 of the 21 significantly affected neurons, the effect size of the difference was small ($r < 2$), in three neurons the effect was small-moderate ($r > 2, < 3$), and in one neuron the effect was moderate ($r > 3$) (exact effect sizes are reported in the Supplementary Data). The difference in spike frequency in the 30 ms response windows is displayed as box plots in **Figure 2D** separately for the three groups of neurons (no significant effect, significant depression, and significant excitation).

Figure 2. Differential impacts of LFP-SPW on tactile evoked responses across neurons. (A) A neuron for which the response to tactile stimulation was depressed when it was preceded by an LFP-SPW. The response is shown as a sample raw trace and as a KDE plot of the average evoked spike response. Left panel corresponds to responses that were not preceded by an LFP-SPW (in the time window shaded in grey). Right panel corresponds to the evoked spike responses when the tactile stimulation was preceded by a LFP-SPW. Red box highlights the time window where the evoked spike responses were counted. (B) Similar display as in A, but for a neuron in which the spike response was enhanced when there was a preceding LFP-SPWs (compare left with right panel). (C) Electrotactile stimulation sites. (D) Summary of LFP-SPW impacts on the tactile evoked spike responses. Neurons were either found to not

have statistically significant effect of preceding LFP-SPWs, to have a significant depression, or to have a significant spike response enhancement.

The impact of SPWs on neuronal spiking

SPWs had a major impact on neuronal spiking (**Figure 3A-C**). When we used SPWs in the spontaneous activity to trigger the spike occurrences, the corresponding PSTHs (here shown as KDEs) illustrated massive spike responses (**Figure 3B,C**). Interestingly, a subset of our neurons had a build-up of activity starting before the occurrence of the SPWs. This suggests they contributed to eliciting the SPWs (as has been suggested for hippocampal neurons for Hipp-SPWs (Stark et al., 2015)). We could separate the neurons into three categories (**Figure 3B**), neurons without preceding spiking activity (N=13), neurons that had spiking activity that preceded the LFP-SPW for a short time window (7 ms or less; N=22) and neurons that that had spiking activity in a long time window (more than 7 ms, maximally 20 ms was observed) preceding the onset of the LFP-SPW (N=13). For the ECoG SPW, the same categories were observed but individual neurons did not necessarily fall into the same category as for the LFP-SPW. When using ECoG-SPWs as the trigger, a total number of 14 neurons showed no preceding activity, 25 neurons showed short preceding activity and 9 neurons showed long preceding activity (Figure 3C). The ECoG-SPWs in general evoked a more powerful spike response than the LFP-SPWs. In 34 neurons, the peak amplitude for the ECoG-SPW KDEs was higher compared to the amplitude for the LFP-SPW KDEs (**Figure 3D**). Welch's t-test indicated that the mean amplitude in the ECoG-SPW group (62.4 ± 34.9 Hz) was significantly higher than the mean amplitude in the LFP-SPW group (41.7 ± 29.5 Hz) with a difference of 20.6 (95% CI; 15.2 to 26.1); t(47) = 3.1, p <0.01 one tailed. Overall, the intense neuron responses indicated that the local network was in a different state when the SPWs occurred.

Figure 3. Spike responses triggered by LFP-SPWs and ECoG-SPWs (A) Superimposed raw data traces for a sample neuron, to illustrate spike responses around LFP-SPWs (top) and concomitant ECoG-SPWs (bottom). (B) Normalized SPW-triggered spike responses sorted by neurons with an absence of a preceding spike response, a short preceding response (up to 7 ms) and a long preceding response (up to 20 ms before the onset of the SPW). Individual neuron responses are shown as KDEs in grey, the average KDEs are shown in light blue (LFP-SPWs). (C) Similar display as in A but with ECoG-SPWs as the trigger, the average

KDE curves are shown in dark red. (D) The difference in peak amplitudes between the LFP-
SPW triggered spike responses and the ECoG triggered spike responses across all individual
neurons.

ECoG-SPWs affected the LFP-SPWs impacts on the evoked spike responses
Considering the more powerful effect of ECoG-SPWs on spontaneous spiking, we next
analyzed if the ECoG-SPWs affected the impact that LFP-SPWs had on evoked responses
when they coincided. We divided all recording traces for each individual neuron into two
classes; traces where preceding LFP-SPWs coincided with an ECoG-SPW and traces where
preceding LFP-SPWs did not coincide with an ECoG-SPW. Example KDEs for normalized
traces from the two groups are shown in the second and thirds columns of **Figure 4A-C** for
three example neurons as dark blue traces (no coinciding ECoG-SPW) and dark red traces
(with a coinciding ECoG-SPW). The first column with black traces illustrates traces that were
not preceded by any LFP-SPW or ECoG-SPW.

As before, we compared the evoked responses in the traces from the two classes using the
Mann-Whitney U statistical test (10 neurons with too few coinciding traces were excluded
from the analysis). In this case, it indicated a significant difference for nine of the neurons.
The effect size of the difference for these nine neurons showed five neurons with a small
effect ($r < 2$) and four neurons with a small-moderate effect ($r > 2, < 3$) (see Supplementary
Material for more detailed test results). Of the nine neurons that showed a significant
difference in the evoked responses, four neurons showed a reduction by the coincidence of an
ECoG-SPW (depressed) and five neurons showed an enhancement (excited) as shown in
Figure 4D.

Figure 4. Effect of coinciding ECoG-SPWs on the LFP-SPW-mediated impacts on evoked
spike responses. (A) An example neuron with a substantial depressing impact of the preceding
LFP-SPWs on the evoked response (dark blue KDE trace) but in which the coincidence of an
ECoG-SPW with the LFP-SPW did not affect that impact (dark red KDE trace), despite a
marked enhancement in the prestimulus spiking activity. (B) A neuron where the preceding
LFP-SPW enhanced the response (dark blue trace) but where the coincident ECoG-
SPW+LFP-SPW depressed that impact (dark red trace). (C) A neuron where the preceding
LFP-SPW depressed the response but where the coincident ECoG-SPW+LFP-SPW inverted
that impact to instead enhance the response above its control level (black trace to the left). (D)
Effect of coinciding ECoG-SPWs on the LFP-SPW-mediated impacts on the evoked spike
responses (i.e. differences between dark blue KDEs and dark red KDEs across all neurons).

Control analysis

Since ECoG-SPWs elicited more intense spike response than LFP-SPWs (**Figure 3**), and
affected the impact of the LFP-SPWs on the evoked responses (**Figure 4**) the next question
we asked was whether the mere presence of a spike in the LFP-SPW had a similar boosting
effect as the ECoG-SPW. Again we divided all recording traces for each individual neuron
into two classes, LFP-SPWs with spikes and LFP-SPWs without any coinciding spike (i.e. no
spike occurring between the onset of the SPW and two times the latency to peak) and
compared the groups using Mann-Whitney U. Example KDEs for the two classes of traces of
one neuron is shown in **Figure 5A**. Note the increased spike activity in the 100 ms time
window preceding the stimulation, indicated by the grey box. As before, we analyzed the

evoked spikes in the 30 ms post-stimulation onset time window (indicated by red numbers in
Figure 5A). In this analysis, only 2/48 neurons had a significant difference between the
evoked responses in the two groups. The effect size for both neurons was small ($r < 3$). See
Supplementary Material for details on Mann-Whitney U tests and effect sizes.

Figure 5. Control analysis if the presence of LFP spiking affected the LFP-SPWs impacts on
the evoked responses. (A) KDEs of the spike responses when the LFP-SPW contained spike
responses (top) or not (bottom). (B). Quantitative analysis indicated that the presence of spike
responses in the LFP-SPWs did not significantly impact the evoked response except for the
two outliers above.

Discussion

Here we reported on a particular category of neocortical LFP event, the LFP SPW, which we
found to have a relatively large and diversified impact on the neuronal responses evoked by
tactile stimulation. This finding can be explained by that the LFP-SPW was associated with a
significant state change, at least in the local circuitry that mediated the tactile input to the
recorded neuron. The LFP-SPW often coincided with an ECoG-SPW recorded some distance
away, which suggests that at least a subset of the LFP-SPWs was associated with a more
widespread cortical state change. The activation dynamics of our cortical SPWs were similar
to SPWs previously recorded in the hippocampus, and a previous study shows that
hippocampal SPWs trigger SPW-like EEG-signatures at least in the prefrontal cortex (Mölle
et al., 2006). Hence, the LFP-SPWs may potentially be part of a widespread coordination of
cortical neuron activity.

We found that the LFP-SPW could have opposing impacts on the tactile-evoked responses in
different neurons. This is in line with previous observations that even adjacent neurons can
encode a given sensory input in a diversified manner (Kristensen and Jörntell, 2023; Oddo et
al., 2017), or have complementary patterns of spontaneous activity (Mogensen et al., 2019;
Stringer et al., 2019; Wahlbom et al., 2021). In relation to the LFP-spike relationship more
generally, a previous study concluded that there is a '…substantial heterogeneity in the timing
and strength of spike-EEG relationships and that these relationships became more diverse
during visual stimulation compared with the spontaneous state.' (Snyder and Smith, 2015).
Our results for the LFP-SPWs are hence in line with that more general EEG relationship, and
further emphasize the diversity of the relationships between individual neurons and
population level features in the brain activity.

The state change signaled by the LFP SPWs appeared to be quite different when the LFP-
SPW coincided with an ECoG-SPW. This was indicated by that the impact that the LFP-SPW
had on the tactile evoked response in a neuron (depression or excitation) could even be
inverted if the LFP-SPW coincided with an ECoG-SPW (**Figure 4**). Although spontaneous
SPWs triggered massive neuronal spike responses on average (**Figure 3**), the mere presence
or absence of spiking in the preceding LFP-SPW overall did not alter the impact it had on the
evoked response (**Figure 5**). This indicates that the spikes themselves were not good
indicators of a state change, in contrast to the SPWs.

The neuronal spiking activity was powerfully driven by the SPWs. This suggests that the
SPWs were associated with powerful synaptic drive to the local neurons. Many neurons also
had activity profiles that indicated that they were activated well in advance of the SPW
initiation (**Figure 4**). This may in addition suggest that the LFP-SPWs were a result of a rapid
but gradual build-up of activity in recurrent excitatory loops that at least partly involved the
local neurons – similar to the current interpretations of how Hipp–SPWs arise (Buzsáki, 2015;
Stark et al., 2015). Since Hipp-SPWs are associated with concomitant intense activity in the
neocortex, including the S1 (Sirota et al., 2003),this in turn raises the question if it is the
hippocampus that paces the neocortex, is it the other way around, or is it predominantly a
distributed contribution across both the hippocampus and the neocortex?

Similarities with Hipp-SPWs
In many respects our LFP-SPWs displayed similarities with Hipp-SPWs. Although the
activation characteristics of Hipp-SPWs are often not analyzed in detail in the literature, many
comparisons can be made based on raw data illustrations. The first similarity is their
approximate duration (Buzsáki, 2015; Mölle et al., 2006). A second similarity is their rate of
occurrence – Hipp-SPWs have been reported to occur at more than 1 Hz (Kanamori, 1985)
and example illustrations indicate rates in the order of up to 2 Hz (Mölle et al., 2006). This is
less than the 3.3 Hz we found for LFP-SPWs, but in line with the rate of our ECoG-SPWs.
However, as discussed below, our detection method may be more sensitive and thus missing
fewer SPW events. In addition, Hipp-SPWs have been reported to be less frequent in awake
active state than in awake relaxed state, sleep and anesthesia (Kanamori, 1985) as was the
case in our experiments. Also, our SPWs could occur in bouts of 2-4 SPWs in a series with
much shorter intervals between them (see **Figure 1**). Unfortunately, there is not a lot of
analysis on SPW IEIs *in vivo* in the literature, but raw data illustrations (i.e. Figure 1 in
Buzsáki (2015) for example) suggest this is the case also for hippocampal SPWs.

What do the SPWs represent and how common are they?
Why were the rates of occurrence of the LFP-SPWs lower than for the ECoG-SPWs? This
might in fact possibly have been due to event detection issues – SPWs were easier to detect in
LFPs, because of the more precise nature of the signal from the patch than from the silver ball
electrode, and because ECoG-SPWs could often be initially positive and then negative, which
made them harder to detect. But this highlights another problem that has not been addressed
in the literature – the detection threshold for SPWs will vary not only depending on the
analysis method but also on what recording method is used. This is likely to apply to
hippocampal recordings as well – one method to detect Hipp-SPWs that includes high-speed
ripples has been to band-pass filter the field potential data to above 50 Hz and then set the
detection threshold to 5 SDs (Stark et al., 2015). It is possible that such an approach will be at
risk of missing SPWs that do not have a significant part of their signal power above that

frequency threshold. This in turn comes back to the question whether SPWs have a strictly unitary character or if these types of events instead are much more gradable. Our detection method did not rely on band-pass-filtering but instead used the dot product obtained with template waveshapes. Even though also our method used an arbitrary detection threshold for the SPWs (**Figure 1D**) their highly variable amplitudes and temporal signatures suggest that the SPWs are extensively gradable, perhaps all the way down to being in continuum with overall baseline activity. Then it may be the case that SPWs occur at a much higher rate than reported in the literature – for example, their lower incidence reported in active animals than in quiet awake animals, as well as sleep (Kanamori, 1985), could possibly be because they are less well synchronized in the active awake state. Lower synchrony could in turn be due to that there is more active processing in the cortex (Poulet and Petersen, 2008) and hippocampus (Gardner et al., 2019). Alternatively, it may be that the higher baseline activity may tend to drown SPW-like phenomena and thus making it more difficult to detect them.

Concluding remarks

Future studies should begin to address the underlying circuitry mechanisms that drive these SPWs in greater detail. Only then can we begin to understand what they represent more specifically. So far perhaps the strongest indicator is that it has been found that Hipp-SPWs correlate with familiarity in visual scenes (Leonard and Hoffman, 2017). But given that there are also neocortical SPW events, could these also represent some type of local familiarity signal? And the fact that we observe them also under anesthesia, both in S1 in the present paper and in the hippocampus (Stark et al., 2015), where information integration is unlikely to be coherent over longer periods of seconds or more, could they represent some type of time-local information-processing completion? The path towards addressing such questions lies in a better understanding of their circuitry-level mechanistic underpinnings and what they correspond to for the cortical information processing mechanisms in general.

485   REFERENCES

Allen, W.E., Chen, M.Z., Pichamoorthy, N., Tien, R.H., Pachitariu, M., Luo, L., and
Deisseroth, K. (2019). Thirst regulates motivated behavior through modulation of brainwide
neural population dynamics. Science 364, 253.

Buzsáki, G. (2015). Hippocampal sharp wave-ripple: A cognitive biomarker for episodic
memory and planning. Hippocampus 25, 1073-1188.

Enander, J.M.D., and Jorntell, H. (2019). Somatosensory Cortical Neurons Decode Tactile
Input Patterns and Location from Both Dominant and Non-dominant Digits. Cell Rep 26,
3551-3560 e3554.

Enander, J.M.D., Spanne, A., Mazzoni, A., Bengtsson, F., Oddo, C.M., and Jörntell, H.
(2019). Ubiquitous Neocortical Decoding of Tactile Input Patterns. Fronties in Cellular
Neuroscience 13, 1-17.

Etemadi, L., Enander, J.M.D., and Jörntell, H. (2022). Remote cortical perturbation
dynamically changes the network solutions to given tactile inputs in neocortical neurons.
iScience 25.

Etemadi, L., Enander, J.M.D., and Jörntell, H. (2023). Hippocampal output profoundly
impacts the interpretation of tactile input patterns in SI cortical neurons. iScience 26.

Gardner, R.J., Lu, L., Wernle, T., Moser, M.B., and Moser, E.I. (2019). Correlation structure
of grid cells is preserved during sleep. Nat Neurosci 22, 598-608.

Kanamori, N. (1985). A spindle-like wave in the cat hippocampus: a novel vigilance level-
dependent electrical activity. Brain Research 334, 180-182.

Kristensen, S.S., and Jörntell, H. (2023). Differential encoding of temporally evolving color
patterns across nearby V1 neurons. Frontiers in Cellular Neuroscience 17, 1249522.

Leonard, T.K., and Hoffman, K.L. (2017). Sharp-Wave Ripples in Primates Are Enhanced
near Remembered Visual Objects. Current Biology 27, 257-262.

Liu, A.A., Henin, S., Abbaspoor, S., Bragin, A., Buffalo, E.A., Farrell, J.S., Foster, D.J.,
Frank, L.M., Gedankien, T., Gotman, J., *et al.* (2022). A consensus statement on detection of
hippocampal sharp wave ripples and differentiation from other fast oscillations. Nat Commun
13, 6000.

Logothetis, N.K., Eschenko, O., Murayama, Y., Augath, M., Steudel, T., Evrard, H.C.,
Besserve, M., and Oeltermann, A. (2012). Hippocampal-cortical interaction during periods of
subcortical silence. Nature 491, 547-553.

Luczak, A., and Barthó, P. (2012). Consistent sequential activity across diverse forms of UP
states under ketamine anesthesia. European Journal of Neuroscience 36, 2830-2838.

Luczak, A., Bartho, P., and Harris, K.D. (2009). Spontaneous events outline the realm of
possible sensory responses in neocortical populations. Neuron 62, 413-425.

Mogensen, H., Norrlid, J., Enander, J.M.D., Wahlbom, A., and Jorntell, H. (2019). Absence
of Repetitive Correlation Patterns Between Pairs of Adjacent Neocortical Neurons in vivo.
Front Neural Circuits 13, 48.

Mölle, M., Yeshenko, O., Marshall, L., Sara, S.J., and Born, J. (2006). Hippocampal sharp
wave-ripples linked to slow oscillations in rat slow-wave sleep. Journal of Neurophysiology
96, 62-70.

Niedermeyer, E., and da Silva, F.H.L. (2005). Electroencephalography:
Basic Principles, Clinical Applications and Related Fields. Encyclopedia of
Psychopharmacology.

Nilsson, M.N.P., and Jörntell, H. (2021). Channel current fluctuations conclusively explain
neuronal encoding of internal potential into spike trains. Physical Review E 103.

Norrlid, J., Enander, J.M.D., Mogensen, H., and Jörntell, H. (2021). Multi-structure Cortical
States Deduced From Intracellular Representations of Fixed Tactile Input Patterns. Frontiers
in Cellular Neuroscience 15.

Oddo, C.M., Mazzoni, A., Spanne, A., Enander, J.M.D., Mogensen, H., Bengtsson, F.,
Camboni, D., Micera, S., and Jörntell, H. (2017). Artificial spatiotemporal touch inputs reveal
complementary decoding in neocortical neurons. Scientific Reports 8.

Poulet, J.F., and Petersen, C.C. (2008). Internal brain state regulates membrane potential
synchrony in barrel cortex of behaving mice. Nature 454, 881-885.

Siapas, A.G., and Wilson, M.A. (1998). Coordinated Interactions between Hippocampal
Ripples and Cortical Spindles during Slow-Wave Sleep. Neuron 21, 1123-1128.

Sirota, A., Csicsvari, J., Buhl, D., and Buzsáki, G. (2003). Communication between neocortex
and hippocampus during sleep in rodents. Proceedings of the National Academy of Sciences
of the United States of America 100, 2065-2069.

Snyder, A.C., and Smith, M.A. (2015). Stimulus-dependent spiking relationships with the
EEG. J Neurophysiol 114, 1468-1482.

Spanne, A., Geborek, P., Bengtsson, F., and Jörntell, H. (2014). Spike generation estimated
from stationary spike trains in a variety of neurons In vivo. Frontiers in Cellular Neuroscience
8.

Stark, E., Roux, L., Eichler, R., and Buzsaki, G. (2015). Local generation of multineuronal
spike sequences in the hippocampal CA1 region. Proc Natl Acad Sci U S A 112, 10521-
10526.

Stringer, C., Pachitariu, M., Steinmetz, N., Reddy, C.B., Carandini, M., and Harris, K.D.
(2019). Spontaneous behaviors drive multidimensional, brainwide activity. Science 364, 255.

Wahlbom, A., Enander, J.M.D., Bengtsson, F., and Jorntell, H. (2019). Focal neocortical
lesions impair distant neuronal information processing. J Physiol 597, 4357-4371.

Wahlbom, A., Mogensen, H., and Jorntell, H. (2021). Widely Different Correlation Patterns
Between Pairs of Adjacent Thalamic Neurons In vivo. Front Neural Circuits 15, 692923.

560
561

LFP-SPW template

10 ms

ECoG-SPW templates

10 ms

Dot product

Dot product

E

LFP-SPW (Patch)

5 mV

ECoG-SPW (ECoG)

50 ms

5 mV

2.5 mV

F

ECoG-SPW

1200

#LFP-SPW

-200   -100   0   100   200
(ms)

G

1200

Inter-LFP-SPW-Interval

0   400   800   1200   1600   2000
(ms)

H

Inter-ECoG-SPW-Interval

0   400   800   1200   1600   2000
(ms)

I

Inter-Spike-Interval

0   10   20   30   40   50
(ms)

J

Inter-Spike-Interval

0   10   20   30   40   50
(ms)

LFP-SPW   ECoG-SPW

LFP-SPW

ECoG-SPW