<u>**Environmental fungi from cool and warm neighborhoods in the urban heat island of**</u>

<u>**Baltimore City show differences in thermal susceptibility and pigmentation**</u>

Daniel F. Q. Smith[1], Madhura Kulkarni[1], Alexa Bencomo[1,2], Tasnim Syakirah Faiez[1,3], J. Marie

Hardwick[1], Arturo Casadevall*[1]

[1] W. Harry Feinstone Department of Molecular Microbiology and Immunology, The Johns

Hopkins Bloomberg School of Public Health, Baltimore, MD, 21205, USA

[2] Krieger School of Arts & Science, Johns Hopkins University, Baltimore, MD 21218, USA

[3] Department of Pathobiology, Johns Hopkins School of Medicine, Baltimore, MD 21205, USA

*Corresponding Author: acasade1@jh.edu

Keywords: Urban fungi, climate change, disaster microbiology, thermotolerance, pathogenic

potential, heat island effect

Abstract

A major barrier for most fungal species to infect humans is their inability to grow at body

temperature (37°C). Global warming and more frequent extreme heat events may impose

selection pressures that allow fungal adaptation to higher temperatures. Cities are heat islands

that are up to 8°C warmer than their suburban counterparts because of mechanical heat

production and reduced greenspace, among other factors, and may be an important reservoir of

fungi that have increased risk of thermotolerance and inhabit environments near humans. Here

we describe a novel and inexpensive technique that was developed to collect fungal samples

from various sites in Baltimore, Maryland using commercially available taffy candy. Our results

show fungal isolates from warmer neighborhoods show greater thermotolerance and lighter

pigmentation relative to isolates of the same species from cooler neighborhoods, suggesting

local adaptation. Lighter pigmentation in fungal isolates from warmer areas is consistent with

known mechanisms of pigment regulation that modulate fungal cell temperature. The

opportunistic pathogen *Rhodotorula mucilaginosa* from warmer neighborhoods had a higher

resistance to gradual exposure to extreme heat than those from cooler neighborhoods. Our

results imply fungal adaptation to increased temperature in an urban environment. The

acquisition of thermotolerance poses a risk for humans if fungal species with pathogenic

potential acquire the capacity to grow at human body temperatures and cause disease.

Introduction

Humans endothermy provides natural protection against many fungal species with

pathogenic potential [1]. Most fungi are unable to survive at or above human body temperature,

and analyses of yeast thermal tolerance has shown that there is a 6% reduction in the number

of fungal species able to survive at each degree above 30°C [1,2].

Natural and human-made disasters are associated with adaptive changes in

environmental microbes that can lead to the emergence of new diseases in humans, plants, and

animals [3]. Changes in the environment have been hypothesized to enable adaptive changes

in fungi and drive the emergence of new fungal pathogens [4], a phenomenon that may have

already occurred with *Candida auris* [4,5]. Global warming and extreme heat (including

prolonged and more frequent heatwaves) may enable environmental fungi with pathogenic

potential to acquire higher thermotolerance and thus overcome the mammalian thermal barrier.

In recent decades, a sharp emergence of new fungal pathogens such as *Candida auris* and

Sporothrix brasiliensis has been proposed to result from environmental pressure to acquire

thermotolerance [4,6]. An analysis of the fungal collection at the Westerdijk Fungal Diversity

Institute showed that fungal isolates collected globally from the environment in recent decades

have a higher maximum growth temperature than those collected in prior decades , consistent

with a possible gain of thermotolerance due to global warming [2]. However, that study is limited

as the isolates in their collection are not randomly collected from the environment and may not

represent the true diversity of fungi present in the environment over the course of a century.

Environments with high thermal selective pressure include urban centers (cities), which

are subjected to the "heat island effect" and tend to be 4-8°C warmer than rural or suburban

neighboring regions [7–10]. The warmer environment in cities is predominantly due to reduced

green spaces, blocking of natural airflow by buildings, higher heat absorbance by building

materials, and anthropomorphic heat production by machines and cars. However, even within

cities, there can be large temperature variation, with some neighborhoods experiencing more

severe heat than others, and the warmer neighborhoods are often the home to marginalized

communities [10–12]. A study published by the National Oceanographic and Atmospheric

Administration (NOAA) reported that in Baltimore City in late August 2018, parks and suburban

neighborhoods are approximately 30.5°C, which is about 8-9°C cooler than the warmest

neighborhoods that reach 39.4°C [13].

Currently, there are few studies investigating the mycobiome of urban centers, including

areas where there are ample bacterial microbiome analyses, partly due to technical difficulties in

performing metagenomic analysis of fungal DNA without adequate reference genomes for

comparison and sequence alignment [14,15]. Few studies have investigated the heat-resistance

and phenotypic profiles of urban fungi. An analysis of filamentous environmental fungi from

urban sites in Louisville, Kentucky showed faster growth rates and higher enzymatic activities at

higher temperatures than the fungi of the same species isolated from rural sites [16].

To understand mechanisms of the emergence of fungal pathogens there is a need for

more functional characterization of urban mycobiome and a more detailed characterization of

how heat affects urban fungal populations. Consequently, we explored the urban fungal census

from sidewalks across thermally diverse neighborhoods in Baltimore, Maryland using a new

technique for collection, and show examples of genotypic and phenotypic characterization of the

fungi present, including pigmentation variation and the isolation and identification of clinically

relevant polyextremophilic yeasts.

Materials and Methods

Sample Site and Date Selection

To select thermally diverse neighborhoods in Baltimore, MD, we used publicly available

data on heat disparities across Baltimore City. Based on 2018 data from the National

Oceanographic and Atmospheric Administration (NOAA) [13], we collected samples from four

sites across Baltimore City. For the two warm neighborhood sites, we chose the corner of N.

Wolfe St and E. Fayette St. ("Fayette St.") and the corner of N. Charles St. and Washington

Park East ("Mt. Vernon"), representative of the warmest areas in Baltimore. For the average

temperature neighborhood site, we chose the corner of S. Wolfe St., and Fell St ("Fells Point"),

and for a cool neighborhood site, we chose the corner of N. Charles St. and St. Paul St

("Guilford"). On August 20[th], 2023, we collected samples from all four sites during sunny

weather, between 12 P.M. and 2 P.M. Additional collections were done on September 4[th], 2023,

June 21[st], 2024, and August 28[th], 2024, between 12 P.M. and 3 P.M, for only the (warmer)

Fayette St. and (cooler) Guilford sites. All dates were chosen due to sunny weather and were at

least three days after the most recent precipitation. The two collection dates in June and August

2024 were for the purpose of performing further mold pigmentation measurements. Thermal

images to record the temperature of each site were taken using a FLIR C2 IR camera, and

images were processed using FIJI (ImageJ) IRimage plugin [17]. Additional temperature

readings were done using a GP-300 infrared thermometer (Duraline Systems, West Nyack, NY)

to confirm infrared camera measurements. Measurements for the thermometer and FLIR

camera were comparable (difference <1°C) (Supplementary Table S1).

Sidewalk Sample Collection

To collect fungal samples from sidewalks, we developed a collection protocol using

Starburst taffy candy (Mars Corporation, VA, USA) (Figure 1A). Taffy is a soft, sticky candy that

easily adapts its contour to that of environmental surfaces when pressed against those surfaces

with hand pressure. The stickiness picks up small particles. To collect environmental fungi found

in the gritty and textured surface of the sidewalk, we rolled a yellow Starburst taffy over an

approximately 100 cm^2 area on the sidewalk for 30 seconds, making sure to press the taffy into

the pavement. The taffy candy was then dissolved in 10 mL PBS containing 60 µl of 25% KOH

solution to neutralize the citric acid in the candy. After the taffy was dissolved, the suspension

was passed through a 100 µm filter to remove debris, and 200 µl aliquots of sidewalk samples

were spread on each of three to five 10 cm Sabouraud dextrose agar (SDA) plates with

penicillin-streptomycin to limit bacterial growth. SDA was chosen as the isolation agar because

it is acidic and inhibits many bacterial species while promoting fungal growth. Samples were

incubated at room temperature (22˚C) for up to 7 d. The Starburst taffy alone did not grow fungi

when treated with these conditions, except for a single mold colony (Supplementary Figure 1).

Dirt Sample Collection

Dirt samples were collected by scraping the top layer of dirt along the tree-wells, garden

patches between sidewalks, and grass-tufted crevices adjacent to the curb. At least 5 mL of

non-compacted dirt was collected at each location. The 5 mL of dirt sample were added to 45

mL PBS, solubilized at room temperature for 15- 30 min, passed through a 100 μm strainer to

remove large debris, plant material, and larger insoluble objects, diluted 1:100 in PBS, and 50 μl

aliquots of diluted dirt samples were spread on 3-5 10 cm SDA plates containing penicillin-

streptomycin and incubated at room temperature (22˚C) for up to 7 d. Isolation scheme

summarized in Figure 1B.

Yeast Colony Isolation

Fungal colonies were isolated from original SDA plates containing material from

sidewalk and dirt samples by re-streaking on SDA plates. Plates were incubated at room

temperature for up to 7 d. Plates were examined daily for fungal growth, both filamentous and

yeast like. Yeast-like colonies (small, circular, opaque, and non-filamentous) were picked with a

pipette tip and streaked to isolation on fresh yeast peptone dextrose (YPD) agar plates, and

incubated at 20˚C. Newly appearing colonies derived from the dirt and sidewalk collection plates

were picked from the YPD plates daily for up to 7 d. Cells from each collected isolate were

observed microscopically to confirm fungal (non-bacterial) identify by cell size, morphology, and

appearance of intracellular organelles. Individual fungal colonies were isolated as they

appeared, transferred to 96-well plates with freezing media (25% glycerol, 75% YPD broth), and

stored at -80˚C. Replicate 96-well plates with sterile PBS were inoculated and stored at 4˚C. All

original plates were wrapped in parafilm and stored at 4˚C.

Growth Area Measurements

All the original samples on SDA agar plates from Fayette St. and Guilford were scanned

using a Canon CanoScan9000F scanner at 600 dpi after 7 days of growth. Using the FIJI

(ImageJ) selection and measure tools, ROIs were manually identified and the total area of

fungal growth per plate was determined for each site.

Pigmentation Analysis

Pigmentation of fungi was determined as previously described [18]. Briefly, pigmentation

of molds growing on the original SDA sample plates was assessed by imaging with a Canon

CanoScan9000F scanner at 600 dpi. To measure the pigmentation of yeast isolates, frozen

stocks of isolated colonies stored at -80˚C were stamped onto YPD agar on an OmniWell agar

plate using a flame-sterilized microplate replicator (Boekel Scientific) and grown at room

temperature for 7 days. Yeast growth (which were in spots of ~ 5 mm in diameter) were

scanned using the CanoScan9000F at 600 dpi. Images were processed and mean gray values

of mold and yeast colonies were measured using FIJI (ImageJ) measure tool.

Thermal Absorbance Analysis

Agar plates with mold cultures were acclimated to room temperature for 1 h (22˚C). Petri

dishes were unlidded and exposed to a pre-warmed 19W 4000K white light bulb at 50 cm for 10

minutes. Following the 10 min exposure, culture plates were imaged immediately for heat

absorption using an FLIR E96 infrared camera. Images were processed using FIJI (ImageJ)

IRimage plugin [17], and temperature measurements were recorded for each mold colony.

Mean gray value measurements were taken using the visible light spectrum image taken by the

FLIR E96 camera at time of experiment.

Heat-Ramp Assay

Heat-Ramp experiments were performed as previously described, with some changes

[19]. Frozen stocks of *R. mucilaginosa* and *C. minutum* collected from Fayette St., Mt. Vernon,

and Guilford were streaked on SDA plates and incubated 48 h at room temperature. Single

colonies of *R. mucilaginosa* and *C. minutum* were scraped from SDA plates, suspended in liquid

SDA and adjusted to a density of 0.1 absorbance at 600 nm and 200 µl of these cultures were

grown overnight at 30°C in stationary 96-well plates. Samples were diluted 1:5 in fresh SDA

liquid medium, and 100 µl of these cultures were directly treated with a linear heat-ramp from

30°C to 55, 56, or 60°C over 8-10 min in a water bath with agitation (LAUDA Scientific,

Germany). Untreated and heat-ramp–treated strains were immediately spotted (5 µL in SDA) in

five-fold serial dilutions on SDA agar and incubated at 30 °C for 48 h. Plates were imaged and

CFUs were counted for each isolate under each treatment.

Internal Transcribed Spacer (ITS) Sequencing

Colonies for each isolate were picked from newly streaked SDA plates of fungal isolates

touched lightly with a pipette tip and transferred and deposited along the inside of an empty

PCR tube in a thin layer. For cultures that were unable to grow following storage at -80°C, we

scraped some of the frozen stock and deposited it inside of a PCR tube. The tubes were

microwaved on high (1100 W) for 2 minutes as previously described [20]. DreamTaq Green

PCR Master Mix (ThermoFisher) was added to each tube with 0.2 µM ITS4 and ITS5 primers to

detect species-specific fungal internal transcribed sequences. The sequences of these primers

are as follows: ITS4 - 5′-TCCTCCGCTTATTGATATGC-3′ and ITS5 - 5′-

GGAAGTAAAAGTCGTAACAAGG-3′. PCR products were amplified according to the

manufacturer's protocol; 3 min initial denaturation at 95°C, 35 cycles with 30 s denaturation at

95°C, 30 s annealing at 55°C, and 1 min extension at 78°C, with a final 5 min extension at 78°C.

PCR products were run on 1% agarose gel to confirm product amplification, and Sanger

sequenced at the Johns Hopkins School of Medicine Sequencing & Synthesis Core facility.

Resulting sequences were analyzed using 4Peaks software and NCBI Blast nucleotide feature.

Results

To investigate the potential effects of urban heat islands on fungi, we collected samples

from four sites (Figure 2A and 2B) across Baltimore City based on 2018 air temperatures from

the National Oceanographic and Atmospheric Administration (NOAA) [13]. Fungal samples

were collected between 12 and 3 P.M. in Baltimore City on four dates with reported air

temperatures of 27-37°C during the summers of 2023 and 2024. Sample sites represented both

warmer and cooler neighborhoods and each site was representative of the environment in its

neighborhood based on NOAA heat island maps. The four sites chosen for the study were: Site

1, (Fells Point) representing an average air temperature neighborhood, Site 2 (Fayette St.)

representing the warmest area within Baltimore City, Site 3 (Mt. Vernon) representing an area

with above average temperature, and Site 4 (Guilford) representing a neighborhood with cooler

temperatures.

At each collection site, we measured sidewalk and dirt surface temperatures

(Supplementary Table 1). The surface temperatures we observed were consistent with patterns

we expected based on heat-island effect air temperatures published by NOAA. On the first and

second collection days (August 20, 2023, and September 4[th], 2023), we collected temperature

data using an FLIR E96 camera to verify the temperature patterns. On the second through

fourth collection dates, surface temperatures were recorded using a GP-300 infrared

thermometer. The raw images from the infrared camera to measure surface temperatures on

August 20[th], 2023 are displayed in the upper panels of Figure 2C and D, with an automatically

scaled color range based on the temperature range in the field of view, which differs in each

frame. To more directly compare average temperatures between the dirt and sidewalk sample

collection sites, the raw image files were converted to the gray scale (Figure 2C and 2D, lower

rows) to reflect the absolute temperature range. The grayscale image allows for direct

comparisons and quantifications, as it is scaled so that the gray value images have 27˚C as the

darkest pixel value and the whitest pixel values to represent 60˚C. The measured average

temperature from the approximate area of the red circle at the center of the image is overlayed.

On August 20[th], 2023, the Fells Point site had a warm sidewalk with an average of

40.7˚C, while a cooler dirt sample ambient temperature (27.2 ˚C) (Figure 2C, D). As expected,

the dirt and sidewalks from the Fayette St. and Mt. Vernon collection sites had the warmest

surfaces tested. Sidewalks in these locations were ~38.4-39.8˚C (Figure 2C), with the dirt

temperature at either site of sample collection reaching up to ~53-57.7˚C on average, with some

areas on the surface reaching up to 60˚C (Figure 2D). Both sidewalk and dirt samples from

Guilford were ~27˚C, which was approximately the same as the reported air temperature at the

time of collection (Supplementary Table 1), and shows a reduced thermal pressure compared to

the high surface temperatures at other areas within the city.

In the subsequent sample collections on September 4[th], 2023, June 21[st], 2024, and

August 28[th], 2024, we collected only from Fayette St. and Guilford sites to simplify the collection

and comparison process on the basis that they are on two ends of the heat-island landscape

within Baltimore City, with Fayette St. being amongst the warmest areas and Guilford being

amongst the coolest areas. In the subsequent fungal sample collection dates at the Fayette St.

and Guilford sites, surface temperatures recorded via infrared thermometers remained

consistent with what we observed on August 20[th], 2023, and September 4[th], 2023, using the

infrared camera (Supplementary Table 1).

We successfully isolated yeast and mold from all four sites on August 20[th], 2023, and

from Guilford and Fayette sites on the remaining collection dates, including the sidewalk and dirt

that reached exceedingly high surface temperatures. While fungi were isolated from samples

across all neighborhoods, the abundance of culturable fungi varied by location. Notably, for the

samples plated from sidewalk and dirt from four collection dates, less area of the petri dish

cultures was covered in mold and yeast in plates plated with material from the warmest (Fayette

St.) samples than the plates from the coolest (Guilford) site (Figure 3). While we were able to

recover viable fungi at these extreme temperature environments in cities, there was a reduced

amount of viable and culturable fungi present in the hotter areas. This method indicates that

there are fewer culturable fungi present in the warmer neighborhoods, but this method does not

account for the presence of species that may have a faster growth rate or species that may

have a naturally larger growth area, causing increased relative coverage of the media plate.

Pigments produced by fungi can absorb and retain heat, which impacts the cell

temperature and supports fungal growth in a heat-deprived environment through a process

described as "thermal melanism" [21]. Dark pigments in fungi are often composed of melanins,

which have a remarkable capacity to absorb electromagnetic radiation and convert it to heat

[18,21]. We compared mold pigmentation across the urban landscape from sidewalk and dirt on

all 4 collection dates and found that molds from the Fayette St. site (warm – 40°C-57°C) were

significantly less pigmented (lighter in color) than those from the Guilford site (cool – 27°C-

33°C), with mean gray values differing by ~10-pixel values (Figure 4A). Similarly, yeast from the

first two collection dates were about 12-pixel values lighter from Fayette St. (warm) compared to

Guilford site (cool) (Figures 4A and B). As an alternative approach, we directly assessed fungal

heat absorbance by mold colonies as measured by an infrared camera. Consistent with having

darker pigmentation, the mold found in the Guilford site (cool) absorbed more heat during a 10-

minute exposure to white light than the mold from Fayette St. (warm) (Figures 4C and D). We

would expect similar effects for yeast, though their small size precluded analyses by this

method. Three mold plates each from samples collected from Fayette St. and Guilford collected

on August 20th, 2023, and September 4th, 2023, were chosen at random. There was a

correlation between the thermal absorbance (temperature) following exposure to white light

(Figure 4C) and the mean gray value of the colonies (Figure 4E). This correlation is suggestive

of thermal melanism/thermal albinism, as these phenomena have been called, to regulate

thermal properties of the fungi [18,21]. These pigmentation patterns have been previously

described for fungi collected at different latitudes, where fungi from arctic regions display

enhanced pigmentation that promotes heat absorbance, while equatorial fungi have less

pigmentation to avoid excessive heat absorbance [18].

To further investigate and compare the thermal properties of the fungi collected from

warmer or cooler sites, we looked for culturable yeast of the same or closely related species by

PCR amplifying and sequencing the ITS sequence of the 107 collected fungal isolates. We

identified the yeast isolates by the species that most closely matched the Internal Transcribed

Spacer (ITS) DNA sequence on NCBI GenBank, usually with high sequence identity. Notably,

the presence of *Cystofilobasidium macerans, Rhodotorula spp., Cystobasidium spp.,* and other

fungi in the Cystobasidiomycetes class was found in all neighborhood sampling sites

(Supplementary Table 2). This class of yeast is ubiquitous in the environment, and some

species are rare human pathogens. The identification of the same fungal species in different

sampling sites allowed us to compare the characteristics of the same species of fungi when

exposed to different environmental pressures. We found *R. mucilaginosa,* a ubiquitous

environmental yeast and rare human pathogen, in both Fayette St. (warm) and Guilford (cool).

We tested whether the *R. mucilaginosa* isolate from the Fayette St. location (Isolate 200S) had

a different susceptibility to heat stress compared to those isolated from Guilford (414S, 423S,

and 432S) in laboratory settings. We found that while none were able to grow at 37°C, the

Fayette St. 200S isolate had greater resistance to killing following exposure to a 30-55°C and

30-56°C heat-ramp stress, remaining more viable compared to the three isolates from Guilford

(Figure 5A, B).

We found fungi belonging to the *Cystobasidium* genus, which includes *C. minutum,* a

rare opportunistic fungal pathogen [22]. These fungi were found at was found at the two

warmest sites, Fayette St and Mt. Vernon. One of the isolates of *C. minutum* recovered from the

38.4˚C sidewalk average temperature in Mt. Vernon (300S) was thermotolerant (able to grow at

37˚C), while another *C. minutum* isolate from Mt. Vernon sidewalk (305S) and a *C. lysinophilum*

isolate from Fayette St. dirt (242D) were unable to grow at the higher temperatures

(Supplementary Figure 2A). We performed similar experiments with *C. minutum* isolated from

Mt. Vernon sidewalk (300S and 305S) and found that the 300S Mt. Vernon isolate was more

resistant to heat-ramp at 55˚C and 60˚C (Supplementary Figure 2B,C). In fact, the 300S *C.*

minutum isolate was the only one able to grow after brief exposure to 30-60˚C heat-ramp.

Together, these data suggest that even within one geographic location, a subset of yeast

belonging to the same species are undergoing adaptation or selection for extreme heat.

Of the 44 yeast-like isolates collected from Fayette St. in our study on August 20[th], 2023

and September 4[th], 2023, 13 (29.5%) were identified as *Aureobasidium pullulans*. The identity of

this fungus was established using ITS sequencing, showing conserved ITS sequence to the

closest sequence match *A. pullulans* PE_11 strain (Supplementary Table 2, Supplementary

Figure 3). *A. pullulans* is notable as it is a polyextremotolerant fungus, including temperature,

saline stress, and nutrient stress, having first been isolated in glacial ice [23]. Microscopically,

we observed *A. pullulans* to have diverse morphology with hyphae, yeast-like structures, and

other abnormal shapes (Supplementary Figure 3). Additionally, 5 of the 44 isolates (11%) were

identified as *Zalaria obscura,* a fungus that is closely related to *A. pullulans* and has similar

resistances to environmental stressors [24]. Conversely, only 2 out of the 51 (4%) of the isolates

from Guilford were identified as *A. pullulans (*Supplementary Table 2*).* Identification of these

species in a neighborhood experiencing the most extreme heat in Baltimore City is thus a

notable finding and places further context in this fungus' role as a potential human pathogen.

Discussion

As areas vulnerable to extreme heat conditions, cities are potential crucibles for the

adaptation of fungi to warmer temperatures, and this can include fungal species with pathogenic

potential that could lead to the emergence of new human fungal pathogens. Here we

investigated the thermal tolerance of fungi in various neighborhoods of a city using a new to

collect and culture environmental fungi. We found that fungi from warmer neighborhoods

manifested different thermal tolerances and pigmentation from those of colder neighborhoods

and our study suggests that the extreme ranges in heat in neighborhoods within urban centers

may be sufficient to drive similar differences. The temperatures of the sidewalk and dirt in the

warmer Fayette St. location were 40˚C and 55˚C, respectively, which are both above the typical

temperature ranges of fungi, and above human body temperature (37˚C), thus providing a

selective pressure for higher heat tolerances that, if it selects for higher thermotolerance, could

allow them to survive at mammalian body temperature. Meanwhile, the temperatures at the

cooler, more wooded and shaded, Guilford neighborhood were around 27˚C, and thus did not

provide a selective pressure to for growth at or above mammalian body temperature. We noted

lower fungal mass growing in agar from warmer sites, consistent with a reduced mycotic flora at

those sites. Given that many fungal species are naturally hypothermic [25], and that the fungal

viability of fungal species declines rapidly about 30˚C [1], one might anticipate that the reduced

fungal census in warmer sites reflects a reduced ability to handle heat stress.

R. mucilaginosa isolated from Fayette St (warm) was better equipped to survive gradual

exposure to extreme heat (55˚C) compared to isolates of the same species isolated from the

Guilford site (cool). This suggests adaptation to the greater temperatures of the sidewalk at

Fayette, implying selection for *R. mucilaginosa* that can withstand extreme thermal

environments for short periods of time although all four were unable to grow at 37˚C. This

potential adaptation in *R. mucilaginosa* to heat is of interest since this fungus causes rare cases

of systemic infection in immunocompromised individuals [26–28], and thus demonstrates some

ability to withstand growth at human body temperature and host immune defenses. The addition

of enhanced thermotolerant properties due to extreme environmental heat may allow this fungus

to be even more equipped to survive within mammalian hosts. A relative of *Rhodotorula,*

Rhodosporidiobolus fluvialis, was recently found in two independent clinical infections in China

[29]. Researchers found that exposure of this fungus to 37°C conditions induced genetic and

phenotypic changes that caused the fungus to have enhanced thermotolerance, drug

resistance, and higher virulence phenotypes. These findings support the idea that fungi related

to *Rhodotorula spp.,* are in a position to adapt to a warming environment, which can lead to

enhanced frequency or severity of cases of disease with this fungus. *Rhodotorula spp.,* is found

in nearly every environmental niche across the globe, and is particularly enriched within the

guano of the lesser black-backed gull *Larus fuscus [30].* This finding is interesting for several

reasons. Firstly, the guano samples were specifically collected from urban sites suggesting that

gulls may disemminate *Rhodotorula* within cities throught their guano, as was with pigeons and

Cryptococcus neoformans [31]. Secondly, the internal temperature of *L. fuscus* is 41.2°C [32],

which is over human body temperature. This indicates that *Rhodotorula* may have a natural

ability to withstand, if not grow, at higher temperatures within an organism, and be primed for

inhabiting warm urban sidewalks upon extrection.

We found a strong presence of the extremotolerant fungus *A. pullulans* in hotter

neighborhoods. This fungus is capable of surviving saline, thermal, and nutrient-poor stresses,

and can also be an opportunistic human pathogen [33–35]. The presence of this fungus is thus

notable as it indicates the selection for extremotolerant fungal populations, which may correlate

to ability to tolerate other stressors including mammalian immune response. Gostinčar et. al.

has noted the potential importance of *A. pullulans* as an emerging pathogen due to its ability to

resist extreme conditions, hyperplasticity, and its presence within the indoor built environment

[23].

The previous comparative study of urban and rural fungi [16], along with the longitudinal

analysis of thermal tolerance in culture collection isolates [2] and our results in this study, are

each consistent with the notion that fungi are adaptingto our changing climate. We have found

evidence that urban fungi on the sidewalk and soil of warmer neighborhoods have adaptations

that allow them to mitigate heat and survive exposure to extreme heat. Molds and yeast in

warmer neighborhoods had reduced pigmentation to avoid absorbing heat, while species like *R.*

mucilaginosa, survive exposure to extreme temperatures (55°C) better than those from cooler

sites. Even within warmer sites, fungi like *C. minutum* appear to show varied adaptation to heat

where some isolates can survive up to a 60°C heat-ramp while another cannot, even though

they are presumably recently related. Within these environments, we also see

polyextremotolerant fungi such as *A. pullulans.* Many of these neighborhoods experiencing

extreme heat vulnerability (and the risk of emerging thermotolerance in fungi), are also often

inhabited by marginalized communities [10,12,36]. This is particularly worrisome, as these

communities are also the ones already most vulnerable to fungal infection due to healthcare

disparities and other social determinants of health [37].

In summary, our study shows differences in thermal tolerance for fungal species within a

city depending on the neighborhoods of origin and their average ambient temperatures. Our

findings warn that the heat islands defined by cities could be sites for rapid fungal adaptation to

higher temperatures, which brings the possibility that some fungal species with pathogenic

potential may be able to overcome human thermal defenses. Given that humans and fungi in

city ecosystems are in close proximity to one another and that the number of

immunocompromised patients are increasing yearly [38], rapid fungal thermal evolution in city

heat islands has the potential to bring us new fungal diseases in the years ahead.

Acknowledgements:

We would like to acknowledge C. Johnson, Gracen Gerbig, and Dr. Jenna Glatzer for the

assistance and support in collecting the fungal samples, Dr. Radames Cordero for the guidance

related to thermal melanism in the fungal isolates. Figure 1 was made using BioRender.com.

A.C. and D.F.Q.S. were supported in part by NIH grants AI162381, AI152078, and HL059842,

AI168539 and AI183596. Funders played no role in the experimental design or outcome of the

project.

References:

1.   Robert VA, Casadevall A. Vertebrate Endothermy Restricts Most Fungi as Potential

Pathogens. The Journal of Infectious Diseases. 2009;200: 1623–1626.

doi:10.1086/644642

2.   Robert V, Cardinali G, Casadevall A. Distribution and impact of yeast thermal tolerance

permissive for mammalian infection. BMC Biology. 2015;13: 18. doi:10.1186/s12915-015-

0127-3

3.   Smith DFQ, Casadevall A. Disaster Microbiology—a New Field of Study. mBio. 2022;13:

e01680-22. doi:10.1128/mbio.01680-22

4.   Casadevall A, Kontoyiannis DP, Robert V. On the Emergence of Candida auris: Climate

Change, Azoles, Swamps, and Birds. mBio. 2019;10. doi:10.1128/mBio.01397-19

5.   Garcia-Solache MA, Casadevall A. Global Warming Will Bring New Fungal Diseases for

Mammals. mBio. 2010;1: e00061-10. doi:10.1128/mBio.00061-10

6.   Valdez AF, Corrêa-Junior D, Bonilla JJA, Zamith-Miranda D, Frases S, Freitas DFS, et al.

A Review on Sporotrichosis and the Emergence of Sporothrix brasiliensis as a Pathogen.

Curr Trop Med Rep. 2023 [cited 20 Oct 2023]. doi:10.1007/s40475-023-00297-6

7.   Learn About Heat Islands | US EPA. [cited 11 May 2022]. Available:

https://www.epa.gov/heatislands/learn-about-heat-islands

8.   Sharma R, Hooyberghs H, Lauwaet D, De Ridder K. Urban Heat Island and Future Climate

Change—Implications for Delhi's Heat. J Urban Health. 2019;96: 235–251.

doi:10.1007/s11524-018-0322-y

9. Sachindra DA, Ng AWM, Muthukumaran S, Perera BJC. Impact of climate change on

urban heat island effect and extreme temperatures: a case-study. Quarterly Journal of the

Royal Meteorological Society. 2016;142: 172–186. doi:10.1002/qj.2642

10. Saverino KC, Routman E, Lookingbill TR, Eanes AM, Hoffman JS, Bao R. Thermal

Inequity in Richmond, VA: The Effect of an Unjust Evolution of the Urban Landscape on

Urban Heat Islands. Sustainability. 2021;13: 1511. doi:10.3390/su13031511

11. Metzger KB, Ito K, Matte TD. Summer Heat and Mortality in New York City: How Hot Is

Too Hot? Environmental Health Perspectives. 2010;118: 80–86. doi:10.1289/ehp.0900906

12. Popovich N, Flavelle C. Summer in the City Is Hot, but Some Neighborhoods Suffer More.

The New York Times. 9 Aug 2019. Available:

https://www.nytimes.com/interactive/2019/08/09/climate/city-heat-islands.html. Accessed 5

Nov 2023.

13. Detailed maps of urban heat island effects in Washington, DC, and Baltimore | NOAA

Climate.gov. [cited 5 Nov 2023]. Available: http://www.climate.gov/news-

features/features/detailed-maps-urban-heat-island-effects-washington-dc-and-baltimore

14. Tessler M, Neumann JS, Afshinnekoo E, Pineda M, Hersch R, Velho LFM, et al. Large-

scale differences in microbial biodiversity discovery between 16S amplicon and shotgun

sequencing. Sci Rep. 2017;7: 6589. doi:10.1038/s41598-017-06665-3

15. Nilsson RH, Anslan S, Bahram M, Wurzbacher C, Baldrian P, Tedersoo L. Mycobiome

diversity: high-throughput sequencing and identification of fungi. Nat Rev Microbiol.

2019;17: 95–109. doi:10.1038/s41579-018-0116-y

16.   McLean MA, Angilletta MJ, Williams KS. If you can't stand the heat, stay out of the city: Thermal reaction norms of chitinolytic fungi in an urban heat island. Journal of Thermal Biology. 2005;30: 384–391. doi:10.1016/j.jtherbio.2005.03.002

17.   Pereyra Irujo G. IRimage: open source software for processing images from infrared thermal cameras. PeerJ Comput Sci. 2022;8: e977. doi:10.7717/peerj-cs.977

18.   Cordero RJB, Robert V, Cardinali G, Arinze ES, Thon SM, Casadevall A. Impact of Yeast Pigmentation on Heat Capture and Latitudinal Distribution. Curr Biol. 2018;28: 2657-2664.e3. doi:10.1016/j.cub.2018.06.034

19.   Stolp ZD, Kulkarni M, Liu Y, Zhu C, Jalisi A, Lin S, et al. Yeast cell death pathway requiring AP-3 vesicle trafficking leads to vacuole/lysosome membrane permeabilization. Cell Rep. 2022;39: 110647. doi:10.1016/j.celrep.2022.110647

20.   Huang MY, Joshi MB, Boucher MJ, Lee S, Loza LC, Gaylord EA, et al. Short homology-directed repair using optimized Cas9 in the pathogen Cryptococcus neoformans enables rapid gene deletion and tagging. Genetics. 2022;220: iyab180. doi:10.1093/genetics/iyab180

21.   Cordero RJB, Casadevall A. Functions of fungal melanin beyond virulence. Fungal Biology Reviews. 2017;31: 99–112. doi:10.1016/j.fbr.2016.12.003

22.   Inácio CP, Diniz MV, Araújo PSR, Barros MS, Andrade MCL, Lima-Neto RG, et al. Bloodstream Infection of a Cancer Patient by Cystobasidium minutum: A Case Report and Literature Review. Mycopathologia. 2020;185: 395–398. doi:10.1007/s11046-019-00415-x

23.   Gostinčar C, Grube M, Gunde-Cimerman N. Evolution of Fungal Pathogens in Domestic Environments? Fungal Biology. 2011;115: 1008–1018. doi:10.1016/j.funbio.2011.03.004

24. Campana R, Fanelli F, Sisti M. Role of melanin in the black yeast fungi *Aureobasidium*

pullulans and *Zalaria obscura* in promoting tolerance to environmental stresses and to

antimicrobial compounds. Fungal Biology. 2022;126: 817–825.

doi:10.1016/j.funbio.2022.11.002

25. Cordero RJB, Mattoon ER, Ramos Z, Casadevall A. The hypothermic nature of fungi. Proc

Natl Acad Sci U S A. 2023;120: e2221996120. doi:10.1073/pnas.2221996120

26. Wirth F, Goldani LZ. Epidemiology of Rhodotorula: An Emerging Pathogen. Interdiscip

Perspect Infect Dis. 2012;2012: 465717. doi:10.1155/2012/465717

27. Jarros IC, Veiga FF, Corrêa JL, Barros ILE, Gadelha MC, Voidaleski MF, et al.

Microbiological and virulence aspects of Rhodotorula mucilaginosa. EXCLI J. 2020;19:

687–704.

28. Ioannou P, Vamvoukaki R, Samonis G. Rhodotorula species infections in humans: A

systematic review. Mycoses. 2019;62: 90–100. doi:10.1111/myc.12856

29. Huang J, Hu P, Ye L, Shen Z, Chen X, Liu F, et al. Pan-drug resistance and hypervirulence

in a human fungal pathogen are enabled by mutagenesis induced by mammalian body

temperature. Nat Microbiol. 2024;9: 1686–1699. doi:10.1038/s41564-024-01720-y

30. Cabodevilla X, Malo JE, Aguirre de Carcer D, Zurdo J, Chaboy-Cansado R, Rastrojo A, et

al. DNA Prevalence of Eukaryotic Parasites with Zoonotic Potential in Urban-Associated

Birds. Birds. 2024;5: 375–387. doi:10.3390/birds5030025

31. Spina-Tensini T, Muro MD, Queiroz-Telles F, Strozzi I, Moraes ST, Petterle RR, et al.

Geographic distribution of patients affected by Cryptococcus neoformans/Cryptococcus

493   gattii species complexes meningitis, pigeon and tree populations in Southern Brazil.

494   Mycoses. 2017;60: 51–58. doi:10.1111/myc.12550

495  32. van der Meer J, van Donk S, Sotillo A, Lens L. Predicting post-natal energy intake of lesser

496   black-backed gull chicks by Dynamic Energy Budget modeling. Ecological Modelling.

497   2020;423: 109005. doi:10.1016/j.ecolmodel.2020.109005

498  33. Bolignano G, Criseo G. Disseminated Nosocomial Fungal Infection by Aureobasidium

499   pullulans var. melanigenum: a Case Report. Journal of Clinical Microbiology. 2003;41:

500   4483–4485. doi:10.1128/jcm.41.9.4483-4485.2003

501  34. Hawkes M, Rennie R, Sand C, Vaudry W. Aureobasidium pullulans infection: Fungemia in

502   an infant and a review of human cases. Diagnostic Microbiology and Infectious Disease.

503   2005;51: 209–213. doi:10.1016/j.diagmicrobio.2004.10.007

504  35. Pikazis D, Xynos ID, Xila V, Velegraki A, Aroni K. Extended fungal skin infection due to

505   Aureobasidium pullulans. Clinical and Experimental Dermatology. 2009;34: e892–e894.

506   doi:10.1111/j.1365-2230.2009.03663.x

507  36. Reckien D, Lwasa S, Satterthwaite D, McEvoy D, Creutzig F, Montgomery M, et al. Equity,

508   Environmental Justice, and Urban Climate Change. 1st ed. In: Rosenzweig C, Solecki WD,

509   Romero-Lankao P, Mehrotra S, Dhakal S, Ali Ibrahim S, editors. Climate Change and

510   Cities. 1st ed. Cambridge University Press; 2018. pp. 173–224.

511   doi:10.1017/9781316563878.013

512  37. Jenks JD, Aneke CI, Al-Obaidi MM, Egger M, Garcia L, Gaines T, et al. Race and ethnicity:

513   Risk factors for fungal infections? PLOS Pathogens. 2023;19: e1011025.

514   doi:10.1371/journal.ppat.1011025

515   38.   Clark C, Drummond RA. The Hidden Cost of Modern Medical Interventions: How Medical

516   Advances Have Shaped the Prevalence of Human Fungal Disease. Pathogens. 2019;8:

517   45. doi:10.3390/pathogens8020045

Figure 1

Figure 1. Sample Collection Methods. A. Fungal sample collection from the sidewalk via Starburst. B. Fungal sample collection from dirt.

Figure 2

Figure 2. **Thermal Properties of Sampling Sites in Baltimore, MD.** A. Map of the four sites sampled in Baltimore, MD (Fells Point, Fayette, Mt. Vernon, Guilford). B. Google StreetView images from the sampling sites taken during summer months in recent years. Infrared images of the sidewalks (C) and dirt (D) on August 20th, 2023. The visible light photo from the FLIR 96E camera overlayed on the generated thermal image (top rows of C and D), and the raw grayscale temperature image standardized to 27˚C (min) and 60˚C (max) with the mean temperature from the center (approximate area shown with a red circle) of the individual image measured overlayed (bottom rows of C and D).

Figure 3

Figure 3. Fungal growth is sparser from Fayette St. (warm) Samples than Guilford (cool) Samples. Less plate surface area is covered by fungal growth from samples collected from Fayette St. compared to those collected at Guilford. Error bars represent 95% CI from the mean. Samples were collected on four collection dates. Each dot represents the percentage of agar area covered in fungal growth from one plate. Statistical significance was calculated with an unpaired t-test.

Figure 4

Figure 4. Fungi collected in thermally diverse neighborhoods exhibit thermal and pigment differences. The mean gray value of mold (A) and yeast (B) isolated from Fayette St. is higher than the mean gray value of the fungi isolated from Guilford, indicating less pigmentation in the Fayette St. fungi. Each dot represents values from individual colonies. Following exposure to light, fungi from the Guilford (cool) samples get warmer than those from the Fayette samples (Panels C and D). Correlation between colony temperature and the mean gray value, where the darker colonies get warmer (Panel E). Panels A and B show mean values with 95% CI and t-tests, while C shows median temperature and non-parametric Mann-Whitney test. Each dot represents mean gray value (Panel A, B, E) or temperature value (Panel C, E) for an individual mold (Panel A, C, E) or yeast colony (Panel B). Samples in (A) were collected from August 20th, 2023, September 4th, 2023, June 21st, 2024, and August 28th, 2024. Samples in (Panels B to E) were collected on August 20th, 2023, and September 4th, 2023.

Figure 5

Figure 5. R. mucilaginosa from warmer neighborhoods has greater resistance to heat ramp stress.
R. mucilaginosa from the sidewalk of Fayette St. (200S) show increased resistance to heat-ramp cell death stress compared to *R. mucilaginosa* strains collected from the cooler Guilford sidewalk (414S, 423S, and 432S) after gradual heat ramp to 55°C (Panels A and B). Significance of Two-Way ANOVA with multiple comparisons to the 200S for A and B at each condition with three biological replicates. Each dot represent CFUs from an individual biological replicate. **** represents p<0.0001, ** represents p<0.01, and ns represents p >0.05.

Supplementary Figure 1

	Technical Rep #1	Technical Rep #2	Technical Rep #3
Starburst Control 1			
Starburst Control 2			
Starburst Control 3			

Supplementary Figure 2

Supplementary Figure 2. **Thermotolerance of Cystobasidium spp. (A)** Cystobasidium spp. isolated from Fayette St. dirt (242D) and Guilford sidewalks (*305*S) were unable to grow at 37˚C, but *one* C. minutum isolate from Mt. Vernon sidewalk (30*0*S) *was* thermotolerant at 37˚C. *C. minutum* from the sidewalk of Mt. Vernon (300S) grows better than the other C. minutum isolated from the Mt. Vernon sidewalk (305S) when exposed to gradual 55˚C heat ramp, while only 300S can grow following exposure to 60˚C heat ramp (Panels B and C). *S*ignificance of Two-Way ANOVA with multiple comparisons to the 300S for B and C at each condition with three biological replicates. Each dot represent CFUs from an individual biological replicate. **** represents p<0.0001, ** represents p<0.01, and ns represents p >0.05.

Supplementary Figure 3

Supplementary Figure 3. **Identification of Aureobasidium pullulans from High-Temperature Environment**. Alignment of three of the 15 A. pullulans cultures isolated from Fayette St. sidewalk (208S, 219S, 220S) to the A. pullulans PE_11 isolate (A), and their phylogenetic relationship (B). Microscopic images of the *A. pullulans* cultures show hyphal, yeast-like, and irregular shaped (C). Scale bar represents 20 *μ*m.