Tonic dopamine and biases in value learning linked through a biologically inspired reinforcement learning model

Sandra Romero Pinto[1,2], Naoshige Uchida[1]

Affiliations:

[1]Department of Molecular and Cellular Biology, Center for Brain Science, Harvard University, Cambridge, MA 02138, USA

[2]Program in Speech and Hearing Bioscience and Technology, Division of Medical Sciences, Harvard Medical School, Boston, MA 02115, USA

Correspondence: uchida@mcb.harvard.edu (N.U); sromeropinto@g.harvard.edu (S.R.P)

Abstract

A hallmark of various psychiatric disorders is biased future predictions. Here we examined the mechanisms for biased value learning using reinforcement learning models incorporating recent findings on synaptic plasticity and opponent circuit mechanisms in the basal ganglia. We show that variations in tonic dopamine can alter the balance between learning from positive and negative reward prediction errors, leading to biased value predictions. This bias arises from the sigmoidal shapes of the dose-occupancy curves and distinct affinities of D1- and D2-type dopamine receptors: changes in tonic dopamine differentially alters the slope of the dose-occupancy curves of these receptors, thus sensitivities, at baseline dopamine concentrations. We show that this mechanism can explain biased value learning in both mice and humans and may also contribute to symptoms observed in psychiatric disorders. Our model provides a foundation for understanding the basal ganglia circuit and underscores the significance of tonic dopamine in modulating learning processes.

Introduction

Our ability to predict the outcomes of our actions is crucial in selecting and motivating appropriate actions. Systematic biases in future predictions or expectations, however, can lead to maladaptive behaviors, such as those observed in patients with various psychiatric disorders[1–4]. For example, overly negative or pessimistic predictions can contribute to major depression[1,5], whereas excessively positive or optimistic predictions may be associated with pathological gambling, addiction, and mania[3,4,6–8] . Despite the importance of understanding the causes of biased future predictions, the biological mechanisms underlying them remain poorly understood.

Our future expectations and decisions are shaped by experiences of positive and negative events. The process of learning from outcomes has been modeled using reinforcement learning (RL) models[9–12], where value predictions are updated based on reward prediction errors (RPEs), that is the discrepancy between received and expected outcomes. In addition to its role in learning,

recent studies have indicated the importance of RPEs in mood; these studies have suggested that

mood depends not on the absolute goodness of outcomes, but rather on the recent history of

RPEs[13,14].

In the brain, dopamine is thought to be a key regulator in this process of learning from positive

and negative outcomes. The dynamics of dopamine are often categorized into two modes: tonic

and phasic. Tonic dopamine refers to "baseline" dopamine that operates on a long timescale such

as tens of seconds or minutes, while phasic activity refers to transient changes that occur at a

much shorter, sub-second timescale, often triggered by external stimuli[15–18]. A significant body

of evidence has shown that phasic responses of dopamine neurons convey RPEs and drive

learning of values and actions[17–20]. On the other hand, changes in tonic dopamine might also

modulate value learning, yet whether and how the level of tonic dopamine modulates learning

remain poorly understood.

Previous studies have reported that patients with psychiatric disorders exhibit biased learning

from positive versus negative outcomes. For one, some studies have shown that patients with

major depression have a reduced sensitivity in learning from rewarding events, while their ability

to learn from negative events remains relatively intact[1,5,21]. Similarly, patients with Parkinson's

disease are better at learning from negative than positive outcomes[22,23]. Analysis of these patients

using RL models has suggested that biases in learning can be explained by alterations in specific

parameters in RL models, such as the learning rate parameters or the sensitivity to positive and

negative outcomes. For example, some studies have suggested that anhedonia in major

depressive disorder may correspond to a reduced learning rate from positive compared to

negative outcomes[1].

Mechanistically, some of these changes in RL parameters can be linked to altered functions of

dopamine. First, it has been shown that dopamine synthesis capacity, an approximate indicator of

baseline dopamine levels, in the striatum, as measured using positron emission tomography

(PET), correlates with learning rate parameters[24]. Second, dopamine medications can change the

balance between learning from positive and negative outcomes[22,24,25]. Third, responses to

positive outcomes in the nucleus accumbens (NAc), as measured based on blood oxygenation-

dependent (BOLD) signals, are reduced in patients with psychiatric disorders such as

depression[26–29] . These observations point to important roles of reinforcement learning processes

and dopamine in regulating value learning. However, the parameters in RL models remain an

abstract entity, and biological processes underlying changes in these parameters are still largely

unknown.

One limitation in most RL models used in previous studies is that they do not reflect key neural

circuit architectures in the brain (but see [30–32]) nor recent findings on intracellular signaling and

plasticity rules that can constrain how dopamine functions in biological circuits[33–35].

Incorporating these key biological factors may lead to better understanding of how changes in

RL parameters may arise in psychiatric disorders. Furthermore, recent studies have found that

the activity of dopamine neurons is consistent with a novel RL algorithm called distributional

RL[36–38]. Distributional RL takes into account the diversity in dopamine signals, and a population

of dopamine neurons together encodes the entire distribution of rewards, not just the average.

Although distributional RL has shown to be efficient in solving various RL problems in artificial

intelligence[37,39], how distributional RL can be implemented in biological neural circuits and how

distributional RL relates to biased value learning remain to be examined.

In this study, we sought to identify potential mechanisms that cause biased value predictions

using biologically inspired RL models. To this goal, we first construct an RL model that

incorporates recent biological findings, such as intracellular signaling and synaptic plasticity

rules as well as the basic circuit architecture in the brain[32]. Based on this model, we propose two

potential biological mechanisms that can cause optimistic or pessimistic biases in value

predictions. We will then show that some existing data can be explained by one of these models.

Finally, we will show how our model can provide an account of how biases in value predictions

arise in psychiatric disorders.

Results

Basic reinforcement learning algorithms

Here we first formulate basic RL algorithms that will become the basis of our later models. In

RL, an agent learns to predict the expectation of future rewards associated with a given state, a

quantity termed as *value*[11]. For simplicity, we will drop the dependency on time here, but note

that the basic results hold even if time is considered (Methods 1.1). Learning of value is driven

by RPEs (δ), the discrepancy between the actual and expected reward (r and V, respectively)

(Eq1). To improve the accuracy of the value prediction, RPEs are utilized to update the estimate

of V. This is done iteratively by adding a fraction (α) of δ (Eq2) where α defines the learning

rate.

$$\delta = r - V \tag{1}$$

$$V \leftarrow V + \alpha \cdot \delta \tag{2}$$

When the magnitude of reward r is fixed (i.e., deterministic environment), the value V learned

through this algorithm (Eq 1 and 2) converges on r and the RPE converges on zero. When the

magnitude of reward r varies stochastically trial-to-trial, the value at convergence fluctuates

around the expected value of the reward distribution (see Methods 1) (Fig. 1a) and the RPE

around zero.

Risk-sensitive RL. In the framework called risk-sensitive RL[40], learning rates are defined

separately for positive and negative RPEs (denoted by α^+, α^-).

$$V \leftarrow V + \alpha^+ \cdot \delta \quad \text{if } \delta > 0 \tag{3}$$

$\qquad V \leftarrow V + \alpha^- \cdot \delta \;\; \text{if } \delta < 0$

In the presence of stochastic rewards, when the learning rates between positive and negative

RPEs are different, the value learned through this algorithm (Eq 1 and 3) does not converge on

the expected value, but instead on a value higher or lower than the expected value depending on

the relative amplitude of the learning rates α^+, α^-. This algorithm, therefore, develops optimistic

or pessimistic value expectations, respectively. This learning algorithm is called "risk-sensitive"

because values of probabilistic (risky) rewards are biased compared to deterministic (certain)

rewards, and, therefore, the agent develops a preference between risky and certain rewards even

when the expected values are the same (Fig. 1b).

Distributional RL. The concept of asymmetric updates has been utilized in a novel RL

framework called distributional RL[36,37,41]. This algorithm allows an agent to learn the entire

probability distribution of rewards, instead of the expected value which is typically the learning

target in traditional RL algorithms (Fig. 1c). In distributional RL, an agent is equipped with a set

of value predictors (V_i), where i corresponds to the index of the value predictor (or "value

neuron"). The value of the i-th neuron (V_i) is updated based on the learning rates (α_i^+, α_i^-) and

the RPE (δ_i) for that neuron i:

$\qquad V_i \leftarrow V_i + \alpha_i^+ \cdot \delta_i \;\; if \; \delta_i > 0$ $\qquad\qquad\qquad\qquad$ (3)

$\qquad V_i \leftarrow V_i + \alpha_i^- \cdot \delta_i \;\;\; if \; \delta_i < 0$

Similar to risk-sensitive RL, the learned value of each value predictor converges on estimates

larger or lower than the expected value, determined by the ratio between α_i^+ and α_i^-.

Mathematically, each V_i converges on the τ_i-th expectile of the distribution (Fig. 1c) where τ_i

(asymmetric scaling factor) is defined by:

$$\text{Asymmetric scaling factor: } \tau_i = \frac{\alpha_i^+}{\alpha_i^- + \alpha_i^+} \qquad\qquad (4)$$

Expectiles are the solutions to asymmetric least squares minimization and generalize the mean of

a distribution (with the mean being the 0.5^{th} expectile) as quantiles generalize the median (with

the median being the 0.5^{th} quantile)[42]. Since a set of expectiles can define a distribution, the

diversity of τ_i across the population enables learning of the entire probability distribution.

Problem. In both risk-sensitive RL and distributional RL, unbalance in learning rate parameters

for positive and negative RPEs gives rise to optimistic and pessimistic biases in learned values.

Importantly, however, the underlying biological mechanism regulating learning rate parameters

(α^+, α^-) and asymmetry thereof (τ) remains unclear.

In the following sections, we will discuss potential biological mechanisms that regulate

asymmetric learning rates (α^+, α^-). We will first modify the above RL algorithms to incorporate

important neural circuit architectures in the brain. We will then propose two key biological

mechanisms that can give rise to asymmetric learning rates (called Model 1 and 2). We will then

show that our model can explain previous experimental data and psychiatric conditions.

Incorporating biological features into RL models

The above RL models provide algorithmic-level formulations, yet they do not recapitulate

fundamental characteristics of the neural circuits thought to perform RL in the brain[43–46]. We

next incorporate some of the important circuit and synaptic properties into the model.

In the brain, it is thought that dopamine neurons in the ventral tegmental area (VTA) broadcast

RPEs[17] and modulate synaptic plasticity in dopamine-recipient areas[33,47]. The striatum is the

major target of dopaminergic projections. It has been thought that spiny projection neurons

(SPNs) in the striatum represent values, and dopamine modulates plasticity of synapses on

SPNs[33,34,47,48] (Fig. 2a). Under this framework, the value representations in SPNs are updated by

dopaminergic RPEs. In most RL models, each value predictor is typically updated by both

positive and negative RPEs. If the value is computed based on a weighted sum of some inputs

(i.e., using linear function approximation[11]), the update rules described above (Eq 3 and 4) are

equivalent to performing a semi-gradient descent that minimizes RPEs[11] (see Methods).

The basic architectural assumptions of these RL models are, however, at odds with the RL

circuitry in the brain. Importantly, in the striatum, there are two major classes of dopamine-

recipient SPNs characterized based on the type of dopamine receptor that they express: D1- or

D2-type dopamine receptors (D1R and D2R)[48]. SPNs expressing D1R and D2R constitute the so-

called direct and indirect pathways and exert opposing effects on downstream "output" neurons,

with each pathway promoting or opposing a certain output (e.g., movement).

In addition to the presence of direct and indirect pathways, there are two additional properties in

these opposing populations that need to be considered[32]. First, D1R and D2R have different

affinities to dopamine: high in D2R and low in D1R (EC$_{50}$ affinity constant is 1 µM for D1R and

10 nM for D2R)[49,50]. The dose-occupancy relationship of D1R and D2R are sigmoidal but they

are shifted with one another with respect to dopamine concentration (Fig. 2b). Importantly, at

normal dopamine levels (approx. 50-100nM)[51,52], D2Rs are mostly occupied while D1Rs are

mostly unoccupied (Fig. 2b). Although whether the affinities of D1R and D2R differ at the

molecular level has been questioned[53], a recent study showed that intracellular signaling through

protein kinase A (PKA) in D1- and D2-SPNs is triggered by a phasic increase and a decrease in

dopamine, respectively, in behaving animals[35]. These results are consistent with (apparent)

difference in affinities of D1R and D2R assumed in previous studies[49], although the exact reason

for the difference remains to be clarified[53].

The second important property pertains to different learning rules in D1- and D2-SPNs which are

predicted from different affinities of the receptors. Consistent with the observed PKA signals in

these cells, recent studies have shown that glutamatergic inputs on D1-SPNs are potentiated by a

transient *increase* in dopamine, whereas those on D2-SPNs are potentiated by a transient

decrease in dopamine[34,35] (Fig. 2c), supporting opposing plasticity rules between D1- and D2-

SPNs.

There have been previous efforts to incorporate in RL models the direct and indirect pathways

(also called "Go" and "NoGo" pathways, respectively) such as Opponent Actor Learning

(OpAL[30], OpAL*[54]) and Actor learning Uncertainty (AU)[32] models. These previous models

were developed as *Actor-Critic models*[11]. Here, we will build on the AU model to focus on the

problem of value learning and extend it to support risk-sensitive RL and distributional RL. Our

model has two separate populations of value predictors corresponding to D1R- and D2R-SPNs,

that store the quantities Pi and N_i respectively (Eq 6, Fig. 2d). Mimicking dopamine's effect on

potentiation, P_i or N_i will increase their estimates if an RPE is positive or negative, respectively,

with the learning rates defined by α_i^+, α_i^- (Eq. 6). Importantly, the value V_i can be obtained

simply by taking the difference between P_i and N_i. (Eq. 7).

$$\text{D1R-SPN: } P_i \leftarrow P_i + \alpha_i^+ \cdot |\delta_i| - \beta \cdot P_i \quad if \ \delta_i > 0 \tag{5}$$

$$\text{D2R-SPN: } N_i \leftarrow N_i + \alpha_i^- \cdot |\delta_i| - \beta \cdot N_i \quad if \ \delta_i < 0 \dots$$

$$\text{Value: } V_i = P_i - N_i \tag{6}$$

where β is a decay parameter which represents synaptic decay in the absence of RPEs. This

model (Eq 6 and 7) preserves various essential properties of the previous RL models: (1)

learning in P and N can be combined to provide a simple update rule for value V, and (2) this

update rule approximates the gradient descent that minimizes RPEs (when $\beta = 0$, the update rule

is equivalent to the gradient descent). Importantly, with $\beta > 0$, we can show that these simple

learning rules guarantee convergence of value, without the need for additional mechanisms to

modulate the learning rates over iterations (Methods 1.3).

For instance, in a stochastic environment where there is a probability p of receiving a reward of a

fixed magnitude $r = 1$, the stochastic fixed point of the learned value V_i (i.e., convergence point)

will be defined by Eq 7.

$$V_i = \frac{\frac{\tau_i}{1-\tau_i} \cdots \frac{p}{1-p}}{\frac{\tau_i}{1-\tau_i} \cdots \frac{p}{1-p} + 1 + C} \cdot r \text{ , where } C = \frac{\beta}{(1-p) \cdot (1-\tau)} \tag{7}$$

Note that Eq.7 contains an additional term C which depends on β and this decay factor β is

important to stabilize the P_i and N_i estimates (avoid infinite increases) (Methods, 1.3.1, Extended

Data Fig. 1).

This formulation now provides a mechanistic model suitable for risk-sensitive RL (when there is

one value predictor) as well as distributional RL (when there are multiple value predictors),

which incorporate the neural circuit architecture and plasticity rules of D1R- and D2R-SNPs

found in the brain.

With this model at hand, we now discuss potential mechanisms that produce an asymmetry in

learning rates α_i^+, α_i^- , which, in turn, causes biases in value predictions. In principle, learning

rate parameters can be a function of (1) the scaling of RPEs, i.e., the slope of dopamine

responses as a function of RPE (δ), and (2) the scaling of value updates, i.e., the efficacy of

dopamine-dependent synaptic plasticity at the level of SPNs. In the following, we discuss each

scenario, emphasizing the role of either tonic or phasic dopamine activity in each of these

mechanisms (Model 1 and 2, respectively). For simplicity, we will start with a model in which

α^+, α^- are equal for all neurons within both P and N populations, equivalent to risk-sensitive

RL. We will then relax this assumption and introduce heterogeneity by allowing α_i^+, α_i^- to vary

across neurons, implementing a form of distributional RL.

Model 1: The role of baseline dopamine in asymmetric learning

As discussed above, D1R and D2R have different affinities to dopamine which leads to different levels of receptors' occupancy at a given baseline dopamine level (Fig. 2b). Crucially, due to the sigmoidal shape of the dose-occupancy curves, the slope of the curve changes with baseline dopamine level, which means that a given dopamine transient leads to a different change in receptor occupancy depending on the baseline dopamine level (Fig. 3a,b). That is, the receptors' *sensitivity* changes with baseline dopamine (Fig. 3c). In addition, a key consequence of the distinct receptors' affinities is that an increase and decrease in baseline dopamine will cause opposite changes in the sensitivity of D1R and D2R. Specifically, an increase in dopamine will decrease D1R sensitivity relative to D2R, whereas a decrease in dopamine will increase D2R sensitivity relative to D1R (Fig. 3c,d).

Building on this insight in Model 1, we postulate that the learning rates for positive and negative RPEs are a function of the D1R and D2R sensitivity, respectively. This is supported by previous studies that have reported that the effect of dopamine transients of a given magnitude in SPNs' plasticity can be modulated by the level of dopamine baseline[34]. In addition, it has been reported that the level of potentiation in SPNs[33,34] or plasticity, which are related to intracellular signals[35], scale with the magnitude of dopamine transients, keeping all else fixed. These observations can be summarized with the following rule:

$$LTP_{D1} \approx \alpha^+ \cdot |DA_{burst}|$$

$$LTP_{D2} \approx \alpha^- \cdot |DA_{pause}|$$

Where α^+, α^- correspond to the receptors' sensitivities and depend on the dopamine baseline level. This rule can be directly related to the update equations for the P and N populations in our model:

$$\Delta P = \alpha^+ \cdot |\delta| \, ... \, \text{if} \ \ \delta \geq 0$$

$$\Delta N = \alpha^- \cdot |\delta| \, ... \, \text{if} \ \ \delta < 0$$

It can been shown that this learning rule is in agreement with the normative solution for the

problem of value learning[11] (Methods 1.6).

In short, in Model 1, a shift in the baseline dopamine level causes asymmetries in scaling of the

value updates for positive versus negatives RPEs via the modulation of receptors' sensitivities,

which leads to value learning biases. This is a direct consequence of the dose occupancy

relationships of D1R and D2R (Fig. 3b-d).

Model 2: Asymmetric scaling of phasic dopamine responses, inspired by distributional RL

In Model 2, we postulate that the learning rates α^+ and α^- are a function of the scaling (i.e.,

'slope') of dopamine responses evoked by positive and negative RPEs, respectively:

$$DA_{burst} = \alpha^+ \cdot \delta \ ... \text{if} \ \ \delta \geq 0$$

$$DA_{pause} = \alpha^- \cdot \delta \ ... \text{if} \ \ \delta < 0$$

This is supported by a previous study on distributional RL that demonstrated that individual

dopamine neurons vary in terms of how the magnitude of reward responses is scaled as a

function of positive and negative RPEs (Fig. 3e)[36].

In the distributional RL framework, individual dopamine neurons vary in terms of their

asymmetric scaling factor τ_i and each of the multiple value predictors (V_i) converges on the τ_i-th

expectile of the reward distribution (Eq. 4). However, in most applications of distributional RL,

action selection is still based on the expected value of the reward distribution. Thus, the quantity

relevant to action selection can be described using the population level average $\tau_{population}$, and

biased value learning at the behavioral level could arise if $\tau_{population}$ is higher or lower than 0.5.

This can occur from a differential loss of optimistic or pessimistic dopamine neurons. Another

possibility is an overall upward or downward shift in the distribution of τ_i across the population

due to, for example, intrinsic factors modulating the gain of dopamine phasic responses.

Risk-sensitive RL can be thought of as a special case of distributional RL which has only one

value predictor. Here, the slope of the average dopamine evoked transient to positive and

negative RPEs, may correspond to the population level learning rates for positive and negative

RPEs (α^+, α^-), respectively. If the asymmetric scaling factor τ is higher or lower than 0.5, value

learning will be biased (Fig. 3e).

Testing for evidence of either model in experimental data

Tian and Uchida (2015).

We next examined whether Model 1 or 2 can explain empirical data obtained in experimental

animals or humans. We first examined the data obtained in mice in our previous study[55]. In this

study, the authors tested the effect of lesioning the habenula, a brain structure implicated in

depression[56–58], on the activity of dopamine neurons and on reward-seeking behavior. Head-fixed

mice were trained in a Pavlovian conditioning task in which odor cues predicted reward with

different probabilities (10%, 50%, 90%). After performing habenula (n=5) or sham (n=7) lesions

(Fig. 4a), the spiking activity of VTA dopamine neurons was recorded while mice performed the

task.

After lesions, mice exhibited an elevated reward-seeking behavior (anticipatory licking) in

response to cues predictive of probabilistic rewards, consistent with an optimistic bias in reward

expectation (Fig. 4b, right). Importantly, anticipatory licking gradually increased over several

sessions after lesions, suggesting that the optimistic bias developed through learning (Fig. 4b,

left). To bring insight into the underlying cause of these biases, we fit two different RL models to

the anticipatory lick responses on a trial-by-trial basis (Extended Data Fig. 2), assuming a linear

relationship between value predictions and anticipatory licking. These models considered either a

change in the sensitivity to rewards (Extended Data Fig. 2b) or asymmetric learning rates

(Extended Data Fig. 2c). This analysis showed that the biases observed in the behavior could be

explained by asymmetric learning rates, but not by reward sensitivity because the reward

sensitivity was unchanged in the lesion group with respect to the control group (Extended Data

Fig. 2c).

Dopamine neurons' responses to reward-predictive cues reflect the increases in value expectation

predicted by the cue with respect to baseline. The overall magnitudes of cue-evoked responses

were not elevated in lesioned animals compared to control animals (Fig. 4d). However, the shape

of the response curve indicated an 'optimistic' bias: although in control animals, cue responses

scaled linearly with the expected value (i.e., reward probability), the response function of the

lesioned animals was convex. In other words, in control animals the response to the 50%-reward

cue was not significantly different from the quantity that results from the linear interpolation

between the responses to 10%- and 90%-reward cues. In lesioned animals, however, the response

to the 50%-reward cue was significantly greater than this quantity, which is indicative of an

optimistic bias in value predictions (Fig. 4d, see Methods 1.3.3 for analysis of value predictions

curve convexity). Such a change was observed at the level of the population average. Further

analysis using individual neurons showed that when calculating a single-cell level metric that

compares the 50%-reward cue to the same linear interpolation point, there was a broad

distribution in this metric below and above the interpolated point both in the control and lesion

groups (Fig. 4e-f). The distribution was, however, shifted in its mean in the lesion group (Fig.

4e). These analyses indicated that both anticipatory licking and dopamine cue responses have an

optimistic bias as characterized by an overvaluation of probabilistic rewards, without still

pointing to the underlying mechanism.

Model 2 cannot explain the optimistic biases in behavior and cue-evoked dopamine

responses after Hb lesions

In Model 2, an optimistic bias in reward expectation can arise if the average of the asymmetric

scaling factor at the population level ($\tau_{population}$) becomes greater than 0.5 (Fig. 5a,b).

To test this idea, we obtained the asymmetric scaling factors (τ_i) from dopamine neurons based

on their outcome responses: for each neuron, we constructed outcome response functions against

the magnitude of RPEs (Fig. 5c, Extended Data Fig. 3a,b). The response functions were obtained

based on (1) whether reward was delivered (positive RPEs) or not (negative RPEs), and on (2)

the magnitude of the reward expectation given by the reward probabilities predicted by each cue

(0.1, 0.5, 0.9) (Extended Data Fig. 3a,b). We then obtained the point at which the responses are

more likely to be below or above baseline (i.e., 'zero-crossing points')[36] (Extended Data Fig. 3c),

and computed α_i^+ and α_i^- as the slopes of the responses in the positive and negative domains

with respect to this zero-crossing point (Extended Data Fig. 3d), respectively. In both control and

lesioned animals, asymmetric scaling factors tiled a wide range between 0 and 1 and presented

other signatures consistent with distributional RL[36] (Extended Data Fig.4). Nonetheless, although

the variance of the distribution of asymmetric scaling factors was greater in lesioned animals, the

mean did not change, indicating a lack of bias between α_i^+ and α_i^- at the population level (Fig.

5d). This was also the case when the asymmetric scaling factor was derived directly from the

population average response (Fig. 5c). Thus, contrary to the conclusion in our previous study[15],

these analyses indicated that changes in reward responses (and the resulting scaling factor τ) do

not explain the optimistic biases in behavior nor cue responses in lesioned animals (Fig. 5e,f).

Model 1 can explain the optimistic biases in behavior and cue-evoked dopamine responses

by Hb lesion

In addition to changes in the magnitude of dopamine RPEs, we observed that the baseline firing

rates of dopamine neurons were elevated in lesioned animals (Fig. 6a). According to Model 1, if

these changes are followed by an increase in the baseline dopamine levels in the striatum, this

should give rise to biased value learning ($\alpha^+ > \alpha^-$) and an optimistic bias in value expectation.

In this way, this change in baseline firing can explain optimistic biases observed in lesioned

animals. However, it remains unclear whether the observed change in baseline firing can result in

functionally relevant levels of changes in the receptor occupancies discussed above.

To quantitatively predict dopamine concentrations in the striatum and resulting receptor

occupancies of D1R and D2R, we used a biophysical model commonly used in the field[59]

(Fig.6a). This model has the firing rate of dopamine neurons as its input, and considers diffusion

of dopamine, dopamine reuptake, and D2-autorreceptor-mediated inhibition of dopamine release

to predict the dopamine concentration in the striatum (Fig. 6b,e). In addition, it considers the

affinities of D1R and D2R to estimate their occupancy levels (Fig. 6c,f). After estimating these

two variables (dopamine concentration and receptor occupancy), we derived the receptor

sensitivities (Fig. 6g-h). The receptor sensitivities were quantified as the slope of the resultant

changes in receptor occupancy given the observed baseline and phasic responses of dopamine

neurons. We then trained Model 1 using the receptor sensitivities as learning rates (α^+ and α^-)

for both control and lesioned animals.

The biophysical model indeed supported that the observed change in dopamine neuron firing can

cause a significant increase in dopamine concentration (Fig. 6e) and in D1 and D2 receptor

occupancies at baseline (Fig. 6g). These changes are expected to cause a significant asymmetry

in receptor sensitivities favoring D1 receptors over D2 receptors (Fig. 6h-i).

These receptor sensitivities were directly used as the asymmetric learning rates in a temporal-

difference (TD) learning version of Model 1 (see Methods 1.3, 3.3). After training, the model

incorporating the predicted asymmetries in learning rates (α^+, α^-) produced optimistic biases in

value predictions and in normalized cue responses, similar to those observed in lesioned animals

(Fig. 6k-l). The model simulating control animals developed no significant biases.

Additionally, the overall decrease in the magnitude of cue responses, observed in lesioned

animals, was reproduced in Model 1 using TD learning (Fig. 6k). This occurs because TD

learning calculates RPEs based on the change in values between before and after cue

presentation, and the "baseline" (pre-cue) reward expectation was also increased by optimistic

value learning (Fig. 6k). These results, together, indicate that Model 1 provides a parsimonious

account of the data: a change in baseline firing of dopamine neurons, rather than changes in

phasic responses, is the likely mechanism that led to optimistic biases in reward-seeking

behavior as well as cue-evoked dopamine responses in habenula lesioned animals.

Model 1 and model 2 play complementary roles in the encoding of asymmetric learning

rates

Although Model 2 did not explain the optimistic biases in the data in habenula-lesioned mice, the

distributional RL version of Model 2 explained other features of the data (Extended Data Fig. 3-

4). As mentioned, in both control and lesioned animals, asymmetric scaling factors tiled a wide

range between 0 and 1[36] (Extended Data Fig.4). Furthermore, cue-evoked responses of individual

neurons showed a wider distribution than what is expected by noise (Figure 4d). Finally, the core

prediction of distributional RL – a positive correlation between the asymmetric scaling factors of

the RPE responses of individual dopamine neurons and their zero-crossing points[36] – was also

present in controls and after Hb lesions. Together these results support that the basic features of

distributional RL are present in a way consistent with Model 2.

To complement this analysis, we tested whether Model 2 could have explained the signatures of

the data if asymmetric scaling factors (τ) derived from dopamine responses were indeed overall

biased (Extended Data Fig. 5). As expected from the model's fixed-point analysis (Methods 1.3),

if we imposed a shift in the mean of the distribution of asymmetric scaling factors (i.e.,

$\tau_{population} > 0.5$), the value predictors indeed exhibited optimistic biases (Extended Data Fig.

5e,f). However, the model did not reproduce the optimistic bias in cue-induced TD errors

observed in the data (Extended Data Fig.5g,h). This is due to an interaction of the biases in

prediction at "baseline" (pre-cue) and the cue, together with the optimistic asymmetry in the

scaling of the TD errors at cue themselves. Importantly, this was found in both versions of Model

2, distributional and risk-sensitive RL (Extended Data Fig. 5a-d and e-h). The difficulty of

explaining biased dopaminergic cue responses further makes the Model 2 an unlikely mechanism

to explain the optimistic biases in the data.

Altogether the data supports a model in which the mechanisms of Model 1 and 2 play

complementary roles in the encoding of asymmetric learning rates. The mechanism of Model 2

explains the variability in single neuron responses, consistent with the expectile code in

distributional RL. On the other hand, the mechanism of Model 1 at the population level,

generating asymmetries in learning rates and biases in value expectations, which might require

context-dependent regulation[60].

Taken together, the above results suggest that Model 1 and 2 coexist in the brain. This can be

formalized as follows:

$$P_i \leftarrow P_i + \hat{\alpha}_i^+ \cdot \delta_i \ldots \text{if } \delta_i \leq 0$$

$$N_i \leftarrow N_i + \hat{\alpha}_i^- \cdot \delta_i \ldots \text{if } \delta_i < 0$$

where:

$$\hat{\alpha}_i^+ = \alpha_P \cdot \alpha_i^+$$

$$\hat{\alpha}_i^- = \alpha_N \cdot \alpha_i^-$$

where α_i^+ and α_i^- correspond to the asymmetric scaling of dopamine RPEs at the single-cell level

(Model 2) and α_P and α_N correspond to the asymmetric scaling of synaptic plasticity in D1R and

D2R at the population level (Model 1).

Linking asymmetric learning and baseline dopamine levels in healthy subjects

Cools et al., (2009)[24].

There have been very few studies that examined the relationship between baseline dopamine

levels and asymmetry in learning from positive and negative outcomes. As a rare case for such

examinations, Cools et al.[24] provided intriguing data in humans. They compared the performance

in reversal learning and the quantity called 'dopamine synthesis capacity'. Dopamine synthesis

capacity is estimated by injecting the positron emission tomography (PET) tracer

[18F]fluorometatyrosine (FMT) and is thought to be correlated with baseline dopamine levels[61,62].

This study found that higher dopamine synthesis capacity was correlated with better learning

from gains but not with learning from losses (Fig. 7b). As a result, in reversal learning, subjects

with higher dopamine synthesis capacity learned from gains than losses, reported as the 'relative

reversal learning (RRL)' index in their study (Fig. 7b). This result, thus, provides direct evidence

supporting our Model 1.

In addition, they found that dopamine synthesis capacity predicts the effectiveness of

bromocriptine (D2 partial agonist) in altering learning rate asymmetry: bromocriptine's ability to

bias learning from gains over losses (i.e., positive change in RRL) was negatively correlated with

dopamine synthesis capacity (Fig. 7c). We found that this result can also be explained by Model

1. For this, we simulated the effects of bromocriptine with the biophysical model used above,

and derived the asymmetric learning rates from the slopes of the D2R occupancy (Fig. 7d,

Extended Data Fig. 6a,b) or activation curves (Fig. 7d, Extended Data Fig. 6c,d). The RRL

parameter reported by Cools et al. corresponds to the asymmetric scaling factor τ, and is

equivalent to $(2\tau - 1)$ (as described in the Methods 4.1). We then computed what would be the

change in this parameter $\Delta(2\tau - 1)$ induced by bromocriptine (Fig. 7e-f, Extended Data Fig. 6e-

l).

This analysis revealed that by considering the asymmetries in learning rates induced by changes

in the baseline occupancy of the receptors, our model can capture their results in a qualitative

manner. Intuitively, the less dopamine there is at baseline, the lower the occupancy of D2R at

placebo conditions. This leads to a larger increase in D2R occupancy induced by D2 agonist in

low dopamine baseline conditions (Fig. 7d, Extended Data Fig. 6a) and, thus, a larger increase in

asymmetry in learning form gains over losses, if D1R occupancy is kept fixed These effects still

hold even if we consider, in addition to bromocriptine's effects in postsynaptic receptors (D2

long or D2l), its effect on inhibition of dopamine release via presynaptic (D2 short or D2s)

autoreceptors[63,64] (Fig. 7d, Extended Data Fig. 6b). This can be simulated as a decrease in

dopamine level, which leads to a shift in the occupancy curves to the right. Finally, we can

consider effect of the *partial* agonism of the drug, that leads to a lower activation level of

receptors even if the occupancy is maximal (Fig. 7d, Extended Data Fig. 6c-d). Even after

considering this last factor, the results remain qualitatively the same as those found in the

original study. These results were robust to a relatively wide range of values in the simulation's

parameters (Extended Data Fig. 7, 8).

Linking psychiatric conditions to baseline dopamine levels

Timmer et al., 2018

Various psychiatric disorders are characterized by abnormal future predictions or mood. Our

Model 1 raise the possibility that an overall decrease in baseline dopamine level in the striatum

would enhance learning from negative outcomes over learning from positive outcomes leading to

persistent pessimistic future value expectations, a hallmark of depressive-like symptoms (Fig.

3a,b). A piece of evidence supporting this in the human literature is the greater learning rates for

losses over gains in patients with Parkinson's disease (PD)[22,25], its comorbidity with

depression[25,65] that can precede the PD diagnosis[65–67], and the reports of decreased dopamine

transporter binding in the ventral striatum in depressed PD patients compared to non-depressed

PD patients[68,69].

In addition, the progression of dopaminergic axonal loss in PD is topographically unbalanced:

the axonal loss is more prominent in the dorsal striatal regions[70] than in the ventral ones. This

leads to uneven dopamine baseline levels across the striatum that would interact with the global

increases in dopamine induced by dopaminergic medications in PD patients. We hypothesize that

a behavioral readout of the degree of this unevenness might be the presence or absence of

depression as a comorbidity: *patients with depression might have lower dopamine levels in the*

ventral striatum. Thus, if indeed baseline dopamine levels are correlated with depression, this

comorbidity could be predictive of the effects of PD medication.

We examined a previous study that provided evidence for this hypothesis [71]. Here, PD patients

with and without depression history were tested in a gambling task, under presence or absence of

medication ('ON' and 'OFF' medication states). The authors fitted a 'loss aversion' parameter to

the behavioral performance, which is equivalent to $1 - \tau$ in our model, under some assumptions

(Methods). Their results were consistent with our model predictions. In the OFF-medication

state, there was a (near-significant) main effect of depression group (with or without depression)

on the learning rate asymmetry: patients with a depression history tended to be more loss averse

than nondepressed patients ($P = 0.052$). This is consistent with a decrease of dopamine levels in

the ventral striatum and thus a regime of $\alpha^+ < \alpha^-$ in value learning. Importantly, in the ON-

medication state, the medication effects on the asymmetry in learning rates were predicted by the

degree of severity of depression: patients with larger depression scores exhibited greater drug-

induced decreases in loss aversion (Fig. 7g), which would correspond to an increase in $\tau =$

$\frac{\alpha^+}{\alpha^+ + \alpha^-}$ in our model This is consistent with our Model 1: higher degrees of depression might be

correlated with lower levels of baseline dopamine, making the D1R sensitivity more susceptible

to an artificial increase in baseline dopamine with L-DOPA medication (Fig. 7h; further details

discussed in Methods).

Discussion

A hallmark of various psychiatric disorders is overly optimistic or pessimistic predictions about

the future. Using RL models, we sought to identify potential biological mechanisms that give rise

to biased value predictions, with a particular focus on the roles of phasic versus tonic dopamine.

Our results demonstrate that variations in tonic dopamine levels can modulate the efficacy of

synaptic plasticity induced by positive versus negative RPEs, thereby resulting in biased value

learning (Model 1). This effect arises due to sigmoidal shapes of the dose-occupancy curves and

different affinities of dopamine receptors (D1R and D2R); alterations in the tonic dopamine level

result in changes in the slope of the dose-occupancy curve (and thus, sensitivity) of dopamine

receptors at the baseline dopamine concentration. We show that this mechanism offers a simple

explanation for how changes in tonic dopamine levels can result in biased value learning in a few

examples of value learning in mice and humans. Additionally, we show that this mechanism may

underlie symptoms of various psychiatric and neurological disorders. Although altered phasic

dopamine responses could have been a natural suspect as a candidate mechanism for biased

value learning[37,38], our study provides a novel mechanism; the interaction between tonic and

phasic dopamine can give rise to biased value learning, even when phasic dopamine responses

remain relatively unchanged.

The impact of properties of dopamine receptors on reinforcement learning (RL)

Our results highlight the importance of considering properties of dopamine receptors and neural

circuit architecture (i.e., direct and indirect pathways) in RL models. Based on different affinities

of dopamine D1 and D2 receptors, it has been proposed that D1- and D2-SPNs play predominant

roles in learning from positive and negative dopamine responses[32,72–75]. In support of this idea,

recent experiments have demonstrated that PKA signaling in D1- and D2-SPNs is primarily

driven by a phasic increase and decrease of dopamine, respectively[35]. Furthermore, LTP-like

changes in D1- and D2-SPNs are triggered by a phasic increase and decrease of dopamine,

respectively[33,34]. These recent pieces of evidence suggest that these plasticity rules are a basic

principle of the RL circuitry in the brain. Here we explored the properties of this RL model and

found the impact of the shape (slope) of receptor occupancy curves and showed that the tonic

dopamine levels can modulate the relative efficacy of learning from positive versus negative

RPEs.

One assumption in our model is that after a change in the tonic dopamine level, intracellular

signaling reaches a steady inactive state, and it is the *change* in receptor occupancy that matters

for inducing synaptic plasticity, rather than the *absolute* level of receptor occupancy reached

during phasic dopamine responses. We note that absolute level might also contribute, yet it is

expected that an increase or decrease in absolute occupancy levels will cause effects in the same

direction as the effects of relative change that we explored in this study.

Additionally, our model, which incorporates the new plasticity rules, the opponent circuit

architecture and properties of D1/D2 dopamine receptors, provides insights into the basic design

principle of the brain's RL circuit. It should be noted that the dose occupancy curves were

plotted as a function of the logarithm of dopamine concentration, which makes the occupancy

curves into sigmoidal shapes (Fig. 3, Extended Data Fig. 9). This logarithmic scaling is

important in two ways. First, considering two sigmoidal curves for D1R and D2R together, the

curves are approximately *symmetric* around the normal baseline dopamine level (Fig. 3a,

Normal). Second, logarithmic scaling means that a fold-change in dopamine concentration will

lead to the same leftward or rightward shift in these plots. It has long been argued that signaling

of RPEs by dopamine neurons is curtailed by the fact that dopamine neurons have relatively low

firing rates (2-8 spikes per second), and inhibitory responses of dopamine neurons tend to be

smaller than excitatory responses[76,77]. Importantly, if we consider logarithmic scaling of

dopamine concentration, the problem of this asymmetry is substantially mitigated (Extended

Data Fig. 10). For example, with the baseline firing of 6 spikes per second, a phasic increase to

18 spikes per second and a phasic decrease to 2 spikes per second will cause the identical *fold-

changes* in spiking (i.e., 3-fold changes in both directions), which would lead to a similar *fold-

changes* in dopamine levels (Extended Data Fig. 11) and similar *percent* increase and decrease in

receptor occupancy in D1R and D2R, respectively (Fig. 3a). Consequently, the system achieves

symmetry in its response to positive and negative dopamine responses of observed magnitudes.

This may help understand *why* the basal ganglia circuit employs the opponent circuit architecture

in the first place. In the model used in the present study, the value is encoded as the difference

between the activity of D1- and D2-SPNs $(V = P - N)$[32]. We propose that this opponent circuit

architecture, together with the logarithmic scaling of dopamine concentration, allows the system

to effectively learn and encode both positive and negative values, which are contributed by the

increase of firing in D1- and D2-SPNs, respectively. This would allow to expand the dynamic

range of value coding, without requiring high baseline firing rates. Thus, at the normal dopamine

baseline, learning from positive and negative dopamine responses is well balanced. When the

tonic dopamine level deviates from the normal level, however, then the symmetry is broken and

value learning becomes biased, as explored in the present study.

The role of tonic dopamine levels in psychiatric disorders

As mentioned above, our modeling results provide an account for biased value predictions

observed in various psychiatric and neurological conditions. For one, our model provides a link

between findings in depressive-like states in animal models and the value learning biases

exhibited by humans.

In a rodent model of depression, it has been reported that spontaneous activity of dopamine

neurons is decreased[78] (but see[79,80]). In addition, decreased spontaneous firing of dopamine

neurons has been observed as a result of chronic pain-induced adaptations that correlate with

anhedonia-like behavior[81]. Furthermore, maternal deprivation, which increases susceptibility to

anhedonia, led to an upregulation of D2R expression in the VTA[82], which is expected to decrease

the excitability of dopamine neurons via its autoreceptor function. Finally, chronic

administration of corticosteroids, a method to mimic anxiety and anhedonia-like states, results in

an increase in somatodendritic dopamine concentration which then decreases dopamine

excitability via D2R hyper-activation[83]. These results of decreased dopamine excitability

correlated with anhedonia-like states are consistent with findings of increased burst firing of

lateral habenula (LHb) neurons[56] and potentiation of glutamatergic inputs onto the habenula[57] in

depression models. This is further supported by reports that depressive-like behavioral

phenotypes can be ameliorated by optogenetic activation of dopamine neurons[84] and the anti-

depressant effects of ketamine might be mediated by the inhibition of bursting in the LHb[58]

The mechanism by which a broad change in dopamine excitability could lead to depressive-like

states remains to be revealed. Just by assuming that a decrease in spontaneous firing leads to a

decrease in baseline dopamine level in the striatum, our model readily predicts that learning from

negative outcomes will be emphasized over learning from positive outcomes (Fig. 3a,b), as has

been reported in some studies of patients with major depressive disorder (MDD)[1]. In addition,

RL agents learning in these conditions exhibit enhanced risk-aversive behavior, pessimistic

outcome expectations, and increased sensitivity to losses compared to gains, all of which are

signatures of depressive-like conditions[1,5,21,85,86]. This contrasts with findings of increased

dopamine synthesis capacity in pathological gambling patients[87], who show the opposite

behavioral signatures[3].

An additional line of research relevant to our proposal is PD patients and pathological gambling

as a comorbidity. Previous work has emphasized the interaction between the degree of

dopaminergic loss and the effects of PD medications[88–90], which can sometimes result in the

development of addictive disorders such as pathological gambling. As mentioned, the loss of

dopaminergic axons in PD patients has been reported to happen predominantly in the dorsal

regions of the striatum[70]. Thus, at the onset of the motor impairment symptoms, which is when

L-DOPA medication tends to be prescribed, dopamine level is expected to be low in the dorsal

striatum while it might be relatively intact in the ventral striatum. This can lead to 'overdose' of

dopamine by medication: while L-DOPA might take dopamine levels in the dorsal striatum back

to its original set-point, it might cause an 'overdose' in the ventral striatum[89,91]. Our model

predicts that this overdose would lead to decreases in D2R sensitivity relative to D1R. Assuming

that the ventral striatal regions have a dominant role in value learning, this would result in

excessive optimistic expectations and risk seeking, two key behavioral features of pathological

gambling and addictive disorders. We provided indirect evidence for this hypothesis; future work

should directly test these predictions.

It should be noted that we did not consider changes in dopamine receptors density, which have

also been related to value learning biases[92] and psychiatric conditions[93]. Future studies should

explore the influence of this additional factor in the encoding of asymmetric learning rates (i.e.,

$(\hat{\alpha}_i^+, \hat{\alpha}_i^-)$.

Tonic dopamine as a modulator of 'mood'

Mood refers to a person's emotional state as it relates to their overall sense of well-being.

Although the exact neural substrate of mood remains unknown, recent studies have indicated that

mood reflects not the absolute goodness of outcomes but rather on the discrepancy between

actual and expected outcomes in recent history[13,14]. That is, mood depends on the cumulative

sum of RPEs that occurred recently[13]. It has also been proposed that mood, in turn, affects the

way we perceive and learn from positive and negative outcomes (RPEs)[13].

Our model provides a unified mechanism for these two aspects of mood; both subjective feeling

of mood and biased learning from positive versus negative outcomes can arise from changes in

baseline dopamine levels which can be modulated by recent history of phasic dopamine

responses. It was proposed that this history dependent modulation of learning is an adaptive

mechanism that allows organisms to adapt quickly to slow changes in environments based on the

"momentum" of whether the situation is changing in a better or worse direction on a slow

timescale (e.g. seasonal change)[13,14]. The models presented in the present study may provide

mechanistic insights into such mood-dependent modulation of learning and perception.

Neural circuits for distributional reinforcement learning (RL)

We examined the possibility that optimistic biases in reward seeking behavior and dopamine cue

responses observed in habenula-lesioned mice can be explained by Model 2, either based on risk-

sensitive RL (the average response) or distributional RL (responses of a diverse set of individual

dopamine neurons). We did not find evidence supporting this possibility. However, the present

study makes two important contributions with respect to distributional RL. First, we can show

that our model, which incorporated direct and indirect pathway architecture, can support

distributional RL (Extended Data Fig. 12, 13). It would be interesting to examine what additional

features and functions could be gained by having this opponent architecture. Second, we largely

replicated the previous results[36] using an independent data set. That is, the signatures of

distributional RL were present in this data set (Extended Data Fig. 3-4), and dopamine cue-

evoked responses did show an optimistic bias. This provides further evidence for a distributional

code in dopamine neurons, and shows that there is an overall elevated distributional

representation in dopamine cue responses in habenula lesioned animals.

Concluding remarks

Taken together, our biologically inspired RL model provides a foundation to link findings in the

brain and formal models of RL. Our work highlights a causal impact of baseline dopamine on

biasing future value predictions, which may underlie mood and some abnormalities observed in

psychiatric patients and could be used to regulate risk sensitive behavior.

Methods

1. Reinforcement learning model

Here we provide formal definitions and the framework of reinforcement learning used in this study. We have focused our model formulations to the problem of *prediction*, in which an agent learns to predict the value function[11]. The problem of *control* (the problem of how an agent selects and executes actions) is not considered. In RL, an agent's objective is to maximize the total cumulative rewards. It does so by learning the value associated with each state in an environment. For now, we will develop the model dropping the dependency on time within each episode. Here, the target to learn is the value function as defined by

$$V(s_i) := \mathbf{E}[r^{(n)}|s^{(n)} = s_i]$$

Where $r^{(n)}$ is the reward experienced in the episode n (i.e., trial) of visiting state s_i. Learning of $V(s)$ is driven by reward prediction errors (RPEs, δ), the discrepancy between the actual and expected reward:

$$\delta^{(n)} = r^{(n)} - V(s_i)$$

The value is updated for the experienced state according to:

$$V^{(n+1)}(s_i) \leftarrow V^{(n)}(s_i) + \alpha \cdot \delta^{(n)}$$

This is also known as the Rescorla-Wagner (RW) delta rule[94]. The reward in each trial is sampled from a reward distribution specific to a given state: $r^{(n)} \sim R(s_i)$. With the learning rule above, the value converges on the expected value of this reward distribution. This can be shown with a stochastic fixed-point approach; the convergence point is derived by obtaining the value of $V(s_i)$ at which the change in $V(s_i)$ from trial n to trial $(n+1)$ is expected to be zero (i.e., is zero on average):

$$\mathbf{E}[V^{(n+1)}(s_i) - V^{(n)}(s_i)] = 0$$

$$\mathbf{E}[\alpha \cdot \delta^{(n)}] = 0$$

$$\mathbf{E}[\alpha \cdot (r^{(n)} - V(s_i)] = 0$$

$$\alpha \cdot \mathbf{E}[r] - \alpha \cdot \mathbf{E}[V(s_i)] = 0$$

$$\mathbf{E}[V(s_i)] = \mathbf{E}[r_t]$$

$$\boldsymbol{V^*}(s_i) = \mathbf{E}[r_t]$$

750   Where $V^*(s_i)$ is the stochastic fixed-point: the value around which $V(s_i)$ is expected to fluctuate after

751   learning and corresponds to the learning target above.

1.1. Temporal difference learning

Now we will consider time and extend the models to the temporal difference (TD) learning framework[11].

Dopamine responses have been shown to present key signatures of TD errors[95]. Therefore, TD learning

models allow us to directly link the model variables to dopamine neural responses.

We can derive TD learning by defining a different environmental structure and learning objective. We

start by considering arbitrary states (s_t), which transition at each time step following a Markov process,

and at each time step the agent samples a random reward from a probability distribution $r_t \sim R(s_t)$.

The learning objective is now the value of a given state $V(s_t)$ defined as the *expected cumulative sum of*

all future rewards starting from state s. Rewards are discounted by a constant discounting factor (γ, with

$0 \leq \gamma \leq 1$) each time step. The expectation is taken over stochastic state transitions and sampled rewards:

762
$$V(s_t) := \mathbf{E}[r_t + \gamma r_{t+1} + \gamma^2 r_{t+2} + \gamma^3 r_{t+3} \dots | s_t = S\}]$$

763   Where s_t is the state at time t, r_t is the reward sampled at time t and $V(s_t)$ is the value of the state s_t.

764   Since the environment and transitions are assumed to follow a Markov process, the equation above can be

765   rewritten in a recursive manner. This is known as the Bellman equation[11] :

766
$$V(s_t) := \mathbf{E}[r_t + \gamma \cdot V(s_{t+1}) | s_t = S]$$

767   The agent approximates the true value $V(s_t)$ with a learned estimate $\hat{V}(s_t)$. With this approximation,

768   before learning converges, the estimates for the left- and right-hand sides are not equal. Thus, after

769   sampling a reward $r_t \sim R(s_t)$ from the environment, the difference between the two terms in the Bellman

770   equation represents the error in value prediction, called the temporal difference reward prediction error

771   (TD RPE, δ below),

772
$$\delta_t = r_t + \gamma \cdot \hat{V}(s_{t+1}) - \hat{V}(s_t)$$

773   With α as the learning rate, the updates for the value estimates are:

774
$$\hat{V}(s_t) \leftarrow \hat{V}(s_t) + \alpha \cdot \delta_t$$

775   With this definition, the TD RPE contains the difference between the estimated value of states evaluated

776   at consecutive time points. If we fix the discounting factor to be $\gamma = 1$, then $\gamma \cdot \hat{V}(s_{t+1}) - \hat{V}(s_t)$ is the

777   temporal derivative of the value function. As a result of this property, unexpected increases and decreases

778   in value result in positive and negative transient changes in TD RPE, respectively[95].

779   If dopamine responses encode TD RPEs, then cue-evoked responses can be formulated as:

780
$$\delta_{cue} = \gamma \cdot \hat{V}(s_{cue}) - \hat{V}(s_b)$$

Where δ_{cue} is the TD RPE induced by the cue, $\hat{V}(s_b)$ is the value prediction at baseline and $\hat{V}(s_{cue})$ is

the value prediction elicited by the cue (which reflects the expected value predicted by each trial type). As

the $\hat{V}(s_b)$ is the same across all trial types and represents the average value predictions across them, then

δ_{cue} is dominated by the expected value of each trial type. This is a useful feature that we used in our

simulations for the habenula lesion experiment.

1.2. Distributional TD learning

In Results, we used a distributional TD learning model to test whether the subtle changes in the

distribution of asymmetric scaling factors observed after lesions could lead to the observed changes in cue

responses after learning.

In distributional TD learning, our learning objective is the entire distribution over cumulative discounted

future rewards, instead of the value defined above[36,37,39]. We will call this the *return distribution*, $Z(s_t)$.

We can thus write an analogue of the Bellman equation, the 'distributional Bellman equation':

793
$$Z(s_t) := R(s_t) + \gamma \cdot Z(s_{t+1})$$

794   The target to learn in distributional TD is now $V_i(s_t)$ that minimizes for the expectile regression loss:

795
$$V_i(s_t) := \underset{v}{\arg\min} \, \mathrm{E}\big[(Z(s_t) - v)^2 \cdot \big(\tau_i - \mathbf{1}_{(Z(s_t)-v)<0}\big)\big]$$

Where $Z(s_t)$ is a random variable, representing the return distribution, and $\mathbf{1}_f$ is the indicator functions

that is equal to 1 if the condition in the subscript $\{f := (Z(s_t) - v) < 0\}$ is met, and 0 otherwise.

Minimizing the expectile regression loss makes $V_i(s_t)$ to converge on the τ_i^{th} expectile of the return

distribution[39].

The target is learned by taking samples from the estimated return distribution[39] $\tilde{z}(s_{t+1}) \sim Z(s_{t+1})$ and

from the reward distribution $r_t \sim R(s_t)$, to compute the TD error:

802
$$\delta_{i,t} := r_t + \gamma \cdot \tilde{z}(s_{t+1}) - V_i(s_t)$$

803   Note that $\tilde{z}(s_{t+1})$ is random so the TD error is also random, and $\delta_{i,t} \neq r_t + \gamma \cdot V_i(s_{t+1}) - V_i(s_t)$. For

804   more information regarding the sampling method employed in the simulations see Methods Section 3.3.

805   In addition, the updates are performed with different learning rates (α_i^+, α_i^-) for positive and negative δ_i.

806   This asymmetry in the weighting of the errors used to update $V_i(s_t)$ is essential to minimize the expectile

807   regression loss.

808
$$\hat{\delta}_{i,t} = \alpha_i^+ \cdot \delta_{i,t} \;...\; \text{if } \delta_{i,t} > 0$$

809
$$\hat{\delta}_{i,t} = \alpha_i^- \cdot \delta_{i,t} \;...\; \text{if } \delta_{i,t} < 0$$

810   The reliance on a single sample for $\tilde{z}(s_{t+1})$ suffers from high variance. Therefore, for performing the

811   updates we average across a set of M updates, each depending on a single sample $\delta_{i,t}$.

812
$$\mathbf{E}[\Delta V_i(s_t)] = \frac{1}{M} \sum_j^M \alpha_i^- \cdot \delta_{i,j} \cdot \mathbf{1}_{\delta_{i,j}<0} + \alpha_i^+ \cdot \delta_{i,j} \cdot \mathbf{1}_{\delta_{i,j}>0}$$

813
$$V_i(s_t) \leftarrow V_i(s_t) + \mathbf{E}[\Delta V_i(s_t)]$$

This learning rule will asymptotically converge to the τ_i-th expectile of the return distribution[39].

1.3. TD learning with D1 and D2 populations

It is straightforward to extend the TD learning algorithm to have separate populations for D1 and D2

SPNs[32]. We employed this model to derive dopamine cue responses with Model 1 (Fig. 6i). In this model,

the same computation of TD RPE of standard TD learning is still used. Yet, this model differs in the

updates and computation of $\hat{V}(s_t)$.

As mentioned previously, the updates in the P_i and N_i populations happen exclusively with positive or

negative TD RPEs, respectively:

822
$$P(s_t) \leftarrow P(s_t) + \alpha^+ \cdot |\delta_t| - \beta \cdot P(s_t) \;...\; \text{if } \delta_t > 0$$

823
$$N(s_t) \leftarrow N(s_t) + \alpha^- \cdot |\delta_t| - \beta \cdot N(s_t) \;...\; \text{if } \delta_t < 0$$

824   Where α^+ and α^- are the learning rates for the P and N populations, that we postulate is modulated by

825   baseline dopamine levels. The variable $\beta \in (0,1)$ is the decay factor, which we keep constant throughout

826   the simulations and serves to stabilize $P(s_t)$, $N(s_t)$.

827   The computation of value estimate $\hat{V}(s_t)$ is given by:

828
$$\hat{V}(s_t) = P(s_t) - N(s_t)$$

1.3.1. Convergence of risk sensitive TD learning

We now discuss the convergence of the proposed TD learning algorithm with D1 and D2 populations.

This analysis builds on the work in risk-sensitive reinforcement learning [40] and the already established

results of convergence for stochastic iterative algorithms (e.g., TD learning) (Bertsekas & Tsitsiklis,

1996[96], Proposition 4.4, p. 156) .

Theorem: The results by Bertsekas & Tsitsiklis (1996)[96] establish that, given a sequence $r_t \in$

$\mathbb{R}^m$ generated by the iterative algorithm:

$$a_{n+1}(s) = \big(1 - \sigma_n(s)\big)a_n(s) + \sigma_n(s)\big((Ha_n)(s) + \omega_n(s)\big) \qquad \forall\, s \in 1, \dots, m \ \ \textbf{Eq. I}$$

The variable a_n converges to the unique solution a^* of the equation: $Ha^* = a^*$ with probability = 1,

assuming the following conditions are fulfilled:

1. The step sizes $\sigma_i(i)$ are non-negative and satisfy:

840
$$\sum_{n=0}^{\infty} \sigma_n(s) = \infty \ \forall\, s \in 1, \dots, m$$

841
$$\sum_{n=0}^{\infty} \sigma_n(s)^2 < \infty \ \forall\, s \in 1, \dots, m$$

2. The noise term $\omega_n(s)$ satisfies:

- $E[\omega_n(s)|\mathcal{F}_n] = 0 \ \forall\, s, n$, where $\mathcal{F}_n$ denotes the history of the process up to and including time

step n

- Given any norm $\|\cdot\|$ on $\mathbb{R}^m$ there exist constants A and B such that: $E[\omega_n^2(s)|\mathcal{F}_n] \le A +$

$B\|r_n\|^2 \ \ \forall\, s, n$

3. The mapping H is a *maximum norm contraction* (see below for definition)

To prove convergence, we will first discuss the case of risk-sensitive TD learning following [40] and then

discuss TD learning with D1 and D2 populations.

We define the risk sensitive TD-learning rule as:

851
$$\hat{V}_n(s) \leftarrow \hat{V}_{n-1}(s) + \sigma \cdot \mathcal{X}^\tau\big(\delta_{s_{n-1}, s_n}\big)$$

Where:

- $\delta_{s_{n-1},s_n} = r_{n-1,n} + \gamma \cdot \hat{V}_{n-1}(s_n) - \hat{V}_n(s_{n-1})$

- The step index is $n \in 0, \ldots, \infty$

- The step size σ is kept constant across iterations.

- For simplicity in calculations we follow [40] and make use of the operator $\mathcal{X}^K$ with $K \in (-1,1)$

$$\mathcal{X}^K(x) = (1 - K) \cdot x \qquad \ldots \text{if } x > 0$$
$$\mathcal{X}^K(x) = (1 + K) \cdot x \qquad \ldots \text{if } x \leq 0$$

It is simple to show that the asymmetric scaling factor used in this paper is a scaled version of

the operator. That is: $\tau = 0.5(1 - K)$ and $1 - \tau = 0.5(1 + K)$.

- In addition, as in [40], given that the function $\mathcal{X}^\tau(x)$ is piece-wise differentiable we can apply

the mean value theorem to show that for each pair of numbers (a, b) there exists a $\mathcal{E}_{a,b,K} \in$

$[1 - |K|, 1 + |K|]$, such that: $\mathcal{E}_{a,b,K} = \frac{\mathcal{X}^\tau(a) - \mathcal{X}^\tau(b)}{a-b}$. This relationship will become useful in

the future.

We will re-format the update rule to better match the iterative algorithm above:

Adding and subtracting $\sigma \cdot \hat{V}_{n-1}(s)/\alpha$

$$\hat{V}_n(s) \leftarrow (1 - \sigma/\alpha)\hat{V}_{n-1}(s) + \sigma/\alpha \left(\alpha \cdot \mathcal{X}^\tau(\delta_{n-1}) + \hat{V}_{n-1}(s) \right)$$

Defining an operator that will become useful:

$$\mathcal{T}_{\alpha K}[V](s) := V(s) + \alpha \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{X}^K \cdot \delta_{ij}$$

Defining the noise term as:

$$\omega_{n-1}(s) = \hat{V}_{n-1}(s) + \alpha \cdot \mathcal{X}^K\left(\delta_{s_{n-1},s_n}\right) - \mathcal{T}_{\alpha K}[\hat{V}_{n-1}](s)$$

Then our update rule above becomes:

$$\hat{V}_n(s) \leftarrow (1 - \sigma/\alpha)\hat{V}_{n-1}(s) + \sigma/\alpha \left(\mathcal{T}_{\alpha K}[\hat{V}_{n-1}](s) + \omega_{n-1}(s) \right)$$

The formulation above can be directly compared to the one of stochastic iterative algorithm theorem

(Eq.I) , and now we can check whether the conditions for convergence are met.

1. The conditions for the learning rate, are a direct consequence of our choice of the parameter

which is a constant in our model and $0 < \alpha < 1$.

2. It has been shown that showed that the conditions for the noise term $\omega_{n-1}(s)$ as formulated

above are satisfied[40].

3. Finally, the operator $\mathcal{T}_{\alpha\tau}[V](s)$ is a contraction mapping as also shown in Bersekas (1996)[40].

Therefore, the variable V_n converges to the unique solution V^* for which:

882
$$V^* = \mathcal{T}_{\alpha K}[V^*] = V^* + \alpha \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{X}^K \cdot \delta_{ij}$$

883 We elaborate now on the proof for the contraction mapping of the operator $\mathcal{T}_{\alpha\tau}[V](s)$, as this will be

884 useful for the proof of the D1 D2 TD learning model.

885 **Definition of contraction mapping.** Let (X, d) be a metric space (a set X, with a notion of distance, d,

886 between points). A mapping $\mathcal{T}: X \to X$ is a contraction mapping if there exists a constant $c: 0 \geq c > 1$

887 such that for all $x \in X$:

888
$$d\big(\mathcal{T}[x_i], \mathcal{T}[x_j]\big) \leq c d\big(x_i, x_j\big)$$

889 That is, a contraction mapping maps points closer together.

890 Elaborating now on the operator $\mathcal{T}_{\alpha\tau}[V](s)$ and using $|\cdot|$ as our distance metric:

891 $|\mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i)|$

892
$$= \left| V_1(i) + \alpha \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{X}^K \cdot \big(r_{i,j} + \gamma\, V_1(j) - V_1(i)\big) - V_2(i) + \alpha \right.$$

893
$$\left. \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{X}^K \cdot \big(r_{i,j} + \gamma\, V_2(j) - V_2(i)\big) \right|$$

894 $\quad | \mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i) |$

895
$$= \left| V_1(i) - V_2(i) \right.$$

896
$$+ \alpha \sum_{i,j \in S} p_{ij} \cdot \left(\mathcal{X}^\tau \cdot \left(r_{i,j} + \gamma V_1(j) - V_1(i) \right) - \mathcal{X}^\tau \cdot \left(r_{i,j} + \gamma V_2(j) - V_2(i) \right) \right) \left. \right|$$

897 $\quad$ Using the relation defined above $\mathcal{E}_{a,b,K} \cdot (a - b) = \mathcal{X}^K(a) - \mathcal{X}^K(b)$

898 $\quad | \mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i) |$

899
$$= \left| V_1(i) - V_2(i) + \alpha \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} \cdot \left(\gamma \left(V_1(j) - V_2(i) \right) - \left(V_1(j) - V_2(i) \right) \right) \right|$$

900 $\quad | \mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i) |$

901
$$= \left| \left(1 - \alpha \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} \right) \cdot \left(V_1(j) - V_2(i) \right) \right.$$

902
$$+ \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} \cdot \left(\gamma \left(V_1(j) - V_2(i) \right) \right) \left. \right|$$

904 $\quad$ Given that $\mathcal{E}_{a,b,K} \in [1 - |K|, 1 + |K|]$ and assuming $\alpha \in (0, (1 + |K|)^{-1})$:

905
$$1 - \alpha \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} > 0$$

906 $\quad$ Taking this term outside the $| \cdot |$ and rearranging:

907
$$| \mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i) | = \left(1 - \alpha \cdot (1 - \gamma) \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} \right) | \left(V_1(j) - V_2(i) \right) |$$

908
$$| \mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i) | = c \cdot \left| \left(V_1(j) - V_2(i) \right) \right|$$

909   Where the term:

$$c = \left(1 - \alpha \cdot (1 - \gamma) \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1, V_2, K}\right)$$

911   To get the upper boundary of c we use the minimum value for the sum, where $\mathcal{E}_{V_1 V_2 K} = 1 - |K| \;\; \forall i, j \in$

912   S. And use the assumption that $\alpha \in (0, (1 + |K|)^{-1})$:

913
$$c \leq (1 - \alpha \cdot (1 - \gamma) \cdot (1 - |K|)$$

914
$$\lim_{\alpha \to 0} c = (1 - \alpha \cdot (1 - \gamma) \cdot (1 - |K|) = 1$$

915   To get the lower boundary of c we use the maximum value for the sum, where $\mathcal{E}_{V_1 V_2 K} = 1 + |K| \;\; \forall i, j \in$

916   S. And use the assumption that $\alpha \in (0, (1 + |K|)^{-1})$:

917
$$c \geq (1 - \alpha \cdot (1 - \gamma) \cdot (1 + |K|)$$

918
$$\lim_{\alpha \to (1 + |K|)^{-1}} c = (1 - (1 + |K|)^{-1} \cdot (1 - \gamma) \cdot (1 + |K|) = \gamma$$

919   Therefore: $\gamma < c < 1$

920
$$|\mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i)| \leq c \cdot |(V_1(j) - V_2(i))|$$

921   And the operator $\mathcal{T}_{\alpha \tau}[V](s)$ is a contraction mapping, under the condition $\alpha \in (0, (1 + |K|)^{-1})$

922   **_1.3.2.   Convergence of TD learning with D1 and D2 populations_**

923   We define the D1-D2 TD-learning rule as:

924
$$\hat{V}_n(s) \leftarrow \hat{V}_{n-1}(s) + \alpha \cdot \mathcal{X}^\tau\left(\delta_{s_{n-1}, s_n}\right) - \beta \hat{V}_{n-1}(s)$$

925   Note this update rule is analogous to the risk sensitive TD learning rule except for the last term that

926   emerges from the decay factor in the P, N populations of our model.

927   Performing the same re-arrangement as above we reach:

928
$$\hat{V}_n(s) \leftarrow (1 - \sigma/\alpha)\hat{V}_{n-1}(s) + \sigma/\alpha \left(\alpha \cdot \mathcal{X}^\tau(\delta_{n-1}) + \hat{V}_{n-1}(s) - \alpha \cdot \beta \hat{V}_{n-1}(s)\right)$$

929   We define a new operator $\mathcal{T}'_{\alpha K}[V](s)$:

930
$$\mathcal{T}'_{\alpha K}[V](s) := (1 - \alpha \cdot \beta) \cdot V(s) + \alpha \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{X}^K \cdot \delta_{ij}$$

931 And the noise then is defined as:

932
$$\omega_{n-1}(s) = \alpha \cdot \mathcal{X}^K\left(\delta_{s_{n-1},s_n}\right) + \hat{V}_{n-1}(s) - \alpha \cdot \beta \cdot \hat{V}_{n-1}(s) - \mathcal{T}'_{\alpha K}\left[\hat{V}_{n-1}\right](s)$$

933 The update becomes:

934
$$\hat{V}_n(s) \leftarrow (1 - \sigma/\alpha)\hat{V}_{n-1}(s) + \sigma/\alpha\left(\mathcal{T}'_{\alpha K}\left[\hat{V}_{n-1}\right](s) + \omega_{n-1}(s)\right)$$

935 The noise term reduces to the same expression as the one of TD learning, and so it fullfils the
936 requirements for the theorem of stochastic iterative algorithms. We will now test whether the operator
937 $\mathcal{T}'_{\alpha K}[V](s)$ also represents a contraction map.

938 $\left|\mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i)\right|$

939
$$= \Bigg| V_1(i) + \alpha \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{X}^K \cdot \left(r_{i,j} + \gamma V_1(j) - V_1(i)\right) - \alpha \cdot \beta \cdot V_1(i) + V_2(i) + \alpha$$

940
$$\cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{X}^K \cdot \left(r_{i,j} + \gamma V_2(j) - V_2(i)\right) - \alpha \cdot \beta \cdot V_2(i) \Bigg|$$

941 $\left|\mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i)\right|$

942
$$= \Bigg| V_1(i) - V_2(i) - \alpha \cdot \beta(V_1(i) - V_2(i))$$

943
$$+ \alpha \sum_{i,j \in S} p_{ij} \cdot \left(\mathcal{X}^\tau \cdot \left(r_{i,j} + \gamma V_1(j) - V_1(i)\right) - \mathcal{X}^\tau \cdot \left(r_{i,j} + \gamma V_2(j) - V_2(i)\right)\right) \Bigg|$$

944 Using the relation defined above $\mathcal{E}_{a,b,K} \cdot (a - b) = \mathcal{X}^K(a) - \mathcal{X}^K(b)$

945 $\left|\mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i)\right|$

946
$$= \Bigg| (1 - \alpha\beta) \cdot (V_1(i) - V_2(i))$$

947
$$+ \alpha \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} \cdot \left(\gamma\left(V_1(j) - V_2(i)\right) - \left(V_1(j) - V_2(i)\right)\right) \Bigg|$$

948 $\quad |\mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i)|$

$$= \left| \left(1 - \alpha \cdot \beta - \alpha \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} + \gamma\alpha \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} \right) \cdot \left(V_1(j) - V_2(i) \right) \right|$$

$$|\mathcal{T}_{\alpha K}[V_1](i) - \mathcal{T}_{\alpha K}[V_2](i)| = c \cdot \left| \left(V_1(j) - V_2(i) \right) \right|$$

951 Where the term:

$$c = \left(1 - \alpha \cdot \beta - \alpha \cdot (1 - \gamma) \cdot \sum_{i,j \in S} p_{ij} \cdot \mathcal{E}_{V_1,V_2,K} \right)$$

953 To get the upper boundary of c we use the minimum value for the sum, where $\mathcal{E}_{V_1 V_2 K} = 1 - |K| \ \forall i,j \in$

954 S, and use the assumption that $\alpha \in (0, (1 + |K|)^{-1})$:

$$c \le 1 - \alpha \cdot \beta - \alpha \cdot (1 - \gamma) \cdot (1 - |K|)$$

$$\lim_{\alpha \to 0} c = 1 - \alpha \cdot \beta - \alpha \cdot (1 - \gamma) \cdot (1 - |K|) = 1$$

957 To get the lower boundary of c we use the maximum value for the sum, where $\mathcal{E}_{V_1 V_2 K} = 1 + |K| \ \forall i,j \in$

958 S, and use the assumption that $\alpha \in (0, (1 + |K|)^{-1})$:

$$c \ge (1 - \alpha \cdot \beta - \alpha \cdot (1 - \gamma) \cdot (1 + |K|)$$

$$\lim_{\alpha \to (1+|K|)^{-1}} c = (1 - (1 + |K|)^{-1} \cdot \beta - (1 + |K|)^{-1} \cdot (1 - \gamma) \cdot (1 + |K|) = \gamma - (1 + |K|)^{-1} \cdot \beta$$

961 Given that we want $c \ge 0$ we can find the parameter ranges to achieve this:

$$c = \gamma - (1 + |K|)^{-1} \cdot \beta \ge 0$$

963 Given that: $K \in (-1,1)$, we use the minimum value of $|K| = 0$ to find the limit of c:

$$\lim_{|K| \to 0} c = \gamma - \beta$$

965 So the condition $\gamma \ge \beta$ needs to be present to keep: $0 \le c < 1$.

966 Under these conditions, the operator $\mathcal{T}'_{\alpha K}[V](s)$ also represents a contraction map.

$$|\mathcal{T}'_{\alpha K}[V_1](i) - \mathcal{T}'_{\alpha K}[V_2](i)| \le c \cdot \left| \left(V_1(j) - V_2(i) \right) \right|$$

Stochastic fixed point for the value estimate:

Having shown convergence of the algorithm, we will now derive the convergent points for our algorithm

using stochastic fixed points. For clarity, we estimate the stochastic fixed point dropping the dependency

on time.

If learning between P and N is symmetric $\alpha^+ = \alpha^- = \alpha$. We derive the convergent estimate of $V(s_t)$

with a fixed-point approach. First, we subtract the P and N update equations, to arrive to the update in

the $\hat{V}(s_t)$ between the (n) and the $(n + 1)$ update:

$$\hat{V}^{(n+1)}(s_i) \leftarrow \hat{V}^{(n)}(s_i) + \alpha \cdot \delta^{(n)} - \beta \cdot \hat{V}^{(n)}(s_i)$$

Where the superscripts indicate the iteration number. We can now derive the stochastic fixed-point

for $\hat{V}(s_t)$:

$$\mathbf{E}\left[\hat{V}^{(n+1)}(s_i) - \hat{V}^{(n)}(s_i)\right] = 0$$

$$\mathbf{E}\left[\alpha \cdot \delta^{(n)} - \beta \cdot \hat{V}^{(n)}(s_i)\right] = 0$$

$$\mathbf{E}\left[\alpha \cdot \left(r^{(n)} - \hat{V}(s_i)\right) - \beta \cdot \hat{V}^{(n)}(s_i)\right] = 0$$

$$\alpha \cdot \mathbf{E}[r] - (\alpha + \beta) \cdot \mathbf{E}\left[\hat{V}(s_i)\right] = 0$$

$$\mathbf{E}\left[\hat{V}(s_i)\right] = \frac{\alpha}{\alpha + \beta}\mathbf{E}[r]$$

$$\boldsymbol{V}^* = \frac{\alpha}{\alpha + \beta}\mathbf{E}[r]$$

Where $\boldsymbol{V}^* = \mathbf{E}\left[\widehat{\boldsymbol{V}}(\boldsymbol{s_i})\right]$ is the value around which $\hat{V}(s_i)$ is expected to fluctuate after convergence.

Throughout this study, we have manipulated the learning rates between P and N to be asymmetric $\alpha^+ \neq$

α^- or, equivalently, $\tau \neq 1 - \tau$. We can find the stochastic fixed point for this more general case:

$$\mathbf{E}\left[\hat{V}^{(n+1)}(s_i) - \hat{V}^{(n)}(s_i)\right] = 0$$

$$\mathbf{E}\left[\tau \cdot \left|\delta^{(n)}\right| \cdot \mathbf{1}_{\delta>0} - (1 - \tau) \cdot \left|\delta^{(n)}\right| \cdot \mathbf{1}_{\delta<0} - \beta \cdot \hat{V}^{(n)}(s_i)\right] = 0$$

To take the expectation we use the definition: $\mathrm{E}[X] = \sum_i p(x_i) \cdot x_i$. For a Bernoulli distribution, $p(x_i)$

takes two values:

• $p(x_i) = p$ if reward is delivered and, thus $r = 1,\ \delta_t > 0$,

• $p(x_i) = (1 - p)$ if reward is not delivered and, thus $r = 0,\ \delta_t < 0$,

Therefore, we can resolve the expectation and expand the RPEs:

$$\mathbf{E}\left[\tau \cdot \left|\delta^{(n)}\right| \cdot \mathbf{1}_{\delta>0} - (1-\tau) \cdot \left|\delta^{(n)}\right| \cdot \mathbf{1}_{\delta<0} - \beta \cdot \hat{V}^{(n)}(s_i)\right] = 0$$

$$\tau \cdot \mathbf{E}\left[\left|r - \hat{V}(s_i)\right| \cdot \mathbf{1}_{\delta>0}\right] - (1-\tau) \cdot \mathbf{E}\left[\left|-\hat{V}(s_i)\right| \cdot \mathbf{1}_{\delta<0}\right] - \beta \cdot \mathbf{E}\left[\hat{V}(s_i)\right] = 0$$

996   Taking the absolute values:

$$\left|r - \hat{V}(s_i)\right| = r - \hat{V}(s_i) \dots \text{ if } \left(r - \hat{V}(s_i)\right) > 0$$

$$\left|-\hat{V}(s_i)\right| = \hat{V}(s_i) \dots \text{ if } \left(-\hat{V}(s_i)\right) < 0$$

$$\tau \cdot \mathbf{E}\left[\left(r - \hat{V}(s_i)\right) \cdot \mathbf{1}_{\delta>0}\right] - (1-\tau) \cdot \mathbf{E}\left[\hat{V}(s_i) \cdot \mathbf{1}_{\delta<0}\right] - \beta \cdot \mathbf{E}\left[\hat{V}(s_i)\right] = 0$$

1000   Replacing stochastic fixed point: $\mathbf{E}\left[\hat{V}(s_t)\right] = V^*$ and taking the expectations:

$$\beta \cdot V^* = \tau \cdot p \cdot (r - V^*) - (1-\tau) \cdot (1-p) \cdot V^*$$

1001

1002   Rearranging and isolating V^*, we obtain:

$$V^* = \frac{\dfrac{\tau}{1-\tau} \cdot \dfrac{p}{1-p} \cdot r}{\dfrac{\tau}{1-\tau} \cdot \dfrac{p}{1-p} + 1 + \dfrac{\beta}{(1-\tau) \cdot (1-p)}}$$

1003

1004   *Stochastic fixed point for P and N populations:*

1005   We have mentioned that the decay term (β) in the update equations serves to stabilize the estimates of the
1006   P and N populations (i.e., avoid infinite increases). We can observe the influence of β by computing the
1007   stochastic fixed points for these variables.

1008   For the P population:

$$\mathbf{E}\left[P^{(n+1)}(s_t) - P^{(n)}(s_t)\right] = 0$$

1009

$$\mathbf{E}\left[\tau \cdot \left|r - V^{(n)}(s_t)\right| \cdot \mathbf{1}_{\delta>0} - \beta \cdot P^{(n)}(s_t)\right] = 0$$

1010

$$p \cdot \tau \cdot (r - V^*) - \beta \cdot P^* = 0$$

1011

$$P^* = \frac{p \cdot \tau}{\beta} \cdot (r - V^*)$$

1012

1013   Similarly, for the N population:

1014
$$\mathbf{E}\big[N^{(n+1)}(s_t) - N^{(n)}(s_t)\big] = 0$$

1015
$$\mathbf{E}\big[(1-\tau) \cdot \big|-V^{(n)}\big| \cdot \mathbf{1}_{\delta<0} - \beta \cdot N^{(n)}(s_t)\big] = 0$$

1016
$$(1-p) \cdot (1-\tau) \cdot V^* - \beta \cdot N^* = 0$$

1017
$$N^* = \frac{(1-p) \cdot (1-\tau)}{\beta} \cdot V^*$$

1018   As it can be seen in the stochastic fixed points P^*, N^*, the term $\frac{1}{\beta}$ is a proportionality constant. Therefore:

1019
$$\lim_{\beta \to 0} P^* = \lim_{\beta \to 0} \left(\frac{p \cdot \tau}{\beta} \cdot (r - V^*) \right) = \text{undefined}$$

1020
$$\lim_{\beta \to 0} N^* = \lim_{\beta \to 0} \left(\frac{(1-p) \cdot (1-\tau)}{\beta} \cdot V^* \right) = \text{undefined}$$

1021   So, $\beta \neq 0$ needs to be met for the stochastic fixed points P^*, N^* to exist. In Extended Data Fig. 1 we show

1022   empirically that the convergence rate is slower as β gets closer to 0, but it is always achieved.

1.3.3.   *Sensitivity of learned variables in D1-D2 model to parameters*

The conditions for the D1-D2 model to reproduce the data from our habenula lesion experiment and some

of the previous studies are that:

1. The bias in V^* induced by the asymmetric learning rates doesn't change the monotonicity of the

learned values as a function of the true expected value of the return distribution $\mathbf{E}[R(s)]$. In other

words, regardless of the level of 'optimism' or 'pessimism', V^* monotonically increases with

$\mathbf{E}[R(s)]$.

2. Asymmetric learning rates change the concavity of V^* as a function of $\mathbf{E}[R(s)]$: 'Optimistic' or

'pessimistic' value functions are concave or convex with respect to $\mathbf{E}[R(s)]$, respectively.

1032

1033   We will now analyze whether these conditions are met, considering the range of parameters of relevance:

1034 $0 < \tau < 1, r \neq 0$ and $0 < \beta < 1$

1035   For the condition 1 to be met, the first derivative of V^* with respect to $\mathbf{E}[R(s)]$ should always be positive.

1036   In the case of Bernoulli return distributions, the derivative of V^* with respect to p(reward) is

1037
$$\frac{\partial V^*}{\partial p} = \frac{\partial}{\partial p} \left(\frac{\frac{\tau}{1-\tau} \cdot \frac{p}{1-p} \cdot r}{\frac{\tau}{1-\tau} \cdot \frac{p}{1-p} + 1 + \frac{\beta}{(1-\tau) \cdot (1-p)}} \right) = \frac{\tau \cdot r \cdot (\tau - \beta - 1)}{(\tau \cdot (2p - 1) + \beta - \tau + 1)^2}$$

We can look at the fixed points of this expression, as they correspond to the value of p at which the

derivative changes the sign. This expression has fixed points at: $\tau = 0, r = 0$, and $\beta = \tau - 1$. Given our

parameters' ranges: $0 < \tau < 1, r \neq 0$ and $0 < \beta < 1$, none of those fixed points are present within those

ranges. In addition, it can be seen that this $(\frac{\partial V^*}{\partial p})$ is positive for the parameter values within those ranges.

Therefore, knowing that the derivative won't reach any fixed point, V^* is always a growing monotonic

function with respect to p.

For the condition 2 to be met, we can analyze the second derivative of V^* with respect to $\mathbf{E}[R(s)]$ as it

indicates the convexity of a function. The conditions to be mat are:

-   V^* is convex if it is 'pessimistic': if $\tau < 0.5 \rightarrow \frac{\partial^2 V}{\partial p^2} > 0$

-   V^* is concave if it is 'optimistic: if $\tau > 0.5 \rightarrow \frac{\partial^2 V}{\partial p^2} < 0$

In the case of Bernoulli return distributions, we take the second derivative of V^* with respect to

p(reward):

1050
$$\frac{\partial^2 V^*}{\partial p^2} = \frac{\partial^2}{\partial p^2} \left(\frac{\frac{\tau}{1-\tau} \cdot \frac{p}{1-p} \cdot r}{\frac{\tau}{1-\tau} \cdot \frac{p}{1-p} + 1 + \frac{\beta}{(1-\tau) \cdot (1-p)}} \right) = \frac{2\tau \cdot (2\tau - 1) \cdot r \cdot (\tau - \beta - 1)}{(\tau \cdot (2p - 1) + \beta - p + 1)^3}$$

1051 We can again look at the fixed points of this expression. These happen at: $\tau = 0, r = 0$, and $\beta = \tau - 1$

1052 and $\tau = 0.5$. Among them, the only fixed point within our parameters range is the latter. In addition, by

1053 replacing τ in the expression above, it is easily shown that it is positive if $\tau < 0.5$ and negative if $\tau > 0.5$.

1054 Thus, given that the ranges for the parameters are such that the second derivative won't reach any other

1055 fixed point, condition 2 will always be met.

1.4. Distributional TD learning with D1 and D2 populations

The signatures of distributional RL were preserved in dopamine neurons firing rates after habenula lesions (Extended Data Fig. 3-4). Therefore, we considered a third alternative to model 1 and 2, that assigns different functions to each of the mechanisms for asymmetric learning rates.

In this model (Extended Data Fig. 13) the single cell asymmetric scaling factors (α_i^+, α_i^-) give rise to a distributional expectile code for value and are implemented at the level of the scaling of RPE-evoked responses of dopamine neurons:

$$\delta_{i,t} := r_t + \gamma \cdot \tilde{z}(s_{t+1}) - V_i(s_t)$$

$$\hat{\delta}_{i,t} = \alpha_i^+ \cdot \delta_{i,t} \ldots \text{ if } \delta_{i,t} > 0$$

$$\hat{\delta}_{i,t} = \alpha_i^- \cdot \delta_{i,t} \ldots \text{ if } \delta_{i,t} < 0$$

The modulation of receptor sensitivities, carried out downstream at the SPN level, gives rise to the global rescaling of the value updates (η^+, η^-) (Extended Data Fig. 13A):

$$P_i(s_t) \leftarrow P_i(s_t) + \eta^+ \cdot \left|\hat{\delta}_i(t)\right| - \beta \cdot P_i(s_t) \ldots \text{ if } \delta_i(t) > 0$$

$$N_i(s_t) \leftarrow N_i(s_t) + \eta^- \cdot \left|\hat{\delta}_{i,}(t)\right| - \beta \cdot N_i(s_t) \ldots \text{ if } \delta_i(t) > 0$$

$$\widehat{V_i}(s_t) = P_i(s_t) - N_i(s_t)$$

These set of update equations are equivalent to a modified version of the update equation of distributional RL:

$$[\Delta V_i(s_t)] = \frac{1}{N} \sum_j^N \eta^+ \cdot \alpha_i^+ \cdot \delta_{i,j} \cdot I_{\delta_{i,j}>0} + \eta^- \cdot \alpha_i^- \cdot \delta_{i,j} \cdot I_{\delta_{i,j}>0}$$

$$V_i(s_t) \leftarrow V_i(s_t) + \mathbf{E}[\Delta V_i(s_t)]$$

Thus, this model can give rise to biases in value learning (Extended Data Fig. 13), while keeping intact information about the value distribution. By employing the results from the biophysical model (Fig. 6), we found that this distributional TD model can parsimoniously explain all aspects of the data in the habenula lesion study (Extended Data Fig. 12B), including the features of a distributional code and the optimistic biases observed in behavior and dopamine cue-evoked responses (Extended Data Fig. 12).

1.5. Dependency of model on assumption: Log vs. linear scaling of receptor occupancy curves

Through this work, we have used the dose-occupancy curves of D1 and D2 receptors to derive the receptor sensitivities that result in the asymmetric scaling factors in Model 1. It is important to note that the slopes of the receptor occupancy curve (= receptor sensitivity) were obtained from the receptor occupancy curves plotted as a function of log of dopamine concentrations.

$$\alpha^+ = \frac{\Delta Occ_{D1}}{\Delta \log(C_{DA+})}$$

$$\alpha^- = \frac{\Delta Occ_{D2}}{\Delta \log(C_{DA-})}$$

To show that this assumption is not essential, we now derive the receptors sensitivities assuming linear changes in dopamine levels due to RPE-evoked responses.

$$\alpha^+ = \frac{\Delta Occ_{D1}}{\Delta \, C_{DA+}}$$

$$\alpha^- = \frac{\Delta Occ_{D2}}{\Delta}$$

As shown in Extended Data Fig. 9, the choice of a linear versus log scale affects the absolute magnitude of the derived receptor sensitivities, but the normalized metric $\tau = \frac{\alpha^+}{\alpha^- + \alpha^+}$ holds the same relationship to baseline dopamine levels with a small shift in the curve (Extended Data Fig.9, right panel). The normalized metric is the factor determining the update asymmetries and, thus, the stochastic fixed points at which the variables converge.

1.6. **Normative motivation for two-factor learning rule**

We have used in the previous models a so-called *two factor learning rule*, where the value updates depend only on the presynaptic activity (i.e., state input) and TD RPEs. Here, we motivate this choice from a normative approach based on previous work[11].

1102 Consider a linear approximation for value, where the value function ($\hat{V}$) is the output of a single linear

1103 neuron. Here, $\hat{V}$ is a linear function of the input feature-vector representing the state $\mathbf{x}(s) =$

1104 $(x_1(s), ..., x_n(s))$, parametrized with a weight vector $\mathbf{w} = (w_1, ..., w_n)$. :

$$\hat{V}(s, \mathbf{w}) = \mathbf{w}^T \mathbf{x}(s)$$

1106 To put it into neural terms, we can think of $x_i(s)$ as the presynaptic activity onto the value neuron $\hat{V}$,

1107 with a synaptic efficacy w_i.

1108 As before, the agent computes the TD error based on this linear approximation for value and the sampled

1109 reward:

$$\delta_t = r_t + \gamma \cdot \hat{V}(s_{t+1}, \mathbf{w}) - \hat{V}(s_t, \mathbf{w})$$

In the problem of *value prediction*, the agent aims to achieve the highest accuracy of prediction. One way

to achieve this is to perform *stochastic gradient descent* (SGD) with respect to the parameters (w) of the

value function to minimize the objective function such as the squared error (δ_t^2). We can define this

optimization problem as: $\text{argmin}_w \left(\frac{1}{2} \delta^2 \right)$ where we have deliberately chosen the constant $\frac{1}{2}$ for clarity,

but it doesn't change the end results.

To perform SGD in this minimization problem, the parameters (w) should be updated in the opposite

direction of the gradient of the loss with respect to the parameters (i.e., opposite to $\nabla_w (\frac{1}{2} \delta^2)$) :

$$\mathbf{w} \leftarrow \mathbf{w} - \alpha \cdot \nabla_w \left(\frac{1}{2} \delta^2 \right)$$

1119 Where α is the learning rate. To compute the gradient, we use the chain rule:

1120 $$\nabla_w \left(\frac{1}{2} \delta^2 \right) = \frac{\partial (1/2 \ \delta^2)}{\partial \hat{V}} \cdot \frac{\partial \hat{V}}{\partial w} = \frac{\partial (1/2 \ (r - \hat{V})^2)}{\partial \hat{V}} \cdot \frac{\partial \mathbf{w}^T \mathbf{x}(s_t)}{\partial w} = \frac{-2(r - V)}{2} \cdot x(s_t) = -\delta \cdot x(s_t)$$

1121 Therefore, the update for the parameters of the value function is:

$$\mathbf{w} \leftarrow \mathbf{w} + \alpha \cdot \delta \cdot x(s_t)$$

The term $\delta \cdot \mathbf{x}(s_t)$ in the equation above is what we call a *two-factor learning rule*, dependent only on the

presynaptic activity and not contingent on the post-synaptic activity.

The development of the TD learning model with D1 and D2 populations (section 1.3) has respected this

learning rule, complying with what is required for SGD in the value prediction problem. Note that we

have implicitly developed our models with a complete serial compound representation (CSC) of the

states[11], where $x(s_t) = 1$ in a single element $x_i(s_t)$ representing the current state and 0 otherwise. It can

be shown that with this representation, the update equation above is equivalent to:

$$\hat{V} \leftarrow \hat{V} + \alpha \cdot \delta$$

1131

2. Computational model of dopamine release and receptor occupancy

To predict changes in dopamine concentrations and receptor occupancies (Fig. 6), we employed a

biophysical model developed elsehwhere[59]. It presents two interacting dynamical systems. The first

system models the change in receptor occupancies while the second the change in dopamine levels per

unit time.

In the first system, the occupancy of receptors is modelled as a binding reaction between dopamine (DA)

and D1 or D2 receptors (R), using the constants for forward and backward reactions (k_{on}, k_{off}).

$$DA + R \;\; {}_{k_{off}}\rightleftarrows^{k_{on}} \;\; DA{:}R$$

This formulation results in the following equation for the change in receptor occupancy $Occ(t)$ per unit

time:

$$\frac{dOcc(t)}{dt} = \left(1 - Occ(t)\right) \times k_{on} \times C_{DA}(t) - Occ(t) \times k_{off}$$

The values used for the association and dissociation constants for each receptor type (k_{on} and k_{off},

respectively) are detailed in Table 1.

In the second system, the change in dopamine concentration ($C_{DA}(t)$) is a function of both dopamine

release and uptake.

$$\frac{dC_{DA}(t)}{dt} = DA_{release}(t) - DA_{uptake}(t)$$

Dopamine release is a product of firing rate ($v(t)$) and release capacity ($\gamma(t)$)

$$DA_{release}(t) = \gamma(t) \cdot v(t)$$

Where:

1. $v(t)$ is the firing rate of dopamine neurons, provided by the neural data.

2. $\gamma(t) = \gamma_{pr_n} \cdot P_r \cdot G_{D2}(t)$ is defined as the increase in $C_{DA}(t)$ by a single synchronized action
potential:

a. $P_r = 1$ release probability in the absence of presynaptic D2-autorreceptors,

b. $\gamma_{pr_n} = 2$ release capacity in the absence of presynaptic D2-autorreceptors. This value was set
to be deliberately high and anticipates a ~50% reduction by terminal feedback.

c. $G_{D2}(t)$ is a multiplicative gain that represents the modulation of dopamine release by D2-
autorreceptors. This is a decaying function of the occupancy of D2-autorreceptors
$(Occ_{D2_a}(t))$ which is modelled by the same binding reaction explained above. The gain is
parametrized by the autoreceptor efficacy, $\alpha = 3$. The smaller the α the less the decay in
release with receptor occupancy.

1162
$$G_{D2}(t) = \frac{1}{1 + \alpha \cdot Occ_{D2_a}(t)}$$

1163   Dopamine uptake is a function of the uptake of dopamine by the dopamine transporter (DAT) and other
1164   non-DAT sources

1165
$$DA_{uptake}(t) = dt \cdot \left(\frac{V_{max}^{pr_n} \cdot C_{DA}(t)}{K_m + C_{DA}(t)} - K_{nonDAT} \right)$$

Where:

- $V_{max}^{pr_n} = 1500 \frac{nM}{sec}$s the maximal uptake capacity assuming approximately 100 terminals in the
near surroundings.

- $K_m = 160\ nM$, is the Michaelis-Menten parameter for uptake mediated by DAT

- $K_{nonDAT} = 0.04\ nM$ is a constant for the dopamine removal not mediated by DAT. For
example, monoamine oxidase (MAO) and noepinephrine transporter (NET) mediated uptake.

The variables of the model reported in Fig. 6 correspond to: $Occ_{D1}(t), Occ_{D2}(t), C_{DA}(t)$. We used as
input to the model the firing rates derived from the electrophysiological recording of optogenetically
identified dopamine neurons conducted in Tian and Uchida (2015)[55]. This modeling, while considering
major processes, does not take into account all of the complexity of the biological environment in the

brain, yet we used this model to obtain an approximate estimate of the order of changes in dopamine

concentrations and receptor occupancies.

3. **Simulation details of habenula lesion data**

3.1. Biophysical model simulations

We used the computational model described previously (methods section 2) [59] to calculate the

extracellular dopamine levels and estimate the occupancy of postsynaptic receptors from the habenula

lesion dataset. The model was driven by the average spike rate of dopamine neurons recorded from

control or lesioned animals. For each recorded dopamine neuron, the simulations were carried on a trial

by trial basis that consisted of a time window [-15, 20] sec with respect to cue onset. A relatively large

window was used to allow for the relevant variables to stabilize in its baseline, as the simulations were

initialized at zero.

For each trial, spikes were first binned with 10-ms windows and then smoothed by a Gaussian kernel

($\sigma = 0.3 \times (ISI_{mean})$). All trials were then averaged across trials, to determine the mean single-cell

response for dopamine release and D1 and D2 receptor activation. Final average dopamine concentrations

and receptor occupancies were obtained from the average of all mean single-cell responses.

Computation of receptors sensitivities from the model results

We computed the receptor sensitivity from the occupancies Occ_{D1}, Occ_{D2} and their theoretical dose-

occupancy curves. Starting from the occupancy at baseline, we derived the change in occupancy as a

function of the transients in dopamine concentration C_{DA} elicited by RPE-evoked dopamine responses, at

the level of the population average.

The ratio between these quantities corresponds to the receptors' sensitives. These are transferred as α^+

and α^- to our reinforcement learning model (model 1):

1198
$$\alpha^+ = \frac{\Delta Occ_{D1}}{\Delta C_{DA}} \quad \dots \quad \text{if } \Delta C_{DA} > 0$$

1199
$$\alpha^- = \frac{\Delta Occ_{D2}}{\Delta C_{DA}} \dots \quad \text{if } \Delta C_{DA} < 0$$

1200 Where $\Delta C_{DA}, \Delta Occ_{D1}, \Delta Occ_{D2}$ are the changes computed with respect to baseline, as: $\Delta x =$

1201 $\bar{x}_{outcome} - \bar{x}_b$, for each variable $x = \{C_{DA}, Occ_{D1}, Occ_{D2}\}$. Where $\bar{x}$ denotes the population average

1202 response for each group. The outcome responses were taken as the average from [0,1] sec after outcome

1203 onset, while the baseline was taken as the average from [-1, 0] sec with respect to cue onset.

3.2. Model 1 simulations

The simulations for Model 1 were carried out with a TD learning model with D1 and D2 populations

(methods section 1.3). We ran the simulations using the resultant receptor sensitivities from the

biophysical model as the population-level asymmetric learning rates in Model 1 (i.e., the learning rates

α^+, α^- for P and N updates). The simulations were run for 3,000 trials on the Pavlovian conditioning task

used in the study[55]. We assumed a uniform distribution of trial types across the session. Each trial

consisted of 4 states (baseline, cue, delay, reward), assuming Markovian dynamics between them. All

variables were initialized at zero. The model had as hyper-parameters a discounting factor of $\gamma = 0.99$

and a decay term $\beta = 0.002$. We report in Fig. 4, Model 1 results assuming a uniform scaling of TD

RPEs across the neuronal population. In Extended Data Fig. 12 we show that this model reproduces key

signatures of the data irrespective of the choice of the decay factor β.

The results are not dependent on a uniform scaling of TD RPEs. Given that distributional RL signatures

were preserved in the data even after habenula lesions, we also considered Model 1 under the

distributional TD learning framework (Extended Data Fig. 13). For this, we used the distribution of single

cell asymmetric scaling factors (α_i^+, α_i^-) derived from the dopamine neurons firing rates. This model also

reproduced key signatures of the data irrespective of the choice of the decay factor β (Extended Data Fig.

12).

3.3. Model 2 simulations

The simulations for Model 2 were carried out with a TD learning model. As with Model 1, simulations

were run for 3,000 trials on the Pavlovian conditioning task[55]. We assumed a uniform distribution of trial

types across the session. Each trial consisted of 4 states (baseline, cue, delay, reward), assuming

Markovian dynamics between them. All variables were initialized at zero. The model had as parameters a

discounting factor of $\gamma = 0.99$.

We used the distribution of single cell asymmetric scaling factors derived from the firing rates of

dopamine neurons as α_i^+, α_i^-. In section 1.2 we emphasized that in order to accurately compute the TD

RPE in distributional TD, we require taking samples from the estimated return distribution

$\tilde{z}_i(s_{t+1}) \sim Z(s_{t+1})$. We did this by running an optimization process where we minimize for the expectile

loss between the taken samples $\tilde{z}_i(s_{t+1})$, $V_i(s_{t+1})$ from the model, and τ_i as estimated from the data.

The problem was defined as $\text{argmin}_{s_i \dots s_m} \mathcal{L}(s, V, \tau)$ where:

$\mathcal{L}(s, V, \tau) = \frac{1}{M} \sum_{m=1}^{M} \sum_{i=1}^{N} |\tau_i - \mathbf{I}_{s_m < V_n}| (\hat{z}_m - V_i)^2$, for N neurons and M samples

In the simulations, we took M samples where M equals the number of neurons (N) and performed an

update taking the expectation across all samples as described in the methods section 1.3.

4. **Simulations details for replications of previous experimental results**

4.1. Cools et al. (2009)

We simulate the results from Cools et al. (2009) (Fig. 7, Extended Data Fig. 6-7) in which they tested the

effects of bromocriptine in altering learning rate asymmetry[24]. In their study, they performed a reversal

learning task and reported a parameter called 'relative reversal learning (RRL)', equivalent to the

difference between the positive and negative learning rates in our model. We computed this as: $\alpha^+ \alpha +$

$+\alpha - \alpha - \alpha + +\alpha - = \tau - 1 - \tau = 2\tau - 1$, reported in Fig. 7 E,F, where the parameters α^+, α^- were

computed from the slopes of the D2l (postsynaptic D2 receptors) and D1 occupancy curves $(2\tau - 1)_{occ}$

or activation curves $(2\tau - 1)_{act}$ The change in relative reversal learning in Fig. 7 H-I was calculated as

taking the difference between drug and the 'control' condition as:

$$\Delta(2\tau - 1) = (2\tau - 1)_{drug} - (2\tau - 1)_{control}.$$

We simulated the effect of bromocriptine using the biophysical model for dopamine release and receptor

occupancy (Section 2, Methods). We added an additional ligand for D2 receptors to the update equations

for occupancy:

$$\frac{dOcc_{DA,r_j}(t)}{dt} = \left(1 - Occ_{DA,r_j}(t)\right) \times k_{on}^{DA,r_j} \times C_{DA}(t) - k_{off}^{DA,r_j}$$

$$\frac{dOcc_{Drug,r_j}(t)}{dt} = \left(1 - Occ_{Drug,r_j}(t)\right) \times k_{on}^{Drug,r_j} \times C_{Drug}(t) - k_{off}^{Drug,r_j}$$

Where $r_j: \{D1, D2s, D2l\}$, and $k_{on}^{Drug,D2s} = 0.02083$, $k_{off}^{Drug,D2s} = 0.1$, $k_{on}^{Drug,D2l} = 0.04$, $k_{off}^{Drug,D2l} = 0.1$

are reported in Table 1 [97].

To calculate the effects of efficiency of the drug, we calculated the activation of D2l and D2s receptors in

the following way:

1256
$$Act_{r_j}(t) = E_{DA,r_j} \cdot Occ_{DA,r_j}(t) + E_{Drug,r_j} \cdot Occ_{Drug,r_j}(t)$$

Where $E_{DA,r_j} = 1$ is the efficiency of dopamine on the receptors activation, and $E_{Drug,r_j} < 1$ the

efficiency of the drug, for r_j: $\{D1, D2s, D2l\}$. The parameter for D1 receptors was kept at $E_{Drug,D1} = 0$

for all simulations.

To simulate the effects of D2s activation by the drug in D2l occupancy in Fig. 7b,e,h we report the effects

of $E_{Drug,D2s} = 0$ (solid lines) and $E_{Drug,D2s} = 0.6$ (dashed lines). To simulate the effect of the drug in

D2s and D2l activation in Fig. 7c,f,i we report the effects of $E_{Drug,D2s} = 0.6, E_{Drug,D2l} = 0.6$.

We show how the qualitative nature of the effects of the drug in relative reversal learning still hold

regardless of whether the parameter τ is computed from the occupancy curves (Extended Data Fig. 7, Fig.

7n,e,h) or the activation curves (Extended Data Fig. 8, Fig. 7c,f,i). In addition, in Supplementary Figure

8-9 we show that the qualitative results still hold regardless of the choice of the efficiency parameters

$E_{Drug,D2s}$ and $E_{Drug,D2l}$.

4.2. Timmer et al. (2018)

In this study[25] they reported a 'loss aversion' parameter (λ in their results).

1270
$$SUG = (1 - \lambda) \cdot p_{gain} \cdot Gain + \lambda \cdot p_{loss} \cdot Loss$$

Where SUG is the 'subjective utility' for a given option, and $p_{gain} = p_{loss}$.

In our formulation, we assume that the task in the study is performed under steady state conditions after

having learned with a learning rate (τ). With this assumption, the SUG at task performance is equivalent to

the convergent V estimate after learning. We will show that at these steady state conditions $(1 - \tau)$ is

equivalent to (λ).

Starting with the solution for V:

1277
$$SUG = V = \frac{\tau \cdot p_{gain} \cdot r_{gain} + (1 - \tau) \cdot (1 - p_{gain}) \cdot r_{loss}}{\tau \cdot p_{gain} + (1 - \tau) \cdot (1 - p_{gain})}$$

1278 Replacing for $p_{gain} = 0.5$:

1279
$$SUG = \frac{\tau \cdot r_{gain} + (1 - \tau) \cdot r_{loss}}{\tau + (1 - \tau)}$$

1280 Given that: $\tau + (1 - \tau) = 1$

$$\mathbf{SUG} = \boldsymbol{\tau} \cdot \boldsymbol{r_{gain}} + (\mathbf{1} - \boldsymbol{\tau}) \cdot \boldsymbol{r_{loss}}$$

1281

Therefore, our model, applied to their task, gives rise to the same SUG computation, with λ equivalent to

$(\mathbf{1} - \boldsymbol{\tau})$.

To generate Fig. 8F, we performed the following steps:

1. We first estimated the theoretical change in baseline DA elicited by the medication. For this, we

computed the equivalent τ for the λ they report in the OFF and ON medication conditions

$(\lambda_{OFF} = 1.51, \lambda_{ON} = 1.19)$, using the relationship: $(1 - \tau) = \lambda$. We then computed the baseline

DA levels that would give rise to the τ_{ON} and τ_{OFF}. With this, we computed the change in

baseline DA (ΔDA) equivalent to the change $\Delta\tau = \tau_{ON} - \tau_{OFF}$. This ΔDA is the theoretical

change in baseline DA elicited by the medication (Fig. 8F).

2. To generate Fig. 8F, we sampled a set of λ from a Gaussian distribution centered at a mean of

$\mu_\lambda = 1.51$ and a standard deviation of $\sigma_\lambda^2 = 3$, to emulate the distribution of λ_{OFF} they report in

the OFF condition. We then computed the equivalent τ for that set of λ with the relationship

above. We will call this the distribution of τ'_{OFF}.

3. We used the derived τ'_{OFF} distribution to compute the equivalent dopamine levels. We imposed a

change in baseline DA equal to the ΔDA computed in the first step and computed the new set of τ

for that set of new baseline DA levels (τ'_{ON}). The 'drug effect in loss aversion' reported in Fig.

8F is the $\tau'_{ON} - \tau'_{OFF}$ for each sample.

5. Details on habenula lesion data

5.1 Animals, surgery and lesions.

The rodent data we re-analyzed here were first reported in Tian and Uchida (2015)[55]. Below we provide a

brief description of the methods. Further methodological details can be found in the original paper. !2

mice were used. Bilateral habenula lesions were performed in five animals. Seven animals were in the

control group including two with sham-lesion operation, one with only small contra-lateral side lesion of

the medial habenula, and four animals without operations in the habenula. During surgery, a head plate

was implanted on the skull, and adeno-associated virus (AAV) that express channelrhodopsin-2 (ChR2)

in a Cre-dependent manner was injected into the VTA (from bregma: 3.1 mm posterior, 0.7 mm lateral,

4–4.2 mm ventral). After recovery from surgery, mice were trained on the conditioning task, after which

mice were randomly selected to be in lesion or sham-lesion group. Electrolytic lesions were made

bilaterally using a stainless-steel electrode (15 kU, MicroProbes, MS301G) with a cathodal current of 150

mA. Each side of the brain was lesioned at two locations (from bregma: 1.6 mm/1.9 mm posterior, 1.15

mm lateral, 2.93 mm depth, with a 14 angle). For sham-lesion operations, no current was applied. In the

same surgery, a microdrive containing electrodes and an optical fiber was implanted in the VTA (from

bregma: 3.1 mm posterior, 0.7 mm lateral, 3.8–4.0 mm ventral)[98].

5.2 Behavioral task

Twelve mice were trained on a probabilistic Pavlovian task. Each trial the animal experienced one of four

odor cues for 1 s, followed by a 1-s pause, followed by a reward (3.75 µl water), an aversive air puff or

nothing. Odor 1 to 3 signaled a 90%, 50% and 10% probability of reward, respectively. Odor 4 signaled a

90% probability of air puff. Odor identities were randomized across trials and included: isoamyl acetate,

eugenol, 1-hexanol, p-cymene, ethyl butyrate, 1-butanol, and carvone (1/10 dilution in paraffin oil). Inter-

trial intervals were exponentially distributed. An infrared beam was positioned in front of the water

delivery spout and each beam break was recorded as one lick event. We report the average lick rate over

the interval 500–2,000 ms after cue onset.

5.3 Electrophysiology

Recordings were made using a custom-built microdrive equipped with 200-µm-fiber optic-coupled with

eight tetrodes. DA neurons were identified optogenetically[98]. A stimulus-associated spike latency test

(SALT) algorithm[99] was used to determine whether light pulses significantly changed a neuron's spike

timing.

5.4 Neural data analysis

Data analyses were performed using MATLAB R2021b (Mathworks). To measure firing rates,

peristimulus time histograms (PSTHs) were constructed using 1-ms bins. These histograms were then

smoothed by convolving with the function $f(t) = (1 - e^{-t}) \cdot e^{-\frac{t}{\tau}}$ where τ was a time constant set to 20

ms as in [18]. 44 dopamine neurons were recorded from lesioned animals (5 animals, 30 sessions), and 45

dopamine neurons were recorded from control animals (7 animals, 35 sessions). We pooled all the cells

across animals in each group for analysis. Cue-evoked responses were defined as the average activity

from 0 to 400 ms after cue onset. Outcome-evoked responses were defined as the average activity from

2000 to 2600 ms after cue onset.

The normalization of cue response shown in Fig. 4 was carried out following a previous work[36] on a per-

cell basis as: $c_{50}^{norm} = \frac{c_{50} - \overline{c_{10}}}{\overline{c_{90}} - \overline{c_{10}}}$, where $\overline{c_{90}}$, $\overline{c_{10}}$ correspond to the mean across trials within a cell for the

90% and 10% probability cure responses. To derive the t-statistics in Fig. 4d, we performed a two-tailed t-

test of the cell's normalized responses to the 50% cue against the average midway point between

responses to the 10% cue and responses to the 90% cue.

The derivation of asymmetric scaling factors from outcome responses (τ_i), was carried out following [36],

with some modifications to adapt it to the task. The procedure is illustrated in Extended Data Fig. 3.

• To compute the reversal points, outcome responses were first aligned to the RPE for each trial

type, computed with the true expected value of each reward distribution. Assuming a fixed reward

value of 1 (arbitrary units), the expected value for the 90%, 50%, 10% reward probability trials

corresponded to 0.1, 0.5, 0.9, respectively. Given this, omission responses from the 90%, 50%,

10% reward probability trials correspond to RPEs of -0.9, -0.5 and -0.1. The rewarded responses

from the 90%, 50%, 10% reward probability trials correspond to RPEs of 0.1, 0.5 and 0.9. The

reward value is arbitrary and doesn't have an effect in this computation as it only shifts the RPE

axis by a fixed amount. The reversal point for each cell (Z_i) was defined as the RPE that

maximized the number of positive responses to RPEs greater than Z_i plus the number of negative

responses to RPEs less than Z_i. The distribution of reversal points is reported in Extended Data

Fig. 4. To obtain statistics for reliability of the computed reversal points, we partitioned the data

into random halves and estimated the reversal point for each cell separately in each half. We

repeated this procedure 1000 times with different random partitions, and we report the

distribution of Pearson's correlation across these 1000 folds (Extended Data Fig. 4).

• After measuring reversal points, we fit linear functions separately to the positive and negative

domains. Given that dopamine's responses are non-linear in the reward space but present a

putative utility function[100], we approximated the underlying utility function from the dopamine

responses to RPEs of varying magnitudes. We used these empirical utilities instead of raw RPEs

for computing the slopes that correspond to α_i^+, α_i^-. We then computed the asymmetric scaling

factors as $\tau_i = \frac{\alpha_i^+}{\alpha_i^+ + \alpha_i^-}$. We performed the same cross-validation procedure used for the reversal

points. The distribution of R value across the 1000 folds are reported in Extended Data Fig. 4.

A key prediction of distributional RL[36] is the presence of a correlation (across cells) between reversal

points Z_i and asymmetric scaling factors τ_i. To elucidate whether signatures of distributional RL were

still present after lesions, we followed the procedure given by Dabney et al. (2020)[36] to compute this

correlation. We first randomly split the data into two disjoint halves of trials. In one half, we first

calculated reversal points Z_i^1 and used them to calculate α_i^+, α_i^-. In the other half, we again calculated the

reversal points Z_i^2. The correlation we report in Extended Data Fig. 4 is between Z_i^2 and $\tau_i = \frac{\alpha_i^+}{\alpha_i^+ + \alpha_i^-}$.

5.5 Model fitting to the anticipatory licking responses

For each trial we computed the average lick rate over the interval 500–2,000 ms after cue onset. For each

model, we fit the free parameters to the lick rates using maximum likelihood estimation. The optimization

was performed using the SciPy optimization toolbox (Python) that minimized the difference between the

predicted lick rates and the ground truth ones, with a uniform prior distribution over the parameters. The

fits were done considering three RL models that had between 2 and 3 parameters. The models, parameters

and bounds used for each of them are detailed in table 2.

1379

1380

1381 **Tables**

1382 **Table 1 – Biophysical model parameters**

1383

Parameter	Abbreviation	Value
DA association constant to D2 autorreceptors	k_{on}^{D2term}	$0.3\ M^{-1}s^{-1}$
DA dissociation constant to D2 autorreceptors	k_{off}^{D2term}	$0.003\ s^{-1}$
DA association constant to D1 receptors	k_{on}^{D1}	$0.01\ M^{-1}s^{-1}$
DA dissociation constant to D1 receptors	k_{off}^{D1}	$10\ s^{-1}$
DA association constant to D2 receptors	k_{on}^{D2}	$0.2\ M^{-1}s^{-1}$
DA dissociation constant to D2 receptors	k_{off}^{D2}	$2\ s^{-1}$
Release probability from terminals	P_r	$1\ a.u.$
Release capacity from terminals	γ_{pr_n}	$2\ a.u.$
D2 autorreceptor efficacy	α	$3\ a.u.$
DAT maximal uptake capacity	$V_{max}^{pr_n}$	$1500\ nMs^{-1}$
Michaelis-Menten parameter DAT-mediated DA uptake	K_m	$160\ nM$
Constant for dopamine removal not mediated by DAT's	K_{nonDAT}	$0.04\ nM$
Bromocriptine association constant to D2 autorreceptors	$k_{on}^{Drug,D2s}$	$0.02083\ M^{-1}s^{-1}$
Bromocriptine dissociation constant to D2 autorreceptors	$k_{off}^{Drug,D2s}$	$0.1\ s^{-1}$
Bromocriptine association constant to D2 receptors	$k_{on}^{Drug,D2l}$	$0.04\ M^{-1}s^{-1}$
Bromocriptine dissociation constant to D2 receptors	$k_{off}^{Drug,D2l}$	$0.1\ s^{-1}$

1384 a.u. = arbitrary units

1385 M = mols

1386 s = seconds

1387

1388

1389   Table 2- Reinforcement learning models fit to the behavioral data from Tian & Uchida

Model	Formulation	Parameters	Parameter bounds
TD learning	$\delta = r - V$ $V \leftarrow V + \alpha \cdot \delta$ $Licking = \beta \cdot V$	α, β	$\alpha \in [.001,1]$ $\beta \in [.1,10]$
TD learning with reward sensitivity	$\delta = \rho \cdot r - V$ $V \leftarrow V + \alpha \cdot \delta$ $Licking = \beta \cdot V$	α, ρ, β	$\alpha \in [.001,1]$ $\rho \in [.001,10]$ $\beta \in [.1,10]$
Risk sensitive TD learning	$\delta = r - V$ $V \leftarrow V + \alpha^+ \cdot \delta \quad \text{if } \delta > 0$ $V \leftarrow V + \alpha^- \cdot \delta \quad \text{if } \delta < 0$ $Licking = \beta \cdot V$	$\alpha^+, \alpha^-, \beta$	$\alpha^+ \in [.001,1]$ $\alpha^- \in [.001,1]$ $\beta \in [.1,10]$

1390

1391

Data availability

1393   The neural data and simulation results reported in this article have been shared in a public

1394   deposit source in: https://osf.io/cr5mv/?view_only=bd13a2d2de1947699b56ce70610b0e9b

1395

Code availability

1397   The accession codes for the data as well as the code for analysis and simulations are available at:

1398   https://github.com/sandraromerop/D1D2_Dopamine

1399

1400

References

References

1. Brown, V. M., Zhu L., Solway A., Wang M., McCurry K., King-Casas B. & Chiu P.
Reinforcement Learning Disruptions in Individuals With Depression and Sensitivity to
Symptom Change Following Cognitive Behavioral Therapy. *JAMA Psychiatry* **78**, 1113–
1122 (2021).

2. Groman, S. M., Thompson, S. L., Lee, D. & Taylor, J. R. Reinforcement learning detuned in
addiction: integrative and translational approaches. *Trends Neurosci.* **45**, 96–105 (2022).

3. Ligneul, R., Sescousse, G., Barbalat, G., Domenech, P. & Dreher, J.-C. Shifted risk
preferences in pathological gambling. *Psychol. Med.* **43**, 1059–1068 (2013).

4. Mason, L., O'Sullivan, N., Bentall, R. P. & El-Deredy, W. Better than I thought: positive
evaluation bias in hypomania. *PLoS One* **7**, e47754 (2012).

5. Pizzagalli, D. A., Iosifescu, D., Hallett, L. A., Ratner, K. G. & Fava, M. Reduced hedonic
capacity in major depressive disorder: evidence from a probabilistic reward task. *J.*
Psychiatr. Res. **43**, 76–87 (2008).

6. Verdejo-Garcia, A., Chong, T. T.-J., Stout, J. C., Yücel, M. & London, E. D. Stages of
dysfunctional decision-making in addiction. *Pharmacol. Biochem. Behav.* **164**, 99–105
(2018).

7. Lim, T. V., Cardinal, R. N., Bullmore, E. T., Robbins, T. W. & Ersche, K. D. Impaired
Learning From Negative Feedback in Stimulant Use Disorder: Dopaminergic Modulation.
Int. J. Neuropsychopharmacol. **24**, 867–878 (2021).

8. Schönfelder, S., Langer, J., Schneider, E. E. & Wessa, M. Mania risk is characterized by an
aberrant optimistic update bias for positive life events. *J. Affect. Disord.* **218**, 313–321
(2017).

9. Dayan, P. & Daw, N. D. Decision theory, reinforcement learning, and the brain. *Cogn. Affect.*
Behav. Neurosci. **8**, 429–453 (2008).

10. Katahira, K. The relation between reinforcement learning parameters and the influence of
reinforcement history on choice behavior. *J. Math. Psychol.* **66**, 59–69 (2015).

11. Sutton, R. S. & Barto, A. G. *Reinforcement Learning: An Introduction*. (Bradford Books,
2018).

12. Maia, T. V. & Frank, M. J. From reinforcement learning models to psychiatric and
neurological disorders. *Nat. Neurosci.* **14**, 154–162 (2011).

13. Eldar, E., Rutledge, R. B., Dolan, R. J. & Niv, Y. Mood as Representation of Momentum.
Trends Cogn. Sci. **20**, 15–24 (2016).

14. Rutledge, R. B., Skandali, N., Dayan, P. & Dolan, R. J. A computational and neural model of
momentary subjective well-being. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 12252–12257 (2014).

15. Floresco, S. B., West, A. R., Ash, B., Moore, H. & Grace, A. A. Afferent modulation of
dopamine neuron firing differentially regulates tonic and phasic dopamine transmission. *Nat.*
Neurosci. **6**, 968–973 (2003).

16. Wang, Y., Toyoshima, O., Kunimatsu, J., Yamada, H. & Matsumoto, M. Tonic firing mode
of midbrain dopamine neurons continuously tracks reward values changing moment-by-
moment. *Elife* **10**, (2021).

17. Schultz, W., Dayan, P. & Montague, P. R. A neural substrate of prediction and reward.
Science **275**, 1593–1599 (1997).

18. Eshel, N., Tian, J., Bukwich, M. & Uchida, N. Dopamine neurons share common response
function for reward prediction error. *Nat. Neurosci.* **19**, 479–486 (2016).

19. Steinberg, E. Keiflin, R., Boivin, J., Witten I., Deisseroth K. & Janak P. A causal link
between prediction errors , dopamine neurons and learning. *Nat. Neurosci.* **16**, 966–973
(2013).

20. Waelti, P., Dickinson, A. & Schultz, W. Dopamine responses comply with basic assumptions
of formal learning theory. *Nature* **412**, 43–48 (2001).

21. Korn, C. W., Sharot, T., Walter, H., Heekeren, H. R. & Dolan, R. J. Depression is related to
an absence of optimistically biased belief updating about future life events. *Psychol. Med.* **44**,
579–592 (2014).

22. Rutledge, R. B., Lazzaro, S. C., Lau, B., Myers, C. E., Gluck, M. A., & Glimcher, P. W.
Dopaminergic drugs modulate learning rates and perseveration in Parkinson's patients in a
dynamic foraging task. *J. Neurosci.* **29**, 15104–15114 (2009).

23. Frank, M. J., Seeberger, L. C. & O'Reilly, R. C. By carrot or by stick: Cognitive

reinforcement learning in Parkinsonism. *Science* **306**, 1940–1943 (2004).

24. Cools, R., Frank M.J., Gibbs S., Miyakawa A., Jagust W. & D'Esposito M. Striatal

dopamine predicts outcome-specific reversal learning and its sensitivity to dopaminergic

drug administration. *J. Neurosci.* **29**, 1538–1543 (2009).

25. Timmer, M. H. M., Sescousse, G., van der Schaaf, M. E., Esselink, R. A. J. & Cools, R.

Reward learning deficits in Parkinson's disease depend on depression. *Psychol. Med.* **47**,

2302–2311 (2017).

26. Gradin, V. B., Kumar, P., Waiter, G., Ahearn, T., Stickle, C., Milders, M., Reid, I., Hall, J.,

& Steele, J. D. Expected value and prediction error abnormalities in depression and

schizophrenia. *Brain* **134**, 1751–1764 (2011).

27. Kumar, P., Waiter, G., Ahearn, T., Milders, M., Reid, I., & Steele, J. D. Abnormal temporal

difference reward-learning signals in major depression. *Brain* **131**, 2084–2093 (2008).

28. Pizzagalli, D. A., Holmes, A. J., Dillon, D. G., Goetz, E. L., Birk, J. L., Bogdan, R.,

Dougherty, D. D., Iosifescu, D. V., Rauch, S. L., & Fava, M. Reduced caudate and nucleus

accumbens response to rewards in unmedicated individuals with major depressive disorder.

Am. J. Psychiatry **166**, 702–710 (2009).

29. Robinson, O. J., Cools, R., Carlisi, C. O., Sahakian, B. J. & Drevets, W. C. Ventral striatum

response during reward and punishment reversal learning in unmedicated major depressive

disorder. *Am. J. Psychiatry* **169**, 152–159 (2012).

30. Collins, A. G. E. & Frank, M. J. Opponent actor learning (OpAL): modeling interactive

effects of striatal dopamine on reinforcement learning and choice incentive. *Psychol. Rev.*

121, 337–366 (2014).

31. Frank, M. J. Dynamic Dopamine Modulation in the Basal Ganglia: A Neurocomputational

Account of Cognitive Decits in Medicated and Non-medicated Parkinsonism. *J. Cogn.*

Neuroci. **17**, 51-72 (2005).

32. Mikhael, J. G. & Bogacz, R. Learning Reward Uncertainty in the Basal Ganglia. *PLoS*

Comput. Biol. **12**, 1–28 (2016).

33. Yagishita, S., Hayashi- Takagi, A., Ellis-Davies, G.C.R., Urakubo, H., Ishii, S. & Kasai, H.
A critical time window for dopamine actions on the structural plasticity of dendritic spines.
Science **345**, 1616–1620 (2014).

34. Iino, Y. Sawada, T., Yamaguchi, K. *et al.* Dopamine D2 receptors in discrimination learning
and spine enlargement. *Nature* **579**, 555–560 (2020).

35. Lee, S. J., Lodder, B., Chen, Y., Patriarchi, T., Tian, L & Sabatini, B. Cell-type-specific
asynchronous modulation of PKA by dopamine in learning. *Nature* **590**, 451–456 (2021).

36. Dabney, W., Kurth-Nelson, Z., Uchida, N., Starkweather, C., Hassabis, D., Munos, R. &
Botvinick, M. A distributional code for value in dopamine- based reinforcement learning.
Nature **1**, (2019).

37. Bellemare, M. G., Dabney, W. & Munos, R. A distributional perspective on reinforcement
learning. *34th International Conference on Machine Learning, ICML 2017* **1**, 693–711
(2017).

38. Lowet, A. S., Zheng, Q., Matias, S., Drugowitsch, J. & Uchida, N. Distributional
Reinforcement Learning in the Brain. *Trends Neurosci.* **43**, 980–997 (2020).

39. Rowland, M., Dadashi, R., Kumar, S., Munos, R., Bellemare, M. G. & Dabney, W. Statistics
and samples in distributional reinforcement learning. *36th International Conference on*
Machine Learning, ICML 2019 **6,** 9727–9750 (2019).

40. Mihatsch, O. & Neuneier, R. Risk-sensitive reinforcement learning. *Mach. Learn.* **49**, 267–
290 (2002).

41. Bellemare, M. G. & Dabney, W. *Distributional reinforcement learning*. (MIT Press, 2023).

42. Jones, M. C. Expectiles and M-quantiles are quantiles. *Stat. Probab. Lett.* **20**, 149–153
(1994).

43. Houk, J., Davis, J., & Beiser, D. *Models of information processing in the basal ganglia*.
(Bradford Books, 2019).

44. Gerfen, C. The neostriatal mosaic: Multiple levels of compartmental organization in the basal
ganglia. *Annu. Rev. Neurosci.* **15**, 285–320 (1992).

45. Smith, Y., Bevan, M. D., Shink, E. & Bolam, J. P. Microcircuitry of the direct and indirect
pathways of the basal ganglia. *Neuroscience* **86**, 353–387 (1998).

46. Kravitz, A. V., Tye, L. D. & Kreitzer, A. C. Distinct roles for direct and indirect pathway
striatal neurons in reinforcement. *Nat. Neurosci.* **15**, 816–818 (2012).

47. Reynolds, J. N. J., Hyland, B. I. & Wickens, J. R. A cellular mechanism of reward-related
learning. *Nature* **413,** 67–70 (2001).

48. Gerfen, C. R. & Surmeier, D. J. Modulation of striatal projection systems by dopamine.
Annu. Rev. Neurosci. **34**, 441–466 (2011).

49. Richfield, E. K., Penney, J. B. & Young, A. B. Anatomical and affinity state comparisons
between dopamine D1 and D2 receptors in the rat central nervous system. *Neuroscience* **30**,
767–777 (1989).

50. Rice, M. E. & Cragg, S. J. Dopamine spillover after quantal release: Rethinking dopamine
transmission in the nigrostriatal pathway. *Brain Res. Rev.* **58**, 303–313 (2008).

51. Gonon, F. G. & Buda, M. J. Regulation of dopamine release by impulse flow and by
autoreceptors as studied by in vivo voltammetry in the rat striatum. *Neuroscience* **14**, 765–
774 (1985).

52. Dodson, P. D., Dreyer, J. K., Jennings, K. A., Syed, E. C. J., Wade-Martins, R., Cragg, S. J.,
Bolam, J. P. & Magill, P. J. Representation of spontaneous movement by dopaminergic
neurons is cell-type selective and disrupted in parkinsonism. *Proc. Natl. Acad. Sci. U. S. A.*
113, E2180–E2188 (2016).

53. Marcott, P. F., Mamaligas, A. A. & Ford, C. P. Phasic dopamine release drives rapid
activation of striatal D2-receptors. *Neuron* **84**, 164–176 (2014).

54. Jaskir, A. & Frank, M. J. On the normative advantages of dopamine and striatal opponency
for learning and choice. *Elife* (2023).

55. Tian, J. & Uchida, N. Habenula Lesions Reveal that Multiple Mechanisms Underlie
Dopamine Prediction Errors. *Neuron* **87**, 1304–1316 (2015).

56. Cui, Y., Yang, Y., Ni, Z., Dong, Y., Cai, G., Foncelle, A., Ma, S., Sang, K., Tang, S., Li, Y.,
Shen, Y., Berry, H., Wu, S. & Hu, H. Astroglial Kir4.1 in the lateral habenula drives
neuronal bursts in depression. *Nature* **554**, 323–327 (2018).

57. Li, B., Piriz, J., Mirrione, M., Chung, C., D. Proulx, C., Schulz, D., Henn, F. & Malinow, R.
Synaptic potentiation onto habenula neurons in the learned helplessness model of depression.
Nature **470**, 535–541 (2011).

58. Yang, Y., Cui, Y., Sang, K., Dong, Y., Ni, Z., Ma, S. & Hu, H. Ketamine blocks bursting in
the lateral habenula to rapidly relieve depression. *Nature* **554**, 317–322 (2018).

59. Dreyer, J. K., Herrik, K. F., Berg, R. W. & Hounsgaard, J. D. Influence of phasic and tonic
dopamine release on receptor activation. *J. of Neurosci.* **30**, 14273–14283 (2010).

60. Mikhael, J. G., Lai, L. & Gershman, S. J. *Rational inattention and tonic dopamine*. *PLoS
Comput. Biol.* **17** e1008659 (2021).

61. Vingerhoets, F. J., Snow, B. J., Schulzer, M., Morrison, S., Ruth, T. J., Holden, J. E., Cooper,
S. & Calne, D. B. Reproducibility of fluorine-18-6-fluorodopa positron emission tomography
in normal human subjects. *J. Nucl. Med.* **35**, 18–24 (1994).

62. Cools, R., Gibbs, S. E., Miyakawa, A., Jagust, W. & D'Esposito, M. Working memory
capacity predicts dopamine synthesis capacity in the human striatum. *J. Neurosci.* **28**, 1208–
1212 (2008).

63. Hoffmann, I. S. & Cubeddu, L. X. Differential effects of bromocriptine on dopamine and
acetylcholine release modulatory receptors. *J. Neurochem.* **42**, 278–282 (1984).

64. Tissari, A. H., Rossetti, Z. L., Meloni, M., Frau, M. I. & Gessa, G. L. Autoreceptors mediate
the inhibition of dopamine synthesis by bromocriptine and lisuride in rats. *Eur. J.
Pharmacol.* **91**, 463–468 (1983).

65. Lieberman, A. Depression in Parkinson's disease -- a review. *Acta Neurol. Scand.* **113**, 1–8
(2006).

66. Leentjens, A. F. G., Van den Akker, M., Metsemakers, J. F. M., Lousberg, R. & Verhey, F.
R. J. Higher incidence of depression preceding the onset of Parkinson's disease: a register
study. *Mov. Disord.* **18**, 414–418 (2003).

67. Nilsson, F. M., Kessing, L. V., Sørensen, T. M., Andersen, P. K. & Bolwig, T. G. Major
depressive disorder in Parkinson's disease: a register-based study. *Acta Psychiatr. Scand.*
106, 202–211 (2002).

68. Remy, P., Doder, M., Lees, A., Turjanski, N. & Brooks, D. Depression in Parkinson's
disease: loss of dopamine and noradrenaline innervation in the limbic system. *Brain* **128**,
1314–1322 (2005).

69. Weintraub, D.*,* Newberg, A. B., Cary, M. S., Siderowf, A. D., Moberg, P. J., Kleiner-Fisman,
G., Duda, J. E., Stern, M. B., Mozley, D. & Katz, I. R. Striatal dopamine transporter imaging
correlates with anxiety and depression symptoms in Parkinson's disease. *J. Nucl. Med.* **46**,
227–232 (2005).

70. Kish, S. J., Shannak, K. & Hornykiewicz, O. Uneven pattern of dopamine loss in the striatum
of patients with idiopathic Parkinson's disease. Pathophysiologic and clinical implications.
N. Engl. J. Med. **318**, 876–880 (1988).

71. Timmer, M. H. M., Sescousse, G., Esselink, R. A. J., Piray, P. & Cools, R. Mechanisms
Underlying Dopamine-Induced Risky Choice in Parkinson's Disease With and Without
Depression (History). *Comput Psychiatr* **2**, 11–27 (2018).

72. Hikida, T., Kimura, K., Wada, N., Funabiki, K. & Nakanishi Shigetada, S. Distinct Roles of
Synaptic Transmission in Direct and Indirect Striatal Pathways to Reward and Aversive
Behavior. *Neuron* **66**, 896–907 (2010).

73. Hikida, T., Yawata, S., Yamaguchi, T., Danjo, T., Sasaoka, T., Wang, Y. & Nakanishi S.
Pathway-specific modulation of nucleus accumbens in reward and aversive behavior via
selective transmitter receptors. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 342–347 (2013).

74. Danjo, T., Yoshimi, K., Funabiki, K., Yawata, S. & Nakanishi, S. Aversive behavior induced
by optogenetic inactivation of ventral tegmental area dopamine neurons is mediated by
dopamine D2 receptors in the nucleus accumbens. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 6455–
6460 (2014).

75. Yamaguchi, T., Goto, A., Nakahara, I., Yawata, S., Hikida, T., Matsuda, M., Funabiki, K &
Nakanishi, S. Role of PKA signaling in D2 receptor-expressing neurons in the core of the
nucleus accumbens in aversive learning. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 11383–11388
(2015).

76. Bayer, H. M. & Glimcher, P. W. Midbrain dopamine neurons encode a quantitative reward
prediction error signal. *Neuron* **47**, 129–141 (2005).

77. Hart, A. S., Rutledge, R. B., Glimcher, P. W. & Phillips, P. E. M. Phasic dopamine release in
the rat nucleus accumbens symmetrically encodes a reward prediction error term. *J.*
Neurosci. **34**, 698–704 (2014).

78. Grace, A. A. Dysregulation of the dopamine system in the pathophysiology of schizophrenia
and depression. *Nat. Rev. Neurosci.* **17**, 524–532 (2016).

79. Anstrom, K. K., Miczek, K. A. & Budygin, E. A. Increased phasic dopamine signaling in the
mesolimbic pathway during social defeat in rats. *Neuroscience* **161**, 3–12 (2009).

80. Razzoli, M., Andreoli, M., Michielin, F., Quarta, D. & Sokal, D. M. Increased phasic activity
of VTA dopamine neurons in mice 3 weeks after repeated social defeat. *Behav. Brain Res.*
218, 253–257 (2011).

81. Markovic, T., Pederson, C. E., Massaly, N., Vachez, Y., Ruyle, B., Murphy, C. A.,
Abiraman, K., Hoon Shin, J., Garcia, J. H., Jean Yoon, H., Alvarez, V. A., Bruchas, M. R.,
Creed, M. C. & Moron, J. A. Pain induces adaptations in ventral tegmental area dopamine
neurons to drive anhedonia-like behavior. *Nat. Neurosci.* **24**, 1601–1613 (2021).

82. Guo, Z., Li, S., Wu, J., Zhu, X. & Zhang, Y. Maternal deprivation increased vulnerability to
depression in adult rats through DRD2 promoter methylation in the ventral tegmental area.
Front. Psychiatry **13**, 827667 (2022).

83. Peng, B., Hu, Q., Liu, J., Guo, S., Borgland, S. L.& Liu, S. Corticosterone attenuates
reward-seeking behavior and increases anxiety via D2 receptor signaling in ventral tegmental
area dopamine neurons. *J. Neurosci.* **41**, 1566–1581 (2021).

84. Tye, K. M., Mirzabekov, J. J., Warden, M. R., Ferenczi, E. A., Tsai, H-C., Finkelstein, J.,
Kim, S-Y., Adhikari, A., Thompson, K. R., Andalman A. S., Gunaydin, L. A., Witten I. &
Deisseroth K. Dopamine neurons modulate neural encoding and expression of depression-
related behaviour. *Nature* **493**, 537–541 (2013).

85. Baek, K., Kwon, J., Chae, J. H., Chung, Y. A., Kralik, J. D., Min, J. A., Huh, H., Choi, K.
M., Jang, K. I., Lee, N. B., Kim, S., Peterson, B. S., & Jeong, J. Heightened aversion to risk
and loss in depressed patients with a suicide attempt history. *Sci. Rep.* **7**, 11228 (2017).

86. Smoski, M. J., Lynch, T. R., Rosenthal, M. Z., Cheavens, J. S., Chapman, A. L., & Krishnan,
R. R. Decision-making and risk aversion among depressive adults. *J. Behav. Ther. Exp.*
Psychiatry **39**, 567–576 (2008).

87. van Holst, R. J., Sescousse, G., Janssen, L. K., Janssen, M., Berry, A. S., Jagust, W. J., &
Cools, R. Increased Striatal Dopamine Synthesis Capacity in Gambling Addiction. *Biol.*
Psychiatry **83**, 1036–1043 (2018).

88. Cools, R., Altamirano, L. & D'Esposito, M. Reversal learning in Parkinson's disease
depends on medication status and outcome valence. *Neuropsychologia* **44**, 1663–1673
(2006).

89. Cools, R., Barker, R. A., Sahakian, B. J. & Robbins, T. W. Enhanced or impaired cognitive
function in Parkinson's disease as a function of dopaminergic medication and task demands.
Cereb. Cortex **11**, 1136–1143 (2001).

90. Cools, R., Barker, R. A., Sahakian, B. J. & Robbins, T. W. L-Dopa medication remediates
cognitive inflexibility, but increases impulsivity in patients with Parkinson's disease.
Neuropsychologia **41**, 1431–1441 (2003).

91. Swainson, R., Rogers, R. D., Sahakian, B. J., Summers, B. A., Polkey, C. E., & Robbins, T.
W. Probabilistic learning and reversal deficits in patients with Parkinson's disease or frontal
or temporal lobe lesions: possible adverse effects of dopaminergic medication.
Neuropsychologia **38**, 596–612 (2000).

92. Cox, S. M., Frank, M. J., Larcher, K., Fellows, L. K., Clark, C. A., Leyton, M., & Dagher, A.
Striatal D1 and D2 signaling differentially predict learning from positive and negative
outcomes. *Neuroimage* **109**, 95–101 (2015).

93. Savitz, J. B. & Drevets, W. C. Neuroreceptor imaging in depression. *Neurobiol. Dis.* **52**, 49–
65 (2013).

94. Rescorla, R. A. & Wagner, A. R. *A theory of Pavlovian conditioning: Variations in the*
effectiveness of reinforcement and nonreinforcement, Classical Conditioning II. 64–99
(1972).

95. Kim, H. R., Malik, A. N., Mikhael, J. G., Bech, P., Tsutsui-Kimura, I., Sun, F., Zhang, Y., Li,

Y., Watabe-Uchida, M., Gershman, S. J. & Uchida, N. A unified framework for dopamine

signals across timescales. *Cell* **183**, 1600-1616.e25 (2020).

96. Bertsekas, D. P. & Tsitsiklis, J. *Neuro-Dynamic Programming*. (Athena Scientific, 1996).

97. Mierau, J., Schneider, F. J., Ensinger, H. A., Chio, C. L., Lajiness, M. E., & Huff, R. M.

Pramipexole binding and activation of cloned and expressed dopamine D2, D3 and D4

receptors. *Eur. J of Pharmac*. **290**, 29-36 (1995).

98. Cohen, J. Y., Haesler, S., Vong, L., Lowell, B. B. & Uchida, N. Neuron-type-specific signals

for reward and punishment in the ventral tegmental area. *Nature* **482**, 85–88 (2012).

99. Kvitsiani, D., Ranade, S., Hangya, B., Taniguchi, H., Huang, J. Z. & Kepec, A. Distinct

behavioural and network correlates of two interneuron types in prefrontal cortex. *Nature* **498**,

363–366 (2013).

100. Stauffer, W. R., Lak, A. & Schultz, W. Dopamine reward prediction error responses reflect

marginal utility. *Curr. Biol.* **24**, 2491–2500 (2014).

1666

1667

1668

1669

1670

Acknowledgements

We thank Ju Tian for providing the data used in this study; Jakob Dreyer for making the biophysical model available for open-source use; Roshan Cools, Peter Dayan and Sam Gershman for valuable feedback on the manuscript; Dan Polley, Mark Andermann and Elizabeth Phelps for mentorship and helpful discussions. We thank Bernardo Sabatini for discussions, and Isobel Green, Paul Masset, Mitsuko Watabe-Uchida and all the rest of the Uchida lab members for helpful comments at the different stages of this work. This work is supported by NIH BRAIN Initiative grants (R01NS116753, U19NS113201), the Simons Collaboration on Global Brain, and the Bipolar Disorder Seed Grant from the Harvard Brain Science Institute.

Author information

Contributions

S.R.P. and N.U. conceived the project. S.R.P performed the modeling work. S.R.P. wrote the first draft and S.R.P and N.U. edited the paper.

Corresponding author

Correspondence to Naoshige Uchida (uchida@mcb.harvard.edu) and Sandra Romero Pinto (sromeropinto@g.harvard.edu).

Competing interest statement

The authors declare no competing interests.

Figures

Figure 1 | Reinforcement learning models.

a. Traditional reinforcement learning with a single learning rate (α) for both positive and negative RPEs (δ) for the value updates (left). This update rule makes value estimate (V) converge on the expected value of the reward distribution (middle). When the reward probability is varied (i.e., for Bernoulli distributions), the V at convergence scales linearly with the reward probability (right).

b. Risk-sensitive reinforcement learning with different learning rates (α^+, α^-) for positive and negative RPEs, respectively (left). This update rule makes value estimate (V) converge on the quantities that are higher or lower than the expected value of the reward distribution (middle). As the reward probabilities are varied, the convexity of the convergent value V changes depending on the asymmetry between α^+ and α^- (Methods 1.3.3). The level of the bias is determined by the asymmetric learning rate parameter τ (right).

c. Distributional reinforcement learning contains a set of value predictors (V_i) each with a given learning rate for positive and negative RPEs (α_i^+, α_i^-, respectively) (left). This makes each value

1712   predictor converge on the quantity equal to the τ_i-th expectile of the reward distribution. Thus,

1713   each value V_i represents an expectile, and together the set of V_i represents the entire distribution

1714   (Methods 1.2) (right).

1715

1716

1717

Figure 2 | Biologically inspired reinforcement learning model.

a. Schematic of the basal ganglia circuitry. Dopaminergic neurons in the VTA modulate plasticity at the level of the cortico-striatal synapses on SPNs in the NAc. The SPNs are subdivided depending on the dopamine receptor type they express (D1R or D2R).

b. Dose-occupancy curves for the D1R and D2R describing receptor occupancies as a function of dopamine concentrations. The curves are shifted between each other due to the different affinities of the receptors. The arrows represent 3-fold increase ("burst") and decrease ("pause") in dopamine concentrations, which causes left-ward or right-ward shifts of the same magnitudes in the log-scale.

c. Schematic of the plasticity rules of VTA-NAc circuitry[33–35]. Transient increases in dopamine, caused by bursts in firing rate of dopamine neurons, generates increases in PKA activity in D1R-expressing SPNs, leading to LTP in the cortico-striatal synapses. Transient decreases in dopamine, caused by pauses in firing rate of dopamine neurons, generates increases in PKA activity in D2R-expressing SPNs, leading to LTP in the cortico-striatal synapses.

d. Schematic and equations of biologically inspired reinforcement learning model[32]

VTA, ventral tegmental area; NAc, nucleus accumbens; SPN, spiny projection neurons; D1R, D1-type dopamine receptor; D2R, D2-type dopamine receptor; PKA, protein kinase A; LTP, long-term potentiation.

1736

Figure 3 | Potential mechanisms for asymmetric learning.

a. Schematic of the mechanism by which increases or decreases in baseline dopamine modulates the degree to which bursts and pauses in dopamine causes changes in D1R and D2R occupancy. Increases in baseline dopamine makes dopamine pauses to cause greater decreases in D2R occupancy than the increases in D1R occupancy caused by dopamine bursts. Conversely, decreases in dopamine, makes dopamine bursts to cause smaller increases in D1R occupancy than the decreases in D2R occupancy caused by dopamine pauses.

b. Schematic of the change in receptor occupancies in D1R and D2R, for a given transient
increase ('burst') or decrease ('pause') in dopamine, receptively. A pause and a burst in
dopamine correspond to $\delta < 0$ and $\delta > 0$ in the model. The slope is modulated by the baseline
dopamine (colormap) and corresponds to the receptor's sensitivity to dopamine transients.

c. Receptor sensitivity for D1R and D2R as a function of baseline dopamine. In Model 1, we
assume that the receptor sensitivity acts as a scaling factor on the PKA activity induced by burst
and pauses. That is, $\mathrm{PKA_{D1}} \propto \alpha^+ \cdot \delta \cdot \mathbf{1}_{\delta>0}$ and $\mathrm{PKA_{D2}} \propto \alpha^- \cdot \delta \cdot \mathbf{1}_{\delta<0}$.

d. Asymmetric scaling factor (τ) as a function of baseline dopamine. Colors depict how
'optimistic' or 'pessimistic' the convergent value estimate will be when learning with a given τ.

e. Model 2. Left, the relationship between dopamine reward responses (spikes/s) and RPEs. The
slopes of these response functions correspond to the asymmetric learning rates (α^+, α^-) for
positive and negative RPEs, respectively. Colors depict how 'optimistic' or 'pessimistic' the
convergent value estimate will be when learning with a given asymmetric scaling factor.

1757

1758

1759

Figure 4 | Habenula lesions leads to optimistic reward-seeking behavior and cue-evoked responses in dopamine neurons.

a. Schematic of the experiment performed by Tian and Uchida (2015)[55]. Animals were trained in a classical conditioning task in which 3 odor cues predicted rewards of different probabilities (10%, 50%, 90%) and one odor cue predicted 80% probability of an air puff. Animals then underwent habenula ($n = 5$) or sham ($n = 7$) lesions and trained on the task again. The neural recordings were performed from optotagged VTA dopamine neurons once behavior stabilized.

b. Anticipatory licking across sessions after lesions (left,). There was a significant increase in anticipatory licking to the 10% (U-statistic = –2.895, $P = 0.003792$, two-sided Mann-Whitney U-test), 50% (U-statistic = –5.579, $P < 1 \times 10^{-9}$, two-sided Mann-Whitney U-test) and 90% (U-

statistic = –3.487, P =0.00048, two-sided Mann-Whitney U-test) cues ($n = 31$ for control $n = 30$

for lesion) that results from progressive changes across sessions. The anticipatory licking curves

show a linear scaling with reward probability in the control group, and a convex curve for the

lesion group (mean ± s.e.m across animals, U-statistic = –6.444, $P < 1 \times 10^{-}$, two-sided Mann-

Whitney U-test for the 50% cue normalized response). These curves are predicted by RL agents

with symmetric and asymmetric ($\alpha^+ > \alpha^-$) learning rates for the control and lesion groups,

respectively, assuming a linear mapping between anticipatory licking and value prediction.

c. RL model fits to the anticipatory licking on a trial-by-trial basis using a risk-sensitive RL

models that allows for separate learning rates of positive and negative RPEs. Each dot represents

a session ($n = 35$ control, $n = 30$ lesion) and each color a mouse ($n = 7$ control, $n = 5$ lesion). The

fits show a significant difference in the learning rates between control and lesion groups (U-

statistic = –4.679, $P < 1.0 \times 10^{-5}$, pooling sessions across mice in each group).

d. Cue-evoked dopamine responses from opto-tagged VTA dopamine neurons (mean ± s.e.m

across neurons, $n = 45$ control group, $n = 44$ lesion group). There was a decrease in the absolute

magnitude of responses to the 90% cue (U-statistic = 3.249, $P = 0.0011$, two-sided Mann-

Whitney U-test) after habenula lesions (left). The normalized cue-evoked responses show the

similar pattern as the normalized anticipatory-licking with a linear and convex function for the

control and lesion groups, respectively, with a significant increase in normalized response to the

50% cue after lesions (U-statistic = –3.824, $P = 0.000131$, two-sided Mann-Whitney U-test)

These curves are predicted by agents with symmetric and asymmetric learning rates for control

and lesion groups, respectively.

e. Distribution of t-statistics comparing the cue-evoked response to the linear interpolation point

between the 90% and 10% cue-evoked responses for each dopamine neuron. The distribution of

t-statistics for the control and lesion cases was wider than what is expected from random noise

(Monte Carlo test for standard deviation different from zero: $P = 0.0222$ control, $P = 0.0217$

lesion, 1000 batches). The distribution was shifted to values larger than 0 in the lesion case

(Monte Carlo test for mean larger than zero: $P = 1$ control, $P = 0.022$ lesion, 1000 batches)

indicative of an optimistic bias in the distribution. The lesion group distribution was also

significantly shifted to higher values with respect to the control group distribution (U-statistic =

–2.815, $P = 0.0024$, single-sided Mann-Whitney U-test). Arrow heads: the mean of the *t*-

statistics.

f. Example of *t*-statistics calculations for dopamine neurons taken from the control group (mean

± s.e.m across trials). A *t*-statistic value close to 0 indicates linear scaling of cue-evoked

responses with reward probability; a *t*-statistics value lower or greater than 0 indicates a concave

or convex function of cue-evoked responses against reward probability, indicative of a

pessimistic or an optimistic bias, respectively.

1806

1807

1808

Figure 5 | Model 2 cannot explain optimistic biases in behavior and cue-evoked dopamine responses of habenula lesioned animals.

a. Possible changes in habenula lesion mice that could explain optimistic biases based on Model 2. At the level of the population dopamine responses, an optimistic bias can be caused by an increase in the slope of the average reward responses to positive RPEs and/or a decrease in the slope of the average reward responses to negative RPEs.

b. At the level of the distribution of individual dopamine neuron responses, an optimistic bias can be caused by an overall increase in the mean of the distribution of asymmetric scaling factors (τ_i), computed from each individual neuron response function.

c. Observed reward responses as a function of RPEs, averaged across the population of dopamine

neurons for the control and lesion groups (left, mean ± s.e.m across neurons, $n = 45$ control

group, $n = 44$ lesion group). There was a significant decrease in the reward responses for the

50% cue (U-statistic = 3.726, $P = 0.000195$, two-sided Mann-Whitney U-test) and 90% cue (U-

statistic = 2.987, $P = 0.00281$, two-sided Mann-Whitney U-test), and for the omission responses

for the 90% cue (U-statistic = -4.940, $P < 10^{-4}$, two-sided Mann-Whitney U-test). Distribution of

asymmetric scaling factors (τ), computed from the average response function over the recorded

neurons for the control and lesion groups (right). The distributions are the result of bootstrapping

by randomly sampling neurons in 5,000 iterations. The distribution of differences between the

obtained asymmetric scaling factors ($\tau_{lesion} - \tau_{control}$) was not significantly larger than zero (5th

percentile = –0.1605).

d. Distribution of asymmetric scaling factors (τ_i), computed from each individual neuron

response function for the control and lesion groups. Each dot represents a single neuron (n = 45

control group, $n = 44$ lesion group), and the neurons were sorted by asymmetric scaling factors

(τ_i). The means were not significantly different (right) (t-statistic = 0.3277, $P = 0.627$, t-test).

e. Value predictions based on a TD learning model trained using the assumptions of Model 2 and

the asymmetric scaling factors derived from the data. The model did not show any optimistic

bias in the value predictors of the model trained with the lesion-derived asymmetric scaling

factors.

f. TD errors at cue show no signs of an optimistic bias in the model trained with the lesion-

derived asymmetric scaling factors.

Centre of box plot shows the median; edges are 25th and 75th percentiles; and whiskers are the

most extreme data points not considered as outliers.

1841

1842

1843

Figure 6 | Model 1 can account for optimistic biases in reward-seeking behavior and cue-evoked dopamine responses.

a. Schematic of the analysis. A biophysical model was used to predict dopamine concentrations, receptor occupancies, and value learning based on firing rates of dopamine neurons recorded in Tian et al. (2015).

b. Average firing rates of dopamine neurons across the population for the control and lesion groups (left, n= 45 control group, n= 44 lesion group). Baseline firing rates were significantly greater in the lesion compared to the control group (right) (U-statistic = –2.429, $P = 0.0151$, single-sided Mann-Whitney U-test).

c. Dopamine concentrations predicted from the firing rates of dopamine neurons based on the
biophysical model of dopamine. Predictions for 90% reward trials are shown.

d. Receptor occupancies predicted by the same biophysical model. Predictions for rewarded
(left) and reward omission (right) trials in 90% reward trials are shown separately for D1R (left)
and D2R (right), respectively ($n = 45$ control group, $n = 44$ lesion group).

e. Mean dopamine concentrations at baseline predicted by the model (U-statistic $= -2.109$, $P =$
0.0175, single-sided Mann-Whitney U-test).

f. Mean receptor occupancies at baseline predicted by the model ($n = 45$ control group, $n = 44$
lesion group). There is a significant increase in occupancies for both the D1R and D2R in the
lesion compared to the control group (U-statistic $= -2.1664$, $P = 0.0151$, U-statistic $= -2.1328$, P
$= 0.0165$ for D1R and D2R respectively, single-sided Mann-Whitney U-test).

g. Schematic showing the model predicted changes in dopamine concentrations and receptor
occupancies for the control (black) and lesion (red) groups. The arrows depict the increase or
decrease in occupancy for a positive or negative dopamine transient of a fixed magnitude.

h. Changes in receptor occupancy as a function of dopamine transients predicted by the model.
The slope for the positive and negative domains correspond to the receptor sensitivities of D1R
and D2R (α^+, α^-), respectively.

i. Asymmetric scaling factors derived from the receptors' sensitivities for the control and lesion
groups (i.e., τ in model 1, $n = 45$ control group, $n = 44$ lesion group). There was a significant
increase in the lesion group with respect to controls (U-statistic $= -12.205$, $P < 1.0 \times 10^{-6}$, single-
sided Mann-Whitney U-test). Note that the increase in the asymmetry was driven mainly due to
decreases in D2R sensitivity (panel h).

j. Value predictions at baseline in a TD learning model trained with the receptor sensitivities
derived from the biophysical. There was a significant increase in the value predictors at baseline
in the model using the lesion group's derived parameters with respect to control. controls (t-
statistic $= -6.417$, $P < 1.0 \times 10^{-6}$, t-test).

k. Value predictions at convergence of a TD learning model trained using the assumptions of
Model 1 and the asymmetric scaling factors derived from receptors' sensitivities predicted by the
biophysical model. The model led to a significant increase in the value predictions for all cues
(U-statistic $= -4.690$, $P < 1.0 \times 10^{-4}$, U-statistic $= -4.734$, $P < 1.0 \times 10^{-4}$, U-statistic $= -4.602$, $P <$

1.0×10^{-4}, single-sided Mann-Whitney U-test, for the 10%, 50% and 90% reward probability

cues) and an optimistic bias in the normalized value prediction to the 50% reward probability cue

(t-statistic $= -5.576$, $P < 1.0 \times 10^{-4}$, t-test) in accordance with the anticipatory licking observed in

the data.

l. Predicted cue responses. There is an overall decrease in RPEs in lesioned animals (left) due to

an increase in the baseline (pre-cue) value prediction (U-statistic $=4.932$, $P < 1.0 \times 10^{-5}$, U-

statistic $= -3.658$, $P = 0.00025$, U-statistic $= 4.734$, $P < 1.0 \times 10^{-4}$, single-sided Mann-Whitney

U-test for the 10%, 50% and 90% reward probability cues), which is consistent with the

decreases in the absolute magnitudes of dopamine cue-evoked responses in the lesion group (Fig.

4c). The normalized TD errors at for the 50% reward probability cue show signs of an optimistic

bias (U-statistic $= -4.624$, $P < 1.0 \times 10^{-4}$, single-sided Mann–Whitney U-test).

Centre of box plot shows the median; edges are 25th and 75th percentiles; and whiskers are the

most extreme data points not considered as outliers.

1896

Figure 7 | Model 1 predicts asymmetric learning rates in healthy humans given inter-individual differences in baseline dopamine, and in Parkinson's disease patients given inter-individual differences in depressive-like symptoms.

a. Schematic of the events occurring at dopaminergic axon terminal. Pre- and post-synaptic sites predominantly express D2s (short) and D2l (long) subtypes, respectively.

b. "Relative reversal learning (RRL)" under placebo conditions as a function of dopamine striatal synthesis capacity measured with PET radio imaging (black dots, left y-axis, bottom x-axis).

Figure taken from Cools et al. (2009)[24]. Positive values of RRL indicate a bias favoring learning

from gains relative to losses, and vice versa for negative values of RRL. There was a positive

relationship between RRL and dopamine synthesis capacity. Model 1 predictions of RRL ($2\tau -$

1 in Model 1) as a function of baseline dopamine using the receptors occupancy curve,

recapitulate the positive relationship shown in the results from Cools et al. (2009)[24] (gray line,

right y-axis, top x-axis).

c. The change in RRL induced by bromocriptine was negatively correlated with striatal

dopamine synthesis capacity. Figure from Cools et al. (2009)[24] (black dots, left y-axis, bottom x-

axis). Model 1 recapitulates qualitatively the effect of bromocriptine in RRL, equivalent to

$\Delta(2\tau - 1)$. The solid light green line represents the $\Delta(2\tau - 1)$ when considering bromocriptine's

effect on *D2l occupancy only*; the dashed line represents the $\Delta(2\tau - 1)$ when both *D2l and D2s*

occupancy was considered; and the dark green line represents the $\Delta(2\tau - 1)$ when both *D2l and*

D2s activation was considered (this includes the fact that bromocriptine is a partial agonist for

the D2l and D2s receptors). The curves were obtained by imposing a concentration of $10^{0.8}$ nM

of bromocriptine in the biophysical model.

d.   Receptor occupancy curves for the D2l receptors at baseline (grey line) and in the presence of

$10^{0.8}$ nM of bromocriptine: Solid light green line corresponds to considering bromocriptine's

effects on D2l receptors occupancy alone; dashed line, corresponds to considering

bromocriptine's effects on both D2l and D2s receptors; solid dark green line corresponds to

considering bromocriptine's effect on the activation curves of both D2l and D2s receptors. The

binding of the drug to the D2l receptors alone causes an increase in occupancy. This happens to a

larger extent when starting from a low dopamine level at baseline than in high dopamine levels.

The binding of the drug to D2s receptors in addition to D2l receptors causes a rightward shift in

the curves. The activation levels are lower than 1 even at the drug levels where occupancy is

close to 1, due to the lower efficiency of bromocriptine in receptor activation (Methos 4.1). See

Extended Figure 6 and 7 for the effect of changing bromocriptine's concentration and efficiency

of activation.

e. Same as in panel **d.** but now reporting $(2\tau - 1)$ calculated from the D2l receptor's occupancy

and activation curves. An increase in $2\tau - 1$ happens to a larger extent when starting from a low

dopamine level than from high dopamine level.

f. Same as in panel **d.** but now reporting $\Delta(2\tau - 1)$ calculated from the D2l receptor's occupancy and activation curves. Model 1 recapitulates qualitatively the effect of bromocriptine on RRL.

g. The effect of PD medication (L-DOPA) on loss aversion is negatively correlated with their off-medication depression score. Figure from Timmer et al. (2018)[71].

h. Model 1 recapitulates qualitatively the effect of PD medication in loss aversion. We assumed that the asymmetry in favor of learning from losses relative to gains $(1 - \tau)_{off}$ scales with the baseline dopamine levels. Given this, we derived a distribution of off-medication baseline dopamine levels centered around the mean $(1 - \tau)_{off}$ derived from the data of Timmer et al. (2018)[71] (see methods 0). We then imposed a fixed increase in baseline dopamine to simulate L-DOPA effects. We derived the new loss-aversion parameter $(1 - \tau)_{on}$ at the shifted baseline dopamine levels. The y-axis shows the change in loss aversion for each sample of the distribution of baseline dopamine levels. If the off-medication depression score is correlated with $(1 - \tau)_{off}$ then model would predict the result in Timmer et al. (2018)[71].

PD: Parkinson's disease.

Extended data Figures

1952 **Extended Data Fig. 1 | Variables of model 1 show convergence irrespective of the value of**

1953 **the decay factor.**

1954 **a**. Value predictors V (left), P population (middle) and N population (right) across trials of

1955 training for an RL agent of model 1. Color of lines denotes the value of the decay factor (β) in

1956 the update rules for the P and N populations. Colormap is the same for all figures (left). All the

1957 model variables show convergence for every value of the decay factor β.

1958 **b**. Difference in the variables estimates between consecutive trials of training, for the value

1959 predictors V (left, ΔV), P population (middle, ΔP) and N population (right, ΔN). All the variables

1960 show convergence for every value of the decay factor β (shown as a $\Delta V, \Delta P, \Delta N$ equal to zero).

a TD learning

$$\delta = r - V$$
$$V \leftarrow V + \alpha \cdot \delta$$
$$Licking = \eta \cdot V$$

b TD learning with reward sensitivity

$$\delta = \rho \cdot r - V$$
$$V \leftarrow V + \alpha \cdot \delta$$
$$Licking = \eta \cdot V$$

c Risk sensitive reinforcement learning

$$\delta = r - V$$
$$V \leftarrow V + \alpha^+ \cdot \delta \quad if \ \delta > 0$$
$$V \leftarrow V + \alpha^- \cdot \delta \quad if \ \delta < 0$$
$$Licking = \eta \cdot V$$

1961

1962

1963 **Extended Data Fig. 2 | RL model fits to the trial-by-trial anticipatory licking responses.**

1964 **a**. TD learning fits reveal no significant difference across groups in the learning rate (left, U-
1965 statistic = –4.954, P =0620, two-sided Mann-Whitney U-test) nor in the linear mapping between
1966 value predictions and anticipatory licking responses (U-statistic = –1.445, $P = 0.148$, two-sided
1967 Mann-Whitney U-test).

1968 **b**. Model fits of TD learning with reward sensitivity reveal no difference across groups in the
1969 learning rate (left, U-statistic = 0.206, $P = 0.836$, two-sided Mann-Whitney U-test) nor in the

1970   linear mapping between value predictions and anticipatory licking responses (middle, U-statistic

1971   = –0.7844, P = 0.4327, two-sided Mann-Whitney U-test), nor in the reward sensitivity (right) (U-

1972   statistic 0.545, P = 0.605, two-sided Mann-Whitney U-test).

1973   **c**. Model fits of TD learning with asymmetric learning rates for positive vs negative RPEs. This

1974   model reveals a significant difference across groups in the asymmetry between α^+ and α^- (U-

1975   statistic = –4.678, $P < 1.0 \times 10^{-5}$, two-sided Mann-Whitney U-test) and a small but significant

1976   difference between the linear mapping between value predictions and anticipatory licking

1977   responses (right, U-statistic = 2.33, P = 0.02, two-sided Mann-Whitney U-test).

1978

1979

Extended Data Fig. 3 | Signatures of distributional reinforcement learning model are preserved after habenula lesions.

a. RPE -evoked responses at outcome as a function of the theoretical RPE for each trial type. The figure shows the average response function across neurons from the control group. The

1985   computation of zero-crossing points and asymmetric scaling factors is carried out in the 'utility

1986   space'(the average response function) as in [36] to account for response nonlinearities.

1987   **b**. Example of the response function of 3 dopamine neurons from the control group ordered by

1988   their asymmetric scaling factors: pessimistic, neutral and optimistic, from top to bottom.

1989   **c**. Computation of zero-crossing points for the neurons in B. The reversal points for each cell (Z_i)

1990   were defined as the point in utility space that maximized the number of positive responses to

1991   points greater than Z_i plus the number of negative responses to points less than Z_i. The y-axis

1992   shows the sum of responses below and above each point in the utility space. The zero-crossing

1993   point is shown as the maxima in this function.

1994   **d**. Computation of asymmetric scaling factors for the neurons in c. Here, the responses functions

1995   in b have been projected to the utility space in A and realigned according to their zero-crossing

1996   points. The asymmetric learning rates (α^+, α^-) are taken to be the slopes of these response

1997   functions.

1998

1999

2000

Extended Data Fig. 4 | Distributional reinforcement learning variables from the Habenula

lesion dataset.

a. Distribution of asymmetric scaling factors for the dopamine neurons from the control (left)

and lesion (right) groups. The error bars were derived by randomly sampling trials to compute

the asymmetric scaling factors for 1000 iterations.

b. Distribution of zero crossing points for the control (left) and lesion(right) groups. The error

bars were derived as in a.

c. Correlation of asymmetric scaling factors (x-axis) and zero-crossing points (y-axis) computed

on disjoint halves of trials for an example partition.

d. Distribution of correlation coefficients between asymmetric scaling factors (x-axis) and zero-

crossing points (y-axis) across disjoint halves of trials for 1000 partitions for the control and

lesion groups.

e. Correlation between zero-crossing points computed on disjoint halves of trials for an example

partition.

f. Distribution of correlation coefficients between zero-crossing points computed on disjoint

halves of trials for 1000 partitions for the control and lesion groups.

g. Correlation between asymmetric scaling factors computed on disjoint halves of trials for an

example partition.

h. Distribution of correlation coefficients between asymmetric scaling factors computed on

disjoint halves of trials for 1000 partitions for the control and lesion groups.

2021

2022

2023

Extended Data Fig. 5 | Biases in cue-evoked responses in the Habenula lesion data cannot be explained by asymmetric scaling of RPEs (Model 2).

a. Value predictors derived from model 1 with TD learning for a set of baseline dopamine levels (colormap). The optimistic and pessimistic biases are present.

b. Cue responses derived from model 1 with TD learning for a set of baseline dopamine levels

(colormap). The optimistic and pessimistic biases are revealed when the responses are

normalized.

c. Value predictors derived from model 2 with TD learning for a set of baseline dopamine levels

(colormap). The optimistic and pessimistic biases are present.

d. Cue responses derived from model 2 with TD learning for a set of baseline dopamine levels

(colormap). The optimistic and pessimistic biases are absent in both the normalized and the raw

TD errors.

e. Mean across the distribution of value predictors derived from model 1 with distributional TD

learning for a set of baseline dopamine levels (colormap). The optimistic and pessimistic biases

are present.

f. Mean across the distribution of cue responses derived from model 1 with distributional TD

learning for a set of baseline dopamine levels (colormap). The optimistic and pessimistic biases

are revealed when the responses are normalized.

g. Mean across the distribution of value predictors derived from model 2 with distributional TD

learning for a set of baseline dopamine levels (colormap). The optimistic and pessimistic biases

are present.

h. Mean across the distribution of cue responses derived from model 2 with distributional TD

learning for a set of baseline dopamine levels (colormap). The optimistic and pessimistic biases

are absent in both the normalized and the raw TD errors.

2048

2049

2050

Extended Data Fig. 6 | Model 1 predicts asymmetric learning rates and the effect of bromocriptine in healthy humans given inter-individual differences in baseline dopamine.

a. Occupancy curves for the D2l receptors at baseline (grey line) and when considering bromocriptine's effects in D2l receptors alone. The binding of the drug to the D2l receptors alone causes an increase in the occupancy. This happens to a larger extent when starting from a low dopamine level at baseline than in high dopamine levels.

b. Occupancy curves for the D2l receptors at baseline (grey line) and when considering bromocriptine's effects in both D2l and D2s receptors. The binding of the drug to D2s receptors causes a rightwards shifts in the curves.

c. Activation curves for the D2l receptors at baseline (grey line) and when considering bromocriptine's effects in D2l receptors alone, including the *partial* quality of the agonism of this drug on the receptor (i.e., efficiency <1).

d. Activation curves for the D2l receptors at baseline (grey line) and when considering bromocriptine's effects in both D2l and D2s receptors, including the *partial* quality of the agonism.

e. Relative reversal learning (RRL) calculated as $2\tau - 1$ in model 1, as a function of baseline dopamine (x-axis) and drug concentration (color) using the receptors occupancy curve, considering only bromocriptine's effect in D2l receptors.

f. Relative reversal learning (RRL) calculated as $2\tau - 1$ in model 1, as a function of baseline dopamine (x-axis) and drug concentration (color) using the receptors occupancy curve, considering bromocriptine's effect in both D2l and D2s receptors.

g. $2\tau - 1$ in model 1, as a function of baseline dopamine (x-axis) and drug concentration (color) using the receptors activation curve, considering only bromocriptine's effect in D2l receptors.

h. $2\tau - 1$ in model 1, as a function of baseline dopamine (x-axis) and drug concentration (color) using the receptors activation curve, considering bromocriptine's effect in both D2l and D2s receptors.

i. The change in $2\tau - 1$ induced by the drug at different concentrations (color) with respect to the baseline condition, as a function of baseline dopamine (x-axis). The curves represent the change when calculating $2\tau - 1$ from the occupancy curves considering only D2l binding.

j. The change in $2\tau - 1$ induced by the drug at different concentrations (color) with respect to the baseline condition, as a function of baseline dopamine (x-axis). The curves represent the change when calculating $2\tau - 1$ from the occupancy curves considering both D2l and D2s binding

k. The change in $2\tau - 1$ induced by the drug at different concentrations (color) with respect to the baseline condition, as a function of baseline dopamine (x-axis). The curves represent the change when calculating $2\tau - 1$ from the activation curves considering only D2l activation.

l. The change in $2\tau - 1$ induced by the drug at different concentrations (color) with respect to the baseline condition, as a function of baseline dopamine (x-axis). The curves represent the

2089   change when calculating $2\tau - 1$ from the activation curves considering both D2l and D2s

2090   activation.

2091   The parameters of efficiency of activation of D2 receptors by the drug $\left(D2l_{eff}, D2s_{eff}\right)$ used in

2092   each column are reported at the bottom of the figure.

2093

2094

Extended Data Fig. 7 | Robustness of the effect of bromocriptine in the relative reversal learning calculated from the occupancy curves to the choice of the drug efficiency parameter.

2099 The qualitative effects on bromocriptine in the change in relative reversal learning $\Delta(2\tau - 1)$

2100 calculated from the D2 *occupancy* curves. Results hold regardless of the choice of the efficiency

2101 of the drug on D2l (rows) or D2s (columns) efficiency.

2102

2103

2104

Extended Data Fig. 8 | Robustness of the effect of bromocriptine in the relative reversal learning calculated from the activation curves to the choice of the drug efficiency parameter.

2108   The qualitative effects on bromocriptine in the change in relative reversal learning $\Delta(2\tau - 1)$

2109   calculated from the D2 *activation* curves. Results hold regardless of the choice of the efficiency

2110   of the drug on D2l (rows) or D2s (columns) efficiency.

2111

Extended Data Fig. 9 | The qualitative aspects of Model 1 are preserved irrespective of the assumption made about the changes in baseline dopamine caused by dopamine transients.

a. Computation of receptor sensitivities (i.e., slope of dose-occupancy curves, α^+, α^-) assuming logarithmic (left) or linear (middle) changes in baseline dopamine induced by dopamine transients (logΔDA, linΔDA, respectively). The absolute magnitude of the slopes differs depending on the assumption made about the changes in baseline dopamine (logarithmic vs linear) but the asymmetric scaling factor presents only a small shift in the curve as a function of baseline dopamine (right). The qualitative aspects of the model (i.e., non-monotonic relationship of the asymmetric scaling factor with baseline dopamine) is preserved regardless on this assumption.

2125

Extended Data Fig. 10 | Change in dopamine firing rates, dopamine concentration and receptor occupancy as a function of RPEs in the linear scale or logarithmic scale.

a. Distributions of the slopes of the change in firing rate derived as a function of RPEs in the positive and negative domains, computed in the linear (left) or logarithmic (right) scale,

calculated at a single neuron level. The slopes are asymmetric if considered in the linear scale,

with the negative transients presenting a shallower slope than the positive ones. The slopes are

symmetric if considered in the logarithmic scale.

b. Distributions of the slopes of the change in dopamine levels derived from the biophysical

model as a function of RPEs in the positive and negative domains, computed in the linear (left)

or logarithmic (right) scale, calculated at a single neuron level. The slopes are again asymmetric

if considered in the linear scale but symmetric if considered in the logarithmic scale.

c. Slope of the change in receptor occupancy derived from the biophysical model for a given

RPE in the positive and negative domains, computed in the linear (left) or logarithmic (right)

scale, calculated at a single neuron level. The slopes are symmetric if considered in the linear

scale but asymmetric if considered in the logarithmic scale.

2141

Extended Data Fig. 11 | Relationship between changes in firing rates and changes in dopamine concentration derived from the biophysical model.

a. Linear fits to the relationship between changes in firing rates and changes in dopamine concentration evoked by the TD error at outcome in the linear scale for the control group. The fits are done separately for each trial type.

b. Same as a, but fits are done in the logarithmic scale.

c. Same as a, but fits are done for the lesion group.

d. Same as c, but fits are done in the logarithmic scale.

e. Distribution of the Pearson correlation coefficients (top) and means squared error (MSE,

bottom) between the predicted change in dopamine concentration by the linear regression and the

ground truth derived from the biophysical model. The coefficients are derived from the fits in

figures a-e, done by pooling all trials for each trial type (each point each trial type, with black for

control and red for lesion group). There was a significant increase in the Pearson correlation

coefficient and a near significant decrease in the MSE if the changes are considered to happen in

the logarithmic scale.

f. Same as E, but the linear regression fits are done for each neuron separately by pooling all

trials. There was a significant increase in the single-cell distribution of Pearson correlation

coefficients (top) and a significant decrease in the MSE distribution (bottom) if the changes are

considered to happen in the logarithmic scale.

2162

2163

2164

Extended Data Fig. 12 | Model 1 captures signatures of the data irrespective of the choice of

the decay factor and is compatible with distributional RL.

a. Model 1 with standard TD learning. Simulations were run using the receptors sensitivities

from the biophysical models and data-derived asymmetric scaling factors (see Methods 3.3).

The model's predictions capture the signatures in cue-evoked dopamine responses (left) and

value predictions (right) irrespective of the choice of the decay factor (β).

b. Model 1 within the distributional RL framework (see Methods 3.3). The model's predictions

also capture the signatures in cue-evoked dopamine responses (left) and value predictions (right)

irrespective of the choice of the decay factor (β). Bottom row shows the distribution of value

predictors for each reward-predictive cue.

2175

$$P_i \leftarrow P_i + \varphi^+ \cdot |\delta_i| - \beta \cdot P_i \quad \text{if } \delta_i > 0$$
$$N_i \leftarrow N_i + \varphi^- \cdot |\delta_i| - \beta \cdot N_i \quad \text{if } \delta_i < 0$$
$$V_i = P_i - N_i$$

$$\delta_i = \alpha_i^+ \cdot (r - V_i) \quad \text{if } \delta_i > 0$$
$$\delta_i = \alpha_i^- \cdot (r - V_i) \quad \text{if } \delta_i < 0$$

2176

Extended Data Fig. 13 | Distributional reinforcement learning with D1 and D2 populations (Model 1).

a. Schematic of the distributional RL model with D1 and D2 populations. The schematic represents three different value predictors (pessimistic, neutral and optimistic from left to right) with their respective P and N neurons. The level of optimism of each individual value predictor is determined by the scaling factors of the individual dopamine RPE-evoked responses (α_i^+, α_i^-, represented by the color in the colormap from purple to pink) and allows the model to encode information about the distribution of rewards (bottom). The global level of 'optimism' or

'pessimism' of the agent is given by the re-scaling of the RPEs by the P and N receptors

sensitivities in the model (ϕ^+, ϕ^-, represented with the color saturation).

b. Example of a Bernoulli distribution, equivalent to the reward distribution predicted by the

50% cue.

c. Distribution of expectiles learnt by the distributional RL model with D1 and D2 population for

the reward distribution in b. The expectiles are sorted based on the asymmetric scaling factor of

each individual dopamine neuron. Colormap represents the level of optimistic or pessimism of

each agent.

d. Samples from the decoded distributions for the set of expectiles in c. The probability density is

bimodal, in accordance with the distribution in b. As the agents goes from pessimism to

optimism, the probability density modes change in their relative magnitude.

2196