Hierarchizing multi-scale environmental effects on agricultural pest population dynamics: a case study on the annual onset of *Bactrocera dorsalis* population growth in Senegalese orchards

Authors: Cécile Caumette[1,2,3], Paterne Diatta[4], Sylvain Piry[1], Marie-Pierre Chapuis[1,2], Emile Faye[3], Fabio Sigrist[5], Olivier Martin[6], Julien Papaïx[6], Thierry Brévault[7], Karine Berthier[8]

Addresses:

[1]CBGP, Montpellier SupAgro, INRAE, IRD, CIRAD, University of Montpellier, Montpellier, France

[2]CIRAD, CBGP, Montpellier, France

[3]CIRAD, UPR Hortsys, F-34398, Montpellier, France

[4]ISRA/CRA de Djibélor, Ziguinchor, Sénégal

[5]Seminar for Statistics, ETH Zurich, Switzerland

[6]INRAE, BioSP, 84914 Avignon, France

[7]CIRAD, UPR AIDA, F-34398, Montpellier, France.

[8]INRAE, Pathologie Végétale, F-84140, Montfavet, France

Corresponding author: cecile.caumette@cirad.fr

Abstract

Implementing integrated pest management programs to limit agricultural pest damage requires an understanding of the interactions between the environmental variability and population demographic processes. However, identifying key environmental drivers of spatiotemporal pest population dynamics remains challenging as numerous candidate factors can operate at a range of scales, from the field (e.g. agricultural practices) to the regional scale (e.g. weather variability). In such a context, data-driven approaches applied to pre-existing data may allow identifying patterns, correlations, and trends that may not be apparent through more restricted hypothesis-driven studies. The resulting insights can lead to the generation of novel hypotheses and inform future experimental work focusing on a limited and relevant set of environmental predictors. In this study, we developed an ecoinformatics approach to unravel the multi-scale environmental conditions that lead to the early re-infestation of mango orchards by a major pest in Senegal, the oriental fruit fly *Bactrocera dorsalis* (BD). We gathered abundance data from a three-year monitoring conducted in 69 mango orchards as well as environmental data (i.e. orchard management, landscape structure and weather variability) across a range of spatial scales. We then developed a flexible analysis pipeline centred on a recent machine learning algorithm (GPBoost), which allows the combination of gradient boosting and mixed-effects models or Gaussian processes, to hierarchize the effects of multi-scale environmental variables on the timing of annual BD population growth in orchards. We found that physical factors (temperature, humidity), and to some extent landscape features, were the main drivers of the spatio-temporal variability of the onset of population growth in orchards. These results suggest that favourable microclimate conditions could provide refuges for small BD populations that could survive, with little or no reproduction, during the mango off-season and, then, recolonize neighbouring orchards at the beginning of the next mango season. Confirmation of such a hypothesis could help to prioritize surveillance and preventive control actions in refuge areas.

Keywords: *Bactrocera dorsalis*, mango crop, weather, landscape, agricultural practices, GPBoost, population dynamics, abundance time series, ecoinformatics, machine learning

Introduction

Limiting pest damage is a major challenge for agriculture that has been mainly addressed through chemical and curative control methods, leading to socio-economic, environmental and human health issues (Deguine et al., 2021; Mutamiswa et al., 2021 ; Brévault & Clouvel 2019; Chaplin-Kramer et al., 2011). The need to develop sustainable forms of agriculture has led to the emergence of the concept of Integrated Pest Management (IPM), which aims to integrate a range of alternative pest control techniques (e.g. biological control, landscape manipulation, changes in cultural practices, use of resistant varieties). However, as highlighted by Deguine et al. (2021), one of the main recurrent issues in the implementation of IPM is the lack of careful consideration for the ecological processes occurring in agroecosystems. Indeed, demographic parameters of pest populations, although dependent on their intrinsic characteristics (e.g. dispersal and reproductive capacities), are also strongly dependent on numerous environmental factors that determine the spatiotemporal availability, accessibility and quality of resources, such as agricultural practices, host plant diversity and phenology, natural enemies, landscape structure and weather (Veres et al., 2013; Kennedy & Storer 2000). Understanding the key environmental drivers of spatiotemporal pest population dynamics remains then challenging, especially in agrosystems that are often highly labile through space and time, notably due to the diversity and phenology of crops and wild hosts as well as farming practices, and where different environmental features influence demographic processes across a range of spatial scales, from the field to regional scales or beyond (Brévault & Clouvel 2019; Kennedy & Storer 2000). Therefore, extensive sampling efforts are required to achieve both population monitoring and environmental data collection, at the relevant spatio-temporal scales and with the appropriate precision.

In this context, a valuable first step in investigating the ecological processes underlying pest population dynamics is to create a composite set of pre-existing data, which may have been collected through various research or management programs, in order to perform correlative statistical analyses. For example, stakeholders often record longitudinal data on pest abundance, crop yields and farming practices, in order to inform real-time pest management decisions (e.g. Rosenheim & Meisner 2013). Open access databases or repositories providing raw or pre-processed data on the variability of environmental features derived from remote sensing technologies or mathematical modelling are also increasingly available (e.g. landscape typologies, weather variables). Such a research framework, termed "ecoinformatics" since the era of big data (Rosenheim & Gratton 2017), can capture multi-year data over large spatial extents and under environmental conditions directly relevant to agriculture and management operations. For example, ecoinformatics research provided important insights on the dependencies between spatiotemporal population dynamics and environmental heterogeneity in several agricultural insect pests such as aphids (Stack Whitney et al., 2016), locusts (Veran et al., 2015) or plant bugs (Rosenheim & Meisner 2013). These studies also give the opportunity to inform future hypothesis-driven experimental research by (i) narrowing down a large number of candidate environmental variables to a limited set of variables relevant for pest population dynamics and

amenable to experimentation and (ii) formulating more focused hypotheses on causal relationships between environment and pest dynamics that can be further tested (Rosenheim et al., 2011; Kelling et al., 2009; Hochachka et al., 2007).

The main objective of the present study was to unravel the environmental conditions that may favour rapid seasonal re-infestation of mango orchards by the oriental fruit fly, *Bactrocera dorsalis* (BD), in Senegal. This invasive species, native from tropical Asia, has emerged as a major pest of mangoes and other tropical fruit crops in Africa in the early 2000 (Ekesi et al., 2006). Direct crop losses are caused by larval feeding in the fruit, but significant indirect losses occur when market access opportunities are lost due to quarantine regulations (Mutamiswa et al., 2021; Ekesi et al., 2011; Vayssières et al., 2008). In the Niayes area, one of the main mango production basins in Senegal, the annual variation in BD abundance is extremely marked, with a striking demographic bottleneck at the end of the mango season raising questions about how orchards get re-infested at the beginning of the next production season. A key factor could be the survival of small demes during the dry season that would constitute discreet sources to initiate local population growth and rapid re-infestation of orchards at the beginning of the production season. Many abiotic and biotic factors have been identified as potentially critical for the survival of BD during the mango off-season in Senegal, including temperature, precipitations, relative humidity, irrigation as well as the abundance, diversity and phenology of alternative host plants within and around orchards (Diallo et al., 2021; Dieng et al., 2019; Boinahadji et al., 2019; Konta et al., 2015; Vayssières et al., 2015; Diatta et al., 2013; Ndiaye et al., 2012; Ndiaye et al., 2008).

Here, we first built a composite dataset from a three-year survey of abundance data previously collected in 69 mango orchards in the Niayes region (Diatta 2016) and environmental data on a large number of candidate predictors at different spatial scales, including cropping systems (Diame et al., 2015; Grechi et al., 2013), landscape structure (Jolivot 2021), and weather variability (Karger et al., 2021; Didan 2015). We then used this dataset to assess the possible source (or sink) effects of environmental features, by investigating their relationship with the onset of annual population growth of *B. dorsalis*. There are specific analytical challenges related to the use of a large number of candidate environmental variables, such as heterogeneity in both their nature (i.e. qualitative and quantitative) and their relationship to pest population dynamics (e.g. nonlinearity), as well as multicollinearity (i.e. high correlations between two or more variables). Thus, we developed a flexible analysis pipeline centred on a recent machine learning algorithm, GPBoost (Sigrist 2022), to integrate multi-scale candidate environmental factors and hierarchize their effects. The results provided insights into the environmental conditions that may favour the rapid annual re-infestation of mango orchards at the beginning of the mango production season, which can also inform on the favourable conditions for BD survival during the dry season. Formulated hypotheses on causal relationships are discussed in relation to published experimental studies and confronted to competing interpretations, in particular the potential influence of any suspected confounding parameters that could not be included in the study. Deciphering the relative role of environmental features on the earliness of the re-

infestation process can help to prioritize future research, but also to adapt possible surveillance and preventive actions for BD control.

Materials & Methods

Figure 1 provides a schematic view of the analytical pipeline detailed in this section. All analyses were performed using the R Statistical Software, version ≥ 4.1.2 (R Core Team 2023).

Study area

The study area encompassed the "Niayes" (Fig. 2), which is the main region of vegetable and fruit production in Senegal (Grechi et al., 2013; De Bon et al., 1997). Mango is the main fruit production, grew either in intensive orchards dedicated to international export or in more traditional and diversified orchards for local markets (Ndiaye et al., 2012; Vayssières et al., 2011). The harvest season in the Niayes, which is subject to a Sahelian climate, is mainly from June to August and coincides with the rainy season (Grechi et al., 2013; Vayssières et al., 2011). Natural vegetation is relatively scarce and forms a landscape mosaic with cultivated lands and urban areas.

Estimation of the starting date of BD population growth within orchards

In this study, we used BD abundance data from Diatta (2016), who monitored 69 mango orchards located roughly among six sites in the Niayes region, between December 2011 and December 2014 (Fig. 2). Within each orchard, an average of three traps were placed in different trees (1 to 2m height). Traps were baited with methyl-eugenol, an attractive parapheromone for BD males, combined with an insecticide (DDVP: dichlorvos). The number of flies caught per trap was counted weekly in each orchard.

For each orchard and sampling date, the number of catches over traps was averaged in order to obtain abundance times series. Then, for each orchard and year, we estimated the starting date of the demographic growth of local BD populations using the POPFIT model (Soulsby et al., 2012; see details in Section 1 of the Supplementary Material). Originally developed for butterfly species, POPFIT can be applied to others insect species with similar annual population dynamics, as observed from BD abundance time series: a phase of zero or almost zero abundance followed by a phase of rapid population growth to a peak and then a decline to zero abundance again (Fig. 3). The initial hypotheses of the POPFIT model were relaxed as we were only interested in estimating the onset of the demographic growth phase (t_0) regardless of the underlying demographic processes (adult survival from one year to the next, local eclosions, migration or a combination of these processes). A mechanistic-statistical framework was used to perform the parameter inference for the POPFIT mechanistic model (Papaïx et al., 2022), conducted within a Bayesian framework

using Nimble (de Valpine et al., 2017) with the R package "nimble" v1.0.1. We visually checked the resulting fits and discarded orchards for which the model was poorly fitted to the data. Finally, for each orchard and year, 500 values of the estimated t_0 were sampled from the posterior distributions of the parameter in order to build 500 replicates of the demographic dataset.

Multi-scale environmental predictors

Within orchards. Between 2010 and 2013, the cropping system of 86 orchards, including the 69 orchards monitored for BD abundance that we analysed in the present study, was characterized using the same methodology (Diatta 2016; Diame et al., 2015; Grechi et al., 2013). Mango producers were interviewed about the management of their orchard, particularly on practices that might affect early BD population growth within orchards: irrigation, sanitation (i.e. removal of aborted mangoes that can host BD larvae) and intercropping with vegetable crops as potential alternative hosts for BD (Diouf et al., 2022; Vayssières et al., 2015; Grechi et al., 2013). These factors were considered as qualitative variables: irrigation and sanitation as factors with three levels (null, moderate or intensive irrigation and null, occasional or regular sanitation) and inter-crops as presence/absence. Host diversity and frequency, which may also influence BD re-infestation dynamics (Diallo et al., 2021; Boinahadji et al., 2019; Diatta 2016; Grechi et al., 2013; Ndiaye et al., 2012; Vayssières et al., 2011), were also estimated from a subset of around 100 fruit trees per orchard. Each selected tree was identified at the species level and for mango trees, the cultivar was also identified.

Here, agricultural practices (irrigation, sanitation, intercropping with vegetables) were kept as categorical variables. Due to the high number of fruit tree species and mango cultivars identified in the orchards and the bias in the representation of some categories (i.e. from only one sample for the rarest category to nearly 2000 for the most frequent category), we classified both the mango cultivars and the alternative host tree species according to their phenology, especially the period during which fruits are available and can potentially host BD larvae, based on data from literature and expert knowledge (from researchers and producers). For mango trees, we individualized the three main cultivars (Kent, Keitt, Boukodiekhal) and grouped the others in three phenological categories (early, medium, late). For other species, we grouped them into three categories according to their potential period of fruit availability: December to April (before the beginning of the mango season), April to November (during and after the mango season), and all year round. Then, for each orchard, we calculated the proportion of each phenological category by dividing the number of trees in a category by the total number of trees sampled in the orchard (details in Supplementary Material Section 2, Tab. S2.1).

Landscape. To quantify the effect of the landscape features surrounding the monitored orchards on the onset of BD population growth, we used a pre-existing typology of 13 classes of land use (Fig. 2) built using 2010-SPOT6 satellite images and time series of 2018-Sentinel 2 satellite images (Jolivot 2021). The effect of

landscape features surrounding plots monitored to record population abundance is often investigated using nested circular buffers or rings of increasing radius, but such approaches have drawbacks such as the high level of correlation of landscape features across the different radii considered and the rather unrealistic assumption that landscape effects are uniform within a given buffer and null outside (Carpentier & Martin 2021; Chandler & Hepinstall-Cymerman 2016). In this work, we used the recent Siland method (Carpentier & Martin 2021), which allows to estimate the spatial scale of influence of a landscape feature on a response variable (here t_0), without any a priori of distance. The method also allows to consider local explanatory variables: here we included the sampling year and site. The spatial influence function (SIF), which models the decreasing influence with distance from the observation points of the landscape features, was a Gaussian function with a mean distance δ, estimated with Siland independently for each of the 13 land use classes. Based on the estimated value of δ, Siland provided the cumulative influence of each land use class at each observation point, i.e. the land contribution, hereafter denoted as *lc*.

We first carried out a sequential tuning step of the Siland hyperparameters: the raster resolution *wd* and the initialisation value for the maximum likelihood optimization procedure *init* (see details in Section 3 of the Supplementary Material). Then, *lc* values were computed independently for each of the 500 replicates of the dataset. Analyses were performed using the R package "siland" v.3.0.2.

Regional weather variability. The spatio-temporal variability of physical factors was analysed over the study area and the period of sampling using two data sources. First, we used raster data of monthly minimal, maximal and mean temperatures (Tmin, Tmax, Tmean), as well as precipitations, obtained from the *CHELSA* model v 2.1 (Karger et al., 2021; Karger et al., 2017) with a spatial resolution of 30 arc-seconds (approximately 1 km). Second, we used rasters of bi-monthly Normalized Difference Water Index (NDWI), an indicator for vegetation water content (Gu et al., 2007; Gao 1996) that we calculated using MODIS/Terra Vegetation Indices 16-Day L3 Global 250m SIN Grid data (Didan 2015; Didan et al., 2015) according to the formula of NDWI$_{2130}$ defined in Chen et al. (2005). NDWI data were averaged monthly and combined with *CHELSA* data to obtain monthly raster time series of 1 km² spatial resolution covering the entire study area. We only kept the rasters for the time period between December and May, which covers the low demographic phase preceding population growth for each sampling year (i.e. Dec 2011–May 2012, Dec 2012–May 2013 and Dec 2013–May 2014, thereafter named 2012, 2013 and 2014 for the sake of simplicity). The final dataset was composed of 30 variables (6 months x 5 variables: Tmin, Tmax, Tmean, precipitations and NDWI), computed for each of the 2863 cells (1 km²) of the spatial raster and for each year; i.e. a matrix of size 30 x 8589. Data were normalized and reduced in a smaller dimensional space using a principal component analysis (PCA). For each monitored orchard and each of the main principal components (PC), selected using the Broken Stick model (MacArthur 1957), we extracted the PC scores of the grid cell corresponding to the spatial location of the orchard in the study area (three values, one per year). The analyses were performed using the R packages "ade4" v. 1.7-19 (Dray & Dufour 2007) and "PCDimension" v. 1.1.13 (Wang et al., 2018).

Effect of multi-scale environmental factors on the onset of local BD population growth

To hierarchize the effects of the multi-scale environmental predictors on the onset of demographic growth of BD populations in orchards (t_0), we used GPBoost, a recent machine learning method that combines gradient boosting and mixed-effects models or Gaussian processes, allowing to consider spatial and temporal dependencies between observations (Sigrist 2022). The GPBoost regression model included the sampling site (S1 to S6) and year (2012, 2013 and 2014) as grouped random effects and all environmental candidate factors as fixed effects: i) the 12 orchard management variables (i.e. irrigation, sanitation, vegetable crops, and phenological groups of mango trees and alternative hosts), ii) the contributions (lc) of the 13 land use classes estimated independently with Siland for each of the 500 replicates and, iii) the scores for each orchard on the retained PCs of the PCA conducted on the physical variables (i.e. minimum, maximum and mean temperatures, precipitations and NDWI) (see Supplementary Material Section 4 for a more detailed description of these variables). Analyses were conducted using the R package "gpboost" v 1.2.3.

GPBoost model hyperparameters (the learning rate, the minimum data number in tree leaves, the maximal depth of trees and the number of trees) were tuned independently on the 500 replicates of the dataset using the grid search procedure implemented in the package "gpboost" and based on a 4-fold cross-validation (see details in Section 5 of the Supplementary Material). For the number of trees, it was automatically optimized by setting the maximum number of iterations to 2000 and the *early_stopping_rounds* parameter to 5, i.e. the process stops if the model's performance on the validation set does not improve for five consecutive iterations. The model was then trained independently for each of the 500 replicates of the dataset with the '*gpboost*' function using the best combination of hyperparameter values identified for each replicate in the tuning step. For each replicate, the predictors were then hierarchized according to their importance in the model, expressed by the SHapley Additive exPlanation (SHAP) values (Lundberg et al., 2018; Lundberg & Lee 2017). SHAP values provide the contribution of each predictor on the predicted values of individual observations. The overall contribution of a given predictor to the model output is obtained by averaging the absolute SHAP values of the observations (hereafter called S_{mean}). For each of the 500 replicates of the dataset, SHAP values were computed using the R package "SHAPforxgboost" v.0.1.1 and the predictors were ranked in decreasing order of the median of the S_{mean} values over all replicates. Based on this ranking, the relationship of the most important predictors with the estimated t_0 was investigated using SHAP dependence plots (i.e. smooth of the individual SHAP values as a function of the predictor values).

Finally, as a validation step, we assessed the predictive performance of the GPBoost model considering either all predictors or the top ranked predictors. In all cases, each of the 500 replicates was partitioned to build training and test datasets (80% and 20% of the data, respectively). Partitioning was done in a way to ensure that all groups for grouped random effects (i.e. year and site) were represented at least

once in the training dataset. For each replicate, the hyperparameters were tuned and the GPBoost model was trained on the training dataset, as previously detailed (see Section 5 of the Supplementary Material). Then, the resulting model was used to predict t_0 from the corresponding test dataset. Model accuracy was assessed by computing the Pearson correlation coefficient between predicted and observed values of t_0 as well as the Root Mean Square Error (RMSE) of the model.

RESULTS

As described in Diatta (2016), the BD abundance time series showed an annual demographic kinetics consistent with the use of the POPFIT model (Fig. 3). Based on visual control of the quality of the POPFIT model fit to the data, we discarded four orchards from the original dataset. The variability of the 500 values of t_0 resampled from the posterior distributions for each of the 65 remaining orchards and for each year is presented Figure 4. On average, for one orchard and year, the difference between the maximal and minimal values of the 500 estimates of t_0 resampled in the posterior distributions was 0.7 weeks (Fig. 4). Since the temporal resolution of the abundance time series is the week, this indicates the good accuracy of the t_0 estimates with the POPFIT framework.

Results of the GPBoost model applied to the 500 replicates of the dataset showed that the median and range values for the error term and grouped random effects (i.e. *year* and *site*), were 0.4 [0; 1.72], 1.86 [0.72; 3.03] and 0.14 [0; 1.64] weeks, respectively. A significant proportion of the variance in t_0 is then expected to be explained by fixed effects. Indeed, within-year variation of t_0 between orchards was substantial, with a difference between the earliest and the latest orchard (i.e. the difference in the median of the within-orchard t_0 values over the 500 replicates) of 12, 17 and 15 weeks in 2012, 2013 and 2014, respectively. For a same orchard, the variation in the median of the t_0 values (over the 500 replicates) between years was, on average over all orchards and years, 3.4 weeks. The results of the SHAP-based ranking over the 500 replicates for the 28 environmental predictors specified as fixed effects in the GPBoost model are presented Figure 5. From this ranking, two groups of predictors stand out as the most meaningful to explain t_0. The first group included the three first principal components (predictors PC3, PC2 and PC1) retained by the broken stick method, that explained 77.8% of the total variance in the PCA performed on the physical variables (i.e. temperatures, precipitations and NDWI; Fig. 6A), as well as the land use class LU13 (urban area). The second group of variable included two additional land use classes, LU7 (shrub savannah) and LU11 (sparsely vegetated ground) as well as an agronomic feature of the studied orchards, i.e. the proportion of potential alternative hosts producing fruits between April and November (AH3).

The result of the PCA indicated that PC3, the top-ranked predictor in the GPboost model over the 500 replicates, was negatively correlated with the vegetation water content index (NDWI) from December to May (Fig. 6A). PC3 predictor mainly reflected fine spatial variation of NDWI values over these months (Fig. 6B). The smoothed SHAP-dependence plot (Fig. 7) showed a positive relationship between PC3 values and the

individual SHAP values (i.e. negative SHAP values for the lowest PC3 values and positive SHAP values for highest PC3 values), meaning that earlier t_0 were associated with lower values of PC3 and then with higher values of NDWI. The second-best predictor, PC2 (Fig. 5), was a temporal dimension which contrasted humidity conditions in the PCA: positive values were correlated with precipitations occurring between February and April and with higher NDWI values between December and May, while negative values were correlated with precipitations in December, January and May (Fig 6A). The former conditions were mainly observed in 2012 and, to a lesser extent, in 2013 while the later mostly corresponded to the year 2014 (Fig 6B). Although the smoothed SHAP-dependence plot for the PC2 predictor showed a sawtooth-kind of relationship (Fig. 7), there was a clear decreasing trend indicating that negative individual SHAP values roughly corresponded to positive PC2 values. This result suggests that earlier t_0 are expected when humidity conditions (precipitations and NDWI) are higher between February and April. The first component of the PCA, i.e. predictor PC1, showed a well-marked spatial gradient in monthly temperature ranges, from positive values in the coastal area, associated with the highest minimum temperatures and the lowest maximum temperatures, to negative values in the inland, characterized by the highest maximum temperatures and lowest minimum temperatures (Fig. 6). Smoothed SHAP-values associated with PC1 exhibited a U-shaped curve (Fig. 7), suggesting that earliest t_0 are associated with intermediate conditions in terms of minimum and maximum monthly temperatures, as observed in the central part of the study area (Fig. 6B). Three other top predictors identified from the ranking of the GPBoost results over the 500 replicates corresponded to landscape classes (Fig. 5), expressed in terms of cumulative influence (*lc*) on each estimate of t_0 as computed with the Siland method (see Tab. S3.1 in Section 3 of the Supplementary Material for a detailed description of Siland results). The smoothed relationships between these landscape predictors and the individual SHAP values (Fig. 7) roughly approximated either: an L-shaped or inverted L-shaped curve for urban area (LU13) and sparsely vegetated ground (LU11) respectively, and a S-shaped curve for shrub savannah (LU7). These relationships suggest that the presence of urban areas (LU13) in the orchard's surrounding is associated with early population growth (i.e. smallest t_0 values). On the contrary, the presence of sparsely vegetated ground (LU11) and shrub savannah (LU7) tend to delay the onset of the demographic growth of BD populations within orchards (i.e. highest t_0 values). Finally, the smoothed SHAP curve for the class of potential alternative hosts AH3 also approximated an inverted L-shaped relationship suggesting that the higher the proportion of these species, the later the onset of BD population growth within orchards.

Lastly, as a validation step we assessed the predictive performance of the GPBoost model considering either all predictors, the seven top ranked predictors (i.e. PC3, PC2, PC1, LU13, LU11, LU7,AH3) or only the four top ranked predictors (i.e. PC3, PC2, PC1 and LU13). The RMSE and Pearson correlation coefficient, averaged over the 500 replicates, were 2.24 and 0.77, respectively, for the model including all predictors, 2.23 and 0.78 for the model including seven predictors and 2.23 and 0.77 for the model including the four top predictors (see details in Section 6 of the Supplementary Material, Fig. S6.1).

DISCUSSION

In this work, we developed a flexible analysis pipeline to hierarchize the effects of multi-scale candidate environmental factors on estimated parameters of pest population dynamics (Fig. 1). At the heart of this pipeline is a recent machine learning method, GPBoost, which allows gradient boosting to be combined with Gaussian process and/or mixed effects models (Sigrist 2022). The method inherently benefits from the advantages of gradient-boosted trees (e.g. handling of nonlinearities, discontinuities, higher order interactions, outliers, multicollinearity between predictors and missing data (Elith et al., 2008)), while allowing to relax the zero prior mean or linearity assumption of Gaussian process and mixed effects models (Sigrist 2022). The possibility to consider grouped random effects, as done in the present study, also provides a unique way to account for the non-independence of the response variable across observations, which is overlooked in most machine learning algorithms. This pipeline allowed us to integrate multi-sources pre-existing data to hierarchize the effects of 28 environmental predictors, assessed from the local to the regional scale, on the annual onset of local population growth of *Bactrocera dorsalis* (t_0), a major invasive pest of the mango crop in Senegal.

Given that the two best environmental predictors were the third and second principal components of the PCA carried out on physical variables, our results clearly suggest that humidity conditions are the primary driver of the spatiotemporal variation in the earliness of local population growth in mango orchards of the Niayes region (i.e. up to 17 weeks of delay between the earliest and latest onset in a year). This result is in line with previous studies reporting humidity as a key component of *B. dorsalis* population dynamics (e.g. Ibrahim et al., 2022; Chuang et al., 2014; Vayssières et al., 2009). The relationship between the demographic estimate t_0 and PC2 indicated that even a very small amount of precipitation (most correlated variable with PC2) occurring between February and April (episodic rainfall events called the "mango rain" in Senegal (Wade et al., 2015)) could be involved in creating favourable microhabitats leading to early development of BD populations in orchards. PC3, that captured both spatial and temporal fine variations during the dry season of the Normalized Difference Water Index (NDWI), index which depends at least partly on precipitations and soil moisture, showed a positive correlation with early population growth. These results are consistent with several experimental studies indicating that humidity is a critical factor for BD survival, especially at the pupal stages. Indeed, the survival of pupae (and so the emergence rate) is significantly affected by soil moisture, which is strongly related to precipitations, with optimal trait values at 10-60% moisture levels (Hou et al., 2006). Desiccation is also an important cause of mortality of third-instar larvae under different climatic conditions (Jackson et al., 1998; Serit & Tan 1990). Furthermore, from observational data in Penang, Malaysia, Serit & Tan (1990) found that the main factors of mortality of *B. dorsalis* immature stages was desiccation or drowning of larvae and pupae in soil (77.8% of mortality for soil-associated immatures). In this way, BD larvae's preference for pupating in shaded areas has already been mentioned (Susanto et al., 2022).

Importantly, the vegetation water content (NDWI) also reflects the vegetation fraction cover (i.e. importance of the canopy). Thus, higher NDWI values may also reflect favourable microclimate conditions for BD adults, with higher moisture levels and higher shading effects that will contribute to moderate temperature variations. Besides, the first component of the physical PCA, which showed a gradient in minimum and maximum monthly temperatures, was also ranked in the top environmental predictors with intermediate conditions of temperature associated with the earliest population growth values. This result would be consistent with previous experimental studies describing performance curves for temperature-dependent development, survival and fecundity traits in *B. dorsalis*. For example, temperatures for optimal immature development ranged around 25-30°C, with development time (or mortality) increasing at lower (or higher) temperatures, preventing from any adult emergence above 35°C (and below 9-10°C) (Dongmo et al., 2021; Rwomushana et al., 2008; Vargas et al., 1996). In addition, adult longevity decreases with increasing temperature, and females can only lay eggs between 15 and 35°C, with the optimal conditions for a higher number of eggs being between 20 and 25°C (Dongmo et al., 2021; Choi et al., 2020; Vargas et al., 1997; Yang et al., 1994). In our study area, the most favourable temperature range for early population development in orchards lies between oceanic conditions in the coastal part and inland conditions where the maximum daily temperature easily exceeds 35°C during the dry season. As temperatures above 35°C challenge all components of BD life history, spatial and inter-annual weather variability in the Niayes region is likely to interact with local factors providing higher levels of humidity and shading (e.g. water bodies and groundwater, vegetation and soil moisture, canopy structure) to create favourable microhabitats allowing BD to mitigate hydric and thermal stress during the dry season (Inskeep et al., 2021; Mutamiswa et al., 2021).

In addition to physical factors, the boosting approach also identified three landscape features that influenced the timing of the annual re-infestation of mango orchards. First, urbanized areas (LU13) would act as a catalyst for early re-infestation. This phenomenon could be attributed to the high frequency of peri-urban farming or even micro-gardens within towns. Intensive irrigation practices and the presence of alternative host plants, like citrus, during the dry season in these areas (Vayssières et al., 2011) might offer conditions favourable to BD survival and reproduction. Another non-exclusive plausible explanation relies on the potential contribution to local population dynamics of flies originating from imported mangoes in the region (Louzeiro et al., 2021; Hong et al., 2015). Before the start of the production season in the Niayes, mangoes are massively imported from southward production basins (e.g. Guinea, Southern Mali and Casamance) to supply the local markets including those in the study area. The arrival of potentially infested mangoes could contribute to the early establishment of a pool of individuals, which may then lead to the rapid re-infestation of nearby orchards. Second, the presence of sparsely vegetated ground (LU11) and shrub savannah (LU7) would conversely delay the onset of local population growth. These landscape classes may be unsuitable habitats for BD due to the absence of host plants and the very low relative humidity, soil moisture and shading. Re-infestation of orchards surrounded by these types of habitats might strongly rely

on BD dispersal from favourable refuges, a process that may be limited under dry conditions, leading to delays in re-infestation.

Although considered to be significant factors impacting BD population dynamics, we did not identify any clear effect suggesting that orchard management could determine the timing of the clear change in BD abundance, either in terms of agricultural practices (irrigation, sanitation and presence of vegetable crops) or host diversity (mango varieties and alternative hosts) and phenology. The only potential effect found was that an increasing proportion of the alternative host class AH3, which produce fruits mainly during the mango season (April to November), would delay the onset of BD population growth within orchards, which may reflect the strong preference of BD for mango. In contrast to previous studies that have investigated BD population dynamics strictly in terms of abundance variation, we specifically focused on the onset of local population growth. Thus, our results suggest that while orchard management may explain differences in abundance, it plays a far less important role in initiating the re-infestation process compared to physical and landscape features. However, one methodological point worth noticing is that the orchard management data we used in our analysis was only available for monitored orchards (no information about practices in the surroundings) and some were coded as categorical variables with a few levels, which may not allow us to properly capture the underlying relationships between categories. Such categorical variables may also be much less informative than continuous predictors to find optimal split points for decision trees in the gradient boosting procedure.

Mango is the preferred cultivated host fruit of BD (Motswagole et al., 2019; Vayssières et al., 2009; Ekesi et al., 2006; Drew et al., 2005), but the species is known to be highly polyphagous (Ndiaye 2009; Ekesi & Billah 2006; Clarke et al., 2005; Allwood et al., 1999), which has led to the assumption that its maintenance during the dry season relies on the presence of alternative hosts to ensure continuous reproduction and larval development (Diallo et al., 2021; Faye et al., 2021; Boinahadji et al., 2019; Diatta 2016; Vayssières et al., 2015; Ndiaye et al., 2012). However, although 34 species of host fruit trees have been reported in the arid to semi-arid environment of the Niayes region (Ndiaye et al., 2012), their availability during the dry season remains relatively erratic, with the exception of cultivated *Citrus spp.*. Thus, one possible explanation for the lack of evidence for a role of alternative hosts in the earliness of orchard re-infestation is that the underlying process involved in BD maintenance during the mango off-season in the Niayes region may not be a continuous reproduction. This explanation is supported by the good performance of the GPBoost model in predicting the onset of the BD population growth in orchards based only on a few environmental factors representing physical and landscape variables. Active dispersal and dormancy are alternative ways of coping with stress during the unfavourable season. Experimental studies have shown that the dispersal of *B. dorsalis* adults is quite spatially restricted, generally upwind and occurs mostly when resources are scarce and temperatures exceed 20-24°C (Chailleux et al., 2021; Makumbe et al., 2020; Froerer et al., 2010). Dormancy is the interruption or reduction of metabolic and developmental activity in an immediate (quiescence) or pre-programmed (diapause) response to unfavourable conditions. While pupal dormancy is a common aridity

survival strategy in Dipteran species (Thorat & Nath 2018), it has not been demonstrated in a *B. dorsalis* desiccation experiment (Hou et al., 2006). Clarke et al. (2022) argue that phenological data on tropical *Bactrocera spp.* strongly suggest an adult reproductive arrest that would allow life span to be extended during the unfavourable dry season when fruits are scarce. This hypothesis of an adaptation to survive desiccation during the dry season fits well with our main findings that habitat characteristics underlying early population growth rates are those that provide milder temperatures and higher humidity and shading.

Our results indirectly provide valuable insights into the spatial and temporal conditions that may lead to the emergence of microhabitats favourable for BD survival during the dry season in the Niayes. Altogether, our results support the hypothesis of localised refuges, with more favourable conditions of temperature (moderate), humidity (high) and shade (high), where small BD populations could survive and re-infest orchards at the beginning of the mango season. If confirmed in future experimental and observational studies, such an information could ultimately be a key step for the design of surveillance programs and preventive control measures. Considering the large delay between the earliest and the latest onset of population growth found in this study (between 12 and 17 weeks depending on the year), locating areas with such favourable environmental conditions could allow preventive control measures to be taken during the dry season to limit the sources of orchard re-infestation. Our results also suggest that agricultural practices, such as tree pruning to reduce the vegetation cover, should be fully considered in IPM implemented to control BD.

It should be noted that the data used in this study have some limitations. First, the data describing the cropping system (i.e. host plant diversity and agricultural practices) may not be precise enough to detect significant effects in statistical analyses. Second, pre-existing demographic data are limited to the abundance of BD in orchards clustered in six sites, which does not allow to investigate whether the variability in the timing of re-infestation also depends on BD dispersal processes in interaction with the environmental matrix. Further research is therefore needed, based on better spatial coverage and finer data collection, to assess environmental heterogeneity (from the orchard to the regional scale) and its impact on the spatio-temporal dynamics of BD. Moreover, the acquisition of longitudinal demographic data would be a crucial advance, allowing the estimation of spatio-temporal interactions between variations in effective densities, population sizes and dispersal patterns. Beyond local processes, we also found elements suggesting potential effects of long-distance passive dispersal due to mango trade, which should be considered in future research.

REFERENCES

Allwood AJ, Chinajariyawong A, Drew RAI, Hamacek EL, Hancock DL, Hengsawad C, Jipanin JC, Jirasurat M, Kong Krong C, Kritsaneepaiboon S, Leong CTS, Vijaysegaran S (1999) Host plant records for fruit flies (Diptera: Tephritidae) in Southeast Asia. Raffles Bulletin of Zoology, **47**, 1-92.

Boinahadji AK, Coly EV, Dieng EO, Diome T, Sembene PM (2019) Interactions between the oriental fruit fly Bactrocera dorsalis (Diptera, Tephritidae) and its host plants range in the Niayes area in Senegal. *Journal of Entomology and Zoology Studies*, **7**, 855-864.

Brévault T, Clouvel P (2019) Pest management: Reconciling farming practices and natural regulations. *Crop Protection*, **115**, 1–6.https://doi.org/10.1016/j.cropro.2018.09.003

Carpentier F, Martin O (2021) Siland a R package for estimating the spatial influence of landscape. *Scientific Reports*, **11**, 7488. https://doi.org/10.1038/s41598-021-86900-0

Chailleux A, Thiao DS, Diop S, Bouvery F, Ahmad S, Caceres-Barrios C, Faye E, Brévault T, Diatta P (2021) Understanding *Bactrocera dorsalis* trapping to calibrate area-wide management. *Journal of Applied Entomology*, **145**, 831–840. https://doi.org/10.1111/jen.12897

Chandler R, Hepinstall-Cymerman J (2016) Estimating the spatial scales of landscape effects on abundance. *Landscape Ecology*, **31**, 1383–1394. https://doi.org/10.1007/s10980-016-0380-z

Chaplin-Kramer R, O'Rourke ME, Blitzer EJ, Kremen C (2011) A meta-analysis of crop pest and natural enemy response to landscape complexity: Pest and natural enemy response to landscape complexity. *Ecology Letters*, **14**, 922–932. https://doi.org/10.1111/j.1461-0248.2011.01642.x

Chen D, Huang J, Jackson TJ (2005) Vegetation water content estimation for corn and soybeans using spectral indices derived from MODIS near- and short-wave infrared bands. *Remote Sensing of Environment*, **98**, 225–236. https://doi.org/10.1016/j.rse.2005.07.008

Choi KS, Samayoa AC, Hwang S-Y, Huang Y-B, Ahn JJ (2020) Thermal effect on the fecundity and longevity of *Bactrocera dorsalis* adults and their improved oviposition model. *PLoS ONE*, **15**, e0235910. https://doi.org/10.1371/journal.pone.0235910

Chuang C-L, Yang E-C, Tseng C-L, Chen C-P, Lien G-S, Jiang J-A (2014) Toward anticipating pest responses to fruit farms: Revealing factors influencing the population dynamics of the Oriental Fruit Fly via automatic field monitoring. *Computers and Electronics in Agriculture*, **109**, 148–161. https://doi.org/10.1016/j.compag.2014.09.018

Clarke AR, Armstrong KF, Carmichael AE, Milne JR, Raghu S, Roderick GK, Yeates DK (2005) Invasive Phytophagous Pests Arising Through A Recent Tropical Evolutionary Radiation: The *Bactrocera dorsalis* Complex of Fruit Flies. *Annual Review of Entomology*, **50**, 293–319. https://doi.org/10.1146/annurev.ento.50.071803.130428

Clarke AR, Leach P, Measham PF (2022) The Fallacy of Year-Round Breeding in Polyphagous Tropical Fruit Flies (Diptera: Tephritidae): Evidence for a Seasonal Reproductive Arrestment in *Bactrocera* Species. *Insects*, **13**, 882. https://doi.org/10.3390/insects13100882

De Bon H, Faye F, Pages J (1997) Development of vegetable cropping systems in the Niayes zone of Senegal. *Experimental Agriculture*, **33**, 83–90. https://doi.org/10.1017/S0014479797000197

Deguine J-P, Aubertot J-N, Flor RJ, Lescourret F, Wyckhuys KAG, Ratnadass A (2021) Integrated pest management: good intentions, hard realities. A review. *Agronomy for Sustainable Development*, **41**, 38. https://doi.org/10.1007/s13593-021-00689-w

de Valpine P, Turek D, Paciorek CJ, Anderson-Bergman C, Lang DT, Bodik R (2017) Programming With Models: Writing Statistical Algorithms for General Model Structures With NIMBLE. *Journal of Computational and Graphical Statistics*, **26**, 403–413. https://doi.org/10.1080/10618600.2016.1172487

Diallo I, Ndiaye A, Faye E, Faye M, Sembene M (2021) Does landscape factors drive the genetic diversity and structure of populations of fruit flies, Senegal? Exploratory study of the case of *Bactrocera dorsalis* (Hendel, 1912) in the Niayes, Senegal. *Journal of Entomology and Zoology Studies*, **9**, 47–52. https://doi.org/10.22271/j.ento.2021.v9.i6a.8883

Diame L, Blatrix R, Grechi I, Rey J-Y, Sane CAB, Vayssières J-F, De Bon H, Diarra K (2015) Relations between the design and management of Senegalese orchards and ant diversity and community composition. *Agriculture, Ecosystems & Environment*, **212**, 94–105. https://doi.org/10.1016/j.agee.2015.07.004

Diatta P (2016) Relations entre la typologie des agro-écosystèmes fruitiers et les fluctuations de la population de la mouche des fruits : Bactrocera dorsalis (Hendel) dans les régions de Dakar et Thiès (Sénégal). *PhD Thesis, Cheikh Anta Diop University*.

Diatta P, Rey J-Y, Vayssières J-F, Diarra K, Coly EV, Lechaudel M, Grechi I, Ndiaye S, Ndiaye O (2013) Fruit phenology of citruses, mangoes and papayas influences egg-laying preferences of *Bactrocera invadens* (Diptera: Tephritidae). *Fruits*, **68**, 507–516. https://doi.org/10.1051/fruits/2013093

Didan K (2015) MOD13Q1 MODIS/Terra vegetation indices 16-day L3 global 250m SIN grid V006 [Data set]. *NASA eosdis land processes daac*, **10**, 415. https://doi.org/10.5067/MODIS/MOD13Q1.006

Didan K, Munoz AB, Solano R, Huete A (2015) MODIS vegetation index user's guide (MOD13 series). *University of Arizona: Vegetation Index and Phenology Lab*, **35**, 2–33.

Dieng EO, Ndiaye S, Faye PD, Balayara A, Badji K, Sembene PM (2019) New inventory of the diversity and seasonal abundance of Tephritid fruit fly species on mango orchards in Senegal. *Journal of Entomology and Zoology Studies*, **7**, 975-986.

Diouf EG, Brévault T, Ndiaye S, Faye E, Chailleux A, Diatta P, Piou C (2022) An agent-based model to simulate the boosted Sterile Insect Technique for fruit fly management. *Ecological Modelling*, **468**, 109951. https://doi.org/10.1016/j.ecolmodel.2022.109951

Dongmo MAK, Fiaboe KKM, Kekeunou S, Nanga SN, Kuate AF, Tonnang HEZ, Gnanvossou D, Hanna R (2021) Temperature-based phenology model to predict the development, survival, and reproduction of the

oriental fruit fly *Bactrocera dorsalis*. *Journal of Thermal Biology*, **97**, 102877. https://doi.org/10.1016/j.jtherbio.2021.102877

Drew RAI, Tsuruta K, White IM (2005) A new species of pest fruit fly (Diptera: Tephritidae: Dacinae) from Sri Lanka and Africa. *African entomology*, **13**, 149-154.

Dray S, Dufour A-B (2007) The ade4 Package: Implementing the Duality Diagram for Ecologists. *Journal of Statistical Software*, **22**, 1–20. https://doi.org/10.18637/jss.v022.i04

Ekesi S, Billah MK (2006) A Field Guide to the Management of Economically Important Tephritid Fruit Flies in Africa. 2nd edition. *Nairobi, Kenya: icipe Science Press.*

Ekesi S, Chabi-Olaye A, Subramanian S, Borgemeister C (2011) Horticultural pest management and the African economy: successes, challenges and opportunities in a changing global environment. *Acta Horticulturae*, **911**, 165-183. https://doi.org/10.17660/ActaHortic.2011.911.17

Ekesi S, Nderitu PW, Rwomushana I (2006) Field infestation, life history and demographic parameters of the fruit fly *Bactrocera invadens* (Diptera: Tephritidae) in Africa. *Bulletin of Entomological Research*, **96**, 379-386. https://doi.org/10.1079/BER2006442

Elith J, Leathwick JR, Hastie T (2008) A working guide to boosted regression trees. *Journal of animal ecology*, **77**, 802–813. https://doi.org/10.1111/j.1365-2656.2008.01390.x

Faye M, Ndiaye A, Diallo I, Sembene PM (2021) Genetic identification of populations of *Bactrocera dorsalis* (Diptera Tephritidae) in the Niayes and Lower Casamance areas in Senegal. *Journal of Applied Biosciences*, **158**, 16351–16362. https://doi.org/10.35759/JABs.158.9

Froerer KM, Peck SL, McQuate GT, Vargas RI, Jang EB, McInnis DO (2010) Long-Distance Movement of *Bactrocera dorsalis* (Diptera: Tephritidae) in Puna, Hawaii: How far can they go? *American Entomologist*, **56**, 88–95. https://doi.org/10.1093/ae/56.2.88

Gao B-C (1996) NDWI—A normalized difference water index for remote sensing of vegetation liquid water from space. *Remote Sensing of Environment*, **58**, 257–266. https://doi.org/10.1016/S0034-4257(96)00067-3

Grechi I, Sane CAB, Diame L, De Bon H, Benneveau A, Michels T, Huguenin V, Malezieux E, Diarra K, Rey J-Y (2013) Mango-based orchards in Senegal: diversity of design and management patterns. *Fruits*, **68**, 447–466. https://doi.org/10.1051/fruits/2013094

Gu Y, Brown JF, Verdin JP, Wardlow B (2007) A five-year analysis of MODIS NDVI and NDWI for grassland drought assessment over the central Great Plains of the United States. *Geophysical research letters*, **34**, https://doi.org/10.1029/2006GL029127

Hochachka WM, Caruana R, Fink D, Munson A, Riedewald M, Sorokina D, Kelling S (2007) Data-Mining Discovery of Pattern and Process in Ecological Systems. *Journal of Wildlife Management*, **71**, 2427-2437. https://doi.org/10.2193/2006-503

Hong SC, Magarey R, Borchert DM, Vargas RI, Souder S (2015) Site-specific temporal and spatial validation of a generic plant pest forecast system with observations of *Bactrocera dorsalis* (oriental fruit fly). *NeoBiota*, **27**, 37–67. https://doi.org/10.3897/neobiota.27.5177

Hou B, Xie Q, Zhang R (2006) Depth of pupation and survival of the Oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae) pupae at selected soil moistures. *Applied Entomology and Zoology*, **41**, 515–520. https://doi.org/10.1303/aez.2006.515

Ibrahim EA, Salifu D, Mwalili S, Dubois T, Collins R, Tonnang HEZ (2022) An expert system for insect pest population dynamics prediction. *Computers and Electronics in Agriculture*, **198**, 107124. https://doi.org/10.1016/j.compag.2022.107124

Inskeep JR, Allen AP, Taylor PW, Rempoulakis P, Weldon CW (2021) Canopy distribution and microclimate preferences of sterile and wild Queensland fruit flies. *Scientific Reports*, **11**, 13010. https://doi.org/10.1038/s41598-021-92218-8

Jackson CG, Long JP, Klungness LM (1998) Depth of Pupation in Four Species of Fruit Flies (Diptera: Tephritidae) in Sand With and Without Moisture. *Journal of Economic Entomology*, **91**, 138–142. https://doi.org/10.1093/jee/91.1.138

Jolivot A (2021) Cartographie de l'occupation du sol de la zone des Niayes (Sénégal) en 2018 (1.5 m de résolution) V1 [Data set]. *CIRAD Dataverse*. https://doi.org/10.18167/DVN1/KJAS6S

Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP, Kessler M (2017) Climatologies at high resolution for the earth's land surface areas. *Scientific Data*, **4**, 170122. https://doi.org/10.1038/sdata.2017.122

Karger DN, Conrad O, Böhner J, Kawohl T, Kreft H, Soria-Auza RW, Zimmermann NE, Linder HP, Kessler M (2021) Climatologies at high resolution for the earth's land surface areas. *EnviDat*. https://doi.org/10.16904/envidat.228.

Kennedy GG, Storer NP (2000) Life Systems of Polyphagous Arthropod Pests in Temporally Unstable Cropping Systems. *Annual Review of Entomology*, **45**, 467–493. https://doi.org/10.1146/annurev.ento.45.1.467

Kelling S, Hochachka WM, Fink D, Riedewald M, Caruana R, Ballard G, Hooker G (2009) Data-intensive Science: A New Paradigm for Biodiversity Studies. *BioScience*, **59**, 613–620. https://doi.org/10.1525/bio.2009.59.7.12

Konta IS, Djiba S, Sane S, Diassi L, Ndiaye AB, Noba K (2015) Etude de la dynamique de *Bactrocera dorsalis* (Hendel) (Diptera: Tephritidae) dans les vergers de mangues en Basse Casamance : influence des facteurs climatiques. *International Journal of Biological and Chemical Sciences*, **9**, 2698–2715. https://doi.org/10.4314/ijbcs.v9i6.15

Louzeiro LRF, Souza-Filho MFD, Raga A, Gisloti LJ (2021) Incidence of frugivorous flies (Tephritidae and Lonchaeidae), fruit losses and the dispersal of flies through the transportation of fresh fruit. *Journal of Asia-Pacific Entomology*, **24**, 50–60. https://doi.org/10.1016/j.aspen.2020.11.006

Lundberg SM, Erion GG, Lee S-I (2018) Consistent individualized feature attribution for tree ensembles. *arXiv preprint arXiv:1802.03888*.

Lundberg SM, Lee S-I (2017) A unified approach to interpreting model predictions. *Advances in neural information processing systems*, **30**, 4765-4774.

MacArthur RH (1957) On the relative abundance of bird species. *Proceedings of the National Academy of Sciences*, **43**, 293–295. https://doi.org/10.1073/pnas.43.3.293

Makumbe LD, Moropa TP, Manrakhan A, Weldon CW (2020) Effect of sex, age and morphological traits on tethered flight of Bactrocera dorsalis (Hendel) (Diptera: Tephritidae) at different temperatures. *Physiological Entomology*, **45**, 110–119. https://doi.org/10.1111/phen.12323

Motswagole R, Gotcha N, Nyamukondiwa C (2019) Thermal biology and seasonal population abundance of Bactrocera dorsalis Hendel (Diptera: Tephritidae): implications on pest management. *International Journal of Insect Science*, **11**, 1-9. https://doi.org/10.1177/1179543319863417

Mutamiswa R, Nyamukondiwa C, Chikowore G, Chidawanyika F (2021) Overview of oriental fruit fly, Bactrocera dorsalis (Hendel) (Diptera: Tephritidae) in Africa: From invasion, bio-ecology to sustainable management. *Crop Protection*, **141**, 105492. https://doi.org/10.1016/j.cropro.2020.105492

Ndiaye M, Dieng EO, Delhove G (2008) Population dynamics and on-farm fruit fly integrated pest management in mango orchards in the natural area of Niayes in Senegal. *Pest Management in Horticultural Ecosystems*, **14**, 1- 8.

Ndiaye O (2009) Plantes hôtes et foyers de réinfestation des mouches des fruits : facteurs phénologiques, morpho physiologiques déterminants sur les infestations de la mangue. *M. Sc. Degree*, *Thiès University.*

Ndiaye O, Vayssieres J-F, Rey J-Y, Ndiaye S, Diedhiou PM, Ba CT, Diatta P (2012) Seasonality and range of fruit fly (Diptera: Tephritidae) host plants in orchards in Niayes and the Thiès Plateau (Senegal). *Fruits*, **67**, 311–331. https://doi.org/10.1051/fruits/2012024

Papaïx J, Soubeyrand S, Bonnefon O, Walker E, Louvrier J, Klein E, Roques L (2022) Inferring Mechanistic Models in Spatial Ecology Using a Mechanistic-Statistical Approach. In: *Statistical Approaches for Hidden Variables in Ecology* , 69–95. John Wiley & Sons, Ltd. https://doi.org/10.1002/9781119902799.ch4

Rosenheim JA, Gratton C (2017) Ecoinformatics (Big Data) for Agricultural Entomology: Pitfalls, Progress, and Promise. *Annual Review of Entomology*, **62**, 399–417. https://doi.org/10.1146/annurev-ento-031616-035444

Rosenheim JA, Meisner MH (2013) Ecoinformatics can reveal yield gaps associated with crop-pest interactions: a proof-of-concept. *PLoS One*, **8**, e80518. https://doi.org/10.1371/journal.pone.0080518

Rosenheim JA, Parsa S, Forbes AA, Krimmel WA, Law YH, Segoli M, Segoli M, Sivakoff FS, Zaviezo T, Gross K (2011) Ecoinformatics for Integrated Pest Management: Expanding the Applied Insect Ecologist's Tool-Kit. *Journal of Economic Entomology*, **104**, 331–342. https://doi.org/10.1603/EC10380

Rwomushana I, Ekesi S, Ogol CKPO, Gordon I (2008) Effect of temperature on development and survival of immature stages of *Bactrocera invadens* (Diptera: Tephritidae). *Journal of Applied Entomology*, **132**, 832–839. https://doi.org/10.1111/j.1439-0418.2008.01318.x

Serit M, Tan K-H (1990) Immature life table of a natural population of *Dacus dorsalis* in a village ecosystem. *Tropical Pest Management*, **36**, 305–309. https://doi.org/10.1080/09670879009371493

Sigrist F (2022) Gaussian process boosting. *The Journal of Machine Learning Research,* **23**, 10565-10610

Soulsby RL, Thomas JA (2012) Insect population curves: modelling and application to butterfly transect data. *Methods in Ecology and Evolution*, **3**, 832–841. https://doi.org/10.1111/j.2041-210X.2012.00227.x

Stack Whitney K, Meehan TD, Kucharik CJ, Zhu J, Townsend PA, Hamilton K, Gratton C (2016) Explicit modeling of abiotic and landscape factors reveals precipitation and forests associated with aphid abundance. *Ecological Applications*, **26**, 2600–2610. https://doi.org/10.1002/eap.1418

Susanto A, Faradilla MG, Sumekar Y, Yudistira DH, Murdita W, Permana AD, Djaya L, Subakti Putri SN (2022) Effect of various depths of pupation on adult emergence of interspecific hybrid of *Bactrocera carambolae* and *Bactrocera dorsalis*. *Scientific Reports*, **12**, 4235. https://doi.org/10.1038/s41598-022-08295-w

Thorat L, Nath BB (2018) Insects With Survival Kits for Desiccation Tolerance Under Extreme Water Deficits. *Frontiers in Physiology*, **9**. https://doi.org/10.3389/fphys.2018.01843

Vargas RI, Walsh WA, Jang EB, Armstrong JW, Kanehisa DT (1996) Survival and Development of Immature Stages of Four Hawaiian Fruit Flies (Diptera: Tephritidae) Reared at Five Constant Temperatures. *Annals of the Entomological Society of America*, **89**, 64–69. https://doi.org/10.1093/aesa/89.1.64

Vargas RI, Walsh WA, Kanehisa D, Jang EB, Armstrong JW (1997) Demography of Four Hawaiian Fruit Flies (Diptera: Tephritidae) Reared at Five Constant Temperatures. *Annals of the Entomological Society of America*, **90**, 162–168. https://doi.org/10.1093/aesa/90.2.162

Vayssières J-F, Vannière H, Gueye PS, Barry O, Hanne AM, Korie S, Niassy A, Ndiaye Mb, Delhove G (2011) Preliminary inventory of fruit fly species (Diptera, Tephritidae) in mango orchards in the Niayes region, Senegal, in 2004. *Fruits*, **66**, 91–107. https://doi.org/10.1051/fruits/2011002

Vayssières J-F, De Meyer M, Ouagoussounon I, Sinzogan A, Adandonon A, Korie S, Wargui R, Anato F, Houngbo H, Didier C, De Bon H, Goergen G (2015) Seasonal Abundance of Mango Fruit Flies (Diptera: Tephritidae) and Ecological Implications for Their Management in Mango and Cashew Orchards in Benin (Centre & North). *Journal of Economic Entomology*, **108**, 2213–2230. https://doi.org/10.1093/jee/tov143

Vayssières J-F, Korie S, Ayegnon D (2009) Correlation of fruit fly (Diptera Tephritidae) infestation of major mango cultivars in Borgou (Benin) with abiotic and biotic factors and assessment of damage. *Crop protection*, **28**, 477–488. https://doi.org/10.1016/j.cropro.2009.01.010

Vayssières J-F, Korie S, Coulibaly O, Temple L, Boueyi SP (2008) The mango tree in central and northern Benin: cultivar inventory, yield assessment, infested stages and loss due to fruit flies (Diptera Tephritidae). *Fruits*, **63**, 335–348. https://doi.org/10.1051/fruits:2008035

Veran S, Simpson SJ, Sword GA, Deveson E, Piry S, Hines JE, Berthier K (2015) Modeling spatiotemporal dynamics of outbreaking species: influence of environment and migration in a locust. *Ecology*, **96**, 737–748. https://doi.org/10.1890/14-0183.1

Veres A, Petit S, Conord C, Lavigne C (2013) Does landscape composition affect pest abundance and their control by natural enemies? A review. *Agriculture, Ecosystems & Environment*, **166**, 110-117. https://doi.org/10.1016/j.agee.2011.05.027.

Wade M, Mignot J, Lazar A, Gaye AT, Carré M (2015) On the spatial coherence of rainfall over the Saloum delta (Senegal) from seasonal to decadal time scales. *Frontiers in Earth Science*, **3**. https://doi.org/10.3389/feart.2015.00030

Wang M, Kornblau SM, Coombes KR (2018) Decomposing the Apoptosis Pathway Into Biologically Interpretable Principal Components. *Cancer Informatics*, **17**. https://doi.org/10.1177/1176935118771082

Yang P, Carey JR, Dowell RV (1994) Temperature Influences on the Development and Demography of *Bactrocera dorsalis* (Diptera: Tephritidae) in China. *Environmental Entomology*, **23**, 971–974. https://doi.org/10.1093/ee/23.4.971

FIGURES

Figure 1 - Summary of the analysis pipeline implemented in this study. All steps presented in the figure are detailed in the section Materials & Methods.

Figure 2 - Study area and sampling sites in the Niayes region, Senegal. The distribution of the 69 orchards monitored for BD abundance among the six main sampling sites (S1 to S6) is 20, 23, 17, 5, 3 and 1, respectively. The background map represents the 13 land use classes (Jolivot 2021) considered in this study**.**

Figure 3 - Illustration of the annual dynamics of BD populations within monitored orchards. The boxplots show the median value (black line), the lower and upper quartiles (upper and lower box limits), the highest and lowest values excluding outliers (vertical lines) and outliers (black dots) of the mean number of flies captured per orchard on a weekly basis over the three years of monitoring of the 69 orchards. For a given week, the mean number of captures per orchard is the average of the number of trapped BD in all the traps set in the orchard.

Figure 4 - Estimates of within-orchard start date of population growth (t_0). For each orchard and year, the graph shows the distribution of the 500 estimates of t_0 drawn from the posterior distributions of the POPFIT model fitted to the abundance time series. The sampling sites (S1 to S6) are shown in grey on the y-axis.

Figure 5 - Ranking of the environmental predictors from the GPBoost model. The ranking is based on the SHAP values resulting from the GPBoost model applied on the 500 replicates of the dataset: for each predictor, the boxplot shows the median, first quartile, third quartile, lowest and highest values (vertical lines) and outliers (black points) of the S_{mean} values (i.e. the average of individual SHAP values) obtained from the 500 replicates.

Figure 6 - Results of the PCA performed on the 30 weather and NDWI variables. A) Variable correlation plots for the first and second PCs (top panel) and first and third PCs (bottom panel): each vector represents a type of input variable (colours) for a given month (numbers) from December (12) to May (05). B) Maps of the grid cell PCA scores, for each PC and each year of sampling.

A)

B)

Figure 7 - SHAP dependence plots. For each of the most important environmental predictors, the gam-smoothed curve of the SHAP values over the 500 replicates is represented as a black line with its 99% confidence interval in purple. Black lines on the x-axis indicate the distribution of the predictor values.