UPR[ER]–immunity axis acts as physiological food evaluation system that promotes aversion behavior in sensing low-quality food

Pengfei Liu, Xinyi Liu, Bin Qi[1,2]

Center for Life Sciences, School of Life Sciences, Yunnan Key Laboratory of Cell Metabolism and Diseases, State Key Laboratory of Conservation and Utilization of Bio-resources in Yunnan, Yunnan University, Southwest United Graduate School, Kunming, China

Footnotes: 1 Corresponding Author (Bin Qi, Email: qb@ynu.edu.cn)

Footnotes: 2 Lead Contact (Bin Qi, Email: qb@ynu.edu.cn)

Summary

To survive in challenging environments, animals must develop a system to assess food quality and adjust their feeding behavior accordingly. However, the mechanisms that regulate this chronic physiological food evaluation system, which monitors specific nutrients from ingested food and influences food-response behavior, are still not fully understood. Here, we established a low-quality food evaluation assay system and found that heat-killed *E. coli* (HK-*E. coli),* a low sugar food, triggers cellular UPR[ER] and immune response. This encourages animals to avoid low-quality food. The physiological system for evaluating low-quality food depends on the UPR[ER] (IRE-1/XBP-1) - Innate immunity (PMK-1/p38 MAPK) axis, particularly its neuronal function, which subsequently regulates feeding behaviors. Moreover, animals can adapt to a low-quality food environment through sugar supplementation, which inhibits the UPR[ER]-PMK-1 regulated stress response by increasing vitamin C biosynthesis.

This study reveals the role of the cellular stress response pathway as
physiological food evaluation system for assessing nutritional deficiencies in
food, thereby enhancing survival in nature environments.

Keywords: aversion behavior; UPR[ER]; immune response; XBP-1; PMK-1;
Food quality evaluation; Food sensing; *C. elegans*;

Introduction

Food is essential for the survival, growth and fitness of all animals. To adapt to
fluctuating environments with a wide range of food sources, animals have
developed a food evaluation system. This system enables them to identify
nutrient-rich food and avoids low-quality or toxic food, thereby maximizing their
survival prospects [1-3]. Various sensory neuron evaluation systems in animals
has evolved to evaluate food quality through vision [4,5], olfactory [3,6-12] and
gustatory senses [13-15]. Besides these sensory systems that facilitate quick
feeding decisions, animals may also initiate cellular stress response programs
to detect nutrition/toxin and trigger food response behaviors [16,17]. This could be
one of physiological food quality evaluation systems that monitor the nutritional
status of consumed food. However, the signaling events in cellular stress
responses involved in evaluating of specific nutrients and the mechanisms that
connect these signaling activities to food behaviors are largely unexplored.
More specifically, while cellular stress response through UPR[ER] [18] and PMK-
1/p38 MAPK [19] dependent immunity in response to pathogens have been
extensively studied, the functions of these cellular stress response in sensing
and evaluating specific nutrients from food remain unclear.

Vitamin C is an essential micronutrient that cannot be synthesized by humans
due to the loss of a key enzyme in the biosynthetic pathway [20]. Animals obtain
Vitamin C from their diet and possibly also from gut microbes [21]. Vitamin C is
an important physiological antioxidant and a cofactor for a family of biosynthetic

and gene regulatory monooxygenase and dioxygenase enzymes. It is also required for the biosynthesis of collagen, L-carnitine, and certain neurotransmitters [22,23]. Vitamin C has been associated with various human diseases including scurvy, immune defect and cardiovascular disease [20]. Therefore, food evaluation systems in animals for sensing Vitamin C could be critical for their fitness in nature. However, the role of the cellular stress response pathway as food evaluation system for sensing and evaluating Vitamin C remains unexplored.

In this study, using the low-quality food evaluation assay system we established [24], we uncovered the mechanism of cellular stress responses pathway as physiological food evaluation system in evaluating D-Glucose deficiency in food and downstream Vitamin C content in animals through the UPR^{ER} (IRE-1/XBP-1) - Innate immunity (PMK-1 p38 MAPK) axis. This mechanism promotes animals to leave low-quality food and is critical for their survival in nature environments.

Results

Low-quality food induces stress response in animals
Our previous studies have shown that Heat-killed *E. coli* (HK-*E. coli*), which lacks certain molecules, is considered a low-quality food that is unable to support animal growth [24,25]. Morevoer, through metabolic-seq analysis, we identified significant changes in a large numbers of metabolites (Figure S1A, Table S1)，including lipids (Figure S1B, Table S1)，amino acid and their metabolites (Figure S1C, Table S1), as well as coenzymes and vitamins (Figure S1D, Table S1). Interestingly, we observed a significant decrease in carboxylic acids and their derivatives (Figure S1E, Table S1) in *E. coli* after being heat-killed (Figure S1F, Table S1). This suggests that HK-*E. coli* is nutritionally deficient for *C. elegnas* when compared to normal *E. coli* food.

Next, we conducted two behavior assays to facilitate the analysis of the food evaluation process in animals by seeding L1 animals in assay plates (Figure 1A and 1B). In the avoidance assay, wild-type animals avoided the HK-*E. coli* food (Figure 1A). Interestingly, in the food choice assay, animals initially showed no preference between the two types of food (1-2h), but eventually exhibited a preference for high-quality food (Live *E. coli*) up until the17h mark (Figure 1B, Figure S1H). This suggests that worms depart from the HK-*E. coli* after recognizing it as low-quality food source through ingestion.

In order to investigate the pathways in animals that response to HK-*E. coli*, we performed transcriptomics analysis on worms that were cultured with both HK-*E. coli* and Live *E. coli*. Gene-expression profiling revealed that stress response genes, including those related to biotic stimulus, immune response and response to stress, are up-regulated in animals fed with HK-*E. coli* (Figure 1C, Table S2). Among these up-regulated genes, we identified 11 out of 62 of UPR[ER] target genes (Figure 1D, Figure S1G and Table S2). Additionally, about 50%-80% of up-regulated genes overlap with genes responding pathogenic bacteria [26,27] (Figure 1E, Table S2), suggesting that UPR[ER] and immune pathways may responded to low-quality food. Consistent with the results of the RNA sequencing (RNA-seq) analysis, the UPR[ER] reporter (*Phsp-4::GFP*) and immunity reporter (*Pirg-5::GFP*) were strongly induced under HK-*E. coli* feeding condition (Figure 1F and 1G). Moreover, UPR[Mt] reporter (*Phsp-6::GFP*) was weakly induced under HK-*E. coli* feeding condition (Figure S1I), and starved worm did not induce UPR[ER] and immunity (Figure S1J and S1K).

Together, these findings suggest that low-quality food (HK-*E. coli*) triggers a stress response pathway in animals, including UPR[ER] and innate immune. This implies that animals may assess the quality of food through UPR[ER] and innate immune pathway.

Animals evaluate food quality through UPR^ER-immune depended physiological food quality evaluation system

To determine whether the UPR^{ER} and innate immune pathways play a role in evaluating low-quality food, we first examined whether the activation of the UPR^{ER} by HK-*E. coli* was dependent on the known signaling components of the UPR^{ER} branches, including IRE/XBP-1, PERK/ATF-4 and ATF-6 [28,29]. We observed no difference in *Phsp-4::GFP* induction with *atf-4* (Figure S2A and S2J) or *atf-6* (Figure S2B and S2J) RNAi-mediated knockdown in animals fed with HK-*E. coli*. However, knockdown of *ire-1/xbp-1* or mutation of *xbp-1* reduced GFP fluorescence (Figure 2A, 2B and S2J). Among the 11 deferentially expressed UPR^{ER} target genes in animals fed with HK-*E. coli* from RNA-seq (Figure 1D), 64% of the genes are IRE-mediated genes (Figure S1G, Table S2). The mRNA level of IRE-1-mediated splicing of *xbp-1* is also induced in animals fed with HK-*E. coli* (Figure S2C). These data suggest that activation of the UPR^{ER} by low-quality food depends on the IRE-1/XBP-1.

To further analyze whether XBP-1-dependent UPR^{ER} activation is critical for animals to leave low-quality food, we tested food avoidance behavior using *xbp-1* mutant. The results show that *xbp-1* mutants had a significantly decreased likelihood of leaving of HK-*E. coli*, which was rescued by expressing *xbp-1* in neuron rather than intestine (Figure 2C). This indicats that XBP-1-dependent UPR^{ER} activation in neuron is critical for animals to evaluate low-quality food.

We then investigated which innate immune pathway is involved in evaluating low-quality food. First, we analyzed HK-*E. coli* induced genes from RNA-seq. Among these up-regulated genes, 82 out of 409 of PMK-1-dependent genes [30] were identified (Figure S2D, Table S2). Second, we confirmed the induction of several well-known PMK-1 target genes in RNA-seq data[31] (Figure S2E) and

reporter analysis (Figure S2F and S2G). Moreover, the induction of *Pirg-5::GFP*

was abolished in *pmk-1* knockdown animals fed with HK-*E. coli* (Figure 2D and

S2K). Third, we found that the phosphorylated PMK-1 (p-PMK-1) level was

prominently increased in wild-type N2 animals fed HK-*E. coli* compared to

feeding *E. coli* OP50 (Figure 2E). Finally, *pmk-1* mutants had a decreased

likelihood of leaving of HK-*E. coli*, which was rescued by expressing *pmk-1* in

neurons rather than intestine (Figure 2F). These data suggest that PMK-1

regulated immune pathway evaluates low-quality food, especially the neuronal

PMK-1 has a critical function for food quality response.

XBP-1 and PMK-1 are in the same pathway for evaluating food quality

Next, we explored the connection between UPR^{ER} (IRE-1/XBP-1) and innate

immunity (PMK-1 p38 MAPK) in food quality evaluation. We found that *Pirg-*

5::GFP induction (Figure S2H and S2K) and PMK-1 activation (Figure 2E) were

decreased in animals with *xbp-1* mutation or knockdown when fed with HK-*E.*

coli, suggesting that XBP-1 could regulate PMK-1 under this condition.

Additionally, *Phsp-4::GFP* induction under HK-*E. coli was* not affected in

animals with *pmk-1* RNAi (Figure S2I), indicating that XBP-1-dependent UPR^{ER}

activation is not regulated by PMK-1. Finally, we constructed a double mutant

of *xbp-1* and *pmk-1* and found that the food avoidance phenotype of the double

mutant was similar to the *pmk-1* mutant (Figure 2G), indicating that PMK-1 is

downstream of XPB-1 in responding to low-quality food.

Sugar deficiency in HK-*E. coli* food induces stress response and

avoidance behavior in animals

We then investigated which nutrients/metabolites are sensed by animals

through the XBP-1-PMK-1 axis for food quality evaluation. First, we hypothesized that the nutrition status is improved in *E. coli* mutant (HK-treatment), which could inhibit URP^{ER} and immune response in animals (Figure S3A). We established a system for screening the *E. coli* mutant keio library (Figure S3A), and identified 20 *E. coli* mutants that did not induce *hsp-4::GFP* through the UPR^{ER} reporter (*irg-5::GFP*) after three rounds of screening (Table S3). From these 20 E. coli mutant, we identified 9 *E. coli* mutants that did not induce *irg-5::GFP* through the immunity reporter (*irg-5::GFP*) screening (Figure S3B, S3C, Table S3). Animals fed HK-*yfbR*, which catalyzes carbohydrate derivative metabolic process[32], had a decreased ability to leave food (Figure 3A, Table S3), indicating that HK-*yfbR* may be a higher quality food for animals compared to HK-K12.

Secondly, we preformed a metabolomics analysis of different quality food (HK-K12, HK-*yfbR* and Live-K12). We found that the level of 13 metabolites were similar between HK-*yfbR* and Live-K12, but significantly changed in HK-K12 (Figure 3B, Figure S3D, and Table S1). We also found that genes involved in glycolysis/gluconeogenesis were up-regulated in animals fed with HK-*E. coli* (Figure S3E), suggesting that glycolysis/gluconeogenesis metabolism is disordered in animals fed with HK-*E. coli,* which may result from changes in sugar/carbohydrate intake. The carbohydrates (D-trehalose, Lactose, and D-(+)-sucrose) were also decreased in HK-*E. coli* (Figure 3B), suggesting that carbohydrate deficiency may induce stress response and avoidance behavior in animals.

Thirdly, to determine which carbohydrate inhibits stress response in animals, we supplemented each metabolite to HK-*E. coli* and found that only Lactose, and D-(+)-sucrose inhibited HK-*E. coli* induced UPR^{ER} (Figure 3C and 3D, Figure S3F). Moreover, we found from our metabolomic data that the sugar level, including Lactose, and D-(+)-sucrose, and D-(+)-Glucose, was also

decreased in HK-*E. coli* (Figure 3B, Table S1). Since Lactose and D-(+)-sucrose are hydrolyzed to produce glucose[33,34], we wondered whether glucose also inhibits the stress response in animals. We found that D-(+)-Glucose supplementation also inhibited HK-*E. coli* induced UPR[ER] (Figure 3E), immune response (Figure 3F, 3G and Figure S3G) and avoidance (Figure 3H). Moreover, sugar supplementation did not affect UPR[ER] and immunity in normal food (OP50) or starved condition (NGM) (Figure S3H and S3I).

Together, these findings suggest that sugar/carbohydrates deficiency in HK-*E. coli* induces a stress response and avoidance behavior in animals, which can be inhibited by D-(+)-Glucose supplementation. This implies that animals may evaluate the sugars deficiency through the activation of UPR[ER] and immune responses.

Animals could overcome a low-quality food environment by sugar supplementation through vitamin C biosynthesis

We discovered that D-(+)-Glucose supplementation inhibited HK-*E. coli* induced UPR[ER] (Figure 3E), immune response (Figure 3F, 3G and Figure S3G) and avoidance (Figure 3H). Simultaneously, vitamin C (VC), which is synthesized by glucuronate pathway using D-Glucose [35,36] (Figure 4A), was found contribute to neuroprotective[37,38], immune defense[39,40], and inhibits inflammatory and ER stress[41,42]. This led us to question whether the Vitamin C biosynthesis pathway is involved in evaluating low-quality food by using D-Glucose.

Firstly, we observed an increase in the vitamin C level in animals when fed with HK-yfbR (Figure 4B), a high carbohydrate food compared to HK-E. coli (Figure 3B and Figure S1E). However, the VC level in bacteria is same (Figure S4A).

The VC level also increased when D-Glucose (Figure S4B) or D-glucuronate (D-GlcA) was added to HK-*E. coli* (Figure 4C), which was abolished by knocking down VC biosynthesis genes (Figure 4C and S4B). This suggests that additional of sugar supplementation promotes VC synthesis in animals fed with HK-*E. coli.*

Secondly, we hypothesized that animals could overcome a low-quality food (HK-*E. coli*) environment by inhibiting the stress response through increasing vitamin C biosynthesis. We found that VC or D-glucuronate (D-GlcA) supplementation inhibited HK-*E. coli* induced UPR^{ER} (Figure 4D), immune response including *irg-5/sysm-1* reporter expression (Figure 4E and Figure S4C) and p-PMK-1 (Figure 4F and 4G), as well as food avoidance (Figure 4H).

Finally, we asked whether inhibition of stress response and avoidance by sugar supplementation depends on the vitamin C biosynthesis pathway. We found that suppression of HK-*E. coli* induced UPR^{ER} (Figure 4I), immune response (Figure 4J and Figure S4D) and food avoidance (Figure 4K) by D-GlcA/sugar supplementation was abolished in animals with RNAi of VC biosynthesis genes. Food selection behavior assays showed that D-GlcA/sugar supplementation increased the preference for heat-killed bacteria, which was also suppressed by knocking down VC biosynthesis genes (Figure S4E). However, VC supplementation still suppressed the UPR^{ER} (Figure 4I), immune response (Figure 4J and Figure S4D) and food avoidance (Figure 4K), and increased the food preference (Figure S4E) in animals with or without RNAi of VC biosynthesis genes. This suggests that VC, as the final metabolite synthesized from D-Glucose, is critical for low-quality food response in animals.

Together, these data indicate that the Vitamin C biosynthesis pathway is critical for evaluating whether food is of higher quality and can be eaten by animals. It also suggests that animals could improve their VC levels to adapt to

267   bad food environment.

**Animals evaluate sugar and vitamin C through neuronal XBP-1 and PMK-
1**

As D-GlcA/sugar and VC supplementation suppressed HK-*E. coli* induced

UPR^{ER}, immune response and food avoidance behavior, we investigated

whether animals evaluate sugar and VC through XBP and PMK-1 dependent

pathways. We performed a food selection behavior assay by adding D-Glc, D-

GlcA or VC to the NGM, *E. coli* and HK-*E. coli* (Figure 5A). The food selection

behavior assays revealed that supplementation with D-Glc, D-GlcA, or VC

inhibits the animals' choice of sugar or VC on *E. coli*-OP50 feeding conditons

(Figure S4F). This suggests that supplementation with D-Glc, D-GlcA, or VC

may alter the metabolites of live bacteria, leading to avoidance by the animals.

There was no preference observed on NGM (no food condition)

supplementation with D-Glc and VC (Figure S4G), indicating that the intake of

sugar or VC alone does not influence animal preference. However, alone D-

GlcA could influence worm physiology which induces preference change

(Figure S4G). Interestingly, D-Glc and D-GlcA (Figure 5B and 5C) or VC (Figure

5D) supplementation increased the preference for heat-killed bacteria, which

was suppressed in *xbp-1* or *pmk-1* mutant animals. However, this preference

was also rescued in *xbp-1* or *pmk-1* mutant animals by expressing XBP-1 or

PMK-1 in neurons rather than intestine (Figure 5B-D), indicating that neuronal

XBP-1 and PMK-1 are critical for physiological food elevation system for

monitoring the level of sugar and VC under low-quality food condition.

Discussion

To better survive, animals must evolve a system to recognize and evaluate the

quality of their food. This includes the sensory neuron evaluation system for

immediate response and feeding decision [2], as well as physiological food

evaluation system for chronic response to ingested food. In our previous study,

we discovered that the TORC1-ELT-2 pathway, acting as master regulators in

intestine, evaluates vitamin B2 deficiency in low-quality food (HK-*E. coli)* and

regulates gut digestive activity to impact animal's food behavior [24]. To further

identified the mechanism by which animals evaluate low-quality food (HK-*E.*

coli), we performed metabolomics and transcriptomics analyses to identify

specific nutrition deficiencies in low-quality food and the cellular response

pathways that are involved in food evaluation pathway. This study identified a

physiological food evaluation mechanism by which animals detect sugar (in

food) and vitamin C (in animals) deficiency through UPR^{ER} (IRE-1/XBP-1) -

Innate immunity (PMK-1/p38 MAPK) regulated cellular stress response

program in neurons that dictates food avoidance and selection behaviors.

Unlike the sensory neuron evaluation system, which permits rapid feeding

decisions through smell and taste, the cellular stress response as physiological

food evaluation system describe here requires a slow and multi-step signal

transduction process after the ingestion of food. The disruption of cellular

homeostasis by ingested of low-quality or toxic food can activate stress

response mechanisms that both increase the cellular ability to withstand and

adapt to this disruption of homeostasis and promote behavioral strategies to

avoid these conditions and lessen their impact on the organism. These cellular

stress response mechanisms include heat shock response, unfolded protein

response, oxidative stress response [43]. Therefore, this slow physiological food

evaluation system is an evolutionary adaptation mechanism for detecting

nutrition deficiencies in food that was not detected by quick sensory neuronal

system.

One of cellular stress response, UPR[ER], is activated by stresses such as

infection and nutrition deficiency that affect homeostasis in the endoplasmic

reticulum (ER) [28]. The activation of the UPR[ER] by expression of *xbp-1s* in the

nervous system has recently been shown to promote changes in feeding and

foraging behavior [44]. The p38 PMK-1 pathway regulates the expression of

secreted innate immune effectors and is required for survival during infection [45].

Moreover, UPR[ER] and innate immunity increased by disrupting in vivo state

would induce worm aversion from food lawn[46-49]. Thus, these two cellular stress

response pathways are critical for animals' survival in changing environment.

However, whether UPR[ER] (IRE-1/XBP-1) - Innate immunity (PMK-1/p38 MAPK)

evaluates food quality under physiological conditions is largely unexplored. Our

study presents evidence that low-quality food (HK-*E. coli*) activate UPR[ER] and

PMK-1, and promotes animals to leave food. It is independent of avoidance

behavior of hyperactivation of kgb-1/MAPK pathway[49] (Figure S4H). Mover, our

study also shows that neuronal UPR[ER] and PMK-1 are critical for evaluating

low-quality food. Therefore, our study established a novel physiological food

evaluation system by activating the cellular stress response program with

UPR[ER] (IRE-1/XBP-1) - Innate immunity (PMK-1/p38 MAPK).

Collectively, this study uncovers the unexpected function of UPR[ER] (IRE-1/XBP-

1) - Innate immunity (PMK-1/p38 MAPK) as a physiological food evaluation

system for evaluating and sensing food quality in animals. It also highlights the

utility of the HK-*E. coli* (low-quality food) - *C. elegans* interaction as a means to

dissect the mechanism of food evaluation system in assessing food. Most

importantly, it reveals that animals are capable of altering their nutrient (Vitamin

C) levels through in vivo synthesis or food intake to adapt to a poor food

environment when better food choices are not available.

Author Contributions

P. L designed, performed experiments, analyzed data. X. L constructed all

transgenic animals. B.Q. designed research, supervised this study, and wrote the paper with inputs from P. L.

Acknowledgments

We thank the Caenorhabditis Genetics Center (CGC) (funded by NIH P40OD010440) for strains; Dr. Zhao Shan for suggestions. This work was supported by the Ministry of Science and Technology of the People's Republic of China (2019YFA0803100，2019YFA0802100), the National Natural Science Foundation of China (32170794), Yunnan Applied Basic Research Projects (202001AV070011, 202001AW070006, 202201AT070196), the Yunnan University Startup Program.

Declaration of interests

The authors declare no competing interests.

Main Figure Legends

Figure 1. The stress response is induced in animals fed low-quality food, HK-*E. coli*.

(A) Schematic drawing and quantitative data of the food aversion assay. Circles indicate the food spot for live (yellow) and HK-OP50 (red) bacteria, respectively. The animals were scored 16-17 hours after L1 worms were placed on the food spot. Data are represented as mean ±SD from three independent experiments，79-129 animals/assay.

(B) Schematic method and quantitative data of the food selection assay. Live (yellow), heat-killed (red) *E. coli* and LB as the buffer for *E. coli* were placed on indicated position. Synchronized L1 worms were place in the center spot. The selection index was calculated at the indicated time. Data are represented as mean ±SD from eight independent experiments, 123-792 animals/assay.

(C) GO enrichment analysis of up-regulated genes in animals fed with HK-*E. coli* vs live *E. coli*.

(D) Venn diagram showing numbers of UPR[ER] target genes and up-regulated genes in animals fed HK-*E. coli*, and their overlap.

(E) Venn diagram showing numbers of induction genes by four pathogenic bacteria and HK-*E. coli* induced genes, and their overlap. The gene expression data was extracted from published data of animals' infection with Pseudomonas aeruginosa (PA) [26], Bacillus thuringiensis (Bthu) [27], Staphylococcus aureus (Saur) [27], and Serratia marcescens (Smar) [27].

(F-G) GFP fluorescence images and bar graph showing that *Phsp-4::GFP* (F) and *Pirg-5::GFP* (G) were induced in animals fed wiht HK-*E. coli.* n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

For all panels, Scale bar shows on indicated figures, 50 μm. * p<0.05, ** p<0.01, *** p < 0.001, **** p<0.0001, ns: no significant difference. Precise P values are provided in Raw Data.

Figure 2. Animals evaluate food quality through UPRER (ire-1/xbp-1) - Innate immunity (pmk-1 MAPK) axis.

(A-B) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Phsp-4::GFP* was decreased in animals with *ire-1* or *xbp-1* RNAi treatment (A) or *xbp-1* mutation (B). n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(C) Food aversion assay showing that *xbp-1* mutation eliminated the discrimination against HK-*E. coli.* However, this effect is rescued by expressing *xbp-1* in neurons rather than intestine. Data are represented as mean ±SD from five independent experiments, 156-763 animals/assay.

(D) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Pirg-5::GFP* was decreased in animals with *pmk-1* RNAi treatment. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(E) Western blot images showing the level of p-PMK-1 in L1 animals (Wild-type N2 and *xpb-1* mutant) fed with OP50 or HK-OP50 for 4 h. The level of p-PMK-1 is induced in animals fed HK-OP50.

(F) Food aversion assay showing that *pmk-1* mutation eliminated the discrimination against HK-*E. coli.* However, this effect is rescued by expressing *pmk-1* in neurons rather than intestine. Data are represented as mean ±SD from five independent experiments, 168-492 animals/assay.

(G) Food aversion assay in wild-type, *xbp-1*, *pmk-1* and double mutant. Data are represented as mean ±SD from five independent experiments, 259-490 animals/assay..

For all panels, Scale bar shows on indicated figures, 50 μm. * $p<0.05$, ** $p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P values are provided in Raw Data.

Figure 3. *HK-E. coli* is low sugar food, which induce stress response and avoidance behavior in animals.

(A) Food aversion assay showing that wild-type animals eliminated the discrimination against HK-*E. coli* when *yfbR* is mutated in *E. coli*. Data are represented as mean ±SD four independent experiments, 251-490 animals/assay.

(B) Heat map showing the thirteen differential metabolites from HK-K12, HK-*yfbR*, and K12 in 4 independent experiments. Color indicates the relative level of each metabolite.

(C-D) The bar graph showing that HK-*E. coli* induced *Phsp-4::GFP* was decreased in animals with lactose (C) or D-(+)-sucrose (D) supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(E-F) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Phsp-4::GFP* (E) and *Pirg-5::GFP* (F) were decreased in animals with D-(+)-

Glucose (D-Glc) supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(G) Western blot images showing the level of p-PMK-1 in L1 animals fed HK-*E. coli* with or without D-(+)-Glucose (D-Glc) supplementation for 4 h. The level of p-PMK-1 is decreased in animals fed HK-OP50+D-Glc.

(H) Food aversion assay showing that wild-type animals eliminated the discrimination against HK-*E. coli* with D-Glc supplementation. Data are represented as mean ±SD six independent experiments, 190-492 animals/assay.

For all panels, Scale bar shows on indicated figures, 50 μm. * $p<0.05$, ** $p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P values are provided in Raw Data.

Figure 4. Vitamin C biosynthesis pathway is critically involved in evaluating sugar in the food

(A) Cartoon illustration of a simplified, Vitamin C biosynthesis pathway in *C. elegans*. The relevant coding genes of enzymes was labeled with red.

(B) The level of total L-ascorbic acid (TAA) in animals fed with Live-K12, HK-K12, or HK-yfbR. Data are represented as mean ±SD from six independent experiments.

(C) The level of total L-ascorbic acid (TAA) in animals (control or knockdown of Vitamin C biosynthesis genes) fed with HK-*E. coli* with or without D-glucuronate (D-GlcA) supplementation. Data are represented as mean ±SD from eight independent experiments.

(D-E) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Phsp-4::GFP* (C) and *Pirg-5::GFP* (D) were decreased in animals with D-GlcA or Vitamin C supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(F-G) Western blot images showing the level of p-PMK-1 in L1 animals fed with HK-*E. coli* with D-GlcA or Vitamin C supplementation for 4 h. The level of p-PMK-1 is decreased in animals with D-GlcA (F) or Vitamin C (G) supplementation.

(H) Food aversion assay showing that wild-type animals eliminated the discrimination against HK-*E. coli* with D-GlcA or Vitamin C supplementation. N is number of independent experiments. Data are represented as mean ±SD from four independent experiments, 153-292 animals/assay.

(I-K) The bar graph showing that suppression of HK-*E. coli* induced *Phsp-4::GFP* (I), *Pirg-5::GFP* (J) and food avoidance (K) by D-GlcA supplementation was abolished in animals with RNAi of VC biosynthesis genes, which was not affect by Vitamin C supplementation. n is the number of worms scored from at least three independent experiments and Data are represented as mean ±SD(I-J). Data are represented as mean ±SD from five independent experiments, 252-537 animals/assay (K).

For all panels, Scale bar shows on indicated figures, 50 μm. * $p<0.05$, ** $p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P values are provided in Raw Data.

Figure 5. Animals evaluate sugar and vitamin C through neuronal XBP-1 and PMK-1

(A) Schematic method of the food selection assay. Heat-killed *E. coli* (red) and HK-*E. coli* with chemical supplementation (green) were placed on indicated position. Synchronized L1 worms were then place in plate. After 16-17hs, the selection index was calculated.

(B-D) Food selection assay showing that *xbp-1* or *pmk-1* mutation eliminated the preference of HK-*E. coli* with D-Glc (B), D-GlcA (C) or Vitamin C (D) supplementation, which was rescued in *xbp-1* or *pmk-1* mutant animals by

expressing XBP-1 or PMK-1 in neurons rather than intestine.

Data are represented as mean ±SD from five independent experiments, 68-647
animals/assay (B).

Data are represented as mean ±SD from six independent experiments, 83-701
animals/assay (C).

Data are represented as mean ±SD from six independent experiments, 67-
1035 animals/assay (D).

For all panels, Scale bar shows on indicated figures, 50 μm. * $p<0.05$, **
$p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P
values are provided in Raw Data.

Figure 6. Schematic model of physiological food evaluation system in
evaluating/sensing sugar and vitamin C through UPR^{ER} (IRE-1/XBP-1) -
Innate immunity (PMK-1/p38 MAPK) axis.

Vitamin C level is low in animals fed low sugar food, HK-*E. coli*. Sugar and
Vitamin C deficiency activate cellular UPR^{ER} and immune response, which
promote animals to leave low-quality food and seek better food for survival. This
cellular stress regulated physiological food evaluation system depends UPR^{ER}
(IRE-1/XBP-1) - Innate immunity (PMK-1/p38 MAPK) axis in neuron.

<u>Supplemental Figure legends</u>

Figure S1. Food selection assay of animals fed HK-*E. coli* or *E. coli*.
Relative to Figure 1.

(A-E) Metabolomics analysis of different quality food (HK-K12, HK-*yfbR* and
Live-K12). Cluster analysis of all metabolites (A), lipids and their derivatives (B),
amino acids and their derivatives (C), coenzymes and vitamins (D), and
carbohydrates and their derivatives (E) from Live-K12, HK-K12, and HK-yfbR.

Color indicates the relative level of each metabolite. HK-K12: heat-killed *E. coli* wild-type K12; HK-*yfbR:* heat-killed *E. coli mutant yfbR;* K12: live *E. coli* wild-type K12. z-score for standardizing data, complete for bi-clustering algorithm, and Euclidean for distance method.

(F) Principal component analysis to test the repeatability of the metabolic experiment. HK-K12: heat-killed *E. coli* wild-type K12; HK-yfbR: heat-killed *E. coli* mutant yfbR; K12: live *E. coli* wild-type K12. LB: LB buffer for culturing *E. coli.*

(G) GO enrichment analysis of UPR^{ER} dependent IRE-1 branch.

(H) The number of worms in each position calculated at the indicated time, indicating that animals initially select both foods (1-2h), but eventually favor high-quality food (Live E. coli) until 17h.

(I) UPR^{mt} Reporter (*Phsp-6:GFP*) was weakly induced in animals fed with HK-E. coli

(J) UPR^{ER} Reporter (Phsp-4:GFP) expression in animals under normal food (OP50), low quality food (HK-OP50), and starved (M9: hatching L1 worm in M9, NGM) condition;

(K) Immunity Reporter (Pirg-5:GFP) expression in animals under normal food (OP50), low quality food (HK-OP50), and starved (M9: hatching L1 worm in M9, NGM).

For all panels, Scale bar shows on indicated figures, 50 μm. * p<0.05, ** p<0.01, *** p < 0.001, **** p<0.0001, ns: no significant difference. Precise P values are provided in Raw Data.

Figure S2. UPRER and Innate immunity pathway in animals are critical for evaluating HK-*E. coli*. Relative to Figure 2.

(A-B) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Phsp-4::GFP* was not affected in animals with *atf-4* (A) or *atf-6* (B) RNAi treatment. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(C) qPCR showing that IRE-1-mediated splicing of *xbp-1* mRNA is induced in animals fed with *HK-E. coli.*

(D) Venn diagram showing the numbers of PMK-1 dependent genes [30] and up-regulated genes in animals fed HK-*E. coli,* and their overlap.

(E) The expression of PMK-1 dependent genes which was extracted from RNA-seq data from animals fed with *HK-E. coli*. The data from average of three independent experiments (N=3)

(F-G) GFP fluorescence images and bar graph showing that *Psysm-1::GFP* (F) and *Pirg-1::GFP* (G) were induced in animals fed *HK-E. coli.* n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(H) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Pirg-5::GFP* was decreased in animals with *ire-1* or *xbp-1* RNAi treatment. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(I) GFP fluorescence images and bar graph showing that *Phsp-4::GFP* was not affected in animals with *pmk-1* RNAi treatment. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(J) UPR^{ER} Reporter (Phsp-4::GFP) expression in animals with candidate RNAi feeding OP50 or HK-OP50

(K) immunity Reporter (Pirg-5::GFP) expression in animals with candidate RNAi feeding OP50 or HK-OP50

For all panels, Scale bar shows on indicated figures, 50 µm. * $p<0.05$, ** $p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P values are provided in Raw Data.

Figure S3. Low sugar food, HK-*E. coli*, induce stress response and avoidance behavior in animals. Relative to Figure 3.

(A) Flow chart of strategy for *E. coli* keio mutant screening. We identified 20 *E. coli* mutants that did not induce *hsp-4::GFP* through the UPR^{ER} reporter (*irg-5::GFP*) after three rounds of screening (Table S3). From these 20 E. coli mutant, we identified 9 *E. coli* mutants that did not induce *irg-5::GFP* through the immunity reporter (*irg-5::GFP*) screening (Table S3).

(B-C) The bar graph showing that HK-*E. coli* induced *Phsp-4::GFP* (B) and *Pirg-5::GFP* (C) was decreased in animals fed mutant *E. coli* (Heat-killed). n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(D) Venn diagram showing the number of deferentially metabolites in HK-*E. coli*-K12, HK-*E. coli-yfbR* and *E. coli*.

(E) KEGG enrichment analysis of deferentially expressed genes in animals fed HK-*E. coli* vs live *E. coli*. We noticed that all most of glycolysis/gluconeogenesis genes are up-regulated in animals fed HK-*E. coli*.

(F) The bar graph showing that HK-E. coli induced *Phsp-4::GFP* was not affected in animals with D-(+)-Trehalose supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean $\pm$ SD. .

(G) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Psym-1::GFP* was decreased in animals with D-(+)-Glucose (D-Glc) supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean $\pm$ SD.

(H) UPR^{ER} Reporter (Phsp-4::GFP) expression animals with D-Glc supplementation under OP50, HK-OP50, or NGM condition.

(I) immunity Reporter (Pirg-5::GFP) expression animals with D-Glc supplementation under OP50, HK-OP50, or NGM condition.

For all panels, Scale bar shows on indicated figures, 50 µm. * $p<0.05$, ** $p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P values are provided in Raw Data.

Figure S4.   Vitamin C biosynthesis pathway is critically for evaluating low sugar. Relative to Figure 4.

(A) The level of total L-ascorbic acid (TAA) in Live-K12, HK-K12, or HK-yfbR. N is number of independent experiments. Data are represented as mean ±SD from six independent experiments.

(B) The level of total L-ascorbic acid (TAA) in animals fed HK-*E. coli* with or without D-Glc supplementation. Data are represented as mean ±SD from from eight independent experiments.

(C) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Psym-1::GFP* was decreased in animals with D-GlcA or Vitamin C supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(D) The bar graph showing that suppression of HK-*E. coli* induced *Psysm-1::GFP* by D-GlcA supplementation was abolished in animals with RNAi of VC biosynthesis genes, which was not affect by Vitamin C supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ±SD.

(E) Food selection assay showing that the preference of HK-*E. coli* with D-GlcA supplementation was abolished in animals by RNAi of Vitamin C biosynthesis genes. Data are represented as mean ±SD from six independent experiments, 427-775 animals/assay.

(F) Food selection assay for OP50 & OP50 + + D-Glc, D-GlcA, or VC, respectively. Data are represented as mean ±SD from five independent experiments, 241-1182 animals/assay.

(G) Food selection assay for buffer & buffer + D-Glc, D-GlcA, or VC, respectively. Data are represented as mean ±SD from five independent experiments, 8-153 animals/assay.

(H) Food avoidance assay for N2 & kgb-1 mutant animals fed with HK-E. coli. Data are represented as mean ±SD from six independent experiments, 348-660 animals/assay.

For all panels, Scale bar shows on indicated figures, 50 μm. * $p<0.05$, ** $p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P values are provided in Raw Data.

Supplemental Tables:

Table S1. Metabolices analysis. Relative to Figure S1A, S1B, S1C, S1D, S1E, S1F, 3B, and S3D.

Table S2. RNA-seq analysis. Relative to Figure 1C, 1D, 1E, S1G, and S2D.

Table S3. Screening data for *E. coli* mutant keio library. Relative to Figure s S3A, S3B, and S3C.

Table S4. Metabolomics-seq data of HK-K12, HK-yfbR and K12.

Table S5. RNA-seq data of animals fed with HK-*E. coli OP50* and *E. coli OP50.*

Raw-data. Raw data for experiments.

Star Methods

Resource Availability

Lead contact

Further information and requests for reagents may be directed to the Lead contact Bin Qi (qb@yun.edu.cn).

Materials availability

All reagents and strains generated by this study are available through request to the lead contact with a completed Material Transfer Agreement.

Data and code availability

Metabolomics-seq data are accessible in Table S4

RNA-seq data are accessible in Table S5.

This paper does not report original code.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Experimental model and subject details

***C. elegans* strains and maintenance**

Nematode stocks were maintained on nematode growth medium (NGM) plates seeded with bacteria (*E. coli OP50*) at 20℃.

1) The following strains/alleles were obtained from the Caenorhabditis Genetics Center (CGC) or as indicated:

N2 Bristol (wild type control strain);

AU78: agIs219 [*T24B8.5p::GFP::unc-54 3' UTR + ttx-3p::GFP::unc-54 3' UTR*];

SJ4005: *Phsp-4::GFP(zcIs4)*;

AY101: acIs101 [*F35E12.5p::GFP + rol-6(su1006)*]IG274: frIs7 [*nlp-29p::GFP +*

col-12p::DsRed] IV;

SJ17: *xbp-1 (zc12)*;

KU25: *pmk-1(km25)*;

AY102: *pmk-1(km25)* IV; acEx102 [*vha-6p::pmk-1::GFP + rol-6(su1006)*];

YNU108: Ex[*Prgef-1::pmk-1::GFP；Podr-1::RFP*][50];

xbp-1(tm2482) [51];

KU21：*kgb-1(km21)*;

AU133: *agIs17 [myo-2p::mCherry + irg-1p::GFP] IV;*

SJ4100: *zcIs13 [hsp-6p::GFP + lin-15(+)]*.

2)The following strains were constructed by this study:

YNU242: *xbp-1(tm2482); pmk-1(km25)* double mutant was constructed by

crossing: *xbp-1(tm2482)* with KU25[*pmk-1(km25)*].

YNU240: ylfEx149 [*xbp-1(tm2482); Prgef-1::xbp-1::GFP; Podr-1::RFP*]

transgene strain was constructed by injecting plasmid *Prgef-1::xbp-1::GFP* with

Podr-1::RFP in *xbp-1(tm2482)* background

YNU241: ylfEx150 [*xbp-1(tm2482); Pges-1::xbp-1::GFP; Podr-1::RFP*]

transgene strain was constructed by injecting plasmid Pges-1:xbp-1::GFP with

Podr-1::RFP in *xbp-1(tm2482)* background

Bacterial strains

E. coli-OP50, *E. coli-K12* (BW25113), and *E. coli-K12* mutant were cultured at

37˚C in LB medium. A standard overnight cultured bacteria was then spread

onto each Nematode growth media (NGM) plate.

Method Details

Generation of transgenes

1）To construct the *C. elegans* plasmid for expression of *xbp-1* in neuron, 3057bp promoter of *rgef-1* and genomic DNA of *xbp-1* was inserted into the PPD95.77 vector. DNA plasmid mixture containing *Prgef-1::xbp-1::GFP* (25ng/ul) and *Podr-1p:RFP* (25ng/ul) was injected into the gonads of adult *xbp-1(tm2482)*.

2）To construct the *C. elegans* plasmid for expression of *xbp-1* in intestine, 2549 bp promoter of ges-1 and genomic DNA of xbp-1 was inserted into the PPD95.77 vector. DNA plasmid mixture containing *Pges-1:xbp-1::GFP* (25ng/ul) and *Podr-1p:RFP (25ng/ul)* was injected into the gonads of adult *xbp-1(tm2482)*.

Preparation and feeding of worm food

We followed an established protocol [24,25] to prepare heat-killed (HK) *E. coli*. Briefly, a standard OD600=0.5-0.6 of *E. coli* OP50 and *E. coli* K12 grown in LB broth was concentrated to 1/20 vol and was then heat-killed at 80℃ for 180 min. About 150ul of the heat-killed bacteria was spread onto each 35mm NGM plate.

Preparation of HK-*E. coli* + carbohydrate or Vitamin C food

1) 100ul of water, 100ul of L-ascorbic acid (dissolved in water at a concentration of 100mg/ml, Sangon Biotech, 100143-0100) or 100 ul of D-Glucuronic acid (dissolved in water at a concentration of 100mg/ml，Adamas, 1102520) was mixed with 500 ul of HK-*E. coli*, then 150ul of the mixture was spread onto 35mm NGM plates.

2) 12.5ul of water or 12.5ul of D-(+)-Glucose (dissolved in water at a concentration of 100mg/ml, Sangon Biotech, A501991-0500) was mixed with 500 ul of HK-*E. coli*, then 150ul of the mixture was spread onto 35mm NGM plates.

Behavioral assay

C. elegans **selection assays**

746 For *C.elegans* to have enough space to evaluate food, we add 18ul of the

747 sample onto a 35mm NGM plate. This creates a round lawn with a radius of

748 0.5mm, which occupies about 8% of the total plate area.

$$\frac{a}{b} = \frac{\pi r_1^2}{\pi \gamma_2^2} = \frac{0.5mm^2}{1.75mm^2} = 8\%$$

a: the area of bacterial lawn
b: the space of worm life (area of culture dish)

1) 18ul of heat-killed OP50, live OP50, and LB broth (as the buffer for bacteria)

was added into 35mm NGM plate in an equilateral triangle pattern. Then,

synchronized L1 worms were seeded in the center of NGM plate for 16-17h at

20℃ (as indicated in Figure1B).

2) 18ul of heat-killed OP50 and heat-killed OP50 with D-GlcA or Vitamin C was

added into 35mm NGM plate in an equilateral triangle pattern., then

synchronized L1 worms were seeded on equilateral triangle of NGM plate for

16-17h at 20℃ (as indicated in Figure 5A).

Here is Selection rate formula:

$$selection\ rate = \frac{\frac{worm\ amount}{one\ lawn\ area}}{\frac{worm\ amount}{lawn1\ area} + \frac{worm\ amount}{lawn2\ area} + \frac{worm\ amount}{lawn3\ area} \dots}$$

C. elegans aversion assays

18ul food was spread out the center of NGM plate, then synchronized L1 worms

were seeded on center of food for 16-17h at 20℃.

$$\text{Aversion index} = \frac{worm\ amount\ of\ out\ of\ lawn}{worm\ amount\ of\ (out + in)\ lawn}$$

Three to ten replicates for each condition were performed for each assay, and

the experiments were duplicated on different days.

Analysis of the fluorescence intensity in worms

The synchronized L1 worms carrying either UPR^{ER} reporter (*Phsp-4::GFP*) or

innate immunity reporter (*Pirg-5::GFP*; Psysm-1p::GFP; *Pirg-1p::GFP*) were

seeded on NGM with indicated food and incubated for 24h at 20℃. For

fluorescence imaging, worms were anesthetized with 25 mM levamisole and

photographed using either an Olympus BX53 microscope or Olympus MVX10

dissecting microscope equipped with a DP80 camera.

The fluorescence intensity in entire intestinal region was quantified using

ImageJ software and normalized to the body area.

E. coli **Keio collection screen**

The whole Keio *E. coli* single mutant collection (Baba et al., 2006) was

screened. Mutant bacteria strains, as well as the wild-type control strain

BW25113, were cultured in LB medium with 50µg/ml kanamycin at 37℃ until an

OD600 of 0.5-0.6 was reached. The bacteria were then heat-killed following our

established protocol [24], and 150 ul of the heat-killed the mutant *E. coli* was

spread onto 3.5 cm NGM plates. Synchronized L1 worms carrying UPR^{ER}

reporter (*Phsp-4::GPF*) were seeded and cultured for 24h at 20℃. The

fluorescence was then examined by using an Olympus MVX10 dissecting

microscope, progressive screening three times. next, an 4th screen was

performed using immune reporter (*Pirg-5::GPF*) animals fed with HK-E. coli

mutants that reduced the *Phsp-4::GPF* fluorescence. Finally, an aversion

behavior assay was performed using HK-*E. coli* mutants that both reduced

Phsp-4::GPF and *Pirg-5::GPF.* HK-*E. coli* mutants that reduced UPR^{ER}, immune and avoidance behavior were identified through this screening.

RNAi treatment

Bacterial clones from either the MRC RNAi library [52] or the ORF-RNAi Library [53] were used. RNAi plates were prepared by adding IPTG to NGM agar to a final concentration of 1 mM. Overnight *E. coli* cultures (LB broth containing 100 ug/ml ampicillin and 100uM IPTG) of specific RNAi strains and the control HT115 strain were seeded onto RNAi feeding plates and cultured at room temperature until adulthood. Synchronized L1 worms were treated RNAi by feeding (Ahringer, Reverse genetics, WormBook 2006) for the first generation and allowed to grow to maturity. The worms were then bleached and hatched in M9 buffer for 18hr. The synchronized L1 worms were then seeded on the indicated feeding plate.

Western blot

To measure the level of p-PMK-1, worms (feeding different food for 4h) were analyzed by standard western blot methods and probed with anti-p38 (dilution = 1:5,000; Cell Signaling, 9212S), anti-p-p38 (dilution = 1:5,000; Cell Signaling, 4511S) and anti-α-tubulin (dilution = 1:10,000; Sigma T5168) as a loading control.

Total content of ascorbic acid (TAA) assay

The total content of ascorbic acid was measured using the kits (Beijing Biotech-Pack-analytical Scientific Co., Ltd., Beijing, China, BKWB132 http://biotech-pack-analytical.foodmate.net/) according to the manufacturer's protocol. Briefly, L1 worms were seeded on the different feeding assay plate and cultured for 4 hours. The worms were then lysed in ice-cold conditions using lysis buffer. Equal amounts of protein were used for the normalization. Here is formula for getting TAA concentration

$\text{TAA(ug/mg prot)} = ((\triangle A - a) \div b)) \div (\text{Cpr} \times V1) \times D$

V1 - the volume of supernatant of for experiment

Cpr – the concentration of supernatant protein

D – Dilution ratio of supernatant

a – the intercept of standard curve

b – the slope of standard curve

the standard curve y = 0.0611x + 0.0003 for Figure 4C; y = 0.0258x + 0.0066

for Figure 4B, S4A and S4B.

Preparation of samples for RNA sequencing

835 RNA-seq was done with three biological replicates that were independently

836 generated, collected, and processed. Adult wild type (N2) worms were

837 bleached and then the eggs were incubated in M9 for 18 hours to obtain

838 synchronized L1 worms. L1 worms were cultured in the NGM plate with *HK-E.*

839 *coli* or *E. coli* for 4hrs at 20℃. L1 worms were then collected for sequencing.

RNA sequencing and data processing

For the RNA sequencing assay, cDNA libraries were constructed, and single-

end libraries were sequenced using the Illumina platform (Novogene, Beijing,

China). HISAT2 [54] was used to map the clean reads to the reference gene

sequence (Species: Caenorhabditis_elegans; Source: NCBI; Reference

Genome Version: GCF_000002985.6_WBcel235), and then "featureCounts"

tool in subread software [55] was used to calculate the gene expression level of

each sample. Read counts were inputted into DESeq2 [56]to calculate differential

gene expression and statistical significance. Differentially expressed genes

(DEGs) were screened using following criteria: |log2(FoldChange)| > 1 &

padj<= 0.05.

Preparation of samples for Metabolome sequencing

Metabolome-seq of bacterial was done with four biological replicates that were independently generated, collected, and processed. Total of three group E. coli sample including: *E. coli K12* (Con), HK-*E. coli K12* (HK), and HK-*E. coli yfbR* mutant (HY). All bacteria were overnight cultured to the same OD (OD600=1). *E. coli K12* and *E. coli yfbR* mutant are heat-killed (80, 180min), *E. coli* K12, HK-*E. coli* K12 and HK-*E. coli yfbR* mutant were then spread out NGM plate for 72hs at room temperature. Finally, sample was collected into 1.5ml tube by using sterile cell scraping.

Metabolome sequencing and data processing

Metabolome were sequenced using the Ultra Performance Liquid Chromatography (UPLC) (ExionLC AD，https://sciex.com.cn/) and Quadrupole-Time of Flight (TripleTOF 6600, AB SCIEX) for Non-targeted; Ultra Performance Liquid Chromatography (UPLC) (ExionLC AD，https://sciex.com.cn/) and Tandem mass spectrometry（MS/MS）（QTRAP®，https://sciex.com/）for Broad targeting (Metware, Wuhan, China). Multiple reaction monitoring (MRM) was used to calculate the expression level of each metabolite. Differential metabolites were screened through Fold change ≥ 2 or Fold change ≤ 0.5 and VIP ≥ 1 (Variable Importance in Projection of OPLS-DA model).

Microscopy

Analysis of fluorescence was performed with an Olympus BX53 microscope or Olympus MVX10 dissecting with a DP80 camera.

QUANTIFICATION AND STATISTICAL ANALYSIS

Quantification

ImageJ software was used for quantifying fluorescence intensity of UPR^{ER} and Innate immunity reporter. ImageJ software was used for counting the number of worms about selection and aversion behavior.

Statistical analysis

All statistical analyses were preformed using Student's t-test in Graphpad prism 8.0. Two-tailed unpaired t test was used for statistical analysis of two groups of samples. Data are presented as Mean±SD, and $p < 0.05$ was considered a significant difference, "*" represents $p < 0.05$, "**" represents $p < 0.01$, "***" is represents < 0.001, "****" represents $p < 0.0001$, "ns" represents no significant difference. For all figures, "n" represents the number of worms scored from at least three independent experiments.

Reference:

1. Filosa, A., Barker, A.J., Dal Maschio, M., and Baier, H. (2016). Feeding State Modulates Behavioral Choice and Processing of Prey Stimuli in the Zebrafish Tectum. Neuron *90*, 596-608. 10.1016/j.neuron.2016.03.014.
2. Florsheim, E.B., Sullivan, Z.A., Khoury-Hanold, W., and Medzhitov, R. (2021). Food allergy as a biological food quality control system. Cell *184*, 1440-1454. 10.1016/j.cell.2020.12.007.
3. McLachlan, I.G., Kramer, T.S., Dua, M., DiLoreto, E.M., Gomes, M.A., Dag, U., Srinivasan, J., and Flavell, S.W. (2022). Diverse states and stimuli tune olfactory receptor expression levels to modulate food-seeking behavior. Elife *11*. 10.7554/eLife.79557.
4. Avery, J.A., Liu, A.G., Ingeholm, J.E., Gotts, S.J., and Martin, A. (2021). Viewing images of foods evokes taste quality-specific activity in gustatory insular cortex. Proc Natl Acad Sci U S A *118*. 10.1073/pnas.2010932118.
5. Melin, A.D., Nevo, O., Shirasu, M., Williamson, R.E., Garrett, E.C., Endo, M., Sakurai, K., Matsushita, Y., Touhara, K., and Kawamura, S. (2019). Fruit scent and observer colour vision shape food-selection strategies in wild capuchin monkeys. Nat Commun *10*, 2407. 10.1038/s41467-019-10250-9.
6. Bargmann, C.I. (2006). Chemosensation in C. elegans. WormBook, 1-29. 10.1895/wormbook.1.123.1.
7. Sengupta, P., Chou, J.H., and Bargmann, C.I. (1996). odr-10 encodes a seven transmembrane domain olfactory receptor required for responses to the odorant diacetyl. Cell *84*, 899-909. 10.1016/s0092-8674(00)81068-5.
8. Troemel, E.R., Kimmel, B.E., and Bargmann, C.I. (1997). Reprogramming chemotaxis responses: sensory neurons define olfactory preferences in C. elegans. Cell *91*, 161-169. 10.1016/s0092-8674(00)80399-2.
9. Fiala, A. (2007). Olfaction and olfactory learning in Drosophila: recent progress. Curr Opin Neurobiol *17*, 720-726. 10.1016/j.conb.2007.11.009.
10. Chalasani, S.H., Chronis, N., Tsunozaki, M., Gray, J.M., Ramot, D., Goodman, M.B., and Bargmann, C.I. (2007). Dissecting a circuit for olfactory behaviour in

Caenorhabditis elegans. Nature *450*, 63-70. 10.1038/nature06292.

11.   Zhang, B., Jun, H., Wu, J., Liu, J., and Xu, X.Z.S. (2021). Olfactory perception of
food abundance regulates dietary restriction-mediated longevity via a brain-
to-gut signal. Nat Aging *1*, 255-268. 10.1038/s43587-021-00039-1.

12.   Ha, H.I., Hendricks, M., Shen, Y., Gabel, C.V., Fang-Yen, C., Qin, Y., Colon-
Ramos, D., Shen, K., Samuel, A.D., and Zhang, Y. (2010). Functional
organization of a neural network for aversive olfactory learning in
Caenorhabditis elegans. Neuron *68*, 1173-1186.
10.1016/j.neuron.2010.11.025.

13.   Scott, K. (2018). Gustatory Processing in Drosophila melanogaster. Annu Rev
Entomol *63*, 15-30. 10.1146/annurev-ento-020117-043331.

14.   Hukema, R.K., Rademakers, S., Dekkers, M.P., Burghoorn, J., and Jansen, G.
(2006). Antagonistic sensory cues generate gustatory plasticity in
Caenorhabditis elegans. Embo J *25*, 312-322. 10.1038/sj.emboj.7600940.

15.   Ni, L., Bronk, P., Chang, E.C., Lowell, A.M., Flam, J.O., Panzano, V.C., Theobald,
D.L., Griffith, L.C., and Garrity, P.A. (2013). A gustatory receptor paralogue
controls rapid warmth avoidance in Drosophila. Nature *500*, 580-584.
10.1038/nature12390.

16.   Jones, D., and Candido, E.P.M. (1999). Feeding is inhibited by sublethal
concentrations of toxicants and by heat stress in the nematode
Caenorhabditis elegans: Relationship to the cellular stress response. J Exp
Zool *284*, 147-157. Doi 10.1002/(Sici)1097-010x(19990701)284:2<147::Aid-
Jez4>3.3.Co;2-Q.

17.   Xie, Z., Zhang, X., Zhao, M., Huo, L., Huang, M., Li, D., Zhang, S., Cheng, X., Gu,
H., Zhang, C., et al. (2022). The gut-to-brain axis for toxin-induced defensive
responses. Cell *185*, 4298-4316 e4221. 10.1016/j.cell.2022.10.001.

18.   Richardson, C.E., Kooistra, T., and Kim, D.H. (2010). An essential role for XBP-1
in host protection against immune activation in C. elegans. Nature *463*, 1092-
U1114. 10.1038/nature08762.

19.   Kim, D.H., Feinbaum, R., Alloing, G., Emerson, F.E., Garsin, D.A., Inoue, H.,
Tanaka-Hino, M., Hisamoto, N., Matsumoto, K., Tan, M.W., and Ausubel, F.M.
(2002). A conserved p38 MAP kinase pathway in Caenorhabditis elegans
innate immunity. Science *297*, 623-626. 10.1126/science.1073759.

20.   Carr, A.C., and Maggini, S. (2017). Vitamin C and Immune Function. Nutrients
9. 10.3390/nu9111211.

21.   Steinert, R.E., Lee, Y.K., and Sybesma, W. (2020). Vitamins for the Gut
Microbiome. Trends Mol Med *26*, 137-140. 10.1016/j.molmed.2019.11.005.

22.   Li, Y., and Schellhorn, H.E. (2007). New developments and novel therapeutic
perspectives for vitamin C. J Nutr *137*, 2171-2184. 10.1093/jn/137.10.2171.

23.   Carr, A.C., and Frei, B. (1999). Toward a new recommended dietary allowance
for vitamin C based on antioxidant and health effects in humans. Am J Clin
Nutr *69*, 1086-1107. 10.1093/ajcn/69.6.1086.

24.   Qi, B., Kniazeva, M., and Han, M. (2017). A vitamin-B2-sensing mechanism
that regulates gut protease activity to impact animal's food behavior and

967   growth. Elife *6*. 10.7554/eLife.26243.

968   25.   Qi, B., and Han, M. (2018). Microbial Siderophore Enterobactin Promotes
969   Mitochondrial Iron Uptake and Development of the Host via Interaction with
970   ATP Synthase. Cell *175*, 571-582.e511. 10.1016/j.cell.2018.07.032.

971   26.   Nakad, R., Snoek, L.B., Yang, W., Ellendt, S., Schneider, F., Mohr, T.G., Rösingh,
972   L., Masche, A.C., Rosenstiel, P.C., Dierking, K., et al. (2016). Contrasting
973   invertebrate immune defense behaviors caused by a single gene, the
974   Caenorhabditis elegans neuropeptide receptor gene npr-1. BMC Genomics
975   *17*, 280. 10.1186/s12864-016-2603-8.

976   27.   Sinha, A., Rae, R., Iatsenko, I., and Sommer, R.J. (2012). System wide analysis
977   of the evolution of innate immunity in the nematode model species
978   Caenorhabditis elegans and Pristionchus pacificus. PLoS One *7*, e44255.
979   10.1371/journal.pone.0044255.

980   28.   Hetz, C., Zhang, K., and Kaufman, R.J. (2020). Mechanisms, regulation and
981   functions of the unfolded protein response. Nat Rev Mol Cell Biol *21*, 421-
982   438. 10.1038/s41580-020-0250-z.

983   29.   Ron, D., and Walter, P. (2007). Signal integration in the endoplasmic reticulum
984   unfolded protein response. Nat Rev Mol Cell Biol *8*, 519-529.
985   10.1038/nrm2199.

986   30.   Fletcher, M., Tillman, E.J., Butty, V.L., Levine, S.S., and Kim, D.H. (2019). Global
987   transcriptional regulation of innate immunity by ATF-7 in C. elegans. Plos
988   Genet *15*. ARTN e1007830

989 10.1371/journal.pgen.1007830.

990   31.   Foster, K.J., McEwan, D.L., and Pukkila-Worley, R. (2020). Measurements of
991   Innate Immune Function in C. elegans. Methods Mol Biol *2144*, 145-160.
992   10.1007/978-1-0716-0592-9_13.

993   32.   Weiss, B. (2007). The Deoxycytidine Pathway for Thymidylate Synthesis in
994   Escherichia coli. Journal of Bacteriology *189*, 7922-7926. 10.1128/jb.00461-07.

995   33.   Franceus, J., and Desmet, T. (2020). Sucrose Phosphorylase and Related
996   Enzymes in Glycoside Hydrolase Family 13: Discovery, Application and
997   Engineering. International Journal of Molecular Sciences *21*.
998   10.3390/ijms21072526.

999   34.   Xing, S., Zhang, L., Lin, H., Mao, Z., Bao, K., Hao, P., Pei, Z., Li, J., and Hu, Z.
1000   (2019). Lactose induced redox-dependent senescence and activated Nrf2
1001   pathway. Int J Clin Exp Pathol *12*, 2034-2045.

1002   35.   Patananan, A.N., Budenholzer, L.M., Pedraza, M.E., Torres, E.R., Adler, L.N.,
1003   and Clarke, S.G. (2015). The invertebrate Caenorhabditis elegans
1004   biosynthesizes ascorbate. Arch Biochem Biophys *569*, 32-44.
1005   10.1016/j.abb.2015.02.002.

1006   36.   Yabuta, Y., Nagata, R., Aoki, Y., Kariya, A., Wada, K., Yanagimoto, A., Hara, H.,
1007   Bito, T., Okamoto, N., Yoshida, S., et al. (2020). L-Ascorbate Biosynthesis
1008   Involves Carbon Skeleton Rearrangement in the Nematode Caenorhabditis
1009   elegans. Metabolites *10*. 10.3390/metabo10080334.

1010   37.   Moritz, B., Schmitz, A.E., Rodrigues, A.L.S., Dafre, A.L., and Cunha, M.P.

1011 (2020). The role of vitamin C in stress-related disorders. The Journal of
1012 Nutritional Biochemistry *85*. 10.1016/j.jnutbio.2020.108459.

38. Rice, M.E. (2000). Ascorbate regulation and its neuroprotective role in the
1014 brain. Trends in Neurosciences *23*, 209-216. https://doi.org/10.1016/S0166-
1015 2236(99)01543-X.

39. Webb, A.L., and Villamor, E. (2007). Update: Effects of Antioxidant and Non-
1017 Antioxidant Vitamin Supplementation on Immune Function. Nutrition
1018 Reviews *65*, 181-217. 10.1111/j.1753-4887.2007.tb00298.x.

40. Maggini, S., Wintergerst, E.S., Beveridge, S., and Hornig, D.H. (2007). Selected
1020 vitamins and trace elements support immune function by strengthening
1021 epithelial barriers and cellular and humoral immune responses. British
1022 Journal of Nutrition *98*, S29-S35. 10.1017/s0007114507832971.

41. Luo, X., Ng, C., He, J., Yang, M., Luo, X., Herbert, T.P., and Whitehead, J.P.
1024 (2022). Vitamin C protects against hypoxia, inflammation, and ER stress in
1025 primary human preadipocytes and adipocytes. Molecular and Cellular
1026 Endocrinology *556*. 10.1016/j.mce.2022.111740.

42. Su, M., Liang, X., Xu, X., Wu, X., and Yang, B. (2019). Hepatoprotective
1028 benefits of vitamin C against perfluorooctane sulfonate-induced liver damage
1029 in mice through suppressing inflammatory reaction and ER stress.
1030 Environmental Toxicology and Pharmacology *65*, 60-65.
1031 10.1016/j.etap.2018.12.004.

43. Ozbey, N.P., Thompson, M.A., and Taylor, R.C. (2021). The regulation of animal
1033 behavior by cellular stress responses. Exp Cell Res *405*, 112720.
1034 10.1016/j.yexcr.2021.112720.

44. Ozbey, N.P., Imanikia, S., Krueger, C., Hardege, I., Morud, J., Sheng, M.,
1036 Schafer, W.R., Casanueva, M.O., and Taylor, R.C. (2020). Tyramine Acts
1037 Downstream of Neuronal XBP-1s to Coordinate Inter-tissue UPR(ER)
1038 Activation and Behavior in C. elegans. Dev Cell *55*, 754-770 e756.
1039 10.1016/j.devcel.2020.10.024.

45. Troemel, E.R., Chu, S.W., Reinke, V., Lee, S.S., Ausubel, F.M., and Kim, D.H.
1041 (2006). p38 MAPK regulates expression of immune response genes and
1042 contributes to longevity in C. elegans. Plos Genet *2*, e183.
1043 10.1371/journal.pgen.0020183.

46. Mouysset, J., Kahler, C., and Hoppe, T. (2006). A conserved role of
1045 Caenorhabditis elegans CDC-48 in ER-associated protein degradation. J Struct
1046 Biol *156*, 41-49. 10.1016/j.jsb.2006.02.015.

47. Sasagawa, Y., Otani, M., Higashitani, N., Higashitani, A., Sato, K., Ogura, T., and
1048 Yamanaka, K. (2009). Caenorhabditis elegans p97 controls germline-specific
1049 sex determination by controlling the TRA-1 level in a CUL-2-dependent
1050 manner. Journal of Cell Science *122*, 3663-3672. 10.1242/jcs.052415.

48. Kim, K., and Li, C. (2004). Expression and regulation of an FMRFamide-related
1052 neuropeptide gene family in Caenorhabditis elegans. J Comp Neurol *475*, 540-
1053 550. 10.1002/cne.20189.

49. Melo, J.A., and Ruvkun, G. (2012). Inactivation of conserved C. elegans genes

engages pathogen- and xenobiotic-associated defenses. Cell *149*, 452-466.
10.1016/j.cell.2012.02.050.

50.   Geng, S., Li, Q., Zhou, X., Zheng, J., Liu, H., Zeng, J., Yang, R., Fu, H., Hao, F.,
Feng, Q., and Qi, B. (2022). Gut commensal E. coli outer membrane proteins
activate the host food digestive system through neural-immune
communication. Cell Host Microbe. 10.1016/j.chom.2022.08.004.

51.   Richardson, C.E., Kinkel, S., and Kim, D.H. (2011). Physiological IRE-1-XBP-1
and PEK-1 signaling in Caenorhabditis elegans larval development and
immunity. PLoS Genet *7*, e1002391. 10.1371/journal.pgen.1002391.

52.   Kamath, R.S., Fraser, A.G., Dong, Y., Poulin, G., Durbin, R., Gotta, M., Kanapin,
A., Le Bot, N., Moreno, S., Sohrmann, M., et al. (2003). Systematic functional
analysis of the Caenorhabditis elegans genome using RNAi. Nature *421*, 231-
237. 10.1038/nature01278.

53.   Rual, J.F., Ceron, J., Koreth, J., Hao, T., Nicot, A.S., Hirozane-Kishikawa, T.,
Vandenhaute, J., Orkin, S.H., Hill, D.E., van den Heuvel, S., and Vidal, M.
(2004). Toward improving Caenorhabditis elegans phenome mapping with an
ORFeome-based RNAi library. Genome Res *14*, 2162-2168.
10.1101/gr.2505604.

54.   Mortazavi, A., Williams, B.A., McCue, K., Schaeffer, L., and Wold, B. (2008).
Mapping and quantifying mammalian transcriptomes by RNA-Seq. Nat
Methods *5*, 621-628. 10.1038/nmeth.1226.

55.   Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general
purpose program for assigning sequence reads to genomic features.
Bioinformatics *30*, 923-930. 10.1093/bioinformatics/btt656.

56.   Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold
change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*. ARTN
550
10.1186/s13059-014-0550-8.

1083

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S1. Food selection assay of animals fed HK-*E. coli* or *E. coli*. Relative to Figure 1.

(A-E) Metabolomics analysis of different quality food (HK-K12, HK-*yfbR* and Live-K12). Cluster analysis of all metabolites (A), lipids and their derivatives (B), amino acids and their derivatives (C), coenzymes and vitamins (D), and carbohydrates and their derivatives (E) from Live-K12, HK-K12, and HK-yfbR. Color indicates the relative level of each metabolite. HK-K12: heat-killed *E. coli* wild-type K12; HK-*yfbR:* heat-killed *E. coli mutant yfbR;* K12: live *E. coli* wild-type K12. z-score for standardizing data, complete for bi-clustering algorithm, and Euclidean for distance method.

(A) Principal component analysis to test the repeatability of the metabolic experiment. HK-K12: heat-killed *E. coli* wild-type K12; HK-yfbR: heat-killed *E. coli* mutant yfbR; K12: live *E. coli* wild-type K12. LB: LB buffer for culturing *E. coli.*

(B) GO enrichment analysis of UPR^{ER} dependent IRE-1 branch.

(C) The number of worms in each position calculated at the indicated time, indicating that animals initially select both foods (1-2h), but eventually favor high-quality food (Live E. coli) until 17h.

(D) UPR^{mit} Reporter (*Phsp-6:GFP*) was weakly induced in animals fed with HK-E. coli

(E) UPR^{ER} Reporter (Phsp-4:GFP) expression in animals under normal food (OP50), low quality food (HK-OP50), and starved (M9: hatching L1 worm in M9, NGM) condition;

(F) Immunity Reporter (Pirg-5:GFP) expression in animals under normal food (OP50), low quality food (HK-OP50), and starved (M9: hatching L1 worm in M9, NGM).

For all panels, Scale bar shows on indicated figures, 50 μm. * p<0.05, ** p<0.01, *** p < 0.001, **** p<0.0001, ns: no significant difference. Precise P values are provided in Raw Data.

Figure S2

Figure S2. UPRER and Innate immunity pathway in animals are critical for evaluating HK-*E. coli*. Relative to Figure 2.

(A-B) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Phsp-4::GFP* was not affected in animals with *atf-4* (A) or *atf-6* (B) RNAi treatment. n is the number of worms scored from at least three independent experiments. Data are represented as mean $\pm$ SD.

(C) qPCR showing that IRE-1-mediated splicing of *xbp-1* mRNA is induced in animals fed with *HK-E. coli.*

(D) Venn diagram showing the numbers of PMK-1 dependent genes [30] and up-regulated genes in animals fed HK-*E. coli,* and their overlap.

(E) The expression of PMK-1 dependent genes which was extracted from RNA-seq data from animals fed with *HK-E. coli*. The data from average of three independent experiments (N=3)

(F-G) GFP fluorescence images and bar graph showing that *Psysm-1::GFP* (F) and *Pirg-1::GFP* (G) were induced in animals fed *HK-E. coli.* n is the number of worms scored from at least three independent experiments. Data are represented as mean $\pm$ SD.

(H) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Pirg-5::GFP* was decreased in animals with *ire-1* or *xbp-1* RNAi treatment. n is the number of worms scored from at least three independent experiments. Data are represented as mean $\pm$ SD.

(I) GFP fluorescence images and bar graph showing that *Phsp-4::GFP* was not affected in animals with *pmk-1* RNAi treatment. n is the number of worms scored from at least three independent experiments. Data are represented as mean $\pm$ SD.

(J) UPR[ER] Reporter (Phsp-4::GFP) expression in animals with candidate RNAi feeding OP50 or HK-OP50

(K) immunity Reporter (Pirg-5::GFP) expression in animals with candidate RNAi feeding OP50 or HK-OP50

For all panels, Scale bar shows on indicated figures, 50 µm. * p<0.05, ** p<0.01, *** p < 0.001, **** p<0.0001, ns: no significant difference. Precise P values are provided in Raw Data.

Figure S3

Figure S3. Low sugar food, HK-*E. coli*, induce stress response and avoidance behavior in animals. Relative to Figure 3.

(A) Flow chart of strategy for *E. coli* keio mutant screening. We identified 20 *E. coli* mutants that did not induce *hsp-4::GFP* through the UPR^{ER} reporter (*irg-5::GFP*) after three rounds of screening (Table S3). From these 20 E. coli mutant, we identified 9 *E. coli* mutants that did not induce *irg-5::GFP* through the immunity reporter (*irg-5::GFP*) screening (Table S3).

(B-C) The bar graph showing that HK-*E. coli* induced *Phsp-4::GFP* (B) and *Pirg-5::GFP* (C) was decreased in animals fed mutant *E. coli* (Heat-killed). n is the number of worms scored from at least three independent experiments. Data are represented as mean $\pm$ SD.

(D) Venn diagram showing the number of deferentially metabolites in HK-*E. coli*-K12, HK-*E. coli-yfbR* and *E. coli*.

(E) KEGG enrichment analysis of deferentially expressed genes in animals fed HK-*E. coli* vs live *E. coli*. We noticed that all most of glycolysis/gluconeogenesis genes are up-regulated in animals fed HK-*E. coli*.

(F) The bar graph showing that HK-E. coli induced *Phsp-4::GFP* was not affected in animals with D-(+)-Trehalose supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD. .

(G) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Psysm-1::GFP* was decreased in animals with D-(+)-Glucose (D-Glc) supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(H) UPR^{ER} Reporter (Phsp-4::GFP) expression animals with D-Glc supplementation under OP50, HK-OP50, or NGM condition.

(I) immunity Reporter (Pirg-5::GFP) expression animals with D-Glc supplementation under OP50, HK-OP50, or NGM condition.

For all panels, Scale bar shows on indicated figures, 50 µm. * $p<0.05$, ** $p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P values are provided in Raw Data.

Figure S4

Figure S4. Vitamin C biosynthesis pathway is critically for evaluating low sugar. Relative to Figure 4.

(A) The level of total L-ascorbic acid (TAA) in Live-K12, HK-K12, or HK-yfbR. N is number of independent experiments. Data are represented as mean ±SD from six independent experiments.

(B) The level of total L-ascorbic acid (TAA) in animals fed HK-*E. coli* with or without D-Glc supplementation. Data are represented as mean ±SD from from eight independent experiments.

(C) GFP fluorescence images and bar graph showing that HK-*E. coli* induced *Psysm-1::GFP* was decreased in animals with D-GlcA or Vitamin C supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ± SD.

(D) The bar graph showing that suppression of HK-*E. coli* induced *Psysm-1::GFP* by D-GlcA supplementation was abolished in animals with RNAi of VC biosynthesis genes, which was not affect by Vitamin C supplementation. n is the number of worms scored from at least three independent experiments. Data are represented as mean ±SD.

(E) Food selection assay showing that the preference of HK-*E. coli* with D-GlcA supplementation was abolished in animals by RNAi of Vitamin C biosynthesis genes. Data are represented as mean ±SD from six independent experiments, 427-775 animals/assay.

(F) Food selection assay for OP50 & OP50 + + D-Glc, D-GlcA, or VC, respectively. Data are represented as mean ±SD from five independent experiments, 241-1182 animals/assay.

(G) Food selection assay for buffer & buffer + D-Glc, D-GlcA, or VC, respectively. Data are represented as mean ±SD from five independent experiments, 8-153 animals/assay.

(H) Food avoidance assay for N2 & kgb-1 mutant animals fed with HK-E. coli. Data are represented as mean ±SD from six independent experiments, 348-660 animals/assay.

For all panels, Scale bar shows on indicated figures, 50 µm. * $p<0.05$, ** $p<0.01$, *** $p < 0.001$, **** $p<0.0001$, ns: no significant difference. Precise P values are provided in Raw Data.