Title

FBXO24 deletion causes abnormal accumulation of membraneless electron-dense granules in sperm flagella and male infertility

Yuki Kaneda[1,2], Haruhiko Miyata[1,*], Zoulan Xu[1,2], Keisuke Shimada[1], Maki Kamoshita[1], Tatsuya Nakagawa[1,2], Chihiro Emori[1], Masahito Ikawa[1,2,3,4,5,*]

[1]Research Institute for Microbial Diseases, Osaka University, 3-1 Yamadaoka, Suita, Osaka 5650871 JAPAN.
[2]Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 5650871 JAPAN.
[3]The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 1088639 JAPAN.
[4]Center for Infectious Disease Education and Research (CiDER), Osaka University, 2-8 Yamadaoka, Suita, Osaka 5650871 JAPAN.
[5]Center for Advanced Modalities and DDS (CAMaD), Osaka University, 2-8 Yamadaoka, Suita, Osaka 5650871 JAPAN.
*Corresponding authors: Haruhiko Miyata and Masahito Ikawa

Contact info
Email: hmiya003@biken.osaka-u.ac.jp (HM); ikawa@biken.osaka-u.ac.jp (MI)

Abstract

Ribonucleoprotein (RNP) granules are membraneless electron-dense structures rich in RNAs and proteins, and involved in various cellular processes. Two RNP granules in male germ cells, intermitochondrial cement and the chromatoid body (CB), are associated with PIWI-interacting RNAs (piRNAs) and are required for transposon silencing and spermatogenesis. Other RNP granules in male germ cells, the reticulated body and CB remnants, are also essential for spermiogenesis. In this study, we disrupted FBXO24, a testis-enriched F-box protein, in mice and found numerous membraneless electron-dense granules accumulated in sperm flagella. *Fbxo24* knockout (KO) mice exhibited malformed flagellar structures, impaired sperm motility, and male infertility, likely due to the accumulation of abnormal granules. The amount and localization of known RNP granule-related proteins were not disrupted in *Fbxo24* KO mice, suggesting that the accumulated granules were distinct from known RNP granules. Further studies revealed that RNAs and two importins, IPO5 and KPNB1, abnormally accumulated in *Fbxo24* KO spermatozoa and that FBXO24 could ubiquitinate IPO5. In addition, IPO5 and KPNB1 were recruited to stress granules, RNP complexes, when cells were treated with oxidative stress or a proteasome inhibitor. These results suggest that FBXO24 is involved in the degradation of IPO5, disruption of which may lead to the accumulation of abnormal RNP granules in sperm flagella.

Keywords

CRISPR/Cas9; Spermiogenesis; Ribonucleoprotein granules

Introduction

Spermatogenesis is a specialized process by which spermatogonia differentiate into spermatocytes, round spermatids, elongating spermatids, and spermatozoa by undergoing meiosis and subsequent spermiogenesis. During spermiogenesis, round spermatids undergo dramatic morphological changes such as nuclear condensation, acrosome formation, flagellum formation, and cytoplasmic removal, leading to the formation of spermatozoa (Leblond and Clermont, 1952). To fertilize eggs, spermatozoa then travel a long distance to reach eggs and pass through the cumulus-cell layer and zona pellucida that encase eggs. The motility apparatus of spermatozoa is the flagellum which is divided into three parts, a midpiece, a principal piece, and an endpiece, in order from proximal to distal (Eddy et al., 2003). The midpiece contains accessory structures called the mitochondrial sheath and outer dense fibers (ODF) that surround the axoneme, a 9+2 arrangement of microtubules, while the principal piece contains a different accessory structure called the fibrous sheath and ODFs surrounding the axoneme. In contrast, the endpiece contains no accessory structures but the axoneme. Abnormal formation of these flagellar structures could lead to male infertility (Nsota Mbango et al., 2019).

Ribonucleoprotein (RNP) granules are membraneless electron-dense structures that assemble through liquid–liquid phase separation and play critical roles in spermatogenesis (Eddy, 1974). Among the RNP granules, intermitochondrial cement (IMC) observed in late spermatocytes and the chromatoid body (CB) found in round spermatids are involved in the PIWI-interacting RNA (piRNA) pathway (Meikar et al., 2011). After meiosis, IMCs are no longer detectable because of mitochondrial dispersion, and CBs become gradually larger in round spermatids. In elongating spermatids, the CBs containing MIWI, the main effector protein of the piRNA pathway, disappear and two other RNP structures called a reticulated body and a CB remnant that contain TSKS, TSSK1, and TSSK2 appear (Shang et al., 2010). Recently, we found that *Tsks* knockout (KO) mice failed to generate reticulated bodies and CB remnants, and were infertile with impaired cytoplasmic removal (Shimada et al., 2023), indicating that the formation and disassembly of RNP granules is highly regulated and important for spermiogenesis.

During spermiogenesis, post-translational modifications of proteins are critical since gene transcription is thought to cease with nuclear condensation. Ubiquitination is one of the post-translational modifications in which a small protein called ubiquitin is attached to target proteins. Ubiquitinated proteins are degraded by a ubiquitin-proteasome system (UPS), which is involved

in various physiological phenomena in cells, such as cell cycle regulation and signal transduction
(Ciechanover and Schwartz, 1998; Mani and Gelmann, 2005; Zheng et al., 2016). Protein
ubiquitination is carried out by a cascade of enzymatic reactions, including ubiquitin-activating
enzyme (E1), ubiquitin-binding enzyme (E2), and ubiquitin ligase (E3), and target proteins
ubiquitinated by E3 ligase are degraded by a large protein complex called the 26S proteasome.
Among the approximately 600 E3 ligase genes estimated to be present in humans (Li et al.,
2008), the mechanism and functions of ubiquitination mediated by the SCF (Skp1-Cul1-F-box
protein) complex has been well studied (Skaar et al., 2013). The SCF complex consists of several
proteins such as SKP1, a cullin protein, an F-box protein, and RBX1, and can ubiquitinate various
substrates by using different F-box proteins that recognize distinct substrates. In humans, at least
38 F-box proteins have been identified, but the functions of most F-box proteins remain unclear
(Kipreos and Pagano, 2000). In this study, we analyzed the function of *Fbxo24* (F-box protein 24),
predominantly expressed in testes, and found that FXBO24 is essential for sperm flagellar
formation and male fertility.

Results
87

***Fbxo24* is expressed predominantly in male germ cells after meiosis and interacts with**
SKP1

We performed RT-PCR using cDNAs obtained from multiple mouse tissues and found a specific
band for *Fbxo24* in testes (Figure 1A). To identify stages in which *Fbxo24* was expressed in
spermatogenesis, we performed RT-PCR using postnatal mouse testes over time. *Fbxo24*
signals increased after birth, with solid signals observed around postnatal day 21 (Figure 1B)
when round spermatids appear (Kluin et al., 1982). Consistent with the RT-PCR results, single-
cell transcriptome data (Hermann et al., 2018) indicates that mouse *Fbxo24* and its human
orthologue, *FBXO24*, are expressed during the post-meiotic stages (Figures S1A and B).
Pairwise sequence alignment of amino acids sequences revealed that FBXO24 is highly
conserved between mice and humans (88% amino acid identity) (Figure S1C) with the F-box
domains located near the N-terminus (Figure S1D). Because it has been reported that F-box
proteins bind to SKP1 via the F-box domain to form the SCF complex (Bai et al., 1996; Schulman
et al., 2000), we investigated the interaction of FBXO24 with SKP1 by co-immunoprecipitation
using HEK293T cells. We constructed vectors that express FBXO24 tagged with 3×FLAG with
(wild-type, WT) or without (ΔF) the F-box domain (Figure 1C). When SKP1-1D4 was
immunoprecipitated with an anti-1D4 antibody, WT FBXO24-FLAG was co-immunoprecipitated,
whereas ΔF FBXO24-FLAG was not (Figure 1D left). Conversely, when an anti-FLAG antibody
was used to immunoprecipitate WT or ΔF FBXO24-FLAG, SKP1-1D4 was co-immunoprecipitated
only with WT FBXO24-FLAG (Figure 1D right). These results suggest that FBXO24 could function
as a component of the SCF complex in mouse testis, during the post meiotic stages of
spermatogenesis.

Lack of *Fbxo24* causes male sterility in mice

To examine the role of *Fbxo24* in spermatogenesis, we generated *Fbxo24* KO mice using the
CRISPR/Cas9 system. Two gRNAs were designed to delete the majority of the coding region of
Fbxo24 (Figure 1E). One hundred fertilized eggs were electroporated, and the resulting 89 two-
cell stage embryos were transferred to the oviducts of pseudopregnant ICR mice. One out of 18
born pups contained the large deletion of the coding region and subsequent mating was
performed to obtain *Fbxo24* KO mice. We found no overt gross defects in development, behavior,
or survival in *Fbxo24* KO mice. We performed genomic PCR using the primers described in
Figure 1E (Figure 1F) and confirmed that *Fbxo24* KO mice had a deletion of 11,575 bp by Sanger
sequencing (Figure 1G). *Fbxo24* KO male mice were then mated with three WT females for three
months to analyze fertility. Although 20 vaginal plugs were detected, no pups were born from the
Fbxo24 KO male mice (Figure 1H), suggesting that *Fbxo24* is indispensable for male fertility.

Disruption of *Fbxo24* results in impaired spermiation and abnormal sperm morphology

We first examined spermatogenesis to determine the cause of male infertility in *Fbxo24* KO mice.
No apparent differences in gross testicular morphology (Figure S2A) or weights (Figure S2B)
were found between controls and *Fbxo24* KO mice. We then performed He-PAS staining of
testicular and epididymis sections (Figures 2A and S2C). Although elongating spermatids can be
found in *Fbxo24* KO testes, step 16 spermatids were still present in Stage IX seminiferous
tubules (Figure 2A), indicating that spermiation was impaired in *Fbxo24* KO testes. In contrast,
we could not find apparent differences in the cross sections of the cauda epididymis between the
two genotypes (Figure S2C). Next, we observed mature spermatozoa collected from the cauda
epididymis. While KO spermatozoa showed comparable head morphology to the controls, KO
spermatozoa exhibited abnormal tail structures such as bent or coiled flagella (Figures 2B and C).
Further analyses of sperm head morphology with immunostaining showed no overt abnormalities
in the acrosome or nucleus (Figure S2D). Since abnormal flagellar morphology could result in
decreased motility, we performed computer-assisted sperm analysis (CASA) after 10 and 120
minutes of incubation in capacitating medium. CASA revealed that the percentages of motile
spermatozoa were significantly reduced in *Fbxo24* KO mice compared to the controls (Figure 2D
and Movies S1 and S2). Furthermore, all velocity parameters such as average path velocity
(VAP), straight-line velocity (VSL), and curvilinear velocity (VCL) were lower in *Fbxo24* KO
spermatozoa (Figures 2E-G). These results indicate that FBXO24 is critical in sperm flagellum
formation during spermiogenesis.

***Fbxo24* KO spermatozoa cannot fertilize eggs *in vitro* and fail to migrate from the uterus into the oviduct**

To further evaluate the fertilizing ability of *Fbxo24* KO spermatozoa, we performed in vitro
fertilization (IVF). Consistent with male sterility observed in vivo (Figure 1H), *Fbxo24* KO
spermatozoa could not fertilize eggs in vitro (Figure 3A). Furthermore, removing cumulus cells
(Figure 3B) or both cumulus cells and the zona pellucida (ZP) (Figure 3C) could not rescue the
fertilization rates, suggesting that *Fbxo24* KO spermatozoa may have defects in not only sperm
motility but also the acrosome reaction, which is a prerequisite for spermatozoa to pass through
the ZP and fuse with eggs (Morohoshi et al., 2023). Therefore, we crossed *Fbxo24* mutant mice
with transgenic (Tg) mice which express EGFP in the acrosome and DsRed2 in the mitochondria
(Red Body Green Sperm, RBGS, mice) (Hasuwa et al., 2010) and analyzed the acrosome
reaction rates. RBGS spermatozoa lose EGFP fluorescent signal when the spermatozoa undergo
the acrosome reaction. First, to test whether *Fbxo24* KO spermatozoa were viable, we performed
propidium iodide (PI) staining and found that the percentages of dead spermatozoa were
significantly higher in *Fbxo24* KO mice (Figure S3A). We then analyzed the acrosome reaction
rates of PI-negative live spermatozoa. While the control spermatozoa underwent the acrosome
reaction after 120 minutes of incubation in a capacitation medium and with Ca^{2+} ionophore
(A23187) treatment, *Fbxo24* KO spermatozoa rarely underwent the acrosome reaction even after
adding A23187 (Figure S3B). Since the SNARE complex is considered critical for the acrosome
reaction (Hutt et al., 2005; Katafuchi et al., 2000; Schulz et al., 1997; Tomes et al., 2002), we
examined the amounts of SNARE-related proteins in mature spermatozoa using mass
spectrometry and immunoblotting (Figure S3C and D), but no notable differences were found in
SNARE-related proteins between the control and *Fbxo24* KO spermatozoa. Further, no significant
differences were found in PLCD4 that has been reported to be important for the acrosome
reaction (Fukami et al., 2001).

Next, we observed sperm migration in the female reproductive tract using RBGS mice. Although
the control spermatozoa migrate through the uterotubal junction (UTJ) four hours after mating,
Fbxo24 KO spermatozoa hardly passed through the UTJ (Figure 3D). Since the processing of
sperm membrane protein, ADAM3, is necessary for sperm migration through the UTJ (Fujihara
et al., 2019, 2018a; Yamaguchi et al., 2006), we performed immunoblotting analyses and found
that ADAM3 was processed correctly even in *Fbxo24* KO spermatozoa (Figure 3E). We further

performed immunoblotting of LY6K because *Ly6k* KO spermatozoa cannot migrate through the
UTJ even with normal ADAM3 processing (Fujihara et al., 2014). However, we did not detect
significant differences in the amounts of LY6K (Figure S3E). Considering that the acrosome
reaction is not essential for sperm migration in the female reproductive tract (Morohoshi et al.,
2023) and defects in sperm motility could cause impaired migration through the UTJ (Chung et
al., 2014; Fujihara et al., 2018a; Miyata et al., 2015; Shimada et al., 2019), these results suggest
that lower viability and decreased sperm motility can be the cause of abnormal sperm migration
and male sterility in vivo.

Next, we performed intracytoplasmic sperm injection (ICSI) to examine whether nuclei of *Fbxo24*
KO spermatozoa have the potential to generate the next generation. We injected *Fbxo24* KO
sperm heads into 91 WT oocytes, and obtained 48 two-cell stage embryos (Figure 3F). By
transplanting the two-cell embryos into the oviduct of pseudopregnant ICR females, we obtained
four pups, which were confirmed to have the heterozygous mutation by PCR (Figures 3G and H).
These results indicate that ICSI can rescue male sterility of *Fbxo24* KO mice.

FBXO24 is required for the sperm midpiece formation

We analyzed mitochondria localization in *Fbxo24* KO spermatozoa because abnormal
mitochondrial sheath structures were found in spermatozoa with bent tails, such as in *Armc12*
KO, *Tbc1d21* KO, and *Gk2* KO mice (Shimada et al., 2021, 2019). We observed midpieces using
RBGS Tg mice and revealed that mitochondria were disorganized in *Fbxo24* KO spermatozoa
(Figure 4A). We also observed SEPT4, a component of the annulus (Kissel et al., 2005), and
found that SEPT4 localized to the proper region (Figure S4A).

We then used a scanning electron microscope (SEM) to observe the formation of mitochondrial
sheaths. During the early step of mitochondrial sheath formation, mitochondria wrap around the
axoneme. In both control and *Fbxo24* KO spermatids, spherical mitochondria were aligned
correctly at this step (Figure 4B, left panels). In the next step, mitochondria became crescent-
shaped and were interlocked in the control spermatids, whereas irregular interlocking was
observed in *Fbxo24* KO spermatids (Figure 4B, middle panels). Subsequently, elongating
mitochondria remain irregularly arranged in *Fbxo24* KO spermatids (Figure 4B, right panels).

To further investigate the morphological abnormality of the midpiece in *Fbxo24* KO mice, we
observed the ultrastructure of the cauda epididymal spermatozoa using transmission electron
microscopy (TEM). Consistent with the SEM results, we found that mitochondria were irregularly
arranged in *Fbxo24* KO spermatozoa (Figures 4C and D). The frequency of abnormal
mitochondria was significantly higher in *Fbxo24* KO spermatozoa (Figure 4E). In addition, the
axoneme (AX) and outer dense fiber (ODF) were disrupted in both midpieces and principal pieces
of *Fbxo24* KO spermatozoa (Figures S4B-E). Furthermore, we frequently observed numerous
membraneless electron-dense granules in the midpieces of *Fbxo24* KO spermatozoa (Figures
4C, D, and F). We found these granules even in the axoneme (Figure. 4C). Despite the low
frequency, we also found membraneless electron-dense granules in the principal pieces of
Fbxo24 KO spermatozoa (Figures S4F and G).

Because membraneless electron-dense granules accumulated in *Fbxo24* KO spermatozoa, we
examined RNP granule-related proteins in *Fbxo24* KO mice. We performed immunoblotting
analyses for ADAD1, ADAD2, MILI, MIWI, RNF-17, YTHDC2, TSKS, and TSSK1, which localized
in germ-cell RNP granules and are essential for spermatogenesis (Lu et al., 2023; Meikar et al.,
2011; Pan et al., 2005; Shang et al., 2010; Shimada et al., 2023; Snyder et al., 2020), but we did
not see any differences in the amounts of these proteins between *Fbxo24* heterozygous and KO
testes (Figure S5A). Further, we found no abnormalities in the localization of MIWI in *Fbxo24* KO
testes (Figure S5B). MIWI was localized in the IMCs of late spermatocytes and CBs of round
spermatids and disappeared in elongating spermatids (Figure S5B). Further, no abnormalities

were found in the immunostaining of TSKS, which localized in the reticulated body and CB
remnant of elongating spermatids (Figure S5C).

IPO5 and KPNB1 amounts increase in *Fbxo24* KO flagella

Since previous studies show that loss of F-box protein causes the accumulation of substrates
(Nakayama et al., 2000; Tsunematsu et al., 2004), we investigated whether certain proteins
accumulated in *Fbxo24* KO spermatozoa. Mass spectrometry analyses revealed that the amounts
of several proteins significantly increased in *Fbxo24* KO mature spermatozoa (Figure 5A). Among
the significantly increased proteins, we focused on IPO5 (importin 5, also known as KPNB3 and
RanBP5) and KPNB1 (karyopherin subunit beta 1, also known as importin β1) because the fold
changes (KO/WT) are the highest in these two proteins. Both IPO5 and KPNB1 are members of
the karyopherin-β (KPNB) family of nuclear transport receptors (NTRs), which play a crucial role
in nucleocytoplasmic transport (Chook and Blobel, 2001; Chook and Süel, 2011; Conti and
Izaurralde, 2001; Kimura and Imamoto, 2014). In rodent testes, KPNB1 localized to the cytoplasm
of spermatogonia, spermatocytes, and Sertoli cells, and IPO5 localized to the cytoplasm of
elongating spermatids (Loveland et al., 2015, 2006). Immunoblotting analyses confirmed that the
amounts of IPO5 and KPNB1 increased in *Fbxo24* KO mature spermatozoa (Figure 5B). In
contrast, the amounts of KPNA2 (Importin α1), a member of karyopherin-α (KPNA) family of
NTRs which localized to the cytoplasm and nucleus of spermatocytes and the cytoplasm of
elongating spermatids (Miyamoto et al., 2013), were comparable between the control and *Fbxo24*
KO spermatozoa. Since the anti-KPNB1 antibody did not work for immunostaining, we performed
immunostaining of mature spermatozoa using an anti-IPO5 antibody. We detected IPO5 in the
Fbxo24 KO flagella but not in control (Figure 5C).

FBXO24 could interact with IPO5

Next, we investigated the proteins that interacted with FBXO24. Because we could not obtain
anti-FBXO24 antibodies that worked for immunoprecipitation, we generated Tg mice expressing
3×FLAG-tagged *Fbxo24* under a testis-specific *Prm1* promoter (Figures S6A and B). This
transgene could not rescue *Fbxo24* KO sperm morphology (Figure S6C). By looking for long non-
coding RNAs, we found *Gm36266* that is located around Exon 9 and 10 of *Fbxo24*, which is
deleted in *Fbxo24* KO mice (Figure 1E); however, a partial deletion of Exon 2 and 3 of *Fbxo24*
showed the same defects in sperm morphology (Figures S6D–G) as mice lacking Exon 3-10
(Figure 2B), suggesting that *Fbxo24*, not *Gm36266*, is responsible for the phenotypes observed
in this study. Abnormal sperm morphology was not rescued by the transgene, likely due to lower
expression of FBXO24 (Figures 5D and S6B) and/or FLAG-tag interfering with FBXO24 function.
Using Tg mice, we could immunoprecipitate FBXO24-FLAG with an anti-FLAG antibody, and
subsequent mass spectrometry analyses detected not only FBXO24 but also IPO5 with the
highest quantitative values (Figure S6H), indicating that these proteins interact in vivo. We also
found SKP1 in this analysis, consistent with the in vitro study (Figure 1D). We confirmed that
IPO5 co-immunoprecipitated with FBXO24-FLAG in mouse testes using immunoblotting analysis
(Figure 5D). Further, we transiently expressed 3×FLAG-tagged *Fbxo24* in HEK293T cells,
performed immunoprecipitation analysis using an anti-FLAG antibody, and found that FBXO24
could interact with endogenous IPO5 but not with endogenous KPNB1 (Figure 5E). To examine if
FBXO24 can ubiquitinate IPO5, we performed a ubiquitination assay using HEK293T cells and
revealed that ubiquitination of IPO5 was upregulated when cells were transfected with WT
FBXO24, but not with ΔF FBXO24 that lacks the F-box domain (Figure 5F). These results
suggest that FBXO24 recognizes and ubiquitinates IPO5 for subsequent protein degradation.

IPO5 is recruited into RNP granules under stress conditions

The amounts of known RNP granule-related proteins were not upregulated in *Fbxo24* KO testes
(Figure S5A) and spermatozoa (Figure 5A). In contrast, the total amount of RNAs was
significantly increased in *Fbxo24* KO spermatozoa (Figures 6A and B) although the

contamination of somatic cells or immature spermatogenic cells were rarely found (Figure S7).
Previous studies showed that various stresses, such as heat shock, arsenite treatment, or
proteasome inhibition, trigger the formation of cytoplasmic stress granules (SGs), membraneless
granules composed of RNPs. Further, it has been shown that SGs could contain KPNB1 (Chang
and Tarn, 2009; Mahboubi et al., 2013). Increased amounts of KPNB1 and RNAs suggest that
RNP granules may be formed in *Fbxo24* KO spermatozoa. We analyzed whether IPO5 can
localize to SGs under stress conditions to explore this possibility. We examined the subcellular
localization of endogenous IPO5 and KPNB1 in COS7 cells treated with arsenite, an oxidative
stress inducer. Immunostaining analyses revealed that IPO5 was predominantly localized to the
cytoplasm without the stress inducer (Figure 6C); however, when cells were treated with arsenite,
IPO5 was detected not only in the cytoplasm but also in cytoplasmic granules that were
colocalized with KPNB1 (Figure 6C). We then analyzed if the stress of proteasome inhibition
could cause a similar response. We performed immunostaining of IPO5 and KPNB1 in COS7
cells using a proteasome inhibitor, MG132. Consistent with arsenite, MG132 caused the
accumulation of cytoplasmic granules that contained IPO5 and KPNB1 (Figure 6D). These results
support the idea that RNP granules that contain KPNB1 and IPO5 are formed in *Fbxo24* KO
spermatozoa.

Discussion
Spermiogenesis is the late stage of spermatogenesis in which round spermatids differentiate into
spermatozoa accompanied by drastic morphological changes such as flagellum formation,
chromatin remodeling, and cytoplasmic removal. Protein ubiquitination of target proteins for
degradation by the UPS is crucial to maintain functional spermatogenesis (Richburg et al., 2014).
In this study, we focused on testis-enriched F-box protein FBXO24. We confirmed that *Fbxo24*
was predominantly expressed in the testis, and its expression level increased in the late stage of
spermatogenesis. The analyses using cultured cells and Tg testes suggested that FBXO24 could
interact with SKP1 to form the SCF complex. Further, we deleted *Fxbo24* using the
CRISPR/Cas9 system and found that *Fbxo24* KO mice exhibited abnormal flagellar structures
and male infertility. *Fxbo24* KO spermatozoa failed to fertilize eggs even with IVF, while ICSI
gave offspring.

More detailed observation of the midpiece of *Fbxo24* KO spermatozoa using electron microscopy
revealed abnormalities in mitochondria coiling, which occurred in the late stages of
spermatogenesis and was consistent with the *Fbxo24* expression timing. Furthermore, *Fbxo24*
KO spermatozoa showed markedly reduced motility. These results suggest that the primary
cause of infertility is impaired sperm migration in the female reproductive tract due to reduced
sperm motility. In addition to the abnormal flagellar morphology, *Fbxo24* KO spermatozoa could
not undergo the acrosome reaction even after the A23187 treatment. We analyzed the amounts
of SNARE-related proteins and PLCD4; however, no overt differences were found between the
control and *Fbxo24* KO spermatozoa. Recently, we found that FER1L5 is important for Ca^{2+}-
dependent acrosome reaction (Morohoshi et al., 2023), but no antibodies are available to analyze
the amount of FER1L5. It is possible that FER1L5 or unknown proteins involved in the acrosome
reaction deteriorated in *Fbxo24* KO spermatozoa. Alternatively, it is possible that acrosome
reaction failure may be caused by impaired sperm capacitation associated with abnormal flagellar
structures.

The TEM analyses observed numerous membraneless electron-dense granules in *Fbxo24* KO
sperm flagella. Although the irregular arrangement of sperm mitochondria has been reported in
KO mice of some mitochondria-associated proteins (Shimada et al., 2021, 2019), numerous
granules have not been observed in these KO spermatozoa. Proteomic analyses revealed that
IPO5 and KPNB1 accumulated in *Fbxo24* KO spermatozoa without significant elevation of
mitochondria-associated proteins. The lack of accumulation of KPNA2, which is part of the
importin family, also suggests that there is selectivity in the accumulated proteins. Abnormal
accumulation of proteins, including IPO5 and KPNB1, may lead to morphological abnormalities in

sperm flagella since the SCF complex is critical for protein degradation and consequently regulate various cellular processes (Skaar et al., 2013). FBXO24 likely plays a role in catalyzing protein degradation during spermiogenesis. Because we found that FBXO24 could interact with IPO5 in testes and catalyze ubiquitination of IPO5, FBXO24 may degrade IPO5 in the testis. Recently, mice lacking both PSME4 and ECPAS, proteasome-associated proteins, were shown to exhibit impaired male fertility with disorganized mitochondrial sheath, but numerous membraneless granules were not observed (Sato et al., 2023), suggesting that the accumulation of the granules is specific to *Fbxo24* deletion rather than simply caused by proteasome defects.

In addition to proteins, the total RNA amount increased in *Fbxo24* KO spermatozoa, suggesting that the abnormal granules observed in *Fbxo24* KO flagella may consist of RNPs. In germ cells, RNP granules have been well studied and shown to be essential for spermatogeneses, such as IMCs, CBs, reticulated bodies, and CB remnants (Chuma et al., 2006; Meikar et al., 2011; Shimada et al., 2023). RNP granules are membraneless electron-dense structures that are similar to the abnormal granules in *Fbxo24* KO; however, the localization of MIWI, an IMC and CB marker, and TSKS, a reticulated body and CB remnant marker, was not disrupted in the KO. These results suggest that electron-dense granules accumulated in *Fbxo24* KO spermatozoa are different from well-studied RNP granules. SGs are also membraneless electron-dense granules that contain RNPs, which assemble through liquid–liquid phase separation in response to cellular stresses (Buchan and Parker, 2009; Protter and Parker, 2016; Taylor et al., 2016). While transient SGs formed in response to oxidative stress or heat shock play roles in translational arrest to ensure cell survival, chronic SGs are relevant to neurodegenerative diseases (Marcelo et al., 2021; Reineke and Neilson, 2019). It has been shown that, under arsenite-induced oxidative stress, KPNB1 relocates to SGs (Chang and Tarn, 2009; Mahboubi et al., 2013). Because we found that not only KPNB1 but also IPO5 can be recruited to SGs after arsenite or proteasome inhibitor treatment, electron-dense granules observed in *Fbxo24* KO spermatozoa may be RNP granules consisting of IPO5 and KPNB1. Our results suggest that FBXO24 is responsible for degrading proteins such as IPO5, which may prevent the accumulation of abnormal RNP granules. While this paper was under review, another group reported analyses of *Fbxo24* KO mice that exhibit not only disorganized mitochondrial coiling but also abnormal head morphology (Li et al., 2024). The differences from our study may be due to different mouse genetic backgrounds. The other group showed that FBXO24 mediates the degradation of MIWI via K48-linked polyubiquitination (Li et al., 2024). FBXO24 may be involved in the degradation of multiple proteins including IPO5 to support proper spermiogenesis.

In conclusion, we reveal that numerous RNP granules accumulated in sperm flagella when FBXO24, which is related to the UPS was depleted. Small numbers of the RNP granules were found even in the control, suggesting that the ability to generate the granules is present in WT testes but may be suppressed by FBXO24. FBXO24 can be a key molecule for understanding the formation and function of the aberrant RNP granules in male fertility. Because amino acid sequences and expression patterns of FBXO24 are conserved, FBXO24 may also play similar roles in human testes. Our research may lead to development of new approaches to infertility treatment and nonhormonal male contraceptives.

Materials and Methods
Animals. All animal experiments were approved by the Animal Care and Use Committee of the Research Institute for Microbial Diseases, Osaka University. Mice were purchased from CLEA Japan (Tokyo, Japan) or Japan SLC (Shizuoka, Japan). WT or *Fbxo24* heterozygous (HET) mice were used as controls. All gene-modified mice generated in this study will be made available through either the RIKEN BioResource Research Center or the Center for Animal Resources and Development (CARD), Kumamoto University.

Isolation of RNA and RT-PCR. Adult mouse multi-tissues and mouse testes at different ages were obtained from C57BL/6N mice. RNA samples were isolated and purified

using TRIzol (Thermo Fisher Scientific, Waltham, MA, USA). RNA was reverse transcribed to cDNA using SuperScript IV First-Strand Synthesis System (Thermo Fisher Scientific) using an oligo (dT) primer. PCR was performed using the KOD Fx Neo DNA Polymerase (Toyobo, Tokyo, Japan). Primers used in this study are listed in Table S1.

Transfection of HEK293T cells and induction of stress. HEK293T cells (Tiscornia et al., 2006) and COS7 cells were maintained in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (Sigma-Aldrich, St. Louis, MO, USA) and 1% Gibco™ penicillin/streptomycin (Thermo Fisher Scientific) at 37°C under 5% CO_2. For transfection, HEK293T cells were transiently transfected with the plasmid DNA using the calcium phosphate transfection method (Tiscornia et al., 2006), and cultured for 24 h before harvesting. For oxidative stress, COS7 cells were treated with 0.5 mM sodium arsenite in medium for 30 min and controls were incubated with water in medium. For proteasome inhibition, COS7 cells were treated with 10 µM sodium MG132 (Sigma-Aldrich) in medium for 3 h and controls were incubated with DMSO in medium.

Generation of *Fbxo24* KO mice. *Fbxo24* KO mice were generated using CRISPR/Cas9. Three crRNAs, 5′- TGTGGAGGCGCATCTGTCGA -3′, 5′- TCCTGAAGGAAGTCGAGCCG -3′ and 5′- TCAGTTGTTCCCCCCAGAGC -3′ were designed using the online source CRISPRdirect (Naito et al., 2015) and annealed to SygRNA™ Cas9 Synthetic tracrRNA (#TRACRRNA05N-5NMOL, Sigma-Aldrich). The gRNAs were mixed with TrueCut™ Cas9 Protein v2 (A36498, Thermo Fisher Scientific) and incubated at 37°C for 5 min to form CRISPR/Cas9 complexes. The complexes were introduced into fertilized eggs which were from super-ovulated WT B6D2F1 females mated with B6D2F1 males. Electroporation was performed using the super electroporator NEPA21 (NEPA GENE, Chiba, Japan) (poring pulse, voltage: 225 V, pulse width: 2 ms, pulse interval: 50 ms, number of pulses: +4, and attenuation: 10%; transfer pulse, voltage: 20 V, pulse width: 50 ms, pulse interval: 50 ms, number of pulses: ±5, and attenuation: 40%). The treated eggs were developed into two-cell-stage embryos by cultivating in KSOM medium (Ho et al., 1995) and transplanted into pseudo-pregnant ICR females. The obtained pups were genotyped by PCR to detect the KO and/or WT allele and then subjected to Sanger sequencing to verify the deleted sequence.

Fertility test. For the in vivo fertility analysis, sexually mature KO male mice or WT male mice were caged with three 8-week-old B6D2F1 female mice for three months and plugs were checked every morning. The number of pups was counted on the day of birth. For the in vitro fertility assay, in vitro fertilization (IVF) analysis was performed as previously described (Morohoshi et al., 2021) with some minor changes. For ZP-free oocytes, sperm insemination was performed at a final density of 2×10^4 spermatozoa/mL.

Histological analysis. PAS staining of sections were performed as previously described (Morohoshi et al., 2020). Testes or cauda epididymis were fixed at 4°C in Bouin's solution (Polysciences, Inc., Warrington, PA, USA) and were processed for paraffin embedding. Paraffin sections were cut at a thickness of 5 µm using an HM325 microtome (Microm, Walldorf, Germany). After rehydrating the sections, they were stained with 1% periodic acid (Nacalai Tesque, Kyoto, Japan) and Schiff's reagent (FUJIFILM WakoPure Chemical, Osaka, Japan) for 20 min each at room temperature. The sections were then counterstained with Mayer's hematoxylin solution (FUJIFILM WakoPure Chemical). The sections were observed with an Olympus BX-53 microscope (Tokyo, Japan).

Morphological and motility analysis of spermatozoa. Spermatozoa extracted from cauda epididymis were suspended in TYH medium (Muro et al., 2016). After 10 min incubation, spermatozoa were collected to observe morphology. Sperm motility was analyzed as previously described (Miyata et al., 2021, 2020). The motility of more than 200 spermatozoa was measured after incubation at 10 and 120 min in TYH medium using CEROS II (software version 1.4; Hamilton Thorne Biosciences, Beverly, MA, USA).

Observation of sperm flagellum ultrastructure using TEM. Cauda epididymis specimens were prepared as previously described (Shimada et al., 2019). The prepared samples were observed using a JEM-1400 plus electron microscope (JEOL, Tokyo, Japan) at 80 kV with a CCD Veleta 2K × 2K camera (Olympus).

Observation of sperm mitochondria during spermatogenesis using SEM. Testes specimens were prepared as previously described (Shimada et al., 2019). The prepared samples were observed using a S-4800 field emission scanning electron microscope (Hitachi, Tokyo, Japan).

Pfam domain search. Pfam domains were detected from amino acid sequences using Simple Modular Architecture Research Tool (SMART) (http://smart.embl-heidelberg.de/). Mouse FBXO24 amino acid sequence (CCDS51674.1) and human FBXO24 (CCDS5698.1) were obtained from the CCDS database (https://www.ncbi.nlm.nih.gov/CCDS/CcdsBrowse.cgi).

Alignment of amino acid sequences. Amino acid sequences of Mouse FBXO24 (CCDS51674.1) and human FBXO24 (CCDS5698.1) were aligned using "Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/)". Aligned sequences were edited using Jalview ver. 2.11.2.0 (https://www.jalview.org/). Domains within the sequences were identified using "SMART (http://smart.embl-heidelberg.de/)".

***In silico* expression data analysis.** Single-cell transcriptome data in the mouse and human testis was obtained from previously published work (Hermann et al., 2018). *Fbxo24* expression in those cells was analyzed using the Loupe Cell Browser 3.3.1 (10X Genomics, Pleasanton, CA, USA).

Plasmids. The open reading frames (ORFs) of *Fbxo24* and *Skp1* were cloned and amplified using cDNA obtained from mouse testis and inserted into the multiple cloning site of pCAG1.1 vector (Addgene; Plasmid #173685). To generate the expression vector coding *Fbxo24(ΔF-box),* inverse PCR was performed with KOD -Plus- Mutagenesis Kit (Toyobo, Tokyo, Japan) according to the manufacturer's instructions. Primers used for inverse PCR are listed in Table S1. HA-human HA-Ubiquitin was a gift from Edward Yeh (Addgene plasmid # 18712; Kamitani et al., 1997)

Observation of spermatozoa migration inside the female reproductive tract. The *Fbxo24* KO mouse line was crossed with B6D2 Tg mice carrying CAG/Su9-DsRed2, Acr3-EGFP (RBGS) (Hasuwa et al., 2010) to label the acrosome with EGFP. B6D2F1 females were superovulated by injecting pregnant mare serum gonadotropin (PMSG, ASKA Pharmaceutical, Tokyo, Japan) followed by human chorionic gonadotropin (hCG, ASKA Pharmaceutical) 48 h appart. The superovulated female was mated with a WT male or *Fbxo24* KO male mice 14 h after the hCG injection. Female mice were sacrificed 4 h after confirming a vaginal plug and the female reproductive tracts were collected. Spermatozoa inside the oviducts were observed using a BZ-X710 microscope (Keyence Japan, Osaka, Japan).

Protein extraction from testes, spermatozoa, and culture cells. HEK293T cells were lysed in 1% Triton X-100 lysis buffer [1% Triton X-100, 50 mM Tris-HCl pH 7.4, 150 mM NaCl, and 1% (v/v) protease inhibitor cocktail (Nacalai Tesque)] at 4°C with end-over-end rotation for 2 h. Testes or spermatozoa were homogenized in 1% Triton X-100 lysis buffer at 4°C with end-over-end rotation for 2 h or sample buffer containing 1 M Tris-HCL pH 6.8, 2% SDS, 10% Glycerol, and 0.005% Bromophenol Blue, and boiled for 5 min. Supernatants were obtained after centrifugation at 15,300 × g for 15 min at 4°C.

Immunoprecipitation. Immunoprecipitation was performed as described previously (Kaneda et al., 2023). Proteins from HEK293T cells were extracted using 1% Triton X-100 lysis buffer [1% Triton X-100, 50 mM Tris-HCl pH 7.4, 150 mM NaCl, and 1% (v/v) protease inhibitor cocktail (Nacalai Tesque)]. The lysates were centrifuged at 15,300 g for 5 min at 4°C to collect

supernatants. Protein lysates were mixed with 20 µL Protein G-conjugated magnetic beads
(#10009D, Thermo Fisher Scientific) preincubated with 1.0 µg antibody. After incubation for 1 h at
4°C, the beads were washed three times with wash buffer (40 mM Tris-HCl pH 7.5, 150 mM
NaCl, 0.1% Triton X-100 and 10% glycerol). For immunoblotting, the immune complexes were
eluted with 2×SDS sample buffer (132 mM Tris HCl pH7.4, 4% SDS, 20% glycerol, and 0.01%
Bromophenol Blue) for 10 min at 70°C. For mass spectrometry (MS), the immune complexes
were eluted with 150 ng/µL 3×FLAG peptide (Sigma-Aldrich) in TBS solution for 30 min at 4°C.
Antibodies used in this study are listed in Table S2.

Immunoblotting. Proteins were separated with sodium dodecyl sulfate polyacrylamide gel
electrophoresis (SDS-PAGE) under reducing conditions using 2-ME and transferred onto
polyvinylidene difluoride (PVDF) membrane using the Trans Blot Turbo system (Bio-Rad,
Hercules, CA, USA). After blocking with 10% skim milk (Becton Dickinson, Franklin Lakes, NJ,
USA) in TBST, the blots were incubated with primary antibody overnight at 4°C, and then
incubated with HRP-conjugated secondary antibodies (1:5000) for 2 h at room temperature.
Chemiluminescence was detected with Chemi-Lumi One Super (#02230, Nacalai Tesque) or
Chemi-Lumi One Ultra (#11644, Nacalai Tesque).

I**n vivo ubiquitination assay**, HEK293T cells were transiently transfected with the plasmid DNA
following the transfection protocol mentioned above. After 36 h of transfection, HEK293T cells
were treated with 10 µM MG132 for 9 h. Immunoprecipitation and immunoblotting were
performed following the above mentioned protocols.

Assessment of sperm viability in vitro. Spermatozoa extracted from cauda epididymis were
dispersed in TYH medium and incubated for 10 and 120 min at 37°C under 5% CO_2. PI was
carefully added to the TYH drop to achieve a final concentration of 10 µg/ml and a small volume
of the sperm suspension was placed on a glass slide. More than 200 spermatozoa were counted
for each trial.

Assessment of acrosome reaction in vitro. The *Fbxo24* KO mouse line was crossed with
B6D2 Tg mice carrying CAG/Su9-DsRed2, Acr3-EGFP (RBGS) (Hasuwa et al., 2010) to label the
acrosome with EGFP. Spermatozoa extracted from cauda epididymis were dispersed in TYH
medium for 10 and 120 min of incubation at 37°C under 5% CO_2. After 120 min of incubation,
spermatozoa were incubated with Ca^{2+} ionophore A23187 (Merck, Rahway, NJ, USA) at 20 µM to
induce the acrosome reaction at 37°C under 5% CO_2. 10 mM stock solution of Ca^{2+} ionophore
A23187 in DMSO was diluted 10 times with TYH medium before carefully adding to the sperm
suspension to reach the final concentration. PI (10 µg/ml at final concentration) was added to the
TYH drop, and a small volume of the sperm suspension was placed on a glass slide. Acrosome-
reacted spermatozoa were determined by observing EGFP signals while distinguishing viable
spermatozoa with PI staining using a BX-53 microscope (Olympus). More than 200 spermatozoa
were counted for each trial.

Intracytoplasmic sperm injection (ICSI). B6D2F1 females were superovulated by injecting
PMSG (ASKA Pharmaceutical) followed by hCG (ASKA Pharmaceutical) 48 h later. MII oocytes
were collected 14 h after injection of hCG. Cumulus cells were removed from the collected eggs
after 10 min treatment with 330 µg/mL hyaluronidase (Sigma-Aldrich). Cumulus-free oocytes
were placed in KSOM medium at 37°C under 5% CO2 until just before performing ICSI. Each
sperm head separated from the tail by applying a few piezo pulses was injected into a cumulus-
free MII oocyte using a piezo manipulator (PrimeTech, Tokyo, Japan). Obtained two-cell embryos
were transferred to pseudopregnant ICR females the next day.

Immunofluorescence of testis cross-sections. Testes were fixed in 4% paraformaldehyde in
PBS for 3 h at 4°C soon after dissection and infused with 15% and 30% sucrose in PBS at 4°C.
The testes were embedded in OCT medium (Tissue-Tek, Miami, FL, USA) and frozen in liquid
nitrogen. Testis blocks were sectioned at 10 µm thickness using a cryostat and the sections were

dried on microscope slides at 42°C to make them adhere. The testis sections were then
processed and imaged in the same manner as sperm immunostaining. After permeabilization with
0.1% Triton X-100 in PBS for 15 min, the sections were blocked with 3% bovine serum albumin
(Sigma-Aldrich), 10% goat serum, and 0.1% Triton X-100 in PBS for 1 h at room temperature.
The sections were then incubated with primary antibody overnight at 4°C. After washing the
sections with PBS for 10 min, the sections were secondarily blocked with 3% bovine serum
albumin (Sigma-Aldrich), and 10% goat serum in PBS for 1 h at room temperature. The sections
were further incubated with fluorophore-conjugated secondary antibodies (1:2000) and Alexa
Fluor 568 Dye PNA Lectin (1:2000) for 2 h at room temperature. After a wash with PBS, the
sections were incubated with Hoechst 33342 (Thermo Fisher Scientific) in PBS (1:5000) for 5 min
at room temperature. Once washed with PBS, the sections were mounted with Shandon Immu-
Mount (Thermo Fisher Scientific) before imaging.

Immunostaining of spermatozoa. Spermatozoa extracted from cauda epididymis were
dispersed in PBS. A small aliquot of the sperm suspension was spotted onto glass slides and
then air-dried at room temperature. Spermatozoa were then fixed with 4% paraformaldehyde in
PBS for 10 min and washed three times with PBS for 5 min each. The slides were blocked with
blocking solution (5% bovine serum albumin and 10% goat serum in PBS) for 1 h at room
temperature. The slides were incubated with primary antibody overnight at 4°C, washed with PBS
three times for 5 min each, and incubated with Alexa Fluor conjugated secondary antibody for 2 h
at room temperature. The slides were washed with PBS three times for 5 min each, incubated
with Hoechst 33342 (2 μg/ml) (Thermo Fisher Scientific) for 5 min, and washed with PBS three
times for 5 min each. The slides were mounted with Shandon Immu-Mount (Thermo Fisher
Scientific) before imaging. Fluorescent images were captured with an Olympus BX-53
microscope (Olympus).

Immunofluorescence of COS7 cells. COS7 cells were fixed with 4% paraformaldehyde in PBS
for 15 min at room temperature. After permeabilization with 0.1% Triton X-100 in PBS for 15 min,
cells were blocked with 1% bovine serum albumin (Sigma-Aldrich) in PBS for 1 h at room
temperature. The cells were then incubated with primary antibody overnight at 4°C. After washing
three times with PBS, the cells were further incubated with fluorophore-conjugated secondary
antibodies (1:1000) for 2 h at room temperature. After washing three times with PBS, the cells
were incubated with Hoechst 33342 (Thermo Fisher Scientific) in PBS (1:5000) for 5 min at room
temperature. After washing three times with PBS, the cells were mounted with Shandon Immu-
Mount (Thermo Fisher Scientific) before imaging.

Extraction of RNA from epididymal spermatozoa. A small cut was made in the cauda
epididymis with ophthalmic scissors and spermatozoa were squeezed out from the cut. Collected
spermatozoa were dispersed in PBS and the number of spermatozoa was counted using a
hemacytometer. Spermatozoa were centrifuged at 15,300 g for 2 min and the sperm pellet was
lysed in TRIzol (Thermo Fisher Scientific). Chloroform was added to the lysate and incubated for
5 min at room temperature. The sample was centrifuged at 15,300 g for 15 min and the aqueous
phase was transferred to a fresh centrifuge tube. To precipitate RNA, an equal volume of
isopropanol was added to the aqueous phase and centrifuged at 15,300 g for 20 min at 4°C. The
supernatant was removed and the pellet was washed with 75% Ethanol. The washed pellet was
dried and resuspended in nuclease-free water. The RNA solution was DNAse treated using
deoxyribonuclease (RT Grade) (NIPPON GENE, Tokyo, Japan) to remove genomic DNA. Then
the RNA solution was incubated at 55°C for 10 min. The sample was stored at -80°C. The
concentration was measured using NanoDrop One (Thermo Fisher Scientific). The
TAE/formamide method (Masek et al., 2005) was used to perform electrophoresis of RNA.

Statistical analysis. All statistical analyses were performed using the two-tailed Welch's t-test
using Microsoft Office Excel 2016 (Microsoft Corporation, Redmond, WA, USA). Differences were
considered significant at $P<0.05$ (*), $P<0.01$ (**), $P<0.001$ (***). Error bars are standard deviation.

Acknowledgments

The authors would like to thank Dr. Julio M. Castaneda and Ferheen Abbasi for the critical reading of the manuscript and Ms. Natsuki Furuta for technical assistance, and the members of both the Department of Experimental Genome Research and the nonprofit organization (NPO) for Biotechnology Research and Development for experimental assistance and discussion. We also thank Dr. Yoichi Miyamoto for kindly providing antibodies against importins (National Institutes of Biomedical Innovation, Health and Nutrition), and Dr. Akinori Ninomiya and Dr. Fuminori Sugihara for MS analysis (Core Instrumentation Facility, Research Institute for Microbial Diseases, Osaka University). This research was supported by the Ministry of Education, Culture, Sports, Science and Technology (MEXT)/Japan Society for the Promotion of Science (JSPS) KAKENHI grants (JP23KJ1523 to Y.K., JP22H03214, JP23K18328 to H.M., JP23K05831 to K.S., JP22K15030 to M.K., and JP19H05750, JP21H04753, JP21H05033 to M.I.); Takeda Science Foundation grant to H.M., K.S., and M.I.; JST FOREST (JPMJFR211F to H.M.); the Eunice Kennedy Shriver National Institute of Child Health and Human Development (P01HD087157 and R01HD088412 to M.I.); and the Bill & Melinda Gates Foundation (Grand Challenges Explorations grant INV-001902 to M.I.).

Author Contributions

Y.K., H.M., and M.I. conceived and designed experiments and wrote the manuscript with input from all authors. All authors performed experiments and analyzed data.

Competing interests

The authors declare no competing interests.

References

Bai C, Sen P, Hofmann K, Ma L, Goebl M, Harper JW, Elledge SJ. 1996. SKP1 Connects Cell Cycle Regulators to the Ubiquitin Proteolysis Machinery through a Novel Motif, the F-Box. *Cell* **86**:263–274. doi:10.1016/S0092-8674(00)80098-7

Buchan JR, Parker R. 2009. Eukaryotic Stress Granules: The Ins and Outs of Translation. *Mol Cell*. doi:10.1016/j.molcel.2009.11.020

Chang WL, Tarn WY. 2009. A role for transportin in deposition of TTP to cytoplasmic RNA granules and mRNA decay. *Nucleic Acids Res* **37**:6600–6612. doi:10.1093/nar/gkp717

Chook Y, Blobel G. 2001. Karyopherins and nuclear import. *Curr Opin Struct Biol* **11**:703–715. doi:10.1016/S0959-440X(01)00264-0

Chook YM, Süel KE. 2011. Nuclear import by karyopherin-βs: Recognition and inhibition. *Biochim Biophys Acta Mol Cell Res*. doi:10.1016/j.bbamcr.2010.10.014

Chuma S, Hosokawa M, Kitamura K, Kasai S, Fujioka M, Hiyoshi M, Takamune K, Noce T, Nakatsuji N. 2006. *Tdrd1/Mtr-1* , a *tudor* -related gene, is essential for male germ-cell differentiation and nuage/germinal granule formation in mice. *Proc Natl Acad Sci U S A* **103**:15894–15899. doi:10.1073/pnas.0601878103

Chung J-J, Shim S-H, Everley RA, Gygi SP, Zhuang X, Clapham DE. 2014. Structurally Distinct
Ca2+ Signaling Domains of Sperm Flagella Orchestrate Tyrosine Phosphorylation and
Motility. *Cell* **157**:808–822. doi:10.1016/j.cell.2014.02.056

Ciechanover A, Schwartz AL. 1998. The ubiquitin-proteasome pathway: The complexity and
myriad functions of proteins death. *Proc Natl Acad Sci U S A* **95**:2727–2730.
doi:10.1073/pnas.95.6.2727

Conti E, Izaurralde E. 2001. Nucleocytoplasmic transport enters the atomic age. *Curr Opin*
Cell Biol **13**:310–319. doi:10.1016/S0955-0674(00)00213-1

Eddy EM. 1974. Fine structural observations on the form and distribution of nuage in germ
cells of the rat. *Anat Rec* **178**:731–757. doi:10.1002/ar.1091780406

Eddy EM, Toshimori K, O'Brien DA. 2003. Fibrous sheath of mammalian spermatozoa.
Microsc Res Tech **61**:103–115. doi:10.1002/jemt.10320

Fujihara Y, Miyata H, Ikawa M. 2018. Factors controlling sperm migration through the
oviduct revealed by gene-modified mouse models. *Exp Anim* **67**:91–104.
doi:10.1538/expanim.17-0153

Fujihara Y, Noda T, Kobayashi K, Oji A, Kobayashi S, Matsumura T, Larasati T, Oura S,
Kojima-Kita K, Yu Z, Matzuk MM, Ikawa M. 2019. Identification of multiple male
reproductive tract-specific proteins that regulate sperm migration through the oviduct in
mice. *Proc Natl Acad Sci U S A* **116**:18498–18506. doi:10.1073/pnas.1908736116

Fujihara Y, Okabe M, Ikawa M. 2014. GPI-Anchored Protein Complex, LY6K/TEX101, Is
Required for Sperm Migration into the Oviduct and Male Fertility in Mice1. *Biol Reprod*
90:60. doi:10.1095/biolreprod.113.112888

Fukami K, Nakao K, Inoue T, Kataoka Y, Kurokawa M, Fissore RA, Nakamura K, Katsuki M,
Mikoshiba K, Yoshida N, Takenawa T. 2001. Requirement of Phospholipase Cδ4 for the
Zona Pellucida-Induced Acrosome Reaction. *Science (1979)* **292**:920–923.
doi:10.1126/science.1059042

HASUWA H, MURO Y, IKAWA M, KATO N, TSUJIMOTO Y, OKABE M. 2010. Transgenic
Mouse Sperm that Have Green Acrosome and Red Mitochondria Allow Visualization of
Sperm and Their Acrosome Reaction in Vivo. *Exp Anim* **59**:105–107.
doi:10.1538/expanim.59.105

Hermann BP, Cheng K, Singh A, Roa-De La Cruz L, Mutoji KN, Chen IC, Gildersleeve H, Lehle
JD, Mayo M, Westernströer B, Law NC, Oatley MJ, Velte EK, Niedenberger BA, Fritze D,
Silber S, Geyer CB, Oatley JM, McCarrey JR. 2018. The Mammalian Spermatogenesis Single-
Cell Transcriptome, from Spermatogonial Stem Cells to Spermatids. *Cell Rep* **25**:1650-
1667.e8. doi:10.1016/j.celrep.2018.10.026

Ho Y, Wigglesworth K, Eppig JJ, Schultz RM. 1995. Preimplantation development of mouse
embryos in KSOM: Augmentation by amino acids and analysis of gene expression. *Mol*
Reprod Dev **41**:232–238. doi:10.1002/mrd.1080410214

Hutt DM, Baltz JM, Ngsee JK. 2005. Synaptotagmin VI and VIII and Syntaxin 2 Are Essential
for the Mouse Sperm Acrosome Reaction. *Journal of Biological Chemistry* **280**:20197–
20203. doi:10.1074/jbc.M412920200

Kamitani T, Kito K, Nguyen HP, Yeh ETH. 1997. Characterization of NEDD8, a
Developmentally Down-regulated Ubiquitin-like Protein. *Journal of Biological Chemistry*
272:28557–28562. doi:10.1074/jbc.272.45.28557

Kaneda Y, Miyata H, Shimada K, Oura S, Ikawa M. 2023. Testis-specific proteins, TSNAXIP1
and 1700010I14RIK, are important for sperm motility and male fertility in mice. *Andrology*
11:799–807. doi:10.1111/andr.13378

Katafuchi K, Mori T, Toshimori K, Iida H. 2000. Localization of a syntaxin isoform, syntaxin
2, to the acrosomal region of rodent spermatozoa. *Mol Reprod Dev* **57**:375–383.
doi:10.1002/1098-2795(200012)57:4<375::AID-MRD10>3.0.CO;2-Z

Kimura M, Imamoto N. 2014. Biological Significance of the Importin-β Family-Dependent
Nucleocytoplasmic Transport Pathways. *Traffic*. doi:10.1111/tra.12174

Kipreos ET, Pagano M. 2000. The F-box protein family. *Genome Biol* **1**:3002.1-3002.7.
doi:10.1186/gb-2000-1-5-reviews3002

Kissel H, Georgescu MM, Larisch S, Manova K, Hunnicutt GR, Steller H. 2005. The Sept4
septin locus is required for sperm terminal differentiation in mice. *Dev Cell* **8**:353–364.
doi:10.1016/j.devcel.2005.01.021

Kluin PM, Kramer MF, de Rooij DG. 1982. Spermatogenesis in the immature mouse
proceeds faster than in the adult. *Int J Androl* **5**:282–294. doi:10.1111/j.1365-
2605.1982.tb00257.x

Leblond CP, Clermont Y. 1952. Spermiogenesis of rat, mouse, hamster and guinea pig as
revealed by the "periodic acid-fuchsin sulfurous acid" technique. *Am Anat* **90**:167–215.
doi:10.1002/aja.1000900202

Li W, Bengtson MH, Ulbrich A, Matsuda A, Reddy VA, Orth A, Chanda SK, Batalov S,
Joazeiro CAP. 2008. Genome-Wide and Functional Annotation of Human E3 Ubiquitin
Ligases Identifies MULAN, a Mitochondrial E3 that Regulates the Organelle's Dynamics and
Signaling. *PLoS One* **3**:e1487. doi:10.1371/journal.pone.0001487

Li Z, Liu X, Zhang Y, Li Y, Zhou L, Yuan S. 2024. FBXO24 modulates mRNA alternative splicing
and MIWI degradation and is required for normal sperm formation and male fertility. *Elife*
12:RP91666. doi:10.7554/eLife.91666.3

Loveland KateL, Major AndrewT, Butler R, Young JuliaC, Jans DavidA, Miyamoto Y. 2015.
Putting things in place for fertilization: discovering roles for importin proteins in cell fate
and spermatogenesis. *Asian J Androl* **17**:537. doi:10.4103/1008-682X.154310

Loveland KL, Hogarth C, Szczepny A, Prabhu SM, Jans DA. 2006. Expression of nuclear
transport importins beta 1 and beta 3 is regulated during rodent spermatogenesis. *Biol*
Reprod **74**:67–74. doi:10.1095/biolreprod.105.042341

Lu Y, Nagamori I, Kobayashi H, Kojima-Kita K, Shirane K, Chang H, Nishimura T, Koyano T,
Yu Z, Castañeda JM, Matsuyama M, Kuramochi-Miyagawa S, Matzuk MM, Ikawa M. 2023.
ADAD2 functions in spermiogenesis and piRNA biogenesis in mice. *Andrology* **11**:698–709.
doi:10.1111/andr.13400

Mahboubi H, Seganathy E, Kong D, Stochaj U. 2013. Identification of Novel Stress Granule
Components That Are Involved in Nuclear Transport. *PLoS One* **8**:e68356.
doi:10.1371/journal.pone.0068356

Mani A, Gelmann EP. 2005. The ubiquitin-proteasome pathway and its role in cancer. *J Clin*
Oncol doi:10.1200/JCO.2005.05.081

Marcelo A, Koppenol R, de Almeida LP, Matos CA, Nóbrega C. 2021. Stress granules, RNA-
binding proteins and polyglutamine diseases: too much aggregation? *Cell Death Dis* **12**:592.
doi:10.1038/s41419-021-03873-8

Masek T, Vopalensky V, Suchomelova P, Pospisek M. 2005. Denaturing RNA
electrophoresis in TAE agarose gels. *Anal Biochem* **336**:46–50.
doi:10.1016/j.ab.2004.09.010

Meikar O, Da Ros M, Korhonen H, Kotaja N. 2011. Chromatoid body and small RNAs in
male germ cells. *Reproduction*. doi:10.1530/REP-11-0057

Miyamoto Y, Baker MA, Whiley PA, Arjomand A, Ludeman J, Wong C, Jans DA, Loveland KL.
2013. Towards delineation of a developmental α-importome in the mammalian male
germline. *Biochim Biophys Acta Mol Cell Res* **1833**:731–742.
doi:10.1016/j.bbamcr.2012.11.005

Miyata H, Abbasi F, Visconti PE, Ikawa M. 2020. CRISPR/CAS9-mediated amino acid
substitution reveals phosphorylation residues of RSPH6A are not essential for male fertility
in mice. *Biol Reprod* **103**:912–914. doi:10.1093/biolre/ioaa161

Miyata H, Oura S, Morohoshi A, Shimada K, Mashiko D, Oyama Y, Kaneda Y, Matsumura T,
Abbasi F, Ikawa M. 2021. SPATA33 localizes calcineurin to the mitochondria and regulates
sperm motility in mice. *Proc Natl Acad Sci U S A* **118**:1–9. doi:10.1073/pnas.2106673118

Miyata H, Satouh Y, Mashiko D, Muto M, Nozawa K, Shiba K, Fujihara Y, Isotani A, Inaba K,
Ikawa M. 2015. Sperm calcineurin inhibition prevents mouse fertility with implications for
male contraceptive. *Science (1979)* **350**:442–445. doi:10.1126/science.aad0836

Morohoshi A, Miyata H, Oyama Y, Oura S, Noda T, Ikawa M. 2021. FAM71F1 binds to
RAB2A and RAB2B and is essential for acrosome formation and male fertility in mice.
Development **148**:1–12. doi:10.1242/dev.199644

Morohoshi A, Miyata H, Shimada K, Nozawa K, Matsumura T, Yanase R, Shiba K, Inaba K,
Ikawa M. 2020. Nexin-Dynein regulatory complex component DRC7 but not FBXL13 is
required for sperm flagellum formation and male fertility in mice. *PLoS Genet* **16**:1–21.
doi:10.1371/journal.pgen.1008585

Morohoshi A, Miyata H, Tokuhiro K, Iida-Norita R, Noda T, Fujihara Y, Ikawa M. 2023.
Testis-enriched ferlin, FER1L5, is required for Ca^{2+}-activated acrosome reaction and male
fertility. *Sci Adv* **9**:eade7607. doi:10.1126/sciadv.ade7607

Muro Y, Hasuwa H, Isotani A, Miyata H, Yamagata K, Ikawa M, Yanagimachi R, Okabe M.
2016. Behavior of mouse spermatozoa in the female reproductive tract from soon after
mating to the beginning of fertilization. *Biol Reprod* **94**:1–7.
doi:10.1095/biolreprod.115.135368

Naito Y, Hino K, Bono H, Ui-Tei K. 2015. CRISPRdirect: Software for designing CRISPR/Cas
guide RNA with reduced off-target sites. *Bioinformatics* **31**:1120–1123.
doi:10.1093/bioinformatics/btu743

Nakayama K, Nagahama H, Minamishima YA, Matsumoto M, Nakamichi I, Kitagawa K,
Shirane M, Tsunematsu R, Tsukiyama T, Ishida N, Kitagawa M, Nakayama KI, Hatakeyama S.
2000. Targeted disruption of Skp2 results in accumulation of cyclin E and p27 (Kip1),
polyploidy and centrosome overduplication. *EMBO Journal* **19**:2069–2081.
doi:10.1093/emboj/19.9.2069

Nsota Mbango JF, Coutton C, Arnoult C, Ray PF, Touré A. 2019. Genetic causes of male
infertility: Snapshot on morphological abnormalities of the sperm flagellum. *Basic Clin*
Androl **29**:1–8. doi:10.1186/s12610-019-0083-9

Pan J, Goodheart M, Chuma S, Nakatsuji N, Page DC, Wang PJ. 2005. RNF17, a component
of the mammalian germ cell nuage, is essential for spermiogenesis. *Development*
132:4029–4039. doi:10.1242/dev.02003

Protter DSW, Parker R. 2016. Principles and Properties of Stress Granules. *Trends Cell Biol*.
doi:10.1016/j.tcb.2016.05.004

Reineke LC, Neilson JR. 2019. Differences between acute and chronic stress granules, and
how these differences may impact function in human disease. *Biochem Pharmacol*.
doi:10.1016/j.bcp.2018.10.009

Richburg JH, Myers JL, Bratton SB. 2014. The role of E3 ligases in the ubiquitin-dependent
regulation of spermatogenesis. *Semin Cell Dev Biol*. doi:10.1016/j.semcdb.2014.03.001

Sato B, Kim J, Morohoshi K, Kang W, Miyado K, Tsuruta F, Kawano N, Chiba T. 2023.
Proteasome-Associated Proteins, PA200 and ECPAS, Are Essential for Murine
Spermatogenesis. *Biomolecules* **13**:586. doi:10.3390/biom13040586

Schulz JR, Wessel GM, Vacquier VD. 1997. The Exocytosis Regulatory Proteins Syntaxin and
VAMP Are Shed from Sea Urchin Sperm during the Acrosome Reaction. *Dev Biol* **191**:80–
87. doi:10.1006/dbio.1997.8712

Schulman BA, Carrano AC, Philip D. J, Zachary B, Elspeth R. E. K, Michael S. F, Stephen J. E,
J. Wade H, Michele P, & Nikola P. P. 2000. Insights into SCF ubiquitin ligases from the
structure of the Skp1-Skp2 complex. *Nature* **408**:381–386.
doi:https://doi.org/10.1038/35042620

Shang P, Baarends WM, Hoogerbrugge J, Ooms MP, Van Cappellen WA, De Jong AAW,
Dohle GR, Van Eenennaam H, Gossen JA, Grootegoed JA. 2010. Functional transformation
of the chromatoid body in mouse spermatids requires testis-specific serine/threonine
kinases. *J Cell Sci* **123**:331–339. doi:10.1242/jcs.059949

Shimada K, Kato H, Miyata H, Ikawa M. 2019. Glycerol kinase 2 is essential for proper
arrangement of crescent-like mitochondria to form the mitochondrial sheath during mouse
spermatogenesis. *J Reprod Dev* **65**:155–162. doi:10.1262/jrd.2018-136

Shimada K, Park S, Miyata H, Yu Z, Morohoshi A, Oura S, Matzuk MM, Ikawa M. 2021.
ARMC12 regulates spatiotemporal mitochondrial dynamics during spermiogenesis and is
required for male fertility. *Proc Natl Acad Sci U S A* **118**:e2018355118.
doi:10.1073/pnas.2018355118

Shimada K, Park S, Oura S, Noda T, Morohoshi A, Matzuk MM, Ikawa M. 2023. TSKS
localizes to nuage in spermatids and regulates cytoplasmic elimination during spermiation.
Proc Natl Acad Sci U S A **120**:e2221762120. doi:10.1073/pnas.2221762120

Skaar JR, Pagan JK, Pagano M. 2013. Mechanisms and function of substrate recruitment by
F-box proteins. *Nat Rev Mol Cell Biol*. doi:10.1038/nrm3582

Snyder E, Chukrallah L, Seltzer K, Goodwin L, Braun RE. 2020. ADAD1 and ADAD2, testis-
specific adenosine deaminase domain-containing proteins, are required for male fertility.
Sci Rep **10**:11536. doi:10.1038/s41598-020-67834-5

Taylor JP, Brown RH, Cleveland DW. 2016. Decoding ALS: From genes to mechanism.
Nature. doi:10.1038/nature20413

Tiscornia G, Singer O, Verma IM. 2006. Production and purification of lentiviral vectors. *Nat
Protoc* **1**:241–245. doi:10.1038/nprot.2006.37

Tomes CN, Michaut M, Blas G De, Visconti P, Matti U, Mayorga LS. 2002. SNARE Complex
Assembly Is Required for Human Sperm Acrosome Reaction. *Dev Biol* **243**:326–338.
doi:10.1006/dbio.2002.0567

Tsunematsu R, Nakayama K, Oike Y, Nishiyama M, Ishida N, Hatakeyama S, Bessho Y,
Kageyama R, Suda T, Nakayama KI. 2004. Mouse Fbw7/Sel-10/Cdc4 Is Required for Notch
Degradation during Vascular Development. *J Biol Chem* **279**:9417–9423.
doi:10.1074/jbc.M312337200

Yamaguchi R, Yamagata K, Ikawa M, Moss SB, Okabe M. 2006. Aberrant distribution of
ADAM3 in sperm from both angiotensin-converting enzyme (Ace)- and calmegin (Clgn)-
deficient mice. *Biol Reprod* **75**:760–766. doi:10.1095/biolreprod.106.052977

Zheng Q, Huang T, Zhang L, Zhou Y, Luo H, Xu H, Wang X. 2016. Dysregulation of Ubiquitin-
Proteasome System in Neurodegenerative Diseases. *Front Aging Neurosci* **8**:303.
doi:10.3389/fnagi.2016.00303

Figures

799
800
Figure 1. *Fbxo24* is predominantly expressed in testes and essential for male fertility. (*A*)
RT-PCR of *Fbxo24* in mouse adult tissues. *Fbxo24* is predominantly expressed in the testis. Br:
brain, Th: thymus, Lu: lung, He: heart, Li: liver, Sp: spleen, Ki: kidney, Te: testis, Ut: uterus, Ov:
ovary, and N.C.: negative control (water). *Actb* was used as a control. (*B*) RT-PCR of *Fbxo24*
using RNAs obtained from mouse testes at various postnatal days. *Actb* was used as a control.
Water was used as a negative control (N.C.). (*C*) Construction of expression vectors for FBXO24
with (WT) or without (ΔF) the F-box domain. (*D*) *Fbxo24 (WT)-FLAG* or *Fbxo24 (ΔF)-FLAG* was
transiently expressed with *Skp1-1D4* in HEK293T cells. Immunoprecipitation (IP) was performed
using anti-1D4 antibody or anti-FLAG antibody. FBXO24-FLAG interacts with SKP1-1D4 via the
F-box domain. α-tubulin was used as a loading control. (*E*) Schematic for generating *Fbxo24* KO
mice using the CRISPR/Cas9 system. White boxes indicate untranslated regions while black
boxes indicate protein coding regions. The gRNAs used are shown. Fw and Rv indicate the
forward and reverse primer used for genotyping, respectively. (*F*) Genotyping of obtained *Fbxo24*
mutant mice. Fw #1-Rv #1 primers for KO allele and Fw #2-Rv #2 primers for WT allele in Fig. 1E
were used. N.C. indicates negative control (water). (*G*) Amplicons of the PCR product using Fw
#1-Rv #1 primers were subjected to direct sequencing and the 11,575 bp deletion was confirmed
in the KO allele. (H) The number of pups born per plug was counted to assess male fertility. Each
WT or KO male was mated with three WT females for three months.

819
820
Figure 2. FBXO24 is essential for sperm flagellar formation and motility. (*A*) The histology of
seminiferous tubules at different stages. An asterisk indicates remaining sperm heads. (*B*)
Morphology of mature spermatozoa obtained from cauda epididymis. White arrowhead indicates
straight spermatozoa. Red arrowhead indicates bent spermatozoa. Black asterisk indicates coiled
spermatozoa. Red asterisk indicates headless spermatozoa. (*C*) Stacked bar graph showing the
frequency of sperm morphology classified as straight, bent, coiled, and headless. $n = 3$
independent experiments. (*D*) Percentages of motile spermatozoa were analyzed at 10 min and
120 min after incubation in capacitation medium. (*E*) VAP (average path velocity) was analyzed at
10 min and 120 min after incubation in capacitation medium. (*F*) VSL (straight line velocity) was
analyzed at 10 min and 120 min after incubation in capacitation medium. (G) VCL (curvilinear
velocity) was analyzed at 10 min and 120 min after incubation in capacitation medium.

832
833
Figure 3. In vitro fertilizing ability and in vivo sperm migration. (*A*) The fertilizing ability of
spermatozoa was analyzed in vitro using cumulus-intact oocytes. (*B*) The fertilizing ability of
spermatozoa was analyzed in vitro using cumulus-free oocytes. (*C*) The fertilizing ability of
spermatozoa was analyzed in vitro using zona-free oocytes. (*D*) Uterus and oviducts of WT
females mated with WT or *Fbxo24* KO males carrying RBGS transgene. Female reproductive
tracts were dissected 4 h after confirming a plug. Right figures are magnified images of the boxes
indicated in the middle panels. (*E*) Immunoblotting of ADAM3 using testis and mature
spermatozoa of *Fbxo24* heterozygous or KO mice. (*F*) ICSI experiment. The number of two-cell
stage embryos and pups developed from WT oocytes injected with WT or *Fbxo24* KO
spermatozoa. (*G*) Pups derived from WT or *Fbxo24* KO spermatozoa. (*H*) Genotyping of pups
obtained from *Fbxo24* KO spermatozoa. N.C. indicates negative control (water).

845
846
Figure 4. Numerous membraneless electron-dense granules were found in *Fbxo24* KO
spermatozoa. (*A*) Epididymal spermatozoa of RBGS Tg mice were stained with Hoechst 33342
(nuclei). Mitochondria were labeled with su9-DsRed2. White arrowhead indicates peeled off
mitochondria. (*B*) Sperm mitochondrial sheath formation during spermiogenesis was observed by
scanning electron microscopy (SEM). (*C*) Cross sections of spermatozoa in the cauda
epididymis. Asterisks indicate electron-dense granules. (*D*) Longitudinal sections of spermatozoa
in the cauda epididymis. An asterisk indicates electron-dense granules. (*E*) Percentages of
morphologically abnormal mitochondria observed with transmission electron microscopy (TEM).
The number of flagellar sections analyzed is shown above each bar. (*F*) Percentages of electron-
dense granules observed in the midpiece cross sections with TEM. The number of flagellar
sections analyzed is shown above each bar.

858
859
Figure 5. IPO5 and KPNB1 accumulate in *Fbxo24* KO spermatozoa. (*A*) Mass spectrometry
analyses of mature spermatozoa. Significantly upregulated proteins are shown with red dots
whereas significantly downregulated proteins are shown with blue dots. (*B*) Immunoblotting
analysis was performed using proteins extracted from testes or mature spermatozoa. IPO5 was
detected using rabbit anti-IPO5 polyclonal antibody. KPNA2 and IZUMO1 were detected as
negative control and loading control, respectively. (*C*) Spermatozoa obtained from the cauda
epididymis were stained for IPO5 (magenta) using rabbit anti-IPO5 polyclonal antibody. Nuclei
were stained with Hoechst 33342 (blue). (*D*) Immunoprecipitation (IP) of FBXO24-FLAG from
Fbxo24-FLAG Tg testes. FBXO24 could interact with IPO5. IPO5 was detected using mouse anti-
IPO5 monoclonal antibody. IPO5 band is slightly larger after IP likely due to different protein
composition in the sample. β-actin was used as a loading control. A picture with high contrast
was shown for the input FLAG band. (*E*) IP of FBXO24-FLAG using HEK293T cells. FBXO24
could interact with IPO5 but not with KPNB1. IPO5 was detected using mouse anti-IPO5
monoclonal antibody. β-actin was used as a loading control. (*F*) In vivo ubiquitination assay of
IPO5 using HEK293T cells. IPO5-PA was immunoprecipitated and the level of ubiquitination was
analyzed using anti-HA antibody. β-actin was used as a loading control.

876
877
Figure 6. KPNB1 and IPO5 are recruited to SGs. (*A*) Total RNA amounts in spermatozoa were
measured by ultraviolet absorption. (*B*) Electrophoresis of RNA extracted from mature
spermatozoa. (*C*) KPNB1 and IPO5 were localized to stress granules (SGs) under exposure to
oxidative stress. COS7 cells were treated with water (upper row) or arsenite (lower row). Nuclei
were stained with Hoechst 33342 (blue). (*D*) KPNB1 and IPO5 were localized to SGs under
exposure to a proteasome inhibitor. COS7 cells were treated with DMSO (upper row) or MG132
(lower row). Nuclei were stained with Hoechst 33342 (blue).