Title

FBXO24 ensures male fertility by preventing abnormal accumulation of membraneless granules in sperm flagella

Yuki Kaneda[1,2], Haruhiko Miyata[1,*], Zoulan Xu[1,2], Keisuke Shimada[1], Maki Kamoshita[1], Tatsuya Nakagawa[1,2], Chihiro Emori[1], Masahito Ikawa[1,2,3,4,5,*]

[1]Research Institute for Microbial Diseases, Osaka University, 3-1 Yamadaoka, Suita, Osaka 5650871 JAPAN.
[2]Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 5650871 JAPAN.
[3]The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 1088639 JAPAN.
[4]Center for Infectious Disease Education and Research (CiDER), Osaka University, 2-8 Yamadaoka, Suita, Osaka 5650871 JAPAN.
[5]Center for Advanced Modalities and DDS (CAMaD), Osaka University, 2-8 Yamadaoka, Suita, Osaka 5650871 JAPAN.
*Corresponding authors: Haruhiko Miyata and Masahito Ikawa

Contact info
Email: hmiya003@biken.osaka-u.ac.jp (HM); ikawa@biken.osaka-u.ac.jp (MI)

Abstract

Ribonucleoprotein (RNP) granules are membraneless electron-dense structures rich in RNAs and proteins, and involved in various cellular processes. Two RNP granules in male germ cells, intermitochondrial cement and the chromatoid body (CB), are associated with PIWI-interacting RNAs (piRNAs) and are required for transposon silencing and spermatogenesis. Other RNP granules in male germ cells, the reticulated body and CB remnants, are also essential for spermiogenesis. In this study, we disrupted FBXO24, a testis-enriched F-box protein, in mice and found numerous membraneless electron-dense granules accumulated in sperm flagella. *Fbxo24* knockout (KO) mice exhibited malformed flagellar structures, impaired sperm motility, and male infertility, likely due to the accumulation of abnormal granules. The amount and localization of known RNP granule-related proteins were not disrupted in *Fbxo24* KO mice, suggesting that the accumulated granules were distinct from known RNP granules. Further studies revealed that RNAs and two importins, IPO5 and KPNB1, abnormally accumulated in *Fbxo24* KO spermatozoa. In addition, IPO5 and KPNB1 were recruited to stress granules, RNP complexes, when cells were treated with oxidative stress or a proteasome inhibitor. These results suggest that FBXO24 plays a critical role in preventing the accumulation of importins and RNP granules in sperm flagella.

Keywords
CRISPR/Cas9; Spermiogenesis; Ribonucleoprotein granules

Introduction

Spermatogenesis is a specialized process by which spermatogonia differentiate into spermatocytes, round spermatids, elongating spermatids, and spermatozoa by undergoing meiosis and subsequent spermiogenesis. During spermiogenesis, round spermatids undergo dramatic morphological changes such as nuclear condensation, acrosome formation, flagellum formation, and cytoplasmic removal, leading to the formation of spermatozoa (Leblond and Clermont, 1952). To fertilize eggs, spermatozoa then travel a long distance to reach eggs and pass through the cumulus-cell layer and zona pellucida that encase eggs. The motility apparatus of spermatozoa is the flagellum which is divided into three parts, a midpiece, a principal piece, and an endpiece, in order from proximal to distal (Eddy et al., 2003). The midpiece contains accessory structures called the mitochondrial sheath and outer dense fibers (ODF) that surround the axoneme, a 9+2 arrangement of microtubules, while the principal piece contains a different accessory structure called the fibrous sheath and ODFs surrounding the axoneme. In contrast, the endpiece contains no accessory structures but the axoneme. Abnormal formation of these flagellar structures could lead to male infertility (Nsota Mbango et al., 2019).

Ribonucleoprotein (RNP) granules are membraneless electron-dense structures that assemble through liquid–liquid phase separation and play critical roles in spermatogenesis (Eddy, 1974). Among the RNP granules, intermitochondrial cement (IMC) observed in late spermatocytes and the chromatoid body (CB) found in round spermatids are involved in the PIWI-interacting RNA (piRNA) pathway (Meikar et al., 2011). After meiosis, IMCs are no longer detectable because of mitochondrial dispersion, and CBs become gradually larger in round spermatids. In elongating spermatids, the CBs containing MIWI, the main effector protein of the piRNA pathway, disappear and two other RNP structures called a reticulated body and a CB remnant that contain TSKS, TSSK1, and TSSK2 appear (Shang et al., 2010). Recently, we found that *Tsks* knockout (KO) mice failed to generate reticulated bodies and CB remnants, and were infertile with impaired cytoplasmic removal (Shimada et al., 2023), indicating that the formation and disassembly of RNP granules is highly regulated and important for spermiogenesis.

During spermiogenesis, post-translational modifications of proteins are critical since gene transcription is thought to cease with nuclear condensation. Ubiquitination is one of the post-translational modifications in which a small protein called ubiquitin is attached to target proteins. Ubiquitinated proteins are degraded by a ubiquitin-proteasome system (UPS), which is involved in various physiological phenomena in cells, such as cell cycle regulation and signal transduction

(Ciechanover and Schwartz, 1998; Mani and Gelmann, 2005; Zheng et al., 2016). Protein
ubiquitination is carried out by a cascade of enzymatic reactions, including ubiquitin-activating
enzyme (E1), ubiquitin-binding enzyme (E2), and ubiquitin ligase (E3), and target proteins
ubiquitinated by E3 ligase are degraded by a large protein complex called the 26S proteasome.
Among the approximately 600 E3 ligase genes estimated to be present in humans (Li et al.,
2008), the mechanism and functions of ubiquitination mediated by the SCF (Skp1-Cul1-F-box
protein) complex has been well studied (Skaar et al., 2013). The SCF complex consists of several
proteins such as SKP1, a cullin protein, an F-box protein, and RBX1, and can ubiquitinate various
substrates by using different F-box proteins that recognize distinct substrates. In humans, at least
38 F-box proteins have been identified, but the functions of most F-box proteins remain unclear
(Kipreos and Pagano, 2000).

In this study, we analyzed the function of *Fbxo24* (F-box protein 24), predominantly expressed in
testes, and found that FXBO24 is essential for sperm flagellar formation and male fertility.
Intriguingly, membraneless and electron-dense granules were found in *Fbxo24* KO sperm
flagella, suggesting that FBXO24 prevents the accumulation of abnormal RNP granules during
spermiogenesis.

Results

Fbxo24 is expressed predominantly in male germ cells after meiosis and interacts with SKP1

We performed RT-PCR using cDNAs obtained from multiple mouse tissues and found a specific
band for *Fbxo24* in testes (Figure 1A). To identify stages in which *Fbxo24* was expressed in
spermatogenesis, we performed RT-PCR using postnatal mouse testes over time. *Fbxo24*
signals increased after birth, with solid signals observed around postnatal day 21 (Figure 1B)
when round spermatids appear (Kluin et al., 1982). Consistent with the RT-PCR results, single-
cell transcriptome data (Hermann et al., 2018) indicates that mouse *Fbxo24* and its human
orthologue, *FBXO24*, are expressed during the post-meiotic stages (Figures S1A and B).
Pairwise sequence alignment of amino acids sequences revealed that FBXO24 is highly
conserved between mice and humans (88% amino acid identity) (Figure S1C) with the F-box
domains located near the N-terminus (Figure S1D). Because it has been reported that F-box
proteins bind to SKP1 via the F-box domain to form the SCF complex (Bai et al., 1996; Schulman
et al., 2000), we investigated the interaction of FBXO24 with SKP1 by co-immunoprecipitation
using HEK293T cells. We constructed vectors that express FBXO24 tagged with 3×FLAG with
(wild-type, WT) or without (ΔF) the F-box domain (Figure 1C). When SKP1-1D4 was
immunoprecipitated with an anti-1D4 antibody, WT FBXO24-FLAG was co-immunoprecipitated,
whereas ΔF FBXO24-FLAG was not (Figure 1D left). Conversely, when an anti-FLAG antibody
was used to immunoprecipitate WT or ΔF FBXO24-FLAG, SKP1-1D4 was co-immunoprecipitated
only with WT FBXO24-FLAG (Figure 1D right). These results suggest that FBXO24 could function
as a component of the SCF complex in mouse testis, during the post meiotic stages of
spermatogenesis.

Lack of *Fbxo24* causes male sterility in mice

To examine the role of *Fbxo24* in spermatogenesis, we generated *Fbxo24* KO mice using the
CRISPR/Cas9 system. Two gRNAs were designed to delete the majority of the coding region of
Fbxo24 (Figure 1E). One hundred fertilized eggs were electroporated, and the resulting 89 two-
cell stage embryos were transferred to the oviducts of pseudopregnant ICR mice. One out of 18
born pups contained the large deletion of the coding region. We found no overt gross defects in
development, behavior, or survival in *Fbxo24* KO mice. We performed genomic PCR using the
primers described in Figure 1E (Figure 1F) and confirmed that *Fbxo24* KO mice had a deletion of
11,575 bp by Sanger sequencing (Figure 1G). *Fbxo24* KO male mice were then mated with three
WT females for three months to analyze fertility. Although 20 vaginal plugs were detected, no

pups were born from the *Fbxo24* KO male mice (Figure 1H), suggesting that *Fbxo24* is
indispensable for male fertility.

Disruption of *Fbxo24* results in impaired spermiation and abnormal sperm morphology

We first examined spermatogenesis to determine the cause of male infertility in *Fbxo24* KO mice.
No apparent differences in gross testicular morphology (Figure S2A) or weights (Figure S2B)
were found between controls and *Fbxo24* KO mice. We then performed He-PAS staining of
testicular and epididymis sections (Figures 2A and S2C). Although elongating spermatids can be
found in *Fbxo24* KO testes, step 16 spermatids were still present in Stage IX seminiferous
tubules (Fig. 2A), indicating that spermiation was impaired in *Fbxo24* KO testes. In contrast, we
could not find apparent differences in the cross sections of the cauda epididymis between the two
genotypes (Figure S2C). Next, we observed mature spermatozoa collected from the cauda
epididymis. While KO spermatozoa showed comparable head morphology to the controls, KO
spermatozoa exhibited abnormal tail structures such as bent or coiled flagella (Figures 2B and C).
Further analyses of sperm head morphology with immunostaining showed no overt abnormalities
in the acrosome or nucleus (Figure S2D). Since abnormal flagellar morphology could result in
decreased motility, we performed computer-assisted sperm analysis (CASA) after 10 and 120
minutes of incubation in capacitating medium. CASA revealed that the percentages of motile
spermatozoa were significantly reduced in *Fbxo24* KO mice compared to the controls (Figure 2D
and Movies S1 and S2). Furthermore, all velocity parameters such as average path velocity
(VAP), straight-line velocity (VSL), and curvilinear velocity (VCL) were lower in *Fbxo24* KO
spermatozoa (Figures 2E-G). These results indicate that FBXO24 is critical in sperm flagellum
formation during spermiogenesis.

Fbxo24 KO spermatozoa cannot fertilize eggs *in vitro* and fail to migrate from the uterus into the oviduct

To further evaluate the fertilizing ability of *Fbxo24* KO spermatozoa, we performed in vitro
fertilization (IVF). Consistent with male sterility observed in vivo (Figure 1H), *Fbxo24* KO
spermatozoa could not fertilize eggs in vitro (Figure 3A). Furthermore, removing cumulus cells
(Figure 3B) or both cumulus cells and the zona pellucida (ZP) (Figure 3C) could not rescue the
fertilization rates, suggesting that *Fbxo24* KO spermatozoa may have defects in not only sperm
motility but also the acrosome reaction, which is a prerequisite for spermatozoa to pass through
the ZP and fuse with eggs (Morohoshi et al., 2023). Therefore, we analyzed the acrosome
reaction rates using transgenic (Tg) mice which express EGFP in the acrosome and DsRed2 in
the mitochondria (Red Body Green Sperm, RBGS, mice) (Hasuwa et al., 2010). RBGS
spermatozoa lose EGFP fluorescent signal when the spermatozoa undergo the acrosome
reaction. First, to test whether *Fbxo24* KO spermatozoa were viable, we performed propidium
iodide (PI) staining and found that the percentages of dead spermatozoa were significantly higher
in *Fbxo24* KO mice (Figure S3A). We then analyzed the acrosome reaction rates of PI-negative
live spermatozoa. While the control spermatozoa underwent the acrosome reaction after 120
minutes of incubation in a capacitation medium and with Ca^{2+} ionophore (A23187) treatment,
Fbxo24 KO spermatozoa rarely underwent the acrosome reaction even after adding A23187
(Figure S3B). These results indicate that *Fbxo24* KO spermatozoa show multiple defects, such as
lower viability, reduced sperm motility, and impaired acrosome reaction.

Next, we observed sperm migration in the female reproductive tract using RBGS mice. Although
the control spermatozoa migrate through the uterotubal junction (UTJ) four hours after mating,
Fbxo24 KO spermatozoa hardly passed through the UTJ (Figure 3D). Since the processing of
sperm membrane protein, ADAM3, is necessary for sperm migration through the UTJ (Fujihara
et al., 2019, 2018; Yamaguchi et al., 2006), we performed immunoblotting analyses and found
that ADAM3 was processed correctly even in *Fbxo24* KO spermatozoa (Figure 3E). Considering
that the acrosome reaction is not essential for sperm migration in the female reproductive tract
(Morohoshi et al., 2023) and defects in sperm motility could cause impaired migration through the

UTJ (Chung et al., 2014; Fujihara et al., 2018; Miyata et al., 2015; Shimada et al., 2019), these
results suggest that lower viability and decreased sperm motility can be the cause of abnormal
sperm migration and male sterility in vivo.

Next, we performed intracytoplasmic sperm injection (ICSI) to examine whether nuclei of *Fbxo24*
KO spermatozoa have the potential to generate the next generation. We injected *Fbxo24* KO
sperm heads into 91 WT oocytes, and obtained 48 two-cell stage embryos (Figure 3F). By
transplanting the two-cell embryos into the oviduct of pseudopregnant ICR females, we obtained
four pups, which were confirmed to have the heterozygous mutation by PCR (Figures 3G and H).
These results indicate that ICSI can rescue male sterility of *Fbxo24* KO mice.

FBXO24 is required for the sperm midpiece formation

We analyzed mitochondria localization in *Fbxo24* KO spermatozoa because abnormal
mitochondrial sheath structures were found in spermatozoa with bent tails, such as in *Armc12*
KO, *Tbc1d21* KO, and *Gk2* KO mice (Shimada et al., 2021, 2019). We observed midpieces using
RBGS Tg mice and revealed that mitochondria were abnormally segmented in *Fbxo24* KO
spermatozoa (Figure 4A). We also observed SEPT4, a component of the annulus (Kissel et al.,
2005), and found that SEPT4 localized to the proper region (Figure S4A).

We then used a scanning electron microscope (SEM) to observe the formation of mitochondrial
sheaths. During the early step of mitochondrial sheath formation, mitochondria wrap around the
axoneme. In both control and *Fbxo24* KO spermatids, spherical mitochondria were aligned
correctly at this step (Figure 4B, left panels). In the next step, mitochondria became crescent-
shaped and were interlocked in the control spermatids, whereas irregular interlocking was
observed in *Fbxo24* KO spermatids (Figure 4B, middle panels). Subsequently, elongating
mitochondria remain irregularly arranged in *Fbxo24* KO spermatids (Figure 4B, right panels).

To further investigate the morphological abnormality of the midpiece in *Fbxo24* KO mice, we
observed the ultrastructure of the cauda epididymal spermatozoa using transmission electron
microscopy (TEM). Consistent with the SEM results, we found that mitochondria were irregularly
arranged in *Fbxo24* KO spermatozoa (Figures 4C and D). The frequency of abnormal
mitochondria was significantly higher in *Fbxo24* KO spermatozoa (Figure 4E). In addition, the
axoneme (AX) and outer dense fiber (ODF) were disrupted in both midpieces and principal pieces
of *Fbxo24* KO spermatozoa (Figures S4B-E). Furthermore, we frequently observed numerous
membraneless electron-dense granules in the midpieces of *Fbxo24* KO spermatozoa (Figures
4C, D, and F). We found these granules even in the axoneme (Figure. 4C). Despite the low
frequency, we also found membraneless electron-dense granules in the principal pieces of
Fbxo24 KO spermatozoa (Figures S4F and G).

Because membraneless electron-dense granules accumulated in *Fbxo24* KO spermatozoa, we
examined RNP granule-related proteins in *Fbxo24* KO mice. We performed immunoblotting
analyses for ADAD1, ADAD2, MILI, MIWI, RNF-17, YTHDC2, TSKS, and TSSK1, which localized
in germ-cell RNP granules and are essential for spermatogenesis (Lu et al., 2023; Meikar et al.,
2011; Pan et al., 2005; Shang et al., 2010; Shimada et al., 2023; Snyder et al., 2020), but we did
not see any differences in the amounts of these proteins between *Fbxo24* heterozygous and KO
testes (Figure S5A). Further, we found no abnormalities in the localization of MIWI in *Fbxo24* KO
testes (Figure S5B). MIWI was localized in the IMCs of late spermatocytes and CBs of round
spermatids and disappeared in elongating spermatids (Figure S5B). Further, no abnormalities
were found in the immunostaining of TSKS, which localized in the reticulated body and CB
remnant of elongating spermatids (Figure S5C).

IPO5 and KPNB1 amounts increase in *Fbxo24* KO flagella

Since previous studies show that loss of F-box protein causes the accumulation of substrates
(Nakayama et al., 2000; Tsunematsu et al., 2004), we investigated whether certain proteins
accumulated in *Fbxo24* KO spermatozoa. Mass spectrometry analyses revealed that the amounts
of several proteins significantly increased in *Fbxo24* KO mature spermatozoa (Figure 5A). Among
the significantly increased proteins, we focused on IPO5 (importin 5, also known as KPNB3 and
RanBP5) and KPNB1 (karyopherin subunit beta 1, also known as importin β1) because the fold
changes (KO/WT) are the highest in these two proteins. Both IPO5 and KPNB1 are members of
the karyopherin-β (KPNB) family of nuclear transport receptors (NTRs), which play a crucial role
in nucleocytoplasmic transport (Chook and Blobel, 2001; Chook and Süel, 2011; Conti and
Izaurralde, 2001; Kimura and Imamoto, 2014). In rodent testes, KPNB1 localized to the cytoplasm
of spermatogonia, spermatocytes, and Sertoli cells, and IPO5 localized to the cytoplasm of
elongating spermatids (Loveland et al., 2015, 2006). Immunoblotting analyses confirmed that the
amounts of IPO5 and KPNB1 increased in *Fbxo24* KO mature spermatozoa (Figure 5B). In
contrast, the amounts of KPNA2 (Importin α1), a member of karyopherin-α (KPNA) family of
NTRs which localized to the cytoplasm and nucleus of spermatocytes and the cytoplasm of
elongating spermatids (Miyamoto et al., 2013), were comparable between the control and *Fbxo24*
KO spermatozoa. Since the anti-KPNB1 antibody did not work for immunostaining, we performed
immunostaining of mature spermatozoa using an anti-IPO5 antibody. We detected IPO5 in the
Fbxo24 KO flagella but not in control (Figure 5C).

FBXO24 could interact with IPO5

Next, we investigated the proteins that interacted with FBXO24. Because we could not obtain
anti-FBXO24 antibodies that worked for immunoprecipitation, we generated Tg mice expressing
3×FLAG-tagged *Fbxo24* under a testis-specific *Prm1* promoter (Figures S6A and B). This
transgene could not rescue *Fbxo24* KO sperm morphology (Figure S6C). By looking for long non-
coding RNAs, we found *Gm36266* that is located around Exon 9 and 10 of *Fbxo24*, which is
deleted in *Fbxo24* KO mice (Figure 1E); however, a partial deletion of Exon 2 and 3 of *Fbxo24*
showed the same defects in sperm morphology (Figures S6D–G) as mice lacking Exon 3-10
(Figure 2B), suggesting that *Fbxo24*, not *Gm36266*, is responsible for the phenotypes observed
in this study. Abnormal sperm morphology was not rescued by the transgene, likely due to lower
expression of FBXO24 (Figures 5D and S6B) and/or FLAG-tag interfering with FBXO24 function.
Using Tg mice, we could immunoprecipitate FBXO24-FLAG with an anti-FLAG antibody, and
subsequent mass spectrometry analyses detected not only FBXO24 but also IPO5 with the
highest quantitative values (Figure S6H), indicating that these proteins interact in vivo. We also
found SKP1 in this analysis, consistent with the in vitro study (Figure 1D). We confirmed that
IPO5 co-immunoprecipitated with FBXO24-FLAG in mouse testes using immunoblotting analysis
(Figure 5D). Further, we transiently expressed 3×FLAG-tagged *Fbxo24* in HEK293T cells,
performed immunoprecipitation analysis using an anti-FLAG antibody, and found that FBXO24
could interact with endogenous IPO5 but not with endogenous KPNB1 (Figure 5E). These results
suggest that FBXO24 may recognize IPO5 for subsequent protein degradation.

IPO5 is recruited into RNP granules under stress conditions

Although the amounts of known RNP granule-related proteins were not upregulated in *Fbxo24*
KO testes (Figure S5A) and spermatozoa (Figure 5A), the total amount of RNAs was significantly
increased in *Fbxo24* KO spermatozoa (Figures 6A and B), suggesting that the electron-dense
granules observed in the *Fbxo24* KO spermatozoa may contain RNAs. Previous studies showed
that various stresses, such as heat shock, arsenite treatment, or proteasome inhibition, trigger the
formation of cytoplasmic stress granules (SGs), membraneless granules composed of RNPs.
Further, it has been shown that SGs could contain KPNB1 (Chang and Tarn, 2009; Mahboubi et
al., 2013). Increased amounts of KPNB1 and RNAs suggest that RNP granules may be formed in
Fbxo24 KO spermatozoa. We analyzed whether IPO5 can localize to SGs under stress
conditions to explore this possibility. We examined the subcellular localization of endogenous
IPO5 and KPNB1 in COS7 cells treated with arsenite, an oxidative stress inducer.

Immunostaining analyses revealed that IPO5 was predominantly localized to the cytoplasm
without the stress inducer (Figure 6C); however, when cells were treated with arsenite, IPO5 was
detected not only in the cytoplasm but also in cytoplasmic granules that were colocalized with
KPNB1 (Figure 6C). We then analyzed if the stress of proteasome inhibition could cause a similar
response. We performed immunostaining of IPO5 and KPNB1 in COS7 cells using a proteasome
inhibitor, MG132. Consistent with arsenite, MG132 caused the accumulation of cytoplasmic
granules that contained IPO5 and KPNB1 (Figure 6D). These results support the idea that RNP
granules that contain KPNB1 and IPO5 are formed in *Fbxo24* KO spermatozoa.

Discussion

Spermiogenesis is the late stage of spermatogenesis in which round spermatids differentiate into
spermatozoa accompanied by drastic morphological changes such as flagellum formation,
chromatin remodeling, and cytoplasmic removal. Protein ubiquitination of target proteins for
degradation by the UPS is crucial to maintain functional spermatogenesis (Richburg et al., 2014).
In this study, we focused on testis-enriched F-box protein FBXO24. We confirmed that *Fbxo24*
was predominantly expressed in the testis, and its expression level increased in the late stage of
spermatogenesis. The analyses using cultured cells and Tg testes suggested that FBXO24 could
interact with SKP1 to form the SCF complex. Further, we deleted *Fxbo24* using the
CRISPR/Cas9 system and found that *Fbxo24* KO mice exhibited abnormal flagellar structures
and male infertility. *Fxbo24* KO spermatozoa failed to fertilize eggs even with IVF, while ICSI
gave offspring.

More detailed observation of the midpiece of *Fbxo24* KO spermatozoa using electron microscopy
revealed abnormalities in mitochondria coiling, which occurred in the late stages of
spermatogenesis and was consistent with the *Fbxo24* expression timing. Furthermore, *Fbxo24*
KO spermatozoa showed markedly reduced motility. These results suggest that the primary
cause of infertility is impaired sperm migration in the female reproductive tract due to reduced
sperm motility. In addition to the abnormal flagellar morphology, *Fbxo24* KO spermatozoa could
not undergo the acrosome reaction even after the A23187 treatment. Because IZUMO1 staining
showed no defects in the acrosome morphology, acrosome reaction failure may be caused by
impaired sperm capacitation associated with abnormal flagellar structures. It is also possible that
FBXO24 is involved in the post-translational modification of proteins required for the acrosome
reaction.

The TEM analyses observed numerous membraneless electron-dense granules in *Fbxo24* KO
sperm flagella. Although the irregular arrangement of sperm mitochondria has been reported in
KO mice of some mitochondria-associated proteins (Shimada et al., 2021, 2019), numerous
granules have not been observed in these KO spermatozoa. Proteomic analyses revealed that
IPO5 and KPNB1 accumulated in *Fbxo24* KO spermatozoa without significant elevation of
mitochondria-associated proteins. The lack of accumulation of KPNA2, which is part of the
importin family, also suggests that there is selectivity in the accumulated proteins. Abnormal
accumulation of proteins, including IPO5 and KPNB1, may lead to morphological abnormalities in
sperm flagella since the SCF complex is critical for protein degradation and consequently
regulate various cellular processes (Skaar et al., 2013). FBXO24 likely plays a role in catalyzing
protein degradation during spermiogenesis. Because FBXO24 could interact with IPO5 in testes,
FBXO24 may catalyze IPO5 degradation directly. Recently, mice lacking both PSME4 and
ECPAS, proteasome-associated proteins, were shown to exhibit impaired male fertility with
disorganized mitochondrial sheath, but numerous membraneless granules were not observed
(Sato et al., 2023), suggesting that the accumulation of the granules is specific to *Fbxo24* deletion
rather than simply caused by proteasome defects.

In addition to proteins, the total RNA amount increased in *Fbxo24* KO spermatozoa, suggesting
that the abnormal granules observed in *Fbxo24* KO flagella consist of RNPs. In germ cells, RNP
granules have been well studied and shown to be essential for spermatogeneses, such as IMCs,

CBs, reticulated bodies, and CB remnants (Chuma et al., 2006; Meikar et al., 2011; Shimada et
al., 2023). RNP granules are membraneless electron-dense structures that are similar to the
abnormal granules in *Fbxo24* KO; however, the localization of MIWI, an IMC and CB marker, and
TSKS, a reticulated body and CB remnant marker, was not disrupted in the KO. These results
suggest that electron-dense granules accumulated in *Fbxo24* KO spermatozoa are different from
well-studied RNP granules. SGs are also membraneless electron-dense granules that contain
RNPs, which assemble through liquid–liquid phase separation in response to cellular stresses
(Buchan and Parker, 2009; Protter and Parker, 2016; Taylor et al., 2016). While transient SGs
formed in response to oxidative stress or heat shock play roles in translational arrest to ensure
cell survival, chronic SGs are relevant to neurodegenerative diseases (Marcelo et al., 2021;
Reineke and Neilson, 2019). It has been shown that, under arsenite-induced oxidative stress,
KPNB1 relocates to SGs (Chang and Tarn, 2009; Mahboubi et al., 2013). Because we found that
not only KPNB1 but also IPO5 can be recruited to SGs after arsenite or proteasome inhibitor
treatment, electron-dense granules observed in *Fbxo24* KO spermatozoa may be RNP granules
consisting of IPO5 and KPNB1. FBXO24 may be responsible for degrading proteins such as
IPO5 to prevent the accumulation of abnormal RNP granules, which may cause deformation of
the sperm flagella.

In conclusion, we reveal that numerous RNP granules accumulated in sperm flagella when
FBXO24, which is related to the UPS was depleted. Small numbers of the RNP granules were
found even in the control, suggesting that the ability to generate the granules is present in WT
testes but may be suppressed by FBXO24. FBXO24 can be a key molecule for understanding the
formation and function of the aberrant RNP granules in male fertility. Because amino acid
sequences and expression patterns of FBXO24 are conserved, FBXO24 may also play similar
roles in human testes. Our research may lead to development of new approaches to infertility
treatment and nonhormonal male contraceptives.

Materials and Methods
Animals. All animal experiments were approved by the Animal Care and Use Committee of the
Research Institute for Microbial Diseases, Osaka University. Mice were purchased from CLEA
Japan (Tokyo, Japan) or Japan SLC (Shizuoka, Japan). WT or *Fbxo24* heterozygous (HET) mice
were used as controls. All gene-modified mice generated in this study will be made available
through either the RIKEN BioResource Research Center or the Center for Animal Resources and
Development (CARD), Kumamoto University.

Isolation of RNA and RT-PCR. Adult mouse multi-tissues and mouse testes at different ages
were obtained from C57BL/6N mice. RNA samples were isolated and purified
using TRIzol (Thermo Fisher Scientific, Waltham, MA, USA). RNA was reverse transcribed to
cDNA using SuperScript IV First-Strand Synthesis System (Thermo Fisher Scientific) using an
oligo (dT) primer. PCR was performed using the KOD Fx Neo DNA Polymerase (Toyobo, Tokyo,
Japan). Primers used in this study are listed in Table S1.

Transfection of HEK293T cells and induction of stress. HEK293T cells (Tiscornia et al., 2006)
and COS7 cells were maintained in Dulbecco's Modified Eagle Medium (DMEM) supplemented
with 10% fetal bovine serum (Sigma-Aldrich, St. Louis, MO, USA) and 1% Gibco™
penicillin/streptomycin (Thermo Fisher Scientific) at 37°C under 5% CO_2. For transfection,
HEK293T cells were transiently transfected with the plasmid DNA using the calcium phosphate
transfection method (Tiscornia et al., 2006), and cultured for 24 h before harvesting. For oxidative
stress, COS7 cells were treated with 0.5 mM sodium arsenite in medium for 30 min and controls
were incubated with water in medium. For proteasome inhibition, COS7 cells were treated with 10
µM sodium MG132 (Sigma-Aldrich) in medium for 3 h and controls were incubated with DMSO in
medium.

Generation of *Fbxo24* KO mice. *Fbxo24* KO mice were generated using CRISPR/Cas9. Three crRNAs, 5′- TGTGGAGGCGCATCTGTCGA -3′, 5′- TCCTGAAGGAAGTCGAGCCG -3′ and 5′-TCAGTTGTTCCCCCCAGAGC -3′ were designed using the online source CRISPRdirect (Naito et al., 2015) and annealed to SygRNA™ Cas9 Synthetic tracrRNA (#TRACRRNA05N-5NMOL, Sigma-Aldrich). The gRNAs were mixed with TrueCut™ Cas9 Protein v2 (A36498, Thermo Fisher Scientific) and incubated at 37°C for 5 min to form CRISPR/Cas9 complexes. The complexes were introduced into fertilized eggs which were from super-ovulated WT B6D2F1 females mated with B6D2F1 males. Electroporation was performed using the super electroporator NEPA21 (NEPA GENE, Chiba, Japan) (poring pulse, voltage: 225 V, pulse width: 2 ms, pulse interval: 50 ms, number of pulses: +4, and attenuation: 10%; transfer pulse, voltage: 20 V, pulse width: 50 ms, pulse interval: 50 ms, number of pulses: ±5, and attenuation: 40%). The treated eggs were developed into two-cell-stage embryos by cultivating in KSOM medium (Ho et al., 1995) and transplanted into pseudo-pregnant ICR females. The obtained pups were genotyped by PCR to detect the KO and/or WT allele and then subjected to Sanger sequencing to verify the deleted sequence.

Fertility test. For the in vivo fertility analysis, sexually mature KO male mice or WT male mice were caged with three 8-week-old B6D2F1 female mice for three months and plugs were checked every morning. The number of pups was counted on the day of birth. For the in vitro fertility assay, in vitro fertilization (IVF) analysis was performed as previously described (Morohoshi et al., 2021) with some minor changes. For ZP-free oocytes, sperm insemination was performed at a final density of 2×10^4 spermatozoa/mL.

Histological analysis. PAS staining of sections were performed as previously described (Morohoshi et al., 2020). Testes or cauda epididymis were fixed at 4°C in Bouin's solution (Polysciences, Inc., Warrington, PA, USA) and were processed for paraffin embedding. Paraffin sections were cut at a thickness of 5 µm using an HM325 microtome (Microm, Walldorf, Germany). After rehydrating the sections, they were stained with 1% periodic acid (Nacalai Tesque, Kyoto, Japan) and Schiff's reagent (FUJIFILM WakoPure Chemical, Osaka, Japan) for 20 min each at room temperature. The sections were then counterstained with Mayer's hematoxylin solution (FUJIFILM WakoPure Chemical). The sections were observed with an Olympus BX-53 microscope (Tokyo, Japan).

Morphological and motility analysis of spermatozoa. Spermatozoa extracted from cauda epididymis were suspended in TYH medium (Muro et al., 2016). After 10 min incubation, spermatozoa were collected to observe morphology. Sperm motility was analyzed as previously described (Miyata et al., 2021, 2020). The motility of more than 200 spermatozoa was measured after incubation at 10 and 120 min in TYH medium using CEROS II (software version 1.4; Hamilton Thorne Biosciences, Beverly, MA, USA).

Observation of sperm flagellum ultrastructure using TEM. Cauda epididymis specimens were prepared as previously described (Shimada et al., 2019). The prepared samples were observed using a JEM-1400 plus electron microscope (JEOL, Tokyo, Japan) at 80 kV with a CCD Veleta 2K × 2K camera (Olympus).

Observation of sperm mitochondria during spermatogenesis using SEM. Testes specimens were prepared as previously described (Shimada et al., 2019). The prepared samples were observed using a S-4800 field emission scanning electron microscope (Hitachi, Tokyo, Japan).

Pfam domain search. Pfam domains were detected from amino acid sequences using Simple Modular Architecture Research Tool (SMART) (http://smart.embl-heidelberg.de/). Mouse FBXO24 amino acid sequence (CCDS51674.1) and human FBXO24 (CCDS5698.1) were obtained from the CCDS database (https://www.ncbi.nlm.nih.gov/CCDS/CcdsBrowse.cgi).

Alignment of amino acid sequences. Amino acid sequences of Mouse FBXO24
(CCDS51674.1) and human FBXO24 (CCDS5698.1) were aligned using "Clustal Omega
(https://www.ebi.ac.uk/Tools/msa/clustalo/)". Aligned sequences were edited using Jalview ver.
2.11.2.0 (https://www.jalview.org/). Domains within the sequences were identified using "SMART
(http://smart.embl-heidelberg.de/)".

In silico **expression data analysis.** Single-cell transcriptome data in the mouse and
human testis was obtained from previously published work (Hermann et al., 2018). *Fbxo24*
expression in those cells was analyzed using the Loupe Cell Browser 3.3.1 (10X
Genomics, Pleasanton, CA, USA).

Plasmids. The open reading frames (ORFs) of *Fbxo24* and *Skp1* were cloned and amplified
using cDNA obtained from mouse testis and inserted into the multiple cloning site of pCAG1.1
vector (Addgene; Plasmid #173685). To generate the expression vector coding *Fbxo24(ΔF-box),*
inverse PCR was performed with KOD -Plus- Mutagenesis Kit (Toyobo, Tokyo, Japan) according
to the manufacturer's instructions. Primers used for inverse PCR are listed in Table S1.

Observation of spermatozoa migration inside the female reproductive tract. The *Fbxo24* KO
mouse line was crossed with B6D2 Tg mice carrying CAG/Su9-DsRed2, Acr3-EGFP (RBGS)
(Hasuwa et al., 2010) to label the acrosome with EGFP. B6D2F1 females were superovulated by
injecting pregnant mare serum gonadotropin (PMSG, ASKA Pharmaceutical, Tokyo, Japan)
followed by human chorionic gonadotropin (hCG, ASKA Pharmaceutical) 48 h appart. The
superovulated female was mated with a WT male or *Fbxo24* KO male mice 14 h after the hCG
injection. Female mice were sacrificed 4 h after confirming a vaginal plug and the female
reproductive tracts were collected. Spermatozoa inside the oviducts were observed using a BZ-
X710 microscope (Keyence Japan, Osaka, Japan).

Protein extraction from testes, spermatozoa, and culture cells. HEK293T cells were lysed in
1% Triton X-100 lysis buffer [1% Triton X-100, 50 mM Tris-HCl pH 7.4, 150 mM NaCl, and 1%
(v/v) protease inhibitor cocktail (Nacalai Tesque)] at 4°C with end-over-end rotation for 2 h.
Testes or spermatozoa were homogenized in 1% Triton X-100 lysis buffer at 4°C with end-over-
end rotation for 2 h or sample buffer containing 1 M Tris-HCL pH 6.8, 2% SDS, 10% Glycerol,
and 0.005% Bromophenol Blue, and boiled for 5 min. Supernatants were obtained after
centrifugation at 15,300 × g for 15 min at 4°C.

Immunoprecipitation. Immunoprecipitation was performed as described previously (Kaneda et
al., 2023). Proteins from HEK293T cells were extracted using 1% Triton X-100 lysis buffer [1%
Triton X-100, 50 mM Tris-HCl pH 7.4, 150 mM NaCl, and 1% (v/v) protease inhibitor cocktail
(Nacalai Tesque)]. The lysates were centrifuged at 15,300 g for 5 min at 4°C to collect
supernatants. Protein lysates were mixed with 20 µL Protein G-conjugated magnetic beads
(#10009D, Thermo Fisher Scientific) preincubated with 1.0 µg antibody. After incubation for 1 h at
4°C, the beads were washed three times with wash buffer (40 mM Tris-HCl pH 7.5, 150 mM
NaCl, 0.1% Triton X-100 and 10% glycerol). For immunoblotting, the immune complexes were
eluted with 2×SDS sample buffer (132 mM Tris HCl pH7.4, 4% SDS, 20% glycerol, and 0.01%
Bromophenol Blue) for 10 min at 70°C. For mass spectrometry (MS), the immune complexes
were eluted with 150 ng/µL 3×FLAG peptide (Sigma-Aldrich) in TBS solution for 30 min at 4°C.
Antibodies used in this study are listed in Table S2.

Immunoblotting. Proteins were separated with sodium dodecyl sulfate polyacrylamide gel
electrophoresis (SDS-PAGE) under reducing conditions using 2-ME and transferred onto
polyvinylidene difluoride (PVDF) membrane using the Trans Blot Turbo system (Bio-Rad,
Hercules, CA, USA). After blocking with 10% skim milk (Becton Dickinson, Franklin Lakes, NJ,
USA) in TBST, the blots were incubated with primary antibody overnight at 4°C, and then
incubated with HRP-conjugated secondary antibodies (1:5000) for 2 h at room temperature.

462 Chemiluminescence was detected with Chemi-Lumi One Super (#02230, Nacalai Tesque) or
463 Chemi-Lumi One Ultra (#11644, Nacalai Tesque).

Assessment of sperm viability in vitro. Spermatozoa extracted from cauda epididymis were
465 dispersed in TYH medium and incubated for 10 and 120 min at 37°C under 5% CO_2. PI was
466 carefully added to the TYH drop to achieve a final concentration of 10 µg/ml and a small volume
467 of the sperm suspension was placed on a glass slide. More than 200 spermatozoa were counted
468 for each trial.

Assessment of acrosome reaction in vitro. The *Fbxo24* KO mouse line was crossed with
B6D2 Tg mice carrying CAG/Su9-DsRed2, Acr3-EGFP (RBGS) (Hasuwa et al., 2010) to label the
acrosome with EGFP. Spermatozoa extracted from cauda epididymis were dispersed in TYH
medium for 10 and 120 min of incubation at 37°C under 5% CO_2. After 120 min of incubation,
spermatozoa were incubated with Ca^{2+} ionophore A23187 (Merck, Rahway, NJ, USA) at 20 µM to
induce the acrosome reaction at 37°C under 5% CO_2. 10 mM stock solution of Ca^{2+} ionophore
A23187 in DMSO was diluted 10 times with TYH medium before carefully adding to the sperm
suspension to reach the final concentration. PI (10 µg/ml at final concentration) was added to the
TYH drop, and a small volume of the sperm suspension was placed on a glass slide. Acrosome-
reacted spermatozoa were determined by observing EGFP signals while distinguishing viable
spermatozoa with PI staining using a BX-53 microscope (Olympus). More than 200 spermatozoa
were counted for each trial.

Intracytoplasmic sperm injection (ICSI). B6D2F1 females were superovulated by injecting
PMSG (ASKA Pharmaceutical) followed by hCG (ASKA Pharmaceutical) 48 h later. MII oocytes
were collected 14 h after injection of hCG. Cumulus cells were removed from the collected eggs
after 10 min treatment with 330 µg/mL hyaluronidase (Sigma-Aldrich). Cumulus-free oocytes
were placed in KSOM medium at 37°C under 5% CO2 until just before performing ICSI. Each
sperm head separated from the tail by applying a few piezo pulses was injected into a cumulus-
free MII oocyte using a piezo manipulator (PrimeTech, Tokyo, Japan). Obtained two-cell embryos
were transferred to pseudopregnant ICR females the next day.

Immunofluorescence of testis cross-sections. Testes were fixed in 4% paraformaldehyde in
PBS for 3 h at 4°C soon after dissection and infused with 15% and 30% sucrose in PBS at 4°C.
The testes were embedded in OCT medium (Tissue-Tek, Miami, FL, USA) and frozen in liquid
nitrogen. Testis blocks were sectioned at 10 µm thickness using a cryostat and the sections were
dried on microscope slides at 42°C to make them adhere. The testis sections were then
processed and imaged in the same manner as sperm immunostaining. After permeabilization with
0.1% Triton X-100 in PBS for 15 min, the sections were blocked with 3% bovine serum albumin
(Sigma-Aldrich), 10% goat serum, and 0.1% Triton X-100 in PBS for 1 h at room temperature.
The sections were then incubated with primary antibody overnight at 4°C. After washing the
sections with PBS for 10 min, the sections were secondarily blocked with 3% bovine serum
albumin (Sigma-Aldrich), and 10% goat serum in PBS for 1 h at room temperature. The sections
were further incubated with fluorophore-conjugated secondary antibodies (1:2000) and Alexa
Fluor 568 Dye PNA Lectin (1:2000) for 2 h at room temperature. After a wash with PBS, the
sections were incubated with Hoechst 33342 (Thermo Fisher Scientific) in PBS (1:5000) for 5 min
at room temperature. Once washed with PBS, the sections were mounted with Shandon Immu-
Mount (Thermo Fisher Scientific) before imaging.

Immunostaining of spermatozoa. Spermatozoa extracted from cauda epididymis were
dispersed in PBS. A small aliquot of the sperm suspension was spotted onto glass slides and
then air-dried at room temperature. Spermatozoa were then fixed with 4% paraformaldehyde in
PBS for 10 min and washed three times with PBS for 5 min each. The slides were blocked with
blocking solution (5% bovine serum albumin and 10% goat serum in PBS) for 1 h at room
temperature. The slides were incubated with primary antibody overnight at 4°C, washed with PBS
three times for 5 min each, and incubated with Alexa Fluor conjugated secondary antibody for 2 h

at room temperature. The slides were washed with PBS three times for 5 min each, incubated
with Hoechst 33342 (2 µg/ml) (Thermo Fisher Scientific) for 5 min, and washed with PBS three
times for 5 min each. The slides were mounted with Shandon Immu-Mount (Thermo Fisher
Scientific) before imaging. Fluorescent images were captured with an Olympus BX-53
microscope (Olympus).

Immunofluorescence of COS7 cells. COS7 cells were fixed with 4% paraformaldehyde in PBS
for 15 min at room temperature. After permeabilization with 0.1% Triton X-100 in PBS for 15 min,
cells were blocked with 1% bovine serum albumin (Sigma-Aldrich) in PBS for 1 h at room
temperature. The cells were then incubated with primary antibody overnight at 4°C. After washing
three times with PBS, the cells were further incubated with fluorophore-conjugated secondary
antibodies (1:1000) for 2 h at room temperature. After washing three times with PBS, the cells
were incubated with Hoechst 33342 (Thermo Fisher Scientific) in PBS (1:5000) for 5 min at room
temperature. After washing three times with PBS, the cells were mounted with Shandon Immu-
Mount (Thermo Fisher Scientific) before imaging.

Extraction of RNA from epididymal spermatozoa. Spermatozoa collected from cauda
epididymis were dispersed in PBS and the number of spermatozoa was counted using a
hemacytometer. Spermatozoa were centrifuged at 15,300 g for 2 min and the sperm pellet was
lysed in TRIzol (Thermo Fisher Scientific). Chloroform was added to the lysate and incubated for
5 min at room temperature. The sample was centrifuged at 15,300 g for 15 min and the aqueous
phase was transferred to a fresh centrifuge tube. To precipitate RNA, an equal volume of
isopropanol was added to the aqueous phase and centrifuged at 15,300 g for 20 min at 4°C. The
supernatant was removed and the pellet was washed with 75% Ethanol. The washed pellet was
dried and resuspended in nuclease-free water. The RNA solution was DNAse treated using
deoxyribonuclease (RT Grade) (NIPPON GENE, Tokyo, Japan) to remove genomic DNA. Then
the RNA solution was incubated at 55°C for 10 min. The sample was stored at -80°C. The
concentration was measured using NanoDrop One (Thermo Fisher Scientific). The
TAE/formamide method (Masek et al., 2005) was used to perform electrophoresis of RNA.

Statistical analysis. All statistical analyses were performed using the two-tailed Welch's t-test
using Microsoft Office Excel 2016 (Microsoft Corporation, Redmond, WA, USA). Differences were
considered significant at $P<0.05$ (*), $P<0.01$ (**), $P<0.001$ (***). Error bars are standard deviation.

Acknowledgments
The authors would like to thank Dr. Julio M. Castaneda for the critical reading of the manuscript
and Ms. Natsuki Furuta for technical assistance, and the members of both the Department of
Experimental Genome Research and the nonprofit organization (NPO) for Biotechnology
Research and Development for experimental assistance and discussion. We also thank Dr.
Yoichi Miyamoto for kindly providing antibodies against importins (National Institutes of
Biomedical Innovation, Health and Nutrition), and Dr. Akinori Ninomiya and Dr. Fuminori Sugihara
for MS analysis (Core Instrumentation Facility, Research Institute for Microbial Diseases, Osaka
University). This research was supported by the Ministry of Education, Culture, Sports, Science
and Technology (MEXT)/Japan Society for the Promotion of Science (JSPS) KAKENHI grants
(JP23KJ1523 to Y.K., JP22H03214, JP23K18328 to H.M., JP23K05831 to K.S., JP22K15030 to
M.K., and JP19H05750, JP21H04753, JP21H05033 to M.I.); Takeda Science Foundation grant to
H.M., K.S., and M.I.; JST FOREST (JPMJFR211F to H.M.); the Eunice Kennedy Shriver National
Institute of Child Health and Human Development (P01HD087157 and R01HD088412 to M.I.);
and the Bill & Melinda Gates Foundation (Grand Challenges Explorations grant INV-001902 to
M.I.).
558
559

560 ## Author Contributions

561   Y.K., H.M., and M.I. conceived and designed experiments and wrote the manuscript with input
562 from all authors. All authors performed experiments and analyzed data.
563

Competing interests

565 The authors declare no competing interests.

566
567

References

Bai C, Sen P, Hofmann K, Ma L, Goebl M, Harper JW, Elledge SJ. 1996. SKP1 Connects Cell Cycle
Regulators to the Ubiquitin Proteolysis Machinery through a Novel Motif, the F-Box. *Cell*
86:263–274. doi:10.1016/S0092-8674(00)80098-7

Buchan JR, Parker R. 2009. Eukaryotic Stress Granules: The Ins and Outs of Translation. *Mol Cell*.
doi:10.1016/j.molcel.2009.11.020

Chang WL, Tarn WY. 2009. A role for transportin in deposition of TTP to cytoplasmic RNA
granules and mRNA decay. *Nucleic Acids Res* **37**:6600–6612. doi:10.1093/nar/gkp717

Chook Y, Blobel G. 2001. Karyopherins and nuclear import. *Curr Opin Struct Biol* **11**:703–715.
doi:10.1016/S0959-440X(01)00264-0

Chook YM, Süel KE. 2011. Nuclear import by karyopherin-βs: Recognition and inhibition. *Biochim*
Biophys Acta Mol Cell Res. doi:10.1016/j.bbamcr.2010.10.014

Chuma S, Hosokawa M, Kitamura K, Kasai S, Fujioka M, Hiyoshi M, Takamune K, Noce T,
Nakatsuji N. 2006. *Tdrd1/Mtr-1* , a *tudor* -related gene, is essential for male germ-cell
differentiation and nuage/germinal granule formation in mice. *Proc Natl Acad Sci U S A*
103:15894–15899. doi:10.1073/pnas.0601878103

Chung J-J, Shim S-H, Everley RA, Gygi SP, Zhuang X, Clapham DE. 2014. Structurally Distinct Ca2+
Signaling Domains of Sperm Flagella Orchestrate Tyrosine Phosphorylation and Motility.
Cell **157**:808–822. doi:10.1016/j.cell.2014.02.056

Ciechanover A, Schwartz AL. 1998. The ubiquitin-proteasome pathway: The complexity and
myriad functions of proteins death. *Proc Natl Acad Sci U S A* **95**:2727–2730.
doi:10.1073/pnas.95.6.2727

Conti E, Izaurralde E. 2001. Nucleocytoplasmic transport enters the atomic age. *Curr Opin Cell*
Biol **13**:310–319. doi:10.1016/S0955-0674(00)00213-1

Eddy EM. 1974. Fine structural observations on the form and distribution of nuage in germ cells
of the rat. *Anat Rec* **178**:731–757. doi:10.1002/ar.1091780406

Eddy EM, Toshimori K, O'Brien DA. 2003. Fibrous sheath of mammalian spermatozoa. *Microsc*
Res Tech **61**:103–115. doi:10.1002/jemt.10320

Fujihara Y, Miyata H, Ikawa M. 2018. Factors controlling sperm migration through the oviduct
revealed by gene-modified mouse models. *Exp Anim* **67**:91–104. doi:10.1538/expanim.17-
0153

Fujihara Y, Noda T, Kobayashi K, Oji A, Kobayashi S, Matsumura T, Larasati T, Oura S, Kojima-Kita
K, Yu Z, Matzuk MM, Ikawa M. 2019. Identification of multiple male reproductive tract-
specific proteins that regulate sperm migration through the oviduct in mice. *Proc Natl Acad*
Sci U S A **116**:18498–18506. doi:10.1073/pnas.1908736116

HASUWA H, MURO Y, IKAWA M, KATO N, TSUJIMOTO Y, OKABE M. 2010. Transgenic Mouse
Sperm that Have Green Acrosome and Red Mitochondria Allow Visualization of Sperm and
Their Acrosome Reaction in Vivo. *Exp Anim* **59**:105–107. doi:10.1538/expanim.59.105

Hermann BP, Cheng K, Singh A, Roa-De La Cruz L, Mutoji KN, Chen IC, Gildersleeve H, Lehle JD,
Mayo M, Westernströer B, Law NC, Oatley MJ, Velte EK, Niedenberger BA, Fritze D, Silber
S, Geyer CB, Oatley JM, McCarrey JR. 2018. The Mammalian Spermatogenesis Single-Cell
Transcriptome, from Spermatogonial Stem Cells to Spermatids. *Cell Rep* **25**:1650-1667.e8.
doi:10.1016/j.celrep.2018.10.026

Ho Y, Wigglesworth K, Eppig JJ, Schultz RM. 1995. Preimplantation development of mouse
embryos in KSOM: Augmentation by amino acids and analysis of gene expression. *Mol*
Reprod Dev **41**:232–238. doi:10.1002/mrd.1080410214

Kaneda Y, Miyata H, Shimada K, Oura S, Ikawa M. 2023. Testis-specific proteins, TSNAXIP1 and
1700010I14RIK, are important for sperm motility and male fertility in mice. *Andrology*
11:799–807. doi:10.1111/andr.13378

Kimura M, Imamoto N. 2014. Biological Significance of the Importin-β Family-Dependent
Nucleocytoplasmic Transport Pathways. *Traffic*. doi:10.1111/tra.12174

Kipreos ET, Pagano M. 2000. The F-box protein family. *Genome Biol* **1**:3002.1-3002.7.
doi:10.1186/gb-2000-1-5-reviews3002

Kissel H, Georgescu MM, Larisch S, Manova K, Hunnicutt GR, Steller H. 2005. The Sept4 septin
locus is required for sperm terminal differentiation in mice. *Dev Cell* **8**:353–364.
doi:10.1016/j.devcel.2005.01.021

Kluin PM, Kramer MF, de Rooij DG. 1982. Spermatogenesis in the immature mouse proceeds
faster than in the adult. *Int J Androl* **5**:282–294. doi:10.1111/j.1365-2605.1982.tb00257.x

Leblond CP, Clermont Y. 1952. Spermiogenesis of rat, mouse, hamster and guinea pig as
revealed by the "periodic acid-fuchsin sulfurous acid" technique. *Am Anat* **90**:167–215.
doi:10.1002/aja.1000900202

Li W, Bengtson MH, Ulbrich A, Matsuda A, Reddy VA, Orth A, Chanda SK, Batalov S, Joazeiro CAP.
2008. Genome-Wide and Functional Annotation of Human E3 Ubiquitin Ligases Identifies
MULAN, a Mitochondrial E3 that Regulates the Organelle's Dynamics and Signaling. *PLoS*
One **3**:e1487. doi:10.1371/journal.pone.0001487

Loveland KateL, Major AndrewT, Butler R, Young JuliaC, Jans DavidA, Miyamoto Y. 2015. Putting
things in place for fertilization: discovering roles for importin proteins in cell fate and
spermatogenesis. *Asian J Androl* **17**:537. doi:10.4103/1008-682X.154310

Loveland KL, Hogarth C, Szczepny A, Prabhu SM, Jans DA. 2006. Expression of nuclear transport
importins beta 1 and beta 3 is regulated during rodent spermatogenesis. *Biol Reprod*
74:67–74. doi:10.1095/biolreprod.105.042341

Lu Y, Nagamori I, Kobayashi H, Kojima-Kita K, Shirane K, Chang H, Nishimura T, Koyano T, Yu Z,
Castañeda JM, Matsuyama M, Kuramochi-Miyagawa S, Matzuk MM, Ikawa M. 2023.
ADAD2 functions in spermiogenesis and piRNA biogenesis in mice. *Andrology* **11**:698–709.
doi:10.1111/andr.13400

Mahboubi H, Seganathy E, Kong D, Stochaj U. 2013. Identification of Novel Stress Granule
Components That Are Involved in Nuclear Transport. *PLoS One* **8**:e68356.
doi:10.1371/journal.pone.0068356

Mani A, Gelmann EP. 2005. The ubiquitin-proteasome pathway and its role in cancer. *J Clin*
Oncol doi:10.1200/JCO.2005.05.081

Marcelo A, Koppenol R, de Almeida LP, Matos CA, Nóbrega C. 2021. Stress granules, RNA-
binding proteins and polyglutamine diseases: too much aggregation? *Cell Death Dis* **12**:592.
doi:10.1038/s41419-021-03873-8

Masek T, Vopalensky V, Suchomelova P, Pospisek M. 2005. Denaturing RNA electrophoresis in
TAE agarose gels. *Anal Biochem* **336**:46–50. doi:10.1016/j.ab.2004.09.010

Meikar O, Da Ros M, Korhonen H, Kotaja N. 2011. Chromatoid body and small RNAs in male
germ cells. *Reproduction*. doi:10.1530/REP-11-0057

Miyamoto Y, Baker MA, Whiley PA, Arjomand A, Ludeman J, Wong C, Jans DA, Loveland KL.
2013. Towards delineation of a developmental α-importome in the mammalian male
germline. *Biochim Biophys Acta Mol Cell Res* **1833**:731–742.
doi:10.1016/j.bbamcr.2012.11.005

Miyata H, Abbasi F, Visconti PE, Ikawa M. 2020. CRISPR/CAS9-mediated amino acid substitution
reveals phosphorylation residues of RSPH6A are not essential for male fertility in mice. *Biol*
Reprod **103**:912–914. doi:10.1093/biolre/ioaa161

Miyata H, Oura S, Morohoshi A, Shimada K, Mashiko D, Oyama Y, Kaneda Y, Matsumura T,
Abbasi F, Ikawa M. 2021. SPATA33 localizes calcineurin to the mitochondria and regulates
sperm motility in mice. *Proc Natl Acad Sci U S A* **118**:1–9. doi:10.1073/pnas.2106673118

Miyata H, Satouh Y, Mashiko D, Muto M, Nozawa K, Shiba K, Fujihara Y, Isotani A, Inaba K, Ikawa
M. 2015. Sperm calcineurin inhibition prevents mouse fertility with implications for male
contraceptive. *Science (1979)* **350**:442–445. doi:10.1126/science.aad0836

Morohoshi A, Miyata H, Oyama Y, Oura S, Noda T, Ikawa M. 2021. FAM71F1 binds to RAB2A and
RAB2B and is essential for acrosome formation and male fertility in mice. *Development*
148:1–12. doi:10.1242/dev.199644

Morohoshi A, Miyata H, Shimada K, Nozawa K, Matsumura T, Yanase R, Shiba K, Inaba K, Ikawa
M. 2020. Nexin-Dynein regulatory complex component DRC7 but not FBXL13 is required

for sperm flagellum formation and male fertility in mice. *PLoS Genet* **16**:1–21.
doi:10.1371/journal.pgen.1008585

Morohoshi A, Miyata H, Tokuhiro K, Iida-Norita R, Noda T, Fujihara Y, Ikawa M. 2023. Testis-
enriched ferlin, FER1L5, is required for Ca $^{2+}$ -activated acrosome reaction and male
fertility. *Sci Adv* **9**:eade7607. doi:10.1126/sciadv.ade7607

Muro Y, Hasuwa H, Isotani A, Miyata H, Yamagata K, Ikawa M, Yanagimachi R, Okabe M. 2016.
Behavior of mouse spermatozoa in the female reproductive tract from soon after mating
to the beginning of fertilization. *Biol Reprod* **94**:1–7. doi:10.1095/biolreprod.115.135368

Naito Y, Hino K, Bono H, Ui-Tei K. 2015. CRISPRdirect: Software for designing CRISPR/Cas guide
RNA with reduced off-target sites. *Bioinformatics* **31**:1120–1123.
doi:10.1093/bioinformatics/btu743

Nakayama K, Nagahama H, Minamishima YA, Matsumoto M, Nakamichi I, Kitagawa K, Shirane
M, Tsunematsu R, Tsukiyama T, Ishida N, Kitagawa M, Nakayama KI, Hatakeyama S. 2000.
Targeted disruption of Skp2 results in accumulation of cyclin E and p27 (Kip1), polyploidy
and centrosome overduplication. *EMBO Journal* **19**:2069–2081.
doi:10.1093/emboj/19.9.2069

Nsota Mbango JF, Coutton C, Arnoult C, Ray PF, Touré A. 2019. Genetic causes of male infertility:
Snapshot on morphological abnormalities of the sperm flagellum. *Basic Clin Androl* **29**:1–8.
doi:10.1186/s12610-019-0083-9

Pan J, Goodheart M, Chuma S, Nakatsuji N, Page DC, Wang PJ. 2005. RNF17, a component of the
mammalian germ cell nuage, is essential for spermiogenesis. *Development* **132**:4029–4039.
doi:10.1242/dev.02003

Protter DSW, Parker R. 2016. Principles and Properties of Stress Granules. *Trends Cell Biol*.
doi:10.1016/j.tcb.2016.05.004

Reineke LC, Neilson JR. 2019. Differences between acute and chronic stress granules, and how
these differences may impact function in human disease. *Biochem Pharmacol*.
doi:10.1016/j.bcp.2018.10.009

Richburg JH, Myers JL, Bratton SB. 2014. The role of E3 ligases in the ubiquitin-dependent
regulation of spermatogenesis. *Semin Cell Dev Biol*. doi:10.1016/j.semcdb.2014.03.001

Sato B, Kim J, Morohoshi K, Kang W, Miyado K, Tsuruta F, Kawano N, Chiba T. 2023. Proteasome-
Associated Proteins, PA200 and ECPAS, Are Essential for Murine Spermatogenesis.
Biomolecules **13**:586. doi:10.3390/biom13040586

Schulman BA, Carrano AC, Philip D. J, Zachary B, Elspeth R. E. K, Michael S. F, Stephen J. E, J.
Wade H, Michele P, & Nikola P. P. 2000. Insights into SCF ubiquitin ligases from the
structure of the Skp1-Skp2 complex. *Nature* **408**:381–386.
doi:https://doi.org/10.1038/35042620

Shang P, Baarends WM, Hoogerbrugge J, Ooms MP, Van Cappellen WA, De Jong AAW, Dohle GR,
Van Eenennaam H, Gossen JA, Grootegoed JA. 2010. Functional transformation of the
chromatoid body in mouse spermatids requires testis-specific serine/threonine kinases. *J
Cell Sci* **123**:331–339. doi:10.1242/jcs.059949

Shimada K, Kato H, Miyata H, Ikawa M. 2019. Glycerol kinase 2 is essential for proper
arrangement of crescent-like mitochondria to form the mitochondrial sheath during mouse
spermatogenesis. *J Reprod Dev* **65**:155–162. doi:10.1262/jrd.2018-136

Shimada K, Park S, Miyata H, Yu Z, Morohoshi A, Oura S, Matzuk MM, Ikawa M. 2021. ARMC12
regulates spatiotemporal mitochondrial dynamics during spermiogenesis and is required
for male fertility. *Proc Natl Acad Sci U S A* **118**:e2018355118.
doi:10.1073/pnas.2018355118

Shimada K, Park S, Oura S, Noda T, Morohoshi A, Matzuk MM, Ikawa M. 2023. TSKS localizes to
nuage in spermatids and regulates cytoplasmic elimination during spermiation. *Proc Natl
Acad Sci U S A* **120**:e2221762120. doi:10.1073/pnas.2221762120

Skaar JR, Pagan JK, Pagano M. 2013. Mechanisms and function of substrate recruitment by F-box
proteins. *Nat Rev Mol Cell Biol*. doi:10.1038/nrm3582

Snyder E, Chukrallah L, Seltzer K, Goodwin L, Braun RE. 2020. ADAD1 and ADAD2, testis-specific
adenosine deaminase domain-containing proteins, are required for male fertility. *Sci Rep*
10:11536. doi:10.1038/s41598-020-67834-5

Taylor JP, Brown RH, Cleveland DW. 2016. Decoding ALS: From genes to mechanism. *Nature*.
doi:10.1038/nature20413

Tiscornia G, Singer O, Verma IM. 2006. Production and purification of lentiviral vectors. *Nat
Protoc* **1**:241–245. doi:10.1038/nprot.2006.37

Tsunematsu R, Nakayama K, Oike Y, Nishiyama M, Ishida N, Hatakeyama S, Bessho Y, Kageyama
R, Suda T, Nakayama KI. 2004. Mouse Fbw7/Sel-10/Cdc4 Is Required for Notch Degradation
during Vascular Development. *J Biol Chem* **279**:9417–9423. doi:10.1074/jbc.M312337200

Yamaguchi R, Yamagata K, Ikawa M, Moss SB, Okabe M. 2006. Aberrant distribution of ADAM3
in sperm from both angiotensin-converting enzyme (Ace)- and calmegin (Clgn)-deficient
mice. *Biol Reprod* **75**:760–766. doi:10.1095/biolreprod.106.052977

Zheng Q, Huang T, Zhang L, Zhou Y, Luo H, Xu H, Wang X. 2016. Dysregulation of Ubiquitin-
Proteasome System in Neurodegenerative Diseases. *Front Aging Neurosci* **8**:303.
doi:10.3389/fnagi.2016.00303

Figures

Figure 1. *Fbxo24* **is predominantly expressed in testes and essential for male fertility.** (*A*)
RT-PCR of *Fbxo24* in mouse adult tissues. *Fbxo24* is predominantly expressed in the testis. Br:
brain, Th: thymus, Lu: lung, He: heart, Li: liver, Sp: spleen, Ki: kidney, Te: testis, Ut: uterus, Ov:
ovary, and N.C.: negative control (water). *Actb* was used as a control. (*B*) RT-PCR of *Fbxo24*
using RNAs obtained from mouse testes at various postnatal days. *Actb* was used as a control.
Water was used as a negative control (N.C.). (*C*) Construction of expression vectors for FBXO24
with (WT) or without (ΔF) the F-box domain. (*D*) *Fbxo24 (WT)-FLAG* or *Fbxo24 (ΔF)-FLAG* was
transiently expressed with *Skp1-1D4* in HEK293T cells. Immunoprecipitation (IP) was performed
using anti-1D4 antibody or anti-FLAG antibody. FBXO24-FLAG interacts with SKP1-1D4 via the
F-box domain. α-tubulin was used as a loading control. (*E*) Schematic for generating *Fbxo24* KO
mice using the CRISPR/Cas9 system. White boxes indicate untranslated regions while black
boxes indicate protein coding regions. The gRNAs used are shown. Fw and Rv indicate the
forward and reverse primer used for genotyping, respectively. (*F*) Genotyping of obtained *Fbxo24*
mutant mice. Fw #1-Rv #1 primers for KO allele and Fw #2-Rv #2 primers for WT allele in Fig. 1E
were used. N.C. indicates negative control (water). (*G*) Amplicons of the PCR product using Fw
#1-Rv #1 primers were subjected to direct sequencing and the 11,575 bp deletion was confirmed
in the KO allele. (H) The number of pups born per plug was counted to assess male fertility. Each
WT or KO male was mated with three WT females for three months.

Figure 2. FBXO24 is essential for sperm flagellar formation and motility. (*A*) The histology of
seminiferous tubules at different stages. An asterisk indicates remaining sperm heads. (*B*)
Morphology of mature spermatozoa obtained from cauda epididymis. White arrowhead indicates
straight spermatozoa. Red arrowhead indicates bent spermatozoa. Black asterisk indicates coiled
spermatozoa. Red asterisk indicates headless spermatozoa. (*C*) Stacked bar graph showing the
frequency of sperm morphology classified as straight, bent, coiled, and headless. *n* = 3
independent experiments. (*D*) Percentages of motile spermatozoa were analyzed at 10 min and
120 min after incubation in capacitation medium. (*E*) VAP (average path velocity) was analyzed at
10 min and 120 min after incubation in capacitation medium. (*F*) VSL (straight line velocity) was
analyzed at 10 min and 120 min after incubation in capacitation medium. (G) VCL (curvilinear
velocity) was analyzed at 10 min and 120 min after incubation in capacitation medium.

Figure 3. In vitro fertilizing ability and in vivo sperm migration. (*A*) The fertilizing ability of spermatozoa was analyzed in vitro using cumulus-intact oocytes. (*B*) The fertilizing ability of spermatozoa was analyzed in vitro using cumulus-free oocytes. (*C*) The fertilizing ability of spermatozoa was analyzed in vitro using zona-free oocytes. (*D*) Uterus and oviducts of WT females mated with WT or *Fbxo24* KO males carrying RBGS transgene. Female reproductive tracts were dissected 4 h after confirming a plug. Right figures are magnified images of the boxes indicated in the middle panels. (*E*) Immunoblotting of ADAM3 using testis and mature spermatozoa of *Fbxo24* heterozygous or KO mice. (*F*) ICSI experiment. The number of two-cell stage embryos and pups developed from WT oocytes injected with WT or *Fbxo24* KO spermatozoa. (*G*) Pups derived from WT or *Fbxo24* KO spermatozoa. (*H*) Genotyping of pups obtained from *Fbxo24* KO spermatozoa. N.C. indicates negative control (water).

789
790
791 **Figure 4. Numerous membraneless electron-dense granules were found in *Fbxo24* KO**
792 **spermatozoa.** (*A*) Epididymal spermatozoa of RBGS Tg mice were stained with Hoechst 33342
793 (nuclei). Mitochondria were labeled with su9-DsRed2. (*B*) Sperm mitochondrial sheath formation
794 during spermiogenesis was observed by scanning electron microscopy (SEM). (*C*) Cross sections
795 of spermatozoa in the cauda epididymis. Asterisks indicate electron-dense granules. (*D*)
796 Longitudinal sections of spermatozoa in the cauda epididymis. An asterisk indicates electron-
797 dense granules. (*E*) Percentages of morphologically abnormal mitochondria observed with
798 transmission electron microscopy (TEM). The number of flagellar sections analyzed is shown
799 above each bar. (*F*) Percentages of electron-dense granules observed in the midpiece cross
800 sections with TEM. The number of flagellar sections analyzed is shown above each bar.

801
802
Figure 5. IPO5 and KPNB1 accumulate in *Fbxo24* KO spermatozoa. (*A*) Mass spectrometry
analyses of mature spermatozoa. Significantly upregulated proteins are shown with red dots
whereas significantly downregulated proteins are shown with blue dots. (*B*) Immunoblotting
analysis was performed using proteins extracted from testes or mature spermatozoa. IPO5 was
detected using rabbit anti-IPO5 polyclonal antibody. KPNA2 and IZUMO1 were detected as
negative control and loading control, respectively. (*C*) Spermatozoa obtained from the cauda
epididymis were stained for IPO5 (magenta) using rabbit anti-IPO5 polyclonal antibody. Nuclei
were stained with Hoechst 33342 (blue). (*D*) Immunoprecipitation (IP) of FBXO24-FLAG from
Fbxo24-FLAG Tg testes. FBXO24 could interact with IPO5. IPO5 was detected using mouse anti-
IPO5 monoclonal antibody. β-actin was used as a loading control. (*E*) IP of FBXO24-FLAG using
HEK293T cells. FBXO24 could interact with IPO5 but not with KPNB1. IPO5 was detected using
mouse anti-IPO5 monoclonal antibody. β-actin was used as a loading control.

815
816
Figure 6. KPNB1 and IPO5 are recruited to SGs. (*A*) Total RNA amounts in spermatozoa were
measured by ultraviolet absorption. (*B*) Electrophoresis of RNA extracted from mature
spermatozoa. (*C*) KPNB1 and IPO5 were localized to stress granules (SGs) under exposure to
oxidative stress. COS7 cells were treated with water (upper row) or arsenite (lower row). Nuclei
were stained with Hoechst 33342 (blue). (*D*) KPNB1 and IPO5 were localized to SGs under
exposure to a proteasome inhibitor. COS7 cells were treated with DMSO (upper row) or MG132
(lower row). Nuclei were stained with Hoechst 33342 (blue).

Supplementary Information for

FBXO24 ensures male fertility by preventing abnormal accumulation
of membraneless granules in sperm flagella

Yuki Kaneda[1,2], Haruhiko Miyata[1,*], Zoulan Xu[1,2], Keisuke Shimada[1], Maki Kamoshita[1], Tatsuya
Nakagawa[1,2], Chihiro Emori[1], Masahito Ikawa[1,2,3,4,5,*]

[1]Research Institute for Microbial Diseases, Osaka University, 3-1 Yamadaoka, Suita, Osaka
5650871 JAPAN.
[2]Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka
5650871 JAPAN.
[3]The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo
1088639 JAPAN.
[4]Center for Infectious Disease Education and Research (CiDER), Osaka University, 2-8
Yamadaoka, Suita, Osaka 5650871 JAPAN.
[5]Center for Advanced Modalities and DDS (CAMaD), Osaka University, 2-8 Yamadaoka, Suita,
Osaka 5650871 JAPAN.
*Corresponding authors: Haruhiko Miyata and Masahito Ikawa
20

21 **Contact info**
22 **Email:** hmiya003@biken.osaka-u.ac.jp (HM); ikawa@biken.osaka-u.ac.jp (MI)

23
24
Figure S1. Characterization of mouse and human FBXO24. (*A*) Expression profile of mouse
Fbxo24 in spermatogenic cells is shown. Expression profiling was obtained from the testis single
cell RNA-seq data set (Hermann et al., 2018). Ud spg: Undifferentiated spermatogonia, A1- A2
Spg: A1 and A2 differentiating spermatogonia, A3-B Spg: A3, A4, In, and B differentiating
spermatogonia, Prele Sc: preleptotene spermatocytes, Le/Zy Sc: leptotene/zygotene
spermatocytes, Pa Sc: pachytene spermatocytes, Di/Se Sc: diplotene/secondary spermatocytes,
Early St: early round spermatids, Mid St: mid round spermatids, Late St: late round spermatids,
SC: Sertoli cells, PTM: peritubular myoid cells, LC: Leydig cells, and EC: Endothelial cells. (*B*)
Expression of human *FBXO24* in spermatogenic cells is shown. Expression profiling was
obtained from the testis single cell RNA-seq data set (Hermann et al., 2018). Dif Spg:
differentiating spermatogonia. (*C*) Pairwise sequence alignment of amino acids between mouse
and human FBXO24. Dark blue highlight indicates identical amino acids. (*D*) The F-box domain of
mouse and human FBXO24 was identified with a protein database search (SMART;
http://smart.embl-heidelberg.de/).

39
40
41 **Figure S2. Histological analyses of *Fbxo24* KO testis and epididymis.** (*A*) Gross morphology
42 of testes obtained from *Fbxo24* heterozygous and KO males. (*B*) Test weight (mg)/body weight
43 (mg) of *Fbxo24* heterozygous and KO mice. (*C*) PAS-hematoxylin staining of testes and cauda
44 epididymis sections. (*D*) Spermatozoa obtained from the cauda epididymis were stained for
45 IZUMO1 (magenta). Nuclei were stained with Hoechst 33342 (blue).

Figure S3. Lack of *Fbxo24* impairs sperm viability and the acrosome reaction. (*A*) Viability
of cauda epididymal spermatozoa was assessed with propidium iodide (PI) at 10 min and 120
min incubation in TYH medium. (*B*) The acrosome reaction rates of cauda epididymal
spermatozoa were assessed using RBGS Tg mice after 10 min and 120 min of incubation in TYH
medium. To induce the acrosome reaction, Ca^{2+} ionophore A23187 was added to the medium
after 120 min of incubation.

53
54
Figure S4. Sperm flagellum ultrastructure was disorganized in *Fbxo24* KO mice. (*A*)
Localization of the annulus in cauda epididymal spermatozoa. Anti-SEPT4 antibody was used to
visualize the annulus (green). Nuclei were detected with Hoechst 33342 (blue). The acrosome
was detected with anti-IZUMO1 antibody (magenta). (*B*) Transmission electron microscopy (TEM)
observation of the principal piece. A white arrowhead indicates a disruption of the axoneme and
outer dense fiber (ODF). (*C*) Percentages of morphologically abnormal fibrous sheaths (FS)
observed with TEM. The number of flagellar sections analyzed is shown above each bar. (*D*)
Percentages of morphologically abnormal axonemes (AX) observed with TEM. The number of
flagellar sections analyzed are shown above each bar. (*E*) Percentages of morphologically
abnormal ODF observed with TEM. The number of flagellar sections analyzed are shown above
each bar. (*F*) TEM observation of the principal piece. A red arrowhead indicates electron-dense
granules. (*G*) Percentages of electron-dense granules observed in principal piece cross sections.
The number of flagellar sections analyzed are shown above each bar.

68
69
Figure S5. No clear differences were found in the amount and localization of RNP granule-related proteins between control and *Fbxo24* KO testes.
(*A*) Immunodetection of ADAD1, ADAD2, MILI, MIWI, RNF-17, YTHDC2, TSKS and TSSK1 in
Fbxo24 heterozygous and KO testes. β-actin was used as loading control. (*B*)
Immunofluorescence observation of MIWI (green) in WT and *Fbxo24* KO testes. Acrosomes were
stained with PNA (red) and nuclei were stained with Hoechst 33342 (white). (*C*)
Immunofluorescence observation of TSKS (green) in WT and *Fbxo24* KO testes. Acrosomes
were stained with PNA (red) and nuclei were stained with Hoechst 33342 (white).

Figure S6. Generation and analyses of *Fbxo24-3×FLAG* Tg mice. (*A*) Schematic of the
Fbxo24-3×FLAG construct used to generate Tg mice. (*B*) Immunodetection of FBXO24-3×FLAG
in the testis from the Tg mice. β-actin was used as a loading control. (*C*) Morphology of cauda
epididymal spermatozoa. (*D*) Schematic for generating *Fbxo24* KO #2 mice using the
CRISPR/Cas9 system. White boxes indicate untranslated regions while black boxes indicate
protein coding regions. The gRNAs used are shown. Fw and Rv indicate the forward and reverse
primer used for genotyping, respectively. (*E*) Genotyping of obtained *Fbxo24* mutant mice. Fw #3-
Rv #3 primers in Fig. S6D were used. N.C. indicates negative control (water). *(F)* Predicted
FBXO24 amino acid sequences of KO #2 mutant mice. The 204 bp deletion resulted in E32D
mutation with a premature stop codon introduced 49 amino acids later. (G) Morphology of mature
spermatozoa obtained from cauda epididymis. (*H*) The list of identified proteins by IP-MS
analysis. The top 15 identified proteins in Tg are shown. The Quantitative Value (Normalized
Total Spectra) was calculated using Scaffold proteome software. IgG-related proteins, which is
likely from the antibody used for IP, were removed from the list.

93   **Table S1. Primers and gRNAs used in this study.**

primer	sequence (5'-3')
Fbxo24 Fw #1	tcctctggcaggtgaagcgc
Fbxo24 Fw #2	gatgtgttgattgtatgagacttcc
Fbxo24 Fw #3	actagcagcaagaagtccagg
Fbxo24 Rv #1	tctcgctcatgatagtcagg
Fbxo24 Rv #2	gaagtatggcaaagtcactgagtgc
Fbxo24 Rv #3	aagaatggcagctctcttcc
Fbxo24 Tg Fw	tgaagtgtgtgatgctgagg
Fbxo24 Tg Rv	gttcccgagtagccaaaaca
Fbxo24 Cloning Fw	atggatccgccgccatggtgaagcgcagctgcccttc
Fbxo24 Cloning Rv	gcgaattcgctctcgggagtcgaggtgtctgg
Actb Fw	catccgtaaagacctctatgccaac
Actb Rv	atggagccaccgatccaca

gRNA	target sequence (5'-3')
Fbxo24 gRNA #1	tgtggaggcgcatctgtcga
Fbxo24 gRNA #2	tcctgaaggaagtcgagccg
Fbxo24 gRNA #3	tcagttgttcccccagagc

97 **Table S2. Antibodies used in this study.**

antibodies	source	identifier
Rabbit anti-KPNA2	Abcam	ab84440
Rabbit anti-β-actin	MBL	PM053
Rabbit anti-human septin4	IBL	#18987
Rabbit anti-IPO5 (H-300)	SantaCruz	sc-11369
Rabbit anti-Miwi (G82)	Cell Sinaling Technology	#2079
Rabbit anti-ADAD1	(Lu et al., 2023)	
Rabbit anti-TSKS	(Shimada et al., 2023)	
Rabbit anti-TSSK1	(Shimada et al., 2023)	
Rat anti-ADAD2	(Lu et al., 2023)	
Rat anti-IZUMO1 monoclonal antibody	(Ikawa et al., 2011)	
Mouse anti-1D4 tag	A kind gift from Dr. Martin M. Matzuk	
Mouse anti-FLAG tag (M2)	Sigma-Aldrich	F1804
Mouse anti-α-tubulin (B-5-1-2)	Sigma-Aldrich	T5168
Mouse anti-acetylated tubulin	Sigma-Aldrich	T7451
Mouse anti-ADAM3 (F-4)	SantaCruz	sc-365288
Mouse anti-IPO5 (B-7)	SantaCruz	sc-55527
Mouse anti-β-actin (AC-15)	Abcam	ab6276
Mouse anti-KPNB1	Abcam	ab2811
Goat anti-rabbit IgG-HRP	Jackson ImmunoResearch	#111-036-045
Goat anti-rat IgG-HRP	Jackson ImmunoResearch	#112-035-167
Goat anti-mouse IgG-HRP	Jackson ImmunoResearch	#115-036-062
Goat anti-rabbit IgG-Alexa Fluor 488	Thermo Fisher Scientific	Cat# A11070
Goat anti-rabbit IgG-Alexa Fluor 546	Thermo Fisher Scientific	Cat# A11071
Goat anti-rat IgG-Alexa Fluor 488	Thermo Fisher Scientific	Cat# A11006
Goat anti-rat IgG-Alexa Fluor 546	Thermo Fisher Scientific	Cat# A11081
Goat anti-mouse IgG-Alexa Fluor 546	Thermo Fisher Scientific	Cat# A11017
Goat anti-mouse IgG-Alexa Fluor 546	Thermo Fisher Scientific	Cat# A11018

Legends for Movies S1 and S2

Movie S1 (separate file). Sperm motion of *Fbxo24* heterozygous mice was recorded after 120 min of incubation in TYH medium.

Movie S2 (separate file). Sperm motion of *Fbxo24* KO mice was recorded after 120 min of incubation in TYH medium.

SI References

Hermann BP, Cheng K, Singh A, Roa-De La Cruz L, Mutoji KN, Chen IC, Gildersleeve H, Lehle JD, Mayo M, Westernströer B, Law NC, Oatley MJ, Velte EK, Niedenberger BA, Fritze D, Silber S, Geyer CB, Oatley JM, McCarrey JR. 2018. The Mammalian Spermatogenesis Single-Cell Transcriptome, from Spermatogonial Stem Cells to Spermatids. *Cell Rep* **25**:1650-1667.e8. doi:10.1016/j.celrep.2018.10.026

Ikawa M, Tokuhiro K, Yamaguchi R, Benham AM, Tamura T, Wada I, Satouh Y, Inoue N, Okabe M. 2011. Calsperin is a testis-specific chaperone required for sperm fertility. *J Biol Chem* **286**:5639–5646. doi:10.1074/jbc.M110.140152

Lu Y, Nagamori I, Kobayashi H, Kojima-Kita K, Shirane K, Chang H, Nishimura T, Koyano T, Yu Z, Castañeda JM, Matsuyama M, Kuramochi-Miyagawa S, Matzuk MM, Ikawa M. 2023. ADAD2 functions in spermiogenesis and piRNA biogenesis in mice. *Andrology* **11**:698–709. doi:10.1111/andr.13400

Shimada K, Park S, Oura S, Noda T, Morohoshi A, Matzuk MM, Ikawa M. 2023. TSKS localizes to nuage in spermatids and regulates cytoplasmic elimination during spermiation. *Proc Natl Acad Sci U S A* **120**:e2221762120. doi:10.1073/pnas.2221762120