Experimental design impacts the vector competence of *Ornithodoros* ticks for African swine fever virus:

a meta-analysis of published evaluations

Jennifer Bernard[1,2], Sara Madeira[3], Joachim Otte[4], Fernando Boinas[3], Marie-Frédérique Le Potier[2], Laurence Vial[1], Hélène Jourdan-Pineau[1*].

[1] CIRAD, UMR ASTRE, F-34398 Montpellier, France.

ASTRE, Univ Montpellier, CIRAD, INRAE, Montpellier, France.

[2] Unité Virologie et Immunologie Porcines, Laboratoire de Ploufragan-Plouzané, Agence Nationale de Sécurité Sanitaire (Anses), Univ Bretagne Loire, BP53, 22440 Ploufragan, France

[3] Faculty of Veterinary Medicine, University of Lisbon, Avenida da Universidade Técnica, 1300-477 Lisbon, Portugal

CIISA—Center of Interdisciplinary Research in Animal Health, 1300-477 Lisbon, Portugal

Associate Laboratory for Animal and Veterinary Sciences (AL4AnimalS), 1300-477 Lisbon, Portugal

[4] Berkeley Economic Advising and Research. 1442A Walnut Street, Berkeley, CA, 94705 USA.

*Corresponding author,

Correspondence: helene.jourdan@cirad.fr

Abstract

African Swine Fever (ASF) is a highly economically devastating viral disease for swine. Soft ticks of the genus *Ornithodoros* are involved in its epidemiology, as vectors and natural reservoirs of African Swine Fever Virus (ASFV). The vector competence of *Ornithodoros* ticks for ASFV has been mainly studied by mimicking natural tick-to-pig transmission pathways through experimental infections in the laboratory. By reviewing the original research studies dating back to the 1960s on the vector competence of *Ornithodoros* for ASFV, we estimated the vector competence of 10 tick species in association with 38 viral strains resulting in 51 tick-virus associations. This assessment emphasized the extensive range of protocol designs employed and their impact on the success of tick infection with ASFV. Our results call for standardised procedures in vector competence experiments to facilitate further investigation and reduce potential experimental bias. In particular, we recommend the use of late nymphs or adult ticks from a laboratory colony to achieve efficient infection rates. In addition, viral inoculation should be carried out by blood meal rather than by injection, and preference should be given to high titre blood. Finally, detection of viral DNA should be performed 2 months after inoculation to distinguish between successful replication and residual virus in the tick.

Keywords: Ornithodoros soft ticks, African swine fever virus, vector competence, experimental methodology, meta-analysis

Introduction

African swine fever (ASF) ranks among the most damaging viral diseases affecting swine farming. The etiologic agent is a large DNA virus belonging to the *Asfaviridae* family (Dixon et al., 2013). African Swine Fever Virus (ASFV) causes haemorrhagic fever in susceptible infected swine, with mortality rates of up to 100% in naïve domestic pigs. In Africa, wild suids (warthogs, and possibly bushpigs and giant forest hogs) are natural vertebrate reservoirs that are resistant to the disease (Jori et al., 2013). In Europe, wild boars are susceptible to ASFV and exhibit clinical symptoms and mortality rates comparable to those of domestic pigs. In certain countries, soft ticks of the genus *Ornithodoros* act as vectors or natural reservoirs of ASFV. The virus can persist in ticks for several years, potentially leading to epizootic outbreaks in pig farms that have become colonized by ticks, even with the implementation of ASF control measures (i.e., pig slaughtering, disinfection, and quarantine) (Boinas et al., 2011; Jori et al., 2013). Once ASFV is introduced into a new free area where the presence of *Ornithodoros* ticks is highly suspected, as it occurred in the Caucasus in 2007 or in the Caribbean in 2021, the potential for long-term persistence of ASFV in tick reservoirs and ASFV transmission via tick bites raises concern (Beltrán-Alcrudo et al., 2008; EFSA et al., 2021). *Ornithodoros* soft ticks feed not only on wild suids but on a large variety of vertebrate hosts (aardvark, porcupine, chicken tortoise…). However, the presence of ASFV in these non-suid hosts has never been detected.

Vector competence generally refers to the vector's capacity to be infected by a pathogen, to maintain and multiply this pathogen, and finally to transmit it to a new susceptible host (Gillespie et al., 2004). The biological steps taking place within the arthropod vector can be broken down into four stages: (i) the ingestion of an infected blood meal, (ii) the infection of the vector upon crossing of its intestinal midgut by the pathogen, (iii) the persistence and multiplication of the pathogen within the vector, and (iv) the dissemination of the pathogen inside the vector by crossing organic barriers and ultimately

reaching organs implicated in the transmission to vertebrate hosts, typically the salivary glands and coxal glands. Sexual

transmission of pathogens between males and females of vectors as well as transovarial transmission from female vectors

to their progeny can occur through the infection of reproductive organs. Although not strictly included in the formal

definition of the vector competence, these pathways of transmission will also be considered in this review.

All these biological steps are observed in the interaction between ASFV and *Ornithodoros* tick although results reported

varied greatly between studies. In addition to historical demonstration of tick vector competence for ASFV in African *O.*

moubata sensu lato and Iberian *O. erraticus*, several other *Ornithodoros* tick species succeed to maintain or transmit ASFV

under experimental conditions (Mellor & Wilkinson, 1985; Endris et al., 1987, 1991; Hess et al., 1987). This suggests that

most *Ornithodoros* ticks might be able to maintain and transmit ASFV (EFSA Panel on Animal Health and Welfare., 2010a).

Based on available results, some researchers have proposed to predict the global role of each tick species as a vector and

reservoir for ASFV (Kleiboeker & Scoles, 2001; EFSA Panel on Animal Health and Welfare., 2010b; a, 2014; Burrage, 2013).

However, these comprehensive reports do not account for the evident disparities in vector competence results, which likely

rely on intricate tick-virus adaptations as proposed by Pereira de Oliveira (Pereira de Oliveira et al., 2019; Pereira De Oliveira

et al., 2020) and other yet-to-be-identified biological and experimental factors. No paper has considered the substantial

diversity of techniques employed to test vector competence except for a study on *O. erraticus* infected with two Portuguese

ASFV strains (Ribeiro et al., 2015). Nonetheless, the importance of experimental methodology has been demonstrated in

other models like mosquitoes with Rift Valley fever, Dengue, Zika or Japanese Encephalitis viruses (Azar, 2019; Auerswald

et al., 2021; Drouin et al., 2022). We hypothesize that the experimental design, including the choice of biological materials

and the methods used to infect ticks and detect the virus, is an important factor in explaining variation in vector

competence, in addition to the influence of tick species, ASFV strains and their combinations.

To investigate this hypothesis, a systematic review of studies published up to 2024 was conducted, examining the vector

competence of *Ornithodoros* ticks for ASFV. The present paper focuses on the diverse experimental factors that may impact

vector competence and determines the overall vector competence of the various associations between *Ornithodoros* tick

and ASFV. We provide recommendations on how experimental trials can be refined to minimize biases and obtain more

comparable and reliable results on *Ornithodoros* vector competence for ASFV.

Material and Methods

Collection of bibliographical resources

The first objective of the study was to identify the bibliographic resources related to vector competence of *Ornithodoros*

ticks for ASFV. For practical issues, we decided to consider only resources published in the English language.. This was

achieved following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Annex

1). Our bibliographic review was conducted until July 2022, using three different bibliographic databases: EThOS

(http://ethos.bl.uk/Home.do), PubMed (http://www.ncbi.nlm.nih.gov/) and Scopus (http://www.scopus.com/). Queries

were built under the keywords "African swine fever" AND "tick". The topics not related with tick competence (e.g.,

immunology, genetic, vaccinology, epidemiology) were excluded during the screening phase. Then, during the eligibility

phase, studies were excluded if quantitative reliable data could not be extracted from publications. Additionally, reviews

lacking original vector competence results and offering merely descriptive information were excluded. In addition to

scientific articles published in peer-reviewed journals, PhD manuscripts were also selected.. All duplicated results were

removed to avoid biases in the analyses. Finally, a review of the citations in the included references was carried out in order

to identify further relevant studies. The final list of included references that were used to extract all data in soft tick vector

competence for ASFV are shown in Table 1.

Table 1 - Included references providing original results on the vector competence of *Ornithodoros* ticks for ASFV, with number attributed to

each paper by year of publication.

N°	Included references
1	Parker J, Plowright W, Pierce MA. The epizootiology of African swine fever in Africa. Vet Rec. 1969;85(24):668-74
2	Plowright W, Perry CT, Peirce MA, Parker J. Experimental infection of the argasid tick, *Ornithodoros moubata porcinus*, with African swine fever virus. Arch Gesamte Virusforsch. 1970;31(1):33-50.
3	Plowright W, Perry CT, Peirce MA. Transovarial infection with African swine fever virus in the argasid tick, *Ornithodoros moubata porcinus,* Walton. Res Vet Sci. 1970 ;11(6):582-4.
4	Greig A. The localization of African swine fever virus in the tick *Ornithodoros moubata porcinus*. Arch Gesamte Virusforsch. 1972;39(1):240-7.
5	Plowright W, Perry CT, Greig A. Sexual transmission of African swine fever virus in the tick, *Ornithodoros moubata porcinus,* Walton. Res Vet Sci. 1974;17(1):106-13.
6	Groocock CM, Hess WR, Gladney WJ. Experimental transmission of African swine fever virus by *Ornithodoros coriaceus*, an argasid tick indigenous to the United States. Am J Vet Res. 1980;41(4):560-4.
7	Thomson GR. The epidemiology of African swine fever: the role of free-living hosts in Africa. Onderstepoort J Vet Res. 1985;52(3):201-9. Review.
8	Mellor PS, Wilkinson PJ. Experimental transmission of African swine fever virus by *Ornithodoros savignyi* (Audouin). Res Vet Sci. 1985;39(3):353-6.
9	Hess WR, Endris RG, Haslett TM, Monahan MJ, McCoy JP. Potential arthropod vectors of African swine fever virus in North America and the Caribbean basin. Vet Parasitol. 1987;26(1-2):145-55.
10	Endris RG, Haslett TM, Geering G, Hess WR, Monahan MJ. A hemolymph test for the detection of African swine fever virus in *Ornithodoros coriaceus* (Acari: Argasidae). J Med Entomol. 1987;24(2):192-7.
11	Hess WR, Endris RG, Lousa A, Caiado JM. Clearance of African swine fever virus from infected tick (Acari) colonies. J Med Entomol. 1989;26(4):314-7.
12	Haresnape JM, Wilkinson PJ. A study of African swine fever virus infected ticks (*Ornithodoros moubata*) collected from three villages in the ASF enzootic area of Malawi following an outbreak of the disease in domestic pigs. Epidemiol Infect. 1989;102(3):507-22.
13	Endris RG, Haslett TM, Hess WR. Experimental transmission of African swine fever virus by the tick *Ornithodoros* (Alectorobius) *puertoricensis* (Acari: Argasidae). J Med Entomol. 1991;28(6):854-8.
14	Endris RG, Hess WR, Caiado JM. African swine fever virus infection in the Iberian soft tick, *Ornithodoros* (Pavlovskyella) *marocanus* (Acari: Argasidae). J Med Entomol. 1992;29(5):874-8.
15	Endris RG, Hess WR. Experimental transmission of African swine fever virus by the soft tick *Ornithodoros* (Pavlovskyella) *marocanus* (Acari: Ixodoidea: Argasidae). J Med Entomol. 1992;29(4):652-6.
16	Endris RG, Haslett TM, Hess WR. African swine fever virus infection in the soft tick, *Ornithodoros* (Alectorobius) *puertoricensis* (Acari: Argasidae). J Med Entomol. 1992;29(6):990-4.

Endris RG, Hess WR. Attempted transovarial and venereal transmission of African swine fever virus by the Iberian soft tick *Ornithodoros* (Pavlovskyella) *marocanus* (Acari: Ixodoidea: Argasidae). J Med Entomol. 1994;31(3):373-81.

Boinas, FS. The role of *Ornithororos erraticus* in the epidemiology of ASFV in Portugal. PhD thesis manuscript. 1994

Rennie L. The Persistence of African swine fever virus in the Argasid Tick *Ornithodoros moubata*. PhD thesis manuscript. 1998.

Anderson EC, Hutchings GH, Mukarati N, Wilkinson PJ. African swine fever virus infection of the bushpig *(Potamochoerus porcus*) and its significance in the epidemiology of the disease. Vet Microbiol. 1998;62(1):1-15.

Kleiboeker SB, Burrage TG, Scoles GA, Fish D, Rock DL. African swine fever virus infection in the argasid host, *Ornithodoros porcinus porcinus*. J Virol. 1998;72(3):1711-24.

Kleiboeker SB, Scoles GA, Burrage TG, Sur J. African swine fever virus replication in the midgut epithelium is required for infection of Ornithodoros ticks. J Virol. 1999;73(10):8587-98.

Rennie L, Wilkinson PJ, Mellor PS. Effects of infection of the tick *Ornithodoros moubata* with African swine fever virus. Med Vet Entomol. 2000;14(4):355-60.

Rennie L, Wilkinson PJ, Mellor PS. Transovarial transmission of African swine fever virus in the argasid tick *Ornithodoros moubata*. Med Vet Entomol. 2001 ;15(2):140-6.

Burrage TG, Lu Z, Neilan JG, Rock DL, Zsak L. African swine fever virus multigene family 360 genes affect virus replication and generalization of infection in *Ornithodoros porcinus* ticks. J Virol. 2004;78(5):2445-53.

Nix RJ. Interaction of ASFV with its tick vector. PhD thesis manuscript. 2006.

Basto AP, Nix RJ, Boinas F, Mendes S, Silva MJ, Cartaxeiro C, Portugal RS, Leitão A, Dixon LK, Martins C. Kinetics of African swine fever virus infection in *Ornithodoros erraticus* ticks. J Gen Virol. 2006;87(7):1863-71.

Vial L, Wieland B, Jori F, Etter E, Dixon L, Roger F. African swine fever virus DNA in soft ticks, Senegal. Emerg Infect Dis. 2007;13(12):1928-31.

Rowlands RJ, Duarte MM, Boinas F, Hutchings G, Dixon LK. The CD2v protein enhances African swine fever virus replication in the tick vector, *Ornithodoros erraticus.* Virology. 2009;393(2):319-28.

Ravaomanana J, Michaud V, Jori F, Andriatsimahavandy A, Roger F, Albina E, Vial L. First detection of African Swine Fever Virus in *Ornithodoros porcinus* in Madagascar and new insights into tick distribution and taxonomy. Parasit Vectors. 2010;3:115.

Boinas FS, Wilson AJ, Hutchings GH, Martins C, Dixon LJ. The persistence of African swine fever virus in field-infected *Ornithodoros erraticus* during the ASF endemic period in Portugal. PLoS One. 2011;6(5):e20383.

Diaz AV, Netherton CL, Dixon LK, Wilson AJ. African swine fever virus strain Georgia 2007/1 in *Ornithodoros erraticus* ticks. Emerg Infect Dis. 2012;18(6):1026-8.

de Carvalho Ferreira HC, Tudela Zúquete S, Wijnveld M, Weesendorp E, Jongejan F, Stegeman A, Loeffen WL. No evidence of African swine fever virus replication in hard ticks. Ticks Tick Borne Dis. 2014;5(5):582-9.

Ribeiro R., Otte J., Madeira S., Hutchings G.H., Boinas F. Experimental Infection of *Ornithodoros erraticus sensu stricto* with Two Portuguese African Swine Fever Virus Strains. Study of Factors Involved in the Dynamics of Infection in Ticks. PLoS One. 2015;10(9):e0137718.

Pereira de Oliveira R, Hutet E, Paboeuf F, Duhayon M, Boinas F, Perez de Leon A, Filatov S, Vial L, Le Potier MF. Comparative vector competence of the Afrotropical soft tick Ornithodoros moubata and Palearctic species, *O. erraticus* and *O. verrucosus*, for African swine fever virus strains circulating in Eurasia. PLoS One. 2019;14(11):e0225657.

Pereira De Oliveira R, Hutet E, Duhayon M, Paboeuf F, Le Potier MF, Vial L. No experimental evidence of co-feeding transmission of african swine fever virus between Ornithodoros soft ticks. Pathogens 2020(a); 9(3), 168.

Pereira De Oliveira R, Hutet E, Duhayon M, Guionnet JM, Paboeuf F, Vial L, Le Potier MF. Successful infection of domestic pigs by ingestion of the European soft tick *O. erraticu*s that fed on African swine fever virus infected pig. Viruses, 2020(b); 12(3), 300.

Pereira De Oliveira R, Hutet E, Lancelot R, Paboeuf F, Duhayon M, Boinas F, Perez de Leon AA, Filatov S, Le Potier MF, Vial L. Differential vector competence of Ornithodoros soft ticks for African swine fever virus: What if it involves more than just crossing organic barriers in ticks? Parasites & vectors 2020©; 13(1), 1-15.

Forth JH, Forth LF, Lycett S, Bell-Sakyi L, Keil GM, Blome S, Calvignac-Spencer S, Wissgott A, Krause J, Höper D, Kampen H, Beer M. Identification of African swine fever virus-like elements in the soft tick genome provides insights into the virus' evolution. BMC biology, 2020; 18(1), 1-18.

Extraction of data about tick competence from bibliographical resources

General bibliometric information on authors and years of publication was collected to describe the general interest in

tick vector competence for ASFV and to identify the community of scientific experts. We considered the experts as the

persons in first and last position of the included references.

Information on tested *Ornithodoros* tick-ASFV associations was systematically reported. Regarding virus, we recorded

the strain, the genotype (on the basis of p72 gene sequences, Qu *et al*. 2022), the geographical origin, and the host (suids

or ticks) from which the ASF virus was isolated. The original names of some ASFV strains were retrieved by crossing

information from the literature, according to known authors' collaborations, or by directly contacting the authors. When

the determination of strains remained impossible, the strain was identified by indicating the country where the ASFV strain

was isolated, the year of the study, and a letter if duplicates existed. One exception was a strain used to infect *O. turicata*

(Hess et al., 1987) without any indication of the country of origin which was named "Unknown". Six recombinant ASFV

strains, genetically modified and obtained from two natural strains (Burrage et al., 2004; Nix, 2006), were excluded from

this review since they cannot infect ticks in nature. Genotypes were not available for some strains of ASFV, but they could

be inferred based on their geographical origin and ASFV genotype distribution published by Lubisi et al. (2005).

Concerning ticks, the species, their geographical origin, their colony origin and type were reported. When ticks were

reared for less than five years in the laboratory, they were considered comparable to ticks sampled from the field due to

the limited number of generations occurring during this period. Soft tick systematics has greatly changed during the last

decades. Tick species identification was updated in accordance with the revision of valid tick names by Guglielmone

(Guglielmone et al., 2010) and the mapping of their known distribution (Vial, 2009). The more recent taxonomy study of

southern African *Ornithodoros* ticks (Bakkes et al., 2018) could not be used, due insufficiently precise location to attribute

a species name among the *O. moubata* complex of species. Finally, when determining tick geographical origin, 6 studies

involving 4 different tick species (*O. coriaceaus, O. parkeri, O. moubata*, and *O. porcinus*) did not report this information

but since *O. parkeri* and *O. coriaceus* are known to be limited to North America, especially the United-States (De La Fuente

et al., 2008), they were assigned to this country.

Authors used various measurements to assess the tick vector competence for ASFV. Some authors reported numbers

of ticks positively detected with ASFV, while others scrutinized the virus titres in ticks and tick organs or monitored ASFV

kinetics for several weeks or months, providing more comprehensive information on the virus persistence within its tick

reservoir. Even if ASFV detection in tick transmission organs (salivary or coxal glands, gonads) suggest possible successful

transmission to pigs or to other ticks, this alone is insufficient to confirm such transmission pathways. Conversely, the

detection of ASFV in transmission secretions (e.g., saliva or coxal fluids) is stronger evidence, as the virus is already excreted

in this case. In these transmission experiments, clinical signs observed in pigs following tick bites and detection of ASFV in

pig blood provide definitive confirmation of the ability of ticks to transmit the disease. Similarly, detection of ASFV in tick

sexual mates, offspring confirms sexual and vertical transmission. The transmission of virus between ticks via co-feeding

has been described in hard ticks, but this phenomenon has not been demonstrated for ASFV in soft ticks (Pereira de Oliveira

et al., 2020).

Based on these data, it was possible to evaluate three major components of the vector competence of the ticks: 1) the

tick infection (revealed by the crossing of the tick midgut by ASFV), 2) the ability of ticks to transmit ASFV to pigs and 3)

their ability to transmit ASFV to other ticks). Transmission to pig through tick bite or from ticks to ticks cannot occur without

the previous infection of the tick, which was thus considered as the main step to be assessed. In addition, tick ability to

transmit ASFV to pigs or other ticks are less frequently reported (N= 160 and N=54 over 900 experiments, respectively)

likely because they are somewhat more difficult to assess than tick infection. Therefore, for each experiment, we computed

a binary variable, the infection status, according to any measurements providing direct information on tick infection as well

as indirect transmission measurements that imply previous tick infection (Figure 1). More precisely, if ASFV transmission to

pigs or ASFV transmission from ticks to other ticks were reported in papers, the ticks were considered to be infected as the

virus has necessarily crossed the tick organic barriers to reach transmission organs and be transmitted.   If

transmission failed or was not tested, the detection of ASFV in ticks was used as an indication of infection status. Obviously

when ASFV was not detected, ticks were considered as not infected. The only exception was when the detection occurred

on the same day as virus exposure, in which case no virus detection indicates a failure of experimental infection or a too

low detection threshold.

*Virus detection in the transmission organs is not a proof of transmission but the detection in the transmission fluids is a proof of transmission.
** we suspect in this case a failure in the experiment.

Figure 1 - Criteria used to determine the infection status of ticks after their virus exposure, with three classes (IS: Infection Status; DPE: Date

post exposure; NA: Not Applicable, 1: Infected, and 0: Not Infected).

Different methods have been used by authors to detect ASFV in ticks, namely ASFV isolation/titration, observation of

ASFV particles, and viral DNA or RNA detection using multiple designs of PCR. In the included references, ASFV

isolation/titration was based on haemadsorption (HAD), which is the ability of many ASFV strains to aggregate on red blood

cells when they infect pig alveolar macrophages and create typical "rosettes" (Malmquist & Hay, 1960). . This technique is

largely used by authors but may suffer from inhibition issues due to cytotoxic effect of tick components on pig macrophages

and its achievement in at least 6 days is very time-consuming (Oura et al., 2013). Virus titration implies that the virus is able

to replicate but does not necessarily confirm that it has crossed the midgut barrier. Similarly, detection of viral RNA by

Reverse Transcription quantitative PCR also indicates active viral replication. Conversely the detection of viral DNA can be

due to residual virus from blood meal and does not mean that the virus is still active. However, we considered that those

methods brought equivalent results to assess infection status once the duration after exposure was taken into account.

Indeed, when ASFV was detected in ticks, a minimum duration of 61 days between tick exposure to ASFV and detection

of ASFV in tick (hereafter DPE for days post exposure) was chosen to distinguish successful viral replication inside the tick

from detection of residual virus after the infectious blood meal or virus inoculation (Figure 1). Indeed, the minimum

duration for blood meal digestion and clearance of ingested blood in soft ticks is approximately 4 weeks (Rowlands et al.,

2009). Furthermore, if tick infection is unsuccessful, ASFV can be detected during 4 weeks by isolation (Ribeiro *et al.*, 2015)

and after at least 49 days by qPCR (de Carvalho Ferreira et al., 2014; see also Bernard, 2015 for a review on ASFV kinetics

inside soft ticks). If tick infection is successful, virus generalization in ticks takes about 30 to 56 days (Plowright et al., 1970;

Kleiboeker et al., 1998; Basto et al., 2006; Burrage, 2013). Therefore, and as a precaution, if ASFV was tested and detected

before 61 DPE, the infection status was considered unknown unless virus was detected in organs other than the midgut. In

this case, this indicates that the virus has crossed the midgut barrier.

Inoculation by injection allows accurate assessment of the virus dose received by the tick. As it delivers the virus directly

into the haemocoel, circumventing the midgut barrier, it offers valuable insights into the molecular compatibility between

the virus and the tick. However, it is acknowledged that the injection can be traumatic for the tick, resulting high mortality

and leads to distribution of the virus throughout the tick (Bonnet & Liu, 2012). In this case, as the detection of virus in the

organs was doubtful, the infection status was recorded as unknown.

The experimental design used in the different studies was also reported, including: i) the exposure route for the tick

infection, ii) the ASFV titres in the inoculum used to infect the ticks (infectious dose), iii) the development stage of the ticks

tested in the study, iv) the DPE, v) the number of ticks tested in the study. In field studies, viral titres in the hosts were not

necessarily reported but they were considered to be high ($>10^{6}$ HAD_{50}/mL of blood) considering the viraemia usually

observed in ASF diseased pigs (Plowright et al., 1968). Tick stage was classified as "early" from larva to nymph 3 and as

"late" from nymph 4 to adult.

Statistical analysis of the tick infection status

A statistical analysis was carried out to identify factors potentially affecting the tick infection status. More precisely, we

investigated if factors related to the nature of tick-virus association or the experimental design influenced the infectious

status (see above for details) of the ticks in published experimental trials. The explanatory variables were: the ASFV strain

and genotype, the tick species, the potential geographical co-occurrence of tick and virus as an indication of possible tick-

virus adaptation (ASFV and ticks from the same country versus from different countries), the viral titre in infectious blood

meal and the exposure route for tick infection, the tick stage, the tick colony type and the host from which the ASFV strain

was sampled. Using a dataset where all ticks were adults, we also tested the influence of tick sex on infectious status (along

with all other factors except tick stage). All statistical analyses were performed on R 4.2.2 (R Core Team, 2022). To assess

which factors influenced the infection status, we applied a generalized mixed model with a binomial family. The tick species,

virus genotype and virus strain were set as random effects to account for the fact that only a random fraction of soft tick

species and virus genotypes and strains were sampled. All other factors were set as fixed effects. The significance of random

effects was tested using a likelihood ratio test (Zuur et al., 2009). We then selected the best parsimonious models (with the

remaining 6 fixed effects) based on the AIC (Akaïke Information Criterion). In this best model, the significance of fixed effects

was tested using Chi² tests between models and post-hoc tests, with Tukey HSD correction, were performed.

Assessment of the overall tick vector competence for ASFV

Although our main assumption is that vector competence results are influenced by the experimental design, we

attempted to assess the overall vector competence of the various *Ornithodoros* tick-ASFV associations. Based on the three

main components described above, we scored each tick-ASFV association, according to the following ranking: "score 0" for

ticks that were unable to become infected or for which it was impossible to conclude (IS = 0 or IS = NA, figure 1) ; "score 1"

when ASFV was able to cross tick midgut and replicate in ticks; "score 2" when transmission from tick to pig was successful

; and, "score 3" for ticks validating all components of vector competence, including ASFV transmission from ticks to other

ticks. The attribution of a score was based on all data available from the different included references related to the

considered tick-ASFV association. Attributing graduate scores of vector competence sounds as if that associated

transmission events are also graduate. This is not the case: even if transmission to host scored 2 and sexual or transovarial

transmissions scored 3, the success of tick-to-tick transmission does not necessarily imply the efficiency of transmission to

host..

<center>**Results**</center>

Bibliometric description

The initial bibliographic search generated 236 references using Scopus and 138 by Pubmed, with 2 articles detected only

by Pubmed. The ETHOS database of thesis manuscripts and our manual bibliography search allowed identifying 8 more

resources. In this pool of bibliographic references, 39 references presenting vector competence original results were

included in the final dataset. They consisted of 5 field studies monitoring natural tick infections, 34 articles of laboratory

experiments on ticks and 2 mixed papers (Table 1). As shown in figure 2, the number of publications pertaining to the vector

competence of *Ornithodoros* ticks for ASFV was consistently low over the years, while the total number of references

related to ticks and ASFV progressively increased from the 1980s to 2022. The community of scientific experts identified is

rather limited with only thirty-nine authors listed as first and last authors of the selected publications. After 2015, 9 out of

39 experts continued to publish in this domain and can be considered the current core of experts on the vector competence

of ticks for ASFV.

Figure 2 - Number of bibliographic references published each year, from 1966 to 2022, on "African swine fever" AND "tick" (black dots = 246

screening references) and on the vector competence of *Ornithodoros* ticks for ASFV (red dots = 39 included references). The names of first and

last authors of the 39 included references were reported in blue boxes at the top of the graphic. The length of boxes indicates their respective

period of expertise, during which they published in the "African swine fever" AND "tick" fields (i.e., among the 246 screening references). List

of the abbreviation author name: AEC: Anderson E.C.; BAP: Basto A.P.; dCF: de Carvalho Ferreira H.C.; GWJ: Gladney W.J.; GCM: Groocock C.M.;

LWL: Loeffen W.L.; McJ: McCoy J.P.; MMJ: Monahan M.J.; RJ: Ravaomanana J.; TGR: Thomson G.R.

Data related to the biological material: ticks and ASFV

A database was created to record all the extracted data. From the 39 included references, 2036 records corresponding

to 900 unique experiments were extracted. For some references, authors detailed the results of their experiment for each

tick or with multiple records over time for the same ticks. Each experiment was characterized by a combination of tick

species, virus strain and experimental conditions (see Annexes 2 and 3). In total, the tick species and ASFV strains

represented 51 tick-virus associations and more than half (34/51) appeared in only one publication.

For the ticks, ten species were tested for their vector competence for ASFV. The Iberian tick *O. erraticus* was the most

studied (representing 44% of the reported experiments), followed by the *O. moubata* complex of species, including *O.*

porcinus and *O. moubata* sensu stricto (43%). The North American *O. coriaceus* (6.4%) were third, then they were far

followed by the other species *O. puertoricensis, O. verrucosus, O. savignyi, O. parkeri, O. sonrai, O. turicata* (3.7% - 1.0% -

0.8% - 0.4% - 0.4% - 0.2 % respectively). The geographical origins of ticks specified in the included references are presented

in figure 3. All *O. erraticus* ticks tested came from Portugal, and most of them from the same region of Alentejo, while *O.*

moubata and *O. porcinus* ticks came from diverse countries, which is representative of their large distribution range in East

and Southern Africa.

A total of 38 different ASFV strains were tested. Most ASFV strains used in the studies were isolated from ticks and pigs

(43.1% and 40.4% of the reported experiments, respectively). Only four strains studied in 16.2% of the experiments were

isolated from warthogs and 0.2% of virus strains are from undetermined hosts. ASFV strains were geographically distributed

over 17 different countries, mostly in Africa, and belonged to 8 genotypes over the 23 genotypes described so far, with a

large representation of the genotype I (64.9%, figure 3). The genotype of the 114a and Transvaal CV strains from South

Africa remains unknown.

Figure 3 - Countries of origin of *Ornithodoros* tick species (coloured according to the species) and ASFV genotypes (indicated by genotype

number associated with each country), involved in tick-ASFV associations studied in included references. The origin of the tick *O. parkeri* was

not indicated but should be the United-States, as described in the literature.

Data related to the experimental design

Of the 900 experiments recorded, half used ticks directly sampled from the field (especially *O. erraticus* and *O. porcinus*)

or reared in insectary for less than 5 years (48.7%). The other experiments used ticks reared in the laboratory for more than

5 years with some reared until 25 years for one study on *O. turicata* (47.2%). It was not possible to find information for the

remaining experiments. The majority of experiments (57.9%) involved ticks in their later developmental stages, ranging

from nymph 4 to adults. Early stages, from larva to nymph 3, were employed in 23.3% of the experiments. In the remaining

experiments, authors either utilized a combination of early and late stages (14.7%) or did not provide information about

the specific developmental stage used (4.1%). The number of ticks tested for each experiment was fairly low especially in

laboratory experiments (median=7 ticks tested, mean=30.03, range=[1;1141]).

Four exposure routes were used to infect ticks. Natural blood feeding on hosts was reported either in field studies, or

in some laboratory experiments using experimentally infected pigs. This method represented almost half of the total

experiments conducted (47.4%). Two artificial feeding methods are reported: the longest established method uses capillary

tubes inserted on tick mouthparts (8.8%) while a more recent approach uses animal skin, parafilm, or silicone membranes

(35.7%). Another artificial exposure route was the syringe inoculation of ASFV directly in the tick haemocoel (8.1%).

ASFV infectious dose used to infect ticks was very heterogeneous and classified in three categories. Most experiments

(78,4%) used medium to high viral doses (between 10^4 and 10^6 or over 10^6 HAD$_{50}$/mL of blood). The DPE was very variable

ranging from 0 day to up to 8 years. A DPE of 0 day is presumably used as a control for successful viral exposure. The longest

reported DPE are from Boinas *et al.* (2011) and represent the duration between an ASFV outbreak and the date of virus

detection in ticks in Portuguese farms, with virus detected in ticks up to 5 years after the outbreak. Nonetheless, more

than half experiments (54.3%) reported a DPE lower than the duration required to differentiate a successful viral replication

inside the tick from a detection of residual virus after the infection (61 days). Among them, one third performed virus

detection in organs or transmission experiments, allowing conclusion on tick infection status.

Factors influencing the likelihood of tick infection

On the initially 900 recorded experiments, we obtained a dataset of 407 observations with full information regarding

tick infection status and experimental conditions (Annexes 4 and 5). Among the random effects tested, the tick species and

virus genotype had no effect on the infection status but the viral strain had (χ^2_1=4.879 p-value= 0.027, see Annex 6 for

details on viral strains). Model selection based on AIC ended up with three variables displaying large and comparable

importance across all models: virus titre, tick stage and tick colony type (Figure 4 left). All those three variables had

significant effects on infectious status (Figure 4 right).

Figure 4 - *Left*: Relative importance of explanatory variables across all models explaining infection status. The importance value for a particular variable is equal to the sum of the weights for the models in which the variable appears. "Virus host" is the host from which ASFV was primarily isolated. "Expos. route" is the exposure route for tick infection (natural, artificial, capillary, injection). *Right*: Effect of tick stage, tick colony and virus titre on the infection status (all three effects are significant). Mean of infection status (i.e., proportion of experiments where ticks were infected). Values above the bars are the number of experiments in each level. Virus titre in HAD_{50}/mL of blood.

In more details, a higher viral titre corresponded to a higher likelihood of tick infection (χ^2_2= 38.493, p-value=4.423×10^{-9}). Bigger nymphs and adults (late stages) were more prone to get infected than small nymphs (χ^2_1=17.880, p-value=2.353×10^{-5}). And ticks from laboratory colonies had higher probability of infection than ticks recently sampled in the field (χ^2_1= 14.031, p-value=1.798×10^{-4}). The potential geographical co-occurrence of tick and virus, the exposure route for tick infection and the host from which ASFV was isolated were not significantly linked with infection status even if they were present in some of the best models (Annex 7). Using the dataset containing only experiments with adult ticks showed that sex had no influence on infection status.

Figure 5 summarizes the vector competence score for each of the 51 associations between soft ticks and ASFV virus tested so far. In more detail, 23.5% of the tick-ASFV associations presented a vector competence score null (score 0) i.e., ticks were unable to become infected or it was impossible to conclude on their infection status. For 27.5% of tick-ASFV associations, only tick infection (score 1) was achieved. Transmission from ticks to pigs was tested in 64.7% of tick-ASFV associations and was successful in 29.4% (score 2). Finally, 33.3% of tick-ASFV associations were assessed for ASFV transmission from ticks to ticks and 19.6% succeeded to transmit (score 3).

Figure 5 - ASFV vector competence scores for the different tick-ASFV associations studied in the included references (score 0: no tick infection or no information to conclude, score 1: successful tick infection, score 2: successful ASFV transmission to pigs, score 3: successful ASFV transmission to other ticks). In each histogram bar, the numbers are related to the included references recorded in Table 1. For each included reference, an individual score is given according to available results in this paper. The overall vector competence score for a given tick-virus association is the best vector competence result reported from all included references related to this tick-ASFV association.

Only 27.5 % of tick-ASFV associations were fully tested for vector competence (including tick infection, transmission to

pig, and transmission to other ticks) and 15.7% validated scores 1 to 3. Conversely, 3 associations reached score 3, without

being tested for pig transmission (O. *erraticus*-Zimbabwe/83/a and O. *erraticus*-Tomar/87) or by failing to transmit to pigs

when tested (O. *erraticus*-Ourt/88/1).

At the species level, most tick species tested were confirmed to be competent vectors (score 2) for at least one ASFV

strain. This competence remains unclear for 3 species that were either tested with only one ASFV strain and failed to

transmit the virus to pig and/or to ticks (O. *parkeri* and O. *verrucosus*) or never tested for transmission *(O. sonrai)*. Regarding

ASFV transmission to other ticks, 6 tick species were tested and 4 (O. *moubata*, O. *porcinus, O. *puertoricensis*. and *O.

erraticus) succeeded for at least one ASFV strain whereas 2 (O. *coriaceus* and O. *verrucosus)* failed. Finally, high

heterogeneity of vector competence scores was observed within a same tick species depending on ASFV strains used (e.g.

score 3 for O. *moubata*-LIV13/33 and score 0 for O. *moubata*-Brazil/78), and sometimes for a same tick-ASFV association

(e.g. scores 0/1 and 2 for two O. *turicata* colonies of different ages with the same ASFV strain).

Discussion

Bibliographic resources

This paper is the first exhaustive review of all the original studies published on the vector competence of soft ticks for

ASFV until 2024. Despite an exhaustive screening of bibliographic resources, only 39 references were considered and the

expert panel had always been limited, as exemplified by only 9 publishing experts since 2015. This relative paucity of

bibliographic resources regarding soft tick competence for ASFV may be attributed to the difficulty of investigating their

vector competence, both in laboratory and in field conditions. Indeed, sampling endophilous *Ornithodoros* ticks and

developing relevant methods to detect ASFV in natural tick populations (Jori et al., 2013; Oura et al., 2013) or rearing soft

ticks and amplifying ASFV in laboratory conditions are long and laborious tasks (Vial, 2009; Carrascosa et al., 2011). In

addition, although ASF is of high veterinary and economic concern, vectorial transmission is seen as occasional with a

relatively minor role in disease spread at least in some regions (Guinat et al., 2016). Nevertheless, soft ticks play a key role

in the maintenance of the virus and this is why characterizing competence for tick-virus associations is required to assess

the risk of ASFV introduction and spillover, as Golnar *et al.* (2019) did for the US.

Vector competence and tick-virus co-adaptation

Although all references included in this study aimed to evaluate the vector competence of soft ticks for ASFV, we noticed

a considerable diversity of vector competence measurements that yield distinct information. As a consequence, vector

competence was only partially assessed for most tick-ASFV associations, with a noticeable decrease of studies along the

successive biological steps of vector competence. In addition, more than half of tick-ASFV associations were characterized

by only one single article. Heterogeneous vector competence scores were observed within species between viral strains.

This could be interpreted either as a lack of sufficient reliable studies or high tick-virus specificity.

Adaptation between arthropod vectors and viruses is based on genetic features defining susceptibility in both organisms

and the selection of susceptibility may be mediated partly by detrimental effects of arboviruses on vector physiology

(Gillespie et al., 2004). In the soft tick-ASFV model, the presence of ASFV MGF360 genes, the lectin-like EP153R gene, or

the CD2-like EP402R gene enhance the entry and the replication of ASFV into ticks (Burrage et al., 2004; Rowlands et al.,

2009). Some ASFV strains naturally lost these genes through prolonged circulation within the population of Iberian domestic

pigs, or as a consequence of repeated passages on pig macrophages for vaccine production (Portugal et al., 2015). MGF 360

has also be shown to suppress the immune response of vertebrate and arthropod hosts against ASFV (Portugal et al., 2015).

Arthropods, meanwhile, are known to display RNA interference as an important immune response against virus infection

(Vijayendran et al., 2013). The involvement of such molecular pathway has recently been suggested in soft ticks, with ASFV-

like integrated elements coding for small interfering RNA in *O. moubata* (Forth et al., 2020). More specifically, it appears

that a greater number of these small RNA matched with ASFV genomes from genotype II than with genotype I, potentially

contributing to the variation in vector competence between viral strains. This is in agreement with our results which showed

significant variation in tick infection status depending on viral strain. However, no effect of virus genotype on tick infection

status was found, which may be due to the very imbalanced dataset regarding virus genotype with genotype I being

overrepresented compared to other genotypes.

We can suppose that field ticks, co-evolving or not with ASFV, may develop a much more heterogeneous and adaptive

response against ASFV infection and show a lower success of infection, as it was already shown in some mosquito-virus

models (Gillespie et al., 2004). Indeed, some authors previously reported very low ASFV infection rates in field ticks

(Plowright et al., 1969; Haresnape & Wilkinson, 1989; Boinas et al., 2011). In the present study, laboratory tick colonies

reared for more than five years were shown to be significantly much more infected by ASFV than field ticks or recently

established laboratory ticks. In addition, ticks from old laboratory colonies may be more prone to efficient artificial blood

feeding, thereby increasing the blood volume and the intake of ASF virus. However, the drawback of using old laboratory

colonies is the risk of genetic drift and laboratory adaptation, leading to divergent phenotypes, compared to wild

populations.

Despite heterogeneous vector competence according to studies and viral strain tested, experimental evidence shows

that 7 *Ornithodoros* species from Africa (*O. moubata, O. porcinus and O. savignyi*), Europe (*O. erraticus*) and North America

(*O. coriaceus, O. turicata and O. puertoricensis*) can be infected by at least one strain of ASFV and then transmit it to pigs.

Effect of experimental factors on tick competence

Our analysis of vector competence references demonstrated the importance of the experimental design on vector

competence results. The ASFV titre in the inoculum for infecting ticks appears as a key parameter in the vectorial

competence experiments, with likelihood of tick infection above 50% when ASFV load was higher than 10^6 HAD$_{50}$/mL.

Accordingly, a threshold of $10^{5.75}$ HAD$_{50}$/mL was previously determined by Ribeiro (Ribeiro et al., 2015) for the specific

ability of *O. erraticus* to maintain and multiply two Portuguese ASFV strains. Forth *et al.* (2020) also demonstrated an impact

of ASFV titre on tick infection probability and load. Soon after ingestion, ASFV has to cross the tick midgut barrier to be

amplified in replication units localized in tick endothelial cells and then enter the tick hemolymph (Burrage, 2013). This

suggests that the tick midgut and other tick membranes act as porous surfaces; the probability of ASFV entry would thus

depend on the number of available pores and on the virus load (Franz et al., 2015). Tick immune system such as RNAi could

also block low level of ASFV infection (Forth et al., 2020).

The development stage of ticks had a significant impact on tick infection status, with better results for late stages (from

nymph 4 to adults). This pattern confirms previous results on *O. erraticus*-Tomar and *O. erraticus*-OURT88/1 from Ribeiro

(Ribeiro et al., 2015). This was explained by higher blood volume intake for late and large stages, and therefore highest

ASFV titre inside the tick after blood meal digestion. In addition, even if female take a larger blood meal than males, we

found no effect of observed sexual dimorphism in adults on the likelihood of tick infection which is in agreement with

Ribeiro *et al.* (2015). In this study, the authors suggest that, in adults, the number of ASFV particles already reached a certain

threshold necessary to cause tick infection.

The route for tick infection was expected to play a role on tick competence. Natural feeding on viraemic pigs may provide

higher viral titre (usually $>10^6$ HAD$_{50}$/mL) than artificial feeding on membrane or with capillary (from 10^4-10^6 HAD$_{50}$/mL)

due to the difficulty for amplifying ASFV on pig macrophages. Despite differences between inoculation methods, this does

not seem to translate into any significant differences on the success rate of infection in our dataset. This could be attributed

to our dataset being incomplete and imbalanced, and with the virus titre being a confounding factor. Similarly, we did not

find any effect of the initial host in which viral strain was isolated on tick competence.

Conclusion

The examination of 39 references on the vector competence of *Ornithodoros* soft ticks for ASFV highlighted factors

influencing vector competence results. For the same tick-ASFV association, vector competence results vary according to

the experimental design employed. The selection of an experimental protocol constitutes a trade-off between scientific

considerations and practical constraints. The findings of this meta-analysis provide a foundation upon which

recommendations can be formulated for optimal protocols to ensure reliable results on vector competence for any soft

tick-ASFV association. First, using late nymphs or adult ticks from a laboratory tick colony will improve the probability of

obtaining engorged ticks, increase the volume of blood intake and allow testing for sexual or vertical transmission. Exposure

should be conducted using a blood meal rather than by injection into the haemocoel. As the viral titre exerts the most

significant influence on tick infection success, preferring blood from viraemic pigs or blood with a high titre will ensure

sufficient virus load. Note that viraemic blood frozen at -80°C seems to result in efficient engorgement and infection rates

(unpublished data). Once ticks are engorged, we recommend that viral DNA detection should not be used until 2 months

after inoculation in order to differentiate between successful viral replication within the tick and detection of residual virus

in the midgut. If viral DNA detection is performed on tick organs before 2 months post exposure, amplification of pig DNA

on ASFV-positive samples is required to control for potential contamination by gut content that may have occurred during

tick dissection (e.g. Bonsergent et al., 2023). Alternatively, the use of RT q-PCR with a reduced time frame is a viable option,

as it detects ASFV gene expression. Such standardized procedures will facilitate further investigations on the vector

competence of *Ornithodoros* ticks for ASFV, both expanding the range of tick species and virus genotypes tested and our

general knowledge on the biological mechanisms involved.

References

AlRyalat SAS, Malkawi LW, Momani SM (2019) Comparing Bibliometric Analysis Using PubMed, Scopus, and Web of Science Databases. *J. Vis.*

Exp., **152**. https://doi.org/10.3791/58494

Auerswald H, Maquart PO, Chevalier V, Boyer S (2021) Mosquito vector competence for japanese encephalitis virus. *Viruses*, **13**, 1–26.

https://doi.org/10.3390/v13061154

Azar SR (2019) Bloodmeal determinants of Zika virus infection of Aedes (Stegomyia) aegypti mosquitoes. PhD Thesis. University of Texas.

Bakkes DK, De Klerk D, Latif AA, Mans BJ (2018) Integrative taxonomy of Afrotropical Ornithodoros (Ornithodoros) (Acari: Ixodida: Argasidae).

Ticks and Tick-borne Diseases, **9**, 1006–1037. https://doi.org/10.1016/j.ttbdis.2018.03.024

Basto AP, Nix RJ, Boinas F, Mendes S, Silva MJ, Cartaxeiro C, Portugal RS, Leitão A, Dixon LK, Martins C (2006) Kinetics of African swine fever

virus infection in Ornithodoros erraticus ticks. *Journal of General Virology*, **87**, 1863–1871. https://doi.org/10.1099/vir.0.81765-0

Beltrán-Alcrudo D, Lubroth J, Depner K, Rocque SD La, Beltran-Alcrudo D, Lubroth J, Depner K, De La Rocque S (2008) African swine fever in the

Caucasus. *FAO Empres watch*, 1–8.

Bernard J (2015) Caractérisation de la compétence vectorielle des tiques Ornithodores pour le virus de la peste porcine africaine et étude de

deux déterminants : la relation souche virale-vecteur et l'influence de la salive de tiques sur l'infection chez le porc domestiqu. PhD Thesis.

University of Montpellier.

Boinas FS, Wilson AJ, Hutchings GH, Martins C, Dixon LJ (2011) The persistence of African swine fever virus in field-infected Ornithodoros

erraticus during the ASF endemic period in Portugal. *PLoS ONE*, **6**. https://doi.org/10.1371/journal.pone.0020383

Bonnet S, Liu X (2012) Laboratory artificial infection of hard ticks: a tool for the analysis of tick-borne pathogen transmission. *Acarologia*, **52**,

453–464. https://doi.org/10.1051/acarologia/20122068

Bonsergent C, Vittecoq M, Leray C, Jouglin M, Buysse M, McCoy KD, Malandrin L (2023) A soft tick vector of Babesia sp. YLG in Yellow-legged

gull (Larus michahellis) nests. *Peer Community Journal*, **3**. https://doi.org/10.24072/pcjournal.316

Burrage TG (2013) African swine fever virus infection in Ornithodoros ticks. *Virus Research*, **173**, 131–139.

https://doi.org/10.1016/j.virusres.2012.10.010

Burrage TG, Lu Z, Neilan JG, Rock DL, Zsak L (2004) African Swine Fever Virus Multigene Family 360 Genes Affect Virus Replication and

Generalization of Infection in Ornithodoros porcinus Ticks . *Journal of Virology*, **78**, 2445–2453. https://doi.org/10.1128/jvi.78.5.2445-

2453.2004

Carrascosa AL, Bustos MJ, de Leon P (2011) Methods for growing and titrating african swine fever virus: Field and laboratory samples. *Current*

Protocols in Cell Biology, 26.14.1-26.14.25. https://doi.org/10.1002/0471143030.cb2614s53

de Carvalho Ferreira HC, Tudela Zúquete S, Wijnveld M, Weesendorp E, Jongejan F, Stegeman A, Loeffen WLA (2014) No evidence of African

swine fever virus replication in hard ticks. *Ticks and Tick-borne Diseases*, **5**, 582–589. https://doi.org/10.1016/j.ttbdis.2013.12.012

Dixon LK, Chapman DAG, Netherton CL, Upton C (2013) African swine fever virus replication and genomics. *Virus Research*, **173**, 3–14.

https://doi.org/10.1016/j.virusres.2012.10.020

Drouin A, Chevalier V, Durand B, Balenghien T (2022) Vector Competence of Mediterranean Mosquitoes for Rift Valley Fever Virus: A Meta-

Analysis. *Pathogens*, **11**. https://doi.org/10.3390/pathogens11050503

EFSA, Nielsen SS, Alvarez J, Bicout DJ, Calistri P, Depner K, Drewe JA, Garin-Bastuji B, Gonzales Rojas JL, Schmidt C, Herskin M, Michel V, Pasquali

P, Roberts HC, Sihvonen LH, Spoolder H, Stahl K, Velarde A, Winckler C, Blome S, Boklund A, Bøtner A, Dhollander S, Rapagnà C, Van der

Stede Y, Miranda Chueca MA (2021) Research priorities to fill knowledge gaps in the control of African swine fever: possible transmission

of African swine fever virus by vectors. *EFSA Journal*, **19**. https://doi.org/10.2903/j.efsa.2021.6676

EFSA Panel on Animal Health and Welfare. (2010a) Scientific Opinion on African Swine Fever. *EFSA Journal*, **8**.

https://doi.org/10.2903/j.efsa.2010.1556

EFSA Panel on Animal Health and Welfare. (2010b) Scientific Opinion on the Role of Tick Vectors in the Epidemiology of Crimean-Congo

Hemorrhagic Fever and African Swine Fever in Eurasia. *EFSA Journal*, **8**. https://doi.org/10.2903/j.efsa.2010.1703

EFSA Panel on Animal Health and Welfare. (2014) Scientific Opinion on African swine fever. *EFSA Journal*, **12**.

https://doi.org/10.2903/j.efsa.2014.3628

Endris RG, Haslett TM, Geering G, Hess WR, Monahan MJ (1987) A Hemolymph Test for the Detection of African Swine Fever Virus in

Ornithodoros Coriaceus (Acari: Argasidae). *Journal of Medical Entomology*, **24**, 192–197. https://doi.org/10.1093/jmedent/24.2.192

Endris RG, Haslett TM, Hess WR (1991) Experimental Transmission of African Swine Fever Virus by the Tick Ornithodoros (Alectorobius)

puertoricensis (Acari: Argasidae). *Journal of Medical Entomology*, **28**, 854–858. https://doi.org/10.1093/jmedent/28.6.854

Endris RG, Hess WR (1994) Attempted Transovarial and Venereal Transmission of African Swine Fever Virus by the Iberian Soft Tick Ornithodoros

(Pavlovskyella)   marocanus   (Acari: Ixodoidea: Argasidae).   *Journal   of   Medical   Entomology*,   **31**,   373–381.

https://doi.org/10.1093/jmedent/31.3.373

Forth JH, Forth LF, Lycett S, Bell-Sakyi L, Keil GM, Blome S, Calvignac-Spencer S, Wissgott A, Krause J, Höper D, Kampen H, Beer M (2020)

Identification of African swine fever virus-like elements in the soft tick genome provides insights into the virus' evolution. *BMC Biology*, **18**,

1–18. https://doi.org/10.1186/s12915-020-00865-6

Franz AWE, Kantor AM, Passarelli AL, Clem RJ (2015) Tissue barriers to arbovirus infection in mosquitoes. *Viruses*, **7**, 3741–3767.

https://doi.org/10.3390/v7072795

Gillespie SH, Smith GL, Osbourn A (2004) Microbe-vector interactions in vector-borne diseases. In:, p. . Cambridge University Press.

Golnar AJ, Martin E, Wormington JD, Kading RC, Teel PD, Hamer SA, Hamer GL (2019) Reviewing the Potential Vectors and Hosts of African

Swine   Fever   Virus   Transmission   in   the   United   States.   *Vector-Borne   and   Zoonotic   Diseases*,   **19**,   512–524.

https://doi.org/10.1089/vbz.2018.2387

Guglielmone AA, Robbins RG, Apanaskevich DA, Petney TN, Estrada-Peña A, Horak IG, Shao R, Barker SC (2010) The Argasidae, Ixodidae and

Nuttalliellidae (Acari: Ixodida) of the world: a list of valid species names. *Zootaxa*, **2528**, 1–28.

Guinat C, Gogin A, Blome S, Keil G, Pollin R, Pfeiffer DU, Dixon L (2016) Transmission routes of African swine fever virus to domestic pigs: Current

knowledge and future research directions. *Veterinary Record*, **178**, 262–267. https://doi.org/10.1136/vr.103593

Haresnape JM, Wilkinson PJ (1989) A study of African swine fever virus infected ticks (Ornithodoros moubata) collected from three villages in

the ASF enzootic area of Malawi following an outbreak of the disease in domestic pigs. *Epidemiology and Infection*, **102**, 507–522.

Hess WR, Endris RG, Haslett TM, Monahan MJ, McCoy JP (1987) Potential arthropod vectors of African swine fever virus in North America and

the Caribbean basin. *Veterinary Parasitology*, **26**, 145–155. https://doi.org/https://doi.org/10.1016/0304-4017(87)90084-7

Jori F, Vial L, Penrith ML, Pérez-Sánchez R, Etter E, Albina E, Michaud V, Roger F (2013) Review of the sylvatic cycle of African swine fever in sub-

Saharan Africa and the Indian ocean. *Virus Research*, **173**, 212–227. https://doi.org/10.1016/j.virusres.2012.10.005

Kleiboeker SB, Burrage TG, Scoles GA, Fish D, Rock DL (1998) African Swine Fever Virus Infection in the Argasid Host, Ornithodoros porcinus

porcinus. *Journal of Virology*, **72**, 1711–1724. https://doi.org/10.1128/jvi.72.3.1711-1724.1998

Kleiboeker SB, Scoles GA (2001) Pathogenesis of African swine fever virus in Ornithodoros ticks. *Animal Health Research Reviews*, **2**, 121–128.

https://doi.org/10.1079/ahrr200133

De La Fuente J, Estrada-Pena A, Venzal JM, Kocan KM, Sonenshine DE (2008) Overview: Ticks as vectors of pathogens that cause disease in

humans and animals. *Frontiers in Bioscience*, **13**, 6938–6946. https://doi.org/10.2741/3200

Lubisi BA, Bastos ADS, Dwarka RM, Vosloo W (2005) Molecular epidemiology of African swine fever in East Africa. *Archives of Virology*, **150**,

2439–2452. https://doi.org/10.1007/s00705-005-0602-1

Malmquist WA, Hay D (1960) Hemadsorption and cytopathic effect produced by African Swine Fever virus in swine bone marrow and buffy coat

cultures. *American Journal of Veterinary Research*, **21**, 104–108. https://doi.org/10.5555/19602202203

Mellor PS, Wilkinson PJ (1985) Experimental transmission of African swine fever virus by Ornithodoros savignyi (Audouin). *Research in veterinary*

science, **39**, 353–361.

Nix RJ (2006) Interaction of African swine fever virus with its tick vector. PhD Thesis. University of Surrey.

Oura CAL, Edwards L, Batten CA (2013) Virological diagnosis of African swine fever-Comparative study of available tests. *Virus Research*, **173**,

150–158. https://doi.org/10.1016/j.virusres.2012.10.022

Pereira de Oliveira R, Hutet E, Duhayon M, Paboeuf F, Le Potier MF, Vial L (2020) No experimental evidence of co-feeding transmission of african

swine fever virus between ornithodoros soft ticks. *Pathogens*, **9**. https://doi.org/10.3390/pathogens9030168

Pereira De Oliveira R, Hutet E, Lancelot R, Paboeuf F, Duhayon M, Boinas F, Pérez de León AA, Filatov S, Le Potier MF, Vial L (2020) Differential

vector competence of Ornithodoros soft ticks for African swine fever virus: What if it involves more than just crossing organic barriers in

ticks? *Parasites and Vectors*, **13**, 1–15. https://doi.org/10.1186/s13071-020-04497-1

Pereira de Oliveira R, Hutet E, Paboeuf F, Duhayon M, Boinas F, de Leon AP, Filatov S, Vial L, Le Potier MF (2019) Comparative vector competence

of the Afrotropical soft tick Ornithodoros moubata and Palearctic species, O. Erraticus and O. Verrucosus, for African swine fever virus

strains circulating in Eurasia. *PLoS ONE*, **14**, 1–16. https://doi.org/10.1371/journal.pone.0225657

Plowright W, Parker J, Peirce MA (1969) African Swine Fever Virus in ticks (Ornithodoros moubata, Murray) collected from animal burrows in

Tanzania. *Nature*, **221**, 1968–1970.

Plowright W, Parker J, Staple RF (1968) The growth of a virulent strain of african swine fever virus in domestic pigs. *Journal of Hygiene*, **66**, 117–

134. https://doi.org/10.1017/S0022172400040997

Plowright W, Perry CT, Peirce MA, Parker J (1970) Experimental Infection of the Argasid Tick , Ornithodoros moubata porcinus , with African

Swine Fever Virus 1. , **31**, 33–50.

Portugal R, Coelho J, Höper D, Little NS, Smithson C, Upton C, Martins C, Leitão A, Keil GM (2015) Related strains of African swine fever virus

with different virulence: Genome comparison and analysis. *Journal of General Virology*, **96**, 408–419.

https://doi.org/10.1099/vir.0.070508-0

R Core Team (2022) R: A language and environment for statistical computing.

Ribeiro R, Otte J, Madeira S, Hutchings GH, Boinas F (2015) Experimental infection of Ornithodoros erraticus sensu stricto with two Portuguese

African swine fever virus strains. Study of factors involved in the dynamics of infection in ticks. *PLoS ONE*, **10**, 1–20.

https://doi.org/10.1371/journal.pone.0137718

Rowlands RJ, Duarte MM, Boinas F, Hutchings G, Dixon LK (2009) The CD2v protein enhances African swine fever virus replication in the tick

vector, Ornithodoros erraticus. *Virology*, **393**, 319–328. https://doi.org/10.1016/j.virol.2009.07.040

Vial L (2009) Biological and ecological characteristics of soft ticks (Ixodida: Argasidae) and their impact for predicting tick and associated disease

distribution. *Parasite*, **16**, 191–202. https://doi.org/10.1051/parasite/2009163191

Vijayendran D, Airs PM, Dolezal K, Bonning BC (2013) Arthropod viruses and small RNAs. *Journal of Invertebrate Pathology*, **114**, 186–195.

https://doi.org/10.1016/j.jip.2013.07.006

Zuur AF, Ieno EN, Walker N, Saveliev AA, Smith GM (2009) *Mixed effects models and extensions in ecology with R*. Springer Science & Business

Media. https://doi.org/10.1007/978-0-387-87458-6

Author contributions

JB coordinated the study, collected all bibliographic resources, extracted the data. HJ-P, SM and JO analysed the data. LV and M-FLP coordinated the study. JB, HJ-P, LV, M-FLP and FB wrote the manuscript.

Acknowledgements

The authors would like to thank Facundo Muñoz for his help with statistics.

Data, scripts and codes availability

All data presented here and the script used for statistical analyses are available https://doi.org/10.18167/DVN1/RYLTJ7.

559 **Appendix**

563   Annex 1: Article selection process according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.

564   Except for 2 articles, all the articles found in PubMed were also found in Scopus, creating 136 duplicates that were removed. The larger number

565   of references obtained in Scopus was expected since Scopus has larger scope and coverage than PubMed (AlRyalat et al., 2019)

568   Annex 2: Number of experiments reported for each virus strain, virus genotype and tick species in the whole dataset (900 experiments)

Virus strain		Virus genotype		Tick species	
asfvp99	30			coriaceus	58
brazil78	4			erraticus	396
chiredzi831	38			moubata	107
dr2	56			parkeri	4
lisbon60	10			porcinus	279
liv1333	46			puertoricensis	33
malta78	9			savignyi	8
mart	5			sonrai	4
netherlands86	4			turicata	2
nhp68	30			verrucosus	9
nu954	5	I	584		
ourt881	110				
ourt882	40				
portugal86	51				
sardinia	10				
senegal06	4				
tomar87	93				
vict901	4				
zimbabwe82	7				
zimbabwe83a	17				
zimbabwe83b	11				
georgia20071	42				
madagascar08	1	II	52		
ukr12zapo	9				
tengani62	19	V	19		
mwLil201	22	VIII	22		
kenrie1	20	IX	10		
ken06bus	10				
kwh12	30				
qet	12	X	176		
qew	17				
uganda61	97				
crocodil961	4	XIX	8		
nooitverwatcht966	4				
pretoriuskop9641	19	XX	19		
114a2	1				
Otu87	2	unknown	10		
transvaalcv	7				

Exposure route		Tick stage		Tick colony		Virus host		Viral titre		Geographic co-occurrence	
artificial	319	early	210	field	438	pig	364	$<10^4$	109	yes	747
capillary	79	late	522	lab	425	tick	389	10^4 to 10^6	296	no	153
natural	427	mix	131	NA's	37	warthog	145	$>10^6$	495		
inoculation	75	NA's	37			NA's	2				

576   Annex 4: Number of experiments reported for each virus strain, virus genotype and tick species in the dataset analysed (407 experiments).

Virus strain		Virus genotype		Tick species	
asfvp99	20			coriaceus	2
chiredzi831	16			erraticus	244
dr2	3			moubata	44
lisbon60	2			parkeri	1
liv1333	29			porcinus	103
malta78	1			puertoricensis	3
nhp68	26			savignyi	4
nu954	5			verrucosus	6
ourt881	61	I	291		
ourt882	21				
portugal86	25				
sardinia	4				
tomar87	69				
vict901	4				
zimbabwe82	1				
zimbabwe83a	2				
zimbabwe83b	2				
georgia20071	20	II	26		
ukr12zapo	6				
tengani62	6	V	6		
mwLil201	6	VIII	6		
kenrie1	4	IX	5		
ken06bus	5				
kwh12	20				
qet	7	X	72		
qew	8				
uganda61	33				
pretoriuskop9641	1	XX	1		

580   Annex 5: Number of experiments reported in each category of the studied experimental factors in the dataset analysed (407 experiments).

Exposure route		Tick stage		Tick colony		Virus host		Viral titre		Geographic co-occurrence	
artificial	169	early	81	field	249	pig	172	$<10^4$	68	yes	369
capillary	46	late	326	lab	158	tick	174	10^4 to 10^6	148	no	38
natural	154					warthog	61	$>10^6$	191		
inoculation	38										

583   Annex 6: Mean tick infection status (i.e., proportion of experiments where ticks were infected) in relationship with virus strain tested.

585 Annex 7: Five best models with the included variables influencing infection status (note that all models have «virus strain» as random effect),

586 corrected Akaike Information Criteria and weight of the model (probability that the model is the best out of all models considered).

Adaptation tick/virus	Exposure	Tick colony	Tick stage	Viral titre	Viral host	AICc	delta	weight
		+	+	+		393.6	0.00	0.506
+		+	+	+		395.6	2.01	0.185
		+	+	+	+	395.7	2.07	0.180
+		+	+	+	+	397.4	3.84	0.074
	+	+	+	+		399.6	5.97	0.026

588