1   Experimental design impacts the vector competence of *Ornithodoros* ticks for African swine fever virus:

2   a meta-analysis of published evaluations

Jennifer Bernard[1,2], Sara Madeira[3], Joachim Otte[4], Fernando Boinas[3], Marie-Frédérique Le Potier[2], Laurence Vial[1], Hélène

Jourdan-Pineau[1*].

[1] CIRAD, UMR ASTRE, F-34398 Montpellier, France.

ASTRE, Univ Montpellier, CIRAD, INRAE, Montpellier, France.

[2] Unité Virologie et Immunologie Porcines, Laboratoire de Ploufragan-Plouzané, Agence Nationale de Sécurité Sanitaire

(Anses), Univ Bretagne Loire, BP53, 22440 Ploufragan, France

[3] Interdisciplinary Centre of Research in Animal Health (CIISA), Faculty of Veterinary Medicine - University of Lisbon, Lisboa,

Portugal

[4] Berkeley Economic Advising and Research. 1442A Walnut Street, Berkeley, CA, 94705 USA.

*Corresponding author,

Correspondence: helene.jourdan@cirad.fr

Abstract

African Swine Fever (ASF) is a highly economically devastating viral disease for swine. Soft ticks of the genus *Ornithodoros* are involved in its epidemiology, as vectors and natural reservoirs of African Swine Fever Virus (ASFV). This constitutes a significant problem as it is currently challenging to eliminate endophilous soft ticks living in inaccessible, sheltered habitats. The vector competence of *Ornithodoros* ticks for ASFV has been mainly studied by mimicking natural tick-to-pig transmission pathways through experimental infections in the laboratory. However, experimental designs are anticipated to exert significant influence on the results, owing to inherent technical limitations of the approaches. By reviewing the original research studies dating back to the 1960s on the vector competence of *Ornithodoros* for ASFV, we estimated the vector competence of 10 tick species in association with 38 viral strains resulting in 51 tick-virus associations. This assessment emphasized the extensive range of protocol designs employed and of vector competence measurements. Those experimental factors have clear impact on the success of tick infection by ASFV. Our results offer valuable insights into how future trials can be refined by curtailing potential experimental biases. They also call for caution when interpreting conclusions drawn from studies on ASFV vector competence.

Keywords: Ornithodoros soft ticks, African swine fever virus, vector competence, methodology, meta-analysis

Introduction

African swine fever (ASF) ranks among the most damaging viral diseases affecting swine farming. The etiologic agent is

a large DNA virus belonging to the Asfaviridae family (Dixon *et al.*, 2013). African Swine Fever Virus (ASFV) causes

haemorrhagic fever in susceptible infected swine, with up to mortality rates of up to 100% in naïve domestic pigs. In Africa,

wild suids (warthogs, and possibly bushpigs and giant forest hogs) are natural vertebrate reservoirs that are resistant to the

disease (Jori et al. 2013). In Europe, wild boars are susceptible to ASFV and exhibit clinical symptoms and mortality rates

comparable to those of domestic pigs. In certain countries, soft ticks of the genus *Ornithodoros* act as vectors or natural

reservoirs of ASFV. The virus can persist for several years, potentially leading to epizootic outbreaks in pig farms that have

become colonized by ticks, even with the implementation of ASF control measures. (i.e., pig slaughtering, disinfection, and

quarantine) (Boinas et al. 2011, Jori *et al.*, 2013). Once ASFV is introduced into a new free area where the presence of

Ornithodoros ticks is highly suspected, as it occurred in the Caucasus in 2007 or more recently in the Caribbean, the

potential for long-term persistence of ASFV in tick reservoirs and ASFV transmission via tick bites raises concern (Beltran

Alcrudo *et al.*, 2008, EFSA, 2021.

Vector competence generally refers to the vector's capacity to be infected by a pathogen, to maintain and multiply this

pathogen, and finally to transmit it to a new susceptible host (Nuttal, 1994; Weaver et al., 2004). The biological steps taking

place within the arthropod vector can be broken down into four stages: (i) the ingestion of an infected blood meal in

sufficient amount, (ii) the infection of the vector upon crossing of its intestinal midgut by the pathogen, (iii) the persistence

and multiplication of the pathogen within the vector despite the vector's immune response, and (iv) the dissemination of

the pathogen inside the vector by crossing organic barriers and ultimately reaching organs necessary for transmission to

vertebrate hosts, typically the salivary glands. Sexual transmission of pathogens between males and females of vectors as

well as transovarial transmission from female vectors to their progeny can occur through the infection of reproductive

organs. Although not strictly included in the formal definition of the vector competence, these pathways of transmission

will also be considered in this review.

All these biological steps are observed in the interaction between ASFV and *Ornithodoros* tick although results reported

varied greatly between studies. In addition to historical demonstration of tick vector competence for ASFV in African *O.*

moubata sensu lato and Iberian *O. erraticus*, several other *Ornithodoros* tick species succeed to maintain or transmit ASFV

under experimental conditions (Endris et al., 1987, Endris et al., 1991, Hess et al., 1987, Mellor & Wilkinson, 1985). This

suggests that most *Ornithodoros* ticks might be able to maintain and transmit ASFV (EFSA, 2010b). Based on available

results, some researchers have proposed to predict the global role of each tick species as a vector and reservoir for ASFV

(Kleiboeker & Scoles, 2001; EFSA, 2010a, 2010b, 2014; Burrage, 2013). However, these comprehensive reports do not

account for the evident disparities in vector competence results, which likely rely on intricate tick-virus adaptations as

proposed by Pereira de Oliveira (2019, 2020c) and other yet-to-be-identified biological and experimental factors. No paper

has considered the substantial diversity of techniques employed to test vector competence except for a study on *O.*

erraticus infected with two Portuguese ASFV strains (Ribeiro et al., 2015). Nonetheless, the importance of experimental

methodology has been demonstrated in other models like mosquitoes with Rift Valley fever, Dengue, Zika or Japanese

Encephalitis viruses. (Azar, 2019; Auerswald, 2021; Drouin, 2022). We hypothesize that the experimental design, including

the choice of biological materials and the methods used to infect ticks and detect the virus, is an important factor in

explaining variation in vector competence, in addition to the influence of tick species, ASFV strains and their combinations.

To investigate this hypothesis, a systematic review of studies published up to 2022 was conducted, examining the vector

competence of *Ornithodoros* ticks for ASFV. The present paper focuses on the diverse experimental factors that may impact

vector competence and determines the overall vector competence of the various associations between *Ornithodoros* tick

and ASFV. We provide recommendations on how experimental trials can be refined to minimize biases and obtain more

comparable and reliable results on *Ornithodoros* vector competence for ASFV.

Material and Methods

Collection of bibliographical resources

The first objective of the study was to identify the bibliographic resources related to vector competence of *Ornithodoros*

ticks for ASFV, published in English, in peer-reviewed journals or under high publication standards. This was achieved

following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (Annex 1). Our

bibliographic review was conducted until July 2022, using three different search systems: EThOS

(http://ethos.bl.uk/Home.do), PubMed (http://www.ncbi.nlm.nih.gov/) and Scopus (http://www.scopus.com/). Queries

were built under the keywords "African swine fever" AND "tick". The topics not related with tick competence (e.g.,

immunology, genetic, vaccinology, epidemiology) were excluded during the screening phase. Then, during the eligibility

phase, studies were excluded if quantitative reliable data could not be extracted from publications. Purely descriptive

reviews without any original vector competence results were also excluded. Apart from published scientific papers, some

authors presented their results in their PhD manuscript. These data were also considered but all duplicated results were

removed to avoid biases in the analyses. Finally, the references cited at the end of the included papers were checked, in

order to identify further relevant studies. The final list of included papers that were used to extract all data in soft tick

vector competence for ASFV are shown in Table 1.

Table 1 - Included papers providing original results on the vector competence of *Ornithodoros* ticks for ASFV, with number attributed to each

paper by years of publication.

N°	Included papers
1	Parker J, Plowright W, Pierce MA. The epizootiology of African swine fever in Africa. Vet Rec. 1969 Dec 13;85(24):668-74
2	Plowright W, Perry CT, Peirce MA, Parker J. Experimental infection of the argasid tick, *Ornithodoros moubata porcinus*, with African swine fever virus. Arch Gesamte Virusforsch. 1970;31(1):33-50.
3	Plowright W, Perry CT, Peirce MA. Transovarial infection with African swine fever virus in the argasid tick, *Ornithodoros moubata* porcinus, Walton. Res Vet Sci. 1970 Nov;11(6):582-4.
4	Greig A. The localization of African swine fever virus in the tick *Ornithodoros moubata porcinus*. Arch Gesamte Virusforsch. 1972;39(1):240-7.
5	Plowright W, Perry CT, Greig A. Sexual transmission of African swine fever virus in the tick, *Ornithodoros moubata* porcinus, Walton. Res Vet Sci. 1974 Jul;17(1):106-13.
6	Groocock CM, Hess WR, Gladney WJ. Experimental transmission of African swine fever virus by *Ornithodoros coriaceus*, an argasid tick indigenous to the United States. Am J Vet Res. 1980 Apr;41(4):560-4.
7	Thomson GR. The epidemiology of African swine fever: the role of free-living hosts in Africa. Onderstepoort J Vet Res. 1985 Sep;52(3):201-9. Review.
8	Mellor PS, Wilkinson PJ. Experimental transmission of African swine fever virus by *Ornithodoros savignyi* (Audouin). Res Vet Sci. 1985 Nov;39(3):353-6.
9	Hess WR, Endris RG, Haslett TM, Monahan MJ, McCoy JP. Potential arthropod vectors of African swine fever virus in North America and the Caribbean basin. Vet Parasitol. 1987 Dec;26(1-2):145-55.
10	Endris RG, Haslett TM, Geering G, Hess WR, Monahan MJ. A hemolymph test for the detection of African swine fever virus in *Ornithodoros coriaceus* (Acari: Argasidae). J Med Entomol. 1987 Mar;24(2):192-7.
11	Hess WR, Endris RG, Lousa A, Caiado JM. Clearance of African swine fever virus from infected tick (Acari) colonies. J Med Entomol. 1989 Jul;26(4):314-7.
12	Haresnape JM, Wilkinson PJ. A study of African swine fever virus infected ticks (*Ornithodoros moubata*) collected from three villages in the ASF enzootic area of Malawi following an outbreak of the disease in domestic pigs. Epidemiol Infect. 1989 Jun;102(3):507-22.
13	Endris RG, Haslett TM, Hess WR. Experimental transmission of African swine fever virus by the tick *Ornithodoros* (Alectorobius) *puertoricensis* (Acari: Argasidae). J Med Entomol. 1991 Nov;28(6):854-8.
14	Endris RG, Hess WR, Caiado JM. African swine fever virus infection in the Iberian soft tick, *Ornithodoros* (Pavlovskyella) *marocanus* (Acari: Argasidae). J Med Entomol. 1992 Sep;29(5):874-8.
15	Endris RG, Hess WR. Experimental transmission of African swine fever virus by the soft tick *Ornithodoros* (Pavlovskyella) *marocanus* (Acari: Ixodoidea: Argasidae). J Med Entomol. 1992 Jul;29(4):652-6.
16	Endris RG, Haslett TM, Hess WR. African swine fever virus infection in the soft tick, *Ornithodoros* (Alectorobius) *puertoricensis* (Acari: Argasidae). J Med Entomol. 1992 Nov;29(6):990-4.
17	Endris RG, Hess WR. Attempted transovarial and venereal transmission of African swine fever virus by the Iberian soft tick *Ornithodoros* (Pavlovskyella) *marocanus* (Acari: Ixodoidea: Argasidae). J Med Entomol. 1994 May;31(3):373-81.
18	Boinas, FS. The role of *Ornithororos erraticus* in the epidemiolohy af ASFV in Portugal. PhD thesis manuscript. 1994
19	Rennie L. The Persistence of African swine fever virus in the Argasid Tick *Ornithodoros moubata*. PhD thesis manuscript. 1998.

Anderson EC, Hutchings GH, Mukarati N, Wilkinson PJ. African swine fever virus infection of the bushpig *(Potamochoerus porcus*) and its significance in the epidemiology of the disease. Vet Microbiol. 1998 Apr 30;62(1):1-15.

Kleiboeker SB, Burrage TG, Scoles GA, Fish D, Rock DL. African swine fever virus infection in the argasid host, *Ornithodoros porcinus porcinus*. J Virol. 1998 Mar;72(3):1711-24.

Kleiboeker SB, Scoles GA, Burrage TG, Sur J. African swine fever virus replication in the midgut epithelium is required for infection of Ornithodoros ticks. J Virol. 1999 Oct;73(10):8587-98.

Rennie L, Wilkinson PJ, Mellor PS. Effects of infection of the tick *Ornithodoros moubata* with African swine fever virus. Med Vet Entomol. 2000 Dec;14(4):355-60.

Rennie L, Wilkinson PJ, Mellor PS. Transovarial transmission of African swine fever virus in the argasid tick *Ornithodoros moubata*. Med Vet Entomol. 2001 Jun;15(2):140-6.

Burrage TG, Lu Z, Neilan JG, Rock DL, Zsak L. African swine fever virus multigene family 360 genes affect virus replication and generalization of infection in *Ornithodoros porcinus* ticks. J Virol. 2004 Mar;78(5):2445-53.

Nix-Rowlands RJ. Interaction of ASFV with its tick vector. PhD thesis manuscript. 2006.

Basto AP, Nix RJ, Boinas F, Mendes S, Silva MJ, Cartaxeiro C, Portugal RS, Leitão A, Dixon LK, Martins C. Kinetics of African swine fever virus infection in Ornithodoros erraticus ticks. J Gen Virol. 2006 Jul;87(Pt 7):1863-71.

Vial L, Wieland B, Jori F, Etter E, Dixon L, Roger F. African swine fever virus DNA in soft ticks, Senegal. Emerg Infect Dis. 2007 Dec;13(12):1928-31.

Rowlands RJ, Duarte MM, Boinas F, Hutchings G, Dixon LK. The CD2v protein enhances African swine fever virus replication in the tick vector, Ornithodoros erraticus. Virology. 2009 Oct 25;393(2):319-28.

Ravaomanana J, Michaud V, Jori F, Andriatsimahavandy A, Roger F, Albina E, Vial L. First detection of African Swine Fever Virus in *Ornithodoros porcinus* in Madagascar and new insights into tick distribution and taxonomy. Parasit Vectors. 2010 Nov 30;3:115.

Boinas FS, Wilson AJ, Hutchings GH, Martins C, Dixon LJ. The persistence of African swine fever virus in field-infected *Ornithodoros erraticus* during the ASF endemic period in Portugal. PLoS One. 2011;6(5):e20383.

Diaz AV, Netherton CL, Dixon LK, Wilson AJ. African swine fever virus strain Georgia 2007/1 in *Ornithodoros erraticus* ticks. Emerg Infect Dis. 2012 Jun;18(6):1026-8.

de Carvalho Ferreira HC, Tudela Zúquete S, Wijnveld M, Weesendorp E, Jongejan F, Stegeman A, Loeffen WL. No evidence of African swine fever virus replication in hard ticks. Ticks Tick Borne Dis. 2014 Sep;5(5):582-9.

Ribeiro R., Otte J., Madeira S., Hutchings G.H., Boinas F. Experimental Infection of Ornithodoros erraticus sensu stricto with Two Portuguese African Swine Fever Virus Strains. Study of Factors Involved in the Dynamics of Infection in Ticks. PLoS One. 2015 Sep 14;10(9):e0137718.

Pereira de Oliveira R, Hutet E, Paboeuf F, Duhayon M, Boinas F, Perez de Leon A, Filatov S, Vial L, Le Potier MF. Comparative vector competence of the Afrotropical soft tick Ornithodoros moubata and Palearctic species, *O. erraticus* and *O. verrucosus*, for African swine fever virus strains circulating in Eurasia. PLoS One. 2019 Nov 27;14(11):e0225657.

Pereira De Oliveira, R., Hutet, E., Duhayon, M., Paboeuf, F., Le Potier, M. F., & Vial, L. No experimental evidence of co-feeding transmission of african swine fever virus between Ornithodoros soft ticks. Pathogens 2020; 9(3), 168.

Pereira De Oliveira, R., Hutet, E., Duhayon, M., Guionnet, J. M., Paboeuf, F., Vial, L., & Le Potier, M. F.. Successful infection of domestic pigs by ingestion of the European soft tick *O. erraticu*s that fed on African swine fever virus infected pig. Viruses, 2020 12(3), 300.

Pereira De Oliveira, R., Hutet, E., Lancelot, R., Paboeuf, F., Duhayon, M., Boinas, F., ... & Vial, L. (2020). Differential vector competence of Ornithodoros soft ticks for African swine fever virus: What if it involves more than just crossing organic barriers in ticks?. Parasites & vectors 2020 13(1), 1-15.

Forth, J. H., Forth, L. F., Lycett, S., Bell-Sakyi, L., Keil, G. M., Blome, S., ... & Beer, M. Identification of African swine fever virus-like elements in the soft tick genome provides insights into the virus' evolution. BMC biology, 2020 18(1), 1-18.

Extraction of data about tick competence from bibliographical resources

Bibliometric general information on authors and years of publication were collected to describe the general interest in tick vector competence for ASFV and to identify scientific experts. We considered the experts as the persons in first and last position of the included papers. Information on tested *Ornithodoros* tick-ASFV associations was systematically reported. A challenge was to deal with missing or incomplete data for ASFV strains, as well as the ambiguities in tick species identification.

Regarding virus, we recorded the strain, the genotype (on the basis of p72 gene sequences, Qu *et al*. 2022), the geographical origin, and the host (vertebrate animals or ticks) from which the ASF virus was isolated. The original names of some ASFV strains were retrieved by crossing information from several papers, according to known authors' collaborations, or by directly contacting the authors. When the determination of strains remained impossible, the strain was identified by indicating the country where the ASFV strain was isolated, the year of the study, and a letter if duplicates existed. One exception was a strain used to infect *O. turicata (*Hess *et al.* 1987*)* without any indication of the country of origin which was named Unknown. Six recombinant ASFV strains, genetically modified and obtained from two natural strains (from papers n°25 and n°26), were excluded from this review since they cannot infect ticks in nature. Genotypes were not available for some strains of ASFV, but they could be inferred based on their geographical origin and ASFV genotype distribution published by Lubisi (Lubisi *et al.*, 2005).

Concerning ticks, the species, their geographical origin, their colony origin and status were reported. When ticks were reared for less than five years in the laboratory, they were considered comparable to ticks sampled from the field due to the limited number of generations occurring during this period. Some *Ornithodoros* ticks belong to species complexes and soft tick systematics has greatly changed during the last decades. Tick species identification was updated in accordance

with the revision of valid tick names by Guglielmone (Guglielmone *et al.*, 2010) and the mapping of their known distribution

(Vial, 2009). The more recent taxonomy study of southern African *Ornithodoros* ticks (Bakkes, 2018) could not be used, due

insufficiently precise location to attribute a species name among the *O. moubata* complex of species. Finally, when

determining tick geographical origin, 6 studies involving 4 different tick species (*O. coriaceaus, O. parkeri, O. moubata*, and

O. porcinus) did not report this information but since *O. parkeri* and *O. coriaceus* are known to be limited to North America,

especially the United-States (de la Fuente et al. 2008), they were assigned to this country.

Authors used various measurements to assess the tick vector competence for ASFV. Some authors reported numbers

of ticks positively detected with ASFV, while others scrutinized the virus titres in ticks or monitored ASFV kinetics in ticks

for several weeks or months, providing more comprehensive information on the virus persistence within its tick reservoir.

Some authors preferred to detect ASFV in tick transmission organs (salivary or coxal glands) or secretions (saliva or coxal

fluid). Such measurements suggest possible successful transmission to pigs although only transmission experiments can

confirm it. In these transmission experiments, clinical signs observed in pigs following tick bites confirm the ability of ticks

to transmit the disease. Detection of ASFV in pig blood is usually associated to confirm infection. Transmission between

ticks is either tested directly by measuring the proportion of transmitting ticks or the proportion of infected tick or indirectly

through the detection of ASFV in male reproductive organs (for sexual transmission) or female ovaries (for transovarial

transmission). However, this was not considered sufficient evidence to confirm such transmission pathways. Conversely,

the detection of ASFV in transmission secretions (e.g., saliva or coxal fluids) was considered sufficient evidence, as the virus

is already excreted in this case.

Based on these data, it was possible to evaluate three major steps of the vector competence of the ticks: 1) the tick

infection (revealed by the crossing of the tick midgut by ASFV), 2) the ability of ticks to transmit ASFV to pigs (named

horizontal transmission) and 3) their ability to transmit ASFV to other ticks through sexual or transovarial vertical

transmission (named tick-to-tick transmission). Although these steps are not necessarily gradual, transmission to vertebrate

hosts through tick bite or from ticks to ticks cannot occur without the previous infection of the tick, which was thus

considered as the main step to be assessed. In addition, tick ability to transmit ASFV to pigs or other ticks are less frequently

reported (N= 160 and N=54 over 900 experiments, respectively) likely because they are somewhat more difficult to assess

than tick infection. Therefore, for each experiment, we computed a binary variable, the infection status, according to any

measurements providing direct information on tick infection as well as indirect transmission measurements that imply

previous tick infection (Figure 1). More precisely, if ASFV transmission to pigs or ASFV transmission from ticks to other ticks

were reported in papers, the ticks were considered to be infected as the virus has necessarily crossed the tick organic

barriers to reach transmission organs and be transmitted.

If transmission failed or was not tested, the detection of ASFV in ticks was used as an indication of infection status.

Obviously when ASFV was not detected, ticks were considered as not infected. The only exception was when the detection

occurred on the same day as virus exposure, in which case no virus detection indicates a failure of experimental infection

or a too low detection threshold.

Different methods have been used by authors to detect ASFV in ticks, namely ASFV isolation/titration, observation of

ASFV particles, and viral DNA or RNA detection using multiple designs of Polymerase Chain Reaction (PCR). In the included

papers, ASFV isolation/titration was based on haemadsorption (HAD), which is the ability of many ASFV strains to aggregate

on red blood cells when they infect macrophages and create typical "rosettes" (Malmquist & Hay 1960); this implies that

the virus is able to replicate but does not necessarily confirm its entry in ticks. Similarly, detection of viral RNA also indicates

active viral replication. Conversely the detection of viral DNA can be due to residual virus from blood meal and does not

mean that the virus is still active. However, we considered that those methods brought equivalent results to assess infection

status once the duration after exposure was taken into account.

Indeed, when ASFV was detected in ticks, a minimum duration of 61 days between tick exposure to ASFV and detection

of ASFV in tick (hereafter DPE for duration post exposure) was chosen to distinguish successful viral replication inside the

tick from detection of residual virus after the infectious blood meal or virus inoculation. Indeed, the minimum duration for

blood meal digestion and clearance of ingested blood in soft ticks is approximately 4 weeks (Rowlands *et al.*, 2009).

Furthermore, if tick infection is unsuccessful, ASFV can be detected during 4 weeks by isolation (Ribeiro et al., 2015) and

after at least 49 days by qPCR (de Carvalho et al., 2014, see also Bernard, 2015, for a review on ASFV kinetics inside soft

ticks). If tick infection is successful, virus generalization in ticks takes about 30 to 56 days (Basto *et al.*, 2006, Burrage, 2013,

Kleiboeker *et al.*, 1998, Plowright *et al.* 1970). Therefore, and as a precaution, if ASFV was tested and detected before 61

DPE, the infection status was considered unknown unless virus was detected in organs other than the midgut. In this case,

this indicates that the virus has crossed the midgut barrier. When inoculation is based on injection into the haemocoel,

virus is spread throughout the tick. In this case, as the detection of virus in the organs was doubtful, the infection status

was recorded as unknown.

The experimental design used in the different studies was also reported, including: i) the exposure route for the tick

infection, ii) the ASFV titres in the inoculum used to infect the ticks (infectious dose), iii) the development stage of the ticks

tested in the study, iv) the DPE, v) the number of ticks tested in the study. Except for the DPE and the number of ticks

monitored, all parameters were categorized in different classes.

In field studies, viral titres in the hosts were not necessarily reported but they were considered to be high (>10[6]

HAD_{50}/mL of blood) considering the viraemia usually observed in ASF diseased pigs (Plowright et al., 1968). Tick stage was

classified as early from larva to nymph 3 and as late from nymph 4 to adult.

*Virus detection in the transmission organs is not a proof of transmission but the detection in the transmission fluids is a proof of transmission.
** we suspect in this case a failure in the experiment.

IS = infection status
DPE = date post exposure

Figure 1 - Criteria used to determine the infection status of ticks after their virus exposure, with three classes (IS: Infection Status; DPE: Date

post exposure; NA: Not Applicable, 1: Infected, and 0: Not Infected).

Statistical analysis of the tick infection status

A statistical analysis was carried out to identify factors potentially affecting the tick infection status. More precisely, we

investigated if factors related to the nature of tick-virus association or the experimental design influenced the infectious

status (see above for details) of the ticks in published experimental trials. The explanatory variables were: the ASFV strain

and genotype, the tick species, the potential geographical co-occurrence of tick and virus as an indication of possible tick-

virus adaptation (ASFV and ticks from the same country versus from different countries), the viral titre in infection blood

meal and the exposure route for tick infection, the tick stage and the tick colony status, the host from which the ASFV strain

was sampled. Using a dataset where all ticks were adults, we also tested the influence of tick sex on infectious status (along

with all other factors except tick stage). All statistical analyses were performed on R 4.2.2 (R Core Team, 2018). To assess

which factors influenced the infection status, we applied a generalized mixed model with a binomial family. The tick species,

virus genotype and virus strain were set as random effects to account for the fact that only a random fraction of soft tick

species and virus genotypes and strains were sampled. All other factors were set as fixed effects. The significance of random

effects was tested using a likelihood ratio test (Zuur et al., 2009). We then selected the best parsimonious models (with the

remaining 6 fixed effects) based on the AIC (Akaïke Information Criterion). In this best model, the significance of fixed effects

was tested using Chi² tests between models and post-hoc tests, with Tukey HSD correction, were performed.

Assessment of the overall tick vector competence for ASFV

Although our main assumption is that vector competence results are influenced by the experimental design, we

attempted to assess the overall vector competence of the various *Ornithodoros* tick-ASFV associations. Based on the three

main steps described above, we scored each tick-ASFV association, according to the following ranking: "score 0" for ticks

that were unable to become infected or for which it was impossible to conclude (see flow chart figure 2) ; "score 1" when

ASFV was able to cross tick midgut and replicate in ticks; "score 2" when transmission from tick to pig was successful ; and,

"score 3" for ticks validating all steps of vector competence, including ASFV transmission from ticks to other ticks. The

attribution of a score was based on all data available from the different included papers related to the considered tick-ASFV

association, and thus the longest period of monitoring and the best vector competence performance for this association.

Attributing graduate steps of vector competence sounds as if that associated transmission events are also graduate.

This is not the case: even if horizontal transmission scored 2 and sexual or transovarial transmissions scored 3, the success

of tick-to-tick transmission does not necessarily imply the efficiency of horizontal transmission. For example, *O. erraticus*

showed vertical transmission with the viral strain OURT88/1 but no transmission to pig (Pereira de Oliveira *et al*. 2019,

Pereira de Oliveira *et al.* 2020)

214   **Results**

Bibliometric description

The initial bibliographic search generated 236 references using Scopus and 138 by Pubmed, with 2 articles detected only

by Pubmed. The ETHOS database of thesis manuscripts and our manual bibliography search allowed identifying 8 more

resources. Among the final 246 references selected, 75 were reviews, 163 scientific papers and 8 thesis manuscripts. In this

pool of bibliographic references, 39 resources presenting vector competence original results were included in the final

dataset. They consisted of 5 field studies monitoring natural tick infections, 34 articles of laboratory experiments on ticks

and 2 mixed papers (Table 1). As shown in figure 1, the number of publications pertaining to the vector competence of

Ornithodoros ticks for ASFV was consistently low over the years, while the total number of papers related to ticks and ASFV

progressively increased from the 1980s to 2022.

Thirty-nine authors were reported as first and/or last authors of the selected publications and were considered as

"experts". Their period of expertise, including all their publications on ASFV & ticks, varied from 1 to 34 years. After 2015,

9 out of 39 experts continued to publish in this domain and can be considered the current core of experts on the vector

competence of ticks for ASFV.

228

Figure 1 - Number of bibliographic resources published each year, from 1966 to 2022, on ticks and ASFV (black dots = 246 screening papers)

and on the vector competence of *Ornithodoros* ticks for ASFV (red dots = 39 included papers). The names of first and last authors of the 39

included papers were reported in blue boxes at the top of the graphic. The length of boxes indicating their respective period of expertise. It is

the period where they appear in the list of authors from the 246 screening papers. List of the abbreviation author name: AEC: Anderson E.C.;

BAP: Basto A.P.; dCF: de Carvalho Ferreira H.C.; GWJ: Gladney W.J.; GCM: Groocock C.M.; LWL: Loeffen W.L.; McJ: McCoy J.P.; MMJ: Monahan

M.J.; RJ: Ravaomanana J.; TGR: Thomson G.R.

Data related to the biological material: ticks and ASFV

A database was created to record all the extracted data. Each record comprises a precise combination of any modalities

of informed indicators concerning the publication, the ticks, the ASFV strain and the experimental design. From the 39

included papers, 2036 records corresponding to 900 unique experiments were extracted. For some papers, authors detailed

the results of their experiment for each tick or with multiple records over time for the same ticks. Each experiment was

characterized by a combination of tick species, virus strain and experimental conditions (see Annexes 2 and 3). In total, the

tick species and ASFV strains represented 51 tick-virus associations and more than half (34/51) appeared in only one

publication.

For the ticks, 10 species were tested for their vector competence for ASFV. The Iberian tick *O. erraticus* was the most

studied (representing 44% of the reported experiments), followed by the *O. moubata* complex of species, including *O.*

porcinus and *O. moubata* sensu stricto (43%). The North American *O. coriaceus* (6.4%) were third, then they were far

followed by the other species *O. puertoricensis, O. verrucosus, O. savignyi, O. parkeri, O. sonrai, O. turicata* (3.7% - 1.0% -

0.8% - 0.4% - 0.4% - 0.2 % respectively). The geographical origins of ticks specified in the included papers are presented in

figure 2. All *O. erraticus* ticks tested came from Portugal, and most of them from the same region of Alentejo, while *O.*

moubata and *O. porcinus* ticks came from diverse countries, which is representative of their large distribution range in East

and Southern Africa.

A total of 38 different ASFV strains were tested. Most ASFV strains used in the studies were isolated from ticks and pigs

(43.1% and 40.4% of the reported experiments, respectively). Only 4 strains studied in 16.2% of the experiments were

isolated from warthogs and 0.2% of virus strains are from undetermined hosts. ASFV strains were geographically distributed

over 17 different countries, mostly in Africa, and belonged to 8 genotypes over the 23 genotypes described so far, with a

large representation of the genotype I (64.9% Figure 2). The genotype of the 114a and Transvaal CV strains from South

Africa remains unknown.

Figure 2 - Countries of origin of *Ornithodoros* tick species (colored according to the species) and ASFV genotypes (indicated by genotype number

associated with each country), involved in natural tick-ASFV associations studied in included papers. The origin of the tick *O. parkeri* was not

indicated but should be the United-States, as described in the literature.

Data related to the experimental design

On the 900 experiments recorded, half used ticks directly sampled from the field (especially *O. erraticus* and *O. porcinus*)

or reared in insectary for less than 5 years (48.7%). The other experiments used ticks reared in the laboratory for more than

5 years with some reared until 25 years for one study on *O. turicata* (47.2%). It was not possible to find information for the

remaining experiments. The majority of experiments (57.9%) involved ticks in their later developmental stages, ranging

from nymph 4 to adults. Early stages, from larva to nymph 3, were employed in 23.3% of the experiments. In the remaining

experiments, authors either utilized a combination of early and late stages (14.7%) or did not provide information about

the specific developmental stage used (4.1%). The number of ticks tested for each experiment was fairly low especially in

laboratory experiments (median=7 ticks tested, mean=30.03, range=[1;1141]).

Four methods were used to perform viral ASF inoculation to the ticks. Natural blood feeding on vertebrate hosts was

reported either in field studies, or in some laboratory experiments using experimentally infected pigs. This method

represented almost half of the total experiments conducted (47.4%). Two artificial feeding methods are reported: the

longest established method uses capillary tubes inserted on tick mouthparts (8.8%) while a more recent approach uses

animal skin, parafilm, or silicone membranes (35.7%). Another artificial infection route was the syringe inoculation of ASFV

directly in the tick heamoceal (8.1%). ASFV infectious dose used to infect ticks was very heterogeneous and thus classified

in three categories. Most experiments (78,4%) used medium to high viral doses (between 10^4 and 10^6 or over10^6 HAD$_{50}$/mL

of blood). The DPE was very variable, ranging from 1 month to up to 8 years (median value=58.0 days and mean value =

115,5 days), meaning that half experiments reported a DPE lower than the duration required to differentiate a successful

viral replication inside the tick from a detection of residual virus after the infection.

Factors influencing the likelihood of tick infection

On the initially 900 recorded experiments, we obtained a dataset of 408 observations with full information regarding

tick infection status and experimental conditions (Annex 4 and 7). Among the random effects tested, the tick species and

virus genotype had no effect on the infection status but the viral strain had (χ^2_1=4.787 p_value= 0.027). Model selection

based on AIC ended up with three variables displaying large and comparable importance across all models: virus titre, tick

stage and tick colony status (Figure 3 left). All those three variables had significant effects on infectious status (Figure 3

right).

In more details, a higher viral titre corresponded to a higher likelihood of tick infection (χ^2_2X²2= 38.497, p-

value=4.369×10-9). Bigger nymphs and adults were more prone to get infected than small nymphs (χ^2_1=17.888,

p_value=2.343×10-5). And ticks from laboratory colonies had higher probability of infection than ticks recently sampled in

the field (χ^2_1= 14.046, p_value=1.784×10-4). The potential geographical co-occurrence of tick and virus, the exposure route

for tick infection and the virus host were not significantly linked with infection status even if they were present in some of

the best models (Annex6. Using the dataset containing only experiments with adult ticks showed that sex had no influence

on infection status.

Figure 3 - *Left*: Relative importance of model terms across all models explaining infection status. The importance value for a particular predictor is equal to the sum of the weights for the models in which the variable appears. *Right*: Effect of tick stage, tick colony and virus titre on the infection status (all three effects are significant). Mean of infection status (i.e., proportion of experiments where ticks were infected). Values above the bars are the number of experiments in each level. Virus titre in HAD_{50}/mL of blood

Vector competence scores for ASFV

Figure 4 summarizes the vector competence score for each of the 51 associations between soft ticks and ASFV virus tested so far. In more details, 23.5% of the tick-ASFV associations presented a vector competence score null (score 0) i.e., ticks were unable to become infected or it was impossible to conclude on their infection status (Figure 2). For 27.5% of tick-ASFV associations, only tick infection (score 1) was achieved. Transmission from ticks to pigs was tested in 64.7% of tick-ASFV associations and was successful in 29.4% (score 2). Finally, 33.3% of tick-ASFV associations were assessed for ASFV transmission from ticks to ticks and 19.6% succeeded to transmit (score 3).

Figure 4 - ASFV vector competence scores for the different tick-ASFV associations studied in the included papers (score 0: no tick infection or

no information to conclude, score 1: successful tick infection, score 2: successful ASFV transmission to pigs, score 3: successful ASFV transmission

to other ticks). In each histogram bar, the number(s) are related to the included papers recorded in the table n°1. For each included paper, an

individual score is given according to available results in this paper. The overall vector competence score for a given tick-virus association is the

best vector competence result reported from all included papers related to this tick-ASFV association.

Only 27.5 % of tick-ASFV associations were fully tested for vector competence (including tick infection, transmission to

pig, and transmission to other ticks) and 15.7% were fully competent, meaning that they reached scores 1, 2 and 3.

Conversely, 3 associations reached score 3, without being tested for horizontal transmission (O. *erraticus*-Zimbabwe/83/a

and O. *erraticus*-Tomar/87) or by failing to transmit to pigs when tested (O. *erraticus*-Ourt/88/1). In one tick-ASFV

association (*O. porcinus*- MwL1/20/1) transmission capacity was suggested by virus secretion in saliva and in coxal fluid only

via virus direct injection into tick hemolymph.

At the species level, when considering only scores 1 and 2, most tick species tested were confirmed to be competent

vectors for at least one ASFV strain. This competence remains unclear for 3 species that were either tested with only one

ASFV strain and failed to transmit the virus to pig and/or to ticks (*O. parkeri* and *O. verrucosus*) or never tested for

transmission *(O. sonrai)*. Regarding ASFV transmission to other ticks, 6 tick species were tested and 4 (*O. moubata, O.*

porcinus, O. puertoricensis. and *O. erraticus)* succeeded for at least one ASFV strain whereas 2 (*O. coriaceus* and *O.*

verrucosus) that were tested with respectively three and one ASFV strains, failed. Finally, high heterogeneity of vector

competence scores was observed within a same tick species depending on ASFV strains used (e.g. score 3 for *O. moubata*-

LIV13/33 and score 0 for *O. moubata*-Brazil/78), and sometimes for a same tick-ASFV association (e.g. scores 0/1 and 2 for

two *O. turicata* colonies of different ages with the same ASFV strain).

336   **Discussion**

Bibliographic resources

This paper is the first exhaustive review of all the original studies published on the vector competence of soft ticks for

ASFV until 2023. Despite an exhaustive screening of bibliographic resources, only 39 papers were considered and the expert

panel had always been limited as exemplified by only 9 publishing experts since 2015. This relative paucity of bibliographic

resources regarding soft tick competence for ASFV may be attributed to the difficulty of investigating their vector

competence, both in laboratory and in field conditions. Indeed, sampling endophilous *Ornithodoros* ticks and developing

relevant methods to detect ASFV in natural tick populations (Jori *et al.*, 2013, Oura 2013) or rearing soft ticks and amplifying

ASF virus in the laboratory conditions are long and laborious tasks (Vial 2009, Carrascosa 2011). In addition, although ASF

is of high veterinary and economic concern, vectorial transmission is seen as occasional with a relatively minor role in

disease spread at least in some regions (Guinat et al., 2016). Nevertheless, soft ticks play a key role in the maintenance of

the virus and this is why characterizing competence for tick-virus associations is required to assess the risk of ASFV

introduction and spill over as Golnar (Golnar et al.2019) did it for the US.

Vector competence and tick-virus co-adaptation

Although all papers included in this study aimed to evaluate the vector competence of soft ticks for ASFV, we noticed a

considerable diversity of vector competence measurements that yield distinct information. As a consequence, vector

competence was only partially assessed for most tick-ASFV associations, with a noticeable decrease of studies along the

successive biological steps of vector competence, from the tick infection to ASFV transmission by tick to pigs, and then by

tick to other ticks. In addition, more than half of tick-ASFV associations were characterized by only one single article.

Heterogeneous vector competence scores were observed within species between viral strains. For instance, *O. moubata*

showed efficient vector competence for only 3 viral strains out of the 12 tested thus far. This could be interpreted either as

a lack of sufficient reliable studies or high tick-virus specificity. This latest hypothesis is in agreement with our results

showing a significant effect of viral strain on tick infection status.

Adaptation between arthropod vectors and viruses is based on genetic features defining susceptibility in both organisms

and the selection of susceptibility may be mediated partly by detrimental effects of arboviruses on vector physiology

(Weaver et al., 2004). In soft tick-ASFV model, the presence of ASFV MGF360 genes, the lectin-like EP153R gene, or the

CD2-like EP402R gene enhance the entry and the replication of ASFV into ticks (Burrage et al., 2004; Rowlands et al., 2009).

Leaving their original sylvatic cycle involving warthogs and ticks, some ASFV strains naturally lost these genes through long-

term circulation among Iberian domestic pigs, or by multiple passages on pig cell cultures for vaccine production (Portugal

et al., 2015). Concerning these vector susceptibility genes, some of them, such as MGF 360, are involved in the suppression

of the immune response of vertebrate and arthropod hosts against ASFV (Portugal et al., 2015). In addition, among the

large panel of immune responses described in arthropods against transmitted pathogens [44], arthropods are known to

display RNA interference as an important immune response against virus infection. The involvement of such molecular

pathway has recently been suggested in soft ticks with ASFV- like integrated elements coding for small interfering RNA in

O. moubata (Forth et al. 2020). More specifically, it appears that a greater number of these small matched with ASFV

genomes from genotype II than with genotype I, contributing to the variation in vector competence between viral strains.

We can suppose that field ticks, coevolving or not with ASFV, may develop a much more heterogeneous and adaptive

response against ASFV infection and show a lower success of infection, as it was already shown in some mosquito-virus

models (Weaver et al., 2004). Indeed, some authors previously reported very low ASFV infection rates in field ticks

(Plowright 1977, Haresnape & Wilkinson 1989, Boinas et al 2011), as well as over-mortality in laboratory ASFV highly

infected ticks (Endris & Hess, 1994, Kleiboeker & Scoles, 2001). In our study, laboratory tick colonies reared for more than

five years were shown to be significantly much more infected by ASFV than field ticks or recently established laboratory

ticks. In addition, tick from old laboratory colonies may be more prone to efficient artificial blood feeding, increasing the

blood volume and ASF virus intake.

Despite heterogeneous vector competence according to studies and viral strain tested, experimental evidence shows

that 7 *Ornithodoros* species from Africa (*O. moubata, O. porcinus and O. savignyi*), Europe (*O. erraticus*) and North America

(*O. coriaceus, O. turicata and O. puertoricensis*) can be infected by at least one strain of ASFV and then transmit it to pigs.

Effect of experimental factors on tick competence

Our analysis of vector competence papers demonstrated the importance of the experimental design on vector competence results. Several factors related to the choices of biological material and methodologies were tested. The ASFV titre in the inoculum for infecting ticks appears as a key parameter in the vectorial competence experiments, with likelihood of tick infection above 50% when ASFV load was higher than 10^6 HAD_{50}/mL. Accordingly, a threshold of $10^{5.75}$ HAD_{50}/mL was previously determined by Ribeiro (Ribeiro *et al.* 2015) for the specific ability of *O. erraticus* to maintain and multiply two Portuguese ASFV strains. Similar assumption was also proposed for the likelihood of ASFV transmission by ticks, with a threshold of 10^4 HAD_{50}/mL as a sufficient virus load per tick to insure the success of ASFV dissemination in ticks and the colonization of tick transmission organs (Kleiboeker & Scoles, 2001, Plowright, 1977, Ribeiro *et al.*, 2015). Forth *et al.* (2020) also demonstrated an impact of ASFV titre on tick infection probability and load. Soon after ingestion, ASFV has to cross the tick midgut barrier to be amplified in replication units localized in tick endothelial cells and then enter the tick hemolymph (Burrage et al., 2013). These results may suggest that the tick midgut and other tick membranes act as porous surfaces; the probability of ASFV entry would thus depend on the number of available pores (Franz *et al.*, 2015, Gutiérrez *et al.*, 2012). Tick immune system such as RNAi could also block low level of ASFV infection (Forth *et al.*, 2020).

The development stage of ticks had a significant impact on tick infection status, with better results for late stages (from nymph 4 to adults). This pattern confirms previous results on *O. erraticus*-Tomar and *O. erraticus*-OURT88/1 from Ribeiro (Ribeiro *et al.* 2015). This was explained by higher blood volume intake for late and large stages, and therefore highest ASFV titre inside the tick after bloodmeal digestion. In addition, we found no effect of observed sexual dimorphism in adults on the likelihood of tick infection which is in agreement with Ribeiro et al (2015). In this study, the authors suggest that, in adults, the number of ASFV particles already reached a certain threshold necessary to cause tick infection.

The route for tick infection was expected to play a role on tick competence. Natural feeding on vireamic pigs may provide

higher viral titre (usually >10^6 HAD$_{50}$/mL) than artificial feeding on membrane or with capillary (from 10^4-10^6 HAD$_{50}$/mL)

due to the difficulty for amplifying ASFV on pig cell cultures. Injection of virus into the haemocoel spread the virus

throughout tick body and remains traumatic for the tick (Bonnet & Liu, 2012). Despite differences between inoculation

methods, this does not seem to translate into any significant differences on the success rate of infection in our dataset. This

could be attributed to our dataset being incomplete and imbalanced, with the virus titre being a confounding factor. The

impact of inoculation methods could be specifically addressed by conducting dedicated studies comparing the various

methods with a similar virus titre. Similarly, we did not find any effect of the initial host in which viral strain was isolated on

tick competence.

Our review highlighted the huge diversity of measurements used in the included papers. They differ in their value for

informing the various steps of tick competence. Apart from PCR based viral detection, all the other methods reported from

included papers show some weakness to assess tick infection. For instance, ASFV isolation and titration on pig alveolar

macrophages was largely used by authors but this technique may suffer from inhibition issues due to cytotoxic effect of tick

components on pig macrophages and its achievement in at least 6 days is very time-consuming (Oura et al., 2013). In our

analysis, the main issue was to differentiate "residual" infection that consisted of remaining virus or viral DNA in the tick

midgut before complete digestion, from "true" infection where active virus crossed the tick midgut barrier and began its

replication in tick cells. Accordingly, we advise to use detection of viral DNA only 2 months after inoculation. Alternatively,

Reverse Transcription quantitative PCR could be used in a shorter delay, as it detects ASFV gene duplication and expression,

indicating ASFV crossing of tick midgut barrier where replication units are located. Finally, favoring oral inoculation of the

virus, instead of virus injection to the heamolymph, would avoid doubtful detection of ASFV and bypassed the immune

system avoiding a counterfeit infection (*O. porcinus*-MwLil/20/1). Since viral titre has the largest effect on tick infection

success, preferring blood from vireamic pigs would insure sufficient virus load and probably an ideal virus hemadsorption

to erythrocytes. Similarly, when feasible, late nymphs or adult ticks should be used since they will allow large blood intake.

430   **Conclusion and perspectives**

The examination of 39 papers on the vector competence of *Ornithodoros* soft ticks for ASFV highlighted factors

influencing vector competence results. For the same tick-ASF association, vector competence results may vary according

to the methodology used for infecting ticks (ASFV titre in the inoculum), the choice of biological material (viral strain, tick

development stage of tick, tick colony), and the techniques used to monitor and assess the likelihood of tick infection.

Although comparisons were difficult due to the scarcity of complete vector competence assessment for many tick-ASFV

associations, our analysis highlighted some standards that could be used for further investigations on the vector

competence of *Ornithodoros* ticks for ASFV.

References

Basto, A. P., R. J. Nix, F. Boinas, S. Mencles, M. J. Silva, C. Cartaxeiro, R. S. Portugal, A. Leitao, L. K. Dixon and C. Martins, 2006: Kinetics of African

swine fever virus infection in *Ornithodoros erraticus* ticks. *Journal of General Virology,* **87,** 1863-1871.

Beltran Alcrudo, D., J. Lubroth, K. Depner and S. De La Rocque, 2008: African swine fever in the Caucasus. *Empress watch***,** 1-8.

Bernard J. 2015 : Caractérisation de la compétence vectorielle des tiques Ornithodores pour le virus de la peste porcine africaine et étude de

deux déterminants : la relation souche virale-vecteur et l'influence de la salive de tiques sur l'infection chez le porc domestique. PhD

Thesis. University of Montpellier. https://www.theses.fr/2015MONTS099

Boinas, F. J. S., 1994: The role of *Ornithodoros erraticus* in the epidemiology of African swine fever in Portugal. University of Reading.

Boinas, F. S., A. J. Wilson, G. H. Hutchings, C. Martins and L. J. Dixon, 2011: The persistence of African swine fever virus in field-infected

Ornithodoros erraticus during the ASF endemic period in Portugal. *PloS one,* **6**.

Bonnet, S., Liu, X. 2012: Laboratory artificial infection of hard ticks: a tool for the analysis of tick-borne pathogen transmission. Acarologia **52**(4),

453-64.

Burrage, T. G., 2013: African swine fever virus infection in Ornithodoros ticks. *Virus Res,* **173,** 131-139.

Burrage, T. G., Z. Lu, J. G. Neilan, D. L. Rock and L. Zsak, 2004: African swine fever virus multigene family 360 genes affect virus replication and

generalization of infection in *Ornithodoros porcinus* ticks. *Journal of virology,* **78,** 2445-2453.

Carrascosa, A. L., Bustos, M. J., & Leon, P. (2011). Methods for growing and titrating African swine fever virus: field and laboratory samples.

Current protocols in cell biology, 26-14

de Carvalho Ferreira HC, Tudela Zúquete S, Wijnveld M, Weesendorp E, Jongejan F, Stegeman A, Loeffen WL. No evidence of African swine fever

virus replication in hard ticks. *Ticks Tick Borne Dis*. 2014 Sep;**5**(5):582-9.

de la Fuente J., Estrada-Peña A., Venzal JM., Kocan KM., Sonenshine, DE. (2008). Overview: ticks as vectors of pathogens that cause disease in

humans and animals. *Frontiers in Bioscience*, **1**(13), 6938-6946

Diaz, A. V., C. L. Netherton, L. K. Dixon and A. J. Wilson, 2012: African swine fever virus strain Georgia 2007/1 in *Ornithodoros erraticus* ticks.

Emerg Infect Dis, **18,** 1026-1028.

Dixon, L. K., D. A. Chapman, C. L. Netherton and C. Upton, 2013: African swine fever virus replication and genomics. *Virus Res,* **173,** 3-14.

EFSA, 2010b: Scientific Opinion on African Swine Fever, EFSA Panel on Animal Health and Welfare. *EFSA journal.*

European Food Safety Authority (EFSA), 2021: Research priorities to fill knowledge gaps in the control of African swine fever: possible

transmission of African swine fever virus by vectors. EFSA Journal, 19(6), e06676..

EFSA, 2014: Scientific Opinion on African Swine Fever Update, EFSA Panel on Animal Health and Welfare. EFSA journal.

Endris, R. G., T. M. Haslett, G. Geering, W. R. Hess and M. J. Monahan, 1987: A hemolymph test for the detection of African swine fever virus in

Ornithodoros coriaceus (Acari: Argasidae). *Journal of Medical Entomology,* **24,** 192-197.

Endris, R. G., T. M. Haslett and W. R. Hess, 1991: Experimental transmission of African swine fever virus by the tick *Ornithodoros (Alectorobius)*

puertoricensis (Acari, Argasidae). *Journal of Medical Entomology,* **28,** 854-858.

Endris, R. G. and W. R. Hess, 1994: Attempted transovarial and venereal transmission of African swine fever virus by the Iberian soft tick

Ornithodoros (Pavlovskyella) marocanus (Acari: Ixodoidea: Argasidae). *Journal of Medical Entomology,* **31,** 373-381.

Forth J.H., Forth L.F., Lycett S., Bell-Sakyi L., Keil G.M., Blome S., Calvignac-Spencer S., Wissgott A., Krause J., Höper D., Kampen H., Beer M.

2020: Identification of African swine fever virus-like elements in the soft tick genome provides insights into the virus' evolution. *BMC*

Biology, **18**, 1–18.

Franz, A. W., A. M. Kantor, A. L. Passarelli and R. J. Clem, 2015: Tissue Barriers to Arbovirus Infection in Mosquitoes. *Viruses,* **7,** 3741-3767.

Guglielmone, A. A., R. G. Robbins, D. A. Apanaskevich, T. N. Petney, A. Estrada-Peña, I. G. Horak, R. Shao and S. C. Barker, 2010: The Argasidae,

Ixodidae and Nuttalliellidae (Acari: Ixodida) of the world: a list of valid species names. *Zootaxa,* **2528,** 1-28.

Guinat, C., Gogin, A., Blome, S., Keil, G., Pollin, R., Pfeiffer, D. U., & Dixon, L. (2016). Review: Transmission routes of African swine fever virus to

domestic pigs: current knowledge and future research directions. *The Veterinary Record*, *178*(11), 262.

Gutiérrez, S., M. Yvon, E. Pirolles, E. Garzo, A. Fereres, Y. Michalakis and S. Blanc, 2012: Circulating virus load determines the size of bottlenecks

in viral populations progressing within a host.

Haresnape, J.M, Wilkinson P,J. Haresnape JM, Wilkinson PJ. A study of African swine fever virus infected ticks (Ornithodoros moubata) collected

from three villages in the ASF enzootic area of Malawi following an outbreak of the disease in domestic pigs. Epidemiol Infect. 1989

Jun;102(3):507-22.

Hess, W. R., R. G. Endris, T. M. Haslett, M. J. Monahan and J. P. McCoy, 1987: Potential arthropod vectors of African swine fever virus in North

America and the Caribbean basin. *Veterinary Parasitology,* **26,** 145-155.

Jori, F., L. Vial, M. L. Penrith, R. Perez-Sanchez, E. Etter, E. Albina, V. Michaud and F. Roger, 2013: Review of the sylvatic cycle of African swine

fever in sub-Saharan Africa and the Indian ocean. *Virus Res,* **173,** 212-227.

Kleiboeker, S. B., T. G. Burrage, G. A. Scoles, D. Fish and D. L. Rock, 1998: African swine fever virus infection in the argasid host, *Ornithodoros*

porcinus porcinus. *Journal of virology,* **72,** 1711-1724.

Kleiboeker, S. B. and G. A. Scoles, 2001: Pathogenesis of African swine fever virus in *Ornithodoros* ticks. *Anim Health Res Rev,* **2,** 121-128.

Kleiboeker, S. B., G. A. Scoles, T. G. Burrage and J. H. Sur, 1999: African swine fever virus replication in the midgut epithelium is required for

infection of *Ornithodoros* ticks. *Journal of virology,* **73,** 8587-8598.

Lubisi, B. A., A. D. Bastos, R. M. Dwarka and W. Vosloo, 2005: Molecular epidemiology of African swine fever in East Africa. Archives of virology,

150, 2439-2452.

Malmquist, W. A. and D. Hay, 1960: Hemadsorption and cytopathic effect produced by African Swine Fever virus in swine bone marrow and

buffy coat cultures. *American journal of veterinary research*, **21**, 104-108.

Mellor, P. S. and P. J. Wilkinson, 1985: Experimental transmission of African swine fever virus by Ornithodoros savignyi (Audouin). Res Vet Sci,

39, 353-356.

OIE - World Organisation for Animal Health, 2012: Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. OIE, Paris, chap 2.8.1, p

1067- 1081

OIE - World Organisation for Animal Health, 2012: Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. OIE, Paris, chap 2.8.1, p

1067- 1081

Oura, C. A., L. Edwards and C. A. Batten, 2013: Virological diagnosis of African swine fever-Comparative study of available tests. Virus Res, 173,

150-158.

Pereira de Oliveira R, Hutet E, Paboeuf F, Duhayon M, Boinas F, Perez de Leon A, Filatov S, Vial L, Le Potier MF, 2019. Comparative vector

competence of the Afrotropical soft tick Ornithodoros moubata and Palearctic species, O. erraticus and O. verrucosus, for African swine

fever virus strains circulating in Eurasia. PLoS One. Nov 27;14(11):e0225657.

Pereira De Oliveira, R., Hutet, E., Duhayon, M., Paboeuf, F., Le Potier, M. F., & Vial, L, 2020. a. No experimental evidence of co-feeding

transmission of african swine fever virus between Ornithodoros soft ticks. Pathogens; 9(3), 168.

Pereira De Oliveira, R., Hutet, E., Duhayon, M., Guionnet, J. M., Paboeuf, F., Vial, L., & Le Potier, M. F, 2020. b. Successful infection of domestic

pigs by ingestion of the European soft tick O. erraticus that fed on African swine fever virus infected pig. Viruses, 12(3), 300.

Pereira De Oliveira, R., Hutet, E., Lancelot, R., Paboeuf, F., Duhayon, M., Boinas, F., ... & Vial, L, 2020. c. Differential vector competence of

Ornithodoros soft ticks for African swine fever virus: What if it involves more than just crossing organic barriers in ticks?. Parasites &

vectors 2020 13(1), 1-15.

Plowright, W., 1977: Vector transmission of African Swine Fever Virus, Brussels.

Plowright W., Parker J. and R. F. Staple, 1968. The growth of a virulent strain of African swine fever virus in domestic pigs. J Hyg. 66(1):117-34.

Plowright, W., J. Parker and M. A. Pierce, 1969: The epizootiology of African swine fever in Africa. *Vet Rec,* **85,** 668-674.

Plowright, W., C. T. Perry, M. A. Peirce and J. Parker, 1970: Experimental infection of the argasid tick, *Ornithodoros moubata porcinus,* with

African swine fever virus. *Arch Gesamte Virusforsch,* **31,** 33-50.

Portugal, R., Coelho, J., Höper, D., Little, N. S., Smithson, C., Upton, C., ... & Keil, G. M. (2015). Related strains of African swine fever virus with

different virulence: genome comparison and analysis. *Journal of General Virology*, **96**(2), 408-419.

Qu H., Ge S., Zhang Y., Wu X., Wang Z. (2022) A systematic review of genotypes and serogroups of African swine fever virus. Virus Genes, 58,

77–87.

Rennie, L. F., 1998: The persistence of african swine fever virus in the argasid tick *ornithodoros moubata*. University of Hertfordshire.

Ribeiro, R., J. Otte, S. Madeira, G. H. Hutchings and F. Boinas, 2015: Experimental Infection of Ornithodoros erraticus sensu stricto with Two

Portuguese African Swine Fever Virus Strains. Study of Factors Involved in the Dynamics of Infection in Ticks. *PloS one,* **10,** e0137718.

Rowlands, R. J., M. M. Duarte, F. Boinas, G. Hutchings and L. K. Dixon, 2009: The CD2v protein enhances African swine fever virus replication in

the tick vector, *Ornithodoros erraticus*. *Virology,* **393,** 319-328.

Teglas, M. B., Drazenovich, N. L., Stott, J., & Foley, J. E. (2006). The geographic distribution of the putative agent of epizootic bovine abortion in

the tick vector, Ornithodoros coriaceus. *Veterinary parasitology*, *140*(3), 327-333.

Versteirt, V., Ducheyne, E., Schaffner, F., & Hendrickx, G. (2013). Systematic literature review on the geographic distribution of rift valley fever

vectors in Europe and the neighbouring countries of the Mediterranean Basin. *EFSA Supporting Publications*, **10**(4).

Vial, L., 2009: Biological and ecological characteristics of soft ticks (Ixodida: Argasidae) and their impact for predicting tick and associated disease

distribution. *Parasite-Journal De La Societe Francaise De Parasitologie,* **16,** 191-202.

Weaver, S. C., (2004): Vector competence. In: C. U. Press (ed.), *Microbe-vector interactions in vector-borne diseases.* Stephen H. Gillespie,

Geoffrey L. Smith, Anne Osbourn

Weiss, B. and S. Aksoy, 2011: Microbiome influences on insect host vector competence. Trends in parasitology, 27, 514-522.

Zuur AF., Ieno E.N., Walker J.N., Saveliev A.A., Smith G.M. (2009 Mixed Effects Models and Extensions in Ecology with R. Springer.

Appendix

546   Annex 1: Article selection process according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines

549 Annex 2: Number of experiments reported for each virus strain, virus genotype and tick species in the whole dataset (900 experiments)

Virus strain		Virus genotype		Tick species	
asfvp99	30			coriaceus	58
brazil78	4			erraticus	396
chiredzi831	38			moubata	107
dr2	56			parkeri	4
lisbon60	10			porcinus	279
liv1333	46			puertorice	33
malta78	9			savignyi	8
mart	5			sonrai	4
netherlands86	4			turicata	2
nhp68	30			verrucosus	9
nu954	5	I	584		
ourt881	110				
ourt882	40				
portugal86	51				
sardinia	10				
senegal06	4				
tomar87	93				
vict901	4				
zimbabwe82	7				
zimbabwe83a	17				
zimbabwe83b	11				
georgia20071	42				
madagascar08	1	II	52		
ukr12zapo	9				
tengani62	19	V	19		
mwLil201	22	VIII	22		
kenrie1	20	IX	10		
ken06bus	10				
kwh12	30				
qet	12	X	176		
qew	17				
uganda61	97				
crocodil961	4	XIX	8		
nooitverwatcht966	4				
pretoriuskop9641	19	XX	19		
114a2	1				
Otu87	2	unknown	10		
transvaalcv	7				

553 Annex 3: Number of experiments reported in each category of the studied experimental factors in the whole dataset (900 experiments). NA:

554 not available.

Exposure route		Tick stage		Tick colony		Virus host		Viral titer		Geographic co-occurrence	
artificial	319	early	210	field	438	pig	364	<10^4	109	yes	747
capillary	79	late	522	lab	425	tick	389	10^4 to 10^	296	no	153
natural	427	mix	131	NA's	37	warthog	145	>10^6	495		
inoculation	75	NA's	37			NA's	2				

556 Annex 4: Number of experiments reported for each virus strain, virus genotype and tick species in the dataset analysed.

Virus strain		Virus genotype		Tick species	
asfvp99	20			coriaceus	2
chiredzi831	16			erraticus	244
dr2	3			moubata	44
lisbon60	2			parkeri	1
liv1333	29			porcinus	104
malta78	1			puertorice	3
nhp68	26			savignyi	4
nu954	5			verrucosus	6
ourt881	61	I	291		
ourt882	21				
portugal86	25				
sardinia	4				
tomar87	69				
vict901	4				
zimbabwe82	1				
zimbabwe83a	2				
zimbabwe83b	2				
georgia20071	20	II	26		
ukr12zapo	6				
tengani62	6	V	6		
mwLil201	6	VIII	6		
kenrie1	4	IX	5		
ken06bus	5				
kwh12	21				
qet	7	X	73		
qew	8				
uganda61	33				
pretoriuskop9641	1	XX	1		

559 Annex 5: Number experiments reported in each category of the studied experimental factors in the dataset analysed.

Exposure route		Tick stage		Tick colony		Virus host		Viral titer		Geographic co-occurrence	
artificial	169	early	81	field	249	pig	172	$<10^4$	68	yes	370
capillary	47	late	327	lab	159	tick	174	10^4 to 10^6	148	no	38
natural	154					warthog	62	$>10^6$	192		
inoculation	38										

562 Annex 6: 5 best models with the included variables influencing infection status (note that all models have « virus strain » as random effect),

563 corrected Akaike Information Criteria and weight of the model (probability that the model is the best out of all models considered).

Adaptation tick/virus	Exposure	Colony type	Tick stage	Viral titer	Viral host	AICc	delta
		+	+	+		393.6	0.00
+		+	+	+		395.6	2.01
		+	+	+	+	395.7	2.07
+		+	+	+	+	397.4	3.84
	+	+	+	+		399.6	5.97