Malaria parasite resistance to azithromycin is not readily transmitted by mosquitoes

Hayley D. Buchanan[1,2*], Robyn McConville[2,3], Lee M. Yeoh[4,5], Michael F. Duffy[4,5], Justin A. Boddey[2,3], Geoffrey I. McFadden[1] & Christopher D. Goodman[1**].

1. School of BioSciences, University of Melbourne, Parkville, 3010, Vic, Australia
2. Walter and Eliza Hall Institute of Medical Research, Parkville, 3052, Vic, Australia
3. Department of Medical Biology, University of Melbourne, Parkville, 3010, Vic, Australia
4. Bio21 Institute, Parkville, 3052, Vic, Australia
5. Department of Microbiology and Immunology, School of Biomedical Sciences, University of Melbourne, Parkville, 3010, Vic, Australia

* Corresponding author (buchanan.h@wehi.edu.au)
** Second corresponding author (deang@unimelb.edu.au)

SUMMARY

Antimalarials are now used in combination with partner drugs to stem parasite drug resistance. Partners are often older, safe, cheap drugs, but resistance is already circulating for many, which raises the risk of selecting for multidrug resistance. If the partner drug(s) could be refractory to the spread of resistance, better resistance control could be implemented. We tested whether resistance to the antibiotic azithromycin, which kills malaria parasites by perturbing prokaryote-like protein synthesis in the apicoplast (relict plastid), had fitness costs to the spread of parasites via mosquitoes where parasites are not under drug pressure. Azithromycin resistance mutations in both rodent and human malaria parasites had a negative impact on the ability of resistant parasites to transmit from one vertebrate host to another via mosquitoes. Azithromycin resistance will therefore be less likely to spread geographically, making it an attractive option as a perennial partner compound to protect appropriate frontline antimalarials.

INTRODUCTION

Drug resistance is a major challenge to controlling, let alone eradicating, malaria[1]. Resistant
parasites emerge, spread geographically and eventually erode drug efficacy. A Sisyphean cycle
of using-then-losing malaria drugs has repeated itself at least four times, starting in the 1950s
with chloroquine and currently playing out with artemisinin, resistance to which has now spread
through the Greater Mekong Delta [2-6] and emerged in Africa [6-8]. Clearly, we need to do things
differently to gain more control of malaria drug resistance and stop relying on one promising
new drug to replace the previous one that we have set up for failure.

Combination therapies have slowed the spread of artemisinin resistance, and any new frontline
drug will require protection by some form of partner compound(s). Triple and even quadruple
combinations are now being considered [9, 10], but where do we turn for the partner drugs? Older,
inexpensive drugs with acceptable safety profiles are one option, but resistance is already
circulating for many, raising the risk of selecting for multidrug resistance [11]. Ideally, the partner
compounds should not only be safe, cheap, compatible, and effective but also be refractory to the
spread of resistance. This would prolong the life of the frontline compound whilst also
minimising the risk of selecting for multidrug resistance.

Atovaquone is an antimalarial refractory to the spread of resistance [12, 13]. Atovaquone resistance
is unable to be transmitted because resistant parasites with mutations in the mitochondrially-
encoded cytochrome *b* suffer fitness deficits during the mosquito phase of their life cycle where
they rely on higher rates of mitochondrial electron transport [13, 14]. In search of further resistance-
refractory drugs, we turned to other drug/target systems with similar life cycle constraints. We
hypothesised that changes to the parasite fitness landscape when switching from vertebrate to
mosquito hosts might also occur in other metabolic pathways that are more active in the insect
and/or liver phases of the life cycle. An obvious candidate is the apicoplast, a relict plastid of
malaria parasites that is substantially more active in mosquito and liver stages [15-17] and for which
a wide selection of safe, cheap inhibitors is known [18-20].

The apicoplast is a non-photosynthetic plastid homologous to the chloroplasts of plants and algae
[21]. Apicoplasts have their own DNA and express proteins encoded by their genome using
housekeeping machinery within the organelle [22]. Apicoplasts, like their chloroplast relatives,
arose by endosymbiosis of a photosynthetic cyanobacterium [23], and contain their own DNA
replication, RNA transcription, and protein translation and modification machinery, which is
bacterial in nature and sensitive to a raft of antibacterials [24]. For this reason, various
antibacterials, including azithromycin, are also parasiticidal [19, 20, 24-29].

Azithromycin is a broad-spectrum macrolide antibiotic that blocks apicoplast translation by
binding to the peptide exit tunnel of the 50S subunit of the apicoplast ribosome [30]. Like many
other drugs targeting the apicoplast, azithromycin has a key drawback as a therapeutic—its lethal
effect is delayed until the second intraerythrocytic after treatment. This "delayed death" response
results primarily from apicoplast metabolic fatigue that prevents parasite feeding during blood
stage, resulting in slow starvation [31]. Despite the delayed drug effects, azithromycin is used as

monotherapy to treat uncomplicated malaria and in combination with chloroquine or artesunate [27-29]. There are calls to increase the use of azithromycin as an antimalarial for combinations [32], and this push is strengthened by the recent discovery that azithromycin has two targets in malaria parasites: one in the apicoplast, and a second that inhibits parasite invasion of the red blood cell [33, 34]. Azithromycin is safe for infants and pregnant women and has a very long half-life in the body, all of which make it attractive as a partner compound [32].

Investigations into the role of the apicoplast in malaria parasites across the two-host life cycle show that the organelle is essential at every stage [35], but the roles of the apicoplast wax and wane across the life cycle [36]. In the blood stage the apicoplast has only two roles: to manufacture the five-carbon isoprenoid precursor isopentenyl diphosphate [18, 37], and to synthesise coenzyme A (CoA) [38]. By contrast, in the mosquito oocyst stage the apicoplast also has to make fatty acids [15] and heme [39-41], although these metabolic requirements are also parasite species-specific [36]. The apicoplast thus has different roles—and hence demands—in the two hosts, which implies that selection constraints on apicoplast efficiency will differ across the life cycle, much as they do with mitochondrial electron transport [13, 14, 42].

We hypothesised that malaria parasites resistant to azithromycin will experience differential selection across the parasite life cycle and incur a deferred fitness cost that restricts their ability to develop in mosquitoes and/or liver stages. Thus, apicoplast mutations allowing parasites to survive under drug pressure in the blood stage should be sub-optimal in mosquito and/or liver stages, when drug pressure is released. To test our hypothesis, we generated azithromycin resistant rodent (*Plasmodium berghei*) and human (*P. falciparum*) malaria parasites and assayed their ability to transmit from vertebrate-to-vertebrate via mosquitoes.

RESULTS

Generation of resistance to azithromycin in vivo in P. berghei and in vitro in P. falciparum

Azithromycin resistance has been reported in malaria parasites [30, 33], but to assay resistance
transmissibility, we generated new azithromycin resistant parasites in genetic backgrounds able
to be transmitted in the laboratory. In *P. berghei* ANKA, several cycles of
treatment→recrudescence→treatment with azithromycin at the ED_{95} (60 mg/kg q.d.; Figure 1A
and B, Figure S1) or ED_{99} (70 mg/kg q.d.; Figure 1A and B, Figure S1) yielded three,
independently derived lines of *P. berghei* azithromycin resistant parasites: PbAZMR_G95D_1,
PbAZMR_G95D_2 (both selected on a 70 mg/kg q.d. drug regimen), and PbAZMR_S89L
(selected on 60 mg/kg q.d.). The delayed-death kinetics of azithromycin make *ex vivo* drug trials
impossible for *P. berghei* so resistance phenotypes were confirmed *in vivo*. Stable resistance was
observed at the maximal allowable azithromycin dose of twice the ED_{99} dose (140 mg/kg q.d.)
(Figure 1C), but animal ethics constraints prevented treatment at higher doses and, therefore,
precluded ED_{50} determinations for these resistant strains. The slow growth rate of
PbAZMR_S89L (Figure S2) prevented *in vivo* confirmation of resistance after multiple attempts
failed to produce high enough blood stage parasitemia and exflagellations.

We selected for azithromycin resistance *in vitro* in the *P. falciparum* mosquito-transmissible
strain NF54 as previously described [30]. The IC_{50} of azithromycin in a 120-hour, delayed-death
assay for the parental line was 0.059 ± 0.017 µM (Figure 1D and E). Two independent cultures
of a clonal line of NF54, and one of a clonal line of the non-transmissible 3D7 strain, were
grown in the presence of 300 nM azithromycin for 196 hours and then maintained in drug-free
culture for 24-days, at which time parasites recrudesced. Resistance to azithromycin in both *P.*
falciparum strains was achieved in a single treatment cycle. Time to recrudescence was similar
for control parasites selected for resistance to pyrimethamine or atovaquone. An increase of drug
to 600 nM azithromycin to azithromycin selected cultures did not inhibit growth (Figure 1D).
Resistance to azithromycin at IC_{50}s ~100-fold greater than the parental line was maintained in all
lines following cloning (Figure 1D and E).

Azithromycin resistance in P. berghei and P. falciparum is apparently conferred by mutations in
Rpl4.

It is well established that azithromycin targets apicoplast translation in malaria parasites,
specifically the 50S ribosomal subunit [20, 30]. Moreover, reported resistance to azithromycin in
malaria parasites is conferred by apicoplast-encoded mutations [30, 33, 43]. Whole genome
sequencing of several clones of each of our independently generated *P. berghei* and *P.*
falciparum azithromycin resistant lines revealed mutations in a conserved region of the
apicoplast-encoded 50S ribosomal protein L4 (Rpl4; PBANKA_API00430 in *P. berghei* and
PF3D7_API01300 in *P. falciparum* respectively; Figure 1F). Two different point mutations in
Rpl4, S89L and G95D, were found in *P. berghei* (Figure 1F). Another mutation, G76V, was
found in Rpl4 of azithromycin resistant *P. falciparum* (Figure 1F). These Rpl4 point mutations
were observed in the absence of any other identifiably relevant point mutations, indels, or copy
number variations.

Azithromycin resistant P. berghei *parasites are severely impaired during mosquito stage*
development
To test the sexual viability of our *P. berghei* azithromycin resistant strains we quantified the
numbers of exflagellating males before each infection. There were no significant differences in
the number of exflagellations seen in PbAZMR_G95D_1, PbAZMR_G95D_2, and the Pb_WT
(Figure 2A). It is unclear if the reduction in PbAZMR_S89L parasites was a specific effect or
resulted from the poor blood stage growth of this mutant (Figure S2). Furthermore, this blood
stage growth defect limited us to only two independent experiments, so statistical analysis was
not robust.

We next infected female *Anopheles stephensi* mosquitoes with our azithromycin resistant *P.*
berghei lines and Pb_WT control parasites by allowing the mosquitoes to bite infected mice. In a
minimum of five independent mosquito infections with PbAZMR_G95D_1, PbAZMR_G95D_2,
and Pb_WT strains, there was no significant difference in the prevalence of mosquito infection,
but the resistant lines developed significantly fewer oocysts with means of 114 ± 12.4, $191\pm$
17.5, and 261 ± 15.8 oocysts per midgut respectively (Figure 2B). The development impairment
in PbAZMR_S89L parasites during mosquito midgut stages likely reflects a mix of specific
mosquito stage effects and very slow blood stage growth producing fewer sexual stages (Figure
2A) and therefore fewer oocysts.

At 20-26 days post blood feed, salivary glands were dissected from mosquitoes infected with
PbAZMR_G95D_1, PbAZMR_G95D_2, PbAZMR_S89L, and Pb_WT. Again, all our
azithromycin resistant lines produced significantly fewer salivary gland sporozoites than Pb_WT
(Figure 2C). The reduction in sporozoite numbers was markedly greater than would be expected
from the lower number of oocysts observed, and suggests a defect in development during the
oocyst stage.

To visualise oocyst stage development in our azithromycin resistant *P. berghei* strains, we
performed an immunofluorescence assay (IFA) on whole, infected mosquito midguts (Figure
2D). *P. berghei* azithromycin resistant oocysts had abnormal, dispersed staining with the
apicoplast-directed antibody to anti-acyl carrier protein (ACP), whereas Pb_WT oocysts showed
the expected branched apicoplast pattern (Figure 2D). ACP is encoded in the parasite nucleus
and targeted to the apicoplast [44, 45] and continues to be produced even if the apicoplast is not
intact. The dispersed ACP staining seen in our PbAZMR oocysts suggests that the apicoplast
structure has been disrupted, and ACP is likely accumulating in vesicles destined for the non-
existent or deformed apicoplast. Although IFA of whole midguts is informative, it is not
quantitative, partly because the epithelial and basal lamina cells of the *A. stephensi* midgut and
the *P. berghei* oocyst wall are significant barriers to consistent reagent entry and prevent
consistent antibody staining of oocysts.

To quantify the developmental defect in the apicoplasts of azithromycin resistant *P. berghei*
oocysts we measured their organllear DNA by quantitative PCR (qPCR), which revealed a loss
of between 74.3% and 82.8% of apicoplast genomes in the *P. berghei* azithromycin resistant
strains compared to Pb_WT (Figure 2E). Loss of the apicoplast genome confirms the

microcsopic observations that azithromycin resistance-conferring mutations in apicoplast-encoded Rpl4 disrupt apicoplast development during mosquito stages in *P. berghei*. The concomitant increase in mitochondrial genome copy number in all the azithromycin resistant strains compared to Pb_WT (Figure 2E) was not expected but could be due to abberent nuclear development (Figure 2D) and disruption of the timing of genome replication, perturbing normalisation of organelle genome number to nuclear gene numbers.

Azithromycin resistant P. falciparum *parasites develop normally in mosquitoes*

To determine if azithromycin resistance mutations impeded *P. falciparum* mosquito stage development, we selected a single clone (Pf_AZMR_C4, hereafter Pf_AZMR) to conduct downstream analyses. We compared the development of Pf_AZMR, and the parent line *P. falciparum* NF54 wildtype (Pf_WT) in female *A. stephensi* mosquitoes following standard membrane feeding. Both Pf_AZMR and Pf_WT parasite lines produced equivalent numbers of mature stage V gametocytes (Figure 2F). Similarly, both Pf_AZMR and Pf_WT parasites produced equivalent numbers of midgut oocysts at day 7 post blood feed, with an equivalent infection prevalence (Figure 2G). Furthermore, Pf_AZMR and Pf_WT parasites had no significant difference in the number of salivary gland sporozoites at day 17 post blood feed (Figure 2H). Thus, Pf_AZMR, despite having a point mutation in Rpl4 near those that severely disrupt mosquito infectivity of Rpl4 mutants in *P. berghei*, is apparently uninhibited in its ability to infect mosquitoes.

Azithromycin resistant P. berghei *sporozoites lose their apicoplast*

Immunofluorescence assays of apicoplast morphology in PbAZMR_G95D_1 and PbAZMR_G95D_2 sporozoites revealed the absence of a defined apicoplast, with only diffuse ACP staining visible throughout the length of the sporozoite (Figure 3A). This constrasts with Pb_WT sporozoites that have a canonical, punctate apicoplast structure adjacent the nucleus as expected (Figure 3A). The diffuse ACP staining seen in azithromycin resistant sporozoites (Figure 3A) is reminiscent of the dispersed apicoplast marker labelling in oocysts of azithromycin resistant parasites and may again indicate futile ACP trafficking. To quantify this aberration, we used IFA microscopy to assess >100 sporozoites from each parasite strain, across three independent mosquito infections (Figure 3B). Only 12.6% of PbAZMR_G95D_1 sporozoites, and 18.8% of PbAZMR_G95D_2 sporozoites exhibited intact apicoplasts, whereas 93% of Pb_WT sporozoites had clearly identifiable, punctate apicoplasts (Figure 3B). Apparently, the disruption of apicoplast development in oocysts that results from the azithromycin resistance-conferring Rpl4 mutations reduces apicoplast integrity during *P. berghei* mosquito stage development, culminating in the production of sporozoites that predominantly lack an apicoplast.

To assess potential viability of these azithromycin resistant sporozoites, we performed a sporozoite gliding motility assay comparing the movement of PbAZMR_G95D_1 and PbAZMR_G95D_2 sporozoites to Pb_WT sporozoites (Figure 3C). PbAZMR_G95D_1 and PbAZMR_G95D_2 sporozoites are somewhat impaired in their gliding motility compared to Pb_WT, but the differences were not statistically significant (Figure 3C). Motility of the PbAZMR_G95D_1 and PbAZMR_G95D_2 sporozoites, despite mostly lacking a distinct apicoplast, is congruent with the fact that they had successfully made the journey from the

midgut to the mosquito salivary glands (Figure 2B). Previous studies have noted, however, that
even a modest reduction in sporozoite gliding can severely impact *in vivo* infectivity of
sporozoites [46, 47]. Thus, the motility differences observed here, although not statistically
significant, may be biologically relevant to the infectivity of azithromycin resistant sporozoites,
particularly *in vivo*.

Azithromycin resistant P. berghei *parasites are critically diminished in their capacity to establish*
an infection in a naïve vertebrate host

After assaying the *P. berghei* azithromycin resistant parasites through mosquito stage
development and showing that they produce fewer sporozoites—most of which lack a distinct
apicoplast—we next tested their ability to establish a patent infection in a naïve vertebrate host,
either by bites from infected mosquitoes or direct intravenous (IV) injection of isolated
sporozoites. Neither PbAZMR_G95D_1 or PbAZMR_G95D_2 was able to establish a patent
infection in naïve mice bitten by at least 15 infected *A. stephensi* mosquitoes (Figure 4A).
Similarly, IV injection of $1x10^3$ PbAZMR_G95D_1 or $1x10^3$ PbAZMR_G95D_2 sporozoites
failed to produce patent infections (Figure 4B), whereas Pb_WT sporozoites reliably produced
infections with a time-to-patency (as observed by microscopy of Giemsa stained thin blood
smears) of 3-4 days (Figures 4A and B). When the inoculum of IV injected sporozoites was
increased to $1x10^4$ sporozoites, PbAZMR_G95D_2 sporozoites were able to establish a patent
blood stage infection in all of six attempts (Figure 4C). It is important to note that patency for the
resistant mutant required 4-13 days (mean 8.5 days ± 3.8 days; Figure 4C), whereas Pb_WT
sporozoites produced a patent infection in 3 days for all of 10 attempts (Figure 4C). DNA
sequencing confirmed that the G95D Rpl4 mutation was present in the newly established
PbAZMR_G95D_2 passage zero (P0) infection (data not shown). PbAZMR_G95D_1
sporozoites were unable to produce a patent infection in a naïve host in five attempts of IV
injection of $1x10^4$ sporozoites (Figure 4C). Thus, through substantial mechanical intervention to
provide a large sporozoite inoculum directly into a vein, we were able to achieve transmission of
PbAZMR_G95D_2 but not PbAZMR_G95D_1. Even then, the PbAZMR_G95D_2 parasites
showed a clear fitness defect over and above their liver stage delay, with blood stage parasites
failing to increase substantially in parasitaemia when monitored over 14 days post sporozoite IV
injection (Figure 4D).

These data again suggest there is a difference in phenotype penetrance between
PbAZMR_G95D_1 and PbAZMR_G95D_2, which recapitulates similar differences during
mosquito stage development. Despite both resistant mutants having the same azithromycin
resistance conferring Rpl4 mutation, there remains a slight fitness advantage in the
PbAZMR_G95D_2 parasites over the PbAZMR_G95D_1 parasites. There is, however, still a
significant deficiency in the PbAZMR_G95D_2 sporozoites in that they are only transmission-
competent when directly IV injected; and then only in massive numbers (Figures 4A-C).

The clear difficulty (or block in the case of PbAZMR_G95D_1) in the ability of our *P. berghei*
azithromycin resistant parasites to establish a patent infection in a naïve host points to liver stage
developmental defects, and the significant delay in patency of P0 infections induced by
PbAZMR_G95D_2 sporozoites is consistent with this. This is not surprising given the
importance of a proper functioning apicoplast during *P. berghei* liver stage development [48-51].

Indeed, while some transmission of PbAZMR_G95D_2 is feasible, the parasites exhibit impaired
development during mosquito stages, liver stages, and transmission to naïve hosts, identifying a
clear fitness cost associated with the G95D Rpl4 mutation.
280
Late liver stage development of azithromycin resistant P. berghei parasites is impeded
To investigate further the apparently aberrant liver stage development of our *P. berghei*
azithromycin resistant strains, we inoculated *in vitro* grown HC-04 hepatocytes with sporozoites
and observed the parasites by IFA to measure size and observe the morphology of the apicoplasts
and nuclei. Firstly, we wanted to confirm whether PbAZMR_G95D_1 sporozoites are even able
to infect liver cells given that they were unable to establish a blood stage infection in a naïve host
by any method attempted. Secondly, while PbAZMR_G95D_2 sporozoites are clearly impeded
in their ability to establish a blood stage infection in a naïve host, we wanted to examine the
reasons for the apparent liver stage developmental delay and requirement for such massive
sporozoite numbers to establish an infection in a naïve vertebrate host at all. Thirdly, we wanted
to observe any associated aberrations in apicoplast morphology and visualise development of
nuclei during liver stage development of PbAZMR_G95D_1 and PbAZMR_G95D_2.
293
294 Preliminary analyses revealed no differences in liver stage schizont size between azithromycin
295 resistant parasites and Pb_WT at 24-hours post inoculation (data not shown). We therefore
296 concentrated our efforts, and limited sporozoite supply, on a later liver stage timepoint of 48-
297 hours.
298
IFA of liver cells inoculated with PbAZMR_G95D_1, PbAZMR_G95D_2 or Pb_WT
sporozoites after 48 hours of development revealed marked difference in morphology (Figure
4E). WT liver stage schizonts, irrespective of size, typically showed reticulated ACP staining,
indicative of normal apicoplast morphology at this point of liver stage development [52] (top two
panels, Figure 4E). Furthermore, the nuclei of Pb_WT liver stage schizonts showed clear
differentiation with each nucleus beginning to be associated with a developing merozoite (top
two panels, Figure 4E). Contrast this with the PbAZMR_G95D_1 and PbAZMR_G95D_2 liver
stage schizonts at 48-hours post infection where the ACP staining marking the apicoplast was
cloudy and diffuse (bottom two panels, Figure 4E), reminiscent of what is seen in oocysts and
sporozoites with these strains. Additionally, the nuclei of PbAZMR_G95D_1 and
PbAZMR_G95D_2 developing liver stage schizonts appeared aggregated, forming unusual
donut-shaped clusters (bottom two panels, Figure 4E). These images are typical of what we
observed for 48-hour liver stages.
312
To quantify these observations, we used maximum projections of 48-hour post infection IFAs
over two biological replicates and measured the size of each liver stage schizont, and we also
quantified the nuclei in each developing parasite (Figure 4F and 4G). Both PbAZMR_G95D_1
(n=31) and PbAZMR_G95D_2 (n=47) liver stage parasites were significantly smaller than
Pb_WT (n=60; Figure 4F) and contained fewer nuclei (Figure 4G). Moreover, this reduction in
nuclei was independent of liver stage parasite size because normalising number-of-nuclei to the
area of the liver stage parasite optical section still showed significantly fewer nuclei in
PbAZMR_G95D_1 and PbAZMR_G95D_2 parasites compared to Pb_WT (Figure 4H). These
late liver stage data essentially recapitulate the phenotype seen during development throughout

mosquito stages; i.e. aberrant apicoplast morphology, and inability to properly differentiate nuclei. In sum, these defects in apicoplast integrity and nuclei biogenesis at both the oocyst and liver stages of the life cycle appear to severely hamper or even abrogate transmission of our *P. berghei* azithromycin resistant parasites.

Phenotypes converge: liver stage development of azithromycin resistant P. falciparum *parasites is impeded similar to* P. berghei

To investigate further the effects of apicoplast-encoded Rpl4 G76V azithromycin resistance mutation on the life cycle progression of *P. falciparum*, we measured the rates of liver cell traversal (3-hours post inoculation) and invasion (18-hours post inoculation) of Pf_AZMR and Pf_WT sporozoites (Figure 5A, 5B). Reflecting the apparently near-normal motility of azithromycin resistant *P. berghei* sporozoites, azithromycin resistant Pf_AZMR sporozoites were able to traverse (Figure 5A) and invade (Figure 5B) *in vitro* cultured HC-04 cells with comparable efficacy to Pf_WT sporozoites. Moreover, Pf_AZMR sporozoites harboured intact apicoplasts (data not shown), unlike the *P. berghei* azithromycin resistant parasites, which mostly lacked a distinct apicoplast (Figure 3B).

To mimic the *in vivo* life cycle phenotyping performed with azithromycin resistant *P. berghei* parasites, we infected humanised mice—with chimeric livers containing mostly human and some mouse hepatocytes—with Pf_AZMR sporozoites. Livers from humanised mice infected with either Pf_WT (n=two mice) or Pf_AZMR (n=three mice) sporozoites were harvested at day 5 post sporozoite IV inoculation and analysed by qRT-PCR to determine i/ the level of liver chimerism, and ii/ parasite loads. All mice had >75% human hepatocytes (data not shown), indicating a uniform infection potential, yet there was an 80% reduction in parasite load of Pf_AZMR parasites compared to Pf_WT (Figure 5C). Thus, Pf_AZMR parasites have a significantly reduced capacity to replicate in liver cells *in vivo*.

IFA on thin sections from these infected humanised mouse livers revealed defects in liver stage schizont development of Pf_AZMR in comparison to Pf_WT parasites (Figure 5D, 5E). The Pf_AZMR liver stage schizonts show little to no staining with either the cytosolic marker anti-HSP70 or the parasitophorous vacuole marker anti-PTEX150 (Figure 5D), indicative of severe overall morphological anomalies. Additionally, Pf_AZMR liver stage schizonts exhibit aberrant nuclear staining with no clear individual nuclei (Figure 5D). Indeed, nuclei often appeared to be aggregated (Figure 5E), reminiscent of the donut-shaped clusters of nuclei in *P. berghei* azithromycin resistant liver stages (Figure 4E). Moreover, whilst the size of the liver stage parasites was the same for Pf_AZMR in comparison to Pf_WT (Figure 5F), the azithromycin resistant parasites had significantly fewer nuclei (Figure 5G). Apicoplast morphology is also aberrant in Pf_AZMR late liver stage schizonts, with anti-ACP staining of Pf_AZMR revealing their apicoplasts to be collapsed and poorly defined in comparison to Pf_WT (Figure 5E).

Based on these severe defects of gross morphology, nuclear proliferation/separation, and apicoplast integrity for the Pf_AZMR parasites at liver stage, they appear to be experiencing severe fitness costs at day 5 of liver stage development attributable to the G76V Rpl4 mutation.

DISCUSSION

Rpl4 is a hotspot for macrolide resistance

Our four azithromycin selection regimens, across two *Plasmodium* species, all retrieved parasites with point mutations in the apicoplast-encoded ribosomal protein Rpl4. Mutations in Rpl4 are also associated with azithromycin resistance in other strains of *P. falciparum* [30, 33, 43], zoonotic infections of people with *Babesia microti* [53], in the plastid of the green alga *Chlamydomonas reinhardtii* [54], and in various bacteria related to the ancestor of plastids [55, 56]. Mutations in Rpl4 thus seem to be a universal mechanism of resistance to azithromycin and closely related macrolides.

Why are apicoplast Rpl4 mutations so prevalent in malaria parasite azithromycin resistance? We previously argued that malaria parasite organelle genomes, which are multi copy [57], are hotspots for resistance generation simply because more chances exist for mutations to arise than do in single copy nuclear genes [14]. Furthermore, organelle genomes could be heteroplasmic, allowing selection of pre-existing mutations by a drug to rapidly generate resistance, as likely occurs with atovaquone [14]. Moreover, several different Rpl4 mutations can confer azithromycin resistance (Figure 1F), somewhat akin to different mutations in mitochondrial cytochrome *b* being able to confer atovaquone resistance [58, 59], which provides higher mathematical probability of resistance arising in a particular gene. Whatever the case, the frequency of Rpl4 mutations in resistance to azithromycin, both in this and other studies, demonstrates that apicoplast Rpl4 is a hotspot for azithromycin resistance. Sequencing Rpl4 should be a first point-of-call for surveillance of malaria parasite azithromycin resistance.

Rpl4 mutations confer fitness advantage to parasites under drug pressure during the asexual, blood stages, but fitness deficits for transmission

Drug pressure is intermittent across the malaria parasite life cycle. Most drugs target the asexual blood phase in the human vertebrate host, yet drug pressure subsides when parasites progress through their sexual cycle in the mosquito host. Pressure can recommence if the parasites make it back into a new human host who is using antimalarials. At this point, a resistance-conferring mutation becomes an obvious selective advantage again, but the resistance allele(s) must transit the mosquito stages and the human liver to achieve transmission (and ultimately spread) of resistance. The fitness landscape for a parasite gene thus changes quite dramatically across the parasite life cycle.

Both *P. berghei* and *P. falciparum* are dependent on their apicoplast during mosquito stage development [15-17, 60, 61], and we set out to determine if mutations in apicoplast genes impacted parasite fitness in mosquitoes. Here, we showed that mutations conferring resistance to azithromycin (selected for by drug pressure during the blood phase) have differing consequences on fitness during mosquito phases. In *P. berghei,* the fitness deficit is severe in mosquito stages such as oocyst development and sporozoite production. However, in *P. falciparum* a different mutation in the same gene created no detectable fitness cost to the human parasite in mosquitoes. The difference between *P. berghei* and *P. falciparum* azithromycin resistant parasites in mosquito stage fitness likely relates to a combination of interacting factors. Firstly, the specific, resistance causing mutations in Rpl4 differ between parsite species (G95D or S89L in *P. berghei*

versus G76V in *P. falciparum*; Figure 1E). Secondly, each parasite species is differentially
reliant on its apicoplast in mosquitoes. For instance, based on the essentiality of apicoplast fatty
acid biosynthesis for mosquito phase development in *P. falciparum* [15] but not rodent malaria
species [48, 49, 62], one might expect *P. berghei* with a sub-optimal apicoplast to cope better in
mosquitoes. However, that was not the case for *P. berghei* Rpl4 mutants phenotyped here, all of
which exhibited marked reduction in fitness in mosquitoes. Differences in the mosquito host-
parasite interaction could also impact the metabolic requirements for sporozoite development. *A.*
stephensi is less susceptible to *P. berghei* than the natural mosquito host, *A. dureni* [63] whereas *A.*
stephensi is an efficient natural host of *P. falciparum* [64]. Thus, the differences in the infection
rates by different parasites might reflect the evolutionary adaptation of each parasite to our
provided mosquito host, as has been observed with immune responses in different
mosquito/parasite interactions [65, 66].

If a mutant parasite is able to complete the mosquito lifecycle stages, and sporozoites accumulate
in the mosquito salivary glands, its next phase—after being delivered by a bite—is to find and
infect the vertebrate liver. Again, drug resistance mutations undergo selection for sufficient
fitness during this liver phase. Interestingly, there is a convergence of phenotypes between *P.*
berghei and *P. falciparum* azithromycin resistant parasites at late liver stage, both having
defective apicoplasts and reduced/abnormal nuclei biogenesis. The malaria parasite liver stage is
tremendously proliferative with extremely high metabolic demands [67], and our phenotyping
indicates that a putatively sub-optimal apicoplast translation apparatus is apparently a major
impediment to liver development of both *P. berghei* and *P. falciparum*, *in vitro* and *in vivo*.

In *P. berghei*, the combined burden of poor/abnormal growth of azithromycin resistant parasites
in mosquitoes and liver blocked transmission, except where we provided massive mechanical
intervention. In *P. falciparum*, a severe detrimental effect in the liver stage *in vivo* (in humanised
mice) predicts a severe impairment of transmission. Whilst we were unable to test whether the
Pf_AZMR parasites are competent to progress from liver stage to blood stage and thus achieve
transmission, it seems likely that the *P. falciparum* azithromycin resistant mutant—despite
performing better in mosquitoes—will nevertheless experience impediments to transmission.
Whether or not the G76V mutation will block or reduce transmission of azithromycin resistance
in a real-world setting is not yet possible to say, but the indications are that the mutation will not
circulate freely, which should ultimately result in reduced transmission of azithromycin
resistance, which gives azithromycin one more desirable characteristic as a potential partner
drug. Refractoriness to spread of resistance would also mean that use of azithromycin for
prophylaxis might pose less of a risk for resistance selection than is commonly assumed.

ACKNOWLEDGEMENTS

Financial support from the National Health and Medical Research Council (GNT2016391, GNT1162550, GNT1143974), the Australian Research Council (FL170100008, LE200100181), and an Australian Postgraduate Scholarship (HB) are gratefully acknowledged. We thank Anton Cozijnsen, Vanessa Mollard and Ryan Steel for technical support.

FIGURE LEGENDS

Figure 1. Summary drug assays performed *in vivo* with *P. berghei* and *in vitro* with *P. falciparum*. A.) Peters' four-day suppressive test *in vivo* with *P. berghei*. Percent suppression of a variable dosing range used to determine both the appropriate clearance dose and potential resistance-generating dose. Parasitaemias of individual mice in each dose cohort were normalised to the mean vehicle control-treated parasitaemia at each time point, and the mean percent-suppression value plotted (error bars represent ±SD). A non-linear regression curve with a variable slope was then fitted to the data points for each dose cohort; B.) End-point (day 4) data for the Peters' four-day suppressive test. Parasitaemias of individual mice in each dose cohort were normalised to the mean vehicle control-treated parasitaemia at day 4 and each individual value plotted. A non-linear regression curve with a variable slope was then fitted to the data points. Where n<4, mice were euthanised prior to the experimental end point for humane reasons; C.) *In vivo* drug assay of PbAZMR_G95D_1 (teal) and PbAZMR_G95D_2 (light blue) grown with a full four-day 70 mg/kg q.d. (solid lines, n=2 per genotype) or 140 mg/kg q.d. (dashed lines, n=2 per genotype) drug course versus WT with the vehicle (no drug; black; n=2) and a full four-day 70 mg/kg q.d. drug course (grey; n=2). Treatment was started on day 1. (Values indicate mean parasitaemia, and error bars represent the SEM); D.) Dose response curves of Pf_AZMR_C4 and Pf_AZMR_D2 clones (both with G76V mutation in *P. falciparum* 50S Rpl4) compared to Pf_WT. Data presented are the mean of three technical replicates and representative of three independent drug trials (error bars represent ±SD); E.) Comparison plot of IC50s of *P. falciparum* azithromycin resistant clones versus WT. Plotted are the mean IC50 values from three independent drug trials. (error bars represent the SEM, **** = $P < 0.0001$, ordinary one-way ANOVA, Dunnett's multiple comparisons test); F.) Alignment of region of Rpl4 associated with azithromycin of erythromycin resistance in bacteria and plastids of parasites, algae and plants. Mutations identified in this study are depicted in red and mutations selected from the literature in blue. Included organisms: P. berghei (*Plasmodium berghei*); P. falciparum (*Plasmodium falciparum*); T. gondii (*Toxoplasma gondii*); B. microti (*Babesia microti*); C. velia (*Chromera velia*), E.coli (*Escherichia coli*); D. radiodurans (*Deinococcus radiodurans*); N. gonorrhoeae (*Neisseria gonorrhoeae*); S. pneumoniae (*Streptococcus pneumoniae*); C. reinhardtii (*Chlamydomonas reinhardtii*);A.thaliana (*Arabidopsis thaliana*). Asterisks denote invariant residues, colons depect conserved residues groups with strongly similar properties, and periods denote semi-conserved residues groups with similar properties. (AZM = azithromycin; ED = effective dose)

Figure 2. Mosquito infection data for both *P. berghei* and *P. falciparum* azithromycin resistant parasites. A.) Exflagellations were counted per 1×10^4 red blood cells (RBCs) and are shown as the mean ±95%CI, with no significant difference between strains (ordinary one-way ANOVA with Dunnett's multiple comparisons; WT [n=17], PbAZMR_G95D_1 [n=20],

PbAZMR_G95D_2 [n=10], PbAZMR_S89L [n=2], where 'n' is the number of independent
biological replicates). At 9-to-14 days post blood feed, a minimum of 10 midguts were dissected
for each 'n' independent infection (WT [n=13], PbAZMR_G95D_1 [n=11], PbAZMR_G95D_2
[n=5], PbAZMR_S89L [n=2]) and assessed for infection prevalence (pie charts below,
percentage indicates the mean; ns= not significant by way of ordinary one-way ANOVA).
Oocysts of infected midguts were counted and pooled, and non-zero values were plotted (error
bars represent the mean ±95%CI). All azithromycin resistant strains had significantly fewer
oocysts than WT (*=P<0.05; ****=P<0.0001; ordinary one-way ANOVA, Dunnett's multiple
comparisons test); B.) Mean number of salivary gland sporozoites per mosquito. Between days
20- and 26- post blood feed, a minimum of 12 mosquitoes (WT) or 20 mosquitoes
(PbAZMR_G95D_1, PbAZMR_G95D_2, and PbAZMR_S89L) were dissected. All
azithromycin resistant strains produce fewer salivary gland sporozoites than WT
(****=P<0.0001, Welch's t-test; error bar represents +95% CI). Results are from 'n'
independent mosquito infections: WT (n=21), PbAZMR_G95D_1 (n=14), PbAZMR_G95D_2
(n=10), PbAZMR_S89L (n=2); C.) Quantitative PCR on genomic DNA isolated from midguts
infected with both *P. berghei* azithromycin resistant parasites and WT strains. Copy numbers of
genes encoded in each organelle were first normalised to nuclear genes, then expressed as a ratio
to WT parasites. There is a clear reduction in apicoplast genome copy number in
PbAZMR_G95D_1 (n=4), PbAZMR_G95D_2 (n=3) and PbAZMR_S89L (n=1), where
n=number of biological replicates. Error bars represent the SEM; D.) Immunofluorescence assay
on midgut oocysts, 14 days post blood feed. Green represents labelling of the *P. berghei*
apicoplast, with anti-ACP antibodies. Hoechst is staining DNA. PbAZMR_G95D_1 oocysts
consistently show dispersed ACP staining (bottom panel), while WT oocysts show a branched
ACP staining (top panel). Images are maximum projections (except brightfield, which shows the
largest single slice). Scale bar=10μm; E.) Quantification of development of blood-feed-ready
stage V gametocytes produced by Pf_WT and Pf_AZMR parasites. No significant difference is
seen (Unpaired t-test, two-tailed P value); F.) At 7 days post blood feed, a minimum of 10
midguts were dissected for each 'n' independent infection (Pf_WT [n=3], Pf_AZMR [n=3]) and
assessed for infection prevalence (pie charts below, percentage indicates the mean; ns= not
significant by way of Unpaired t-test, two-tailed P value). No significant difference was observed
between the number of midgut oocysts in mosquitoes infected with either Pf_WT or Pf_AZMR
parasites (Unpaired t-test, two-tailed P value); G.) At day 17 post blood feed, a minimum of 12
mosquitoes were dissected per genotype. No significant difference was observed between the
number of salivary gland sporozoites in mosquitoes infected with either Pf_WT or Pf_AZMR
parasites (Unpaired t-test, two-tailed P value).
529
Figure 3. Immunofluorescence assay on PbAZMR_G95D_1, PbAZMR_G95D_2 and WT
sporozoites reveals that azithromycin resistant sporozoites lose their apicoplast. A.) Sporozoites
were stained with anti-circumsporozoite protein (CSP; red) to mark the outer surface of the
sporozoite, anti-ACP (green) to mark the apicoplast, and Hoechst (blue) to mark the nucleus. WT
sporozoites have a clear, punctate apicoplast (top panel), while azithromycin resistant
sporozoites have diffuse apicoplast marker staining (bottom two panels). Images are
representative of the calculated majority; B.) Quantification of apicoplast presence (API+, green)
and absence (API-, red) in WT (n=2068), PbAZMR_G95D_1 (n=445), and PbAZMR_G95D_2
(n=855) sporozoites, where n=number of total sporozoites counted. A minimum of 110
sporozoites were counted for each parasite strain, for each of three independent mosquito

infections. Green bars indicate the mean percentage apicoplast positive apicoplast, error bar
represents +95% CI. Both azithromycin resistant strains produce sporozoites with significantly
fewer intact apicoplasts than WT (P<0.0001, Fisher's exact test); C.) Quantification of
sporozoite motility trails from PbAZMR_G95D_1 (n=159), PbAZMR_G95D_2 (169), and WT
(n=175) sporozoites, from two independent mosquito infections. There was no significant
difference in types of trails observed between each parasite line (chi-square test).
546
Figure 4. Transmission of azithromycin resistant *P. berghei* parasites to a naïve host is severely
impaired, and *in vitro* liver stage development appears stalled. A.) Kaplan-Meier curves of P0
infections by biting. At least 15 infected mosquitoes were allowed to feed on naïve mice for at
least 15 minutes and mice were monitored for 14 days by Giemsa smear (one mouse infected
with PbAZMR_G95D_1 was only monitored until day 7, indicated by the initial triangle
marker). Number of transmission attempts per strain were WT, n=8; PbAZMR_G95D_1, n=4;
PbAZMR_G95D_2, n=4. Naïve mice infected with azithromycin strains by biting failed to
establish an infection, and this was significantly different than WT (P=0.0007, Mantel-Cox test)
B.) Kaplan-Meier curves of P0 infections by $1x10^3$ IV of sporozoites. Mice were monitored for
14 days by Giemsa smear. Number of transmission attempts per strain were WT, n=6;
PbAZMR_G95D_1, n=1; PbAZMR_G95D_2, n=4. Naïve mice infected with azithromycin
strains by $1x10^3$ IV failed to establish an infection, and this was significantly different than WT
(P=0.01, Mantel-Cox test). C.) Kaplan-Meier curves of P0 infections by $1x10^4$ IV of sporozoites.
Mice were monitored for 14 days by Giemsa smear. Number of transmission attempts per strain
were WT, n=10; PbAZMR_G95D_1, n=5; PbAZMR_G95D_2, n=6. Only naïve mice infected
with either WT or PbAZMR_G95D_2 by $1x10^4$ IV were able to establish an infection, but
infection with PbAZMR_G95D_2 sporozoites showed delayed patency (mean 8.5 days ± 3.8
days [SD]) and impaired growth. This was significantly different than WT (P<0.0001, Mantel-
Cox test). D.) Growth curve comparison of P0 infections established by WT and
PbAZMR_G95D_2 by $1x10^4$ IV of sporozoites. Mice infected with PbAZMR_G95D_2
sporozoites grow slowly. Points shown indicate mean parasitaemia (±SEM), WT, n=9;
PbAZMR_G95D_2, n=6, where 'n' represents the number of infected mice monitored for the
period; E.) Immunofluorescence assay of liver cells infected with WT, PbAZMR_G95D_1 and
PbAZMR_G95D_2 sporozoites in vitro and fixed at 48 hours post infection. The top two panels
show HC-04 cells infected with WT sporozoites, the middle two panels show HC- 04 cells
infected with PbAZMR_G95D_1 sporozoites and the bottom two panels show HC-04 cells
infected with PbAZMR_G95D_2 sporozoites. Azithromycin resistant liver stage schizonts only
show dispersed ACP (apicoplast) staining. Nuclei (blue) in azithromycin resistant liver stage
schizonts appear clustered and donut-shaped. Red staining indicates anti-Pb, marking the outside
of the liver stage parasite. All images are maximum projections and scale bars represent 10 μm;
F.) Quantification of parasite size ($μm^2$) at 48-hours post sporozoite inoculation of HC-04 cells *in
vitro* pooled from two independent experiments, with WT (n=60), PbAZMR_G95D_1 (n=31),
and PbAZMR_G95D_2 (n=47), where 'n'= number of individual liver stage schizonts measured.
PbAZMR_G95D_1 and PbAZMR_G95D_2 liver stages are significantly smaller than WT
(P<0.0001, Ordinary one-way ANOVA with Dunnett's multiple comparisons test; bars represent
the mean ±SD); G.) Pooled quantification of nuclei counts from the same liver stage schizonts
quantified in F. PbAZMR_G95D_1 and PbAZMR_G95D_2 liver stages have significantly fewer
nuclei than WT (P<0.0001, Ordinary one-way ANOVA with Dunnett's multiple comparisons
test; error bars represent the mean ±SD); H.) Nuclei counts, normalised to the size of each

individual liver stage parasite described in F (number of nuclei per μm^2). PbAZMR_G95D_1 and
PbAZMR_G95D_2 liver stages have significantly fewer nuclei than WT even when normalised
to liver stage parasite size (P<0.0001, Ordinary one-way ANOVA with Dunnett's multiple
comparisons test; error bars represent the mean ±SD).
590
Figure 5. Liver stage development of *P. falciparum* azithromycin resistant parasites is impaired.
A.) HC-04 cell traversal by Pf_WT and Pf_AZMR sporozoites at multiplicity of infection (MOI)
0.3, measured by FITC-Dextran uptake and counted by FACS. Cells were fixed at 3-hours post
inoculation and show no significant difference between Pf_WT and Pf_AZMR indicating normal
sporozoite motility in azithromycin resistant parasites (ns = not significant, Paired t-test two-
tailed P value); B.) HC-04 cell invasion by Pf_WT and Pf_AZMR sporozoites measured by
FACS after incubation for 18-hours using CSP-positive antibody staining of fixed permeabilized
cells. No significant difference was seen in invasion of HC-04 between Pf_WT and Pf_AZMR
sporozoites (ns = not significant, Paired t-test two-tailed P value); C.) Quantification of parasite
liver load in humanised mice infected by i.v. injection of 8 x 10^5 Pf_WT sporozoites (n=2 mice)
or Pf_AZMR sporozoites (n=3 mice) by qRT-PCR. Livers of infected humanised mice were
harvested at day 5 post sporozoite i.v. and tissue homogenised for subsequent loads analysis and
determination of the degree of chimerism. Pf_AZMR infected mice showed a significantly lower
liver stage parasite load than mice infected with Pf_WT sporozoites (Unpaired t-test, two-tailed
P value, P=0.0023, error bars represent mean ±SD); D.) Immunofluorescence assay of liver
sections from livers of humanised mice infected with either Pf_WT or Pf_AZMR sporozoites.
Liver stage schizonts were stained with anti-HSP70 (cytosolic marker) and anti-PTEX150
(parasitophorous vacuole marker) and DAPI DNA stain. Pf_AZMR liver stage schizonts show
little to no staining with either anti-HSP70 or anti-PTEX150 and show aberrant nuclear staining,
with individual nuclei unclear. Scale bar 10 μm; E.) Immunofluorescence assay of liver stage
schizonts and stained with anti-ACP (apicoplast marker), anti-CSP (parasite marker) and DAPI
DNA stain. Pf_AZMR livers stage schizonts show a loss of anti-ACP and anti-CSP stain
compared to WT, and again show aberrant nuclear staining, with DAPI appearing to aggregate
around holes. Scale bar 10 μm; F.) Quantification of liver stage schizont size (μm^2) from liver
sections from livers of humanised mice infected with either Pf_WT or Pf_AZMR sporozoites.
Sections from independent mice were selected and individual 'n' liver stage schizonts measured,
Pf_WT (n=10), Pf_AZMR (n=10). No significant difference in liver stage schizont size was seen
(ns = not significant, Unpaired t-test, two-tailed P value); G.) Quantification of nuclear centres
counted from liver stage schizonts imaged in D and E. Pf_AZMR liver stage schizonts show
significantly fewer nuclei than Pf_WT, indicating impaired liver stage development (Unpaired t-
test, two-tailed P value, P<0.0001; error bars represent mean ±SD).
622
623
624   **MATERIALS & METHODS**
625
Experimental animals and parasites
Male Swiss Webster mice, between 3 and 12 weeks old, were used in all experiments. Animals
were sourced from Monash Animal Research Platform. All animal experiments were in
accordance to the Prevention of Cruelty to Animals Act 1986 and the Prevention of Cruelty to
Animals Regulations 2008 and reviewed and were permitted by the Melbourne University
Animal Ethics Committee (Ethics IDs 1613928.1, 1714169, 1914889.1). Animals were

anaesthetised using ketamine/xylazine and euthanised via slow fill carbon dioxide followed by
cervical dislocation. *Plasmodium berghei* ANKA strain parasites were used in all experiments
both as a reference wild type (WT) line and to generate all subsequent drug resistant parasite
lines. *P. falciparum* NF54 strain parasites were used to generate all *P. falciparum* drug resistant
parasite lines. Adult female *Anopheles stephensi* mosquitoes (MR4) were used in all experiments
and reared under standard insectary conditions. Adult mosquitoes were grown at 27°C in 80%
humidity, with a light/dark cycle of 14:10 hours. Mosquito infections were performed in dark
conditions, and infected mosquitoes were maintained on 10% (v/v) sucrose.
640

Drugs
Azithromcyin was obtained from Sigma-Aldrich (Merck, NSW Australia), as pure
pharmaceutical secondary standards. Azithromycin was provided as a dihydrate (CAS Number:
117772-70-0) and administered once per day (q.d.) for 4 days as a suspension in PEG400 (MP
Biomedicals, Ohio, CAS Number 25322-68-3) via oral gavage (OG). Drug dilutions were made
fresh for each dosing experiment from identical drug batches as determined by LOT numbers.
647

Activity of drugs *in vivo*
To test for the curative dose and to confirm the efficacy and tolerance of each drug, an initial *in*
vivo drug dosing assay was performed. Briefly, a donor mouse was pre-infected with *P. berghei*
ANKA parasites via intraperitoneal (IP) injection of frozen stock. Once the donor mouse reached
5% parasitaemia, it was euthanised and infected blood collected via cardiac puncture. Infected
blood was diluted in RPMI 1640 medium (Gibco, Cat no. 61870-036) + 10% foetal bovine
serum (Gibco, Cat no. 10100147) to a concentration of $1x10^7$ infected red blood cells (iRBC) per
200 µL. Two cohorts of three mice (placebo/vehicle treated [n=3] and drug treated [n=3]) were
infected with $1x10^7$ iRBC via intravenous (IV) tail vein injection on day 0. Once mice reached
2% parasitaemia (approximately day 3), they were treated with the placebo/vehicle or drug by
OG or IP injection (on days 3-6) and monitored daily by Giemsa (10% [v/v]) stained thin blood
smears to determine parasite clearance. Mice were further monitored for health by daily weight
measurements.
661

To evaluate the reported schizontocidal activity, dosing range and dosing methods of
azithromycin, and to determine the 50% and 90% effective doses (ED50 and ED90 respectively)
a Peters' four-day suppressive test [68] was performed. Two donor mice were pre-infected with
WT *P. berghei* ANKA parasites via IP injection of fresh blood from a P0 infection. Once the
donor mice reached approximately 5% parasitaemia, they were euthanised and infected blood
collected via cardiac puncture. Infected blood from both mice was mixed and diluted in RPMI
1640 medium + 10% foetal bovine serum to a concentration of $1x10^7$ iRBC per 200 µL. Seven
cohorts of four mice were infected with $1x10^7$ iRBC via IV tail vein injection on day 0. All seven
cohorts were administered the vehicle (200 µL PEG400) as a control or a given concentration of
azithromycin (5 mg/kg, 15 mg/kg, 30 mg/kg, 45 mg/kg, 60 mg/kg, 70 mg/kg in 200 µL PEG400)
q.d. by OG on days 0-3, with the initial dose on day 0 given 4 hours post-infection (pi).
Parasitaemia was measured on day 4 by Giemsa-stained thin blood smears to determine
percentage suppression in comparison to the untreated control. Mice were also monitored daily
for health and to confirm parasites were being cleared. Percentage suppression (PS) of
parasitaemia was calculated using the formula: PS=100 x ([C_{AV}-T]/C_{AV}), where C_{AV}= average
parasitaemia in the control group, and T=parasitaemia of each treated replicate. Each PS

replicate was then plotted against the dose, and a non-linear regression curve was fitted using
GraphPad Prism 9 (Version 9.1.0) to determine the ED50 and ED90 doses of azithromycin.
680

Generating drug resistant *P. berghei* parasites
Selection of resistant *P. berghei* parasites *in vivo* was performed in the same way for each drug.
After inoculation via IV injection with $1x10^7$ iRBCs from a WT *P. berghei* ANKA infected
donor mouse, mice were treated with a clearance dose of the drug (60 mg/kg or 70 mg/kg
azithromycin) administered as described above. Mice were monitored daily by Giemsa-stained
thin blood smears, and once recrudescence was observed and parasitaemia reached
approximately 1-2%, mice were treated with another clearance dose of the drug. Parasites were
occasionally passaged into naïve mice should an immune response appear to have been
suppressing growth. This recrudescence→treatment→recrudescence→treatment cycle was
repeated until parasitaemia failed to reduce in the presence of the drug, at which point parasites
were deemed resistant and parasite lines were cloned by limiting dilution. Cloned lines were then
confirmed to be resistant to the clearance dose by performing an *in vivo* drug assay as described
above.
694

Generating drug resistant *P. falciparum* parasites
NF54 and 3D7 parasites were cloned by limiting dilution prior to resistance generation. Two
independent clones of NF54 and one of 3D7 were used for selection. Parasite cultures containing
$5x10^8$ parasites were treated continuously for 96 hours with 300 nM azithromycin or 72 hours
with 30 nM atovaquone, or 200 nM pyrimethamine. All three parasite lines were also grown
without drug as a sequencing control. Following treatment, cultures were allowed to recover in
drug free media until recrudescence (28 days post azithromycin treatment, 30-38 days post
atovaquone treatment, and 24-29 days post pyrimethamine treatment). Drug was reapplied to
recrudescent parasites at the initial concentration. For azithromycin resistance, parasites grew
normally at 300 nM azithromycin and continued to grow at concentrations up to 1.2 µM.
Atovaquone and pyrimethamine resistant parasites required from two to five cycles of treatment
and recovery before stable drug resistance at three times the initial concentrations was obtained.
Resistance parasites were cloned by limiting dilution, and drug resistance confirmed by standard
SYBRgreen drug assays. DNA was prepared from independent clones for whole genome
sequening.
710

Genomic DNA preparation and initial sequence analysis
Whole blood was collected via cardiac puncture from mice infected with clonal drug resistant *P.*
berghei parasite lines, and 10 µL aliquots were transferred into 190 µL dPBS for a total volume
of 200 µL. Genomic DNA was extracted using an Isolate II Genomic DNA Kit (Bioline,
Meridian Bioscience Australia), following the manufacturer's instructions. Known loci
potentially involved in resistance to azithromycin amplified by polymerase chain reaction (PCR)
for targeted sequencing (See Supplementary Table 1. for primer sequences and PCR conditions).
Correct sized amplicons were then purified using an Isolate II PCR and Gel Kit (Bioline,
Meridian Biosciences, Australia) following the manufacturer's instructions for PCR clean-up and
sent for Sanger sequencing at the Victorian Clinical Genetics Services (VCGS; Murdoch
Children's Research Institute, Royal Children's Hospital, Parkville, Victoria) following their
purified DNA criteria. Alignment and analysis of sequenced amplicons was done using
Sequencher (Gene Codes Corporation, USA).

Sample preparation and genomic DNA extraction for Illumina sequencing

To obtain a concentrated, pure parasite sample, free of contaminant mouse leucocytes, platelets and other material, we performed a *P. berghei* schizont preparation. Whole blood was collected via cardiac puncture from mice infected with the clonal drug resistant *P. berghei* parasite lines, and placed into a 50 mL overnight culture in complete schizont culture medium (70% [v/v] RPMI 1640 medium, 30% [v/v] FBS and gentamycin (Gibco) $4x10^{-2}$ mg/mL). Approximately 20 hours post culture, schizonts were purified using a CS column (Miltenyi Biotec, Germany) assembled in a VarioMACS magnetic cell separator (Miltenyi Biotec, Germany) and eluted in RPMI 1640 medium. Purified schizonts were spun down in a swing-bucket centrifuge at 1500 RPM for 10 minutes, resuspended in 1 mL of dPBS, centrifuged again at 1500 RPM for 10 minutes and resuspended in 1 mL dPBS. Each 1 mL washed schizont preparation was then distributed into five 200 µL aliquots and genomic DNA was extracted immediately using a QIAmp DNA Blood Mini Kit (Qiagen) following the manufacturer's instructions with one exception. After lysis at 56°C, preparations were centrifuged at maximum (13000 RPM) and supernatant was placed into a new 1.5 mL tube to remove digested haemoglobin. After this step, the protocol was followed verbatim. Samples were sent to the VCGS (Murdoch Children's Research Institute, Royal Children's Hospital, Parkville, Victoria) for Illumina Nextera XT library preparation and whole genome sequencing on either an Illumina HiSeq4000 or NovaSeq2 (150 bp paired-end reads).

Bioinformatic analysis

Analyses were performed using the High Performance Computing Services at The University of Melbourne (Parkville, Victoria) or personal computers. Raw FASTQ files were mapped to the *P. berghei* ANKA or *P. falciparum* 3D7 reference genome (Version 43 (*P. berghei*), Version 44 (*P. falciparum*), PlasmoDB [69]) using the Burrows-Wheeler Aligner [70] and the quality checked with FastQC [71] and QualiMap [72]. Mapped reads were visualised using Integrated Genomics Viewer (IGV, version 2.8.4 [73]). Duplicate reads were marked using Picard tools (version 2.18.27) MarkDuplicates tool. Variant calling was performed following the Genome Analysis Toolkit (GATK)[74] best practices pipeline [75], using the HaplotypeCaller (in GVCF mode) and GenotypeGVCFs tools. Raw variants were then annotated using SnpEff [76] and filtered using SnpSift [77] based on quality scores, read depth and genome position (i.e. intergenic versus coding regions). Further manual filtering and analysis was performed using GNU/Linux coreutils and Awk programming language [78] based on biological relevance.

Infection of mosquitoes with *P. berghei*

Female *A. stephensi* mosquitoes were infected with clonal *P. berghei* drug resistant lines and WT *P. berghei* ANKA to investigate the fitness, phenotype and transmissibility of drug resistant parasites. Approximately 200 µL frozen blood stock was used to infect a donor mouse by IP injection. Once parasitaemia reached between 5-8% parasite readiness for infection was confirmed by checking for exflagellation of microgametes by microscopy. Donor mice were then anaesthetised, and individually placed on an infection cage containing 100 female *A. stephensi* mosquitoes, which were allowed to feed for up to 40 minutes in a 20°C incubator at 80% humidity. The donor mouse was euthanised post feed. Infected mosquitoes were maintained on 10% (v/v) sucrose for up to 28 days, or when the experiment was ended.

Oocyst quantification and immunofluorescence assays

Whole midguts were dissected from infected mosquitoes between 9- and 14-days post blood feed and either stained with 0.2% (v/v) mercurochrome to assess oocyst numbers, or prepared for immunofluorescence assay. In 24-well plates, whole midguts were dissected in dPBS and then fixed in 4% (v/v) paraformaldehyde in dPBS for 1 hour at room temperature. After washing in 1 mL fresh dPBS three times for 5 minutes on an orbital shaker, midguts were blocked and permeabilised overnight at 4°C in midgut blocking buffer (0.25% [v/v] Triton X-100, 5% [w/v] BSA [Sigma-Aldrich, Cat no. A4503-50G] in dPBS). Midguts were incubated with primary antibodies (apicoplast marker acyl carrier protein [ACP] anti-serum [79] at 1:250) diluted in midgut blocking buffer for a minimum of 4 hours at room temperature on an orbital shaker. After washing three times in 1 mL dPBS, midguts were incubated at room temperature for 2 hours on an orbital shaker with secondary antibodies (goat anti-rabbit IgG Alexa Fluor 488 [Abcam] at 1:4000) diluted in midgut blocking buffer. Midguts were then washed three times for 5 minutes in 1 mL dPBS on an orbital shaker, with the second wash containing Hoechst 33342 diluted to 1:10 000 and mounted on glass slides in Dako fluorescence mounting medium. Immunofluorescence assays were imaged on a Nikon C2 confocal microscope and images assembled in ImageJ Fiji (Version 2.1.0/1.53c) [80].

Quantitative PCR for organellar genome detection

Primers designed to target genes encoded in the nuclear, apicoplast and mitochondrial genomes (See Supplementary Table 1. for primer sequences and qPCR conditions) were used to perform quantitative PCR (qPCR) on oocyst stage parasites isolated as described above. All primers were first tested to ensure similar amplification efficiency under the assay conditions by generating standard curves (Supplementary Figure 1.). Genomic DNA was extracted from parasite lines using an Isolate II Genomic DNA Kit (Bioline, Meridian Bioscience Australia), following the manufacturer's instructions, with one exception. Whole midguts were first dissected into 180 µL lysis buffer GL (Bioline) and 20 µL Proteinase K and incubated at 56°C for 1 hour to ensure efficient lysis of oocysts, the original protocol was then followed verbatim. All qPCR assays were performed on a CFX Connect Real-Time PCR Detection System (Bio-Rad). Genome copy number was calculated using the $\Delta\Delta C_T$ method [81], first normalising mean C_T values of apicoplast and mitochondrial genes within samples to the mean C_T of the nuclear genes, then using the WT *P. berghei* ANKA samples as a calibrator for the drug resistant lines.

Sporozoite quantification and immunofluorescence motility assays

Eight-well chamber slides (Merck Millipore Cat no. PEZGS0816) were coated with either mouse-derived monoclonal circumsporozoite protein (CSP) anti-serum (1:1000) [82] or mouse-derived polyclonal *P. berghei* anti-serum (1:1000; made in our laboratory) diluted in dPBS, and incubated at 37°C for 1 hour. Salivary glands were dissected in 1640 medium + 10% foetal bovine serum from infected mosquitoes between 20- and 24-days post blood feed, and purified sporozoites were counted. After removing the coating antibody solution from the chamber slide, up to $2x10^4$ sporozoites from each drug resistant *P. berghei* parasite line or WT *P. berghei* ANKA were added to individual wells and allowed to settle and glide for 1 hour at 37°C. The supernatant was then removed and parasites fixed with 150 µL of 4% (v/v) paraformaldehyde in dPBS for 20 minutes at room temperature. After washing gently with dPBS three times for 5 minutes, blocking buffer (0.2% [v/v] Triton X-100, 3% [w/v] BSA in dPBS) was added to each well and left to incubate for 20 minutes at room temperature. Primary antibodies were then added

to each well (mouse anti-CSP [1:500], mouse anti-*P. berghei* [1:500] or rabbit anti-ACP [1:250]
as described above) diluted in 3% (w/v) BSA in dPBS. Wells were again washed three times for
5 minutes with dPBS before incubating with secondary antibodies (goat anti-rabbit IgG Alexa
Fluor 488 [Abcam]; goat anti-mouse IgG Alexa Fluor 568 [Abcam]) diluted in 3% (w/v) BSA
(all 1:2000). Wells were then washed three times for 5 minutes in dPBS, with the second wash
containing Hoechst 33342 diluted to 1:10000, and mounted with Dako fluorescence mounting
medium. Sporozoites were then imaged on a Leica DM6000B compound fluorescence
microscope and images assembled in ImageJ Fiji (Version 2.1.0/1.53c) [80].
824
Mosquito infection and analysis of parasite development
Seven-day old female *Anopheles stephensi* mosquitoes (originally provided by M. Jacobs-
Lorena, John Hopkins University) were allowed to feed on asynchronous gametocytes at 0.6%
stage V gametocytemia using water jacketed glass membrane feeders. Mosquitos were then
sugar starved for 48 hours to kill any that had not fed or males. Those that survived were given
di-ionised water through cotton wicks and sugar cubes. For oocyst quantification; midguts were
dissected from ethanol killed mosquitos 7 days post feed and stained with 0.1% mercurochrome.
Salivary glands were dissected for mosquitos 16 days post feed, crushed with a pestle, and then
filtered through glass wool. Sporozoites were counted using a hemocytometer.
834
Infection of naïve mice with *P. berghei* sporozoites and analysis of infectivity
Naïve mice were infected with drug resistant or WT *P. berghei* ANKA sporozoites either by IV
tail vein injection or by direct blood feeding. For IV infection, salivary glands were dissected
from infected mosquitoes between days 20 and 24 post blood feed directly into RPMI 1640
medium + 10% (v/v) FBS. Sporozoites were then purified and counted prior to injecting either
$1x10^3$ or $1x10^4$ sporozoites per mouse. For direct blood feed, infected mosquitoes were allowed
to bite anaesthetised naïve mice. A total of 15 mosquitoes were allowed to blood feed for 15
minutes or until fed. Parasitaemia of mice was monitored by Giemsa-stained thin blood smears
from day 2 post IV or mosquito bite to assess blood stage patency.
844
Liver stage development analysis *in vitro*
To investigate the development of parasites through the liver stage, human hepatoma HepG2
(ATCC No. HB-8065) or HC-04 (ATCC MRA-975) cell lines were cultured at 37°C and 5%
CO_2 in a T-25 culture flask (Corning) in HepG2 medium (Advanced MEM minus L-Glutamine
[Gibco] + 10% [v/v] HI-FBS [Gibco] + 1% [v/v] penicillin-streptomycin + 2 mM Glutamax
[Gibco] and 0.1% [v/v] amphotericin B). Once the HepG2 or HC-04 cells were grown to the
desired concentration, $1x10^5$ to $2x10^5$ cells were seeded onto coverslips in a 24-well plate in 1
mL HepG2 medium per well. The following day, sporozoites were dissected directly into HepG2
medium, purified and counted and up to $2x10^4$ sporozoites were added into each well and
allowed to settle and invade for 2 hours at 37°C and 5% CO_2. Media was changed after the 2-
hour invasion process, and then twice daily thereafter until cells were fixed. Infected HepG2 or
HC-04 cells were fixed on the coverslips for 20 minutes with 4% (v/v) paraformaldehyde in
dPBS, at either 24- or 48-hours post sporozoite infection. Coverslips were washed with dPBS
three times for 5 minutes in the 24-well plate before adding 1 mL ice-cold methanol for storage
(no more than 3 weeks) at 20°C. After removing the methanol, coverslips were washed three
times for 5 minutes with dPBS before cells were permeabilised with 0.1% (v/v) Triton X-100 in
dPBS for 20 minutes at room temperature. Washing steps were repeated, and cells were then

blocked for 30 minutes at room temperature with 3% (w/v) BSA in dPBS. Coverslips were
incubated with primary antibodies (mouse anti-*P. berghei* [1:500] and rabbit anti-ACP [1:250])
and secondary antibodies (goat anti-rabbit IgG Alexa Fluor 488 [Abcam]; goat anti-mouse IgG
Alexa Fluor 568 [Abcam]) in 3% BSA in dPBS for one hour at room temperature, washing three
times for 5 minutes with dPBS in between each incubation. Following the probing with
secondary antibodies, coverslips were washed three times for 5 minutes in dPBS, with the second
wash containing Hoechst 33342 diluted to 1:10000 and mounted with Dako fluorescence
mounting medium on glass slides and sealed with clear nail polish. Infected HepG2 and HC-04
liver cells were imaged on a Nikon C2 confocal microscope, and images assembled and
measured in ImageJ Fiji (Version 2.1.0/1.53c) [80].
872
Humanised mice processing
FRG humanised mice were obtained from Yecuris. An inoculum of 8×10^5 *P. falciparum* control
NF54 sporozoites and 8×10^5 Azithromycin resistant sporozoites freshly isolated from mosquito
salivary glands were injected into each humanised mouse (2 mice infected with NF54 and 3 mice
with Azithromycin resistant lines).
Livers were obtained 5.5 days post infection from CO_2-euthanized mice, and individual lobes
were cut as described [83]. Half of the lobes were embedded in OCT and frozen in liquid nitrogen
for IFA analysis, while half was emulsified into a single-cell suspension and frozen and -80 for
subsequent gDNA extraction.
For quantification of parasite load in the chimeric livers of each mouse; gDNA was isolated from
the single-cell liver suspensions, and Taqman probe-based quantitative PCRs were performed as
previously described [83-86] using primers that would specifically amplify the 18S gene in each
parasite line 18S_fwd (5' -GTAATTGGAATGATAGGAATTTACAGGT-3') and 18S_rev (5'-
TCAACTACGAACGTTTTAACTGCAAC-3'). Human and mouse genomes were quantified
using oligonucleotides specific for prostaglandin E receptor 2 (PTGER2) from each species, as
described [84]. All probes were labelled 5' with the fluorophore 6-carboxy-fluorescein (FAM) and
contain a double quencher that includes and internal ZEN quencher and a 3' Iowa Black
quencher from IDT.

The following probes were used:

18S: 5'FAM TGCCAGCAG/ZEN/CCGCGGTA-3IABkFQ

hPTGER2: FAM/TGCTGCTTC/ZEN/TCATTGTCTCG/3IABkFQ,

mPTGER2: FAM/CCTGCTGCT/ZEN/TATCGTGGCTG/3IABkFQ
895
896   Standard curves were prepared by titration of from a defined number of DNA copies for *P.*
897   *falciparum* NF54, human and mouse controls. PCRs were performed on a Roche LC80 using
898   LightCycler 480 Probes Master (Roche).
899
Immunofluorescence assay
Purified NF54 and azithromycin resistant sporozoites were fixed with 4% paraformaldehyde
(PFA)/ 0.0075% glutaraldehyde in PBS at room temperature for 20 minutes. Sporozoites were

permeabilised in 0.1% Triton X 100 for 10 minutes before being blocked in 3% BSA PBS for 1
hour at room temperature. Samples were incubated with primary antibodies; rabbit anti-ACP
antibody (1:500), and mouse anti-CSP 2A10 (1:1000), for 1 hour in 3% BSA PBS followed by
secondary antibodies; anti-mouse 488 and anti-rabbit 594 (1:1000, Invitrogen). Images of NF54
or azithromycin sporozoites were taken at random based on CSP and DAPI and presence or
absence of apicoplast was then determined by staining with ACP.

For liver stage immunofluorescence analysis; livers were perfused with 1x PBS and fixed in 4%
PFA PBS overnight before being exchanged for sucrose and snap frozen in OCT (Invitrogen).
Sections of 8 mm were cut on the cryostat apparatus HMM550. Samples were permeabilised
using ice cold 10% methanol 90% acetone and blocked using 3% BSA PBS. Sections were then
incubated with primary antibodies; mouse anti-CSP (2A10; 1:1000), mouse anti-HSP70 (1:500),
rabbit anti-PTEX150 (1:500), mouse anti-EXP2 (1:500), rabbit anti-ACP (1:500) diluted in 3%
BSA PBS. Secondary antibodies were goat anti-rabbit Alexa 594 and anti-mouse Alexa 488
(1:1000; Invitrogen). All samples were then incubated with DAPI (4',6'- diamidino-2-
phenylindole) at 1 ug/ml in PBS to visualize DNA and mounted in Prolong Gold mounting
media (Invitrogen). Images were acquired on the Zeiss LSM 980 microscope.

Z stacks of both NF54 and azithromycin resistant liver stages were acquired using the same
settings to allow quantitative analysis between different samples. The number of nuclear centres
were counted by eye based on DAPI signal at the centre of each Z stack. For analysis of parasite
size; a region of interest was drawn around each parasite based on DAPI signal and then the
membrane area was calculated using FIJI software.

Traversal assay

Cell traversal was measured using a cell wounding assay [87]. HC-04 hepatocytes ($1x10^5$) were
seeded onto the bottom of a 96-well plate using IMDM (Life Technologies, 11360-070).
Sporozoites were added at an MOI of 0.3 and left to traverse hepatocytes for 2.5 hours in the
presence of 1 mg ml-1 FITC-labelled dextran (10,000 MW, Sigma Aldrich). Cells were
trypsinized to obtain a single cell suspension for FACS analysis. For each condition, triplicate
samples of 10,000 cells were counted by flow cytometry in each of the three independent
experiments.

Invasion assay

Assay was performed as previously described [87]. Briefly, HC-04 hepatocytes ($1x10^5$) were
seeded onto the bottom of a 96-well plate using IMDM (Life Technologies, 11360-070).
Sporozoites were added at an MOI of 0.3 and left to invade hepatocytes for either 5 or 18 hours
before being trypsinized to obtain a single cell suspension and transferred to a 96 well round
bottom plate. Cells were fixed and permeabilised (BD bioscience) and stained with Alexa fluor
647 conjugated mouse anti-CSP antibody in 3% BSA permeabilizing solution. Cells were then
washed in PBS and analysed by Flow cytometry. For each condition, triplicate samples of 10,000
cells were counted in each of the three independent experiments.

SUPPLEMENTAL FIGURES

Figure S1. Generation of azithromycin resistant *P. berghei* parasites *in vivo*. PbAZMR_G95D_1 and PbAZMR_G95D_2 were both generated off a 70 mg/kg q.d. dosing regimen, and PbAZMR_S89L was generated off a 60 mg/kg q.d. dosing regimen. The initial four-day dose is represented by the shaded rectangle, and the start of each dose thereafter is indicated by a shaded box to avoid rectangle overlaps. Parasites were deemed resistant when they failed to reduce in parasitaemia on the third day post treatment start (with the exception of PbAZMR_S89L, which grew slowly so was monitored differently).

Figure S2. Blood stage growth of *P. berghei* azithromycin parasites is impaired compared to WT without drug pressure. Growth assay of PbAZMR_G95D_1 (n=3) and PbAZMR_S89L (n=3) compared to WT (n=3). Parasitaemia was monitored by Giemsa-stained thin blood smears. Dot points represent mean parasitaemia, and error bars indicate ±SD. Growth of both azithromycin resistant parasite lines is delayed compared to WT, and eventually fails to increase.

REFERENCES

1.   White, N. J., Can new treatment developments combat resistance in malaria? *Expert Opinion on Pharmacotherapy* **2016,** *17* (10), 1303-1307.
2.   Ashley, E. A.; Dhorda, M.; Fairhurst, R. M.; Amaratunga, C.; Lim, P.; Suon, S.; Sreng, S.; Anderson, J. M.; Mao, S.; Sam, B., et al., Spread of Artemisinin Resistance in Plasmodium falciparum Malaria. *New England Journal of Medicine* **2014,** *371* (5), 411-423.
3.   Dondorp, A. M.; Fairhurst, R. M.; Slutsker, L.; MacArthur, J. R.; M.D., J. G. B.; Guerin, P. J.; Wellems, T. E.; Ringwald, P.; Newman, R. D.; Plowe, C. V., The Threat of Artemisinin-Resistant Malaria. *New England Journal of Medicine* **2011,** *365* (12), 1073-1075.
4.   Imwong, M.; Hien, T. T.; Thuy-Nhien, N. T.; Dondorp, A. M.; White, N. J., Spread of a single multidrug resistant malaria parasite lineage (PfPailin) to Vietnam. *The Lancet Infectious Diseases* **2017,** *17* (10), 1022-1023.
5.   Das, S.; Saha, B.; Hati, A. K.; Roy, S., Evidence of Artemisinin-Resistant Plasmodium falciparum Malaria in Eastern India. *New England Journal of Medicine* **2018,** *379* (20), 1962-1964.
6.   Balikagala, B.; Fukuda, N.; Ikeda, M.; Katuro, O. T.; Tachibana, S. I.; Yamauchi, M.; Opio, W.; Emoto, S.; Anywar, D. A.; Kimura, E., et al., Evidence of Artemisinin-Resistant Malaria in Africa. *N Engl J Med* **2021,** *385* (13), 1163-1171.
7.   Conrad, M. D.; Asua, V.; Garg, S.; Giesbrecht, D.; Niaré, K.; Smith, S.; Namuganga, J. F.; Katairo, T.; Legac, J.; Crudale, R. M., et al., Evolution of Partial Resistance to Artemisinins in Malaria Parasites in Uganda. *N Engl J Med* **2023,** *389* (8), 722-732.
8.   Fola, A. A.; Feleke, S. M.; Mohammed, H.; Brhane, B. G.; Hennelly, C. M.; Assefa, A.; Crudal, R. M.; Reichert, E.; Juliano, J. J.; Cunningham, J., et al., Plasmodium falciparum resistant to artemisinin and diagnostics have emerged in Ethiopia. *Nat Microbiol* **2023,** *8* (10), 1911-1919.
9.   van der Pluijm, R. W.; Amaratunga, C.; Dhorda, M.; Dondorp, A. M., Triple Artemisinin-Based Combination Therapies for Malaria – A New Paradigm? *Trends in Parasitology* **2021,** *37* (1), 15-24.
10.   Koehne, E.; Adegnika, A. A.; Held, J.; Kreidenweiss, A., Pharmacotherapy for artemisinin-resistant malaria. *Expert Opinion on Pharmacotherapy* **2021,** *22* (18), 2483-2493.
11.   Ménard, D.; Fidock, D. A., Accelerated evolution and spread of multidrug-resistant Plasmodium falciparum takes down the latest first-line antimalarial drug in southeast Asia. *The Lancet Infectious Diseases* **2019,** *19* (9), 916-917.
12.   Balta, V. A.; Stiffler, D.; Sayeed, A.; Tripathi, A. K.; Elahi, R.; Mlambo, G.; Bakshi, R. P.; Dziedzic, A. G.; Jedlicka, A. E.; Nenortas, E., et al., Transmissibility of clinically relevant atovaquone-resistant Plasmodium falciparum by anopheline mosquitoes. *bioRxiv* **2023**, 2023.02.07.527535.
13.   Goodman, C. D.; Siregar, J. E.; Mollard, V.; Vega-Rodriguez, J.; Syafruddin, D.; Matsuoka, H.; Matsuzaki, M.; Toyama, T.; Sturm, A.; Cozijnsen, A., et al., Parasites resistant to the antimalarial atovaquone fail to transmit by mosquitoes. *Science (New York, N.Y.)* **2016,** *352* (6283), 349-53.
14.   Goodman, C. D.; Buchanan, H. D.; McFadden, G. I., Is the mitochondrion a good malaria drug target? *Trends in parasitology* **2017,** *33* (3), 185-193.
15.   van Schaijk, B. C. L.; Santha Kumar, T. R.; Vos, M. W.; Richman, A.; van Gemert, G.-J.; Eappen, A. G.; Williamson, K. C.; Morahan, B. J.; Fishbaugher, M.; Kennedy, M., et

al., Type II fatty acid biosynthesis is essential for *Plasmodium falciparum* sporozoite development in the midgut of *Anopheles* mosquitoes. *Eukaryotic Cell* **2014,** *13* (5), 550-559.

16.   Biddau, M.; Santha Kumar, T. R.; Henrich, P.; Laine, L. M.; Blackburn, G. J.; Chokkathukalam, A.; Li, T.; Lee Sim, K.; King, L.; Hoffman, S. L., et al., *Plasmodium falciparum* LipB mutants display altered redox and carbon metabolism in asexual stages and cannot complete sporogony in *Anopheles* mosquitoes. *International Journal for Parasitology* **2021,** *51* (6), 441-453.

17.   Charan, M.; Choudhary, H. H.; Singh, N.; Sadik, M.; Siddiqi, M. I.; Mishra, S.; Habib, S., [Fe-S] cluster assembly in the apicoplast and its indispensability in mosquito stages of the malaria parasite. *The FEBS Journal* **2017,** *284* (16), 2629-2648.

18.   Uddin, T.; McFadden Geoffrey, I.; Goodman Christopher, D., Validation of Putative Apicoplast-Targeting Drugs Using a Chemical Supplementation Assay in Cultured Human Malaria Parasites. *Antimicrobial Agents and Chemotherapy* **2018,** *62* (1), e01161-17.

19.   Goodman, C. D.; Su, V.; McFadden, G. I., The effects of anti-bacterials on the malaria parasite *Plasmodium falciparum. Molecular and Biochemical Parasitology* **2007,** *152* (2), 181-191.

20.   Dahl, E. L.; Rosenthal, P. J., Multiple Antibiotics Exert Delayed Effects against the *Plasmodium falciparum* Apicoplast. *Antimicrobial Agents and Chemotherapy* **2007,** *51* (10), 3485-3490.

21.   McFadden, G. I.; Reith, M. E.; Munholland, J.; Lang-Unnasch, N., Plastid in human parasites. *Nature* **1996,** *381*, 482.

22.   Wilson, R. J.; Denny, P. W.; Preiser, P. R.; Rangachari, K.; Roberts, K.; Roy, A.; Whyte, A.; Strath, M.; Moore, D. J.; Moore, P. W., et al., Complete gene map of the plastid-like DNA of the malaria parasite Plasmodium falciparum. *Journal of Molecular Biology* **1996,** *261* (2), 155-172.

23.   van Dooren, G. G.; Striepen, B., The Algal Past and Parasite Present of the Apicoplast. *Annual Review of Microbiology* **2013,** *67* (1), 271-289.

24.   Botté, C. Y.; Dubar, F.; McFadden, G. I.; Maréchal, E.; Biot, C., Plasmodium falciparum Apicoplast Drugs: Targets or Off-Targets? *Chemical Reviews* **2012,** *112* (3), 1269-1283.

25.   Goodman, C. D.; McFadden, G. I., Targeting apicoplasts in malaria parasites. *Expert Opinion on Therapeutic Targets* **2013,** *17* (2), 167-177.

26.   Dahl, E. L.; Rosenthal, P. J., Apicoplast translation, transcription and genome replication: targets for antimalarial antibiotics. *Trends in parasitology* **2008,** *24* (6), 279-284.

27.   Chandra, R.; Ansah, P.; Sagara, I.; Sie, A.; Tiono, A. B.; Djimde, A. A.; Zhao, Q.; Robbins, J.; Penali, L. K.; Ogutu, B., Comparison of azithromycin plus chloroquine versus artemether-lumefantrine for the treatment of uncomplicated *Plasmodium falciparum* malaria in children in Africa: a randomized, open-label study. *Malaria Journal* **2015,** *14* (1), 1-10.

28.   Sagara, I.; Oduro, A. R.; Mulenga, M.; Dieng, Y.; Ogutu, B.; Tiono, A. B.; Mugyenyi, P.; Sie, A.; Wasunna, M.; Kain, K. C., Efficacy and safety of a combination of azithromycin and chloroquine for the treatment of uncomplicated *Plasmodium falciparum* malaria in two multi-country randomised clinical trials in African adults. *Malaria Journal* **2014,** *13*, 1-10.

29.   van Eijk, A. M.; Terlouw, D. J., Azithromycin for treating uncomplicated malaria. *Cochrane database of systematic reviews* **2011,** (2).

30.   Sidhu, A. B. S.; Sun, Q.; Nkrumah, L. J.; Dunne, M. W.; Sacchettini, J. C.; Fidock, D.
A., In vitro efficacy, resistance selection, and structural modeling studies implicate the malarial
parasite apicoplast as the target of azithromycin. *Journal of Biological Chemistry* **2007,** *282* (4),
2494-2504.
31.   Kennedy, K.; Cobbold, S. A.; Hanssen, E.; Birnbaum, J.; Spillman, N. J.; McHugh, E.;
Brown, H.; Tilley, L.; Spielmann, T.; McConville, M. J., et al., Delayed death in the malaria
parasite *Plasmodium falciparum* is caused by disruption of prenylation-dependent intracellular
trafficking. *PLoS Biol* **2019,** *17* (7), e3000376-28.
32.   Rosenthal, P. J., Azithromycin for Malaria? *The American journal of tropical medicine
and hygiene* **2016**, 16-0332.
33.   Wilson, D. W.; Goodman, C. D.; Sleebs, B. E.; Weiss, G. E.; de Jong, N. W. M.;
Angrisano, F.; Langer, C.; Baum, J.; Crabb, B. S.; Gilson, P. R., et al., Macrolides rapidly
inhibit red blood cell invasion by the human malaria parasite, *Plasmodium falciparum. BMC
Biology* **2015,** *13* (1), 1.
34.   Burns, A. L.; Sleebs, B. E.; Siddiqui, G.; De Paoli, A. E.; Anderson, D.; Liffner, B.;
Harvey, R.; Beeson, J. G.; Creek, D. J.; Goodman, C. D., et al., Retargeting azithromycin
analogues to have dual-modality antimalarial activity. *BMC Biology* **2020,** *18* (1), 133.
35.   McFadden, G. I.; Yeh, E., The apicoplast: now you see it, now you don't. *International
Journal for Parasitology* **2017,** *47* (2), 137-144.
36.   Buchanan, H. D.; Goodman, C. D.; McFadden, G. I., Roles of the apicoplast across the
life cycles of rodent and human malaria parasites. *Journal of Eukaryotic Microbiology* **2022,** *69*
(6), e12947.
37.   Yeh, E.; DeRisi, J. L., Chemical Rescue of Malaria Parasites Lacking an Apicoplast
Defines Organelle Function in Blood-Stage *Plasmodium falciparum. PLoS Biol* **2011,** *9* (8),
e1001138.
38.   Swift, R. P.; Rajaram, K.; Liu, H. B.; Prigge, S. T., Dephospho-CoA kinase, a nuclear-
encoded apicoplast protein, remains active and essential after *Plasmodium falciparum* apicoplast
disruption. *EMBO J* **2021,** *40*, e107247.
39.   Ke, H.; Sigala, P. A.; Miura, K.; Morrisey, J. M.; Mather, M. W.; Crowley, J. R.;
Henderson, J. P.; Goldberg, D. E.; Long, C. A.; Vaidya, A. B., The heme biosynthesis pathway
is essential for *Plasmodium falciparum* development in mosquito stage but not in blood stages. *J
Biol Chem* **2014,** *289* (50), 34827-37.
40.   Nagaraj, V. A.; Sundaram, B.; Varadarajan, N. M.; Subramani, P. A.; Kalappa, D. M.;
Ghosh, S. K.; Padmanaban, G., Malaria Parasite-Synthesized Heme Is Essential in the Mosquito
and Liver Stages and Complements Host Heme in the Blood Stages of Infection. *PLOS
Pathogens* **2013,** *9* (8), e1003522.
41.   Rathnapala, U. L.; Goodman, C. D.; McFadden, G. I., A novel genetic technique in
Plasmodium berghei allows liver stage analysis of genes required for mosquito stage
development and demonstrates that de novo heme synthesis is essential for liver stage
development in the malaria parasite. *PLoS Pathogens* **2017,** *13* (6), e1006396-18.
42.   Jacot, D.; MacPherson, D. A., Apicomplexan Energy Metabolism: Carbon Source
Promiscuity and the Quiescence Hyperbole. *Trends in parasitology* **2016,** *32* (1), 56-70.
43.   Goodman, C. D.; Useglio, M.; Peiru, S.; Labadie, G. R.; McFadden, G. I.; Rodriguez,
E.; Gramajo, H., Chemobiosynthesis of New Antimalarial Macrolides. *Antimicrobial Agents and
Chemotherapy* **2013,** *57* (2), 907-913.

44. Waller, R. F.; Keeling, P. J.; Donald, R. G. K.; Striepen, B.; Handman, E.; Lang-
Unnasch, N.; Cowman, A. F.; Besra, G. S.; Roos, D. S.; McFadden, G. I., Nuclear-encoded
proteins target to the plastid in *Toxoplasma gondii* and *Plasmodium falciparum*. *Proceedings of*
the National Academy of Sciences **1998,** *95* (21), 12352-12357.
45. Waller, R. F.; Reed, M. B.; Cowman, A. F.; McFadden, G. I., Protein trafficking to the
plastid of *Plasmodium falciparum* is via the secretory pathway. *The EMBO Journal* **2000,** *19* (8),
1794-1802.
46. Ejigiri, I.; Ragheb, D. R. T.; Pino, P.; Coppi, A.; Bennett, B. L.; Soldati-Favre, D.;
Sinnis, P., Shedding of TRAP by a Rhomboid Protease from the Malaria Sporozoite Surface Is
Essential for Gliding Motility and Sporozoite Infectivity. *PLOS Pathogens* **2012,** *8* (7),
e1002725.
47. Hopp, C. S.; Chiou, K.; Ragheb, D. R. T.; Salman, A. M.; Khan, S. M.; Liu, A. J.;
Sinnis, P., Longitudinal analysis of *Plasmodium* sporozoite motility in the dermis reveals
component of blood vessel recognition. *eLife* **2015,** *4*, e07789.
48. Yu, M.; Kumar, T. R. S.; Nkrumah, L. J.; Coppi, A.; Retzlaff, S.; Li, C. D.; Kelly, B.
J.; Moura, P. A.; Lakshmanan, V.; Freundlich, J. S., et al., The fatty acid biosynthesis enzyme
FabI plays a key role in the development of liver-stage malarial parasites. *Cell Host & Microbe*
2008, *4* (6), 567-78.
49. Vaughan, A. M.; O'Neill, M. T.; Tarun, A. S.; Camargo, N.; Phuong, T. M.; Aly, A. S.
I.; Cowman, A. F.; Kappe, S. H. I., Type II fatty acid synthesis is essential only for malaria
parasite late liver stage development. *Cellular Microbiology* **2009,** *11* (3), 506-20.
50. Falkard, B.; Kumar, T. R. S.; Hecht, L.-S.; Matthews, K. A.; Henrich, P. P.; Gulati, S.;
Lewis, R. E.; Manary, M. J.; Winzeler, E. A.; Sinnis, P., et al., A key role for lipoic acid
synthesis during *Plasmodium* liver stage development. *Cellular Microbiology* **2013,** *15* (9),
1585-1604.
51. Stanway, R. R.; Bushell, E.; Chiappino-Pepe, A.; Roques, M.; Sanderson, T.; Franke-
Fayard, B.; Caldelari, R.; Golomingi, M.; Nyonda, M.; Pandey, V., et al., Genome-Scale
Identification of Essential Metabolic Processes for Targeting the *Plasmodium* Liver Stage. *Cell*
2019, *179* (5), 1112-1128.e26.
52. Stanway, R. R.; Witt, T.; Zobiak, B.; Aepfelbacher, M.; Heussler, V. T.,
GFP☐targeting allows visualization of the apicoplast throughout the life cycle of live malaria
parasites. *Biology of the Cell* **2009,** *101* (7), 415-435.
53. Marcos, L. A.; Wormser, G. P., Relapsing Babesiosis With Molecular Evidence of
Resistance to Certain Antimicrobials Commonly Used to Treat *Babesia microti* Infections. *Open*
Forum Infect Dis **2023,** *10* (8).
54. Bowers, A. K.; Keller, J. A.; Dutcher, S. K., Molecular markers for rapidly identifying
candidate genes in *Chlamydomonas reinhardtii*: ERY1 and ERY2 encode chloroplast ribosomal
proteins. *Genetics* **2003,** *164* (4), 1345-1353.
55. Canu, A.; Malbruny, B.; Coquemont, M.; Davies, T. A.; Appelbaum, P. C.; Leclercq,
R., Diversity of ribosomal mutations conferring resistance to macrolides, clindamycin,
streptogramin, and telithromycin in *Streptococcus pneumoniae*. *Antimicrobial Agents and*
Chemotherapy **2002,** *46* (1), 125-131.
56. Ma, K. C.; Mortimer, T. D.; Duckett, M. A.; Hicks, A. L.; Wheeler, N. E.; Sánchez-
Busó, L.; Grad, Y. H., Increased power from conditional bacterial genome-wide association
identifies macrolide resistance mutations in *Neisseria gonorrhoeae*. *Nature Communications*
2020, *11* (1), 5374.

57.   Wilson, R. J.; Fry, M.; Gardner, M. J.; Feagin, J. E.; Williamson, D. H., Subcellular fractionation of the two organelle DNAs of malaria parasites. *Current Genetics* **1992,** *21*, 405-408.

58.   Siregar, J. E.; Syafruddin, D.; Matsuoka, H.; Kita, K.; Marzuki, S., Mutation underlying resistance of Plasmodium berghei to atovaquone in the quinone binding domain 2 (Qo2) of the cytochrome b gene. *Parasitology International* **2008,** *57* (2), 229-232.

59.   Syafruddin, D.; Siregar, J. E.; Marzuki, S., Mutations in the cytochrome b gene of Plasmodium berghei conferring resistance to atovaquone. *Molecular and Biochemical Parasitology* **1999,** *104* (2), 185-194.

60.   Korbmacher, F.; Drepper, B.; Sanderson, T.; Martin, P.; Stach, T.; Maier, A. G.; Matuschewski, K.; Matz, J. M., An apicoplast□resident folate transporter is essential for sporogony of malaria parasites. *Cellular Microbiology* **2020**, 1-35.

61.   Rizopoulos, Z.; Matuschewski, K.; Haussig, J. M., Distinct Prominent Roles for Enzymes of *Plasmodium berghei* Heme Biosynthesis in Sporozoite and Liver Stage Maturation. *Infect. Immun.* **2016,** *84* (11), 3252-3262.

62.   Pei, Y.; Tarun, A. S.; Vaughan, A. M.; Herman, R. W.; Soliman, J. M. B.; Erickson-Wayman, A.; Kappe, S. H. I., *Plasmodium* pyruvate dehydrogenase activity is only essential for the parasite's progression from liver infection to blood infection. *Molecular Microbiology* **2010,** *75* (4), 957-71.

63.   Yoeli, M., *Plasmodium berghei*: mechanisms and sites of resistance to sporogonic development in different mosquitoes. *Experimental parasitology* **1973,** *34* (3), 448-458.

64.   Tadesse, F. G.; Ashine, T.; Teka, H.; Esayas, E.; Messenger, L. A.; Chali, W.; Meerstein-Kessel, L.; Walker, T.; Wolde Behaksra, S.; Lanke, K., et al., *Anopheles stephensi* Mosquitoes as Vectors of *Plasmodium vivax* and *falciparum*, Horn of Africa, 2019. *Emerg Infect Dis* **2021,** *27* (2), 603-607.

65.   Cohuet, A.; Osta, M. A.; Morlais, I.; Awono-Ambene, P. H.; Michel, K.; Simard, F.; Christophides, G. K.; Fontenille, D.; Kafatos, F. C., *Anopheles* and *Plasmodium*: from laboratory models to natural systems in the field. *EMBO Rep* **2006,** *7* (12), 1285-9.

66.   Molina-Cruz, A.; Canepa, G. E.; Kamath, N.; Pavlovic, N. V.; Mu, J.; Ramphul, U. N.; Ramirez, J. L.; Barillas-Mury, C., *Plasmodium* evasion of mosquito immunity and global malaria transmission: The lock-and-key theory. *Proc Natl Acad Sci U S A* **2015,** *112* (49), 15178-83.

67.   Lahree, A.; Mello-Vieira, J.; Mota, M. M., The nutrient games—*Plasmodium* metabolism during hepatic development. *Trends in Parasitology* **2023**.

68.   Peters, W., Drug resistance in *Plasmodium berghei* Vincke and Lips, 1948. I. Chloroquine resistance. *Experimental parasitology* **1965,** *17* (1), 80-89.

69.   Aurrecoechea, C.; Brestelli, J.; Brunk, B. P.; Dommer, J.; Fischer, S.; Gajria, B.; Gao, X.; Gingle, A.; Grant, G.; Harb, O. S., et al., PlasmoDB: a functional genomic database for malaria parasites. *Nucleic Acids Res* **2009,** *37* (Database issue), D539-43.

70.   Li, H.; Durbin, R., Fast and accurate short read alignment with Burrows–Wheeler transform. *bioinformatics* **2009,** *25* (14), 1754-1760.

71.   Andrews, S., FastQC: a quality control tool for high throughput sequence data. Babraham Bioinformatics, Babraham Institute, Cambridge, United Kingdom: 2010.

72.   Okonechnikov, K.; Conesa, A.; García-Alcalde, F., Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* **2016,** *32* (2), 292-294.

73.   Thorvaldsdóttir, H.; Robinson, J. T.; Mesirov, J. P., Integrative Genomics Viewer (IGV):
high-performance genomics data visualization and exploration. *Briefings in bioinformatics* **2013,**
14 (2), 178-192.
74.   McKenna, A.; Hanna, M.; Banks, E.; Sivachenko, A.; Cibulskis, K.; Kernytsky, A.;
Garimella, K.; Altshuler, D.; Gabriel, S.; Daly, M., The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*
2010, *20* (9), 1297-1303.
75.   Van der Auwera, G. A.; Carneiro, M. O.; Hartl, C.; Poplin, R.; del Angel, G.; Levy-
Moonshine, A.; Jordan, T.; Shakir, K.; Roazen, D.; Thibault, J., et al., *From FastQ Data to*
High-Confidence Variant Calls: The Genome Analysis Toolkit Best Practices Pipeline. John
Wiley & Sons, Inc.: Hoboken, NJ, USA, 2013; Vol. 467, p 11.10.1-11.10.33.
76.   Cingolani, P.; Platts, A.; Wang, L. L.; Coon, M.; Nguyen, T.; Wang, L.; Land, S. J.;
Lu, X.; Ruden, D. M., A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2;
iso-3. *Fly* **2012,** *6* (2), 80-92.
77.   Ruden, D. M.; Cingolani, P.; Patel, V. M.; Coon, M.; Nguyen, T.; Land, S. J.; Lu, X.,
Using *Drosophila melanogaster* as a model for genotoxic chemical mutational studies with a
new program, SnpSift. *Frontiers in genetics* **2012,** *3*, 35.
78.   Aho, A. V.; Kernighan, B. W.; Weinberger, P. J., Awk—a pattern scanning and
processing language. *Software: Practice and Experience* **1979,** *9* (4), 267-279.
79.   Waller, R. F.; Reed, M. B.; Cowman, A. F.; McFadden, G. I., Protein trafficking to the
plastid of *Plasmodium falciparum* is via the secretory pathway. *EMBO J* **2000,** *19* (8), 1794-
1802.
80.   Schindelin, J.; Arganda-Carreras, I.; Frise, E.; Kaynig, V.; Longair, M.; Pietzsch, T.;
Preibisch, S.; Rueden, C.; Saalfeld, S.; Schmid, B., Fiji: an open-source platform for biological-
image analysis. *Nature Methods* **2012,** *9* (7), 676-682.
81.   Livak, K. J.; Schmittgen, T. D., Analysis of Relative Gene Expression Data Using Real-
Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **2001,** *25* (4), 402-408.
82.   Yoshida, N.; Nussenzweig, R. S.; Potocnjak, P.; Nussenzweig, V.; Aikawa, M.,
Hybridoma produces protective antibodies directed against the sporozoite stage of malaria
parasite. *Science* **1980,** *207* (4426), 71-73.
83.   Foquet, L.; Hermsen, C. C.; van Gemert, G.-J.; Libbrecht, L.; Sauerwein, R.;
Meuleman, P.; Leroux-Roels, G., Molecular detection and quantification of *Plasmodium*
falciparum-infected human hepatocytes in chimeric immune-deficient mice. *Malaria journal*
2013, *12* (1), 1-8.
84.   Alcoser, S. Y.; Kimmel, D. J.; Borgel, S. D.; Carter, J. P.; Dougherty, K. M.;
Hollingshead, M. G., Real-time PCR-based assay to quantify the relative amount of human and
mouse tissue present in tumor xenografts. *BMC biotechnology* **2011,** *11*, 1-8.
85.   Yang, A. S.; Lopaticki, S.; O'Neill, M. T.; Erickson, S. M.; Douglas, D. N.;
Kneteman, N. M.; Boddey, J. A., AMA1 and MAEBL are important for *Plasmodium falciparum*
sporozoite infection of the liver. *Cellular microbiology* **2017,** *19* (9), e12745.
86.   Lopaticki, S.; Yang, A. S.; John, A.; Scott, N. E.; Lingford, J. P.; O'Neill, M. T.;
Erickson, S. M.; McKenzie, N. C.; Jennison, C.; Whitehead, L. W., Protein O-fucosylation in
Plasmodium falciparum ensures efficient infection of mosquito and vertebrate hosts. *Nature*
communications **2017,** *8* (1), 561.

87.   Kaushansky, A.; Rezakhani, N.; Mann, H.; Kappe, S. H., Development of a quantitative
flow cytometry-based assay to assess infection by *Plasmodium falciparum* sporozoites.
Molecular and biochemical parasitology **2012,** *183* (1), 100-103.
1235

A.

	anti-CSP	anti-ACP	Hoechst	Merge	Brightfield
Pb_WT					
PbAZMR_G95D_1					
PbAZMR_G95D_2					

B.

% of total counted

**** ****

API +
API -

Pb_WT   PbAZMR_G95D_1   PbAZMR_G95D_2

C.

Pb_WT

PbAZMR_G95D_1

PbAZMR_G95D_2

% of total counted

Irregular trails   ≤2 circular trails   3-20 circular trails   >20 circular trails

S1.

S2.