The Parkinson's drug entacapone disrupts gut microbiome homeostasis via iron sequestration

Fátima C. Pereira[a,b,+,*], Xiaowei Ge[c,+], Jannie Munk Kristensen[a], Rasmus H. Kirkegaard[a,e], Klara Maritsch[a], Yifan Zhu[d], Marie Decorte[a], Bela Hausmann[e,f], David Berry[a], Kenneth Wasmund[a,§], Arno Schintlmeister[a], Thomas Boettcher[a,g], Ji-Xin Cheng[c,h*], Michael Wagner[a,i*]

[a]Centre for Microbiology and Environmental Systems Science, Department of Microbiology and Ecosystem Science, University of Vienna, Vienna, Austria.
[b]School of Biological Sciences, University of Southampton, Southampton, United Kingdom.
[c]Department of Electrical & Computer Engineering, Boston University, Boston, Massachusetts, USA.
[d]Department of Chemistry, Boston University, Boston, Massachusetts, USA.
[e]Joint Microbiome Facility of the Medical University of Vienna and the University of Vienna, Vienna, Austria.
[f]Department of Laboratory Medicine, Medical University of Vienna, Vienna, Austria.
[g]Faculty of Chemistry, Department of Biological Chemistry, University of Vienna, Vienna, Austria.
[h]Photonics Center, Boston University, Boston, Massachusetts, USA.
[i]Department of Chemistry and Bioscience, Aalborg University, Aalborg, Denmark.

[+] These authors contributed equally
[§] Present address: School of Biological Sciences, University of Portsmouth, Portsmouth, United Kingdom.

Corresponding authors: f.c.pereira@soton.ac.uk, michael.wagner@univie.ac.at, jxcheng@bu.edu

Abstract

Increasing evidence shows that many human-targeted drugs alter the gut microbiome, leading to implications for host health. However, much less is known about the mechanisms by which drugs target the microbiome and how drugs affect microbial function. Here we combined quantitative microbiome profiling, long-read metagenomics, stable isotope probing and single-cell chemical imaging to investigate the impact of two widely prescribed nervous system-targeted drugs on the gut microbiome. *Ex vivo* supplementation of physiologically relevant concentrations of entacapone or loxapine succinate to faecal samples significantly impacted the abundance of up to one third of the microbial species present. Importantly, we demonstrate that the impact of these drugs on microbial metabolism is much more pronounced than their impact on abundances, with low concentrations of drugs reducing the activity, but not the abundance of key microbiome members like *Bacteroides, Ruminococcus* or *Clostridium* species. We further demonstrate that entacapone impacts the microbiome due to its ability to complex and deplete available ferric iron, and that microbial growth can be rescued by replenishing levels of microbiota-accessible iron. Remarkably, entacapone-induced iron starvation selected for iron-scavenging organisms carrying antimicrobial resistance and virulence genes. Collectively, our study unveils the impact of two under-investigated drugs on whole microbiomes and identifies metal sequestration as a mechanism of drug-induced microbiome disturbance.

Main

Drugs initially designed to specifically target human cells often can affect microbes as well[1]. As a result of poor gastrointestinal absorption and/or biliary secretion, many of these drugs reach the large intestine where they encounter - and potentially interact with - hundreds to thousands of different microbial species that play important roles in various aspects of human physiology[1,2,3]. Indeed, several cohort studies have reported significant associations between the use of medication and shifts in gut microbial composition and function[4,5,6,7]. While the cross-sectional nature of these large cohort studies hinders the establishment of causality, *in vitro* studies have been pivotal for systematically evaluating direct effects of human-targeted drugs on gut microbes. A landmark study assessing the antimicrobial effect of 835 human-targeted drugs against a panel of 40 cultured gut microbes revealed that a significant proportion (24%) of drugs could inhibit the growth of at least one gut bacterial strain *in vitro*[1]. An additional study testing the effect of a smaller panel of drugs on faecal samples by metaproteomics demonstrated selective anti- and/or pro-microbial activity for the great

majority of drugs tested, with a significant fraction of drugs also shifting microbiome
composition[8]. Of note, this study demonstrated that bacterial function could shift in response
to drugs without a change in taxon abundance, thus highlighting the need for using metrics
other than abundance when investigating the impact of drugs. Importantly, the interaction
between the microbiome and drugs is bidirectional, with many studies clearly demonstrating
that gut microbes can also actively metabolize[9,10,11,12], and under certain circumstances
bioaccumulate[13], pharmaceutical drugs. While several studies to date have shed light on the
nature and extent of microbe-driven drug transformations[9,10,11], mechanistic details for human-
targeted drug impact on the microbiome remain to be elucidated.

One hypothesis is that drugs can change intestinal microenvironments such as pH or
osmolarity, and by doing so, directly affect bacterial growth[14]. Another explanation is that
drugs interact with structural analogues of their human targets within bacteria, thus interfering
with cellular processes also in microbes[15]. Drug-microbiome interactions have been shown to
modulate the therapeutic effect of the drug, contribute to its side effects, or both[11,16,17,18].
Importantly, in certain cases, drug-induced microbiome changes might also contribute to other
diseases. For instance, proton pump inhibitors (PPIs) cause major shifts in the gut
microbiome[19], leading to decreased resistance to colonisation by enteric pathogens such as
Clostridioides difficile, *Campylobacter* and *Salmonella*[20]. Furthermore, PPIs-induced shifts in
the microbiome during early childhood have been linked to obesity[21].

Among the human-targeted drugs tested *in vitro*, compounds that target the nervous system
seem to exhibit stronger anti-commensal activity against gut bacteria compared to other
tested drug classes[1]. Indeed, several studies have reported a strong and selective inhibitory
activity of antipsychotics and antidepressants on gut microbial strains and microbiomes[22,23,24].
This is concerning, given the growing number of studies implicating the microbiome in many
neuropsychiatric disorders[25], and the widespread and rising use of this class of
pharmaceuticals worldwide[26]. Thus, a better mechanistic understanding of drug-microbiome
interactions in the context of nervous system-targeted medications may facilitate novel ways
to improve efficacy and/or minimize side effects of therapies for such disorders.

Here, we investigate the effects of two nervous system-targeted drugs on whole gut
microbiomes using a suite of complementary functional microbiome approaches. These aimed
at investigating the effects of the drugs in the context of whole microbial communities, which
more closely resembles their effects *in situ*, and to therefore examine most key members of
the gut microbiome. We studied two commonly used drugs: i) entacapone, a catechol-O-
methyltransferase (COMT) inhibitor that acts by preventing the degradation of levodopa, the
main drug used in the treatment of Parkinson's disease[27]; ii) loxapine succinate, a tricyclic
antipsychotic medication primarily used in the treatment of schizophrenia[28]. These drugs,

whose yearly total prescription exceeds 30 million tablets only in the US[29], belong to different

therapeutic classes and were shown to be selective in their anti-commensal activity against a

panel of 40 cultured gut strains, with entacapone predominantly targeting Gram-positive

organisms of the Firmicutes phylum and loxapine succinate only taxa within the Gram-

negative order Bacteroidales[1]. We demonstrate that the impact of either drug on microbiome

composition and/or activity extends beyond the taxa initially detected *in vitro* in pure culture,

with entacapone causing strong shifts in microbiome composition. This prompted us to look

for the cause of these shifts. We identified microbial iron deprivation, driven by the ability of

entacapone to complex iron, as the main mechanism behind entacapone's strong modulatory

effect. These results advance our understanding of the impact of nervous-system targeted

drugs on whole microbial communities and reveal micronutrient-deprivation as a mechanism

through which entacapone disrupts the microbiome.

Results

Nervous-system targeted drugs affect microbiome composition and abundance

To evaluate the impact of entacapone (ENT) and loxapine succinate (LOX) on whole gut

microbiome communities, we incubated freshly collected faecal samples from healthy adult

individuals in sM9 medium with two different concentrations of each drug dissolved in

dimethylsulfoxide (DMSO) (Fig.1a, see Methods). The low drug concentration (ENT-Low,

LOX-Low, 20 μM) was previously used in a screening aimed at determining drug effects on

pure culture isolates, while the high concentrations (ENT-Hi, 1965 μM, LOX-Hi, 100 μM) were

based on estimated colon concentrations for each drug, and were included to better reflect

the exposure of gut microbes to these drugs in the large intestine[1] (see Methods). After short

incubation times (6 and 24 hours) under anaerobic conditions at 37 °C, samples were

collected and processed to determine: i) changes of the total microbial loads; ii) microbial

community profile dynamics based on 16S rRNA gene amplicon sequencing; iii) reconstructed

microbial genomes based on long-read metagenomics and iv) single-cell microbial activity

changes via tracing of the incorporation of deuterium from isotopically-labelled heavy water

(D_2O) into single cells of microbiota by chemical imaging based on stimulated Raman

scattering spectroscopy (SRS)[30] (Fig.1a).

Counts of total microbial cell loads by flow cytometry in samples incubated with or without

drugs revealed that supplementation of a physiologically relevant concentration of

entacapone (ENT-Hi) significantly reduced the numbers of microbial cells over time, when

compared to all other tested conditions (Fig.1b, Supplementary Table 1). Major shifts in the

microbial community composition, as determined by 16S rRNA gene amplicon sequencing

analyses, were detected in response to ENT-Hi, LOX-Hi and LOX-Low treatments (Fig.1c),
and in the case of ENT-Hi these were also accompanied by shifts in alpha diversity (Extended
Data Fig.1a,b; Supplementary Table 2,3).

Figure 1. Drug-supplementation of faecal samples impacts biomass accumulation and microbiota
composition. a. Schematic representation of faecal sample incubations with the drugs entacapone (ENT)
and loxapine (LOX). As indicated, two different concentrations of each drug were used (Low and Hi).
Incubations were performed in the presence of 50% heavy water (D_2O) and 2% DMSO in the medium (either
sM9 or BHI), except for the H_2O control that consisted of medium without D_2O or DMSO. After 6 or 24 hours
of incubation, samples were collected for further analyses (see Methods). All incubations were performed in
triplicates. **b.** Total microbial cell loads in faecal sample incubations described in (A), at the start of the
incubation (time=0 hours) and after 6 and 24 hours, respectively. Total cell loads were assessed by flow
cytometry. **p<0.01; ***p<0.001; unpaired two-sample t-test with "No drug" used as the reference. **c, d.**
Relative (**c**) and absolute (**d**) genus abundance profiles of faecal microbial communities incubated for 6 or
24 hours as described in (**a**) and assessed by 16S rRNA gene amplicon sequencing. The community

composition at 0 hours (inoculum) is also shown. All genera present at a relative abundance below 0.025 or
absolute abundance below 6×10^6 cells.ml^{-1} were assigned to the category "Other". "unclass": unclassified.
e. Heatmap showing the absolute abundance, expressed in $\log_{10}$(cells+1), of the 25 most abundant ASVs
detected across all samples. Each column displays data from one replicate. **f**. Metagenome-assembled
genomes (MAGs) with 16S rRNA gene sequences matching ASVs shown in (**e**). MAGs were retrieved by
metagenomic sequencing of the initial faecal samples using Oxford Nanopore sequencing (see Methods
and Supplementary Table 7). 'NA" indicates no match of the ASV sequence to each MAG 16S rRNA gene
sequence.

By integrating total microbial counts with 16S rRNA gene amplicon sequencing data[31], we

determined absolute abundances for all detected taxa (Fig.1d). Importantly, absolute

abundance data confirmed that the employed incubation conditions enabled an increase in

abundance for nearly all the taxa initially present in faecal samples (Fig.1e: No drug versus

inoculum; Supplementary Table 4) and thus allowed tracking drug-induced activity and

abundance changes of microbiome members. Absolute abundances of genera such as

Bacteroides or *Clostridium sensu stricto* 1 decreased in both ENT-Hi and LOX-Hi samples

compared to the DMSO control (Fig.1d). However, many of the detected effects were drug-

specific, with ENT-Hi decreasing and LOX-Hi increasing total abundances of the genera

Anaerostipes, Fusicatenibacter, *Ruminococcus torques* group, *Eubacterium hallii* group,

Erysipelotrichaceae group UG-003 and *Roseburia*. Abundances of several genera were

significantly altered in response to ENT-Hi only: *Escherichia-Shigella* and *Ruminococcus*

increased in abundance, and genera such as *Alistipes*, *Streptococcus* or *Blautia* decreased

(Fig.1d,e; Supplementary Table 5). A similar impact of ENT-Hi on microbial biomass

accumulation and on the overall community composition could also be observed when we

used a different, nutrient rich medium (BHI) for the incubations (Extended Data Fig.1c-g).

Differential abundance analysis indicated that the abundance of 29.4% of all 16S rRNA gene

amplicon sequencing variants (ASVs) were significantly impacted by ENT-Hi after 24 hours

of incubation, 11.8% of ASVs were impacted by LOX-Hi, and only 3.6% and 6.0% of ASVs

were impacted by ENT-Low and LOX-Low, respectively (see Methods, Supplementary Table

5). Interestingly, LOX-Low resulted in growth inhibition patterns that differ from the ones

observed when the same drug was supplemented to gut members grown under isolation[1],

where it only specifically inhibits growth of Bacteroidales strains. Our results show that other

Gram-negatives and several Gram-positive species are affected by LOX-Low in the context

of whole microbiome communities (Supplementary Table 5). These include

Erysipelotrichaceae spp., *Oscillospiraceae* spp. and *Lachnospiraceae* spp., suggesting a

cross-sensitization to loxapine in the context of the community. For ENT-Low, only a very low

number of organisms were significantly impacted and were mostly Firmicutes, which agrees

with previous reports[1] (Supplementary Table 5).

Using long-read metagenomic sequencing of the starting faecal sample material, we retrieved
a total of 1049 metagenome-assembled genomes (MAGs), 11 of which are complete
genomes and 255 are medium- or high-quality genomes (Supplementary Tables 6 and 7).
BLASTn[32] analysis enabled us to link 16S rRNA gene sequences of ASVs to MAGs that
contained 16S rRNA genes and to follow drug-driven community shifts at a higher taxonomic
resolution (Fig.1e,f). This revealed that ASVs classified as *Escherichia* and *Ruminococcus*
taxa thriving in ENT-Hi give exact hits to the 16S rRNA genes in genomes of *E. coli* and *R.*
bromii. These analyses also indicated that LOX-Hi conditions selectively promoted the growth
of taxa like *Mediterraneibacter faecis*, *Faecalibacillus faecis* and *Blautia_A*. Species such as
Clostridium sp900539375 (*Clostridium sensu stricto* 1 based on SILVA taxonomy) and several
Bacteroides species *(B. uniformis, B. stercoris, Phocaeicola dorei* - formerly *B. dorei,* and *P.*
vulgatus - formely *B. vulgatus*) were totally or partially inhibited by the presence of high
concentrations of either drug, but the effect of ENT-Hi is much more pronounced than that of
LOX-Hi (Fig.1e,f). On the other hand, *Prevotella* sp900546535 and several Gram-positive
organisms such as *Streptococcus lutetiensis*, *Collinsella* sp003487125, *Bifidobacterium*
longum, *Mediterraneibacter faecis* or *Faecalibacillus faecis* showed growth inhibition by ENT-
Hi only (Fig.1e,f). All together these results reveal a strong but distinct impact of entacapone
and loxapine at physiological concentrations on microbiota composition and abundance, with
entacapone having a much more pronounced effect than loxapine.
226

Nervous system-targeted drugs can alter microbial metabolism without impacting
microbial abundance

Next, we evaluated the effect of these drugs on microbial activity at the single-cell level and
explored whether drug exposure that did not cause major shifts in microbiome abundance
within 24 hours of incubation did, nevertheless, impact their activity. To determine drug-
induced changes in microbial activity, we added D_2O as universal metabolic tracer[33,34,35] to
our incubations (Fig.1a). In complex microbial communities, all metabolically active cells will
incorporate deuterium (D) from D_2O into their biomass during synthesis of new
macromolecules[34]. The newly formed carbon-deuterium (C-D) bonds can then be used as a
read-out of microbial activity. Detection and quantification of C-D levels in single microbial
cells can be achieved using SRS, a method that efficiently excites the Raman active
vibrational modes coherently with two synchronized ultrafast lasers[36,37,38]. We have
successfully combined SRS with fluorescence *in situ* hybridization (FISH) in the past to
determine gut microbiome response to sugars with high-throughput[30]. Using an optimized
SRS-FISH platform that provides even higher throughput and sensitivity than the previous
setup (Supplementary Information text, Extended Data Fig.2), we measured around thirty

thousand individual microbiota cells after 6 and 24 hours of incubation in the presence of the drugs and in controls (Fig.2a,b). An unexpected, non-vibrational signal was detected in samples that were incubated with ENT-Hi and thus these samples were excluded from SRS-based activity measurements and the origin of this signal was further explored as detailed in the next section. As expected, in the absence of the drugs, nearly all of the analyzed cells were detected as active in the incubation medium, with 90% and 98% of cells displaying %CD$_{SRS}$ values above threshold at 6 hours and 24 hours of incubation, respectively (Fig.2b, Supplementary Table 8). Addition of either ENT-Low, LOX-Low or LOX-Hi resulted in a significantly reduced fraction of total active cells, as well as in a significant reduction of single-cell metabolic activity. This reduction was more pronounced for LOX-Hi, followed by LOX-Low and ENT-Low (Fig.2b). Thus, these drugs clearly impacted microbial activity within communities, even after short incubation times and under conditions where no impacts on their overall abundances were detected (Fig.1b).

To examine the activity of specific populations of the microbiome within the complex communities, we targeted six distinct abundant taxa of interest using SRS-FISH, which enabled us to determine the activity of individual cells from these taxa. Both published as well as newly designed and optimized oligonucleotide probes targeting 16S and 23S rRNA sequences predicted from MAGs retrieved by metagenomics were used for this purpose (Supplementary Table 9). Targeted taxa included organisms whose abundances were both negatively and positively impacted by drugs. We then evaluated the levels of activity of the targeted populations using picosecond SRS (Fig.2c, Extended Data Fig.2). Both LOX-Low and LOX-Hi incubations significantly reduced the activity of all targeted gut microbiota members except for *E. coli* (Fig.2d). Interestingly, *Clostridium* sp900539375, *B. uniformis*, *Ruminococcus bromii* and *P. dorei* show reduced activity in LOX treatments, but only the abundance of *P. dorei* and *Clostridium* sp900539375 was negatively impacted (Fig.2d,e; Supplementary Table 10). LOX-Low and LOX-Hi seem to strongly inhibit *P. dorei* growth within 6 hours (Fig. 1e), probably rendering most cells of this taxon undetectable by FISH, as the low ribosome content of non-active cells hinders FISH detection. We speculate that only a few drug-resistant *P. dorei* cells remained active enough to be detected by FISH, and these cells were not strongly impacted in activity (Fig.2c,d). A comparable decrease in activity was detected for *Streptococcus* spp., but in this case this decrease was surprisingly accompanied by a slight increase in abundance at 6 hours (Fig.2d,e). However, from 6 to 24 hours, we detect a decrease of the population represented by *Streptococcus lutentiensis* MAG in the presence of LOX compared to the control (Fig. 1e,f), which could be an effect of the lower activities detected by SRS in the LOX conditions at 6 hours. In summary, SRS-FISH provided new insights into the impact of ENT and LOX on the activities of specific microbiome

 members, which are often masked when only considering abundance data from relatively
 short incubation experiments.

Figure 2. Metabolic activity of drug-incubated single-microbiome cells measured by deuterium incorporation via SRS. a. Representative SRS images of faecal samples incubated with the indicated drugs for 6 hours. The measure %CD$_{SRS}$ indicates different levels of microbial activity. %CD$_{SRS}$ distribution images are displayed on top and C-H for biomass visualization (log scaled) on the bottom. %CD$_{SRS}$ scaling: minimum 0, maximum 20%. **b.** Single-cell %CD$_{SRS}$ values in each analyzed sample. Violin plots illustrating summary statistics (median, first and third quartiles, with the extended lines representing the minimum and maximum values within 1.5 interquartile ranges from the first and third quartiles). **p=0.0043, ***p=5.2e-12, **** p<2e-16, Wilcoxon test using "No drug" as the reference. **c.** Drug-incubated faecal samples were hybridized with fluorescently-labeled oligonucleotide probes targeting *E. coli* (Ecol_268), *Streptococcus* and *Lactococcus* species (Strc_493), *Clostridium* sp900539375 (*Clostridium sensu stricto 1,* Clo_648), *Ruminococcus bromii* (Brom_2036), *B. uniformis* (Buni_1001), and *Phocaeicola dorei* (Bado_374) (all shown in cyan, Supplementary Table 9). For each targeted group, top rows show representative images obtained by overlay of transmitted light (grey) and fluorescence intensity (cyan). Bottom rows show the corresponding SRS images (displaying %CD$_{SRS}$) for the FISH-targeted microbes (%CD$_{SRS}$ values of other microbes are not displayed for the sake of visibility). %CD$_{SRS}$ scaling: minimum 0, maximum 20%. Scale bar, 10 µm. **d.** Bubble plot denoting the fold change (FC, represented as Log$_2$FC) in activity levels (calculated as %CD$_{SRS}$) for the taxa targeted by FISH and incubated with drugs relative to "No drug" incubations. **e.** Bubble plot denoting the FC in absolute abundances for the taxa targeted by FISH and incubated with drugs relative to "No drug" incubations, as determined by DeSeq2.

Entacapone bioaccumulates in microbiota cells

Gut microbes have been shown to bioaccumulate some drugs leading to depletion of the drug from the surrounding environment[13,39]. As ENT-Hi samples showed a strong, Raman unspecific signal during SRS pump-probe detection, we explored the origin of this signal and concluded that this is a photothermal (PT) signal originating from entacapone bioaccumulation within microbial cells (Supporting Information Text). By mapping the intensity of this PT signal from ENT-Hi samples and controls (Fig.3a), we were able to show that the signal from entacapone occurred in a large fraction of microbial cells that had been incubated with the drug and washed before being fixed for analysis, but only in very few cells that had been fixed prior to incubation with the drug (i.e., dead cells) (Fig.3b,c, Supplementary Table 11).

Figure 3. Photothermal imaging of entacapone bioaccumulation by microbiota cells. a. Schematic illustration of the time-dependent signal obtained from a solution of 10 mM entacapone in DMSO, with photothermal (w/PT) and without photothermal (w/o PT) detection. **b.** Photothermal signal intensity distribution from entacapone, shown in green (PT, determined by photothermal imaging), and C-H signal, shown in red, of live and dead (PFA fixed) microbiota cells incubated in the presence (+) or absence (-) of ENT-Hi for 6 hours. PT channel contrast: min 0 max 1.8. C-H signals are represented on a log scale. Scale bar, 10 µm. **c.** Single-cell photothermal signal intensity distribution in samples shown in (a). p-values were determined using an unpaired two-sample Wilcoxon test. **d.** Representative images of faecal samples incubated with entacapone for 6 hours followed by hybridization with fluorescently-labeled oligonucleotide probes targeting *E. coli* (Ecol), *Streptococcus* and *Lactococcus* (Strc), *Ruminococcus bromii* (Rumi), *B. uniformis* (Buni), and *Phocaeicola dorei* (Pdor) (see Supplementary Table 9). FISH-probe signals from

hybridized cells (in cyan) were detected using widefield fluorescence microscopy (top panel). Photothermal
signal maps from entacapone (PT, in green) are shown in the middle (cells targeted by FISH are shown
with respective white contour lines). Entacapone PT signals overlayed with SRS C-H signals (in red) are
shown in the bottom panel. PT channel contrast: min 0 max 1.8. C-H signals are represented on a log scale.
Scale bar, 8 µm. **e.** Single-cell photothermal signal distribution in samples shown in (c). p-values were
determined using the unpaired two-sample Wilcoxon test. **f.** Time series of the absolute abundance of the
respective taxa targeted by FISH in No drug, Ent-Low and ENT-Hi incubations. In **c**, **e** boxes represent the
median, first and third quartile. Whiskers extend to the highest and lowest values that are within one and a
half times the interquartile range.

To identify the main drug accumulating taxa, we further looked at the entacapone distribution

among populations targeted with the FISH probes described above (Fig.3d, Supplementary

Table 9). All of the targeted populations displayed entacapone signals to different intensities.

While *P. dorei* and, to a lesser extent *Streptococcus* spp. and *E. coli,* seemed to be strong

entacapone accumulators, only a small percentage of *R. bromii* and part of the *B. uniformis*

population accumulated entacapone (Fig.3d,e). High entacapone signals were also detected

in cells not targeted by FISH (Fig.3e). We further confirmed the ability of *P. dorei* to

accumulate entacapone when grown in pure culture (Extended Data Fig.5). Interestingly,

while entacapone accumulation drastically inhibited the growth of *P. dorei* as a microbiome

community member and in pure culture*,* it did not affect *Streptococcus* growth in the

community to the same extent and showed even growth promotion for *E. coli* (Fig.3e). Thus,

bioaccumulation of the drug led to growth inhibition for certain taxa, whereas others were

unaffected or even stimulated in growth. This is in line with findings reported for the well-

studied antidepressant duloxetine, which was also found to be bioaccumulated by gut

microbiota taxa without impacting their growth[13].

Entacapone chelates iron and induces iron starvation in whole microbiome

populations

Entacapone's nitrocatechol group can act as a bidentate ligand, chelating and forming stable

complexes with transition metal ions such as iron (Fe) through the catecholate oxygen

atoms[40]. Using an assay widely applied to detect siderophores and other strong Fe-chelating

agents in solution, we confirmed entacapone's ability to chelate ferric iron (Fe(III)) (Fig.4a).

Binding of entacapone to metal ions such as Fe(III) has been proposed to occur via formation

of a tris complex[40]. The stability constant of entacapone's association with Fe (pFe(III)=19.3)

has been demonstrated to be similar to constants described for other known iron chelators

such as 2,2′-bipyridyl (pFe(III)=21.5)[41], but lower than reported for the siderophore

enterobactin (pFe(III)=49)[42]. Entacapone is also predicted to complex ferrous iron (Fe(II)) with

rather high affinity, but it is not predicted to form strong complexes with any other metal

cations[41].

Figure 4. Iron supplementation rescues the impact of entacapone on the gut microbiome. a.
Siderophore detection assay showing the change of color of the indicator complex from blue to orange/pink
(see Methods) in the presence of 0.1 mM of entacapone (in DMSO), but not in the presence of DMSO alone.
The indicator complex changed back to its original color (blue) after addition of an excess of ferric iron
($FeCl_3$, added at 0.1 to 1.0 mM). **b.** Growth of *Bacteroides thetaiotaomicron* in the absence of drug or in the
presence of: ENT-Hi; ENT-Hi preloaded with 1 mM $FeSO_4$ (ENT-Hi:Fe(II)), ENT-Hi, ENT-Hi preloaded with
1 mM $FeCl_3$ (ENT-Hi:Fe(III)); or ENT-Hi supplemented with the indicated concentrations of Fe(II) ($FeSO_4$)
or Cu(II) ($CuCl_2$)(see Methods). The box plots on the right refer to the same data as displayed on the left
plot, but split by the time of incubation (24 and 48 hours), for better visualization. **c.** Total cell loads in faecal
samples incubated for 6 and 24 hours with either ENT-Hi, 1mM $FeSO_4$ (No drug + 1.0 mM Fe), both (ENT-
Hi + 1.0 mM Fe), or ENT-Hi pre-incubated with 1 mM of $FeSO_4$ (ENT-Hi: Fe). Total cell loads were assessed
by flow cytometry. **d.** Relative (left panels) and absolute (right panels) genus abundance profiles of faecal
sample microbial communities incubated as described in c and assessed by 16S rRNA gene amplicon
sequencing. All genera present at relative abundances below 0.01 or absolute abundances below 1.5×10^6
cells.ml^{-1} were assigned to the category "Other". **e.** Alpha diversity metrics (Observed ASVs, Shannon index
and Inverse Simpson's diversity index) in gut microbial communities described in d. **f.** Bubble plot denoting
the fold change (FC, represented as Log_2FC) in absolute abundance relative to control incubations for the
25 most abundant ASVs that are affected by ENT-Hi treatment, as assessed by DeSeq2 analyses. All ASVs
have a base mean >50000, fold change of >1 or <-1 (relative no drug control), and $p_{adjusted}<0.05$. In **b**, **e**
boxes represent the median, first and third quartile. Whiskers extend to the highest and lowest values that
are within one and a half times the interquartile range. In **b**, **c** and **e:** *p<0.05; **p<0.01; ***p<0.001,
****p<0.0001; unpaired two-samples t-test with "No drug" used as a reference group. Only statistically
significant differences are indicated.

Iron is a limiting nutrient in the gut and essential for growth of most gut microbes[43]. Iron

concentrations in stool are estimated to be around 60 µM[44], and the medium used here for

faecal sample incubations contained similar concentrations of iron (31 µM). As the estimated

concentration for entacapone in the large intestine is approximately 2 orders of magnitude

higher (1965 µM), we postulated that the inhibitory effect of ENT-Hi on microbial growth could

be directly related to its ability to deprive gut microbes from iron via chelation, similarly to what

has been documented for other Fe(II) and Fe(III) chelators. To test this hypothesis, we grew

Bacteroides thetaiotaomicron, a gut commensal severely impacted by Ent-Hi (log$_2$-fold

change: -4.16, adjusted p-value: 0.016, Supplementary Table 5: ASV_g1t_iba) in the

presence of ENT-Hi or Fe-loaded ENT-Hi (Fig.4b). Fe-loaded entacapone was obtained by

addition of 1 mM of Fe(II) or Fe(III) to ENT-Hi followed by removal of any excess iron cations

via phosphate precipitation[45] (Fig.4b; see Methods). While ENT-Hi inhibited the growth of *B.*

thetaiotaomicron, ENT-Hi pre-complexed with either Fe(III) or Fe(II) did not, enabling *B.*

thetaiotaomicron to grow normally (Fig.4b). This is in agreement with previous studies

reporting that supplementation of the iron chelator 2,2′-bipyridyl inhibits growth of *Bacteroides*

fragilis, but this effect is largely diminished when bipyridyl is saturated with Fe(II)[46]. This is

because iron-saturated chelators are no longer able to complex free Fe(II) present in the

medium (or intracellularly), thus enabling bacteria to access iron and grow normally. As we

expected iron to be mostly present in the lower oxidation state under anaerobic conditions,

we used Fe(II) in all subsequent incubations. Supplementation of ENT-Hi alone followed by

addition of increasing amounts of Fe(II) also alleviated the inhibitory effect of ENT-Hi on *B.*

thetaiotaomicron, but only when Fe(II) was added at concentrations of 0.5 mM or above

(Fig.4b). Addition of similar amounts of cupric (Cu(II)) ions did not reverse growth inhibition

by ENT-Hi (Fig.4b), confirming that the observed effect is specific for iron. In summary, these

results demonstrate that supplementation of Fe-salts alone or of an ENT-Hi:Fe complex

rescues the inhibitory effect of ENT-Hi on *B. thetaiotaomicron,* strongly indicating that

entacapone drives iron limitation.

To determine whether the results above also apply to a complex microbiota, we incubated

faecal samples under equivalent conditions as described in Fig.1a with ENT-Hi, ENT-Hi

preloaded with Fe(II) or ENT-Hi followed by addition of Fe(II). In agreement with the results

obtained for *B. thetaiotaomicron* alone, supplementation of the whole microbiome with iron-

loaded ENT-Hi or with ENT-Hi followed by 1 mM of ferrous iron resulted in a complete or near

complete reversal of the inhibitory effect on microbial biomass accumulation (Fig. 4c,d,

Supplementary Table 12). Importantly, we confirmed that this effect is not due to lower cellular

uptake of iron-complexed entacapone, as ENT-Hi:Fe(II) bioaccumulates in microbiota cells to

the same level or higher than ENT-Hi alone (Extended Data Fig.6), thus suggesting that the

ENT-Hi:Fe(II) complex behaves similarly to entacapone alone in terms of its ability to penetrate and accumulate in cells. Supplementation of ferrous iron rescues the impact of ENT-Hi on the community alpha diversity metrics as well as on the growth of nearly all of the top 25 most affected taxa (exceptions are Firmicutes such as *Anaerostipes* and *Blautia*) (Fig4e,f; Supplementary Tables 13 and 14). Iron supplementation does not only enable taxa negatively impacted by entacapone to grow, but it also seems to restrict the accelerated expansion of organisms such as *E. coli*, prompted by the presence of ENT-Hi alone (Fig.1e and Fig.4d). All together these results strongly suggest that complexation of the limiting nutrient iron by entacapone is the primary mechanism behind the strong inhibitory effect of entacapone on the microbiome.

Entacapone promotes growth of iron-scavenging *E. coli*

Next, we interrogated the capability of the few indigenous microbiome members to thrive under the iron-limiting conditions induced by ENT-Hi. Several commensal and pathogenic *Enterobacteriaceae*, including *E. coli*, are known to synthesize and release siderophores which bind to Fe(III) with high affinity, enabling them to scavenge iron when it becomes limiting and in this way, enhance their gut colonization[47]. The most dominant organism in ENT-Hi incubations is an *E. coli* strain classified as *E. coli*_D (according to GTDB[48]), for which we were able to recover a complete MAG (bin.187, Supplementary Table 7). A search for genes involved in siderophore synthesis within the *E. coli*_D MAG led us to identify an entire gene cluster coding proteins necessary for synthesis, export, and import of the siderophore enterobactin (Fig.5a). Thus, we postulate that the ability of *E. coli*_D to produce enterobactin enables it to grow under iron-limiting conditions induced by ENT-Hi. Indeed, after isolating this *E. coli* strain from glycerol-preserved ENT-Hi incubations (isolate E2 described in methods), we could confirm its ability to grow in minimal medium prepared without iron (Fig.5b). We further demonstrated that its growth is not impacted (neither inhibited nor promoted) by the presence of ENT-Hi when grown under isolation (Fig.5b). Enterobactin chelates ferric iron with much higher efficiency than entacapone[41], and our results suggest that it enables *E. coli* to scavenge and acquire enough iron to sustain its growth under the iron-limiting conditions induced by the presence of entacapone (Fig.5c). The Ton-B siderophore receptor FepA may also enable *E. coli* to uptake and scavenge iron from exogenous catecholate:iron complexes[49], possibly even directly from iron-complexed entacapone. We could not find any genes involved in the production of known siderophores in the MAG of the ENT-Hi thriving *Ruminococcus*, and thus we speculate it expands in the entacapone supplemented medium by importing iron-loaded siderophores produced by *E. coli*, or by producing a yet unidentified siderophore or other high-affinity iron binding proteins (Fig.5c).

Figure 5. The iron-limiting conditions induced by entacapone select for siderophore producers and drive an increase in AMR and virulence potential in the faecal microbiome. a. Genetic organization of the enterobactin biosynthesis locus in *E. coli* D bin.187. Genes encoding enzymes involved in enterobactin biosynthesis (*entABCDEFH*) and export (*entS*) are highlighted in orange. The steps of enterobactin synthesis catalysed by the products of these genes are shown on the bottom. Genes involved in enterobactin import and iron release are highlighted in blue. **b.** Growth of an *E. coli* D bin.187 isolate determined by optical density measurements of cultures supplemented or not with 1965 μM entacapone (ENT-Hi) and/or 100 μM of $FeSO_4$. **c.** Schematic representation of the working hypothesis. In the absence of entacapone (left panel), enough iron is available to most gut microbiome members. In ENT-Hi conditions, entacapone complexes available iron (ENT:Fe), and only organisms able to produce siderophores (represented with blue and pink diamonds) for iron scavenging are able to grow and thrive (right panel). **d.** Increase in the AMR and virulence index (see Methods) in no drug controls and entacapone (ENT-Hi) treated samples. Indicated p-values were obtained from unpaired two-sample t-tests.

Siderophore production and iron scavenging are commonly observed in pathogenic or pathobiont strains that also tend to encode other virulence traits[50]. A screening of our MAG catalog for the presence of virulence factors and antimicrobial resistance genes[51] identified a diversified panel of AMR and virulence genes to be present in the retrieved gut MAGs (Supplementary Table 15), even though these samples originated from healthy individuals. These include genes encoding resistance to, among others, beta-lactams, tetracyclines, macrolides and aminoglycosides, as well as more general antimicrobial resistance genes such as efflux pumps, and virulence genes. ENT-Hi drives an increase in the abundance of AMR and virulence within faecal microbiomes (Fig.5d). This increase is driven in great part by the increase in abundance of *E. coli_D*, whose genome encodes a total of 14 AMR and virulence genes, in addition to at least one siderophore production cluster, thus suggesting a

483 pathogenic potential of this organism (Supplementary Table 15). Overall, these results reveal
484 that ENT-Hi promotes the growth of iron scavenging organisms with an associated pathogenic
485 potential.
486

Discussion

Gaining a deeper insight into the interactions between drugs and the gut microbiome is
essential for revealing and predicting how the microbiome might influence the availability,
efficacy and toxicity of pharmaceuticals. Here, we evaluated the impact of the nervous-system
targeted drugs loxapine succinate and entacapone on whole microbiomes derived from faecal
samples. Both drugs caused major shifts in microbial communities at physiologically relevant
concentrations, inhibiting the growth of many taxa while stimulating others. Using single-cell
activity measurements, we further revealed fine changes in the activity of specific community
members even at low drug concentrations and that were not captured by commonly used 16S
rRNA gene sequencing and abundance profiling. For instance, we did detect reduced *B.*
uniformis and *Clostridium* sp900539375 activities at 6 hours, but their abundance is not
affected until 24 hours (Fig.1e, Fig 2d). This is a major advantage of our approach as it
captures drug-induced changes in short incubation times, which are ideal for *ex vivo* systems
like the one in use. Importantly, our results show that loxapine succinate exerts an effect on
a broader range of taxa in the context of the community than its effects on microbes grown
under isolation[1]. This highlights the necessity of using complex microbial communities for a
better assessment of drug-microbiota interactions. One way by which some taxa may
sensitize or protect others to a particular drug is through chemical conversion or accumulation
of the drug[10]. However, previous studies have shown that the gut microbiota does neither
significantly bioaccumulate nor transform loxapine succinate[10]. Thus, we presume that cross-
sensitization to loxapine is likely due to drug-induced changes in microbial metabolites that
are involved in interspecies interactions.

The gut microbiome produces metabolites that can signal to the host and influence several
aspects of host physiology, including brain function[52]. Recent studies suggest that modulation
of the microbiome by drugs may contribute to the therapeutic effect of antipsychotics[1].
Interestingly, our results demonstrate that loxapine succinate promotes the growth of several
key gut species, including *Lachnospiraceae* (e.g. *Anaerostipes*) and *Butyricicoccaceae*
species, which have been shown to produce neuroactive metabolites such as short chain fatty
acids (SCFAs)[53]. SCFAs might directly or indirectly communicate along the microbiota-gut-
brain axis by activating G protein-coupled receptors or inhibiting histone deacetylases[54].
Additionally, they can modulate the blood-brain barrier, activate the vagus nerve, facilitate the

secretion of other hormones or neurotransmitters, as well as interfere with the immune response[54], all these factors contributing to its neuromodulatory effects. Thus, it may be plausible to consider changes in the microbiome as an additional mode of action of loxapine succinate. This may also help to explain why it may take weeks for this antipsychotic to act entirely[55], as its therapeutic effects may first require changes in the gut microbiome.

The impact of entacapone on the microbial community was very pronounced, selecting for the growth of organisms with pathogenic potential, such as *E. coli*. An increased abundance of *Enterobacteriaceae* was previously reported for Parkinson's disease (PD) patients taking entacapone[56], while other studies reported increases in *Enterococcaceae, Bifidobacteriaceae, and Clostridiales* family X[57]. These inconsistencies across studies can be partially attributed to the cross-sectional and rather small number of patients involved in each study. An increase in *Enterobacteriaceae* has also been linked to the side effects of other drugs[58] and may also help to explain diarrhoea experienced by some patients taking entacapone. In addition, entacapone is always prescribed in combination with levodopa to inhibit its off-site metabolism, and frequently also in combination with carbidopa, which inhibits levodopa off-site decarboxylation[27]. While levodopa has been shown to exert no effect on the distal gut microbiota, probably due to its high absorption in the upper gastrointestinal (GI) tract, future studies would be valuable to investigate the effects of a simultaneous supplementation of entacapone and levodopa to whole microbiome communities, especially in the upper GI tract. Biotic conversion of levodopa to dopamine by *Enterococcus* spp. and *Lactobacillus* spp. tyrosine decarboxylases in the small intestine significantly reduces its bioavailability to the host[59]. Given the strong inhibitory effect of entacapone on the microbiota, it would be interesting to evaluate whether entacapone can inhibit taxa that metabolize levodopa in the upper gastrointestinal tract, thereby contributing also in this way to increase the bioavailability of levodopa.

Like levodopa, entacapone has also been shown to be metabolized by gut microbial species[10]. Here, we demonstrate that entacapone is also bioaccumulated, with *Bacteroides, Phocaeicola, Streptococcus* and *Escherichia* spp. being able to accumulate sufficient amounts of entacapone for photothermal detection (Fig.3). This accumulation likely results in depletion of entacapone in the surrounding environment overtime, thus explaining the slight alleviation of entacapone's inhibitory effect between 6 and 24 hours of incubation, when compared to the first 6 hours (Fig.1b). Interestingly, using single-cell chemical imaging we further show that this bioaccumulation is heterogeneous, with some cells within a particular taxon accumulating varying amounts of the drug. In addition, we further show that entacapone bioaccumulation does not necessarily lead to growth inhibition (Fig.3e, f), similarly to what has been described before for other bioaccumulated drugs. It remains to be determined

whether entacapone bioaccumulation can be linked to its metabolism. Bacterial metabolism of entacapone occurs mostly by means of nitroreduction, which is prone to occur under anaerobic conditions, with several *Bacteroides* spp. encoding genes linked to entacapone nitroreduction[10]. Given the extensive conversion of entacapone by gut taxa observed previously, we predict a large fraction of entacapone's nitrocatechol group to be in its reduced form in our incubations and in the gut. Despite its bioaccumulation and/or conversion, we demonstrate that entacapone exerts a strong effect on the gut community due to its ability to complex iron via the catechol group.

Iron represents an essential enzyme cofactor in most bacteria[43]. Iron complexation by entacapone led to a decrease in biomass accumulation and in the abundance of most of the top abundant taxa detected in these samples, an effect that was reversed by supplementation of iron (Fig.4c-f). Taxa not found to be significantly impacted by entacapone were presumably able to grow by relying on intracellularly accumulated iron or on high-affinity surface-associated iron transporters that are ubiquitous in bacteria. Among stimulated taxa, the siderophore-producing *E. coli*_D strain present in our incubations greatly benefitted from entacapone's presence, but only in the context of the community, as entacapone supplementation to the isolate alone did not cause any significant boost in *E.coli*_D growth (Fig.5b). Thus, taxa stimulated by entacapone likely acquired iron via the mechanisms mentioned above, or via siderophores, and expanded in the community at the cost of nutrients released by dead cells or supplied via cross-feeding from other species. Treatment of communities with other iron complexing agents has shown somewhat similar effects on the gut community[60,61].

The expansion of an organism with siderophore synthesis, AMR and virulence potential in the presence of entacapone is concerning. Most successful gut pathogens tend to encode siderophore production systems and are therefore also expected to take advantage of the entacapone-induced remodeling of the gut community. It would be important to determine if entacapone treatment increases the likelihood of intestinal infections, similarly to what has been observed in patients taking PPIs[19]. As oral iron supplementation reduces levodopa and entacapone absorption in the small intestine[62,63], the effect of entacapone on the microbiome could be circumvented in the future by delivering iron specifically to the colon of patients taking the drug. If administered in appropriate amounts, this new adjuvant therapy could be expected to aid in preservation of microbiome homeostasis for patients taking entacapone, and presumably also for patients taking other catechol-containing drugs that might reach relevant concentrations in the large intestine (e.g. opicapone, apomorphine, fenoldopam mesylate) and that might affect the microbiome via similar mechanisms. Colon-targeted adsorbents, such as DAV132, could also be tested, as these have shown the ability to sequester antibiotic drugs

590 and reduce antibiotic-induced effects on the microbiome[64]. Our results advance our
591 understanding of the impact of antipsychotic and antidyskinetic drug therapies on microbiome
592 homeostasis, their mechanisms of action, and can direct future optimization of such therapies.
593

Methods

Ex-vivo gut microbiome incubations with drugs

Human faecal samples were collected from six healthy adult individuals (two males and four
females between the ages of 25 to 39) who had not received antibiotics in the prior 3 months.
Study participants provided informed consent and self-sampled using an adhesive paper-
based feces catcher (FecesCatcher, Tag Hemi, Zeijen, NL) and a sterile polypropylene tube
with the attached sampling spoon (Sarstedt, Nümbrecht, DE). The study protocol was
approved by the University of Vienna Ethics Committee (reference #00161). All meta(data) is
100% anonymized and compliant with the University's regulations. Samples were transferred
into an anaerobic tent (Coy Laboratory Products, USA) within 30 min after sampling, and all
sample manipulation and incubations were performed under anaerobic conditions (5% H_2,
10% CO_2, 85% N_2). Each sample was suspended in M9 mineral medium supplemented with
0.5 mg.mL^{-1} D-glucose (Merck), 0.5% v/v of vitamin solution (DSMZ Medium 461) and trace
minerals, herein referred to as sM9. Samples were suspended in sM9 to yield a 0.05 g.mL^{-1}
faecal slurry. At this point one aliquot of each sample was collected, pelleted, and stored at -
80°C for metagenomic analysis. The homogenate was left to settle for 10 minutes, and the
supernatant (devoid of any large faecal particles) was transferred into a new flask, where
supernatants from the six different donors were combined. This combined sample was further
diluted 1:10 in sM9 medium (as described above) or in supplemented Brain Heart Infusion
(BHI) medium containing either 0% or 55% D_2O (99.9% atom % (at%) D; Merck) for final 0%
(control) or 50% D_2O in incubation medium (Fig.1a). Supplemented BHI medium consisted of
37 g.L^{-1} of brain heart infusion broth (Oxoid), 5 g.L^{-1} yeast extract (Oxoid), 1 g.L^{-1} L-cysteine
(Merck) and 1 g.L^{-1} NaHCO$_3$ (Carl Roth GmbH, Germany). Incubation tubes were
supplemented with dimethylsulfoxide (DMSO, from Merck), entacapone (Prestwick
Chemicals) or loxapine succinate (Prestwick Chemicals) pre-dissolved in DMSO. The final
concentration was 2% w/v of DMSO in all vials (except for the H_2O control, where water was
added instead of DMSO). A subset of vials was supplemented with 20 µM or 1965 µM
entacapone (ENT-Low and ENT-Hi, respectively) and another subset was supplemented with
20 µM or 100 µM loxapine succinate (LOX-Low and LOX-Hi, respectively). The colon
concentration estimated for entacapone is 1965 µM[1], while for loxapine no estimate was
available. We predicted that loxapine would reach similar colon concentrations as its chemical

and therapeutic analogues amoxapine and clozapine, which are estimated to reach colon

concentrations of 138 µM and 153 µM, respectively[1]. Using these values as a reference, we

predict that loxapine succinate should be present in the colon at concentrations of at least

100 µM and chose it as the LOX-Hi concentration. At time 0, and after an incubation time of

6 or 24 hours at 37°C under anaerobic conditions, two sample aliquots from each incubation

and controls were collected by centrifugation. One aliquot was washed with 1× PBS and then

fixed in 3% paraformaldehyde solution for 2 h at 4°C. Samples were finally washed two times

with 1 ml of PBS and stored in PBS:Ethanol (50% v/v) at -20°C until further use. The second

pelleted aliquot was stored at -20°C until further processing. A third aliquot was collected into

sealed anaerobic vials containing 40% glycerol (Carl Roth GmbH, Germany) in PBS for a final

cell 50% v/v cell suspension in 20% glycerol and stored at -80°C until further use.

For iron rescue experiments, fresh faecal samples received from the same individuals (except

for one male that was traveling at the time of the experiment) were collected and processed

as described above. To establish appropriate controls for imaging of entacapone

bioaccumulation by microbiota cells, an aliquot of the freshly prepared 0.05 g.mL^{-1} faecal

slurry was immediately fixed with either 3% PFA solution or ethanol (50% v/v) at 4°C for 2

hours. Fixed faecal samples were washed with 1xPBS as described above and incubations

with fixed samples and entacapone or entacapone:iron (see below) were conducted in parallel

with incubations using live samples. Incubation vials were then supplemented with DMSO 2%

v/v with or without 1965 µM entacapone, in the presence or absence of supplemented iron

(1mM $FeSO_4$, Merck; Fig.5). Additional incubation vials (triplicates) were treated with

entacapone pre-complexed with iron: briefly, entacapone and iron ($FeSO_4$ or $FeCl_3$) powder

were mixed and resuspended in 2 mL of DMSO yielding a final concentration of 1965 µM

entacapone and 1 mM $FeSO_4$ (or $FeCl_3$) and stored overnight under anaerobic conditions.

The next day, 120µL of sodium phosphate monobasic (0.5M) was added and the samples

were mixed well. After 20 minutes, samples were centrifuged for 5 min, 14000 g to remove

unbound iron precipitated by the addition of sodium phosphate[45]. The supernatant containing

the iron-complexed entacapone was collected into a new eppendorf tube and supplemented

to the faecal incubation vials. Incubations were sampled as described above.

Cell counts from *ex-vivo* microbiome incubations

Microbial loads in faecal incubation vials were determined using flow cytometry and counting

beads as detailed below. Samples preserved in glycerol were diluted 200 to 800 times in 1x

PBS (Supplementary Table 1). To remove any additional debris from the faecal incubations,

samples were transferred into a flow cytometry tube by passing the sample through a snap

cap containing a 35 µm pore size nylon mesh. Next, 500 µL of the microbial cell suspension

was stained with the nucleic acid dye SYTO™ 9 (Thermo Fisher Scientific, 0.5 µM in DMSO)

for 15 min in the dark. The flow cytometry analysis of the microbial cells present in the

suspension was performed using a BD FACSMelody™ (BD Biosciences), equipped with a BD

FACSChorus™ software (BD). Briefly, background signals from the instrument and the buffer

solution (PBS) were identified using the operational parameters forward scatter (FSC) and

side scatter (SSC). Microbial cells were then displayed using the same settings in a scatter

plot using the forward scatter (FSC) and side scatter (SSC) and pre-gated based on the

presence of SYTO™ 9 signals. Singlets discrimination was performed. Absolute counting

beads (CountBright™, ThermoFisher Scientific) added to each sample were used to determine

the number of cells per mL of culture by following the manufacturer's instructions.

Fluorescence signals were detected using the blue (488 nm – staining with SYTO™ 9 and

CountBright™ beads) and yellow-green (561 nm - CountBright™ beads only) optical lasers.

The gated fluorescence signal events were evaluated on the forward–sideways density plot,

to exclude remaining background events and to obtain an accurate microbial cell count

(Supplementary Table 1). Instrument and gating settings were identical for all samples (fixed

staining–gating strategy).

678   Nucleic acid isolation and 16S rRNA gene amplification and sequencing

Pellets of microbiome incubation samples were resuspended in 600 µl of lysis solution RL

(InnuPREP DNA/RNA mini kit, Analytik Jena) and subjected to bead beating for 30 seconds

at 6.5 m/s in lysis matrix E (MPBiomedicals) tubes. After pelleting cell debris for 10 minutes

at 8000g, supernatants were transferred into the InnuPREP DNA/RNA mini kit spin filter tubes

(Analytik Jena) and DNA and RNA were extracted according to the manufacturer's protocol.

Amplification of bacterial and archaeal 16S rRNA genes from DNA extracts was performed

with a two-step barcoding approach[65] (UDB-H12).

In the first-step PCR, the primers 515F[66] (5′-GTGYCAGCMGCCGCGGTAA-3′) and 806R[67]

(5′-GGACTACNVGGGTWTCTAAT-3′), including a 5′-head sequence for 2-step PCR

barcoding, were used. PCRs, barcoding, library preparation and Illumina MiSeq sequencing

were performed by the Joint Microbiome Facility (Vienna, Austria) under project numbers

JMF-2208-05 and JMF-2103-29. First-step PCRs were performed in triplicate (12.5 µl vol per

reaction) with the following conditions: 1X DreamTaq Buffer (Thermo Fisher), 2 mM MgCl2

(Thermo Fisher), 0.2 mM dNTP mix (Thermo Fisher), 0.2 µM of forward and reverse primer

each, 0.08 mg ml−1 Bovine Serum Albumin (Thermo Fisher), 0.02 U Dream Taq Polymerase

(Thermo Fisher), and 0.5 µl of DNA template. Conditions for thermal cycling were: 95°C for

3 min, followed by 30 cycles of 30 s at 95°C, 30 s at 52°C and 50 s at 72°C, and finally 10 min

at 72°C. Triplicates were combined for barcoding (with eight PCR cycles). Barcoded samples

were purified and normalized over a SequalPrep Normalization Plate Kit (Invitrogen) using a
Biomek NXP Span-8 pipetting robot (Beckman Coulter), and pooled and concentrated on
PCR purification columns (Analytik Jena). Indexed sequencing libraries were prepared with
the Illumina TruSeq Nano Kit as described previously[68], and sequenced in paired-end mode
(2× 300 bp; v3 chemistry) on an Illumina MiSeq following the manufacturer's instructions. The
workflow systematically included four negative controls (PCR blanks, i.e., PCR-grade water
as template) for each 90 samples sequenced. The 16S rRNA gene sequences were deposited
in the NCBI Sequence Read Archive (SRA) as BioProject Accession PRJNA1033532.
705

Analysis of 16S rRNA gene amplicon sequences

Amplicon pools were extracted from the raw sequencing data using the FASTQ workflow in
BaseSpace (Illumina) with default parameters[65]. Demultiplexing was performed with the
python package demultiplex (Laros JFJ, github.com/jfjlaros/demultiplex) allowing one
mismatch for barcodes and two mismatches for linkers and primers. DADA2[69] R package
version 1.16.0 (https://www.r-project.org/, R 4.0.2) was used for demultiplexing amplicon
sequencing variants (ASVs) using a previously described standard protocol[70]. FASTQ reads
were trimmed at 150 nt with allowed expected errors of 2. Taxonomy was assigned to 16S
rRNA gene sequences based on SILVA taxonomy[71] (release 138) using the DADA2 classifier.

Samples were analyzed using the vegan (v2.5-.6; https://CRAN.R-
project.org/package=vegan) and phyloseq[72] (v1.30.0) packages of the software R
(https://www.r-project.org/, R 4.0.2). For samples subjected to different drug treatments,
sequencing in parallel of two extraction controls (without adding faecal samples) yielded 10
(control 1) and 189 reads (control 2). In control 1, 9 of the 10 reads were assigned to
Cyanobacteria or chloroplast and were not detected in the samples. Likewise, in control 2,
more than 90% of the reads originated from either Cyanobacteria ASVs or a single
Comamonadaceae (*Aquabacterium*) not detected in any of the samples. These ASVs were
removed from analysis. The remaining negative control reads (control 1: 1 read, control 2: 8
reads) were assigned to taxa typically found in the gut that were also detected in the samples
and were therefore retained for subsequent analyses. We assume these low number of reads
to originate from a low level of cross-contamination that can occur when multiple samples are
handled in parallel. After quality filtering and removal of contaminant sequences, a total of
1132 ASVs were retained. The average read number per sample was 11176 ± 3087 high-
quality sequences, and sample coverage was above 98% (Supplementary Table 2). For alpha
and beta diversity analysis, sequence libraries were rarefied to 4681 reads per sample. For
samples referring to the entacapone and iron supplementation experiment, sequencing in
parallel of two extraction controls yielded 2 (control 1) and 134 reads (control 2). After quality

filtering and removal of contaminant sequences (using the rationale described above), a total of 716 ASVs were retained. The average read number per sample was 18733 ± 4427 high-quality sequences and the sample coverage was above 99% (Supplementary Table 2). For alpha diversity analysis, sequence libraries were rarefied to 9729 reads per sample. For quantitative microbiome analyses, relative abundances of each taxon in a sample were calculated after correcting for the different number of copies of the 16S rRNA gene, according to *rrn*DB (version 5.7). For this correction, we classified ASVs using DADA2 and the RDP[73] taxonomy 18, release 11.5 (https://doi.org/10.5281/zenodo.4310151), by applying default parameters. These corrected relative abundances were then multiplied by the total microbial loads obtained from flow cytometry (Supplementary Table 1), yielding the total abundance of each taxon per sample.

DESeq2[74] (v1.26.0) implemented in phyloseq was used to identify significant differences in ASV abundances between drug treatments. Only ASVs that had in total ≥10 reads (relative abundance microbial profile) or 5.0e5 reads (quantitative microbial profile) were considered for comparisons by DESeq2 analyses. All statistical analysis of microbiome data was carried out with the software R (R 4.0.2). The applied significance tests and obtained p-values are referred to in the main text and figure legends.

Long-read sequencing

DNA for long-read sequencing was isolated using the DNeasy PowerSoil Pro Kit (Qiagen), according to the manufacturer's instructions. A pool of 6 DNA extracts was prepared for sequencing using the ligation sequencing kit (SQK-LSK112, Oxford Nanopore Technologies) following the manufacturer's protocol. The DNA was sequenced on a Promethion P24 (Oxford Nanopore Technologies) on a R10.4 flowcell (FLO-PRO112, Oxford Nanopore Technologies). The DNA sequencing was carried out using Minknow (v. 21.10.8, Oxford Nanopore Technologies).

Shotgun metagenomic sequencing

The same 6 samples were individually sequenced in an Illumina Novaseq 6000 platform by the Joint Microbiome Facility (Vienna, Austria), under project number JMF-2110-04. The illumina reads were trimmed using cutadapt[75] (v. 3.1). Illumina reads were mapped to the assemblies using Minimap2[76] (v. 2.17).

Metagenomic analysis

The Nanopore reads were assembled using flye[77] (v. 2.9-b1768) with "–nano-hq," polished three times with Minimap2[76] (v. 2.17) and Racon[78] (v. 1.4.3), followed by two rounds of

polishing with Medaka (v. 1.4.4, github.com/nanoporetech/medaka). Illumina and Nanopore

reads (Supplementary Table 6) were mapped to the assemblies using Minimap2 (v. 2.17) and

read mappings were converted using SAMtools[79] (v. 1.12). Read coverage and automatic

binning was performed using MetaBAT2[80] (v. 2.15). Contigs labeled as circular by the

assembler were extracted as independent bins before the automated binning process. The

quality of the recovered metagenome-assembled genomes (MAGs) was checked using

QUAST[81] (v. 5.0.2) and CheckM[82] (v. 1.1.1), and genomes were classified using GTDBtk[48] (v.

1.5.1).   rRNA   genes   were   detected   using   Barrnap   (v.   0.9,

https://github.com/tseemann/barrnap.) and tRNA genes were detected using trnascan[83] (v.

2.0.6). MAGs with a completeness >90 % but where barrnap did not pick up a 5S rRNA gene

were checked for 5S rRNA genes using INFERNAL[84] (v. 1.1.3).

All MAGs were searched for anti-microbial resistance (AMR) and virulence genes using

AMRFinderPlus[51] (v.3.10.21). The AMR and virulence index (Fig.5a) was calculated as

follows: the total copies of AMR and virulence genes found to be present in each MAG were

multiplied by the absolute abundance of the MAG (abundance of the ASV matching the 16S

rRNA gene of the MAG) in the sample. The same was repeated for all MAGs for which

AMRFinderPlus identified AMR or virulence genes and by summing these we were able to

predict the total number of copies of AMR and virulence genes for each sample, per mL of

culture. The resulting values were then normalized to the total biomass per mL of each sample

in order to obtain an AMR and virulence index per sample.

FISH probe design and optimization

Phylogenetic analysis and FISH probe design were performed using the software ARB v. 7.0[85].

By analysis of the 16S and 23S rRNA gene, phylogenetic trees were calculated with IQ-

TREE[86] (v 1.6.12) using the RAxML GTR algorithm with 1,000 bootstraps within ARB. For

abundant groups, 4 FISH probes were designed for this study and 2 additional published

probes were used (Supplementary Table 9). The probes were validated *in silico* with

mathFISH to test the *in-silico* hybridization efficiency of target and non-target sequences. The

number of non-target sequences was assessed using the probe match function in ARB and

the mismatch analysis function in mathFISH. All probes were purchased from biomers

(Biomers.net, Ulm, Germany) and were double labeled with indocarbocyanine (Cy3) or sulfo-

cyanine5 (Cy5) fluorochromes.

Pure cultures of *Escherichia coli* K-12, *Phocaeicola dorei* 175 (DSM17855) and *Bacteroides*

thetaiotaomicron VPI-5482 (DSM2079) were grown in supplemented BHI until the mid-

exponential phase and harvested by centrifugation. Pure cultures were fixed for 2 h by addition

of 3 volumes of 4% (w/v) paraformaldehyde solution at 4 °C. After washing once with PBS,

cells were stored in a 1:1 mixture of PBS and 96% (v/v) ethanol at − 20 °C. Where pure

cultures were not available, fixed faecal samples with a high relative abundance (as

determined by amplicon sequencing) of the specific target taxon was used.

To evaluate probe dissociation profiles, cells obtained from fixed pure cultures or faecal

incubation samples (Supplementary Table 9) were spotted onto microscopy slides (Paul

Marienfeld EN). FISH was performed as described before[87,88], with 3 or 5h hybridization to

obtain fluorescence signals with sufficient intensity. The optimal hybridization formamide

concentration was found using formamide dissociation curves, obtained by application of

formamide concentration series in the range from 0 to 70% in 5% increments[89]. After a

stringent washing step and counterstaining using 4′,6-diamidino-2-phenylindole, samples

were visualized using a Leica Thunder Epifluorescence microscope with an APO 100x/1,40

Leica oil immersion objective. Probe EUB338[90], which is complementary to a region of the

16S rRNA conserved in many members of the domain Bacteria, was used as a positive control

and a nonsense NON-EUB probe was applied to samples as a negative control for binding.

Images for inferring probe dissociation profiles were recorded using the same microscopy

settings and exposure times. The probe dissociation profiles were determined based on the

mean fluorescence signal intensities of at least 100 probe-labeled cells and evaluated by the

ImageJ software (v 1.53t). From the calculated average values, a curve was plotted and the

respective value right before a decline on each curve was defined as the optimal formamide

concentration.

FISH in solution

Fixed cells (100 µl) were pelleted at 14000 g for 10 min, resuspended in 100 µl 96% analytical

grade ethanol and incubated for 1 min at room temperature for dehydration. Subsequently,

the samples were centrifuged at 14000g for 5 min, the ethanol was removed, and the cell

pellet was air-dried. For SRS-FISH analysis, cells were hybridized in solution (100 µl) for 3 h

at 46°C. The hybridization buffer consisted of 900 mM NaCl, 20 mM TRIS HCl, 1 mM EDTA,

0.01% SDS and contained 100 ng of the respective fluorescently labelled oligonucleotide as

well as the required formamide concentration to obtain stringent conditions (Supplementary

Table 9). After hybridization, samples were immediately transferred into a centrifuge with a

rotor pre-heated at 46°C and centrifuged at 14000 g for 15 min at maximum allowed

temperature (40°C), to minimize unspecific probe binding. Samples were washed in a buffer

of appropriate stringency for 15 min at 48°C, cells were centrifuged for 15 min at 14 000 g

and finally resuspended in 20 µl of PBS. Cells (5 µl) were spotted on Poly-L-lysine coated

cover glasses No. 1.5H (thickness of 170 μm ± 5 μm, Paul Marienfeld EN) and allowed to dry

overnight at 4°C under protection from light. Salt precipitates were removed by dipping the

coverslips 2× in ice-cold Milli-Q water and allowed to dry at room temperature under protection

from light.

Picosecond stimulated Raman scattering (SRS) with widefield fluorescence
microscopy

An 80-MHz pulsed laser (InSight DeepSee+; Spectra-Physics) emitting two synchronized

femtosecond beams was used (Extended Data Fig.2a). One beam was tunable in wavelength

from 680 nm to 1300 nm, while the other beam had a fixed wavelength of 1040 nm. The time

delay between single pulses of the two beams is adjustable by a motorized delay line on the

1040 nm beam. To implement the picosecond stimulated Raman scattering (SRS) (Extended

Data Fig.2), the fixed beam (termed Stokes beam) is intensity modulated at 2.5 MHz by an

acousto-optic modulator (1205c; Isomet Corporation) and co-aligned with the tunable beam

(termed pump beam), by a dichroic mirror (DMLP1000; Thorlabs). Both beams are chirped by

SF57 rods to two picoseconds pulse-width and directed towards the lab-built upright

microscope frame. Then, a four-focal system and a flip mirror conjugate a pair of galvo mirrors

to the back aperture of a 60× water objective (UPlanApo 60XW, numerical aperture = 1.2;

Olympus) or a 100× oil objective (UAPON 100XOTIRF, numerical aperture = 1.49; Olympus),

allowing the collinear pump and Stokes beams to raster-scan the sample via synchronized

movement of the galvo mirrors. A 1.4 numerical aperture oil condenser (Aplanat Achromat

1.4; Olympus) collects the output beams, which are then reflected by a flip mirror and filtered

by a short pass filter (DMSP950; Thorlabs). Finally, the filtered-out pump beam is focused

onto a silicon photodiode connected to a resonant amplifier effective at a resonant frequency

around 2.5 MHz. The output alternative current signal is further amplified by a lock-in amplifier

(UHFLI; Zurich Instrument) at the frequency and in phase (x channel detection) with the

modulation. The output direct current signal is recorded for normalization. A data acquisition

card (PCIe-6363; National Instruments) collects the output signal for image generation.

To perform widefield fluorescence imaging for FISH-visualization of the identical sample areas

analyzed by SRS and photothermal imaging, two flip mirrors were flipped off (Extended Data

Fig.2a). A halogen lamp (12V100WHAL; Olympus) provides Kohler illumination of the sample

from the condenser side. Then, the objective and the tube lens conjugate the sample plane

to the camera (CS505CU; Thorlabs). To enable imaging of various fluorophores, different

excitation and emission filter sets were inserted between lamp and condenser, and in front of

the camera. For Cy3 imaging, two 530/10 nm bandpass filters (FBH530-10; Thorlabs) were

used as excitation filters and two 570/20 nm bandpass filters (ET570/20x; Chroma) were used

as emission filters. For Cy5 imaging, two 640 nm bandpass filters (FBH640-10; Thorlabs)
were used as excitation filters and two 670/20 nm bandpass filters (ET670/50m; Chroma)
were used as emission filters.
878
Chemical imaging of entacapone accumulation

By utilizing the multiphoton absorption of entacapone, we detected the photothermal signal
originating from optical absorption to generate entacapone distribution maps. The
experimental setup was identical with picosecond SRS, but with detection of the lock-in signal
by the y channel, which exhibits a $\pi/2$ phase delay relative to the intensity modulation by the
AOM (Extended Data Fig.4b). With this orthogonal phase detection, the interference of the
photothermal signal with the signals emerging from cross-phase modulation and SRS was
minimized (Extended Data Fig.4d).
887
Image acquisition and processing

Samples were prepared by drying fixed bacterial cells spotted onto poly-L-lysine coated
coverslips (VistaVision cover glasses, No. 1; VWR) in a 4°C refrigerator and subsequent
dipping into water three times to dissolve precipitates from the growth media. Then the
bacteria were immersed in 5 µL of water and sandwiched by another coverslip with a 0.11
mm thick spacer in between. To acquire deuterium incorporation profiles of microbiome
members labeled by FISH probes, widefield fluorescence was performed first. For different
fluorophores, corresponding excitation and emission filters were applied. The signal and color
gain of the camera were set to 5. Then the exposure time was adjusted to 0.5 to 5 seconds
depending on the fluorescence signal intensity. The widefield transmission image was
acquired by minimizing the condenser aperture and removing the filters.

To acquire the deuterium incorporation profile of the FISH-visualized cells, two flip mirrors
were inserted into the beam path to guide the pump and Stokes lasers to the sample. Three
SRS images, specific for Raman active vibrational modes of carbon-deuterium (C-D) bonds,
carbon-hydrogen (C-H) bonds as well as the off-resonance background signal were recorded
by tuning the wavelength of the pump beam to 849 nm, 796 nm and 830 nm, respectively.
These wavelength values correspond to spectral wavenumbers of 2163 cm^{-1} (C-D), 2947 cm^{-1}
^{1} (C-H) and 2433 cm^{-1} (silent region). Signal intensities were accumulated over increments of
20 cm^{-1}. Images were acquired sequentially within the identical field of view of 32 x 32 µm^2
with a raster step size of 106.8 nm. The per-pixel dwell time was set to 10 µs and, depending
on the signal intensity level, 1~10 image cycles were recorded to achieve a signal-to-noise
ratio (SNR) of >5 for single bacterial cells in the C-H spectral region.

For acquisition of entacapone distribution maps, the pump laser was tuned to 849 nm and the
signal detection was switched to the y channel of the lock-in amplifier. All images were
recorded utilizing the identical scanning parameters as applied for SRS.

To process the image data sets, firstly, the illumination patterns were corrected for both
widefield images (FISH) and point-scan images (SRS and entacapone distribution). Then, the
widefield images and point-scan images were co-localized via a calibrated projective
transform matrix. The fluorescence images were utilized to generate a single cell mask for
inference of the single cell activity and the drug accumulation level. The single cell activity is
expressed as $\%CD_{SRS} = (I_{CD}-I_{off})/(I_{CD}+I_{CH}-2I_{off})$, where the symbols I_{CD}, ICH and I_{off} refer to the
SRS signal intensities detected at the spectral positions assigned to C-D bonds, C-H bonds
and the silent region (off-resonance background). All intensity values were normalized to the
direct current intensity level detected at the photodiode. The relative entacapone
accumulation level is expressed as the signal intensity level detected in the y channel of the
lock-in amplifier. intensity outliers (>mean±2 standard deviations) observed in the SRS signals
and the photothermal signal of the human gut microbiome samples, were rejected from the
single cell masks. All imaging data analysis was performed with CellProfiler and Matlab.
926

Siderophore assay

The iron binding capacity of entacapone was tested using the SideroTec Assay[TM] (Accuplex
Diagnostics, Ireland), a colorimetric test for the detection of siderophores, according to the
manufacturer's instructions. Wells were read photographically and on a microplate reader at
630 nm wavelength (MultiskanTM GO Microplate Spectrophotometer, Thermo Fisher
Scientific).
933

Isolation and sequencing of *Escherichia coli* human gut isolate

Escherichia coli isolate E2 was isolated from glycerol-preserved faecal samples after
incubation in sM9 medium supplemented with 1965 µM entacapone for 48 hours under
anaerobic conditions. An aliquot of the glycerol stock was serially diluted in PBS, and 10^{-4} and
10^{-5} dilutions were plated in BHIs agar. Isolated colonies were re-streak 3 times to purity and
submitted to colony PCR using the 16S rRNA gene primers 616V (5'- AGA GTT TGA TYM
TGG CTC AG-3') and 1492R (5'- GGT TAC CTT GTT ACG ACT T -3'). Single colonies were
picked with an inoculation loop, resuspended in 50 µl of nuclease free water and boiled at
95°C for 10 minutes to lyse the cells and release cell contents. After a short spin to pellet cell
debris, 2 µl of the supernatant was added to a PCR reaction mix (final concentrations; Green
1X Dream Taq Buffer, dNTPs 0.2 mM, BSA 0.2 mg/ml, Taq polymerase 0.05 U/µl, primers 1
µM) prepared in a final volume of 50 µl per reaction. The amplification cycles were as follow:

initial denaturation at 95 °C for 3 min, followed by 30 cycles at 95 °C for 30 s, 52 °C for 30 s,
72 °C for 1.5 min, and a final elongation at 72 °C for 10 min. PCR products were visualized
on 1.5% agarose gel electrophoresis, and subsequently cleaned and concentrated on
columns (innuPREP PCRpure Kit, Analytik Jena) according to the manufacturer´s
instructions. Concentrations were measured by Nanodrop, and samples were sent for Sanger
sequencing at Microsynth AG (Vienna, Austria). Sequencing results were analyzed by
BLASTn[32] against the 16S rRNA gene sequences retrieved by metagenomics. The near full
16S rRNA gene sequences of isolate E2 obtained were 99% identical to the 16S rRNA copies
of *E. coli* D bin.187 and 100% identical to ASV_9ec_9n4.

Growth of microbial pure cultures in the presence of entacapone and/or iron

Bacteroides thetaiotaomicron (DSM 2079) cells were grown anaerobically (85% N_2, 10% CO_2,
5% H_2) at 37°C in *Bacteroides* minimal medium (BMM) containing 27.5 µM of iron ($FeSO_4$,
Merck)[91]. To test the effect of iron on rescuing the inhibitory effect of entacapone on *B.*
thetaiotaomicron growth, entacapone-treated (1965 µM) cultures were supplemented with
100 µM, 500 µM or 1mM of iron. Growth rescue was observed when cultures were
supplemented with either 500 µM $FeSO_4$ (Fe(II)) or $FeCl_3$ (Fe(III)), and for all subsequent
experiments only $FeSO_4$ was used. Cu(II)SO_4 (500 µM or 1mM - Merck) or entacapone pre-
complexed with iron (1965 µM, prepared as described above) were also supplemented to *B.*
thetaiotaomicron cultures. Samples for total cell counts were taken at 0, 24, and 48 hours of
growth under anaerobic conditions. *Phocaeicola dorei* (DSM 17855) was grown in BMM in
the presence or absence of 1965 µM entacapone. After 24 and 48 hours of growth, an aliquot
was collected and fixed with paraformaldehyde as described above, and stored at -20°C in
PBS:EtOH 50% v/v until further analyses. To test if growth of the *Escherichia coli* faecal isolate
E2 was dependent on the presence of iron, the isolate was grown in MOPS medium without
or without 10 µM of $FeSO_4$ in the presence or absence of entacapone for 24 hours at 37°C.
To avoid the transfer of any iron from the pre-inoculum, pre-inoculum cells were washed
thoroughly in MOPS medium without iron, prior to inoculation.

Determination of total cell loads in pure cultures incubated with entacapone and iron

To determine microbial cell loads in pure culture, samples were collected at 0, 24 or 48 hours
of growth and stained (either undiluted or after a dilution of 10 times in 1x PBS) with the
QUANTOM™ Total Cell Staining Kit (Logos Biosystems, Korea). Total cell counts were
determined using a QUANTOM Tx™ Microbial Cell Counter (Logos Biosystems, Korea),
according to the manufacturer's instructions.

Data availability

The 16S rRNA gene sequences data have been deposited in the National Center for Biotechnology Information (NCBI) Sequence Read Archive (accession number PRJNA1033532).

Code availability

CellProfiler pipelines and Matlab codes have been deposited in GitHub (https://github.com/buchenglab/srs-fish-drugs).

References

1. Maier, L. *et al.* Extensive impact of non-antibiotic drugs on human gut bacteria. *Nature* **555**, 623–628 (2018).
2. Weersma, R. K., Zhernakova, A. & Fu, J. Interaction between drugs and the gut microbiome. *Gut* **69**, 1510–1519 (2020).
3. Lynch, S. V. & Pedersen, O. The human intestinal microbiome in health and disease. *N Engl J Med* **375**, 2369–2379 (2016).
4. Falony, G. *et al.* Population-level analysis of gut microbiome variation. *Science* **352**, 560–564 (2016).
5. Zhernakova, A. *et al.* Population-based metagenomics analysis reveals markers for gut microbiome composition and diversity. *Science* **352**, 565–569 (2016).
6. Jackson, M. A. *et al.* Gut microbiota associations with common diseases and prescription medications in a population-based cohort. *Nat Commun* **9**, 2655 (2018).
7. Vich Vila, A. *et al.* Impact of commonly used drugs on the composition and metabolic function of the gut microbiota. *Nat Commun* **11**, 362 (2020).
8. Li, L. *et al.* RapidAIM: a culture- and metaproteomics-based Rapid Assay of Individual Microbiome responses to drugs. *Microbiome* **8**, 33 (2020).
9. Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R. & Goodman, A. L. Separating host and microbiome contributions to drug pharmacokinetics and toxicity. *Science* **363**, eaat9931 (2019).
10. Zimmermann, M., Zimmermann-Kogadeeva, M., Wegmann, R. & Goodman, A. L. Mapping human microbiome drug metabolism by gut bacteria and their genes. *Nature* **570**, 462–467 (2019).
11. van Kessel, S. P., Bullock, A., van Dijk, G. & El Aidy, S. Parkinson's disease medication alters small intestinal motility and microbiota composition in healthy rats. *mSystems* **7**, e01191-21 (2022).
12. Wallace, B. D. *et al.* Alleviating cancer drug toxicity by inhibiting a bacterial enzyme. *Science* **330**, 831–835 (2010).
13. Klünemann, M. *et al.* Bioaccumulation of therapeutic drugs by human gut bacteria. *Nature* **597**, 533–538 (2021).
14. Tropini, C. *et al.* Transient osmotic perturbation causes long-term alteration to the gut microbiota. *Cell* **173**, 1742-1754.e17 (2018).
15. Fung, T. C. *et al.* Intestinal serotonin and fluoxetine exposure modulate bacterial colonization in the gut. *Nat Microbiol* **4**, 2064–2073 (2019).
16. Bhatt, A. P. *et al.* Targeted inhibition of gut bacterial β-glucuronidase activity enhances anticancer drug efficacy. *Proc Natl Acad Sci USA* **117**, 7374–7381 (2020).
17. Wu, H. *et al.* Metformin alters the gut microbiome of individuals with treatment-naive type 2 diabetes, contributing to the therapeutic effects of the drug. *Nat Med* **23**, 850–858 (2017).

18. Sun, L. *et al.* Gut microbiota and intestinal FXR mediate the clinical benefits of metformin.
Nat Med **24**, 1919–1929 (2018).
19. Jackson, M. A. *et al.* Proton pump inhibitors alter the composition of the gut microbiota.
Gut **65**, 749–756 (2016).
20. Buffie, C. G. & Pamer, E. G. Microbiota-mediated colonization resistance against intestinal
pathogens. *Nat Rev Immunol* **13**, 790–801 (2013).
21. Stark, C. M., Susi, A., Emerick, J. & Nylund, C. M. Antibiotic and acid-suppression
medications during early childhood are associated with obesity. *Gut* **68**, 62–69 (2019).
22. Ait Chait, Y., Mottawea, W., Tompkins, T. A. & Hammami, R. Unravelling the antimicrobial
action of antidepressants on gut commensal microbes. *Sci Rep* **10**, 17878 (2020).
23. Cussotto, S. *et al.* Differential effects of psychotropic drugs on microbiome composition
and gastrointestinal function. *Psychopharmacology* **236**, 1671–1685 (2019).
24. Zhang, W., Qu, W., Wang, H. & Yan, H. Antidepressants fluoxetine and amitriptyline
induce alterations in intestinal microbiota and gut microbiome function in rats exposed to
chronic unpredictable mild stress. *Transl Psychiatry* **11**, 1–16 (2021).
25. Fang, P., Kazmi, S. A., Jameson, K. G. & Hsiao, E. Y. The microbiome as a modifier of
neurodegenerative disease risk. *Cell Host Microbe* **28**, 201–222 (2020).
26. Pratt, L. A., Brody, D. J. & Gu, Q. Antidepressant use among persons aged 12 and
over:United States,2011-2014. *NCHS Data Brief* 1–8 (2017).
27. Myllylä, V. V., Sotaniemi, K. A., Illi, A., Suominen, K. & Keränen, T. Effect of entacapone,
a COMT inhibitor, on the pharmacokinetics of levodopa and on cardiovascular responses
in patients with Parkinson's disease. *Eur J Clin Pharmacol* **45**, 419–423 (1993).
28. Schiele, B. C. Loxapine succinate: a controlled double-blind study in chronic
schizophrenia. *Dis Nerv Syst* **36**, 361–364 (1975).
29. Rome, B. N., Egilman, A. C., Patel, N. G. & Kesselheim, A. S. Using multiple authorized
generics to maintain high prices: the example of entacapone. *Value in Health* **26**, 370–
377 (2023).
30. Ge, X. *et al.* SRS-FISH: A high-throughput platform linking microbiome metabolism to
identity at the single-cell level. *Proc Natl Acad Sci USA* **119**, e2203519119 (2022).
31. Vandeputte, D. *et al.* Quantitative microbiome profiling links gut community variation to
microbial load. *Nature* **551**, 507–511 (2017).
32. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
search tool. *J Mol Biol* **215**, 403–410 (1990).
33. Kopf, S. H. *et al.* Trace incorporation of heavy water reveals slow and heterogeneous
pathogen growth rates in cystic fibrosis sputum. *Proc Natl Acad Sci USA* **113**, E110–E116
(2016).
34. Berry, D. *et al.* Tracking heavy water (D2O) incorporation for identifying and sorting active
microbial cells. *Proc Natl Acad Sci USA* **112**, E194–E203 (2015).
35. Wegener, G., Kellermann, M. Y. & Elvert, M. Tracking activity and function of
microorganisms by stable isotope probing of membrane lipids. *Curr Opin Biotechnol* **41**,
43–52 (2016).
36. Hong, W. *et al.* Antibiotic susceptibility determination within one cell cycle at single-
bacterium level by stimulated Raman metabolic imaging. *Anal Chem* **90**, 3737–3743
(2018).
37. Zhang, L. *et al.* Spectral tracing of deuterium for imaging glucose metabolism. *Nat Biomed*
Eng **3**, 402–413 (2019).
38. Heuke, S. *et al.* Shot-noise limited tunable dual-vibrational frequency stimulated Raman
scattering microscopy. *Biomed Opt Express* **12**, 7780 (2021).
39. Zimmermann, M., Patil, K. R., Typas, A. & Maier, L. Towards a mechanistic understanding
of reciprocal drug–microbiome interactions. *Mol Syst Biol* **17**, e10116 (2021).
40. Orama, M., Tilus, P., Taskinen, J. & Lotta, T. Iron(III)-chelating properties of the novel
catechol O-methyltransferase inhibitor entacapone in aqueous solution. *J Pharm Sci* **86**,
827–831 (1997).

41. Tosato, M. & Di Marco, V. Metal chelation therapy and Parkinson's disease: a critical review on the thermodynamics of complex formation between relevant metal ions and promising or established drugs. *Biomolecules* **9**, 269 (2019).

42. Loomis, L. D. & Raymond, K. N. Solution equilibria of enterobactin and metal-enterobactin complexes. *Inorg Chem* **30**, 906–911 (1991).

43. Mayneris-Perxachs, J., Moreno-Navarrete, J. M. & Fernández-Real, J. M. The role of iron in host-microbiota crosstalk and its effects on systemic glucose metabolism. *Nat Rev Endocrinol* **18**, 683–698 (2022).

44. Lund, E. K., Wharf, S. G., Fairweather-Tait, S. J. & Johnson, I. T. Oral ferrous sulfate supplements increase the free radical-generating capacity of feces from healthy volunteers. *Am J Clin Nutr* **69**, 250–255 (1999).

45. Stintzi, A., Barnes, C., Xu, J. & Raymond, K. N. Microbial iron transport via a siderophore shuttle: A membrane ion transport paradigm. *Proc Natl Acad Sci USA* **97**, 10691–10696 (2000).

46. Rocha, E. R., de Uzeda, M. & Brock, J. H. Effect of ferric and ferrous iron chelators on growth of Bacteroides fragilis under anaerobic conditions. *FEMS Microbiol Lett* **84**, 45–50 (1991).

47. Ellermann, M. & Arthur, J. C. Siderophore-mediated iron acquisition and modulation of host-bacterial interactions. *Free Radic Biol Med* **105**, 68–78 (2017).

48. Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2020).

49. Seyoum, Y., Baye, K. & Humblot, C. Iron homeostasis in host and gut bacteria – a complex interrelationship. *Gut Microbes* **13**, 1874855 (2021).

50. Ellermann, M. & Arthur, J. C. Siderophore-mediated iron acquisition and modulation of host-bacterial interactions. *Free Radic Biol Med* **105**, 68–78 (2017).

51. Feldgarden, M. *et al.* AMRFinderPlus and the Reference Gene Catalog facilitate examination of the genomic links among antimicrobial resistance, stress response, and virulence. *Sci Rep* **11**, 12728 (2021).

52. Lynch, J. B. & Hsiao, E. Y. Microbiomes as sources of emergent host phenotypes. *Science* **365**, 1405–1409 (2019).

53. Deleu, S., Machiels, K., Raes, J., Verbeke, K. & Vermeire, S. Short chain fatty acids and its producing organisms: An overlooked therapy for IBD? *eBioMedicine* **66**, (2021).

54. Dalile, B., Van Oudenhove, L., Vervliet, B. & Verbeke, K. The role of short-chain fatty acids in microbiota–gut–brain communication. *Nat Rev Gastroenterol Hepatol* **16**, 461–478 (2019).

55. Singh, A. N., Barlas, C., Saeedi, H. & Mishra, R. K. Effect of loxapine on peripheral dopamine-like and serotonin receptors in patients with schizophrenia. *J Psychiatry Neurosci* **28**, 39–47 (2003).

56. Weis, S. *et al.* Effect of Parkinson's disease and related medications on the composition of the fecal bacterial microbiota. *npj Parkinsons Dis.* **5**, 1–9 (2019).

57. Fu, S.-C. *et al.* A pilot study exploring the association ofentacapone, gut microbiota, and the subsequent side efects in patients with Parkinson's disease. *Front Cell Infect Microbiol* **12**, 837019 (2022).

58. Zhang, M. *et al.* Effects of metformin, acarbose, and sitagliptin monotherapy on gut microbiota in Zucker diabetic fatty rats. *BMJ Open Diab Res Care* **7**, e000717 (2019).

59. van Kessel, S. P. *et al.* Gut bacterial tyrosine decarboxylases restrict levels of levodopa in the treatment of Parkinson's disease. *Nat Commun* **10**, 310 (2019).

60. Dostal, A., Fehlbaum, S., Chassard, C., Zimmermann, M. B. & Lacroix, C. Low iron availability in continuous in vitro colonic fermentations induces strong dysbiosis of the child gut microbial consortium and a decrease in main metabolites. *FEMS Microbiol Ecol* **83**, 161–175 (2013).

61. Celis, A. I., Relman, D. A. & Huang, K. C. The impact of iron and heme availability on the healthy human gut microbiome in vivo and in vitro. *Cell Chem Biol* **30**, 110-126.e3 (2023).

62. Campbell, N. R. & Hasinoff, B. Ferrous sulfate reduces levodopa bioavailability: chelation as a possible mechanism. *Clin Pharmacol Ther* **45**, 220–225 (1989).

63. Novartis Pharmaceuticals. Comtan® (entacapone) tablets: Prescribing information (US).
New Jersey: Novartis Pharmaceuticals Corporation 2011.

64. Vehreschild, M. J. G. T. *et al.* An open randomized multicentre Phase 2 trial to assess the
safety of DAV132 and its efficacy to protect gut microbiota diversity in hospitalized patients
treated with fluoroquinolones. *J Antimicrob Chemother* **77**, 1155–1165 (2022).

65. Pjevac, P. *et al.* An Economical and Flexible Dual Barcoding, Two-Step PCR Approach
for Highly Multiplexed Amplicon Sequencing. *Front Microbiol* **12**, 669776 (2021).

66. Parada, A. E., Needham, D. M. & Fuhrman, J. A. Every base matters: assessing small
subunit rRNA primers for marine microbiomes with mock communities, time series and
global field samples. *Environ Microbiol* **18**, 1403–1414 (2016).

67. Apprill A, McNally S, Parsons R, & Weber L. Minor revision to V4 region SSU rRNA 806R
gene primer greatly increases detection of SAR11 bacterioplankton. *Aquat Microb Ecol*
75, 129–137 (2015).

68. Herbold, C. W. *et al.* A flexible and economical barcoding approach for highly multiplexed
amplicon sequencing of diverse target genes. *Front Microbiol* **6**, 731 (2015).

69. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat Methods* **13**, 581–583 (2016).

70. Callahan, B. J., Sankaran, K., Fukuyama, J. A., McMurdie, P. J. & Holmes, S. P.
Bioconductor Workflow for Microbiome Data Analysis: from raw reads to community
analyses. *F1000Res* **5**, 1492 (2016).

71. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res* **41**, D590–D596 (2012).

72. McMurdie, P. J. & Holmes, S. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217 (2013).

73. Cole, J. R. *et al.* Ribosomal Database Project: data and tools for high throughput rRNA
analysis. *Nucleic Acids Res* **42**, D633-642 (2014).

74. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

75. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet J **17**, 10–12 (2011).

76. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–
3100 (2018).

77. Kolmogorov, M., Yuan, J., Lin, Y. & Pevzner, P. A. Assembly of long, error-prone reads
using repeat graphs. *Nat Biotechnol* **37**, 540–546 (2019).

78. Vaser, R., Sović, I., Nagarajan, N. & Šikić, M. Fast and accurate de novo genome
assembly from long uncorrected reads. *Genome Res* **27**, 737–746 (2017).

79. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).

80. Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient
genome reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

81. Gurevich, A., Saveliev, V., Vyahhi, N. & Tesler, G. QUAST: quality assessment tool for
genome assemblies. *Bioinformatics* **29**, 1072–1075 (2013).

82. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. *Genome Res* **25**, 1043–55 (2015).

83. Lowe, T. M. & Eddy, S. R. tRNAscan-SE: a program for improved detection of transfer
RNA genes in genomic sequence. *Nucleic Acids Res* **25**, 955–964 (1997).

84. Nawrocki, E. P. & Eddy, S. R. Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics **29**, 2933–2935 (2013).

85. Ludwig, W. *et al.* ARB: a software environment for sequence data. *Nucleic Acids Res* **32**,
1363–1371 (2004).

86. Minh, B. Q. *et al.* IQ-TREE 2: new models and efficient methods for phylogenetic inference
in the genomic era. *Mol Biol Evol* **37**, 1530–1534 (2020).

87. Wagner, M., Horn, M. & Daims, H. Fluorescence in situ hybridisation for the identification
and characterisation of prokaryotes. *Curr Opin Microbiol* **6**, 302–309 (2003).

88. Yilmaz, L. S., Parnerkar, S. & Noguera, D. R. mathFISH, a web tool that uses thermodynamics-based mathematical models for in silico evaluation of oligonucleotide probes for fluorescence in situ hybridization. *Appl Environ Microbiol* **77**, 1118–1122 (2011).

89. Manz, W., Amann, R., Ludwig, W., Wagner, M. & Schleifer, K.-H. Phylogenetic oligodeoxynucleotide probes for the major subclasses of Proteobacteria: problems and solutions. *Syst Appl Microbiol* **15**, 593–600 (1992).

90. Daims, H., Brühl, A., Amann, R., Schleifer, K. H. & Wagner, M. The domain-specific probe EUB338 is insufficient for the detection of all Bacteria: development and evaluation of a more comprehensive probe set. *Syst Appl Microbiol* **22**, 434–444 (1999).

91. Varel, V. H. & Bryant, M. P. Nutritional features of Bacteroides fragilis subsp. fragilis. *Appl Microbiol* **28**, 251–257 (1974).

Acknowledgements

Research reported in this manuscript was funded by NIH Awards R35GM136223 (to J.-X.C.) and R01AI141439 (to J.-X.C.) and supported by the Boston University Micro and Nano Imaging Facility and Office of the Director, NIH Award S10OD024993. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. Funding for the presented research was also provided via the Young Independent Research Group Grant ZK-57 (to F.C.P.) of the Austrian Science Fund (FWF) and the FWF-Wittgensteinaward Z383-B (to M.W.) as well as the FWF-funded Cluster of Excellence "Microbiomes drive Planetary Health" (COE 7). We thank Jasmin Schwarz, Gudrun Kohl, Petra Pjevac and Joana Seneca Silva from the Joint Microbiome Facility of the Medical University of Vienna and the University of Vienna for assisting with amplicon and metagenomic sequencing, as well as repositing of sequencing data.

Author contributions

F.C.P., X.G., J.-X.C. and M.W. designed this study. F.C.P, X.G., J.M.K., K.M. and M.D. performed the experiments with input from T.B, Y.Z., D.B. and A.S.. F.C.P, J.M.K., R.H.K, B.H., and K.W. analyzed sequencing data and performed bioinformatic analysis. F.C.P. and X.G. wrote the manuscript with input from all co-authors and all authors read and approved the final manuscript.

Ethics declarations

The authors declare no competing interests.