Carboxydovores from the Pseudomonadota colonise volcanic soils during succession

Robin A. Dawson[1], Patricia Aguila[2], Gary M. King[3], Marcela Hernández[1]*

[1]School of Biological Sciences, University of East Anglia, Norwich, NR4 7TJ, UK

[2]Laboratorio de Microbiología Molecular, Escuela de Tecnología Médica, Universidad Austral

de Chile, Juan Soler Manfredini 1771, Puerto Montt, Chile

[3]Department of Biological Sciences, Louisiana State University, LA 70803, USA

Running title:

CO oxidation by bacteria in volcanic soils

*Corresponding author: marcela.hernandez@uea.ac.uk

ABSTRACT

Carbon monoxide (CO) degrading microorganisms are present in volcanic deposits

throughout succession, with vegetation and soil influencing the communities present. The

carboxydovores are a subset of CO degraders that use CO only as an energy source, raising

the question of how the physiological and metabolic features of the carboxydovores can make

these bacteria more competitive in harsh volcanic ecosystems. An enrichment strategy was

modified, which enabled the isolation of two carboxydovore representatives from genera that

were abundant in the native soils, *Cupriavidus* sp. CV2^T (92.3% ANI vs. *Cupriavidus basilensis*

DSM 11853) and a putative strain of *Paraburkholderia terrae* (*Pb. terrae* COX) (96.42% ANI vs.

Pb. terrae KU-64^T). These isolates oxidise CO across a very broad range of concentrations, and

genome sequence analysis indicated that they use form-I carbon monoxide dehydrogenase

(CODH) to do so. *Cupriavidus* sp. CV2^T and *Pb. terrae* COX each oxidised CO specifically at

stationary phase, but the conditions for induction of CODH expression were distinct.

Cupriavidus sp. CV2^T expressed CODH only in the presence of CO, while *Pb. terrae* COX

expressed CODH regardless of the presence of CO. Based on metabolic and phylogenetic

analyses, *Cupriavidus* sp. CV2^T is recommended as a novel species within the genus

Cupriavidus. Therefore, we propose the name *Cupriavidus ulmosensis* sp. nov. for the type

strain CV2^T (= NCIMB 15506^T, = CECT 30956^T). This study provides valuable insights into the

physiology and metabolism of carboxydovores, which colonise volcanic ecosystems during

succession.

Importance

Volcanic ecosystems harbour many bacteria that contribute to the environmentally important

process of carbon monoxide (CO) oxidation. We demonstrate a modified method for isolating

bacteria, which consume CO at very low concentrations as a supplementary energy source

(carboxydovory), leading to the isolation of two novel strains (*Cupriavidus* sp. CV2^T and

Paraburkholderia terrae COX) from volcanic strata that formed in 1917 and 2015,

respectively. The conditions under which CO consumption occurs were investigated; each

strain consumed CO during stationary phase, but *Pb. terrae* COX consumed CO regardless of

the prior growth conditions while *Cupriavidus* sp. CV2 was more controlled. *Cupriavidus* sp.

CV2 is a type strain of a new species, *Cupriavidus ulmosensis* str. CV2, which demonstrates

relatively high tolerance for CO. These strains provide the basis for further study of the

physiology, metabolism, and genetics of CO oxidation by carboxydovores, and will help us to
understand how bacteria colonise harsh volcanic ecosystems.
59

INTRODUCTION

Carbon monoxide (CO) is a toxic gas, which is emitted to the atmosphere in large quantities
each year (~2,600 Tg yr^{-1} emitted) (Khalil and Rasmussen, 1990), causing significant changes
to atmospheric chemistry and climate. Levels of CO in non-urban environments typically
range from 60-300 ppbv (King, 2003b), but anthropogenic sources can increase urban CO
concentrations into the low ppmv range (Wright *et al.*, 1975; Saadi *et al.*, 2021), causing issues
with air quality. Up to 2,100 Tg CO yr^{-1} is removed abiotically from the atmosphere through
reactions with hydroxyl radicals, hindering the subsequent removal of polluting compounds
(e.g. methane) (Crutzen and Gidel, 1983) and exacerbating the effects of global warming.
69

Soils act as a significant sink for CO due to microbial activity (Conrad and Seiler, 1980;
Bartholomew and Alexander, 1981), accounting for removal of ~250-300 Tg yr^{-1} (King and
Weber, 2007; Cordero *et al.*, 2019). Both aerobic and anaerobic microorganisms are capable
of CO oxidation due to the use of a Mo-Cu or Ni-Fe carbon monoxide dehydrogenases (CODH),
respectively (King, 2003b; Depoy *et al.*, 2021), although limited evidence is available to
suggest that anaerobic CO degraders play a role in atmospheric CO consumption. CO
degraders colonise diverse ecological niches, and isolates have been retrieved from
terrestrial, aquatic and even clinical samples (King and Weber, 2007; Stott *et al.*, 2008; King
and King, 2014). CO degraders can be broadly classified into two groups according to the
specific usage of CO; carboxydotrophs and carboxydovores. Carboxydotrophs can grow using
CO as the sole source of carbon and energy, with many isolates able to use high
concentrations of CO (e.g. 10% (v/v) (King and Weber, 2007), and can tolerate up to 90% (v/v)
headspace CO (Cypionka and Meyer, 1982; Mörsdorf *et al.*, 1992), although some
carboxydotrophs perform better with lower CO concentrations (e.g. 3% v/v) (Cypionka and
Meyer, 1982). It is unclear whether carboxydotrophs are able to oxidise atmospheric
concentrations of CO, although early studies suggested that they cannot (Conrad, 1996).
CODH catalyses the oxidation of CO to CO_2, meaning that carboxydotrophs can utilise a
pathway for carbon fixation such as the Calvin-Benson-Basham cycle to metabolise the
generated CO_2 while also gaining ATP and NAD(P)H + H$^+$ (Siebert *et al.*, 2020). Conversely,

while carboxydovores may oxidise higher concentrations of CO, they can also oxidise

atmospheric levels, but only benefit from this reaction by acquiring energy in the form of

electrons (King, 2003b; Cordero *et al.*, 2019). Carboxydovores often lack the genes required

for fixation of CO_2, but some reports indicate that these CO degraders are actually inhibited

by the high concentrations of CO required by carboxydotrophs for growth (Hardy and King,

2001; Weber and King, 2007).

The physiology, biochemistry and genetics of carboxydotrophs have been studied in detail,

with many studies focussing on the model bacterium *Oligotropha carboxidovorans* OM5[T]

(Schubel *et al.*, 1995; Nam *et al.*, 2003; Wilcoxen and Hille, 2013; Siebert *et al.*, 2020).

However, relatively few publications are available that focus on bacterial carboxydovores in

similar detail (King, 2003b; King and King, 2014; Cordero *et al.*, 2019; Islam *et al.*, 2019).

Recent studies have begun to demonstrate that carboxydovores may be abundant and

environmentally important bacteria (Quiza *et al.*, 2014), especially as this group are able to

oxidise atmospheric concentrations of CO (Islam *et al.*, 2019), but the contribution of

carboxydovores to CO uptake *in situ* remains to be established. We require a greater diversity

of cultured representatives to fully appreciate the contribution of carboxydovores to the

consumption of atmospheric CO. In this study, a targeted enrichment strategy was modified

and employed on volcanic soils, leading to the isolation of two novel carboxydovores from

the Pseudomonadota (formerly Proteobacteria). According to 16S rRNA amplicon sequence

analysis, these isolates are representatives of abundant taxa within the native volcanic

deposits. Physiological, biochemical, and genomic analyses revealed the metabolic flexibility

of these bacteria, as well as their distinctive use of CO in response to carbon limitation.

RESULTS AND DISCUSSION

Targeted enrichment and isolation of carboxydovores

In this study, an enrichment strategy was modified (from Weber and King, 2017) to isolate

carboxydovores from volcanic strata formed after eruptions in 1917 (regenerated soil) and

2015 (mostly rock). The enrichment conditions were designed to promote mixotrophy, using

0.5 mM pyruvate to facilitate growth and 100 ppmv CO to provide supplementary energy for

CO degrading bacteria. 100 ppmv CO was deemed to be too low to enrich carboxydotrophs,

as very high concentrations of CO (e.g. 10-20% (v/v)) are used by these bacteria for growth

(King, 2003b). Although heterotrophic bacteria would have been able to utilise the pyruvate, this enrichment strategy relied on the ability of carboxydovores to remain competitive by acquiring supplementary energy through CO oxidation, as suggested previously (Weber and King, 2010a). 0.5 mM pyruvate was selected as higher concentrations of organic carbon may inhibit CO consumption (Weber and King, 2007, 2012), but concentrations of pyruvate around 0.4 mM still permitted CO oxidation by members of the Burkholderiales (Weber and King, 2012). Similarly, *Labrenzia aggregata* consumed CO when grown with glucose concentrations ≤ 0.5 mM, suggesting that low concentrations of heterotrophic carbon would be suitable for a range of CO degraders (Weber and King, 2007). The enrichment conditions permitted CO oxidation by the microbial communities in Calbuco volcanic soils and sediments, as ~100 ppmv CO was consumed after 48 and 72 hours by the 1917 and 2015 stratifications, respectively (data not shown). CO was consumed by slurry enrichments and sub-cultured enrichments over the course of 700-hour incubations.

The carboxydovores, *Cupriavidus* sp. CV2[T] and *Pb. terrae* COX, were isolated from 1917 and 2015 volcanic stratifications, respectively. *Cupriavidus* spp. have previously been isolated from volcanic ecosystems due to their ability to oxidise H_2, but they have not been isolated as carboxydovores (Sato *et al.*, 2006). Previously, *Cupriavidus necator* was noted for containing a form-II CODH, but this form is not certain to oxidise CO as a physiological substrate (King and Weber, 2007). *C. necator* H16 grows autotrophically with CO_2 and H_2 and, although it was able to tolerate high CO concentrations, it could not oxidise CO to CO_2 (Wickham-Smith *et al.*, 2023). Conversely, CO oxidation by *Burkholderia* and *Paraburkholderia* species from volcanic soils has been well established (Weber and King, 2010b, 2017), although *Paraburkholderia terrae* has not been previously identified as a CO degrader.

Cupriavidus sp. CV2[T] and *Pb. terrae* COX are both members of the Pseudomonadota, a group that was previously noted for being highly abundant in vegetated volcanic deposits (Dunfield and King, 2004; Weber and King, 2010a; Hernández *et al.*, 2020a). The Pseudomonadota may be important to environmental CO cycling as, in vegetated volcanic soils, an increase in Pseudomonadota *coxL* sequences correlated with higher CO uptake activity (Weber and King, 2010a). The 2015 samples were almost entirely rock, with only traces of newly-formed soil present, likely necessitating the use of reduced gases such as CO_2, CO and H_2 by the resident

microbial community (Fierer *et al.*, 2010; Hernández *et al.*, 2020b). Recently formed soil on

volcanic sediments, such as the 1917 soils, provides a wealth of organic carbon to the resident

microbial community, coinciding with increased Pseudomonadota sequences (Weber and

King, 2010a). Weber & King (2010a) posited that CO oxidisers were able to compete with

other heterotrophs for organic carbon during ecosystem development, and CO is supplied to

these communities via plant roots (King and Crosby, 2002). However, vegetated Hawaiian

volcanic deposits were net emitters of atmospheric CO (King and Weber, 2008) in spite of the

high potential CO uptake potential in these soils, possibly due to abiotic CO production

associated with high organic matter concentrations. Grasses colonised the vertical surface of

the Calbuco stratification (Figure S1), potentially providing a source of CO to the resident

microbial community. However, it must be noted that *in situ* CO fluxes were not measured in

this study.

Relative abundance of the Pseudomonadota in recent volcanic deposits

The bacterial communities that inhabited the 2015 and 1917 volcanic deposits were largely

dominated by the Pseudomonadota (26.08% and 28.99% average relative abundance (RA),

respectively), Actinomycetota (29.09% and 5.21% RA, respectively) and Acidobacteriota

(10.79% and 26.02% RA, respectively) (Figure 1A). Other groups of note included the

Chloroflexota, previously noted for its abundance in recent volcanic deposits (Weber and

King, 2010a; Hernández *et al.*, 2020a; Dragone *et al.*, 2023) and its potential for CO oxidation

(King and King, 2014; Islam *et al.*, 2019; Hernández *et al.*, 2020b). The Chloroflexota increased

from a RA of 0.96% in the 2015 deposit to 8.01% in the 1917 deposit, indicating that this group

was less suited to colonising the very earliest volcanic deposit (2015), but was more

competitive during soil succession. The Acidobacteriota have been detected in relatively high

abundance in vegetated and unvegetated volcanic deposits from Llaima and Kilauea

volcanoes (Weber and King, 2010a; Hernández *et al*., 2020a). The Pseudomonadota were the

most abundant group in recent deposits from the Krafla volcanic field (Iceland) (Byloos *et al.*,

2018) and in deposits from Miyake-Jima (Japan) (Guo *et al.*, 2014). Members of the

Pseudomonadota were also abundant in deposits from the older/revegetated Llaima and

Kilauea volcanoes (Weber and King, 2010a; Hernández *et al.,* 2020a). This contrasts with the

relative abundance of the Pseudomonadota observed in the unvegetated 2015 deposit in

Calbuco (Figure 1A). The reason for the differing abundance of the Pseudomonadota in young

volcanic deposits is unclear, although it was previously suggested that the abundance of this group increased with increasing availability of organic carbon (Weber and King, 2010a). As *Cupriavidus* sp. CV2^T and *Pb. terrae* COX are members of the Pseudomonadota, the abundance of this group was studied in greater detail to determine the potential importance of these strains *in situ*.

Alphaproteobacteria dominated both the 2015 and 1917 volcanic deposits (32.97% and 34.99% RA, respectively), followed by Betaproteobacteria (12.22% and 18.64% RA, respectively) (data not shown). Within the Betaproteobacteria, Burkholderiaceae was dominated by the genus *Paraburkholderia* in the 2015 volcanic deposit (73.81% average RA), with a decrease in abundance in the 1917 deposit (41.67% RA) (Figure 1B). These data indicate that *Pb. terrae* is a member of a well-represented group bacteria *in situ*. The genera *Burkholderia* and *Paraburkholderia* were previously suggested to be important CO oxidisers within the Pseudomonadota (Weber and King, 2010a, 2017), and this study provides valuable evidence that CO-oxidising members of this group can be present in young, unvegetated volcanic deposits. *Cupriavidus* spp. were less abundant in the 2015 volcanic deposit (11.17% RA), but increased to 33.64% RA in the 1917 volcanic deposit (Figure 1B), indicating that *Cupriavidus* sp. CV2^T is relatively well-represented in the older soil. These data are expected, as *Cupriavidus*-related bacteria, many of which oxidised CO_2 and H_2, comprised 25% of culturable bacteria from volcanic deposits from Mt. Pinatubo (the Philippines) (Ogiwara *et al.*, 1999; Sato *et al.*, 2004, 2006).

Genome sequence analysis of novel carboxydovores
Strains *Cupriavidus* sp. CV2^T and *Pb. terrae* COX have similarity to *Cupriavidus basilensis* strain DSM 11853^T (99.33% nucleotide ID with 16S rRNA, BioSample ID SAMN12697569, Bioproject PRJNA563568) and *Paraburkholderia terrae* strain KU-64^T (NBRC 100964, 99.73%, Biosample ID SAMD00391168, Bioproject PRJNA33175). The genome of *Cupriavidus* sp. CV2^T has a completeness of 99.5 % with 2.3% contamination. The genome of *Paraburkholderia terrae* COX has a completeness of 99.7% with 2.4% contamination. Additional genome information is listed in Table S1. It is interesting that *Paraburkholderia terrae* strain COX was isolated from a recently deposited volcanic soil (lava 2015), as Weber and King (2010b) did not detect *Burkholderia*-like *coxL* sequences in bare Hawaiian samples.

In order to determine whether either carboxydovore was a member of a new species, genome-based taxonomic analysis was conducted using TYGS (Figures S2, S3), Average Nucleotide Identity (ANI) and digital DNA-DNA hybrisisation (dDDH) as well as autoMLST. *Cupriavidus* sp. CV2[T] showed an ANI of 92.3% and a dDDH of 47.5% to their closest type strain *Cupriavidus basilensis* strain DSM 11853 (data not shown). *Pb. terrae* COX showed an ANI of 96.42% and a dDDH of 71.1% to their closest type strain *Paraburkholderia terrae* strain KU-64 (NBRC 100964, data not shown). *Cupriavidus* sp. CV2[T] was determined to be a member of a new species, one that was closely related to the type strain *Cupriavidus basilensis* strain DSM 11853. *Pb. terrae* COX, however, was confirmed to be a *Paraburkholderia terrae*. CO-oxidising *Paraburkholderia* spp. have previously been isolated from soils from Kilauea volcano (1959 deposit, vegetated) (Weber and King, 2017).

The genome of *Cupriavidus* sp. CV2[T] is over 10 Mbp with a GC content of 64.8% (Table S1) compared with 8.9 Mbp and a GC content of 65.0% for the type strain *Cupriavidus basilensis* DSM 11853 (Salvà-Serra *et al.*, 2021). The genome of *Pb. terrae* COX is over 10 Mbp with a GC content of 62.0%, similar to the type strain *Paraburkholderia terrae* strain KU-64 (NBRC 100964), which has a genome measuring 9.9 Mbp with a GC content of 61.5%. The genomes of *Cupriavidus* sp. CV2[T] and *Pb. terrae* COX contain a large number of coding sequences (10,380 and 10,390, respectively), and subsystems analysis indicated that many of these genes were devoted to metabolism (Table S1). *Cupriavidus* sp. CV2[T] contains more genes with predicted roles involved in membrane transport and fatty acid metabolism, while *Pb. terrae* COX encoded more genes with predicted roles in iron acquisition and metabolism and carbohydrate metabolism (Table S1). The environmental source of each strain provides valuable context when considering their metabolic potential; *Cupriavidus* sp. CV2[T] was likely to encounter more organic material in its natural environment, while *Pb. terrae* COX would have to employ a versatile metabolism in order to take the fullest advantage of transient nutrient sources and trace gases. CO flux measurements were not taken at Calbuco volcano, but previous studies have demonstrated that significant quantities of CO are produced by plant roots and through abiotic processes (King and Crosby, 2002; King and Weber, 2008).

Consistent with the isolation of *Cupriavidus* sp. CV2[T] and *Pb. terrae* COX during enrichment

with CO, visually identical gene clusters were identified on each genome that were highly

likely to encode a form-I CODH (*coxMSL*) (Figure 2). *coxMSL* encodes the medium (FAD-

binding), small (2 x [2Fe-2S]) and large (Mo-Cu) subunits of carbon monoxide dehydrogenase,

respectively, with CoxL containing the active site and critical molybdopterin cystosine

dinucleotide cofactor (Johnson *et al.*, 1990; Meyer *et al.*, 1993). The form of *coxL* was

determined by identifying the active site motif AYXCSFR, characteristic of form-I CODH

(Dunfield and King, 2004), during alignment of the translated amino acid sequences with

known form-I CoxL sequences. CoxL amino acid sequences for *Cupriavidus* sp. CV2[T] and *Pb.*

terrae COX were most closely related to other Pseudomonadota CoxL sequences (Figure 3).

tBLASTn analysis of the *coxL* gene from *Cupriavidus* sp. CV2[T] against the genomes of

Cupriavidus spp. type strains yielded varying results; *Cupriavidus basilensis* DSM 11853[T] had

no *coxL* genes or *coxMSL* gene clusters, while *Cupriavidus necator* N-1[T] and *Cupriavidus*

oxalaticus strain Ox1[T] each encoded form-II CODH gene clusters (*coxSLM*), indicating that CO

oxidation using form-I CODH may be unusual in the genus *Cupriavidus*.

Cupriavidus sp. CV2[T] and *Pb. terrae* COX only possess a limited selection of the accessory *cox*

genes (*coxGDEF*), which were initially identified on the classic CODH-encoding gene cluster

from *O. carboxidovorans* OM5 (Figure 2). Previous studies in *O. carboxidovorans* OM5

demonstrated that the accessory Cox proteins (CoxBC, CoxDEFGHIK) have roles in the

maturation and anchoring of CODH (CoxMSL) (Meyer *et al.*, 2000; Nam *et al.*, 2003). The

translated CoxGDEF homologs identified in these novel carboxydovores only had low identity

(27.9-56.9%) with the corresponding translated amino acid sequences from strain OM5,

indicating that these proteins may have differing roles. Two distinct genes were present in

the novel strains when compared to strain OM5; *rcoM*, which encodes a putative heme-

containing CO-sensing transcription factor (Kerby *et al.*, 2008; Kerby and Roberts, 2012), and

mocA, which encodes a molybdenum cofactor cytidyltransferase that facilitates the assembly

of the molybopterin cytosine dinucleotide (MCD) cofactor required by CODH (Neumann *et al.*,

2009) (Figure 2). The presence of all accessory *cox* genes is not universally essential to CO

oxidation, as *Haliangeum ochraceum* is capable of expressing functional CODH with only

coxMSL and a single accessory gene (Quiza *et al.*, 2014; Weber and King, 2017). Further study

would benefit greatly from the use of targeted gene deletions to verify the genes that are
essential to CO oxidation by carboxydovores.

The genomes of *Cupriavidus* sp. CV2^T and *Pb. terrae* COX indicated that each bacterium lacked
a complete CO_2 fixation pathway. However, each genome encodes a phosphoenolpyruvate
carboxylase, meaning that assimilation of a small amount of carbon from CO through
anaplerotic CO_2 fixation cannot be discounted (Braun *et al.*, 2021). Carboxydovory was
confirmed as incubation of each strain with elevated concentrations of CO, with or without
organic carbon, failed to facilitate growth (Figure S4; Figures 3A, 3C). The genome of
Cupriavidus sp. CV2^T revealed a capacity for trace gas utilisation beyond CO alone. A gene
cluster encoding a putative Ni-Fe hydrogenase was detected, which had a high level of identity
at the amino acid level with the functionally verified hydrogenase gene cluster from
Cupriavidus necator H16 (Schwartz *et al.*, 2003) and, additionally, shared the same genetic
organisation (Figure S5). The ability to gain energy from the oxidation of both CO and H_2 may
provide a competitive advantage to carboxydovores such as *Cupriavidus* sp. CV2^T over other
mixotrophs during the colonisation of volcanic deposits (King, 2003a). Hydrogenases are
widely distributed in nature and their activity has been observed in volcanic cinders (King and
King, 2012). Hydrogenases also support microbial productivity when organic carbon is limited
(Greening *et al.*, 2016; Islam *et al.*, 2020). *Pb. terrae* COX lacked any recognisable
hydrogenase-encoding genes on its genome, suggesting that it employed other methods of
surviving in nutrient-poor unvegetated volcanic deposits. Evidence from genomes and studies
with cultured isolates have demonstrated the use of both CO and H_2 by many
carboxydotrophs and carboxydovores (Meyer and Schlegel, 1978; Mörsdorf *et al.*, 1992; King,
2003c; King and Weber, 2007; Islam *et al.*, 2019; Hernández *et al.*, 2020b), suggesting that the
use of both trace gases may be a relatively common strategy.

Physiological and metabolic characterisation
Cupriavidus sp. CV2^T and *Pb. terrae* COX grow using a relatively wide range of organic carbon
sources (Table S2). *Cupriavidus* sp. CV2^T was unable to use most of the sugars tested in this
study, with only weak growth observed with arabinose. Similarly, *Cupriavidus basilensis* DSM
18853 was unable to utilise glucose (Steinle *et al.*, 1998). *Pb. terrae* COX, however, grew on a
range of organic compounds. The growth substrate ranges observed in these novel strains

may reflect their native environments. *Pb. terrae* COX was isolated from a recent volcanic deposit (lava 2015), lacking in soil or organic matter, possibly explaining the metabolic versatility of this strain (Table S2). *Cupriavidus* sp. CV2^T, however, was isolated from soil from the 1917 volcanic stratum. This may allow the resident microorganisms to streamline their metabolism to make use of a smaller substrate range. With this in mind, it is curious to note that *Cupriavidus* sp. CV2^T still has a 10 Mbp genome containing many genes with predicted roles in metabolic pathways (Table S1). Although the genus *Cupriavidus* can use a wide range of substrates, including sugars in some cases (Brigham *et al.*, 2010; Park *et al.*, 2011; Pearcy *et al.*, 2022), *Cupriavidus* sp. CV2^T may be unable to use sugars such as glucose due to the lack of a gene encoding phosphofructokinase, as determined by KEGG analysis on the MicroScope annotation platform.

Cupriavidus sp. CV2^T grew on 5 mM pyruvate between pH 5.0-8.0 (Figure S6), and *Pb. terrae* COX grew between pH 5.0-7.0 (Figure S7). A relatively broad pH tolerance may be a beneficial trait to volcanic bacteria in general, as volcanic ash can significantly contribute to soil acidification (Lubis *et al.*, 2021). After 24 hours, *Cupriavidus* sp. CV2^T had grown to a statistically similar OD_{600} at pH 6.0, 7.0 and 8.0, with a small but significant decrease in growth at pH 5.0 when compared to the latter three conditions ($p \leq 0.05$) (Figure S6). No statistically significant difference was detected after 48 hours of growth ($p > 0.05$). Additionally, no growth was observed at pH 4.0 over the course of the experiment. *Pb. terrae* COX grew to a significantly higher final OD_{600} at pH 7.0 than at any other pH ($p \leq 0.01$), and remained significantly higher over the course of the experiment (Figure S7). No statistically significant difference in growth was observed between pH 5.0 or 6.0 over the course of the experiment, and no growth was observed at pH 4.0 or 8.0. These data indicated that *Cupriavidus* sp. CV2^T has a broader pH tolerance than *Pb. terrae* COX, possibly making it more suited to surviving in more changeable environments, but both strains are unable to grow under more acidic conditions.

CO consumption and tolerance of carboxydovores
Both *Cupriavidus* sp. CV2^T and *Pb. terrae* COX consumed CO when grown in liquid culture with 5 mM pyruvate (Figure 4A-4D). A significant difference in culture density for either strain during incubations 0-100,000 ppmv CO was only detected immediately after inoculation

(ANOVA, $p \leq 0.01$), with no significant difference in growth detected at any other timepoint. These data strongly supported the use of carboxydovory by *Pb. terrae* COX and *Cupriavidus* sp. CV2^T, as these strains were likely to be acquiring energy for maintenance from CO oxidation (King and Weber, 2007). 200 ppm CO was consumed below detection by *Pb. terrae* COX after 7 days (Figure 5), with an initial lag period of approximately 2 days before CO was consumed. Similarly, *Cupriavidus* sp. CV2^T did not consume CO for the first 72 hours of incubation, with 200 ppm CO consumed below detection after 7 days (Figure 4C). For each strain, CO consumption did not occur while growth on pyruvate was occurring (Figures 4 and 5), consistent with previous reports that carboxydovores switch to trace gas oxidation during stationary phase (Islam *et al.*, 2019). It was previously reported that carboxydovores were unable to grow using CO as the sole source of carbon and energy due to inhibition of CO oxidation at high concentrations of CO (Hardy and King, 2001; Weber and King, 2007). For example, CO uptake by *S. aggregata* was inhibited by concentrations of CO above 1000 ppmv (Weber and King, 2007). *S. aggregata* is unlikely to encounter high concentrations of CO *in situ* as 0.1 Tg C CO yr^{-1} is actually released to the top 10 m of the ocean (Conte *et al.*, 2019). The thermophilic carboxydovore *Thermomicrobium roseum* was able to oxidise CO completely at 50,000 ppmv and still exhibited the ability to consume CO at 200,000 ppmv, although it could not oxidise all of the CO (Wu *et al.*, 2009). *Cupriavidus* sp. CV2^T and *Pb. terrae* COX were isolated at an enrichment concentration of ~100 ppmv, lower than the affinities/requirements for carboxydotrophs (Cypionka and Meyer, 1982; Mörsdorf *et al.*, 1992; King and Weber, 2007), but the tolerance of these strains for higher concentrations of CO was unknown. Geothermal activity may cause transient changes in CO concentration in volcanic ecosystems, particularly as CO production by soils appears to increase with rising temperature (Inman *et al.*, 1971; Conrad and Seiler, 1980), but much of the CO available in established soils would be provided by plant roots (King and Crosby, 2002) and abiotic degradation of biomass (Conrad and Seiler, 1980). As transient increases in CO concentrations may occur in volcanic (and other geothermal) environments, the question of how *Cupriavidus* sp. CV2^T and *Pb. terrae* COX would respond to elevated CO was investigated.

CO uptake was observed between 100-10,000 ppmv CO (Figure 4), but not at 100,000 ppmv CO (data not shown), and notable differences in the lag period before CO consumption began were observed for each strain at different concentrations of CO (0-10,000 ppmv). *Cupriavidus*

sp. CV2^T consumed a significant amount of CO from the headspace of vials with 200 ppmv CO after 94 hours (one-way ANOVA: p≤0.001), 168 hours for 1,000 ppmv CO (p≤0.001), and 210 hours for 5,000 ppmv CO (p=5.68x10^{-9}) and 10,000 ppmv CO (1% v/v) (p=3.32x10^{-5}). *Pb. terrae* COX consumed a significant quantity of headspace CO after 70 hours with an initial concentration of 200 ppmv (one-way ANOVA: p≤0.001), 168 hours for 1000 ppmv (p≤0.01), 190 hours for 5,000 ppmv (p=2.47x10^{-7}) and did not consume a significant quantity of CO from the headspace with 10,000 ppmv (1% v/v) CO (p>0.05). The increasing lag before CO consumption with increasing initial CO concentrations, without any significant detriment to growth that could indicate broader toxicity to cellular processes (Figures 4A, 4C), indicated that elevated CO concentrations are inhibitory to the activity of CODH in *Cupriavidus* sp. CV2^T and *Pb. terrae* COX (Figures 4B, 4D).

Cupriavidus sp. CV2^T appears to have a greater tolerance for elevated CO than *Pb. terrae* COX, as *Cupriavidus* sp. CV2^T exhibited a shorter lag period before consuming a significant quantity of CO at 5,000-10,000 ppmv CO (Figures 4B, 4D), and *Pb. terrae* COX did not consume any CO from the 100,000 ppmv headspace during the test period. *Pb. terrae* COX exhibited the shortest lag period before consuming a significant quantity of CO at 200 ppmv initial concentration, indicating that this strain may be more effective at utilising smaller concentrations of CO. The levels of CO experienced by *Cupriavidus* sp. CV2 and *Pb. terrae* COX *in situ* are unknown. However, these data can be justified when considering the environmental differences between the sites of isolation for each strain, as the lack of soil in the deposit from 2015 may have required *Pb. terrae* COX to rely more heavily on reduced gases such as CO for persistence. *Cupriavidus* sp. CV2^T and *Pb. terrae* COX have a similar tolerance for CO when compared to carboxydovores such as *T. roseum* (Wu *et al.*, 2009), and a greater tolerance than marine CO degraders such as *S. aggregata* (Weber and King, 2007). The autotrophic non-CO degrader *Cupriavidus necator* H16 was able to tolerate 50% (v/v) CO following laboratory evolution, but the wild-type strain was inhibited by ≥15% (v/v) CO (Wickham-Smith *et al.*, 2023). As CO inhibited autotrophic growth in *C. necator* H16 rather than CODH activity, it is unknown whether tolerance to elevated CO could occur in a similar laboratory evolution study with *Cupriavidus* sp. CV2^T. Little information is available in the literature to indicate the typical level of tolerance to CO in *Paraburkholderia* spp. Overall, these data demonstrate that *Cupriavidus* sp. CV2 and *Pb. terrae* COX exhibit lower CO

tolerance compared to many carboxydotrophs (Cypionka and Meyer, 1982; Mörsdorf *et al.*,

1992). However, their tolerance levels are still higher than what appears necessary when

considering ambient CO concentrations ranging from 60-300 ppbv (King, 2003b), even with

transient increases in CO due to abiotic processes. Further studies are required to better

understand whether CO tolerance in these strains depends on CODH activity, whether these

strains benefit from CO-insensitive *o*-type cytochromes (Meyer *et al.*, 1986), and whether CO

tolerance is influenced by the ecology of CO-degrading bacteria.

Regulation of CO consumption by growth conditions and growth phase

The lack of CO consumption during the initial period of growth on pyruvate (Figure 4A-4D)

indicated that each carboxydovore switched from heterotrophic growth on organic

compounds (pyruvate) to chemolithoheterotrophic energy acquisition through CO oxidation

when organic carbon became limited. This would be consistent with previous observations by

Islam *et al*. (2019) who showed that the thermophilic carboxydovore *T. roseum* underwent

extensive metabolic remodelling during the shift to stationary phase, favouring processes

such as H_2 and CO oxidation. Mixotrophic use of CO during heterotrophic growth has also

been reported in carboxydotrophs (Kiessling and Meyer, 1982; Kim and Kim, 1989),

confirming that such metabolism is not limited to the carboxydovores. Additionally, CO-

induced *cox* gene transcription by the carboxydotroph *O. carboxidovorans* OM5 occurred

specifically in the absence of organic substrates (Santiago *et al.*, 1999). The conditions under

which *Cupriavidus* sp. CV2^T and *Pb. terrae* COX consume CO were investigated by growing

each strain with either pyruvate or a combination of pyruvate and CO, then testing the rate

of CO uptake during exponential or stationary phase (Figure 5A, 5B). The rate of CO oxidation

increased significantly during stationary phase compared to exponential phase for cells that

were grown with a combination of pyruvate and CO (one-way ANOVA: $p \leq 0.01$). Little to no

CO uptake was detected in cells harvested at exponential phase for either strain, consistent

with the use of CO oxidation as a starvation response by carboxydovores. A key difference

was seen between *Cupriavidus* sp. CV2^T and *Pb. terrae* COX during the shift to stationary

phase. *Cupriavidus* sp. CV2^T oxidised very little CO during stationary phase after growth with

pyruvate alone (Figure 5A), likely indicating that CODH expression requires the presence of

CO. *Pb. terrae* COX, however, oxidised CO at a very similar rate during stationary phase

regardless of the growth condition (one-way ANOVA $p \geq 0.05$) (Figure 5B), indicating that CO

uptake by *Pb. terrae* COX may not be regulated by the presence of CO but, rather, by the

presence or absence of other organic carbon sources.

The regulation of CODH expression is highly variable between different carboxydotrophs and

carboxydovores. CODH expression is constitutive in some carboxydotrophs such as

Pseudomonas carboxydoflava and *Hydrogenophaga pseudoflava* (Kiessling and Meyer, 1982;

Meyer *et al.*, 1986; Kang and Kim, 1999), although *P. carboxydoflava* begins to use gases such

as CO and H_2 only as supplementary energy sources (mixotrophy) when supplied with

heterotrophic substrates. This was suggested to support *P. carboxydoflava* in assimilating

surplus organic carbon (Kiessling and Meyer, 1982). Conversely, other carboxydotrophs have

demonstrated inducible expression of CODH (Kiessling and Meyer, 1982; Santiago *et al.*,

1999). Repression of CODH activity by organic carbon sources was previously demonstrated,

as CO uptake in the carboxydovore *S. aggregata* and carboxydotroph *O. carboxidovorans* was

reduced or eliminated (Santiago *et al.*, 1999; Weber and King, 2007). Weber and King (2007)

suggested that glucose (or glucose metabolites) exerted allosteric repression on CODH

expression in *S. aggregata,* even though this bacterium could not grow on glucose.

Carboxydovores such as *T. roseum* and *Mycobacterium smegmatis* switch to CO oxidation

during starvation (Cordero *et al.*, 2019; Islam *et al.*, 2019), suggesting that the regulation of

CODH expression is dependent on levels of organic carbon. Novel carboxydovores *Cupriavidus*

sp. CV2[T] and *Pb. terrae* COX both demonstrate inducible expression of CODH, as activity was

only observed during stationary phase (Figures 5A, 5B). The difference in CO uptake between

growth substrates for each strain is curious. Consistent with previous observations that

carboxydovores derive supplementary energy from CO oxidation during nutrient limitation

(Cordero *et al.*, 2019; Islam *et al.*, 2019), *Pb. terrae* COX, itself a resident of a nutrient-poor

unvegetated volcanic deposit, appeared to have developed a general CO scavenging response

during the shift to stationary phase (Figure 5B). *Cupriavidus* sp. CV2[T], isolated from a 1917

volcanic deposit with higher organic matter abundance, seemed to adopt a more conservative

approach, expressing CODH only during the stationary phase if CO was present (Figure 5A).

Detailed analysis of the metabolism and physiology of other carboxydovores may provide

greater insight into the mechanisms of control of CO oxidation.

Conclusions

This study demonstrates an effective strategy for enriching and isolating carboxydovore bacteria from volcanic soils. In this work two new strains are presented; *Cupriavidus* sp. CV2[T] and *Pb. terrae* COX. These strains consumed CO specifically during stationary phase, with *Cupriavidus* sp. CV2[T] employing a more conservative regulation strategy where CODH expression required the presence of CO, while *Pb. terrae* COX expressed CODH at stationary phase regardless of the presence of CO. Putative transcriptional regulators were identified with the *coxMSL* gene clusters, which could be studied with the aim of better understanding the environmental triggers of CO uptake. The tolerance of these strains to elevated CO concentrations is similar to other carboxydovores from geothermal environments, and their metabolic versatility and pH tolerance likely contribute to the successful colonisation of hostile volcanic sediments.

Further genomic comparisons of the isolate CV2[T] using MIGA, dDDH, ANI, and a set of 77 house-keeping genes using autoMLST, revealed similarities below the recommended ANI species-level thresholds (95%, Richter and Rosselló-Móra, 2009) and DDH values (70%, Tindall *et al.*, 2010) when the type strain genome sequences were used as query. *Cupriavidus* sp. CV2[T] is the representative of a novel species of the genus *Cupriavidus* within the order *Burkholderiales* (Phylum *Pseudomonadota*). Based on this characterisation, we propose *Cupriavidus* sp. CV2[T] as a new species, *Cupriavidus ulmosensis* sp. nov.

Description of *Cupriavidus ulmosensis* sp. nov

Cupriavidus ulmosensis sp. nov. (ul.mo.sen'sis. N. L. masc n. *ulmosensis*, of the Parque Valle Los Ulmos, referring to Parque Valle Los Ulmos in Chile, park named after the ulmo tree in Calbuco volcano, where the strain was isolated)

The type strain *Cupriavidus ulmosensis* CV2[T] (= NCIMB 15506[T], = CECT 30956[T]), was isolated from soil of Calbuco volcano in the Los Lagos Region, Chile. The genome is characterized by a size of 10.31 Mb and has a G + C content of 64.8 mol%. Gram-negative, rod-shaped aerobic heterotrophic bacterium. The isolate contains CODH-related genes and can grow in the presence of CO (200 – 100,000 ppmv CO). Cells grow at 30°C, at pH 4.0-8.0 (optimally at pH 8) and with 5 mM of pyruvate in VL55 medium (pH 5.5).

MATERIALS AND METHODS

Targeted enrichment and isolation of carboxydovores from volcanic soil

Soil samples were collected in February 2022 from Calbuco volcano (41.3304° S, 72.6087° W), Chile. Samples were taken from a lava stratification (Fig S1) formed by eruptions in 1893, 1917, 1961 and 2015 (the most recent eruption), characterised by different levels of soil development. Physical details of the lava stratification have been reported previously (Romero *et al.*, 2021). Soil samples (*in situ* pH ~5.5) were collected in triplicate at a 30 cm horizontally into the vertical surface of the lava strata except for the topmost layer, which had only volcanic rocks. Grasses and weeds (Figure S1) were removed before sampling. In this recent site (2015, approximately 5 cm of volcanic rocks were removed before sampling to a depth that did not cross the horizon of the 1961 stratum. Samples were stored in polyethylene bags (ziplok bags) until their transport to the laboratory in Chile, where they were refrigerated at 4 °C before shipping them to our laboratory in the UK. The carboxydovores discussed in this study were isolated from samples after the lava event in 2015 and 1917.

Slurries were prepared by mixing volcanic soils/sediments with VL55 medium (DSMZ recipe 1266) at a ratio of 1g:1ml, supplemented with 1 µl ml^{-1} vitamin solution (DSMZ recipe 1266). Carbon sources were ~100 ppmv CO and 0.5 mM pyruvate to support mixotrophy (supplied as sodium pyruvate). The slurries were incubated at 30 °C with shaking at 100 rpm. Headspace CO was measured daily as described above until it became undetectable, then the headspace was amended with approximately 100 ppmv CO. Following the consumption of ~300 ppm CO in total, 1 ml of the enrichment medium (without soil matter) was transferred to 11 ml of fresh VL55 with 0.5 mM pyruvate and 100 ppmv CO. Headspace analysis and amendments with CO were conducted as described above until a further 300 ppm CO was consumed. The enrichment medium was diluted across a 10-fold series to 10^{-5} and spread on VL55 plates, solidified with 1.5% (w/v) Bacto agar (Fisher Scientific) and amended with 0.5 mM pyruvate.

Colonies were screened for the presence of form-I *coxL*, the active site-containing component of CODH, using primers OMPf (5'-GGCGGCTT[C/T]GG[C/G]AA[C/G]AAGGT-3') and O/BR (5'-[C/T]TCGA[T/C]GATCATCGG[A /G]TTGA-3') reported by King (2003). The PCR protocol was

conducted as described previously (King, 2003b) with modifications to use colony biomass as the template. 5% (w/v) DMSO and 0.23% (w/v) BSA were added to the DreamTaq PCR mastermix (ThermoScientific, Waltham, MA) and the initial denaturation was maintained at 94 °C for 10 minutes to lyse cells. Colonies that contained form-I *coxL* were re-inoculated in sterile VL55 medium with 2 mM pyruvate and 100 ppmv CO and incubated at 30 °C for up to 2 weeks with shaking at 100 rpm. Cultures that consumed CO were serially diluted across a 10-fold series to 10^{-5} and spread on solid VL55 agar with 2 mM pyruvate. Colony morphology was examined for evidence of contamination and colonies were re-tested for the presence of form-I *coxL*. This process was repeated until axenic cultures of CO degrading bacteria were obtained. Two isolates were further characterised in this study (see below) and named *Cupriavidus* sp. CV2^T and *Paraburkholderia terrae* COX. *Cupriavidus* sp. CV2^T and *Pb. terrae* COX were cultivated in VL55 medium (pH 5.5, matching the pH of the original soils). Cells were grown at 30 °C in 120 ml vials using 20 ml medium with shaking at 150 rpm. CO consumption was measured by gas chromatography.

Extraction of DNA from volcanic soils and sediments

Total DNA from the soil samples was extracted using a DNeasy PowerSoil Pro kit (Qiagen) according to the manufacturer's instructions. DNA was amplified with primers 341F (5'-CCTAVGGGRBCCASCAG-3') and 806R (5'-GGACTACNNGGGTATCTAAT-3') and 16S rRNA gene sequencing was done using Illumina PE250 at Novogene, UK. The identification of Amplicon Sequence Variants (ASV), removal of chimeras and taxonomic assignment using the RDP classifier (threshold 80%) were carried out with the DADA2 algorithm (Callahan *et al.*, 2016) using Lotus2 software package (version 2.23, Özkurt *et al.*, 2022) on Galaxy Europe (https://usegalaxy.eu).

Cultivation of aerobic carboxydovores

The growth substrate ranges of each strain were tested by adding 5 mM of a defined carbon source to VL55 medium as the sole source of carbon and energy. Carbon sources were prepared as 1M stock solutions in distilled water and sterilised by passage through 0.2 μm syringe filters. Culture densities were recorded at 600 nm using a UV-1800 spectrophotometer (Shimadzu, UK) using two biological replicates. The pH range that supported VL55 medium was buffered to pH 4.0 using citric acid-Na_2HPO_4 buffer (0.1 M and

0.2 M, respectively), pH 5.0-6.0 using MES (2-(N-morpholino)ethanesulfonic acid), and pH 7.0-

8.0 using Trizima base. pH was adjusted using HCl or NaOH. To confirm that our isolates are

carboxydovores (i.e. unable to grow using CO as the sole source of carbon and energy), each

strain was initially grown using 5 mM pyruvate and 100 ppmv CO to induce CODH expression

(see below). CO was added to the headspace of 120 ml vials, sealed using butyl rubber

stoppers and aluminium crimp caps, from a 5000 ppmv stock prepared in N_2. Cultures were

sub-cultured into fresh VL55 medium with 1% (v/v) or 10% (v/v) CO as the sole carbon source,

using three biological replicates per condition. These cultures were incubated for 17 days.

Measurement of headspace CO

CO (CK Isotopes, UK) was measured by injecting 100 µl of headspace gas into an Agilent 7890A

gas chromatograph fitted with an Agilent HP-Molsieve PLOT (Porous Layer Open Tubular)

column (30 m length, 0.53 mm bore, 25 µm film, 7 inch cage,) at an initial oven temperature

of 50 °C, programmed at a rate of 10 °C/min to 100 °C with no initial hold time, injector at 250

°C (1:2 split ratio) and flame ionization detector at 300 °C (carrier gas He, 4 ml min^{-1}).

Headspace CO was quantified relative to standards containing a known quantity of CO (8.33

ppmv – 110,000 ppmv) prepared in N_2.

Tolerance of carboxydovores to elevated CO concentrations

The two isolates were inoculated in VL55 medium with 5 mM pyruvate to support growth.

The headspace of 120 ml vials was amended with 200 ppmv, 1000 ppmv, 10,000 ppmv (1%

v/v) or 100,000 ppmv (10% v/v) CO, using three biological replicates per condition. Headspace

CO concentrations were measured by gas chromatography as described above for a maximum

of 14 days, or until all detectable CO was consumed. Culture density (OD_{600}) was measured

using a spectrophotometer every 24 hours until growth ceased.

Differential CO uptake activity under different growth conditions and stages of growth

The isolates were cultivated in 400 ml VL55 medium in 2 L flasks with 5 mM pyruvate ± 200

ppmv CO, using three biological replicates per condition. 10 ml aliquots of cultures were

harvested at exponential phase (18-24 hours) or stationary phase (72 hours) by centrifuging

at 4000 *g* for 10 minutes at 4 °C, followed by resuspension to an OD_{600} of 4.0 in 1 ml of fresh

VL55 medium. Cell suspensions were kept on ice in sealed 30 ml vials during transfer to a

water bath heated to 30 ºC with shaking at 150 rpm. Vials were allowed to pre-warm for 3

minutes, then 200 ppmv CO was added to the headspace from a 5000 ppmv stock of CO in

N$_2$. After a further 1 minute, a headspace sample was measured by gas chromatography as

described above. Headspace samples were measured every 7.5 minutes until 6 samples had

been taken. Abiotic VL55 medium and heat-killed cell controls were tested under the same

conditions.

Statistical analysis

Statistically significant differences between conditions were compared by one-way analysis

of variance (ANOVA) followed by a Tukey post hoc test on R studio v.4.1.3.

Genome sequencing, annotation, and analysis

DNA was extracted from the bacterial strains grown in VL55 medium with 5 mM pyruvate

using the Qiagen Genomic-tip 100/G DNA isolation kit and associated DNA buffers (Qiagen)

according to the manufacturer's instructions. Genomic DNA concentration was quantified

using a Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific).

The strains were initially identified by amplifying the 16S rRNA gene using primers 27F/1492R

(Lane, 1991), and Sanger sequencing was conducted by Eurofins Genomics (UK).

Subsequently, whole genome sequencing was performed by MicrobesNG (Birmingham, UK).

Genomic DNA libraries were prepared using the Nextera XT Library Prep Kit (Illumina, San

Diego, USA) following the manufacturer's protocol with the following modifications: input

DNA was increased 2-fold, and PCR elongation time was increased to 45 seconds. DNA

quantification and library preparation were carried out on a Hamilton Microlab STAR

automated liquid handling system (Hamilton Bonaduz AG, Switzerland). Libraries were

sequenced on an lllumina NovaSeq 6000 (Illumina, San Diego, USA) using a 250 bp paired end

protocol. Reads were adapter trimmed using Trimmomatic version 0.30 (Bolger *et al.*, 2014)

with a sliding window quality cutoff of Q15. De novo assembly was performed on samples

using SPAdes version 3.7 (Bankevich *et al.*, 2012), and contigs were annotated using Prokka

1.11 (Seemann, 2014). The final sequences were uploaded to Microscope

(https://mage.genoscope.cns.fr/microscope (accessed 05/06/2023)) for annotation (Vallenet

et al., 2019). The form-I CODH-encoding genes (*coxMSL*) from the type strain *Oligotropha*

carboxidovorans OM5 (Santiago *et al.*, 1999) were used as queries in tBLASTn analysis against the newly sequenced genomes to identify putative *cox* gene clusters. The translated CoxL component was aligned against a database of corresponding CoxL sequences using MEGA7 (Kumar *et al.*, 2016) to confirm the presence of the form-I CoxL active site motif (AYXCSFR) (Dunfield and King, 2004). To determine whether our two isolates were members of previously undescribed species, the genome sequence data were uploaded to the Type (Strain) Genome Server (TYGS) (https://tygs.dsmz.de) for whole genome-based taxonomic analysis. Additionally, Microbial Genomes Atlas Online (MIGA), average nucleotide identity (ANI) and average amino acid sequence identity (AAI) was calculated using the MIGA (http://microbial-genomes.org) and Enveomics (Rodriguez-R and Konstantinidis, 2016) platforms, respectively, and confirmed with the outputs from automated multi-locus species tree analysis (autoMLST) (Alanjary *et al.*, 2019).

ACKNOWLEDGEMENTS

The authors would like to thank Barbara Corrales, Pablo Saumann, and the sampling team at Parque Volcánico Valle Los Ulmos, Chile, for their assistance with sampling at Calbuco volcano and for their warm hospitality during our visit. The authors also wish to thank Professor Aharon Oren and Professor Ramon Rosselló-Móra for their valuable discussions on bacterial identification. Robin A. Dawson was supported by Royal Society Research Fellows Enhanced grants awarded to Marcela Hernández (RF\ERE\210050 and RF\ERE\231066). Marcela Hernández was supported by a Royal Society Dorothy Hodgkin Research Fellowship (DHF\R1\211076).

SUPPLEMENTARY MATERIAL

Supplementary material for this article can be found in the online version.

DATA AVAILABILITY STATEMENT

The genome sequences of the isolates have been deposited in the NCBI GenBank under the accession number PRJNA1001293 (SAMN36798297 for *Cupriavidus* sp. CV2^T and SAMN36798260 for *Paraburkholderia terrae* COX). The 16S rRNA gene Sanger sequences have also been deposited in NCBI under the accession numbers OR536588 for *Cupriavidus* sp. CV2T^T and OR536592 for *Pb. terrae* COX. The amplicon-sequencing data of the soils were

deposited in the NCBI Sequence Read Archive (SRA) under the bioproject accession number

PRJNA1036796**.** Both strains have been deposited in culture collections under accession

numbers NCIMB 15506^T and CECT 30956^T for Cupriavidus sp. CV2^T and NCIMB 15505

for *Paraburkholderia terrae* COX. The CoxL database was assembled by Prof. Gary King and is

available upon request.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

MH conceived the project and secured the funding; MH and PA collected the samples; RAD

performed laboratory experiments; MH performed 16S rRNA gene amplicon sequence

analysis; GK provided advice on the enrichment and isolation of bacteria; RAD and MH

analysed the data; RAD and MH drafted the paper; all authors contributed to its revisions and

approved the final version.

REFERENCES

Alanjary, M., Steinke, K., and Ziemert, N. (2019) AutoMLST: an automated web server for

generating multi-locus species trees highlighting natural product potential. *Nucleic*

Acids Res **47**: W276-W282. doi: 10.1093/nar/gkz282.

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., et al. (2012)

SPAdes: a new genome assembly algorithm and its applications to single-cell

sequencing. *J Comput Biol* **19**: 455-477. doi: 10.1089/cmb.2012.0021.

Bartholomew, G.W. and Alexander, M. (1981) Soil as a sink for atmospheric carbon

monoxide. *Science (80-)* **212**: 1389–1391.

Bolger, A.M., Lohse, M., and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**: 2114-2120. doi: 10.1093/bioinformatics/btu170.

Braun, A., Spona-Friedl, M., Avramov, M., Elsner, M., Baltar, F., Reinthaler, T., et al. (2021)

Reviews and syntheses: Heterotrophic fixation of inorganic carbon - Significant but

invisible flux in environmental carbon cycling. *Biogeosciences* **18**: 3689-3700. doi:

10.5194/bg-18-3689–2021.

Brigham, C.J., Budde, C.F., Holder, J.W., Zeng, Q., Mahan, A.E., Rha, C., and Sinskey, A.J.

(2010) Elucidation of beta-oxidation pathways in Ralstonia eutropha H16 by

examination of global gene expression. *J Bacteriol* **192**: 5454-5464. doi:

10.1128/JB.00493–10.

Byloos, B., Monsieurs, P., Mysara, M., Leys, N., Boon, N., and Van Houdt, R. (2018)

Characterization of the bacterial communities on recent Icelandic volcanic deposits of

different ages. *BMC Microbiol* **18**: 122.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes, S.P.

(2016) DADA2: High-resolution sample inference from Illumina amplicon data. *Nat*

Methods **13**: 581–583.

Conrad, R. (1996) Soil microorganisms as controllers of atmospheric trace gases (H2, CO,

CH4, OCS, N2O, and NO). *Microbiol Rev* **60**: 609-640. doi: 10.1128/mmbr.60.4.609-

640.1996.

Conrad, R. and Seiler, W. (1980) Role of microorganisms in the consumption and production

of atmospheric carbon monoxide by soil. *Appl Environ Microbiol* **40**: 437-445. doi:

10.1128/aem.40.3.437-445.1980.

Conte, L., Szopa, S., Séférian, R., and Bopp, L. (2019) The oceanic cycle of carbon monoxide

and its emissions to the atmosphere. *Biogeosciences* **16**: 881-902. doi: 10.5194/bg-16-

881–2019.

Cordero, P.R.F., Bayly, K., Man Leung, P., Huang, C., Islam, Z.F., Schittenhelm, R.B., et al.

(2019) Atmospheric carbon monoxide oxidation is a widespread mechanism supporting

microbial survival. *ISME J* **13**: 2868–2881.

Crutzen, P.J. and Gidel, L.T. (1983) A two-dimensional photochemical model of the

atmosphere: 2. The tropospheric budgets of the anthropogenic chlorocarbons CO, CH4,

CH3Cl and the effect of various NO x sources on tropospheric ozone. *J Geophys Res*

Ocean **88**: 6641-6661. doi: 10.1029/JC088iC11p06641.

Cypionka, H. and Meyer, O. (1982) Influence of carbon monoxide on growth and respiration

of carboxydobacteria and other aerobic organisms. *FEMS Microbiol Lett* **15**: 209-214.

doi: 10.1016/0378-1097(82)90056–8.

Depoy, A.N., King, G.M., and Ohta, H. (2021) Anaerobic carbon monoxide uptake by

microbial communities in volcanic deposits at different stages of successional

development on o-yama volcano, Miyake-jima, Japan. *Microorganisms* **9**: doi:

10.3390/microorganisms9010012.

Dragone, N.B., Whittaker, K., Lord, O.M., Burke, E.A., Dufel, H., Hite, E., et al. (2023) The

Early Microbial Colonizers of a Short-Lived Volcanic Island in the Kingdom of Tonga.

MBio **14**: e0331322. doi: 10.1128/mbio.03313-22.

Dunfield, K.E. and King, G.M. (2004) Molecular analysis of carbon monoxide-oxidizing

bacteria associated with recent Hawaiian volcanic deposits. *Appl Environ Microbiol* **70**:

4242-4248. doi: 10.1128/AEM.70.7.4242-4248.2004.

Fierer, N., Nemergut, D., Knight, R., and Craine, J.M. (2010) Changes through time:

integrating microorganisms into the study of succession. *Res Microbiol* **161**: 635-642.

doi: 10.1016/j.resmic.2010.06.002.

Greening, C., Biswas, A., Carere, C.R., Jackson, C.J., Taylor, M.C., Stott, M.B., et al. (2016)

Genomic and metagenomic surveys of hydrogenase distribution indicate H2 is a widely

utilised energy source for microbial growth and survival. *ISME J* **10**: 761-777. doi:

10.1038/ismej.2015.153.

Guo, Y., Fujimura, R., Sato, Y., Suda, W., Kim, S., Oshima, K., et al. (2014) Characterization of

early microbial communities on volcanic deposits along a vegetation gradient on the

island of Miyake, Japan. *Microbes Environ* **29**: 38–49.

Hardy, K.R. and King, G.M. (2001) Enrichment of High-Affinity CO Oxidizers in Maine Forest

Soil. *Appl Environ Microbiol* **67**: 3671-3676. doi: 10.1128/AEM.67.8.3671-3676.2001.

Hernández, M., Calabi, M., Conrad, R., and Dumont, M.G. (2020) Analysis of the microbial

communities in soils of different ages following volcanic eruptions. *Pedosphere* **30**: 126-

134. doi: 10.1016/S1002-0160(19)60823–4.

Hernández, M., Vera-Gargallo, B., Calabi-Floody, M., King, G.M., Conrad, R., and Tebbe, C.C.

(2020) Reconstructing genomes of carbon monoxide oxidisers in volcanic deposits

including members of the class ktedonobacteria. *Microorganisms* **8**: doi:

10.3390/microorganisms8121880.

Inman, R.E., Ingersoll, R.B., and Levy, E.A. (1971) Soil: A natural sink for carbon monoxide.

Science (80-) **172**: 1229-1231. doi: 10.1126/science.172.3989.1229.

Islam, Z.F., Cordero, P.R.F., Feng, J., Chen, Y.J., Bay, S.K., Jirapanjawat, T., et al. (2019) Two

Chloroflexi classes independently evolved the ability to persist on atmospheric

hydrogen and carbon monoxide. *ISME J* **13**: 1801-1813. doi: 10.1038/s41396-019-

0393–0.

Islam, Z.F., Welsh, C., Bayly, K., Grinter, R., Southam, G., Gagen, E.J., and Greening, C. (2020)

A widely distributed hydrogenase oxidises atmospheric H2 during bacterial growth.

ISME J **14**: 2649-2658. doi: 10.1038/s41396-020-0713–4.

Johnson, J.L., Rajagopalan, K. V., and Meyer, O. (1990) Isolation and characterization of a

second molybdopterin dinucleotide: Molybdopterin cytosine dinucleotide. *Arch*

Biochem Biophys **283**: 542-545. doi: 10.1016/0003-9861(90)90681-N.

Kang, B.S. and Kim, Y.M. (1999) Cloning and molecular characterization of the genes for

carbon monoxide dehydrogenase and localization of molybdopterin, flavin adenine

dinucleotide, and iron-sulfur centers in the enzyme of Hydrogenophaga pseudoflava. *J*

Bacteriol **181**: 5581-5590. doi: 10.1128/jb.181.18.5581-5590.1999.

Kerby, R.L. and Roberts, G.P. (2012) *Burkholderia xenovorans* RcoMBx-1, a transcriptional

regulator system for sensing low and persistent levels of carbon monoxide. *J Bacteriol*

194: 5803–5816.

Kerby, R.L., Youn, H., and Roberts, G.P. (2008) RcoM: A new single-component

transcriptional regulator of CO metabolism in bacteria. *J Bacteriol* **190**: 3336–3343.

Khalil, M.A.K. and Rasmussen, R.A. (1990) The global cycle of carbon monoxide: Trends and

mass balance. *Chemosphere* **20**: 227-242. doi: 10.1016/0045-6535(90)90098-E.

Kiessling, M. and Meyer, O. (1982) Profitable oxidation of carbon monoxide or hydrogen

during heterotrophic growth of Pseudomonas carboxydoflava. *FEMS Microbiol Lett* **13**:

333-338. doi: 10.1111/j.1574-6968.1982.tb08283.x.

Kim, Y.J. and Kim, Y.M. (1989) Induction of carbon monoxide dehydrogenase during

heterotrophic growth of *Acinetobacter* sp. strain JC1 DSM 3803 in the presence of

carbon monoxide. *FEMS Microbiol Lett* **59**: 207–210.

King, C.E. and King, G.M. (2014) Description of Thermogemmatispora carboxidivorans sp.

nov., a carbon-monoxide oxidizing member of the class Ktedonobacteria isolated from

a geothermally heated biofilm, and analysis of carbon monoxide oxidation by members

of the class Ktedonobacter. *Int J Syst Evol Microbiol* **64**: 1244-1251. doi:

10.1099/ijs.0.059675–0.

King, C.E. and King, G.M. (2012) Temperature responses of carbon monoxide and hydrogen

uptake by vegetated and unvegetated volcanic cinders. *ISME J* **6**: 1558-1565. doi:

10.1038/ismej.2011.206.

King, G.M. (2003a) Contributions of atmospheric CO and hydrogen uptake to microbial

dynamics on recent Hawaiian volcanic deposits. *Appl Environ Microbiol* **69**: 4067-4075.

10.1128/AEM.69.7.4067-4075.2003.

King, G.M. (2003b) Molecular and Culture-Based Analyses of Aerobic Carbon Monoxide

Oxidizer Diversity. *Appl Environ Microbiol* **69**: 7257-7265. doi:

10.1128/AEM.69.12.7257-7265.2003.

King, G.M. (2003c) Uptake of Carbon Monoxide and Hydrogen at Environmentally Relevant

Concentrations by Mycobacteria. *Appl Environ Microbiol* **69**: 7266-7272. doi:

10.1128/AEM.69.12.7266-7272.2003.

King, G.M. and Crosby, H. (2002) Impacts of plant roots on soil CO cycling and soil-

atmosphere CO exchange. *Glob Chang Biol* **8**: 1085-1093. doi: 10.1046/j.1365-

2486.2002.00545.x.

King, G.M. and Weber, C.F. (2007) Distribution, diversity and ecology of aerobic CO-oxidizing

bacteria. *Nat Rev Microbiol* **5**: 107-118. doi: 10.1038/nrmicro1595.

King, G.M. and Weber, C.F. (2008) Interactions between bacterial carbon monoxide and

hydrogen consumption and plant development on recent volcanic deposits. *ISME J* **2**:

195-203. doi: 10.1038/ismej.2007.101.

Kumar, S., Stecher, G., and Tamura, K. (2016) MEGA7: molecular evolutionary genetics

analysis version 7.0 for bigger datasets. *Mol Biol Evoution* **33**: 1870-1874. doi:

10.1093/molbev/msw054.

Lane, D.J. (1991) 16S/23S rRNA sequencing. In *Nucleic Acid Techniques in Bacterial*

Systematics. Stackebrant, E. and Goodfellow, M. (eds). New York: John Wiley & Sons,

pp. 115-175. doi: 10.1002/jobm.3620310616.

Lubis, R.L., Juniarti, Rajmi, S.L., Armer, A.N., Hidayat, F.R., Zulhakim, H., et al. (2021)

Chemical properties of volcanic soil after 10 years of the eruption of Mt. Sinabung

(North Sumatera, Indonesia). *IOP Conf Ser Earth Environ Sci* **757**: doi: 10.1088/1755-

1315/757/1/012043.

Meyer, O., Frunzke, K., and Morsdorf, G. (1993) Biochemistry of the aerobic utilization of

carbon monoxide. In *Microbial growth on C1 compounds*. Murrell, J.C. and Kelly, D.P.

(eds). Andover, MA: Intercept Limited, pp. 433-459. doi: 10.1007/978-94-009-3539–6.

Meyer, O., Gremer, L., Ferner, R., Ferner, M., Dobbek, H., Gnida, M., et al. (2000) The role of

Se, Mo and Fe in the structure and function of carbon monoxide dehydrogenase. *Biol*

Chem **381**: 865-876. doi: 10.1515/BC.2000.108.

Meyer, O., Jacobitz, S., and Krüger, B. (1986) Biochemistry and physiology of aerobic carbon

monoxide-utilizing bacteria. *FEMS Microbiol Lett* **39**: 161-179. doi: 10.1016/0378-

1097(86)90445–3.

Meyer, O. and Schlegel, H.G. (1978) Reisolation of the carbon monoxide utilizing hydrogen

bacterium Pseudomonas carboxydovorans (Kistner) comb. nov. *Arch Microbiol* **118**: 35-

43. doi: 10.1007/BF00406071.

Mörsdorf, G., Frunzke, K., Gadkari, D., and Meyer, O. (1992) Microbial growth on carbon

monoxide. *Biodegradation* **3**: 61-82. doi: 10.1007/BF00189635.

Nam, G.H., Ahn, K., Bae, J.H., Cho, B.W., Park, K. Do, Lee, H.K., et al. (2003) Complete

nucleotide sequence of the circular megaplasmid pHCG3 of Oligotropha

carboxidovorans: function in the chemolithoautotrophic utilization of CO, H2 and CO2.

Gene **322**: 67-75. doi: 10.1016/j.gene.2012.04.005.

Neumann, M., Mittelstädt, G., Seduk, F., Iobbi-Nivol, C., and Leimkühler, S. (2009) MocA is a

specific cytidylyltransferase involved in molybdopterin cytosine dinucleotide

biosynthesis in *Escherichia coli*. *J Biol Chem* **284**: 21891–21898.

Ogiwara, K., Sekiguchi, M., Waranabe, M., Yoshida, M., and Ohta, H. (1999) Vegetation

recovery and soil microorganisms in volcanic mudflow deposits from Mt. Pinatubo, the

Philippines. In *Abstracts of the Annual Meeting, Japanese Society of Soil Science and*

Plant Nutrition. Vol. 45. p. 49.

Özkurt, E., Fritscher, J., Soranzo, N., Ng, D.Y.K., Davey, R.P., Bahram, M., and Hildebrand, F.

(2022) LotuS2: an ultrafast and highly accurate tool for amplicon sequencing analysis.

Microbiome **10**: 176.

Park, J.M., Kim, T.Y., and Lee, S.Y. (2011) Genome-scale reconstruction and in silico analysis

of the Ralstonia eutropha H16 for polyhydroxyalkanoate synthesis, lithoautotrophic

growth, and 2-methyl citric acid production. *BMC Syst Biol* **5**: doi: 10.1186/1752-0509-

5-101.

Pearcy, N., Garavaglia, M., Millat, T., Gilbert, J.P., Song, Y., Hartman, H., et al. (2022) A

genome-scale metabolic model of Cupriavidus necator H16 integrated with TraDIS and

transcriptomic data reveals metabolic insights for biotechnological applications. *PLoS*

Comput Biol **18**: e1010106. doi: 10.1371/journal.pcbi.1010106.

Quiza, L., Lalonde, I., Guertin, C., and Constant, P. (2014) Land-use influences the

distribution and activity of high affinity CO-oxidizing bacteria associated to type I-coxL

genotype in soil. *Front Microbiol* **5**: doi: 10.3389/fmicb.2014.00271.

Richter, M. and Rosselló-Móra, R. (2009) Shifting the genomic gold standard for the

prokaryotic species definition. *Proc Natl Acad Sci U S A* **106**: 19126-19131. doi:

10.1073/pnas.0906412106.

Rodriguez-R, L.M. and Konstantinidis, K.T. (2016) The enveomics collection: a toolbox for

specialized analyses of microbial genomes and metagenomes. *PeerJ Prepr* **4**: e1900v1.

doi: 10.7287/peerj.preprints.1900v1.

Romero, J.E., Alloway, B. V., Gutiérrez, R., Bertín, D., Castruccio, A., Villarosa, G., et al. (2021)

Centennial-scale eruptive diversity at Volcán Calbuco (41.3°S; Northwest Patagonia)

deduced from historic tephra cover-bed and dendrochronologic archives. *J Volcanol*

Geotherm Res **417**: doi: 10.1016/j.jvolgeores.2021.107281.

Saadi, D., Tirosh, E., and Schnell, I. (2021) The Relationship between City Size and Carbon

Monoxide (CO) Concentration and Their Effect on Heart Rate Variability (HRV). *Int J*

Environ Res Public Health **18**: doi: 10.3390/ijerph18020788.

Salvà-Serra, F., Donoso, R.A., Cho, K.H., Yoo, J.A., Lee, K., Yoon, S.-H., et al. (2021) Complete

Multipartite Genome Sequence of the *Cupriavidus basilensis* Type Strain, a 2,6-

Dichlorophenol-Degrading Bacterium. *Microbiol Resour Announc* **10**:.

Santiago, B., Schübel, U., Egelseer, C., and Meyer, O. (1999) Sequence analysis,

characterization and CO-specific transcription of the cox gene cluster on the

megaplasmid pHCG3 of Oligotropha carboxidovorans. *Gene* **236**: 115-124. doi:

10.1016/S0378-1119(99)00245–0.

Sato, Y., Nishihara, H., Yoshida, M., Ohta, H., Watanabe, M., and Rondal, J.D. (2004)

Occurrence of hydrogen-oxidizing *Ralstonia* species as primary microorganisms in the

Mt. Pinatubo volcanic mudflow deposits. *Soil Sci Plant Nutr* **50**: 855–861.

Sato, Y., Nishihara, H., Yoshida, M., Watanabe, M., Rondal, J.D., Concepcion, R.N., and Ohta,

H. (2006) *Cupriavidus pinatubonensis* sp. nov. and *Cupriavidus laharis* sp. nov., novel

hydrogen-oxidizing, facultatively chemolithothrophic bacteria isolated from volcanic

mudflow deposits from Mt. Pinatubo in the Philippines. *Int J Syst Evol Microbiol* **56**:

973-978. doi: 10.1099/ijs.0.63922–0.

Schubel, U., Kraut, M., Morsdorf, G., and Meyer, O. (1995) Molecular characterization of the

gene cluster coxMSL encoding the molybdenum-containing carbon monoxide

dehydrogenase of Oligotropha carboxidovorans. *J Bacteriol* **177**: 2197-2203. doi:

10.1128/jb.177.8.2197-2203.1995.

Schwartz, E., Henne, A., Cramm, R., Eitinger, T., Friedrich, B., and Gottschalk, G. (2003)

Complete nucleotide sequence of pHG1: A Ralstonia eutropha H16 megaplasmid

encoding key enzymes of H2-based lithoautotrophy and anaerobiosis. *J Mol Biol* **332**:

369-383. doi: 10.1016/S0022-2836(03)00894–5.

Seemann, T. (2014) Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**: 2068-

2069. doi: 10.1093/bioinformatics/btu153.

Siebert, D., Busche, T., Metz, A.Y., Smaili, M., Queck, B.A.W., Kalinowski, J., and Eikmanns,

B.J. (2020) Genetic Engineering of Oligotropha carboxidovorans Strain OM5-A

Promising Candidate for the Aerobic Utilization of Synthesis Gas. *ACS Synth Biol* **9**:

1426-1440. doi: 10.1021/acssynbio.0c00098.

Steinle, P., Stucki, G., Stettler, R., and Hanselmann, K.W. (1998) Aerobic mineralization of

2,6-dichlorophenol by *Ralstonia* sp. strain RK1. *Appl Environ Microbiol* **64**: 2566–2571.

Stott, M.B., Crowe, M.A., Mountain, B.W., Smirnova, A. V., Hou, S., Alam, M., and Dunfield,

P.F. (2008) Isolation of novel bacteria, including a candidate division, from geothermal

soils in New Zealand. *Environ Microbiol* **10**: 2030-2041. doi: 10.1111/j.1462-

2920.2008.01621.x.

Tindall, B.J., Rosselló-Móra, R., Busse, H.J., Ludwig, W., and Kämpfer, P. (2010) Notes on the

characterization of prokaryote strains for taxonomic purposes. *Int J Syst Evol Microbiol*

60: 249-266. doi: 10.1099/ijs.0.016949–0.

Vallenet, D., Calteau, A., Dubois, M., Amours, P., Bazin, A., Burlot, L., et al. (2019)

MicroScope: an integrated platform for the annotation and exploration of microbial

gene functions through genomic, pangenomic and metabolic comparative analysis.

Nucleic Acids Res **48**: D579–D589. doi; 10.1093/nar/gkz926.

Weber, C.F. and King, G.M. (2010a) Distribution and diversity of carbon monoxide-oxidizing

bacteria and bulk bacterial communities across a succession gradient on a Hawaiian

volcanic deposit. *Environ Microbiol* **12**: 1855-1867. doi: 10.1111/j.1462-

2920.2010.02190.x.

Weber, C.F. and King, G.M. (2007) Physiological, ecological, and phylogenetic

characterization of Stappia, a marine CO-oxidizing bacterial genus. *Appl Environ*

Microbiol **73**: 1266-1276. doi: 10.1128/AEM.01724–06.

Weber, C.F. and King, G.M. (2010b) Quantification of Burkholderia coxL Genes in Hawaiian

Volcanic Deposits. *Appl Environ Microbiol* **76**: 2212-2217. doi: 10.1128/aem.01861–09.

Weber, C.F. and King, G.M. (2012) The phylogenetic distribution and ecological role of

carbon monoxide oxidation in the genus Burkholderia. *FEMS Microbiol Ecol* **79**: 167-

175. doi: 10.1111/j.1574-6941.2011.01206.x.

Weber, C.F. and King, G.M. (2017) Volcanic soils as sources of novel CO-oxidizing

Paraburkholderia and Burkholderia: Paraburkholderia hiiakae sp. nov.,

Paraburkholderia metrosideri sp. nov., Paraburkholderia paradisi sp. nov.,

Paraburkholderia peleae sp. *Front Microbiol* **8**: doi: 10.3389/fmicb.2017.00207.

Wickham-Smith, C., Malys, N., and Winzer, K. (2023) Improving carbon monoxide tolerance

of Cupriavidus necator H16 through adaptive laboratory evolution. *Front Bioeng*

Biotechnol **11**: doi: 10.3389/fbioe.2023.1178536.

Wilcoxen, J. and Hille, R. (2013) The hydrogenase activity of the molybdenum/copper-

containing carbon monoxide dehydrogenase of Oligotropha carboxidovorans. *J Biol*

Chem **288**: 36052-36060. doi: 10.1074/jbc.M113.522441.

Wright, G.R., Jewczyk, S., Onrot, J., Tomlinson, P., and Shephard, R.J. (1975) Carbon

monoxide in the urban atmosphere: hazards to the pedestrian and the street-worker.

Arch Environ Health **30**: 123-129. doi: 10.1080/00039896.1975.10666660.

Wu, D., Raymond, J., Wu, M., Chatterji, S., Ren, Q., Graham, J.E., et al. (2009) Complete

genome sequence of the aerobic CO-oxidizing thermophile *Thermomicrobium roseum*.

PLoS One **4**: doi: 10.1371/journal.pone.0004207.

Figure legends

Figure 1. Relative abundance of 16S rRNA gene sequences in Calbuco stratifications originating from eruptions in 2015 and 1917. A) Relative abundance of 16S rRNA divided by phylum. B) Relative abundance of members of the Burkholderiaceae. Where the abundance of unclassified was lower than 0.8%, it was grouped into "Others".

Figure 2. Form-I CODH-encoding gene clusters from *Cupriavidus* sp. CV2^T and *Paraburkholderia terrae* COX. Translated amino acid identities (%) were calculated using BLASTp against homologous query sequences from the model type strain carboxydotroph *Oligotropha carboxidovorans* OM5. Asterisk (*) indicates a hypothetical gene.

Figure 3. Evolutionary relatedness of translated CoxL amino acid sequences. The tree was drawn using the Maximum Likelihood method with 500 Bootstrap replicates in MEGA7. MAGs were retrieved from Hernández *et al*. (2020).

Figure 4. Growth of, and CO consumption by, bacterial strains. When determining statistically significant consumption of CO during CO tolerance experiments, one-way ANOVA was used to compare all timepoints within a given concentration. A) Growth of *Cupriavidus* sp. CV2^T with 5 mM pyruvate combined with varying concentrations of headspace CO (0-100,000 ppmv) (OD$_{600}$). B) Headspace CO concentrations (ppm) during growth with 5 mM pyruvate by *Cupriavidus* sp. CV2^T, Y-axis values are presented on a Log$_{10}$ scale. C) Growth of *Pb. terrae* COX with 5 mM pyruvate combined with varying concentrations of headspace CO (0-100,000 ppmv) (OD$_{600}$). D) Headspace CO concentrations (ppmv) during growth with 5 mM pyruvate by *Pb. terrae* COX, Y-axis values are presented on a Log$_{10}$ scale. Bars represent mean values with standard deviation of independent triplicate incubations for each substrate.

Figure 5. Rates of CO uptake by (A) *Cupriavidus* sp. CV2^T and (B) *Pb. terrae* COX at stationary phase or exponential phase, varied according to the growth substrate. Bars represent mean values with standard deviation of independent triplicate incubations for each substrate. Different letters above the bars indicate significant differences determined by one-way ANOVA followed by a Tukey post hoc test.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5