Genome assembly and genomic architecture of a prominent cold-resistant rapeseed

germplasm

Runing title：Genome assembly of a cold-resistant rapeseed germplasm

Zefeng Wu[1], Guoqiang Zheng[1], Yali Sun[1], Xiaoyun Dong[1], Ying Wang[1], Hui Li[1], Jiaping Wei[1],

Junmei Cui[1], Yan Fang[1], Yinin Niu[1], Zhen Huang[2], Jihong Hu[2], Zigang Liu[1*]

[1] State Key Laboratory of Aridland Crop Science, Gansu Agricultural University, Lanzhou,

730070, Gansu, China

[2] State Key Laboratory of Crop Stress Biology for Arid Areas, College of Agronomy,

Northwest A&F University, Yangling, 712100, Shaanxi, China.

*Correspondence: lzgworking@163.com (Zigang Liu)

Graphical Abstract

The genomic landscape of cold-tolerant winter rapeseed (*Brassica napus*, L) has been poorly

characterized. We assembled a high-quality reference genome of a prominent cold-tolerant

winter rapeseed cultivar, NTS57, and performed phylogenetic and pan-genomic analyses by

integrating other reported *Brassica napus* accessions. The transcriptome analysis revealed that

microtubule-associated biological pathways were much more active in NTS57 under cold

stress than in the cold-sensitive variety. Whole genome methylation data analysis revealed

that DNA demethylation on protein-coding genes and repetitive elements, especially at CHH

sites, is essential for cold response in winter rapeseed.

To the editor，

Winter rapeseed is an important oilseed crop that is often subjected to low temperature stress

and consequent yield loss at high latitudes. Therefore, identification of highly cold-resistant

winter type rapeseed and elucidation of the genomic architecture are crucial for genetic

improvement of the cold-resistant trait. Here, we assembled a chromosome-level reference

genome of the prominent cold-resistant winter rapeseed cultivar NTS57 by integrating

Illumina, PacBio and Hi-C sequencing data. Phylogenetic and comparative genomic analyses

were performed to discover the relationships and genomic variations between NTS57 and

other rapeseed accessions. Transcriptomic analysis of NTS57 and a cold-sensitive cultivar

under cold stress identified > 10,000 differentially expressed genes that were enriched in

regulation of circadian rhythm, photosynthesis, and cold response. Many important molecular

components, including *DREBs*, *ZAT12*, *CAMTA3/5*, and *CORs*, in the classical cold signaling

pathway were also discovered. Importantly, genes or biological pathways associated with

microtubules were much more active in NTS57 under cold stress relative to the cold-sensitive

cultivar, underlying its prominent cold resistance. We also showed that methylation levels

were significantly decreased under cold stress, suggesting that demethylation may be required

for the cold stress response. The high-quality NTS57 rapeseed reference genome and other

omics data will be valuable genetic resources for understanding cold tolerance in rapeseed.

RESULTS AND DISCUSSIONS

Genome assembly and annotation of a cold tolerant rapeseed NTS57

Brassica napus NTS57 (2n = 19, AACC) is a cultivated rapeseed developed by crossing

winter rapeseed with *B. rapa*. It has strong cold resistance and can survive in the field at

−26°C in northwest China. To explore the genetic architecture that contributes to the cold

resistance of NTS57, we performed DNA sequencing and de novo assembly of the NTS57

genome by integrating 60 × PacBio high-fidelity (HiFi) long-read sequencing, 100× Hi-C

sequencing, and 30 × Illumina paired-end sequencing data. First, K-mer analysis of the

Illumina sequencing reads estimated the genome size of NTS57 to be 1.01–1.29 Gb (Figure

S1). Then, the PacBio reads were assembled and polished into a 1.06-Gb genome assembly

with contig N50 of 21.81 Mb, and > 93.6% (995 Mb) of the sequences were anchored on 19

chromosomes with scaffold N50 of 58.64 Mb after Hi-C scaffolding (Figure 1A, B).

Centromere sequences were successfully identified for all 19 NTS57 chromosomes (Figure

1B and Table S1). The high completeness of the genome assembly was evidenced by a

BUSCO recovery score of 99.75%. The assembly quality of the present sequenced genome is

higher than those of the published rapeseed reference genomes (Table S2). We identified

101,438 protein-coding genes in the NTS57 genome using ab initio prediction, homology

search, and transcriptome mapping strategies. Among them, 96,922 (95.5%) were located on

19 chromosomes; 44,411 on the A subgenome and 52,511 on the C subgenome, which is close

to the gene numbers in the Darmor-*bzh* and ZS11 genomes (Table S3). By homology search,

97.89% were annotated by at least one of the protein-related databases (Figure S2). In

addition, 36,552 non-coding RNAs were annotated in the NTS57 genome (Table S4).

Annotation of the NTS57 genome showed that > 55.6% of the genomic content was repetitive

elements (Table S5); 80% of them were TEs, including retrotransposons and DNA

transposons. The Gypsy and Copia represented the highest proportion of the retrotransposons

(Table S5). Genomic visualization showed that Gypsy and Copia had overall similar

distribution patterns along the chromosomes and an anti-correlation with gene density;

however, they displayed an opposite pattern near centromeres with high GC content and DNA

methylation level (Figure 1B), which is similar to the findings of a previous study [1].

Chromosomal synteny analysis based on protein-coding gene arrangements detected strong

intraspecific collinearity between homologous chromosomes from the two subgenomes

(Figure 1B), demonstrating the homology of chromosomes that were inherited from two

ancestral diploid genomes.

Phylogeny and pan-genomic analysis of rapeseed varieties

To determine the phylogeny of NTS57 with other rapeseed cultivars, we integrated nine

publicly sequenced rapeseed accessions with different ecotypes, namely three winter type

(Darmor-*bzh*, Quinta, and Tapidor), four semi-winter type (ZS11, Shengli, Zheyou, and

Gangan), and two spring-type (Westar and No2127) rapeseed cultivars. The maximum

likelihood tree constructed using 2,268 single-copy orthologous genes showed that the A

subgenome of NTS57 was closely related to the semi-winter rapeseed accessions, whereas the

C subgenome formed a clade with the winter type rapeseed accessions (Figure 1C). This

result indicated that NTS57 and semi-winter rapeseed bred in China had different breeding

histories, and the cold tolerance may have been inherited or retained from winter type

rapeseed. We then investigated the pan-genome architecture of rapeseed by integrating

NTS57 and nine other rapeseed accessions. By clustering protein-coding genes, we identified

43,734 and 54,091 pangene families from the A and C subgenomes, respectively. The number

of gene families identified in the A and C subgenomes increased rapidly when more

subgenomes or species were incorporated (Figure 1D), implying that the 10 rapeseed

genomes have rich gene pools and that a single reference genome cannot capture the full

genetic diversity of rapeseed. For the A subgenome, more than 18,375 gene families were

conserved among the 10 rapeseed accessions, and were considered to represent the core gene

families. Dispensable gene families that were present in 2–10 accessions accounted for 55.69%

of the pangene families, and 2.3% of the pangene families were categorized as genome

specific (Figure S3A). For the C subgenome, similar percentages of core, dispensable, and

genome-specific pangene families were identified, although there was a slightly lower

percentage of core gene families and a slightly higher percentage of dispensable gene families

(Figure S3B). Gene function enrichment analysis showed that the core gene families and the

indispensable gene families have different functions in rapeseed growth and development

(Figure S4).

The high-quality NTS57 reference genome allowed the SVs associated with cold

tolerance to be evaluated by comparative genomic analysis. A total of 51,511 insertions,

58,775 deletions, 29,645 translocations, 15,654 duplications, and 4,047 inversions were

identified in the nine rapeseed accessions with NTS57 as the reference genome (Figure 1E).

The artificially synthesized rapeseed line No2127 had the highest number of SVs, and Tapidor

had the smallest number of SVs. The total sizes of the SVs were 113.5–153.6 Mb in the

different accessions, with the length per SV ranging from a few tens of bp to hundreds of kb

or even above the Mb scale (Figure S5). We also observed that insertions and deletions tended

to be enriched at both chromosome ends and depleted near the centromeres, with a

distribution similar to that of gene density and opposite to that for GC content. Unlike the

deletions and insertions, the inversions, duplications, and translocations were randomly

distributed across the 19 chromosomes, although some enrichment was detected near

centromeres (Figure 1F and Figure S6). These findings were confirmed by calculating the

correlations between the chromosomal distributions of the different SVs (Figure 1F).

Gene dynamic expression and DNA methylation of NTS57 under cold stress

To identify genes associated with cold tolerance in NTS57, we analyzed the transcriptomic

data of plants under normal and cold stress conditions that had been generated in a previous

study [2]. Compared with the control, we identified 15,888 and 22,479 DEGs in the T1 (12h

after cold) and T2 (24h after cold) samples, respectively (Figure 2A and Table S6, S7). As

expected, the number of identified cold-responsive genes was higher at the longer time after

cold stress. We identified 12,346 DEGs that were common at both time points after cold stress

(Figure S7), suggesting that these genes may be core molecular components in NTS57 in

response to cold stress. We found that many DEGs were directly involved in plant cold

response or cold acclimation processes, including genes that encoded superoxide dismutase

(SOD) [3], VERNALIZATION INSENSITIVE 3 (VIN3) [4], transcription factor LUX [5],

RNA-binding protein CP29B [6], flavanone 3-dioxygenase 3 [7], and dehydration-responsive

element-binding protein 2E (DREB2E) [8]. Furthermore, many DEGs were annotated as cold-

responsive or cold-induced genes (Table S6,S7). We then performed GO and KEGG pathway

enrichment analysis. The DEGs in T1 were annotated mainly with regulation of circadian

rhythms, cold response, and xyloglucan metabolism terms under the biological process

category, MCM complex, cell wall, and microtubule under the cellular component category,

and UDP-glycosyltransferase, protein serine/threonine phosphatase activity, and ubiquitin

protein ligase activity under the molecular function category (Figure 2B). The DEGs in T2

were enriched with the same GO terms as the DEGs in T1, as well as photosynthesis and

monopolar cell growth under the biological process category (Figure S8), suggesting that

more biological processes are affected in the response to cold stress with longer stress

duration. The KEGG pathway enrichment results showed that the DEGs were involved

mainly in MAPK signaling pathway, circadian rhythm, flavonoid biosynthesis, and

photosynthesis, which is similar to the results of the GO enrichment analysis (Figure S9). To

identify potential genes and pathways specific to the prominent cold tolerance rapeseed

NTS57, we performed similar gene expression analysis on the cold-sensitive rapeseed NF

cultivar, which was found to have an extremely low overwintering survival rate based on

previous phenotype observation and physiological measurements [2]. Principal component

analysis showed that NTS57 and NF samples under cold stress were separated from each

other, suggesting that accession-specific transcriptomic changes occurred upon cold treatment

(Figure S10). The comparative transcriptomic analysis at the whole genome level showed that

there were high correlations in gene expression changes between the NTS57 and NF

accessions in both T1 ($R = 0.68$) and T2 ($R = 0.79$) samples (Figure S11). We identified

15,558 and 20,988 high-quality DEGs in the T1 and T2 samples of NF, respectively (Table S8,

S9), which is comparable with that in the NTS57. Besides, many DEGs in the NF were

overlapped with those identified in the NTS57 (Figure S12). Similar gene function

enrichments were also observed for the DEGs of NF derived from the two stress time points

(Figure 2B and Figures S8, S13), implying that these biological processes are essential or

conserved across rapeseed accessions under cold stress. Despite the high similarity of

transcriptomic changes between the two accessions, a number of genes were exclusively

expressed in NTS57 but not in NF in T1 or T2 samples (Figure S11), which may explain the

difference in cold tolerance between the two accessions. The enrichment analysis results

showed that genes involved in enriched microtubule-associated processes under the biological

process, cellular component, and molecular function GO categories were all specifically

altered in NTS57 upon cold stress (Figure S14). Gene expression profiles of microtubule-

associated genes also changed more dramatically in NTS57 than they did in NF under cold

stress, especially *A05G015370* (*TUBB7*), *A06G029560* (*TOR1L3*), and *C05G002130* (*TPX2*)

(Figure 2C). These finding indicate that microtubule-associated processes were more active in

NTS57 than they were in NF, and likely contributed to the cold tolerance in NTS57.

Cold stress has been reported to induce genomic demethylation in many plant species [9].

By calculating the genome-wide DNA methylation levels of the NTS57 genome, we found a

remarkable decrease in methylation levels across the gene body and flanking regions after

cold stress in all cytosine sequence contexts, namely CpG, CHG, and CHH (Figure 2D). In

particular, the decrease in CHH methylation was most pronounced, followed by CHG and

CpG methylation, implying that CHH methylation was more sensitive to cold stress. When

the A and C subgenomes were treated independently, similar trends were observed (Figure

S15), which is consistent with the findings of previous studies [10, 11]. In plants, DNA

methylation is enriched over heterochromatic TEs and repetitive elements [12]. As expected,

there was high enrichment of DNA methylation on repeat sequences, including long terminal

repeats, DNA transposons, and tandem repeats in the NTS57 genome (Figure 2E). However,

we noticed two exceptions for CHG and CHH methylation in tandem repeats, which showed

some deletions compared with the surrounding regions (Figure 2E). We also found that

methylation levels were decreased in all the repeat sequences and in all methylation contexts

in NTS57 under cold stress, which is very similar to that of the genic regions, suggesting that

methylase acts on genes and repeats in a similar way in the plant cold stress response.

AUTHOR CONTRIBUTIONS

Zefeng Wu: Conceptualization, Funding acquisition, Formal analysis, Methodology,

Visualization, Writing - original draft. **Guoqiang Zheng**: Conceptualization, Formal analysis,

Investigation. **Yali Sun:** Conceptualization, Formal analysis. **Xiaoyun Dong:** Visualization,

Software, Validation. **Ying Wang:** Software, Investigation. **Hui Li:** Investigation, Validation.

Jiaping Wei: Investigation, Validation. **Junmei Cui:** Visualization, Software. **Yan Fang:**

Investigation, Validation. **Yinin Niu:** Investigation, Validation. **Zhen Huang:** Investigation,

Writing – review & Editing. **Jihong Hu:** Conceptualization, Writing – review & Editing.

Zigang Liu: Conceptualization, Funding acquisition, Project administration, Supervision.

ACKNOWLEDGMENTS

This work was supported by the Research Program Sponsored by State Key Laboratory of

Aridland Crop Science, Gansu Agricultural University (GSCS-2023-03), the Scientific

Research Starting Foundation of Gansu Agricultural University (GAU-KYQD-2020-27), the

Natural Science Foundation of Gansu Province (22JR5RA862 and 23JRRA1421), the Gansu

CONFLICT OF INTEREST STATEMENT

The authors have declared no competing interests.

DATA AVAILABILITY STATEMENT

All the raw sequencing data and the genome assembly generated for this research are

available in the NCBI BioProject database under accession number PRJNA1012021. Genome

annotation files and other related documents are available in figshare

(https://doi.org/10.6084/m9.figshare.24098538.v1) or in the public database

(http://oilseed.online:8000/). The data and scripts used are saved in GitHub

https://github.com/Zefeng2018/NTS57_genome_assembly. Supplementary materials

(methods, figures, tables, graphical abstract, slides, videos, Chinese translated version and

update materials) may be found in the online DOI or iMeta Science

http://www.imeta.science/imetaomics/.

ETHICS STATEMENT

No animals or humans were involved in this study.

REFERENCES

1. Chen, Xuequn, Chaobo Tong, Xingtan Zhang, Aixia Song, Ming Hu, Wei Dong, Fei Chen, et al. 2021. "A high-quality Brassica napus genome reveals expansion of transposable elements, subgenome evolution and disease resistance." *Plant Biotechnol J* 19: 615-630. https://doi.org/10.1111/pbi.13493

2. Wei, Jiaping, Guoqiang Zheng, Xingwang Yu, Sushuang Liu, Xiaoyun Dong, Xiaodong Cao, Xinling Fang, et al. 2021. "Comparative Transcriptomics and Proteomics Analyses of Leaves Reveals a Freezing Stress-Responsive Molecular Network in Winter Rapeseed (*Brassica rapa* L.)." *Front Plant Sci* 12: 664311. https://doi.org/10.3389/fpls.2021.664311

3. Lin, Kuan-Hung, Sin-Ci Sei, Yu-Huei Su, Chih-Ming Chiang. 2019. "Overexpression of the Arabidopsis and winter squash superoxide dismutase genes enhances chilling tolerance via ABA-sensitive transcriptional regulation in transgenic Arabidopsis." *Plant Signal Behav* 14: 1685728. https://doi.org/10.1080/15592324.2019.1685728

4. Zhao, Yusheng, Rea L Antoniou-Kourounioti, Grant Calder, Caroline Dean, Martin Howard. 2020. "Temperature-dependent growth contributes to long-term cold sensing." *Nature* 583: 825-829. https://doi.org/10.1038/s41586-020-2485-4

5. Chow, Brenda Y, Sabrina E Sanchez, Ghislain Breton, Jose L Pruneda-Paz, Naden T Krogan, Steve A Kay. 2014. "Transcriptional regulation of LUX by CBF1 mediates cold input to the circadian clock in Arabidopsis." *Curr Biol* 24: 1518-1524. https://doi.org/10.1016/j.cub.2014.05.029

6. Kupsch, Christiane, Hannes Ruwe, Sandra Gusewski, Michael Tillich, Ian Small, Christian Schmitz-Linneweber. 2012. "Arabidopsis chloroplast RNA binding proteins CP31A and CP29A associate with large transcript pools and confer cold stress tolerance by influencing multiple chloroplast RNA processing steps." *Plant Cell* 24: 4266-4280. https://doi.org/10.1105/tpc.112.103002

7. Wu, Lishuang, Jian Tian, Yao Yu, Lixia Yuan, Yujiong Zhang, Hao Wu, Furong Wang, Xin Peng. 2023. "Functional characterization of a cold related flavanone 3-hydroxylase from Tetrastigma hemsleyanum: an in vitro, in silico and in vivo study." *Biotechnol Lett* https://doi.org/10.1007/s10529-023-03440-5

8. Agarwal, Pradeep K, Kapil Gupta, Sergiy Lopato, Parinita Agarwal. 2017. "Dehydration responsive element binding transcription factors and their applications for the engineering of stress tolerance." *J Exp Bot* 68: 2135-2148. https://doi.org/10.1093/jxb/erx118

9. Peng, Hai, Jing Zhang. 2009. "Plant genomic DNA methylation in response to stresses: Potential applications and challenges in plant breeding." *Progress in Natural Science* 19: 1037-1045. https://doi.org/10.1016/j.pnsc.2008.10.014

10. Bird, Kevin A, Chad E Niederhuth, Shujun Ou, Malia Gehan, J Chris Pires, Zhiyong Xiong, Robert VanBuren, Patrick P Edger. 2021. "Replaying the evolutionary tape to investigate subgenome dominance in allopolyploid Brassica napus." *New Phytol* 230: 354-371. https://doi.org/10.1111/nph.17137

11. Chalhoub, Boulos, France Denoeud, Shengyi Liu, Isobel A P Parkin, Haibao Tang, Xiyin Wang, Julien Chiquet, et al. 2014. "Early allopolyploid evolution in the post-Neolithic *Brassica napus* oilseed genome." *Science* 345: 950-953. https://doi.org/10.1126/science.1253435

12. Gallego-Bartolome, Javier. 2020. "DNA methylation in plants: mechanisms and tools for targeted manipulation." *New Phytol* 227: 38-44. https://doi.org/10.1111/nph.16529

SUPPORTING INFORMATION

The online version contains supplementary figures and tables available.

Figure S1. Genome size estimation based on survey sequencing.

Figure S2. Venn diagram of gene function annotations obtained using several protein-related

databases.

Figure S3. Percentage of core genes, dispensable genes, and accession-specific genes in the A

and C subgenomes of rapeseed.

Figure S4. Gene function enrichment for core and dispensable genes of NTS57.

Figure S5. Length statistics for structural variations (SVs) in different rapeseed accessions

using NTS57 as the reference genome.

Figure S6. Synteny and rearrangement plot for 19 chromosomes of NTS57 and ZS11.

Figure S7. Number of DEGs identified in NTS57 under cold stress.

Figure S8. Gene ontology (GO) functional enrichment for the DEGs identified in T2 samples

relative to those in the normal condition samples.

Figure S9. KEGG pathway enrichment for the DEGs identified in T1 or T2 samples relative

to those in normal condition (CK) samples.

Figure S10. Principal component analysis of gene expression profiles from accessions NTS57

and NF under cold stress.

Figure S11. Gene expression changes in NTS57 and NF under cold stress.

Figure S12. Veen diagram showing the intersection of the DEGs in NTS57 and NF in the T1

and T2 samples.

Figure S13. Gene functional enrichment of the DEGs in NF in the T1 and T2 samples.

314 **Figure legends**

Figure 1 Genome assembly of NTS57 and pangenome analysis of rapeseed. (A) Genome-wide

Hi-C heat map of the NTS57 genome. The diagonal pattern indicates strong Hi-C intra-

chromosomal interactions of the A and C subgenomes of NTS57. The color bar indicates the log-

transformed interaction frequencies of the Hi-C chromosome links. (B) Circos plot showing the

genome assembly and genomic features of NTS57. The letters (*a–i*) indicate the ideograms of 19

NTS57 chromosomes, gene density, GC content, Copia transposons, Gypsy transposons, CpG

methylation, CHG methylation, CHH methylation, and chromosomal collinearity, respectively.

The black rectangles in the ideograms indicate the centromeres. (C) Maximum likelihood tree

constructed using 2,268 single-copy orthologous genes shows the phylogenetic relationships of

NTS57 and other public rapeseed accessions. Bootstrap values inferred from 1000 replicates are

given as percentages on the branches of the consensus tree. (D) Modelling of pan-genome and

core genome sizes by including additional rapeseed accessions. (E) Number of different structural

variation (SV) types in the genome of each rapeseed accession using NTS57 as the reference

genome. DEL, DUP, INS, INV, and TRANS, deletion, duplication, insertion, inversion, and

translocation, respectively. (F) SV landscape in the NTS57 genome. The heat map shows the

correlations between the different SVs, GC content, and gene density (Gs). The correlation

significance is marked in the corresponding cells.

Figure 2 Gene dynamic expression and DNA methylation of NTS57 under cold stress. (A)

Sankey plot showing the up- and down-regulated genes of NTS57 after 12 h (T1) and 24 h (T2)

cold stress at −4°C. CK, normal conditions at 25°C. (B) Gene ontology (GO) function enrichment

of DEGs in the T1 samples. BP, biological process; CC, cellular component; MF, molecular

function. (C) Heat map of the expression change of the NTS57-specific DEGs involved in

microtubule process in accessions NTS57 and NF under cold stress. (D) Profiles of methylation

levels in the gene body, and upstream and downstream regions in the NTS57 genome under

normal and cold stress. CK, normal condition at 25°C; T1 and T2, cold stress at –4°C for 12 h and

24 h, respectively; TSS, transcription start site; TTS, transcription termination site; LTR, long

terminal repeat. The gene body was divided into 100 bins of equal size. For each bin in each gene,

the methylation level was calculated by dividing the total number of methylated Cs by the total

number of Cs, and the average of methylation level in the window across all protein-coding genes

was used to generate the profiles. The three methylation profiles for the CpG, CHG, and CHH

contexts were analyzed independently. (E) Profiles of methylation levels in the repetitive elements

of NTS57. CK, T1, and T2 are as defined in (D).