Genome assembly and genomic architecture of a prominent cold-resistant rapeseed germplasm

Zefeng Wu[1], Guoqiang Zheng[1], Yali Sun[1], Xiaoyun Dong[1], Ying Wang[1], Hui Li[1], Jiaping Wei[1], Junmei Cui[1], Yan Fang[1], Yinin Niu[1], Zhen Huang[2], Jihong Hu[2], Zigang Liu[1*]

[1]State Key Laboratory of Aridland Crop Science, Gansu Agricultural University, Lanzhou, 730070, Gansu, China

[2]State Key Laboratory of Crop Stress Biology for Arid Areas, College of Agronomy, Northwest A&F University, Yangling, 712100, Shaanxi, China.

[*]Corresponding author: lzgworking@163.com

Runing title：Genome assembly of a cold-resistant rapeseed germplasm

Abstract

Cold stress is one of the most limiting factors affecting crop production. Winter rapeseed, as an important oilseed crop, is often subjected to low temperature stress and consequent yield loss at high latitudes. Therefore, identification of highly cold-resistant winter type rapeseed and elucidation of the genomic architecture are crucial for genetic improvement of the trait. Here, we report a chromosome-level reference genome of a prominent cold-resistant winter rapeseed cultivar, NTS57. Phylogenetic analysis revealed that the A subgenome was more closely related to that of other semi-winter rapeseed accessions, while the C subgenome was in a clade with that of winter rapeseed accessions. Comparative genomic analyses revealed extensive structural variation between the NTS57 and other public rapeseed accessions, some of which are involved in plant temperature sensing or vernalization. Gene expression profiling under cold stress identified more than ten thousand differentially expressed genes which are enriched in the regulation of circadian rhythms, photosynthesis, cold response, etc. Gene co-expression analysis was performed and identified three functional modules which is associated with cold response in NTS57 rapeseed. DNA methylation profiling revealed that the methylation levels were significantly decreased at both protein-coding genes and repetitive elements, suggesting that demethylation is required for cold stress response. Overall, this research provided a basis for genetic dissection and improvement of cold tolerance in rapeseed.

Keywords

Cold stress; rapeseed; genome assembly; RNA-Seq; methylation.

1. Introduction

Cold stress, including chilling (0-15□) and freezing (< 0□), is one of environmental factors affecting plant growth, productivity, and geographical distributions of species (Ding et al., 2019). Most temperate plants have evolved the ability to adapt to low temperatures, referred to as cold acclimation, and show improved freezing tolerance after exposure to non-freezing or non-lethal low temperatures (Browse and Xin, 2001).

During cold acclimation or chilling stress, plants undergo a number of physiological and biochemical changes, including transcriptional, translational, and metabolic changes that increase or decrease the levels of specific proteins, metabolites, and phytohormones (Ding et al., 2019; Guo et al., 2018; Kidokoro et al., 2022). First, cold stress affects membrane rigidification, which is one of the most prominent physiological indicators and further triggers downstream cold stress responses in plants (Orvar et al., 2000). This situation results in the extravasation of cell sap and an increase in relative electrical conductivity (REC) (Lu et al., 2017). Thus, REC has become a commonly used index to assess membrane permeability and measure the effect of low temperature on plants (Cui et al., 2020; Guan et al., 2019; Yu et al., 2021). Second, chilling stress disrupts the stability of proteins or protein complexes and reduces the activities of enzymes such as ROS-scavenging enzymes (Ding et al., 2019; Yu et al., 2020). These physiological changes may be mirrored by biochemical responses. For example, many up-regulated genes during cold acclimation are involved in the synthesis of protective substances, such as soluble sugars and proline, and proteins including late embryogenesis abundant (LEA) proteins, anti-freezing proteins (AFP), and cold shock proteins (CSPs), while the down-regulated genes have functions in photosynthesis, such as genes encoding photosystem I subunits (*Psa*) and photosystem II subunits (*Psb*) (Kidokoro et al., 2022). Although much progress has been made in the last two decades in identifying the crucial molecular components involved in cold stress tolerance and in dissecting their regulatory mechanisms, we are still far from a good understanding of the molecular mechanism underlying cold signal perception and transduction in plants, especially for important economic crops.

Rapeseed (*Brassica napus* L.) is one of the most important economic crops in the world. There are three major ecotypes of rapeseed, including spring, winter, and semi-winter types, which are the result of evolution and long-term cultivation to adapt to different climates (Wei et al., 2017). Among them, winter-type rapeseed is one of the few crops that can survive frosty winters in temperate regions (Liu et al., 2022). The Yangtze River basin is the most important ecological region for rapeseed production in China. Winter- and semi-winter rapeseed are cultivated in the Yangtze

River basin, with seeds sown from late September to mid-October, which means that the growth and production of rapeseed are always threatened by the cold temperature (Ke et al., 2020). More importantly, in northern China, where winters are extremely cold, most winter rapeseed cultivars could not successfully overwinter until some highly cold-tolerant rapeseeds were bred. "NTS57" is currently the most cold-tolerant variety of winter rapeseed bred by our research group, as it can survive temperatures as low as -26 □ with an overwintering rate of over 90% without snow cover (Mi et al., 2021; Wei et al., 2021). Unraveling the molecular mechanisms of these highly cold-tolerant rapeseeds would not only help us better understand the cold stress adaptation of plants, but is also of great agricultural importance for the northward expansion of winter rapeseed in China and other countries.

Currently, the genomes of Darmor-*bzh* and ZS11, representatives of two different ecotypes of rapeseed, have been assembled and serve as the excellent reference for functional genomics study of rapeseed (Chalhoub et al., 2014; Chen et al., 2021; Sun et al., 2017). More recently, the germplasm of eight rapeseed varieties has also been de novo assembled, providing a pan-genomic view and a valuable genomic resource for rapeseed breeding (Song et al., 2020). However, none of these varieties has been recognized as a prominent cold-resistant germplasm, which hinders the improvement and promotion of cold-resistant varieties in cold regions. In the present study, we assembled a chromosome-level reference genome of winter rapeseed, NTS57. The genome assembly enabled us to explore the molecular mechanism of cold resistance through pan-genome comparison, phylogenetic, transcriptomic and epigenomic analyses. Our work provides a high-confidence framework for studying and breeding crops with strong cold tolerance.

2. Materials and Methods

2.1. Plant materials and DNA extraction for genome assembly

To perform genome sequencing and assembly of rapeseed NTS57, the seeds were successfully germinated, and plants were grown in the greenhouse of Gansu Agricultural University under the growth condition of 20□/18□ and 16/8 h days. Leaf

tissue was collected from 4-week-old plants, flash-frozen and stored at -70□. High

molecular weight genomic DNA was extracted using the CTAB method, followed by

purification using the Grandomics Genomic Kit for regular sequencing according to

the manufacturer's standard operating procedure. The DNA degradation and

contamination of the extracted DNA was checked on 0.75% agarose gels. DNA purity

was then detected using NanoDrop™ One UV-Vis spectrophotometer (Thermo Fisher

Scientific, USA), of which OD260/280 was between 1.8 and 2.0 and OD 260/230 is

between 2.0-2.2. Finally, the DNA concentration was further measured using Qubit®

3.0 Fluorometer (Invitrogen, USA).

2.2. PacBio HiFi library preparation and DNA sequencing

For long-read sequencing, the extracted DNA was sequenced on a PacBio Sequel II

platform (Pacific Biosciences, Menlo Park, CA, USA) using the circular consensus

sequencing (CCS) model from two SMRT cells using 15kb preparation solutions. The

main steps for library preparation are: gDNA shearing, DNA damage repair, end

repair and A-tailing, ligation with hairpin adapters from the SMRTbell Express

Template Prep Kit 2.0 (Pacific Biosciences), nuclease treatment of the SMRTbell

library with the SMRTbell Enzyme Cleanup Kit, size selection, and binding to

polymerase.

Briefly, a total amount of 8 μg DNA per sample was used for the DNA library

preparation. The genomic DNA sample was first sheared with g-TUBEs (Covaris,

USA) according to the expected size of the fragments for the library. Single strand

overhangs were removed, and DNA fragments were damage repaired, end repaired,

A-tailed, and ligated with the hairpin adaptor for PacBio sequencing. The library was

treated by nuclease with SMRTbell Enzyme Cleanup Kit and purified with AMPure

PB Beads. Target fragments were screened using PippinHT (Sage Science, USA). The

SMRTbell library was purified with AMPure PB Beads and Agilent 2100 Bioanalyzer

(Agilent Technologies, USA) was used to determine the size of the library fragments.

Finally, sequencing was performed on a PacBio Sequel II instrument using

Sequencing Primer V5 and Sequel II Binding Kit 2.2 at the Genome Center of

Grandomics (Wuhan, China). After sequencing, the resulting BAM files were

submitted to the pbccs program (https://ccs.how/) with default parameters, resulting in

65.16Gb HiFi pass reads with an average length of 15.6 kb.

137　2.3. Preparation of short reads libraries and DNA sequencing

Short-read libraries were prepared using the MGIEasy Universal DNA Library Prep

Kit V1.0 (CAT#1000005250, MGI) according to the standard protocol. Briefly, 1 μg

of genomic DNA was randomly fragmented by Covaris. The fragmented DNA was

selected to an average size of 200-400 bp using MGIEasy DNA Clean Beads

(CAT#1000005279, MGI). The selected fragments were then end-repaired, 3'

adenylated and then ligated with adapters. The DNA samples were amplified by PCR

and the products were purified using MGIEasy DNA Clean Beads (CAT#1000005279,

MGI). The double-stranded PCR products were heat denatured and circularized using

the splint oligo sequence in the MGIEasy Circularization Module (CAT#1000005260,

MGI). The single-strand circle DNA (ssCir DNA) was formatted as the final library

and qualified by QC. The qualified libraries were sequenced on DNBSEQ-T7RS

platform.

150　2.4. Transcriptome library preparation and sequencing

Different tissues at different developmental stages were sampled to perform RNA-Seq

with three biological replicates (Supplementary Table S9). Briefly, total RNA was

extracted for each sample by grinding the tissue in TRIzol reagent (TIANGEN) on

dry ice and processed according to the manufacturer's protocol. Poly-A RNAs were

enriched from total RNA using Dynabeads mRNA Purification Kit (Cat. No. 61006,

Invitrogen) and fragmented into small pieces using fragmentation reagent in MGIEasy

RNA Library Prep Kit V3.1 (Cat# 1000005276, MGI). First-strand cDNA was

synthesized using random primers and reverse transcriptase, followed by

second-strand cDNA synthesis. The synthesized cDNA was end-repaired, A-tailed and

ligated to the sequencing adapters according to the library construction protocol. The

cDNA fragments were amplified by PCR and purified using MGIEasy DNA Clean

Beads (CAT#1000005279, MGI). The library was analyzed on the Agilent

Technologies 2100 Bioanalyzer. The double-stranded PCR products were heat

denatured and circularized using the splint oligo sequence in the MGIEasy

Circularization Module (CAT#1000005260, MGI). The single-strand circle DNA

(ssCir DNA) were formatted as the final library. The qualified libraries were

sequenced on DNBSEQ-T7RS platform.

2.5. Hi-C library preparation and sequencing

Freshly harvested leaves from the 4-week-old plants were cut into 2 cm pieces and

vacuum infiltrated in nuclear isolation buffer supplemented with 2% formaldehyde.

Cross-linking was stopped by the addition of glycine and additional vacuum

infiltration. The fixed tissue was then ground to powder and resuspended in nuclear

isolation buffer to obtain a nuclear suspension. The purified nuclei were digested with

100 units of DpnII and labeled by incubation with biotin-14-dCTP. Biotin-14-dCTP

was removed from unligated DNA ends due to the exonuclease activity of T4 DNA

polymerase. The ligated DNA was sheared into 300-600 bp fragments, and then was

blunt-end repaired and A-tailed, followed by purification by

biotin-streptavidin-mediated pull-down. Finally, the Hi-C libraries were quantified

and sequenced using the MGI-2000 platform.

2.6. Genome size and heterozygosity estimation

To understand the genomic characteristics of the NTS57, the k-mer analysis was

performed using Illumina DNA data prior to genome assembly to estimate the genome

size and heterozygosity. Briefly, quality-filtered reads were subjected to 25-mer

frequency distribution analysis using the KMC program (Kokot et al., 2017). By

analyzing the 25-mer depth distribution from the clean reads using FindGSE (Sun et

al., 2018) and GenomeScope2.0 (Ranallo-Benavidez et al., 2020), we estimated the

genome size using the following equation: G = K-num/K-depth (where K-num is the

total number of 25-mers, K-depth is the k-mer depth, and G is the genome size).

Finally, the heterozygosity and repeat content of the NTS57 genome were estimated

by combining the simulation data results of Arabidopsis with different heterozygosity

and the frequency peak distribution of 25-mers.

2.7. Genome denovo assembly and completeness assessment

The high quality HiFi reads were first used to assemble the initial contigs using the Hifiasm (v.0.13-r308) package with default parameters (Cheng et al., 2021). To improve the accuracy of the assembly, the assembled contigs were polished with Nextpolish (Hu et al., 2020) using short reads with default parameters. Finally, redundancies were removed using Haplomerger2 (Huang et al., 2017) to generate the primary contigs.

Next, Hi-C reads were used to support the assembly of the genome at the chromosome level. For this, the low-quality Hi-C reads (quality scores<20), adaptor sequences, and sequences shorter than 30 bp were filtered out using fastp (Chen et al., 2018), and then the clean paired-end reads were mapped to the assembled contig sequences using bowtie2 (v2.3.2) (Langmead and Salzberg, 2012) (-end-to-end --very-sensitive -L 30) to obtain the uniquely mapped paired-end reads. Valid interaction paired reads were identified and rerained by HiC-Pro (v2.8.1) (Servant et al., 2015) from the unique mapped paired-end reads for further analysis. Invalid read pairs, including dangling-end, self-cycle, re-ligation, and dumped products were filtered. Scaffolds were further clustered, ordered, and oriented on chromosomes using LACHESIS (Burton et al., 2013) with parameters CLUSTER_MIN_RE_SITES=100, CLUSTER_MAX_LINK_DENSITY=2.5, CLUSTER NONINFORMATIVE RATIO = 1.4, ORDER MIN N RES IN TRUNK=60, ORDER MIN N RES IN SHREDS=60. Finally, placement and orientation errors with obvious discrete chromatin interaction patterns were manually adjusted.

The completeness of the genome assembly was evaluated using BUSCO v4.0.5 (Benchmarking Universal Single Copy Orthologs) and CEGMA (Core Eukaryotic Gene Mapping Approach) (Parra et al., 2007; Simao et al., 2015). To assess the accuracy of the assembly, all the Illumina paired-end reads were mapped to the assembled genome using BWA software (Li and Durbin, 2009), and the mapping rate and genome coverage of the sequencing reads were assessed using SAMtools v0.1.1855 (Li et al., 2009). Furthermore, the base accuracy of the assembly was

calculated using bcftools (Danecek and McCarthy, 2017). In addition, the coverage of

potentially expressed genes of the assembly was examined by aligning RNA-Seq data

to the assembly using HISAT2 with default parameters (Kim et al., 2019). To avoid

including mitochondrial sequences in the assembly, the draft genome assembly was

submitted to the NCBI (NT) library and the aligned sequences were eliminated.

2.8. Repetitive DNA elements and gene model annotation

Tandem repeats were first annotated using the software GMATA (Wang and Wang,

2016) and Tandem Repeats Finder (TRF) (Benson, 1999), where GMATA identifies

the simple sequence repeats (SSRs) and TRF recognizes all tandem repeat elements in

the entire genome. Transposable elements (TEs) in the NTS57 genome were then

identified using a combination of de novo and homology-based strategies. For de

novo prediction, three de novo prediction programs, MITE-hunter (Han and Wessler,

2010), RepeatModeler (Flynn et al., 2020), and LTR_retriever (Ou and Jiang, 2018),

were used with default parameters. The results from these programs were integrated

into a de novo database, which was then aligned to Repbase

(http://www.girinst.org/repbase) and classified into different types. Using the resulting

database, repetitive sequences were identified by homologous sequence search using

RepeatMasker with default parameters (Tarailo-Graovac and Chen, 2009).

Overlapping repeat elements belonging to the same repeat class were merged as a

final non-redundant repeat sequence annotation. For centromere sequences

identification, the quarTeT program was used with the NTS57 genome sequences as

the input with default parameters (Lin et al., 2023). The top candidates were selected

based on the length and periods of repetitive sequences. Meanwhile, the gene density

and LTR density distribution across chromosomes were also used to help to determine

the centromere positions in chromosomes.

For gene model prediction, three independent approaches, including ab initio

prediction, homology search, and reference-guided transcriptome assembly, were used

to predict genes in the repeat-masked genome. Specially, GeMoMa (Keilwagen et al.,

2019) was used to align the homologous peptides from related species to the assembly

and then obtain the gene structure information, which was the homology prediction.

For RNA-Seq-based gene prediction, filtered RNA-Seq reads were mapped to the

reference genome using STAR with default parameters (Dobin et al., 2013). The

transcripts were then assembled using stringtie (Pertea et al., 2015) and open reading

frames   (ORFs)   were   predicted   using   PASA

(https://github.com/PASApipeline/PASApipeline). For the de novo prediction,

RNA-Seq reads were de novo assembled using stringtie (Pertea et al., 2015) and

analyzed using PASA to generate a training set. Augustus with default parameters

(Stanke et al., 2004) was then used for ab initio gene prediction with the training set.

Finally, EVidenceModeler (EVM) (Haas et al., 2008) was used to generate an

integrated gene set from which genes with TE were removed using the TransposonPSI

package (http://transposonpsi.sourceforge.net/) and the miscoded genes were further

filtered. Untranslated regions (UTRs) and alternative splicing regions were

determined using PASA based on RNA-seq assemblies. We retained the longest

transcripts for each locus, and regions outside of the ORFs were designated as UTRs.

2.9. Functional annotation of gene models

Functional information of genes, motifs and domains of their proteins were assigned

by comparison with public databases including Swiss-Prot (Boutet et al., 2007),

nonredundant protein database NR (Pruitt et al., 2007), KEGG (Kanehisa and Goto,

2000), KOG (Tatusov et al., 2001) and Gene Ontology (Gene Ontology, 2015). The

putative domains and GO terms of genes were identified using the InterProScan

program with default parameters (Quevillon et al., 2005). For the other four databases,

BLASTp (Altschul et al., 1990) was used to compare the EvidenceModeler-integrated

protein sequences against the four well-known public protein databases with an

E-value cutoff of 1e-05, and the results with the hit with the lowest E-value were

retained. The results from the above five database searches were concatenated.

2.10. Annotation of non-coding RNAs (ncRNAs)

Two strategies were used to obtain the ncRNA (non-coding RNA): database search

and model prediction. Transfer RNAs (tRNAs) were predicted using tRNAscan-SE

with eukaryote parameters (Chan et al., 2021). MicroRNA, rRNA, small nuclear RNA, and small nucleolar RNA were detected using Infernal cmscan to search the Rfam database (Nawrocki and Eddy, 2013). The rRNAs and their subunits were predicted using RNAmmer (Lagesen et al., 2007).

2.11. Synteny analysis between NTFS57 and other rapeseeds

MCscan (Python version) (Tang et al., 2008) was used to identify collinear blocks of at least of 30 genes in NTS57 and other rapeseed genomes.

2.12. Phylogenetic analysis

To determine the phylogenetic relationship between NTS57 and other published rapeseed cultivars, the phylogenetic tree was reconstructed using 10 rapeseed species and two ancestral reference genomes, *B. rapa* Chiifu (v4.0) and *B. oleracea*, with *A. thaliana* as the outgroup species. For this, the single-copy orthogroups of the above 13 species were firstly identified using amino acid sequences by OrthoFinder v2.0.9 with the parameter "-M msa" (Emms and Kelly, 2019). Then, the multiple sequence alignment in each orthogroup was performed using mafft v7.453 (Katoh and Standley, 2013), and the resulting alignments were concatenated into a superalignment. Finally, the phylogenetic tree was determined by the maximum likelihood (ML) method as implemented in IQ-TREE v1.6.9 with 1000 bootstrap tests (Nguyen et al., 2015), and the best-fit model was determined as being JTT+F+R4 by ModelFinder with free rate heterogeneity (Kalyaanamoorthy et al., 2017).

2.13. Gene expression analysis

For both the newly generated RNA-seq datasets and published RNA-seq datasets, Trimmomatic (v0.36) (Bolger et al., 2014) with default parameters was used to remove adapters and low-quality reads from the raw RNA-seq reads. The clean RNA-seq reads were mapped to the corresponding reference genome using Hisat2 with default settings (Kim et al., 2019). Gene expression abundance was estimated using FPKM (fragments per kilobase of exon per million fragments calculated by StringTie (v2.1.6) (Pertea et al., 2015) for each transcript. The gene co-expression

network was constructed using the R package WGCNA (Langfelder and Horvath,

2008) by using 51 RNA-Seq samples from different tissues and developmental stages

as well as cold stress treatment.

2.14. Analysis of whole-genome bisulfite sequencing data

Whole genome methylation data for leaf tissues of NTS57 under normal or low

temperature (-4□) were reanalyzed from our previous study (NCBI accession:

PRJNA685002) (Wei et al., 2022). Briefly, fastp was first used to perform quality

filtering, FastQC v.0.11.7 (Wingett and Andrews, 2018) was used for quality control

and Bismark (v.0.23.0) (Krueger and Andrews, 2011) was used to align the trimmed

reads to our assembled reference genome of NTS57. The DNA methylation levels for

individual CpGs, CHG and CHH were calculated separately using methyKit (v.1.22.0)

(Akalin et al., 2012), and we retained only cytosine sites with ≥10 covered sequencing

reads. Methylated reads (containing Cs) and unmethylated reads (containing Ts) at

each cytosine site were counted, and the percentage of methylated reads among the

total reads covering the corresponding cytosine was calculated to quantify the DNA

methylome for each sample at the single base resolution. To identify the DMRs, we

used a window-based method to calculate the overall methylation level between

samples for comparisons. For this purpose, 1,000 bp continuous windows were

generated across 19 chromosomes of the NTS57 genome, and the windows covering

no less than 10 reads that showed ≥25% methylation level difference with q-value

≤0.01 were considered as DMRs using methyKit (v.1.22.0) (Akalin et al., 2012).

To compare the overall methylation level for different genomic features, such as

upstream, downstream, the gene body of protein-coding genes, and repeat elements,

the mean CpG, CHG, and CHH methylation levels for 2 kb upstream and downstream

of each type of the genomic features were binned into 100 equal-length windows, and

the mean methylation levels in each window were calculated using a custom R script.

2.15. SV identification

For SV calling, minimap2 (v.2.21-r1071) (Li, 2018) was used to generate pairwise

genome alignments between each of the nine rapeseed accessions and the NTS57

genome using the parameters "-ax asm5 -eqx". The alignment results were then passed to SyRI (Goel et al., 2019) and SVs consisting of insertions, deletions, inversions (<1 Mb in size), translocations (>50 bp in length) and duplications were retained. SVs containing 'N' sequences and the ambitious alignment margins and/or poor synteny alignment around the breakpoint were filtered. SVs between different genomes were visualized using plotsr (Goel and Schneeberger, 2022).

3. Results

3.1. Genome assembly and annotation of a cold tolerant rapeseed NTS57

B. napus NTS57 (2n=19, AACC) is a cultivated rapeseed developed by crossing winter rapeseed with *Brassica rapa*. It has strong cold resistance that can survive in the field at -26 ☐ in northwest China. For example, NTS57 plants grew and developed well under normal conditions, similar to ZS11 and Westar (Figure 1a). However, leaves of

the ZS11 and Westar accessions began to wilt after 12 hours at -4 ☐, while NTS57 appeared normal and healthy (Figure 1a). At 24 h after treatment, NTS57 was still growing better than the other two accessions, and only some of its leaves were curled or wilted (Figure 1a). When all plants were transferred to normal temperature and left for a period of time, NTS57 could fully recover its phenotype, while the other two accessions could not or only partially recover (Figure 1a). Apparently, NTS57 can survive cold stress and will be an important germplasm for the cold tolerance research in rapeseed.

To explore the genetic architecture contributing to the cold resistance trait of NTS57, we performed DNA sequencing and de novo assembly of the NTS57 genome by integrating 60x PacBio high-fidelity (HiFi) long-read sequencing, 100x Hi-C (high-throughput chromosome conformation capture) sequencing, and 30x Illumina paired-end sequencing data. First, K-mer analysis of the Illumina sequencing reads estimated the genome size of NTS57 to be *c.* 1.01~1.29 Gb (Figure S1). The PacBio reads were then assembled and polished into a 1.06 Gb genome assembly with contig N50 21.81 Mb, and more than 93.6% (995 Mb) of the sequences were anchored on 19

chromosomes with scaffold N50 58.64 Mb after Hi-C scaffolding (Figure 1b and c).

Centromere sequences were successfully identified for all 19 chromosomes of NTS57

(Figure 1c; Table S1). The high completeness of the assembly was also evidenced by

a BUSCO recovery score of 99.75%. Accordingly, the assembly quality of the present

sequenced genome is higher than the published rapeseed reference genomes (Table 1).

Based on ab initio prediction, homology search and transcriptome mapping strategies,

we obtained a total of 101,438 protein-coding genes in the NTS57 genome. Of the

gene set, 96,922 (95.5%) were located on 19 chromosomes, with 44,411 in the A

subgenome and 52,511 in the C subgenome, which was close to the gene number in

Darmor-*bzh* and ZS11 genomes (Table S2). Among them, we identified 24,217

homologous gene pairs based on reciprocal blast search, and the gene expression

analysis revealed that 6,994 (28.9%) gene pairs on the homologous chromosomes

showed at least a twofold gene expression difference (Figure S2). Although there

were 936 more gene pairs where the C-homeolog was more highly expressed than the

A-homeolog, no evidence for remarkable subgenome dominance was observed

considering the overall strong gene expression correlations (PCC = 0.81; *P*-value <

2.2e-16) between two subgenomes (Figure S2).

Of the 101,438 predicted genes, more than 78% were supported by

transcriptomic data. By homology search, 97.89% were well annotated by at least one

of the following protein-related databases: NCBI nr database, the Swiss-Prot protein

database, the KOG database, KEGG database, and the GO database (Figure S3). In

addition, 36,552 non-coding RNAs, including ribosomal RNAs (rRNAs), transfer

RNAs (tRNAs), microRNAs (miRNAs) and small nuclear RNAs (snRNAs), were

annotated in the NTS57 genome (Table S3). These results indicated the high quality

of the NTS57 genome assembly.

The amplification of various repetitive DNA elements is one of the key drivers of

plants evolution. Annotation of the NTS57 genome revealed that more than 55.6% of

the genomic content was repetitive elements (Table S4). Among these DNA repeats,

80% were TEs, including retrotransposons and DNA transposons. The Gypsy and

Copia represented the highest proportion of the retrotransposons, accounting for 8.6%

and 5.1% of the whole genome, respectively (Table S4). Genomic visualization analysis revealed that the two types of transposons had overall similar distribution patterns along the chromosomes and an anti-correlation with gene density (Figure. 1b). However, we noticed that they displayed an opposite pattern at some local positions, especially near centromeres with high GC content and DNA methylation level (Figure 1c), which is similar to a previous study on the ZS11 genome (Chen et al., 2021). In addition, chromosomal synteny analysis based on protein-coding gene arrangements revealed that there is strong intraspecific collinearity between homologous chromosomes from two subgenomes (Figure 1c), further demonstrating the homology of chromosomes inherited from two ancestral diploid genomes.

3.2. Phylogeny and pan-genomic analysis of rapeseed varieties

To reveal the phylogeny of NTS57 with other rapeseed cultivars, we integrated an additional 10 publicly sequenced rapeseeds with different eco-types, including three winter-type (Darmor-*bzh*, Quinta, and Tapidor), four semi-winter-type (ZS11, Shengli, Zheyou, and Gangan), and two spring-type (Westar and No2127) rapeseed cultivars. To reduce the clustering bias of the orthologous genes derived from two subgenomes, all the investigated rapeseed genomes were split into A and C subgenomes, resulting in 22 subgenomes for further analysis. In addition, Arabidopsis and two diploid progenitors, *Brassica rapa* and *Brassica oleracea*, were added as outgroup species to infer the species tree. The max-likelihood species tree inferred based on 2,268 single-copy orthologous genes revealed that the A subgenome of NTS57 were closely related to that of semi-winter rapeseed, while the C subgenome formed a clade with that of winter-type rapeseed (Figure 2a). This result indicated that the NTS57 and semi-winter rapeseed bred in China had a different breeding history, and the cold tolerance may inherit or kept from winter type rapeseeds.

While research on pan-genomes for rapeseed has been reported (Song et al., 2020), there is currently limited research on the cold tolerance of rapeseed at the pan-genome level. To this end, we investigated the pan-genome architecture by integrating NTS57 and 10 other rapeseed accessions. We identified a total of 43,734

pangene families by clustering protein-coding genes from the A sub-genomes and 54,091 from the C sub-genomes. The number of gene families in both A subgenome and C subgenome increased rapidly when more subgenomes or species were added (Figure 2b), suggesting that the 11 rapeseed genomes have rich gene pools and that a single reference genome cannot capture the full genetic diversity of rapeseed. For the A subgenome, more than 18,375 gene families were conserved among the 11 rapeseed accessions, representing the core gene families. The dispensable gene families that present in two to 10 accessions occupied 55.69% of the pan-gene families, and 2.3% of the pan-gene families were categorized as genome-specific types (Figure 2c). For the C subgenome, a similar percentage of three kinds of pangene families was observed, although there was a slightly lower percentage of core gene families and a slightly higher percentage of dispensable gene families (Figure 2d).

Gene function enrichment analysis revealed that the core gene families were mainly involved in cytokinin-activated signaling pathway, glucose metabolism process, response to bacterium, mismatch repair, phosphorelay signal transduction system, and intracellular protein transport; while the indispensable gene families were associated with phosphate ion transport, pollen recognition, cell surface receptor signaling pathway, positive regulation of ubiquitin protein ligase activity, regulation of gene expression, telomere maintenance and DNA integration (Figure S4), suggesting the different functions of the two classes of genes in rapeseed growth and development. However, there were no enriched GO terms with the same significance level for the species-specific gene families. Gene expression analysis on NTS57 revealed that the expression levels of core genes were significantly higher than those of dispensable genes in different tissues, and the specific genes had the lowest gene expression levels. (Figure S5, $P < 2.2 \times 10^{-16}$, Wilcoxon rank-sum test), reflecting the functional importance of the three groups of genes.

3.3. Extensive structural variations between NTS57 and other rapeseed varieties

The specific characteristics of NTS57 prompted us to further explore structural variations (SVs) at the genomic level with other rapeseed genomes. For this, purpose,

we performed pairwise genome alignments between each of the nine rapeseed

genomes and the NTS57 genome. We identified a total of 51,511 insertions, 58,775

deletions, 29,645 translocations, 15,654 duplications, and 4,047 inversions in the nine

rapeseed accessions using NTS57 as the reference genome (Figure 3a). Among these

different genomes, the artificially synthesized rapeseed line No2127 had the most SVs

with the NTS57 reference genome, while the Tapidor had the least SVs. The total size

of the SVs varied between 113.5 Mb and 153.6 Mb in the different accessions

compared, with the length per SV ranging from a few tens of bp to hundreds of kb or

even above the Mb scale (Figure S6). We found highly consistent patterns when using

an alternative reference genome, ZS11, as the reference (Figure S7 and Figure S8).

We also observed that insertions and deletions tended to be enriched at both

chromosome ends and depleted near centromeres, with a distribution similar to gene

density and opposite to GC content. In contrast to deletions and insertions, inversions,

duplications and translocations appeared to be randomly distributed across the 19

chromosomes, although some enrichment was observed near centromeres (Figure 3b

and Figure S9). These observations were also confirmed by calculating correlations

between the chromosomal distributions of different SVs (Figure 3b).

Numerous studies have shown that SVs play an important role in plant growth

and development (Alonge et al., 2020; Cai et al., 2021; Yuan et al., 2021). To reveal

the potential functional influence of SVs in rapeseed, we mapped these SVs to NTS57

reference genes based on chromosomal positions and found that 39.2.6% (39,820) of

protein-coding genes overlapped at least one SV across different species. GO analysis

of these genes within SV hotspot regions revealed enrichment for peptidase activity,

cysteine-type endopeptidase activity, and DNA integration, suggesting an important

role of SVs in adaptation of the investigated rapeseed to abiotic stresses.

In addition to the SVs identified at the level of genomic fragments, the potential

SVs based on the genome collinearity analysis at the level of protein-coding genes

were also investigated. Overall, there are strong collinearity relationships between

NTS57 and other rapeseed varieties, which do not reflect a large variation in

chromosomal structure among different rapeseeds. However, we found a

chromosomal fragment translocation from chromosome A03 to C03 in the NTS57

genome (Fig 3C). This translocation could not be a mis-assembly and this can be

confirmed by chromosomal interaction and long read sequencing data analysis (Figure

S10). Further inspection revealed that there are 1,283 protein-coding genes are located

in this region. Among these genes, many are involved in plant temperature sensing or

vernalization, including *VERNALIZATION INSENSITIVE 3* (*VIN3*) (Zhao et al., 2020),

Temperature-induced lipocalin-1(*TIL-1*) (Huang et al., 2021), *Heat shock 70 kDa*

protein 10 (*HSP70-10*), *HSP90-2* (Jahan et al., 2023), and *Protein HEAT*

INTOLERANT 4 (*HIT4*) (Probst and Mittelsten Scheid, 2015), *etc*. (Table S5).

3.4. Gene expression of cold-stress regulated genes in NTS57

To identify the genes associated with cold tolerance in NTS57, we performed

transcriptomic data analysis under normal and cold stress conditions that had been

generated in a previous study (Wei et al., 2021). For this, the leaf RNA-Seq data from

normal condition (CK), 12 h (T1), and 24 h (T2) after 4□ were mapped onto the

NTS57 reference genome and gene expression levels of all protein-coding genes were

calculated. Compared with the normal condition, we identified 15,888 and 22,479

differentially expressed genes (DEGs) in T1 and T2, respectively (Figure 4a, Table S6

and S7). Obviously, the number of identified cold-responsive genes increased with

increasing duration of cold stress. These DEGs were highly reproducible by using

different statistical analysis programs (Figure S11). Among these DEGs, there are

12,346 common DEGs at both time points after cold stress (Figure 4b), suggesting

that these genes may represent core molecular components in response to cold stress.

Next, to test whether the DEGs are closely related to cold response, we checked

their possible gene functions through gene function annotation and literature mining.

Indeed, we observed that many of them are directly involved in plant cold response or

cold acclimation process, including genes encoding superoxide dismutase (SOD) (Lin

et al., 2019), VERNALIZATION INSENSITIVE 3(VIN3) (Zhao et al., 2020),

transcription factor LUX (Chow et al., 2014), RNA-binding protein CP29B (Kupsch

et al., 2012), flavanone 3-dioxygenase 3 (Wu et al., 2023), and

dehydration-responsive element-binding protein 2E (DREB2E) (Agarwal et al., 2017),

etc. Also, many genes annotated as cold response or cold induced genes are observed

among these DEGs (Table S6 and S7). For example, there were 84 genes directly

annotated as cold response or cold induced genes in the NTS57 reference genome, of

which 38 and 49 were identified as DEGs in T1 and T2, respectively, implying that

these DEGs were enriched in the high-confidence reference gene set for rapeseed cold

resistance research. In addition, by combining the pan-genome analysis, we identified

83 and 82 DEGs in T1 and T2 samples, respectively, which are only present in the

NTS57 genome, representing NTS57-specific genes functioning in cold response

(Table S8).

To further understand the possible biological functions of the DEGs, we

performed Gene Ontology and KEGG pathway enrichment analysis. It revealed that at

the biological process (BP) level of DEGs in T1 are involved in the regulation of

circadian rhythms, photosynthesis, cold response and xyloglucan metabolism (Figure

4c). At the cellular component (CC) level, these DEGs are associated with MCM

complex, photosystem and cell wall, etc. (Figure S12). At the molecular function (MF)

level, these DEGs play roles in UDP-glycosyltransferase, protein serine/threonine

phosphatase activity, and ADP binding, etc. (Figure S12). For DEGs in T2 sample, we

observed a similar enrichment result, suggesting that cold-responsive genes are

relatively stable with increasing cold treatment time (Figure S12). The KEGG

pathway enrichment results showed that the DEGs were mainly involved in MAPK

signaling pathway, circadian rhythm, flavonoid biosynthesis, photosynthesis, etc.,

which were similar to the results of GO enrichment analysis (Figure S13).

To further find the possible functional modules controlling cold resistance in

NTS57, a weighted gene co-expression network analysis (WGCNA) was performed

by integrating the above cold stress RNA-Seq data and a newly generated gene

expression atlas of NTS57 from different tissues and developmental stages (Figure

S14, S15 and Table S9). Based on the WGCNA, a total of 25 distinct co-expression

modules with 24 to 15,396 genes per module were identified, and 15,496 uncorrelated

genes were assigned to a grey module that was ignored in the following study

(Figure 4d). We then mapped the 12,346 union DEGs onto these modules and tested

whether these DEGs are overrepresented in some particular modules. Indeed, there

are 10 modules, except the gray module, that are enriched in DEGs (hypergeometric

test; $P < 0.05$, Figure 4e), suggesting that the DEGs function cooperatively with the

gene in these modules. To confirm that these 10 modules are associated with cold

stress, we performed a module–trait relationships analysis and found that the

"darkgray" ($r = 0.55$, $P = 3e-05$), "darkgreen" ($r = 0.45$, $P = 0.001$), "lightcyan" ($r =

0.59$, $P = 6e-6$), and "brown" ($r = -0.67$, $P = 8e-08$) modules were highly correlated

with the cold stress (Figure 4f). Thus, we focus on the common modules, including

"darkgreen", "lightcyan", and "brown", which were both enriched in DEGs and

correlated with cold stress. The high correlation of these three modules with cold

stress was also confirmed by gene expression heat map analysis (Figure S16-18).

Gene function enrichment analysis revealed that genes in these modules are involved

in protein folding and heat response, fatty acid biosynthetic process, guard cell

differentiation, transcription, DNA-templated, and regulation of transcription by RNA

polymerase II or epigenetic modification, etc. (Figure S19). Collectively, the diverse

biological functions of these modules suggest that the cold response of rapeseed is

involved in different biological processes beyond our previous knowledge.

3.5. DNA hypomethylation contributes to cold resistance in the NTS57

Most of the existing research has documented that cold stress induces genomic

demethylation in many plant species, including rape, maize,

wheat, Arabidopsis, and *Antirrhinum majus* (Peng and Zhang, 2009). However, it is

unclear whether this rule applies to the materials of interest in this study. Therefore,

we calculated genome-wide DNA methylation levels around protein-coding genes in

the NTS57 genome under cold stress and normal conditions. The results showed a

remarkable decrease in methylation levels across the gene body and flanking regions

after cold stress compared to normal conditions in all cytosine sequence contexts CpG,

CHG and CHH (H for A, T or C) (Figure 5a). In particular, the decrease in CHH

methylation was most pronounced, which was followed by CHG and CpG

methylation, suggesting that CHH methylation is more sensitive to cold stress. Furthermore, when the A and C subgenomes were treated independently, the similar trends were observed, although the C subgenome had an overall higher methylation level compared to A, which was independent of the treatment conditions of the samples (Figure S20), which is consistent with previous studies (Bird et al., 2021; Chalhoub et al., 2014).

In plants, DNA methylation is enriched over heterochromatic transposable elements (TEs) and repetitive elements (Gallego-Bartolome, 2020). We next analyzed DNA methylation changes in repeat sequences and transposons in the NTS57 genome. First, as expected, there was a high enrichment of DNA methylation on repetitive sequences including LTR and DNA transposons and tandem repeats in general, although we noticed two exceptions that CHG and CHH methylation in tandem repeats showed some deletions compared to the surrounding regions (Fig 5b). We also found that methylation levels were decreased in all repeats and in all methylation contexts under cold stress, which was very similar to that of the genic regions, suggesting that methylase acts on genes or repeats in a similar way under plant cold stress response.

To further understand the methylation dynamics upon cold stress, we identified differentially methylated regions (DMRs) between the two cold treatment samples (T1 and T2) and the CK. A total of 631-9,004 DMRs were identified in all sequence contexts upon cold stress (Figure 5c). Obviously, the number of hypo-methylation regions is higher than that of the hyper-methylation regions in both T1 and T2 samples, which is consistent with the observation from methylation profiles that the methylation level decreased after cold treatment (Figure 5a). In addition, we observed that 47%-61% of these DMRs were located in the distal intergenic regions, followed by promoter regions of genes ranging from 24.12%-35.47% (Figure 5d), suggesting that the non-coding regions are the potential methylation target sites that response to cold stress in rapeseed. In particular, the DMRs in the CpG context were more likely to be deposited in promoter regions than those in the CHH and CHG in turn. On the contrary, DMRs in the CHG context were more frequently located in the distal

intergenic region that in the CHH and CpG contexts in turn. Based on these DMRs, we obtained 24,947 associated protein-coding genes, hereafter referred to as DMR-associated genes (Figure S21). Consistent with the DMRs number, the percentage of hypomethylated genes (63.5%-72.5%) was much higher than the number of hypermethylated genes in all three methylation contexts in the two samples of cold treatment. After comparing these DMR-linked genes with DEGs from two cold treatment samples, there are 506~2,381 common genes in three sequence contexts (Figure S22). We then determined the fraction of genes showing differential (hypo- or hyper-) methylation in different sequence contexts and their direction of differential expression (up or down-regulated) upon cold stress. In general, there was no consistent relationship between differential methylation in different sequence contexts and the direction of differential gene expression (Figure S23), suggesting that there is a complex DNA methylation regulatory mechanism regarding to DEGs at low temperature stress in rapeseed.

4. Discussion

Cold tolerance is one of the most important traits in crop domestication and breeding. In this study, we generated an improved high-quality genome assembly of a prominent cold-resistant rapeseed cultivar, NTS57. Various genome assembly indexes were used to confirm the high genome completeness. For example, the contig N50 of the present assembly is 21.81 Mb, almost 10 times longer than that of previous studies (Song et al., 2020), with the BUSCO score as high as 99.19%. Moreover, the number of assembly gaps is remarkably reduced in the present reference genome. Thus, this chromosome-level reference genome can allow us to systemically analyze the cold tolerance of rapeseed.

At the phylogeny level, the A and C subgenomes of the NTS57 have different clustering patterns, where the A subgenome was more likely to form a clade with that of semi-winter rapeseed, but the C subgenome was more closely related to the winter type rapeseed. This incongruity could be explained by the unique breeding history of the cold-tolerant NTS57. Basically, NTS57 underwent an additional artificial

interspecific hybridization between winter rapeseed and *Brassica rapa* after being bred from winter type rapeseed. This interspecific hybridization led to a high sequence similarity between the A subgenome of NTS57 and that of semi-winter rapeseed. However, the C subgenome showed sequence conservation with that of the winter rapeseed. At the pan-genome level, by aligning different rapeseed accessions to the newly assembled genome of NTS57, we detected numerous genomic structures, including indels, translocations, inversions, and duplications in NTS57 (Figure 3). These variations and associated genes are valuable for understanding the specific phenotypes of NTS57. Integration of the newly assembled genome into existing ones improves the quality of the rapeseed reference genome, which will promote the domestication and improvements in cold tolerance of rapeseed.

Studies have shown that Arabidopsis can respond to cold stress through either DREB1/CBF or DREB1/CBF-independent signaling pathways (Kidokoro et al., 2022). In the classical DREB1/CBF pathway, DREB1/CBF were considered as key transcription factors that involved in cold-responsive gene expression (Shi et al., 2018). By analyzing the transcriptomic data, there are at least 16 *DREBs* homologous genes in NTS57 that are induced by cold stress (Table S6 and S7). ICE1 has been considered as one of many transcription factors that can regulate the *DREB1* expression (Guo et al., 2018). However, we did not observe a significant change in the expression of *ICE1* in the NTS57 after cold stress, suggesting that ICE1 may not regulate the expression of the *DREB1* genes, or at least ICE1 is not induced by low temperature. Therefore, the ICE1-DREB1 regulatory module needs to be further confirmed (Kidokoro et al., 2022). In addition to the DREB1/CBF pathway, other transcription factors such as RVEs, ZAT12, HSFA1, BZR1, etc. are also differentially expressed under cold stress in the NTS57, suggesting a lot of crosstalk between DREB1/CBF and the DREB1/CBF-independent signaling pathways. Also, several uncharacterized genes such as *A07G034200*, *C08G048350*, *C01G004880* showed extremely significant expression changes upon cold stress. Whether they belong to DREB1/CBF or DREB1/CBF-independent signaling pathways needs further research, and these genes will be new potential targets for genetic modification of rapeseed to

improve cold tolerance.

DNA methylation is a conserved epigenetic modification that is important for gene regulation in plant responses to biotic and abiotic stress conditions (Zhang et al., 2018). Previous studies in many plant species, including rapeseed, maize, wheat, Arabidopsis, and *Antirrhinum majus*, have documented that cold stress induces genomic demethylation (Peng and Zhang, 2009). Consistent with these findings, we found that the upstream and downstream methylation levels of protein-coding genes in the NTS57 were significantly decreased under cold stress (Figure 5). The similar results were observed for repetitive elements, especially for tandem repeats. These results suggested that the decrease in methylation level under cold stress may trigger the transcription of cold responsive genes.

In summary, we have dissected the genomic architecture of cold tolerance by generating a high-quality reference genome of the NTS57 and performing phylogenetic, pan-genomic, DNA methylomic, and transcriptomic analyses of cold-responsive genes. These resources and findings provide valuable information that can be used in the breeding of rapeseed with high cold tolerance.

Acknowledgements

This work was supported by the Scientific Research Staring Foundation of Gansu Agricultural University (GAU-KYQD-2020-27), the Natural Science Foundation of Gansu Province (22JR5RA862), the Gansu Province Higher Education Institutions Young Doctoral Fund (2023QB-126), the Gansu Provincial Science and Technology Major Project (22ZD6NA009), the Central-Guided Local Science and Technology Development Fund (ZCYD-2021-17), and the National Natural Science Foundation of China (32360520).

CRediT authorship contribution statement

Zefeng Wu: Conceptualization, Funding acquisition, Formal analysis, Methodology, Visualization, Writing - original draft. **Guoqiang Zheng**: Conceptualization, Formal analysis, Investigation. **Yali Sun:** Conceptualization, Formal analysis. **Xiaoyun Dong:** Visualization, Software, Validation. **Ying Wang:** Software, Investigation. **Hui Li:**

690 Investigation, Validation. **Jiaping Wei:** Investigation, Validation. **Junmei Cui:**

691 Visualization, Software. **Yan Fang:** Investigation, Validation. **Yinin Niu:**

692 Investigation, Validation. **Zhen Huang:** Investigation, Writing – review & Editing.

693 **Jihong Hu:** Conceptualization, Writing – review & Editing. **Zigang Liu:**

694 Conceptualization, Funding acquisition, Supervision.

Declaration of competing interest

696 The authors declare that they have no known competing financial interests or personal

697 relationships that could have appeared to influence the work reported in this paper.

Data availability

All the raw sequencing data and the genome assembly generated for this research are

archived at NCBI under accession number PRJNA1012021. Genome annotation files

and other related documents are available in figshare

(https://doi.org/10.6084/m9.figshare.24098538.v1) or the database

(http://oilseed.online:8000/). All the materials and related analysis scripts from this

study are available upon request.

References

[1] Agarwal, P.K., Gupta, K., Lopato, S. and Agarwal, P. (2017) Dehydration responsive element
binding transcription factors and their applications for the engineering of stress tolerance. *J*
Exp Bot **68**, 2135-2148.
[2] Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F.E., Figueroa, M.E., Melnick, A. and
Mason, C.E. (2012) methylKit: a comprehensive R package for the analysis of genome-wide
DNA methylation profiles. *Genome Biol* **13**, R87.
[3] Alonge, M., Wang, X., Benoit, M., Soyk, S., Pereira, L., Zhang, L., Suresh, H., *et al.* (2020) Major
Impacts of Widespread Structural Variation on Gene Expression and Crop Improvement in
Tomato. *Cell* **182**, 145-161 e123.
[4] Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990) Basic local alignment
search tool. *J Mol Biol* **215**, 403-410.
[5] Benson, G. (1999) Tandem repeats finder: a program to analyze DNA sequences. *Nucleic*
Acids Res **27**, 573-580.
[6] Bird, K.A., Niederhuth, C.E., Ou, S., Gehan, M., Pires, J.C., Xiong, Z., VanBuren, R., *et al.* (2021)
Replaying the evolutionary tape to investigate subgenome dominance in allopolyploid
Brassica napus. *New Phytol* **230**, 354-371.
[7] Bolger, A.M., Lohse, M. and Usadel, B. (2014) Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120.
[8] Boutet, E., Lieberherr, D., Tognolli, M., Schneider, M. and Bairoch, A. (2007)

725   UniProtKB/Swiss-Prot. *Methods Mol Biol* **406**, 89-112.

726   [9]   Browse, J. and Xin, Z. (2001) Temperature sensing and cold acclimation. *Curr Opin Plant Biol* **4**,
727   241-246.

728   [10]   Burton, J.N., Adey, A., Patwardhan, R.P., Qiu, R., Kitzman, J.O. and Shendure, J. (2013)
729   Chromosome-scale scaffolding of de novo genome assemblies based on chromatin
730   interactions. *Nat Biotechnol* **31**, 1119-1125.

731   [11]   Cai, X., Chang, L., Zhang, T., Chen, H., Zhang, L., Lin, R., Liang, J.*, et al.* (2021) Impacts of
732   allopolyploidization and structural variation on intraspecific diversification in Brassica rapa.
733   *Genome Biol* **22**, 166.

734   [12]   Chalhoub, B., Denoeud, F., Liu, S., Parkin, I.A., Tang, H., Wang, X., Chiquet, J.*, et al.* (2014)
735   Early allopolyploid evolution in the post-Neolithic Brassica napus oilseed genome. *Science*
736   **345**, 950-953.

737   [13]   Chan, P.P., Lin, B.Y., Mak, A.J. and Lowe, T.M. (2021) tRNAscan-SE 2.0: improved detection and
738   functional classification of transfer RNA genes. *Nucleic Acids Res* **49**, 9077-9096.

739   [14]   Chen, S., Zhou, Y., Chen, Y. and Gu, J. (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor.
740   *Bioinformatics* **34**, i884-i890.

741   [15]   Chen, X., Tong, C., Zhang, X., Song, A., Hu, M., Dong, W., Chen, F.*, et al.* (2021) A high-quality
742   Brassica napus genome reveals expansion of transposable elements, subgenome evolution
743   and disease resistance. *Plant Biotechnol J* **19**, 615-630.

744   [16]   Cheng, H., Concepcion, G.T., Feng, X., Zhang, H. and Li, H. (2021) Haplotype-resolved de novo
745   assembly using phased assembly graphs with hifiasm. *Nat Methods* **18**, 170-175.

746   [17]   Chow, B.Y., Sanchez, S.E., Breton, G., Pruneda-Paz, J.L., Krogan, N.T. and Kay, S.A. (2014)
747   Transcriptional regulation of LUX by CBF1 mediates cold input to the circadian clock in
748   Arabidopsis. *Curr Biol* **24**, 1518-1524.

749   [18]   Cui, P., Li, Y., Cui, C., Huo, Y., Lu, G. and Yang, H. (2020) Proteomic and metabolic profile
750   analysis of low-temperature storage responses in Ipomoea batata Lam. tuberous roots. *BMC*
751   *Plant Biol* **20**, 435.

752   [19]   Danecek, P. and McCarthy, S.A. (2017) BCFtools/csq: haplotype-aware variant consequences.
753   *Bioinformatics* **33**, 2037-2039.

754   [20]   Ding, Y., Shi, Y. and Yang, S. (2019) Advances and challenges in uncovering cold tolerance
755   regulatory mechanisms in plants. *New Phytol* **222**, 1690-1704.

756   [21]   Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.*, et al.* (2013)
757   STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21.

758   [22]   Emms, D.M. and Kelly, S. (2019) OrthoFinder: phylogenetic orthology inference for
759   comparative genomics. *Genome Biol* **20**, 238.

760   [23]   Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C. and Smit, A.F. (2020)
761   RepeatModeler2 for automated genomic discovery of transposable element families. *Proc*
762   *Natl Acad Sci U S A* **117**, 9451-9457.

763   [24]   Gallego-Bartolome, J. (2020) DNA methylation in plants: mechanisms and tools for targeted
764   manipulation. *New Phytol* **227**, 38-44.

765   [25]   Gene Ontology, C. (2015) Gene Ontology Consortium: going forward. *Nucleic Acids Res* **43**,
766   D1049-1056.

767   [26]   Goel, M. and Schneeberger, K. (2022) plotsr: visualizing structural similarities and
768   rearrangements between multiple genomes. *Bioinformatics* **38**, 2922-2926.

[27] Goel, M., Sun, H., Jiao, W.B. and Schneeberger, K. (2019) SyRI: finding genomic
rearrangements and local sequence differences from whole-genome assemblies. *Genome*
Biol **20**, 277.

[28] Guan, S., Xu, Q., Ma, D., Zhang, W., Xu, Z., Zhao, M. and Guo, Z. (2019) Transcriptomics
profiling in response to cold stress in cultivated rice and weedy rice. *Gene* **685**, 96-105.

[29] Guo, X., Liu, D. and Chong, K. (2018) Cold signaling in plants: Insights into mechanisms and
regulation. *J Integr Plant Biol* **60**, 745-756.

[30] Haas, B.J., Salzberg, S.L., Zhu, W., Pertea, M., Allen, J.E., Orvis, J., White, O., *et al.* (2008)
Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to
Assemble Spliced Alignments. *Genome Biol* **9**, R7.

[31] Han, Y. and Wessler, S.R. (2010) MITE-Hunter: a program for discovering miniature
inverted-repeat transposable elements from genomic sequences. *Nucleic Acids Res* **38**, e199.

[32] Hu, J., Fan, J., Sun, Z. and Liu, S. (2020) NextPolish: a fast and efficient genome polishing tool
for long-read assembly. *Bioinformatics* **36**, 2253-2255.

[33] Huang, Q.X., Liao, X.Q., Yang, X.H., Luo, Y.C., Lin, P., Zeng, Q.H., Bai, H.R., *et al.* (2021) Lysine
crotonylation of DgTIL1 at K72 modulates cold tolerance by enhancing DgnsLTP stability in
chrysanthemum. *Plant Biotechnology Journal* **19**, 1125-1140.

[34] Huang, S.F., Kang, M.J. and Xu, A.L. (2017) HaploMerger2: rebuilding both haploid
sub-assemblies from high-heterozygosity diploid genome assembly. *Bioinformatics* **33**,
2577-2579.

[35] Jahan, K., Nie, H. and Yan, X. (2023) Revealing the potential regulatory relationship between
HSP70, HSP90 and HSF genes under temperature stress. *Fish & Shellfish Immunology* **134**,
108607.

[36] Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A. and Jermiin, L.S. (2017)
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**,
587-589.

[37] Kanehisa, M. and Goto, S. (2000) KEGG: kyoto encyclopedia of genes and genomes. *Nucleic*
Acids Res **28**, 27-30.

[38] Katoh, K. and Standley, D.M. (2013) MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772-780.

[39] Ke, L., Lei, W., Yang, W., Wang, J., Gao, J., Cheng, J., Sun, Y., *et al.* (2020) Genome-wide
identification of cold responsive transcription factors in Brassica napus L. *BMC Plant Biol* **20**,
62.

[40] Keilwagen, J., Hartung, F. and Grau, J. (2019) GeMoMa: Homology-Based Gene Prediction
Utilizing Intron Position Conservation and RNA-seq Data. *Methods Mol Biol* **1962**, 161-177.

[41] Kidokoro, S., Shinozaki, K. and Yamaguchi-Shinozaki, K. (2022) Transcriptional regulatory
network of plant cold-stress responses. *Trends Plant Sci* **27**, 922-935.

[42] Kim, D., Paggi, J.M., Park, C., Bennett, C. and Salzberg, S.L. (2019) Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-915.

[43] Kokot, M., Dlugosz, M. and Deorowicz, S. (2017) KMC 3: counting and manipulating k-mer
statistics. *Bioinformatics* **33**, 2759-2761.

[44] Krueger, F. and Andrews, S.R. (2011) Bismark: a flexible aligner and methylation caller for
Bisulfite-Seq applications. *Bioinformatics* **27**, 1571-1572.

[45] Kupsch, C., Ruwe, H., Gusewski, S., Tillich, M., Small, I. and Schmitz-Linneweber, C. (2012)

Arabidopsis chloroplast RNA binding proteins CP31A and CP29A associate with large
transcript pools and confer cold stress tolerance by influencing multiple chloroplast RNA
processing steps. *Plant Cell* **24**, 4266-4280.

[46] Lagesen, K., Hallin, P., Rodland, E.A., Staerfeldt, H.H., Rognes, T. and Ussery, D.W. (2007)
RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* **35**,
3100-3108.

[47] Langfelder, P. and Horvath, S. (2008) WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics* **9**, 559.

[48] Langmead, B. and Salzberg, S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat*
Methods **9**, 357-359.

[49] Li, H. (2018) Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
3094-3100.

[50] Li, H. and Durbin, R. (2009) Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760.

[51] Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., *et al.* (2009) The
Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079.

[52] Lin, K.H., Sei, S.C., Su, Y.H. and Chiang, C.M. (2019) Overexpression of the Arabidopsis and
winter squash superoxide dismutase genes enhances chilling tolerance via ABA-sensitive
transcriptional regulation in transgenic Arabidopsis. *Plant Signal Behav* **14**, 1685728.

[53] Lin, Y., Ye, C., Li, X., Chen, Q., Wu, Y., Zhang, F., Pan, R., *et al.* (2023) quarTeT: a
telomere-to-telomere toolkit for gap-free genome assembly and centromeric repeat
identification. *Hortic Res* **10**, uhad127.

[54] Liu, Z., Dong, X., Cao, X., Xu, C., Wei, J., Zhen, G., Wang, J., *et al.* (2022) QTL mapping for cold
tolerance and higher overwintering survival rate in winter rapeseed (Brassica napus). *J Plant*
Physiol **275**, 153735.

[55] Lu, Y., Hu, Y. and Li, P. (2017) Consistency of electrical and physiological properties of tea
leaves on indicating critical cold temperature. *Biosystems Engineering* **159**, 89-96.

[56] Mi, W., Liu, Z., Jin, J., Dong, X., Xu, C., Zou, Y., Xu, M., *et al.* (2021) Comparative proteomics
analysis reveals the molecular mechanism of enhanced cold tolerance through ROS
scavenging in winter rapeseed (Brassica napus L.). *PLoS One* **16**, e0243292.

[57] Nawrocki, E.P. and Eddy, S.R. (2013) Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics **29**, 2933-2935.

[58] Nguyen, L.T., Schmidt, H.A., von Haeseler, A. and Minh, B.Q. (2015) IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*
32, 268-274.

[59] Orvar, B.L., Sangwan, V., Omann, F. and Dhindsa, R.S. (2000) Early steps in cold sensing by
plant cells: the role of actin cytoskeleton and membrane fluidity. *Plant J* **23**, 785-794.

[60] Ou, S. and Jiang, N. (2018) LTR_retriever: A Highly Accurate and Sensitive Program for
Identification of Long Terminal Repeat Retrotransposons. *Plant Physiol* **176**, 1410-1422.

[61] Parra, G., Bradnam, K. and Korf, I. (2007) CEGMA: a pipeline to accurately annotate core
genes in eukaryotic genomes. *Bioinformatics* **23**, 1061-1067.

[62] Peng, H. and Zhang, J. (2009) Plant genomic DNA methylation in response to stresses:
Potential applications and challenges in plant breeding. *Progress in Natural Science* **19**,
1037-1045.

[63]   Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.C., Mendell, J.T. and Salzberg, S.L. (2015)
858   StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat*
859   *Biotechnol* **33**, 290-295.

860   [64]   Probst, A.V. and Mittelsten Scheid, O. (2015) Stress-induced structural changes in plant
861   chromatin. *Curr Opin Plant Biol* **27**, 8-16.

862   [65]   Pruitt, K.D., Tatusova, T. and Maglott, D.R. (2007) NCBI reference sequences (RefSeq): a
863   curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic*
864   *Acids Res* **35**, D61-65.

865   [66]   Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R. and Lopez, R. (2005)
866   InterProScan: protein domains identifier. *Nucleic Acids Res* **33**, W116-120.

867   [67]   Ranallo-Benavidez, T.R., Jaron, K.S. and Schatz, M.C. (2020) GenomeScope 2.0 and
868   Smudgeplot for reference-free profiling of polyploid genomes. *Nat Commun* **11**, 1432.

869   [68]   Servant, N., Varoquaux, N., Lajoie, B.R., Viara, E., Chen, C.J., Vert, J.P., Heard, E., *et al.* (2015)
870   HiC-Pro: an optimized and flexible pipeline for Hi-C data processing. *Genome Biol* **16**, 259.

871   [69]   Shi, Y., Ding, Y. and Yang, S. (2018) Molecular Regulation of CBF Signaling in Cold Acclimation.
872   *Trends Plant Sci* **23**, 623-637.

873   [70]   Simao, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V. and Zdobnov, E.M. (2015)
874   BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs.
875   *Bioinformatics* **31**, 3210-3212.

876   [71]   Song, J.M., Guan, Z., Hu, J., Guo, C., Yang, Z., Wang, S., Liu, D., *et al.* (2020) Eight high-quality
877   genomes reveal pan-genome architecture and ecotype differentiation of Brassica napus. *Nat*
878   *Plants* **6**, 34-45.

879   [72]   Stanke, M., Steinkamp, R., Waack, S. and Morgenstern, B. (2004) AUGUSTUS: a web server for
880   gene finding in eukaryotes. *Nucleic Acids Res* **32**, W309-312.

881   [73]   Sun, F., Fan, G., Hu, Q., Zhou, Y., Guan, M., Tong, C., Li, J., *et al.* (2017) The high-quality
882   genome of Brassica napus cultivar 'ZS11' reveals the introgression history in semi-winter
883   morphotype. *Plant J* **92**, 452-468.

884   [74]   Sun, H.Q., Ding, J., Piednoël, M. and Schneeberger, K. (2018) findGSE: estimating genome size
885   variation within human and Arabidopsis using k-mer frequencies. *Bioinformatics* **34**, 550-557.

886   [75]   Tang, H., Bowers, J.E., Wang, X., Ming, R., Alam, M. and Paterson, A.H. (2008) Synteny and
887   collinearity in plant genomes. *Science* **320**, 486-488.

888   [76]   Tarailo-Graovac, M. and Chen, N. (2009) Using RepeatMasker to identify repetitive elements
889   in genomic sequences. *Curr Protoc Bioinformatics* **Chapter 4**, 4 10 11-14 10 14.

890   [77]   Tatusov, R.L., Natale, D.A., Garkavtsev, I.V., Tatusova, T.A., Shankavaram, U.T., Rao, B.S.,
891   Kiryutin, B., *et al.* (2001) The COG database: new developments in phylogenetic classification
892   of proteins from complete genomes. *Nucleic Acids Res* **29**, 22-28.

893   [78]   Wang, X. and Wang, L. (2016) GMATA: An Integrated Software Package for Genome-Scale SSR
894   Mining, Marker Development and Viewing. *Front Plant Sci* **7**, 1350.

895   [79]   Wei, D., Cui, Y., He, Y., Xiong, Q., Qian, L., Tong, C., Lu, G., *et al.* (2017) A genome-wide survey
896   with different rapeseed ecotypes uncovers footprints of domestication and breeding. *J Exp*
897   *Bot* **68**, 4791-4801.

898   [80]   Wei, J., Shen, Y., Dong, X., Zhu, Y., Cui, J., Li, H., Zheng, G., *et al.* (2022) DNA methylation
899   affects freezing tolerance in winter rapeseed by mediating the expression of genes related to
900   JA and CK pathways. *Front Genet* **13**, 968494.

[81] Wei, J., Zheng, G., Yu, X., Liu, S., Dong, X., Cao, X., Fang, X., *et al.* (2021) Comparative
Transcriptomics and Proteomics Analyses of Leaves Reveals a Freezing Stress-Responsive
Molecular Network in Winter Rapeseed (Brassica rapa L.). *Front Plant Sci* **12**, 664311.
[82] Wingett, S.W. and Andrews, S. (2018) FastQ Screen: A tool for multi-genome mapping and
quality control. *F1000Res* **7**, 1338.
[83] Wu, L., Tian, J., Yu, Y., Yuan, L., Zhang, Y., Wu, H., Wang, F., *et al.* (2023) Functional
characterization of a cold related flavanone 3-hydroxylase from Tetrastigma hemsleyanum:
an in vitro, in silico and in vivo study. *Biotechnol Lett*.
[84] Yu, J., Cang, J., Lu, Q., Fan, B., Xu, Q., Li, W. and Wang, X. (2020) ABA enhanced cold tolerance
of wheat 'dn1' via increasing ROS scavenging system. *Plant Signal Behav* **15**, 1780403.
[85] Yu, T., Zhang, J., Cao, J., Cai, Q., Li, X., Sun, Y., Li, S., *et al.* (2021) Leaf transcriptomic response
mediated by cold stress in two maize inbred lines with contrasting tolerance levels. *Genomics*
113, 782-794.
[86] Yuan, Y., Bayer, P.E., Batley, J. and Edwards, D. (2021) Current status of structural variation
studies in plants. *Plant Biotechnol J* **19**, 2153-2163.
[87] Zhang, H., Lang, Z. and Zhu, J.K. (2018) Dynamics and function of DNA methylation in plants.
Nat Rev Mol Cell Biol **19**, 489-506.
[88] Zhao, Y., Antoniou-Kourounioti, R.L., Calder, G., Dean, C. and Howard, M. (2020)
Temperature-dependent growth contributes to long-term cold sensing. *Nature* **583**, 825-829.
920

Figures and their captions

923   **Figure 1. Cold tolerance phenotypes and genome assembly of NTS57. (a)** Seedling survival

924   of NTS57, ZS11 and Westar. Seedlings were photographed before or 12h and 24h after treatment

925   at -4 ☐, and 12h after returning to normal conditions at 25 ☐. **(b)** Genome-wide Hi-C heat map of

the NTS57 genome. The heatmap shows the diagonal pattern of strong Hi-C intra-chromosomal

interactions of NTS57. The color bar indicates the log-transformed interaction frequencies of the

Hi-C chromosome links. **(c)** Circos plot showing the genome assembly and genomic features of

NTS57. The letters (*a-i*) in the circos indicate the ideograms of 19 chromosomes of the NTS57

genome, gene density, GC content, Copia transposons, Gypsy transposons, CpG methylation,

CHG methylation, CHH methylation, and chromosomal collinearity, respectively. The

centromeres are highlighted in black in the ideograms.

Figure 2. Phylogenetic relationships and pan-genomic analysis of NTS57 and other rapeseed

accessions. (a) Maximum likelihood tree constructed using 2,268 single-copy genes showing the

phylogenetic relationships of NTS57 and other public rapeseed accessions. Bootstrap values are

given as percentages on the branches of the consensus tree inferred from 1000 replicates. **(b)**

Modelling of pangenome and coregenome sizes by including additional rapeseed accessions. **(c)**

and **(d)** show the percentage of core genes, dispensable genes and accession-specific genes in the A and C subgenomes of rapeseed, respectively.

Figure 3. Genome structure variation between NTS57 and other rapeseed accessions. (a) Number of different SV types within each rapeseed accession using NTS57 as the reference genome. "TRANS", "INV", "INS", "DUP", and "DEL" stand for translocation, inversion, insertion, duplication, and deletion, respectively. **(b)** SV landscape in the NTS57 genome. The heat map shows the correlations between different SVs, GC content and gene density, with the correlation significance was marked in the corresponding cells. **(c)** Chromosomal synteny for A03 and C03 between Darmor-*bzh*, ZS11 and NTS57. The green and blue ribbons show the synteny blocks of the leftmost region (~6.6 Mb) of C03 in NTS57 with that of A03 and C03 of ZS11, respectively.

Figure 4. Gene expression analysis of NTS57 under cold stress. (a) Sankey plot showing the up- and down-regulated genes of NTS57 after 12h (T1) and 24h (T2) cold stress. **(b)** Shared DEGs in T1 and T2 samples. **(c)** Gene function enrichment of DEGs in T1 samples. **(d)** Co-expression gene modules identified by WGCNA analysis, with different colors denoting the different gene

modules. **(e)** Enrichment of DEGs in each of the co-expression gene modules. **(f)** Heat map

showing the module-trait relationships. The color bar denotes the correlation of the modules with

each trait.

Figure 5. Methylation dynamics of NTS57 under cold stress. (a) Profiles of methylation

levels in the gene body, upstream and downstream regions in the NTS57 genome under normal

and cold stress. The gene body was divided into 100 bins of equal size. For each bin in each gene,

the methylation level was calculated by dividing the total number of methylated Cs by the total

number of Cs, and the average of methylation level in the window across all protein genes was

used to generate the profiles. The three methylation profiles for CpG, CHG and CHH contexts

were analyzed independently. **(b)** Similar to (a) but showing the profiles of methylation levels in

the repetitive elements of TNS57. T1 and T2 represent 12 and 24 h after cold treatment, and the

CK represents the normal condition. **(c)** Number of differentially methylated regions (DRMs) in

T1 and T2 samples relative to CK. DMRs were classified into hypermethylation (hyper) and

hypomethylation (hypo). **(d)** The percentage of DMRs in different genomic contexts.

Tables

Table 1. Summary of genome assembly and annotation for NTS57 and other public rapeseed genomes.

Accessions	Assembly size (MB)	Anchored percentage (%)	Contig N50 (Mb)	No. of predicted genes	Repeats (%)	BUSCO (%)	No. of gaps	Reference
Darmor-*bzh* (v10)	924	93.84	11.48	108,190	53.86	98.60	268	Rousseau-Gueutin et al.
Westar	1007	93.54	3.13	97514	55.78	98.00	2164	Song et al.
No2127	1011	92.18	2.70	95385	56.86	98.00	2073	Song et al.
Zheyou7	1015	91.03	2.10	96209	56.39	98.30	3027	Song et al.
Gangan	1033	90.41	2.71	96843	56.58	98.30	1976	Song et al.
Shengli	1001	91.98	2.82	94586	56.29	98.10	2241	Song et al.
Tapidor	1014	91.91	2.85	96117	56.59	98.00	1871	Song et al.
Quinta	1003	93.02	2.80	95492	56.29	98.30	2132	Song et al.
ZS11	1010	95.33	2.70	100,919	55.62	98.10	5460	Song et al.
ZS11	921	93.10	1.64	106,059	61.83	98.30	361	Chen et al.
NTS57	1062	93.60	21.81	101,438	55.60	99.75	66	This study

Supporting Information

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Figure S1. Genome size estimation based on survey sequencing.

Figure S2. Bias in gene expression between two sub-genomes of NTS57.

Figure S3. Venn diagram showing the results of gene function annotations using different databases.

Figure S4. Gene function enrichment for core genes and dispensable genes of NTS57.

Figure S5. Comparisons of gene expression levels for three classes of genes in NTS57.

Figure S6. Length statistics for structural variations in different rapeseed accessions using NTS57 as the reference genome.

Figure S7. Number of structural variations in different rapeseed accessions using ZS11 as the reference genome.

Table S3. Summary of non-coding RNAs in the NTS57 genome.

Table S4. Summary of repetitive elements in the NTS57 genome.

Table S5. Genes associated with SVs on the chromosome C03 of NTS57

Table S6. Summary of differentially expressed genes after 12 hours of cold
exposure.

Table S7. Summary of differentially expressed genes after 24 hours of cold
exposure.

Table S8. NTS57 specific genes that were differentially expressed in cold stress.

Table S9. Gene expression levels of NTS57 across different developmental stages
and tissues.

1032