Single cell RNA-sequencing suggests a novel lipid associated mast cell population following weight cycling

Heather L. Caslin[1,2], Matthew A. Cottam[3], Alexa M. Betjemann[1], Mona Mashayekhi[4], Heidi J Silver [5,6], Alyssa H. Hasty[1,6]

[1] Department of Molecular Physiology and Biophysics, Vanderbilt University
[2] Department of Health and Human Performance, University of Houston
[3] Department of Surgery, Division of Surgical Oncology & Endocrine Surgery, Vanderbilt University Medical Center
[4] Department of Medicine, Division of Diabetes, Endocrinology, and Metabolism, Vanderbilt University Medical Center
[5] Department of Medicine, Division of Gastroenterology, Hepatology & Nutrition, Vanderbilt University Medical Center
[6] Veterans Affairs, Tennessee Valley Healthcare System

Corresponding Author:
Heather Caslin, PhD
3875 Holman St.
Rm 104 Garrison
Houston, TX 77204-6015
hcaslin@central.uh.edu

Running Title: Adipose mast cells in weight cycling

Keywords: weight cycling, weight loss, weight regain, adipose, mast cells, immunometabolism, lipid associated macrophages

Abstract

      Our recent study showed weight cycled mice have increased adipose mast cells compared to obese mice by single cell RNA-sequencing. Here, we aimed to confirm and elucidate these changes. Further analysis of our dataset showed that our initial mast cell cluster could subcluster into two unique populations: one with very high expression of classical mast cell markers and another with elevated lipid handling and antigen presentation genes. This new mast cell cluster accounted for most of the mast cells in the weight cycled group although it was not possible to detect the different populations by new studies with flow cytometry or Toluidine blue staining in mice, possibly due to a downregulation in classical mast cell genes. Interestingly, a pilot study in humans did suggest the existence of two mast cell populations in subcutaneous adipose tissue from obese women that appear similar to the murine populations detected by sequencing; one of which was significantly correlated with weight variance. Together, these data suggest that weight cycling may induce a unique population of mast cells similar to lipid associated macrophages. Future studies will focus on isolation of these cells to better determine their lineage, differentiation, and functional roles.

Abbreviations

BMI   Body mass index

CD   Cluster of differentiation marker

FMO   Fluorescence minus one

IFN   Interferon

IL-   Interleukin

MFI   Mean fluorescent intensity

MHC   Major histocompatibility complex

PBS   Phosphate buffered saline

scRNAseq   Single cell RNA-sequencing

1. Introduction

Risk for cardiometabolic diseases is higher with excess adipose tissue, and thus, weight loss is often prescribed to reduce risk. Attempts at weight loss are common; the most recent NHANES data indicates that ~50% of adults in the United States attempted weight loss between 2013-2016[1]. Other reports suggest that up to 76% of adults in the United States have attempted weight loss in their lifetime[2]. Weight regain is also a common occurrence as most individuals regain most of the lost weight within just a few years[3–8]. This process of repeatedly gaining and losing weight, termed weight cycling, further increases the risk for type 2 diabetes, cardiovascular disease, and hypertension beyond that of stable long-term obesity[9–13].

During weight gain, and specifically adipose expansion, immune cell recruitment to the adipose tissue promotes adipocyte lipolysis, fibrosis, and insulin resistance[14]. These changes contribute to the development of type 2 diabetes and other cardiometabolic diseases. With adipose expansion, type 1 immune subsets like T helper 1 cells, CD8$^+$ T cells, and neutrophils dominate, and macrophages shift from a tissue regulatory phenotype to a lipid handling, pro-inflammatory phenotype[15,16]. However, less is known about how immune cells change and contribute to systemic metabolism with weight cycling. We recently used single-cell RNA-sequencing (scRNAseq) to classify differences in proportion and phenotype of gonadal adipose tissue immune cells from lean, obese, weight loss, and weight cycled mice[17]. As previously published[18], weight cycled mice had worsened glucose tolerance compared with their weight stable obese counterparts. Unexpectedly, these data also showed that weight cycling substantially increased the proportion of adipose mast cells.

Mast cells are tissue-resident cells in connective and mucosal tissues around the body[19]. Primarily considered type 2 immune cells, mast cells are well-studied for their protective role against extracellular parasites and their detrimental role in allergic disease and asthma[20,21]. However, mast cells also have many physiological functions in the defense against bacteria and viruses[22,23] and contribute to autoimmunity and atherosclerosis[24]. While mast cells have been shown to increase with weight gain, their role in adipose tissue homeostasis and expansion is not well understood, and there are contradictory results in the literature due to the use of different mast cell knockout models and different concentrations of mast cell stabilizers[25–29]. Therefore, the purpose of this study was to elucidate how adipose mast cells change under the conditions of weight gain, weight loss, and weight cycling.

2. Methods

2.1 Animals

Male C57BL/6J mice were purchased from Jackson Labs (Bar Harbor, ME, #000664) at 7 weeks of age. At 8 or 9 weeks of age, male C57BL/6J mice were placed on 9-week cycles of 60% high fat diet (Research Diets, New Brunswick, NJ, #D12492, 5.21 kcal/g food) or 10% low fat diet (Research Diets #D12450B, 3.82 kcal/g) for a total of 27 weeks as published[17,18] and as visualized in **Figure 1A**. Food and water were provided *ad libitum*, and body weight and food intake were recorded weekly.

After 26 weeks on diet, mouse body fat and muscle mass were measured by nuclear magnetic resonance whole body composition analysis using a Minispec (Bruker,

Billerica, MA). The following day, mice were fasted for 5-h and briefly anesthetized with isoflurane to obtain blood via tail snip. After a 1-h recovery, basal blood glucose levels were measured using a hand-held glucometer (Contour Next EZ, Bayer, Pittsburg, PA). Mice were injected intraperitoneal with 1.5 g dextrose/kg lean mass and blood glucose was measured at 15, 30, 45, 60, 90, and 120 min.

All procedures were approved and performed in compliance with the Vanderbilt University Institutional Animal Care and Use Committee. Vanderbilt University is accredited by the Association for Assessment and Accreditation of Laboratory Animal Care International.

2.2 Single cell analysis

We used our previously published single cell sequencing dataset to further assess adipose mast cells (GEO accession number GSE182233)[17]. In this previous dataset, all cluster annotations were performed using common cell type markers and confirmed via SingleR V1.6. Here, we subset the data on what was initially clustered as mast cells and performed further clustering using the FindClusters function in Seurat V4[30] with a resolution of 0.1. Differential expression analysis was performed in R version 4.1 with Seurat V4 using Wilcoxon Ranked Sum tests with the FindMarkers function to compare the new subclusters. GO Biological Processes pathway analysis was conducted on differential expression with ShinyGO 0.76.2[31] using FDR cutoff of 0.05 (last date of access 10/17/2022).

2.3 Histology and toluidine blue staining

Epididymal adipose tissue was fixed in 4% paraformaldehyde for 24-h, then washed with phosphate buffered saline (PBS). Samples were embedded in paraffin and then sectioned with the help of the Translational Pathology Shared Resource core at Vanderbilt University. Slides were the deparaffinized in xylene and rehydrated, then stained in 10% Toluidine blue for 5 min. After staining, slides were dehydrated, cleared in xylene, and mounted. Slides were imaged with an inverted DMi8 microscope and Flexacam color camera (Leica, Teaneck, NJ) at 20x. Multiple images were taken using the tilescan function to collect 104 sections/image. Mast cells were counted per 104 frames for two tissue sections per sample and counts for each sample were averaged.

2.4 Murine adipose immune cell isolation

Epididymal and subcutaneous adipose tissue were collected as previously described[32]. Following isoflurane overdose and cervical dislocation, mice were perfused with 20 ml PBS through the left ventricle. Adipose fat pads were collected, minced, and digested in 6 ml of 2 mg/mL type IV collagenase (Worthington Biomedical, Lakewood, NJ, #LS004186) for 30 min at 37°C. The collagenase was diluted with cold PBS and digested tissue was vortexed and filtered through 100 µm filters. The pelleted stromal vascular fraction was then lysed with ACK buffer to remove red blood cells and filtered through a 35 µm filter to produce a single cell suspension.

2.5 Human data and adipose samples

Human subcutaneous adipose tissue samples were collected from an ongoing clinical study (NCT04907214). The study was approved by the Vanderbilt Institutional Review Board and conducted according to the Declaration of Helsinki. All participants provided written informed consent. We obtained adipose samples paired with deidentified demographic information and self-reported weight history from four participants. Inclusion criteria was women with obesity (BMI ≥ 30kg/m^2) and pre-diabetes (fasting glucose of 100-125mg/dL, or two-hour plasma glucose of 140-199 mg/dL during a 75-gram oral glucose tolerance test, or hemoglobin A1c of 5.7-6.4%). Additionally, all participants completed a weight history questionnaire asking about weight over the past 6 months, one year, two years, five years, and ten years along with any noted diet attempts at weight loss. As an indicator of weight cycling, we calculated sample weight variance (standard deviation squared) for each subject.

Participants underwent bedside liposuction for collection of adipose tissue from the periumbilical region using a Tulip^{TM} Medical closed syringe system (Tulip Medical Products, San Diego, CA). Under aseptic conditions and local lidocaine anesthesia, a small incision was made in the skin. The GEMS Johnnie Snap lock was placed inside a 60-cc syringe with the Tulip liposuction cannula attached and was inserted at an angle through the incision to below Scarpa's fascia. Suction was applied until the syringe activated the clicker lock. The needle was oscillated at a rate of approximately 1 Hz without breaking suction with a twisting motion. The sampling continued until 10-15 grams of tissue was removed. The tissue sample was placed in 20-30 cc of cold saline and immediately processed for enrichment of the stromal vascular fraction.

Harvested adipose tissue was rinsed with PBS and weighed, then digested in gentleMACS C tubes with a Dissociator (Miltenyi Biotec, Gathersburg, MD). Collagenase D (Roche, Indianapolis, IN) was added at 2 mg/mL and the sample was shaken at 37 °C degrees at 150 rpm for 40 min. The digested tissue was rinsed, filtered through a 70 micron filter, and the floating adipocyte fraction removed. The remaining immune-rich stromal vascular fraction was pelleted and washed, then used for staining for flow cytometry.

2.6 Flow cytometry

Human and murine stromal vascular cells were prepared for flow cytometry. FcBlock (1:200) was added for 10 min. Surface proteins were stained with fluorescent antibodies at 1:200 for 30 min at 4 °C using the antibody panels in **Supplemental Table 1**. Samples were then stained with DAPI prior to acquisition (1:5000). Data was acquired on a MACSQuant10 (Miltenyi) or 3 laser Fortessa (BD Biosciences, San Jose, CA) and analyzed on FlowJo using single color compensation controls and fluorescence minus one (FMO) controls for gating.

2.7 RNA isolation and real time PCR

RNA was isolated with a Qiagen RNeasy Mini Kit. Following quantification with a Nanodrop, the RNA was converted to cDNA using iScript Reverse Transcription Supermix for RT-qPCR (Bio-rad, Hercules, CA, #1708840) or Sensifast cDNA Synthesis kit (Meridian Biosciences, Memphis, TN, #BIO-65054). Real Time PCR was completed

using Taqman primer-probe sets from Thermofischer and iQ Supermix (Bio-Rad #1708860). Gene expression was normalized via ΔΔCt method to a housekeeping gene and the lean control.

2.8 Statistical analyses

Statistical analyses were performed using GraphPad Prism. Student's *t*-tests were run for comparisons between two groups, one-way analysis of variances (ANOVA)s were used for comparisons between more than two groups, and two-way ANOVAs were conducted for >2 groups over >2 timepoints. For significant main effects, *post-hoc* pairwise comparisons using Tukey or Sidak corrections were used to determine statistical differences. Simple regression was used for correlations. All data are presented as mean ± standard error of the mean (SEM). A p-value (or adjusted p-value) of <0.05 was used to determine significance.

3. Results

3.1 scRNAseq data show increased murine adipose mast cells with weight cycling

Adipose immune cell populations change with weight gain; however, less is known about how these populations change with weight loss and weight cycling. Therefore, to examine changes in adipose immune populations with weight loss and weight cycling, we recently completed scRNAseq in a mouse model of weight cycling and published the initial findings broadly focused on T cells, dendritic cells, and macrophages. In the mouse model used by our research group, male mice are placed on low fat or high fat diets for 27 weeks total as shown in **Figure 1A**. Strikingly, in our

scRNAseq study, we found that mast cell counts were significantly elevated in the weight cycled mice compared to all other diet conditions, and for which we provide new analyses for here (**Figure 1B&C** and [17]). Additionally, our previous report showed that classical mast cell genes such as chymase (*Cma1*) and the IL-33 receptor (*Il1r1*) were reduced in the weight cycled cluster while lipid handling genes such as *Trem2* and *Fabp5* were increased in the weight cycled cluster [17]. Here, we show these changes by violin plot **(Figure 1D&E**).

3.2 Standard techniques do not identify increased mast cells in weight cycling

To confirm the previous scRNAseq data, we attempted to detect mast cells using complementary techniques in a new cohort of mice. As seen previously in our sequencing as well as other published studies[17,18,33,34], glucose clearance was significantly lower in the weight cycled group compared with the obese group during an intraperitoneal glucose tolerance test, despite similar body weight at the end of 27 weeks (**Figure 2A&B**). For initial mast cell analysis, fixed sections of epididymal adipose tissue were stained with toluidine blue, a cationic dye that stains nuclei pink and mast cells dark purple (example image shown in **Figure 2C**). Supporting the sequencing data, there was a significant increase in mast cells in the obese and weight loss groups; however, the weight cycled group did not show increased mast cells compared with any other group (**Figure 2D**). Flow cytometry was next used to quantify mast cells in epididymal and subcutaneous adipose tissue by the expression of canonical mast cell markers, cKit and FcεR1 (**Figure 2E&F**). While there was an increase in the percent and number of mast cells/gram epididymal fat in weight cycled

animals compared to the lean mice, weight loss animals showed the highest number of mast cells (**Figure 2F**). We also analyzed the expression of ST2, the IL-33 receptor, and CD107a and CD63, two lysosomal proteins which can appear on the surface of mast cells following degranulation and detected an increase in CD63 in the obese and weight cycled groups compared to the lean and weight loss groups (**Supplemental Figure 1**).

Similar to the epididymal adipose tissue, there was a significant increase in mast cells in the subcutaneous adipose tissue of the weight loss animals compared with the lean and weight cycled groups by flow cytometry (**Figure 2G&H**). However, the number of mast cells/gram fat was more variable in this fat pad, and there were no significant differences. Real time PCR analysis was also performed on whole subcutaneous adipose tissue for *Fcer1* (the gene for FcεR1), the proteases *Mcpt4, Tpsb2,* and *Cma1 (*found in mast cell granules) and the tryptophan hydroxylase gene *Tph1 (*shown in adipose mast cells to be important for the metabolic adaptations to thermoneutrality[35]). *Fcer1* was significantly elevated in both the obese and weight cycled groups compared to the lean animals (**Supplemental Figure 2**). Weight loss increased *Mcpt4, Tpsb2, Cma1,* and *Tph1*, but this was only significant compared to the lean group for *Tpsb2* and Tph1 (**Supplemental Figure 2**). These results suggest that we cannot detect the increase in adipose mast cells during weight cycling by standard methods of detection.

3.3 scRNAseq suggests weight cycling induces a distinct cluster of lipid handling mast cells

Because our sequencing data were not supported by standard methods, we returned to our published scRNAseq data. First, we subset our data on the initial "mast cell" cluster which expresses high levels of mast cell specific genes (*Mcpt4*, *Cma1*, and *Tpsb2*) when compared to all other cells. We then recalculated the highly variable genes among the remaining cells and clustered the cells using a new uniform manifold approximation and projection (UMAP) based on the first 30 principal components. Following reclustering using the Louvain implementation in Seurat, two distinct subclusters were identified (**Figure 3A**). There was a significant increase only in cluster 2 in the weight cycled group compared with the other diet groups (**Figure 3B**). Pathway analysis of differentially expressed genes in cluster 2 compared to cluster 1 suggested a reduction in many leukocyte differentiation, hematopoiesis, and activation genes and an increase in biosynthetic and metabolic genes (**Supplemental Figure 3**). Moreover, there was a significant decrease in many surface and protease genes specifically related to mast cells and an increase in some of the lysosomal, surface receptor genes often associated with macrophages in cluster 2 (**Figure 3C and Supplemental Table 2**). It's important to note that mast cell genes were still increased in cluster 2 relative to other populations in our data. We show this relative to all macrophage populations (**Supplemental Figure 4**). Additionally, there was a significant increase in genes involved in lipid handling and antigen presentation in cluster 2 as compared to cluster 1 (**Figure 3D and Supplemental Table 2**). All significant differences between clusters are reported in Supplemental Table 2. Markedly, these cells appear very similar to lipid associated macrophages, first reported in 2019[36], though they are a unique population by sequencing (**Supplemental Figure 4A**) and due to the expression of

mast cell associated genes (**Supplemental Figure 4B**). Together, these new analyses suggest that weight cycling specifically increases a novel population of lipid handling mast cells.

3.4 Weight loss and weight cycling affect proteins involved with lipid handling and antigen presentation on adipose mast cells

To assess lipid handling and antigen presentation in the mast cells from weight cycled mice, we used flow cytometry for CD36 and CD9 (lipid handling) and MHCII (I-A/ I-E) and CD74 (antigen presentation). Of the proteins involved in lipid handling, there was a significant increase in the percent of mast cells positive for CD36 and CD9 in the weight loss group (**Figure 4A**). Additionally, there was an increase in the percent of mast cells positive for CD9 in the weight cycled group compared to the lean group. Of the proteins involved with antigen presentation, there was an increase the percent of mast cells positive for MHCII (I-A/ I-E) and an increase in the surface protein expression level of MHCII (I-A/ I-E) in mast cells in the weight loss compared to the lean and obese groups (**Figure 4B**). Moreover, there was an increase in the percent of cells positive for MHCII (I-A/ I-E) and CD74 as well as the surface expression level of CD74 in the weight cycled mast cells compared to the lean group. While we could not detect different groups of cells by flow cytometry, these data suggest that weight loss and weight cycling may differentially affect lipid handling and antigen presentation-related surface markers on adipose mast cells, which should be confirmed by other methods and functional analyses in the future.

3.5 Weight variance correlates with a cKit intermediate mast cell population in humans

Finally, to determine whether mast cell populations in humans are affected by weight cycling, we collected pilot data from four women with obesity and prediabetes. Immune cells were obtained from the subcutaneous adipose tissue and weight history was collected via questionnaire. Baseline participant characteristics include age 56.3±12.0 years, weight 109.6±27.7 kg, BMI 40.9±9.6 kg/m2, and fasting glucose 102.5±5.4 mg/dL. Using FMO controls, we gated on cKit$^+$ FcϵR1$^+$ cells in the adipose tissue samples (**Figure 5A**). Interestingly, there appeared to be two diverse populations of mast cells, one with very high cKit expression (cKit^{hi} FcϵR1$^+$) and one with more intermediate cKit expression (cKit^{int} FcϵR1$^+$). The cKit^{int} FcϵR1$^+$ population was more prevalent than the cKit^{hi} FcϵR1$^+$ population (**Figure 5B**). The cKit^{hi} FcϵR1$^+$ population had increased surface expression of CD9 by MFI. By comparison, the cKit^{int} FcϵR1$^+$ population had a greater percentage of CD68$^+$ cells and greater CD68 surface expression by MFI. Additionally, there was a trend towards increased MFI of CD36 in the cKit^{int} FcϵR1$^+$ cells.

We next determined whether these mast cell populations were different based on the weight history of the four study participants (**Figure 5D**). The participant who had most recently lost weight had the greatest population of the cKit^{hi} FcϵR1$^+$ cells, although there was no significant relationship between weight change in the past year and cKit^{hi} FcϵR1$^+$cells (**Figure 5E**). To examine weight cycling, we calculated weight variance (standard deviation squared) in all participants. There was a significant correlation between weight variance and the cKit^{int} FcϵR1$^+$ cells (**Figure 5F**). These data suggest that women who weight cycle have adipose mast cells with a different expression

pattern of mast cell identity markers, lipid handling, and antigen presentation related surface proteins.

4. Discussion

Many studies show that immune cell populations change with adipose tissue expansion[37–39]. However, despite the high prevalence of weight loss and weight regain in the US population, only a few studies measure the proportion and function of adipose immune cells in these conditions[17,38,40–43]. Our data suggest that adipose mast cells change with weight loss and weight cycling, though our tools to identify these lipid handling mast cells in the adipose tissue are currently limited.

Using new scRNAseq analysis, we identified two distinct mast cell clusters in the adipose tissue of male mice from our larger mast cell cluster from data previously published by our group [17]. The first is a cluster found in all mice that highly expresses the canonical mast cell surface receptors and proteases such as *Fcer1, Kit, Cma1,* and *Tpsb2* among many others. We consider these the classical mast cell population. Additionally, there is a second cluster of cells that also express classical mast cell markers at a high level relative to all other immune cells, but at a lower level compared to the classical mast cell cluster. These cells also have increased expression of genes that play a role in lipid handling and antigen presentation. Interestingly, many of these genes are also elevated in a cluster of adipose macrophages following weight gain, first identified and named "lipid associated macrophages", or LAMs, in 2019[36]. In our dataset, the LAMs and lipid handling mast cells cluster nearby on the UMAP, but are two distinct groups based on marker expression[17]. Importantly, the macrophage

populations do not express mast cell genes and the lipid handling mast cells still have evident (though reduced) expression level of mast cell associated genes. Thus, we refer to these cells as lipid associated mast cells in the remainder of the discussion.

Unfortunately, we cannot detect these two populations of adipose mast cells using conventional techniques like flow cytometry or Toluidine blue staining in our murine groups. Flow cytometry is difficult in adipose mast cells, as the collagenase that is used for tissue dissociation can cleave the mast cell surface proteins (**Supplemental Figure 5**). This is one reason we believe there is very limited data reported on murine adipose mast cells by flow cytometry. To our knowledge, this is only the second paper to attempt to quantify murine adipose mast cells by flow cytometry[44]. There is also substantial autofluorescence in adipose samples[45], contributing to measurement noise. Thus, more sensitive techniques are needed to better study adipose mast cell populations.

Nevertheless, we found that the percent of cells expressing CD36, CD9, MHCII, and CD74 changes with weight loss and weight cycling, suggesting that weight stability status may influence adipose mast cell lipid handling and antigen presentation. One group has previously shown the presence of lipid droplets in mast cells following 6 days of insulin treatment[46,47]. However, the specific expression of these lipid handling genes was not reported in the studies and little work has been done to show a mechanism beyond ER stress. Interestingly, we did not observe increased plasma insulin in weight cycled mice, and actually detect reduced insulin secretion from the pancreatic beta cells in response to maximal stimulation[34], suggesting that weight cycling may be different from chronic hyperinsulinemia in our cells. Future work could

examine how different signals such as insulin, glucose, fatty acids, and leptin within the adipose environment affect lipid breakdown, storage, and biosynthesis in mast cells. In macrophages, activation with signals like LPS or IFNγ increase triglyceride storage and lipid droplet formation[48,49], while macrophages from obese fat upregulate both lipid droplet storage and lysosomal degradation of lipids[50]. Thus, there may be different lipid signaling and metabolism programs in both mast cells and macrophages under different activation states. Further functional analyses should be used to identify the different lipid programming profiles of adipose tissue mast cells.

Excitingly, two adipose mast cell populations were observed in subcutaneous adipose samples from obese women tested in our pilot work, which increase the potential translation of our findings. The population of cells with intermediate cKit expression was significantly associated with the weight variance of these subjects, similar to the lipid associated mast cells observed by scRNAseq. Moreover, this population had higher expression of CD68, and a trend towards increased CD36, which was also observed in the scRNAseq data. However, there was lower expression of CD9 in the cKit intermediate mast cells than the population with higher Kit expression. We should emphasize that our murine data was primarily collected from male epididymal adipose tissue while our human data was collected from female subcutaneous adipose tissue, and thus differences may derive from differences in sex, adipose depots, and/or organism. Interestingly, a study by Lopez-Perez et al. showed a change in adipose mast cells in adults with obesity and type 2 diabetes. In their first cohort, a reduction in classical adipose mast cells from omental fat was observed[51]. Upon follow-up, the authors noticed that omental adipose mast cells had reduced surface expression of

CD45, cKit, and FcεR1[52]. However, it is not clear if they also observed an intermediate population in their participants or if the effects of weight cycling are required to induce this intermediate population.

Together, our data suggest that future studies and new techniques are needed to understand the role of adipose mast cells in obesity and weight cycling, as changes in lipid and antigen presentation gene expression suggest there may be different functional programs. The role of mast cells in obesity is still unclear because the use of different mast cell knockout models and different concentrations of mast cell stabilizers have produced contradictory results[25–29]. Moreover, flow cytometry and microscopy do not allow us to see if there are multiple populations of mast cells in the adipose tissue of weight cycled animals. Future plans include developing an in vitro model of lipid handling and using a reporter model system to identify, trace, and isolate mast cells without the reliance on antibodies. Moreover, we plan to measure and isolate the mast cell populations in adipose tissue from more human samples. Functional studies can then be performed to determine if and how lipid associated mast cells differ from classical mast cells and macrophages. Moreover, future studies should collect the adipose immune fractions from multiple adipose depots in both women and men. In sum, we are hopeful that future studies can isolate this novel population of lipid handling mast cells and help determine if adipose mast cells contribute to the impairment in systemic metabolism following weight cycling in mice and humans.

Authorship

HLC conceptualized the study, obtained and analyzed the data, drafted the manuscript, and is the guarantor of the work. MAC assisted with sequencing data collection, analysis, and interpretation. AMB assisted with culture model troubleshooting, data collection, and analysis. HJS provided the weight cycling questionnaire and MM provided the human samples. Both assisted with study design and data analysis. AHH assisted with study design and data interpretation and provided funding. All authors contributed to manuscript revisions.

Acknowledgements

This project was funded by a Veterans Affairs Merit Award 5I01BX002195 to AHH. AHH is also supported by a Career Scientist Award from the Veterans Affairs (IK6 BX005649). HLC was funded by an American Heart Association Postdoctoral Fellowship (20POST35120547), MAC was funded by an NIH F31 Predoctoral Fellowship (1F31DK123881), and both were previously supported by the Molecular Endocrinology Training Program (T32 DK07563). Additionally, the human mast cell work was funded by The Vanderbilt Institute for Clinical and Translational Research (VICTR) grants to HLC and MM and a NIH KL2 (KL2TR002245) to MM. VICTR is funded by the National Center for Advancing Translational Sciences (NCATS) Clinical Translational Science Award (CTSA) Program, Award Number 5UL1TR002243-03. Figure schematics were created with Biorender.com.

Conflict of Interest Disclosure

The authors declare no conflicts of interest.

References

[1] C. Martin, K. Herrick, N. Sarafrazi, CL Ogden, Attempts to Lose Weight Among Adults in the United States, 2013–2016, NCS Data Briefs. 313 (2018). https://www.cdc.gov/nchs/products/databriefs/db313.htm (accessed September 23, 2022).

[2] D.M. Quinn, R.M. Puhl, M.A. Reinka, Trying again (and again): Weight cycling and depressive symptoms in U.S. adults, PLOS ONE. 15 (2020) e0239004. https://doi.org/10.1371/journal.pone.0239004.

[3] R.R. Wing, J.O. Hill, Successful weight loss maintenance, Annu. Rev. Nutr. 21 (2001) 323–341. https://doi.org/10.1146/annurev.nutr.21.1.323.

[4] A. Fildes, J. Charlton, C. Rudisill, P. Littlejohns, A.T. Prevost, M.C. Gulliford, Probability of an Obese Person Attaining Normal Body Weight: Cohort Study Using Electronic Health Records, Am J Public Health. 105 (2015) e54-59. https://doi.org/10.2105/AJPH.2015.302773.

[5] D. Crawford, R.W. Jeffery, S.A. French, Can anyone successfully control their weight? Findings of a three year community-based study of men and women, Int. J. Obes. Relat. Metab. Disord. 24 (2000) 1107–1110.

[6] A. Stunkard, The Results of Treatment for Obesity: A Review of the Literature and Report of a Series, A.M.A. Archives of Internal Medicine. 103 (1959) 79. https://doi.org/10.1001/archinte.1959.00270010085011.

[7] J.P.H. Wilding, R.L. Batterham, M. Davies, L.F. Van Gaal, K. Kandler, K. Konakli, I. Lingvay, B.M. McGowan, T.K. Oral, J. Rosenstock, T.A. Wadden, S. Wharton, K. Yokote, R.F. Kushner, S. 1 S. Group, Weight regain and cardiometabolic effects after withdrawal of semaglutide: The STEP 1 trial extension, Diabetes, Obesity and Metabolism. 24 (2022) 1553–1564. https://doi.org/10.1111/dom.14725.

[8] M. Lahti-Koski, S. Männistö, P. Pietinen, E. Vartiainen, Prevalence of Weight Cycling and its Relation to Health Indicators in Finland, Obesity Research. 13 (2005) 333–341. https://doi.org/10.1038/oby.2005.45.

[9] S. Bangalore, R. Fayyad, R. Laskey, D.A. DeMicco, F.H. Messerli, D.D. Waters, Body-Weight Fluctuations and Outcomes in Coronary Disease, New England Journal of Medicine. 376 (2017) 1332–1340. https://doi.org/10.1056/NEJMoa1606148.

[10] J.-P. Montani, Y. Schutz, A.G. Dulloo, Dieting and weight cycling as risk factors for cardiometabolic diseases: who is really at risk?, Obes Rev. 16 Suppl 1 (2015) 7–18. https://doi.org/10.1111/obr.12251.

[11] E.-J. Rhee, Weight Cycling and Its Cardiometabolic Impact, Journal of Obesity & Metabolic Syndrome. 26 (2017) 237–242. https://doi.org/10.7570/jomes.2017.26.4.237.

[12] E.-J. Rhee, J.H. Cho, H. Kwon, S.E. Park, C.-Y. Park, K.-W. Oh, S.-W. Park, W.-Y. Lee, Increased risk of diabetes development in individuals with weight cycling over 4 years: The Kangbuk Samsung Health study, Diabetes Research and Clinical Practice. 139 (2018) 230–238. https://doi.org/10.1016/j.diabres.2018.03.018.

[13] P. Rzehak, C. Meisinger, G. Woelke, S. Brasche, G. Strube, J. Heinrich, Weight change, weight cycling and mortality in the ERFORT Male Cohort Study, Eur. J. Epidemiol. 22 (2007) 665–673. https://doi.org/10.1007/s10654-007-9167-5.

[14] W.R. Bolus, A.H. Hasty, Contributions of Innate Type 2 Inflammation to Adipose Function, Journal of Lipid Research. (2018) jlr.R085993. https://doi.org/10.1194/jlr.R085993.

[15] A.W. Ferrante, Macrophages, fat, and the emergence of immunometabolism, Journal of Clinical Investigation. 123 (2013) 4992–4993. https://doi.org/10.1172/JCI73658.

[16] L. Russo, C.N. Lumeng, Properties and functions of adipose tissue macrophages in obesity, Immunology. 155 (2018) 407–417. https://doi.org/10.1111/imm.13002.

[17] M. Cottam, H. Caslin, N. Winn, A. Hasty, Multiomics reveals persistence of obesity-associated immune cell phenotypes in adipose tissue during weight loss and subsequent weight regain, Nature Communications. 13 (2022) 2950. https://doi.org/10.1038/s41467-022-30646-4.

[18] E.K. Anderson, D.A. Gutierrez, A. Kennedy, A.H. Hasty, Weight cycling increases T-cell accumulation in adipose tissue and impairs systemic glucose tolerance, Diabetes. 62 (2013) 3180–3188. https://doi.org/10.2337/db12-1076.

[19] M. Krystel-Whittemore, K.N. Dileepan, J.G. Wood, Mast Cell: A Multi-Functional Master Cell, Frontiers in Immunology. 6 (2016). https://doi.org/10.3389/fimmu.2015.00620.

[20] S.J. Galli, M. Tsai, IgE and mast cells in allergic disease, Nature Medicine. 18 (2012) 693–704. https://doi.org/10.1038/nm.2755.

[21] M.R. Hepworth, M. Maurer, S. Hartmann, Regulation of type 2 immunity to helminths by mast cells, Gut Microbes. 3 (2012) 476–481. https://doi.org/10.4161/gmic.21507.

[22] S.N. Abraham, R. Malaviya, Mast cells in infection and immunity., Infection and Immunity. 65 (1997) 3501.

[23] M. Urb, D.C. Sheppard, The role of mast cells in the defence against pathogens, PLoS Pathog. 8 (2012) e1002619.

[24] I. Bot, G.-P. Shi, P.T. Kovanen, Mast Cells as Effectors in Atherosclerosis, Arteriosclerosis, Thrombosis, and Vascular Biology. 35 (2015) 265–271. https://doi.org/10.1161/ATVBAHA.114.303570.

[25] J. Liu, A. Divoux, J. Sun, J. Zhang, K. Clément, J.N. Glickman, G.K. Sukhova, P.J. Wolters, J. Du, C.Z. Gorgun, A. Doria, P. Libby, R.S. Blumberg, B.B. Kahn, G.S. Hotamisligil, G.-P. Shi, Genetic deficiency and pharmacological stabilization of mast cells reduce diet-induced obesity and diabetes in mice, Nat Med. 15 (2009) 940–945. https://doi.org/10.1038/nm.1994.

[26] A. Tanaka, Y. Nomura, A. Matsuda, K. Ohmori, H. Matsuda, Mast cells function as an alternative modulator of adipogenesis through 15-deoxy-delta-12, 14-prostaglandin J $_2$, American Journal of Physiology-Cell Physiology. 301 (2011) C1360–C1367. https://doi.org/10.1152/ajpcell.00514.2010.

[27] D.A. Gutierrez, S. Muralidhar, T.B. Feyerabend, S. Herzig, H.-R. Rodewald, Hematopoietic Kit Deficiency, rather than Lack of Mast Cells, Protects Mice from Obesity and Insulin Resistance, Cell Metabolism. 21 (2015) 678–691. https://doi.org/10.1016/j.cmet.2015.04.013.

[28] J. Chmelař, A. Chatzigeorgiou, K.-J. Chung, M. Prucnal, D. Voehringer, A. Roers, T. Chavakis, No Role for Mast Cells in Obesity-Related Metabolic Dysregulation, Front. Immunol. 7 (2016). https://doi.org/10.3389/fimmu.2016.00524.

[29] S.M. El-Haggar, W.F. Farrag, F.A. Kotkata, Effect of ketotifen in obese patients with type 2 diabetes mellitus, Journal of Diabetes and Its Complications. 29 (2015) 427–432. https://doi.org/10.1016/j.jdiacomp.2015.01.013.

[30] Y. Hao, S. Hao, E. Andersen-Nissen, W.M. Mauck, S. Zheng, A. Butler, M.J. Lee, A.J. Wilk, C. Darby, M. Zager, P. Hoffman, M. Stoeckius, E. Papalexi, E.P. Mimitou, J. Jain, A. Srivastava, T. Stuart, L.M. Fleming, B. Yeung, A.J. Rogers, J.M. McElrath, C.A. Blish, R. Gottardo, P. Smibert, R. Satija, Integrated analysis of multimodal single-cell data, Cell. 184 (2021) 3573-3587.e29. https://doi.org/10.1016/j.cell.2021.04.048.

[31] S.X. Ge, D. Jung, R. Yao, ShinyGO: a graphical gene-set enrichment tool for animals and plants, Bioinformatics. 36 (2020) 2628–2629. https://doi.org/10.1093/bioinformatics/btz931.

[32] J.S. Orr, A.J. Kennedy, A.H. Hasty, Isolation of Adipose Tissue Immune Cells, JoVE (Journal of Visualized Experiments). (2013) e50707. https://doi.org/10.3791/50707.

[33] H.L. Caslin, M.A. Cottam, J.M. Piñon, L.Y. Boney, A.H. Hasty, Weight cycling induces innate immune memory in adipose tissue macrophages, (2022) 2022.07.02.498553. https://doi.org/10.1101/2022.07.02.498553.

[34] N.C. Winn, M.A. Cottam, M. Bhanot, H.L. Caslin, J.N. Garcia, R. Arrojo E Drigo, A.H. Hasty, Weight Cycling Impairs Pancreatic Insulin Secretion but Does Not Perturb Whole-Body Insulin Action in Diet-Induced Obese Mice, Diabetes. (2022) db220161. https://doi.org/10.2337/db22-0161.

[35] J.M. Yabut, E.M. Desjardins, E.J. Chan, E.A. Day, J.M. Leroux, B. Wang, E.D. Crane, W. Wong, K.M. Morrison, J.D. Crane, W.I. Khan, G.R. Steinberg, Genetic deletion of mast cell serotonin synthesis prevents the development of obesity and insulin resistance, Nat Commun. 11 (2020) 463. https://doi.org/10.1038/s41467-019-14080-7.

[36] D.A. Jaitin, L. Adlung, C.A. Thaiss, A. Weiner, B. Li, H. Descamps, P. Lundgren, C. Bleriot, Z. Liu, A. Deczkowska, H. Keren-Shaul, E. David, N. Zmora, S.M. Eldar, N. Lubezky, O. Shibolet, D.A. Hill, M.A. Lazar, M. Colonna, F. Ginhoux, H. Shapiro, E. Elinav, I. Amit, Lipid-Associated Macrophages Control Metabolic Homeostasis in a Trem2-Dependent Manner, Cell. 178 (2019) 686-698.e14. https://doi.org/10.1016/j.cell.2019.05.054.

[37] G.G. Davanzo, G. Castro, P.M.M. Moraes-Vieira, Immunometabolic regulation of adipose tissue resident immune cells, Current Opinion in Pharmacology. 58 (2021) 44–51. https://doi.org/10.1016/j.coph.2021.03.004.

[38] A.W. Ferrante, The immune cells in adipose tissue, Diabetes, Obesity and Metabolism. 15 (2013) 34–38. https://doi.org/10.1111/dom.12154.

[39] M. Mraz, M. Haluzik, The role of adipose tissue immune cells in obesity and low-grade inflammation, Journal of Endocrinology. 222 (2014) R113–R127. https://doi.org/10.1530/JOE-14-0283.

[40] A. Kosteli, E. Sugaru, G. Haemmerle, J.F. Martin, J. Lei, R. Zechner, A.W. Ferrante, Weight loss and lipolysis promote a dynamic immune response in murine adipose tissue, J Clin Invest. 120 (2010) 3466–3479. https://doi.org/10.1172/JCI42845.

[41] E.K. Anderson-Baucum, A.S. Major, A.H. Hasty, A possible secondary immune response in adipose tissue during weight cycling: The ups and downs of yo-yo dieting, Adipocyte. 3 (2014) 141–145. https://doi.org/10.4161/adip.27556.

[42] S. Thillainadesan, S. Madsen, D.E. James, S.L. Hocking, The impact of weight cycling on health outcomes in animal models: A systematic review and meta-analysis, Obes Rev. (2022) e13416. https://doi.org/10.1111/obr.13416.

[43] J. Zou, B. Lai, M. Zheng, Q. Chen, S. Jiang, A. Song, Z. Huang, P. Shi, X. Tu, D. Wang, L. Lu, Z. Lin, X. Gao, CD4+ T cells memorize obesity and promote weight regain, Cellular & Molecular Immunology. 15 (2018) 630–639. https://doi.org/10.1038/cmi.2017.36.

[44] S. Hirai, C. Ohyane, Y.-I. Kim, S. Lin, T. Goto, N. Takahashi, C.-S. Kim, J. Kang, R. Yu, T. Kawada, Involvement of mast cells in adipose tissue fibrosis, American Journal of Physiology-Endocrinology and Metabolism. 306 (2014) E247–E255. https://doi.org/10.1152/ajpendo.00056.2013.

[45] K.W. Cho, D.L. Morris, C.N. Lumeng, Flow Cytometry Analyses of Adipose Tissue Macrophages, Methods in Enzymology. 537 (2014) 297. https://doi.org/10.1016/B978-0-12-411619-1.00016-1.

[46] J.T. Aldan, C. Jansen, M. Speck, K. Maaetoft-Udsen, E.A. Cordasco, M. Faiai, L.M.N. Shimoda, W.E. Greineisen, H. Turner, A.J. Stokes, Insulin-induced lipid body accumulation is accompanied by lipid remodelling in model mast cells, Adipocyte. 8 (2019) 265–279. https://doi.org/10.1080/21623945.2019.1636624.

[47] W.E. Greineisen, K. Maaetoft-Udsen, M. Speck, J. Balajadia, L.M.N. Shimoda, C. Sung, H. Turner, Chronic Insulin Exposure Induces ER Stress and Lipid Body Accumulation in Mast Cells at the Expense of Their Secretory Degranulation Response, PLoS ONE. 10 (2015) e0130198. https://doi.org/10.1371/journal.pone.0130198.

[48] X.A.M.H. van Dierendonck, F. Vrieling, L. Smeehuijzen, L. Deng, J.P. Boogaard, C.-A. Croes, L. Temmerman, S. Wetzels, E. Biessen, S. Kersten, R. Stienstra, Triglyceride breakdown from lipid droplets regulates the inflammatory response in macrophages, Proceedings of the National Academy of Sciences. 119 (2022) e2114739119. https://doi.org/10.1073/pnas.2114739119.

[49] M. Rosas-Ballina, X.L. Guan, A. Schmidt, D. Bumann, Classical Activation of Macrophages Leads to Lipid Droplet Formation Without de novo Fatty Acid Synthesis, Front Immunol. 11 (2020) 131. https://doi.org/10.3389/fimmu.2020.00131.

[50] X. Xu, A. Grijalva, A. Skowronski, M. van Eijk, M.J. Serlie, A.W. Ferrante, Obesity Activates a Program of Lysosomal-Dependent Lipid Metabolism in Adipose Tissue Macrophages Independently of Classic Activation, Cell Metab. 18 (2013) 816–830. https://doi.org/10.1016/j.cmet.2013.11.001.

[51] D. Lopez-Perez, A. Redruello-Romero, J. Garcia-Rubio, C. Arana, L.A. Garcia-Escudero, F. Tamayo, J.D. Puentes-Pardo, S. Moreno-SanJuan, J. Salmeron, A. Blanco, J. Galvez, J. Leon, Á. Carazo, In Patients With Obesity, the Number of Adipose Tissue Mast Cells Is Significantly Lower in Subjects With Type 2 Diabetes, Frontiers in Immunology. 12 (2021). https://www.frontiersin.org/articles/10.3389/fimmu.2021.664576 (accessed October 25, 2022).

[52] D. Lopez-Perez, A. Redruello-Romero, J. Garcia-Rubio, C. Arana, L.A. Garcia-Escudero, F. Tamayo, J. Salmeron, J. Galvez, J. Leon, Á. Carazo, In Obese Patients With Type 2 Diabetes, Mast Cells in Omental Adipose Tissue Decrease the Surface Expression of CD45, CD117, CD203c, and FcℇRI, Front. Endocrinol. 13 (2022) 818388. https://doi.org/10.3389/fendo.2022.818388.

Figure 1: Weight cycling increases adipose mast cells. (A) Schematic of weight cycling mouse model commonly used by our group. (B) UMAP of mast cell cluster in each diet group. (C) Mast cell counts (number of cells sequenced). Violin plots of (D) *Cma1* and *Il1rl1* and (E) *Trem2* and *Fabp5* expression in each diet group. All data were derived from new analyses of our previously published dataset [17].

Figure 2: Histology and flow cytometry do not confirm an increase in adipose mast cells with weight cycling. (A) Body weight over time from a new murine cohort measured weekly with diet switch indicated by dashed lines in lean, obese, weight loss (WL), and weight cycled (WC) mice (n=12/group). (B) Blood glucose during an intraperitoneal glucose tolerance test (1.5 g dextrose/kg lean mass) at 27 weeks (n=12/group). (C) Representative image of Toluidine blue staining on adipose mast cell section. (D) Quantification of adipose mast cells per 104 frames (n=4-6/group). (E)

Representative flow cytometry plot of cKit+Fcer1+ mast cells in epididymal adipose tissue. (F) Quantification of epididymal adipose tissue mast cells by % and #/ gram fat (n=14-18/group). (G) Representative flow cytometry plot of cKit+Fcer1+ mast cells in subcutaneous adipose tissue. (H) Quantification of subcutaneous adipose tissue mast cells by % and #/g fat (n=8-12/group). Data are means ± SEM in 1-2 new cohorts of mice. *$p<0.05$, **$p < 0.01$, *** $p <0.001$, ****$p < 0.0001$.

Figure 3: Weight cycling induces a distinct cluster of lipid handling mast cells. (A) UMAP of two distinct adipose mast cell clusters after subsetting on mast cells from [17]. (B) Mast cell counts (number of cells sequenced) and proportion (mast cells sequenced/ total cells) of both mast cell clusters. Violin plots of (C) mast cell and macrophage genes, as well as (D) lipid handling and antigen presentation related genes in each cluster. Expression level of all genes were significantly different between cluster 1 and cluster 2 by adjusted p-value with differential expression. All data were derived from new analyses of our previously published dataset [17].

Figure 4: Weight loss and regain change lipid handling and antigen presentation proteins. Quantification of percent of mast cells and MFI by flow cytometry. for (A) the lipid handling receptors CD36 and CD9 and (B) the antigen presentation proteins MHCII (I-A/ I-E) and CD74 on mast cells in a new cohort of mice. Data are means ± SEM in n=13-17/ group from 2 independent experiments. *p<0.05, **p < 0.01, *** p <0.001

Figure 5: Weight variability correlates cKit intermediate mast cells in humans. (A) Representative flow plots for cKit⁺ FcεR1⁺, cKit^hi FcεR1⁺, and cKit^int FcεR1⁺ mast cell populations from human stromal vascular fraction. (B) Quantification of each mast cell population. (C) Quantification of percent of mast cells and MFI for the lipid handling receptors CD36 and CD9 and the antigen presentation proteins CD74 and CD68 on each mast cell population by flow cytometry. (D) Weight change plotted over time for each subject. (E) Correlation between the weight change in the previous year with the % cKit^hi FcεR1⁺ cells. (F) Correlation between weight variance and the % cKit^int FcεR1⁺ mast cells. Data are means ± SEM in n=4. *p<0.05, **p < 0.01.