Main text

The genetic architecture of the load linked to dominant and recessive self-
incompatibility alleles in _Arabidopsis halleri_ and _A. lyrata_

Le Veve Audrey[1,2], Genete Mathieu[1], Lepers-Blassiau Christelle[1], Ponitzki Chloé[1], Céline
Poux[1], Vekemans Xavier[1], Durand Eleonore[1], Castric Vincent[1*]

[1]_Univ. Lille, CNRS, UMR 8198 – Evo-Eco-Paleo, F-59000 Lille, France_

[2]_Current address: Department of Botany, Faculty of Science, Charles University, Benátská 2,_
CZ-128 01 Prague, Czechia

* Correspondence : vincent.castric@univ-lille.fr

Abstract

The long-term balancing selection acting on mating-types or sex determining genes is expected to
lead to the accumulation of deleterious mutations in the tightly linked chromosomal segments that
are locally "sheltered" from purifying selection. However, the factors determining the extent of this
accumulation are poorly understood. Here, we take advantage of the large number of alleles at the
sporophytic self-incompatibility system of the Brassicaceae along a complex dominance hierarchy to
evaluate how the pace at which linked deleterious mutations accumulate varies with the intensity of
balancing selection. We first experimentally measured the phenotypic manifestation of the linked
load at three different levels of the dominance hierarchy. We then sequenced and phased
polymorphisms in the chromosomal regions linked to 126 distinct copies of _S_-alleles in two
populations of _Arabidopsis halleri_ and three populations of _A. lyrata_. We find that linkage to the _S_-
locus locally distorts phylogenies over about 10-30kb along the chromosome. The more intense
balancing selection on dominant _S_-alleles results in greater fixation of deleterious mutations, while
recessive _S_-alleles accumulate more linked deleterious mutations that are segregating. Hence, the
structure rather than the overall magnitude of the linked genetic load differs between dominant and
recessive _S_-alleles. Our results have consequences for the long-term evolution of new _S_-alleles, the
evolution of dominance modifiers between them, and raise the question of why the non-
recombining regions of some sex and mating type chromosomes expand over evolutionary times
while others, such as that the _S_-locus of the Brassicaceae, remain restricted to small chromosomal
regions.

Keywords: Supergene, sheltered load, mating types, balancing selection, genetic dominance, S-locus.

Introduction

Sexual reproduction typically involves mating between individuals that belong to separate groups of reciprocal reproductive compatibility. This partition among conspecific individuals is obvious in species where the male and female reproductive functions are performed by distinct categories of individuals, but can also be present in purely hermaphroditic species, where compatibility among individuals is governed by the segregation of various numbers of mating types in the total absence of sexual specialisation. The existence of sexes or mating-types leads to one the strongest forms of long-term balancing selection, and is often associated with clusters of polymorphisms around sex/mating-type controlling regions kept together by structural rearrangements. In some cases, such rearrangements can span almost entire chromosomes (*e.g.* sex chromosomes in mammals[1] or mating-type chromosomes in ascomycete fungi[2]), while in others they remain limited to relatively small genomic regions (e.g. chromosomal inversions controlling male reproductive morphs in the ruff[3], mating-type loci in some basidiomycete fungi, segregating indels controlling pin *vs.* thrum floral morphs in Primula[4]). The long-term balancing selection acting on these systems is expected to lead to the accumulation of deleterious mutations in the tightly linked chromosomal segments that are "sheltered" from purifying selection by the presence of the balanced polymorphism[5,6]. These deleterious mutations can have drastic short- and long-term consequences for the evolution of the species, and determining the processes by which they accumulate is crucial to understand how the rearranged regions can either expand along the chromosomes or conversely remain restricted to limited genomic tracts[7,8].

Self-incompatibility (SI) is a genetic mechanism allowing recognition and rejection of self-pollen by hermaphrodite individuals, thereby preventing inbreeding and promoting outcrossing in hermaphroditic plant species[9]. In the Brassicaceae family, SI is controlled by a single non-recombining chromosomal region, the *S*-locus[10,11]. SI is one of the most prominent examples of long-term balancing selection[12], and as such deleterious mutations are expected to accumulate in very close genetic linkage to the *S*-alleles because of the indirect effects of linked selection[6]. Population genetics models predict that deleterious variants should accumulate within specific *S*-allele lineages[12,13], and should then be reshuffled among them by recombination. However, due to the technical difficulty of phasing polymorphisms, this process has rarely been characterised in detail[14].

A key feature of sporophytic SI systems, also shared by sex chromosomes, is the existence of dominance interactions between *S*-alleles. While most individuals are heterozygous at the *S*-locus and thus carry two different *S*-alleles, only one of them is generally expressed at the phenotypic level, especially for the pollen specificity, following a complex genetic dominance hierarchy[15,16]. Evolutionary properties of *S*-alleles vary along the dominance hierarchy because balancing selection is predicted to act more strongly on dominant than on recessive *S*-alleles, as the latter are often masked at the phenotypic level[17]. As a result, Llaurens et al.[13] and Goubet et al.[18] suggested that the dynamics of accumulation of deleterious variation may differ in close linkage with dominant *vs* recessive *S*-alleles, similarly to Y *vs.* X chromosomes, respectively. Specifically, recessive *S*-alleles can form homozygous combinations in natural populations more often than dominant *S*-alleles[17], such that recombination can occur occasionally between distinct gene copies of the same recessive *S*-allele, providing the opportunity for the linked recessive deleterious mutations to be purged from within the *S*-locus itself. In addition, because recessive *S*-alleles reach higher population

frequencies[15,17], purifying selection on linked deleterious variants is expected to have higher efficacy
among gene copies of recessive than dominant S-alleles. This is expected to result in a higher fixation
probability of deleterious variants linked to the class of dominant S-alleles than to the class of
recessive S-alleles[13]. Empirical support for this simple prediction has been conflicting, though. Based
on phenotypic measurements in A. halleri, Llaurens et al.[13] observed a decrease of fitness associated
by enforced homozygosity for one of the most dominant S-alleles (Ah15) but not for the most
recessive S-allele of the allelic series (Ah01). In contrast, Stift et al.[19] observed no effect of
dominance on the genetic load linked to three dominant vs. recessive S-alleles in a natural
population of the closely related A. lyrata. Hence, the data available so far are inconclusive, but are
restricted to very small numbers of S-alleles. They are also based on inherently limited phenotypic
measurements, seriously limiting the power of the comparisons, and preventing proper
generalisation of the effect of dominance on the accumulation of linked deleterious variation.

In this study, we combined phenotypic, genomic and theoretical approaches to finely dissect the
patterns of accumulation of deleterious variation linked to the S-locus supergene in A. halleri and A.
lyrata. We first extended the phenotypic approach of Llaurens et al.[13] to a series of additional S-
alleles from the same local A. halleri population to evaluate the effect of S-allele dominance on the
sheltered load. We then used parent-offspring trios and targeted genome re-sequencing to directly
quantify the accumulation of putative deleterious mutations linked to phased dominant vs. recessive
S-alleles in two A. halleri and three A. lyrata natural populations. Finally, we used stochastic
simulations to refine the theoretical predictions about the effect of S-allele dominance on the
dynamics of linked deleterious mutations. Overall, our results provide a more nuanced view of the
effect of the intensity of balancing selection on the sheltered load, in which the structure of the
sheltered load rather than its magnitude differs among S-alleles from different dominance classes.

Results

The genetic load linked to the S-locus varies among S-alleles, but is not correlated with dominance.

We first expanded the experimental approach of Llaurens et al.[13] to phenotypically evaluate the
effect of S-allele dominance on the intensity of the sheltered load. The previous study focused on
three S-alleles (Ah01, Ah02 and Ah15)[13]. Here we included two S-alleles from the same local
population (Nivelle, France): Ah03 and Ah04, and included Ah01 again for comparative purposes. In
the Arabidopsis genus, S-alleles have been shown to form a complex dominance hierarchy[15,16]. This
hierarchy is largely associated with the phylogeny of S-alleles[15], and at least four phylogenetic classes
(I, II, III and IV) have been described, from the most recessive (class I) to the most dominant of S-
alleles (class IV). Dominance interactions also exist among S-alleles within classes, such that these
five S-alleles form the following dominance hierarchy[15,16]: Ah01<Ah03<Ah02<Ah04<Ah15, from the
most recessive (Ah01) to the most dominant (Ah15). To reveal the linked load, we enforced
homozygosity at the S-locus using controlled crosses between parental individuals sharing a given S-
allele that was masked by different dominant S-alleles (e.g., to obtain Ah_xAh_x homozygotes we
deposited pollen from a Ah_xAh_y plant, where $Ah_y>Ah_x$, on pistils of a Ah_xAh_z plant where $z≠y$, or on a
Ah_xAh_x pistil when available; table S1). We obtained 399 offspring from a total of six such crosses.
Note that our experimental procedure differs slightly from that of Llaurens et al.[13] in that their
procedure required a CO_2 treatment to bypass the SI system and obtain selfed offspring, while here
we took advantage of the dominance interactions to obtain outcrossed S-locus homozygous

individuals that we phenotypically compared to their full-sibs with S-locus heterozygous genotypes.
Note also that the S-locus homozygous offspring we obtained contain distinct gene copies of a given
S-allele lineage. Hence, they could in principle carry distinct suites of linked deleterious mutations in
case these mutations segregate within S-allele lineages.

We first tested whether homozygosity at the S-locus affected survival by measuring for each cross
the proportion of homozygotes at the S-locus reaching the reproductive stage for three S-alleles (in
two replicate families per S-allele, Table S1). The proportion of Ah01/Ah01 and Ah04/Ah04
homozygotes surviving to the reproductive stage was consistent with mendelian expectations in
their respective families. However, we observed a significant decrease of Ah03/Ah03 homozygotes
at the reproductive stage compared with Mendelian expectations (Table 1), whereas the observed
proportion of the Ah03 S-allele among heterozygous individuals did not depart from expectations
(2/3=0.67; Table 1). Thus, the increased mortality is associated with Ah03 homozygosity, rather than
with a lower performance of individuals carrying the Ah03 S-allele itself. Overall, a genetic load was
thus observed linked to the Ah03 S-alleles, which is at an intermediate level of dominance, but
neither to the most dominant (Ah04) nor to the most recessive (Ah01) S-allele. Hence, these
observations do not support a positive correlation between S-allele dominance and the magnitude
of the sheltered load.

Table 1: Proportion of S-locus homozygous offspring having reached the reproductive stage for
three different S-alleles. *The test is performed relative to the expected proportion of homozygous*
genotypes in the offspring (25% when both parents are heterozygous; 50% when one of the parents
is homozygous and the other heterozygous).

S-allele	Level of dominance	Number of seedlings having reached the reproductive stage	Observed proportion of homozygotes	Ratio of the observed to expected proportion of homozygous genotypes (p-values[a])	Observed proportion of heterozygotes with the S-allele (p-values compared to expected)
Ah01	I	35	0.29	1.14 (0.70; 0.76)	19 (0.40; 0.90)
Ah03	II	27	**0.074**	**0.3 (0.04; 0.02)**	17 (1; 0.69)
Ah04	III	96	0.479	0.96 (0.75; 0.39)	50

[a] *p-values were obtained by binomial test and after 10,000 random permutations, respectively.*
Values departing from Mendelian expectations are figured in bold. For Ah04, the maternal parent
was Ah04/Ah04, so all heterozygous offspring carried the S-allele.

Next we measured thirteen vegetative and reproductive traits in the resulting families and compared
offspring that were homozygous for their S-alleles with their full sibs that were heterozygous (Fig. 1).
We first used permutations to test whether the mean trait value of homozygotes differed from that
in heterozygotes. Overall, with a single exception, we found no effect of homozygosity at the S-locus
on variation of the traits measured (Fig. 1; Table S2). The maximum length of flowering stems was
the exception to this general pattern, with longer reproductive stems for S-locus homozygous than
heterozygous genotypes, hence in the opposite direction from our expectation of lower fitness in

homozygotes. For this trait, there was significant variation among replicate families for homozygotes
of the recessive allele Ah01 but not of the dominant allele Ah04 (Table S3). We then used
generalised linear models (GLM) to evaluate the effect of dominance (as a fixed effect) on the mean
phenotypic value of homozygotes compared to heterozygotes for each trait (Table S4; treating
family of origin, attacks by phytopathogens, phytophagous and oxidative stress as random effects
whenever necessary). We also observed no effect of S-allele dominance on the contrast between S-
locus homozygotes and heterozygotes for any of these traits. A single of the thirteen traits was an
exception to this general pattern, but again the effect was in the opposite direction from our
expectation, with an earlier rather than delayed appearance of the first leaf for homozygotes of
more dominant S-alleles; Table S4). Overall, our phenotypic results confirmed the presence of a
detectable linked load on some phenotypic traits (survival; time to produce the first leaf), but we
could not replicate the observation of Llaurens et al.[13] that dominant S-alleles carry a more severe
deleterious load than recessive S-alleles, even though our samples were obtained from the same
local population.

Figure 1: Effect of homozygosity at the S-locus on 13 phenotypic traits compared to heterozygotes.
For each trait, the phenotypic values in homozygotes (in grey) were normalised relative to the mean
phenotypic values in heterozygotes (in black). The distributions were obtained by 10,000 random
permutations. SD: Standard Deviation.

S-alleles are associated with specific sets of tightly linked mutations.

The model of the sheltered load assumes that distinct S-allele lineages carry specific sets of linked
deleterious mutations, but to our knowledge this prediction was never tested directly. We combined
a parent-offspring trio approach with sequencing of the S-locus flanking regions to phase the
mutations segregating in the S-locus flanking regions with their respective S-alleles. Briefly, we used
a previously developed sequence capture protocol specifically targeting the nucleotide sequences
over 75 kb on each side of the S-locus along with a series of 100 control regions from throughout the
genome[20], and we analysed nucleotide sequence polymorphism (including only invariant and biallelic
SNPs), based on the A. lyrata reference genome[21]. We define a haplotype as a unique combination
of mutations along the phased chromosome, and a S-allele lineage as the collection of gene copies of
a given functional S-allele (different functional S-alleles are distinguished based on their strong
sequence divergence at the S-locus pollen and pistil genes). Different gene copies within an S-allele
lineage can thus be associated with distinct linked haplotypes in the flanking regions. The S-alleles
were identified based on short reads sequences according to a previously published method[41]. We
analysed two closely related A. halleri populations from Europe (Nivelle and Mortagne) and three
allogamous A. lyrata populations from North America (IND, PIN and TSS[22]). Overall, we were able to
reconstruct 34 haplotypes linked to a total of 12 distinct S-allele lineages in Nivelle, 38 haplotypes
linked to 11 distinct S-allele lineages in Mortagne and 16, 22 and 16 haplotypes associated with 6, 7
and 5 distinct S-allele lineages in populations IND, PIN and TSS, respectively (Table S5). Nine of the S-
alleles were shared between the two A. halleri populations (Ah01, Ah03, Ah04, Ah05, Ah12, Ah20,
Ah24, Ah25 and Ah59). In the populations of A. lyrata, four S-alleles were shared between PIN and
TSS (Ah01*, Ah03*, Ah18* and Ah63*), five S-alleles were shared between PIN and IND (Ah01*,
Ah03*, Ah46* and Ah63*), four S-alleles were shared between IND and TSS (Ah01*, Ah03*, Ah31*
and Ah63*), and three were shared across all three (Ah01*, Ah03* and Ah63*). Note that for
convenience, we used A. halleri notations (with the addition of a *) to refer to the trans-specifically
shared A. lyrata S-alleles. Altogether, we were able to obtain the phased flanking sequences of 126
S-locus haplotypes, comprising a total of 4,854 variable sites. This provides considerable power to
evaluate the local accumulation of linked mutations across S-alleles of different levels of dominance
and to examine their patterns of conservation between populations and between species.

Mutations in the S-locus flanking regions can be exchanged between S-alleles by recombination, and
between local populations by migration[23]. The relative time scale of these two processes
(recombination vs. migration) determines the distribution of the linked mutations. To capture the
chromosomal extent of this effect of linkage to S-alleles, we developed a new phylogenetic method
comparing the likelihood of two contrasted topologies of interest in overlapping windows along the
chromosome: (1) the topology clustering haplotypes by the populations where they came from vs.
(2) the topology clustering them by the S-allele to which they are linked (Fig 2A). This allowed us to
evaluate the progressive shift from a predominant topology by S-alleles close to the S-locus to a
topology by populations further along the chromosome and in unlinked control regions (Fig. 2B). The
difference in log likelihood between the two topologies decreased significantly with distance to the
S-locus (Pearson coefficient = -0.015 and -0.010 for A. halleri and A. lyrata respectively; p-values $<2^{e-16}$
). In A. halleri, the topology grouping haplotypes by populations became more likely than the
topology grouping them by S-alleles at a distance of around 30kb from the S-locus, but even at a
distance of 50kb the phylogenetic structure was still different from that in regions unlinked to the S-
locus used as controls for the genomic background[20] (Fig 2B). In A. lyrata, the shift was even more
rapid (within 10-15kb), although we note that the phylogenetic structure of the control regions was

less resolved (Fig 2B). To evaluate these patterns more directly we first examined the data using a
Major Component Analysis (MCA, a modified version of PCA adapted to binary data, Fig S1 and S2)
and using simple phylogenetic reconstructions (Fig S3, S4, S5 and S6). We confirmed that haplotypes
linked to a given S-allele tended to cluster together in the most tightly linked region, and that this
grouping by S-alleles was progressively lost in favour of a grouping by population of origin in the
most distant regions. Following Kamau et al.[24], we compared the fixation index F_{ST} among local
populations and among S-alleles in *A. lyrata* and *A. halleri*. In both *A. halleri* and *A. lyrata*, F_{ST} values
among S-alleles were high in regions close to the S-locus and quickly decreased to reach the
background level (Fig. S7) as the distance from the S-locus increased. In parallel, the differentiation
among populations followed roughly the opposite pattern, i.e. it was initially low in regions close to
the S-locus (as expected under strong balancing selection) and increased up to background level
within the first few kilobases (Fig. S7). In line with our phylogenetic analysis, differentiation between
populations started to exceed differentiation between S-alleles much closer to the S-locus in the *A.*
lyrata than in the *A. halleri* populations (Fig S3, S4, S5 and S6). Finally, we explored the fine-scale
patterns of association within populations between individual S-alleles and SNP in the linked and the
control regions (Fig S8). As expected, the vast majority of significant associations were found for the
most closely linked SNPs. With a single exception, all S-alleles were associated with unique SNPs in
the 50kb region around the S-locus, albeit with substantial heterogeneity among S-alleles in the
patterns and extent of associations that they show (Fig S8). Overall, our results indicate that due to
limited recombination, the S-alleles carry a specific set of polymorphic sites in the linked region. This
association fades away for more distant sites over a few kilobases, where population structure
becomes predominant, as in the rest of the genome. Hence, different S-alleles are associated with
specific sets of tightly linked mutations, but only within 10-30kb.

Figure 2: Linkage to the S-locus locally distorts the phylogenetic relationships. *A. The two
topologies of interest cluster haplotypes either by the S-allele to which they are linked (top) or by the
populations where they came (bottom). Different S-alleles are represented by symbols of different
colours, different populations of origin are represented by symbols of different shapes. B. Difference
in log likelihood of the two topologies of interest. Dots correspond to the difference in log likelihood
for overlapping series of 50 SNPs around the S-locus for A. halleri (top panel) and A. lyrata (bottom
panel). Positive values correspond to chromosomal positions where the topology by S-alleles explains
the phylogeny of haplotypes better than the topology by populations. The right panels show the
difference in log likelihood in the control regions. 2.5 and 97.5 percentiles of the distribution in the
control regions are indicated by dashed lines.*

No overall evidence that dominant S-alleles accumulate more linked deleterious mutations.

Llaurens et al.[13] predicted a positive correlation between the dominance of *S*-alleles and their
tendency to fix linked deleterious mutations. Thus, we investigated the correlation between the
level of dominance of the *S*-alleles and their total number of 0-fold degenerate mutations (S_{0f}) or the
ratio of 0-fold to 4-fold mutations (S_{0f}/S_{4f}) for the phased haplotypes, assuming that the vast majority
of 0-fold degenerate mutations are deleterious. Based on the results presented above and the
results of our previous study[20], for the rest of our analyses we focused on the phased haplotypes
over the first 25 kb on either side of the *S*-locus. We found no overall effect of dominance on S_{0f} (p-
values= 0.54 and 0.07 for *A. halleri* and *A. lyrata* respectively; Fig. 3; Table S6) or S_{0f}/S_{4f} (p-values=
0.54 and 0.07 for *A. halleri* and *A. lyrata* respectively; Table S6). Extending the analysis to all non-
synonymous mutations led to identical conclusions (Table S6). Overall, our genomic results did not
confirm the prediction that dominant *S*-alleles accumulate a larger number of putatively deleterious
mutations in their linked regions. We note that the particular *S*-allele whose sheltered load was
quantified in Llaurens et al.[13] (Ah15, red arrow on Fig 3A) appears to be one of the *S*-alleles
associated with the highest number of 0-fold degenerate mutations among all *S*-alleles of the most
dominant class (class IV).

Figure 3: No overall effect of S-allele dominance on the total number of 0-fold degenerate mutations (S_{0f}) in the linked genomic regions within 25kb. Each dot represents the mean number of mutations observed among haplotypes linked to one S-allele in one population. The correlations evaluated by a GLM are represented by lines, with confidence intervals represented in grey. A: A. halleri. Black dots correspond to the Nivelle population, grey dots to the Mortagne population. The red arrow points to the copy of Ah15, corresponding to the S-allele whose sheltered load was phenotypically characterised by Llaurens et al.[13] in Nivelle. B: A. lyrata. Red dots correspond to the IND population, black dots to the PIN population and blue dots to the TSS population.

The structure of the linked genetic load differs between dominant and recessive S-alleles.

Theory predicts that dominant S-alleles should fix linked recessive deleterious mutations with a higher probability than recessive S-alleles[13], but in natural populations we observed no difference in the total number of putatively deleterious linked to dominant *vs.* recessive S-alleles. To clarify this discrepancy, we took advantage of our sequencing of multiple copies of S-alleles to consider separately the fixed and the segregating mutations linked to each of the S-allele lineages. For each population, we included only mutations that were segregating, and excluded those that were locally fixed. In agreement with the prediction of Llaurens et al.[13], we observed that lineages of dominant S-alleles do indeed tend to fix deleterious mutations more readily (Fig. 4). The fact that they do not accumulate a larger total number of deleterious mutations is explained by the fact that the structure of the genetic load differs between dominant and recessive S-alleles: the dominant S-alleles tend to have more fixed deleterious mutations, but the recessive S-alleles compensate by accumulating a larger number of segregating mutations, resulting in similar numbers of deleterious mutations overall in most of the populations.

Figure 4: The number of 0-fold degenerate mutations fixed in the 25kb regions flanking the S-locus
increases with dominance of the S-allele associated. Each dot represents the value obtained for
haplotypes linked to one S-allele in one population. The correlations evaluated by a GLM are
represented by lines, with confidence intervals represented in grey. A: A. halleri. Black dots
correspond to the Nivelle population, grey dots to the Mortagne population. B: A. lyrata. Red dots
correspond to the IND population, black dots to the PIN population, blue dots to the TSS population.

Motivated by these empirical observations, we extended the model proposed by Llaurens et al.[13] to
predict the dynamics of accumulation of recessive deleterious mutations linked to S-alleles, focusing
not only on fixed deleterious mutations but also on those that are segregating within allelic lineages.
These stochastic simulations confirmed that, at equilibrium, dominant S-alleles tend to accumulate a
larger number of deleterious mutations that are fixed among gene copies within S-allele lineages
(Fig. 5A). In contrast, the number of segregating linked mutations was higher for recessive than for
dominant S-alleles (Fig. 5B). Our model predicted that these two effects eventually compensate each
other, such that in the end the total number of linked deleterious mutations was not expected to
change with dominance of the S-alleles (Fig. 5C). These predictions are in line with our genomic
observations and suggest that the dominance level of S-alleles modifies the structure of the genetic
load they shelter: dominant S-alleles accumulate more fixed deleterious mutations, but recessive S-
alleles accumulate more segregating mutations, resulting in an equivalent total load.

Figure 5: Stochastic simulations confirm the contrasted architecture of the load of deleterious mutations linked to dominant vs. recessive S-alleles. *Number of fixed (A), segregating (B) and total (C) deleterious mutations linked to S-alleles at four different levels of dominance (I<II<III<IV). The means (bold lines) were estimated per S-allele dominance classes over 100 replicate simulations after discarding an initial burn-in of 100,000 generations. h=0. s=0.01.*

Discussion

The genetic load linked to the S-locus is detectable and is manifested on different phenotypes

Our results contribute to a still restricted but growing body of evidence confirming that the accumulation of deleterious mutations linked to strongly balanced allelic lines can be substantial, and that their effect can be detected at the phenotypic level[13,19,25-28]. An interesting observation is that the phenotypes on which the load was revealed varied among these studies. Here, the effect of homozygosity at the *S*-locus was apparent on juvenile survival and on the length of the longest flowering stem, but we detected no effect on any other morphological measurements, including leaf and rosette traits. In the same population of *A. halleri,* Llaurens et al.[13] detected an effect on juvenile survival and on leaf size. A study in North American outcrossing populations of *A. lyrata*[19] detected an effect on juvenile survival, but not on any other traits that they measured. In the horsenettle *Solanum carolinense*, the load was associated with reduced seed viability, flower number and germination[25,28]. Hence, the most consistent pattern seems to be a decrease of overall juvenile survival, possibly because it is a highly integrative measurement of fitness, whereas other morphological or life history traits can be associated with more specific components of overall fitness.

A unique genetic load associated with each allele in each population.

The model of the sheltered load posits that each *S*-allele should be associated with a specific set of linked mutations[13]. In line with this prediction, another consistent observation is that the magnitude of the *S*-linked load varied among *S*-alleles, as the load linked to some *S*-alleles was phenotypically detectable, while for others it was not. This variation of the genetic load is expected since deleterious mutations associated with the different alleles are likely to hit different linked genes, and affect different phenotypic traits with different effects on fitness. Also in line with the model of the sheltered load, our phasing of a large number of variants linked to *S*-haplotypes in several natural populations revealed that the same suite of linked mutations was consistently associated among different copies of a given allele when sampled from within the same population, in particular for the dominant *S*-allele lineages under more intense balancing selection. As expected for outcrossing populations with short-scale linkage disequilibrium, this association was lost when examining sites at increasing genetic distances from the *S*-locus along the chromosome (see also Le Veve et al.[20]). However, a proper model of sheltered load taking into account recombination among *S*-alleles is still missing. Finally, the association with linked sites was further lost when comparing gene copies of *S*-alleles sampled from different local populations, suggesting that recombination within populations decouples alleles from their linked sites faster than migration can homogenise the genetic composition among these natural populations. We note that the patterns of association and phylogenetic structure differed among populations, possibly due to their contrasted demographic histories. Indeed, the *A. lyrata* populations colonised North America from ancestral European populations about 20-30.000 years ago[29,30], and are less diverse overall than the *A. halleri* populations we studied, who colonised the north of France during the last century from ancestral German populations[31]. The progressive decoupling between alleles and their linked sites leads to the simple prediction that *S*-locus homozygous genotypes formed by crossing individuals carrying identical alleles from distinct populations should not reveal as much load as when they are formed

by crossing individuals within populations. Hence, the S-locus region could contribute to overall
hybrid vigour. Testing this prediction will be an interesting next step.

Different properties of the linked load according to S-allele dominance

The question of whether S-allele dominance, which modifies the intensity of balancing selection in
the S-locus region, could explain variation of the linked load has received conflicting support in the
literature. In line with Stift et al.[19], but in contradiction with Llaurens et al.[13], we observed no overall
effect of dominance on the magnitude of the load. Several technical and biological reasons could
explain the contrasted results obtained in these different studies. First, phenotypic quantification of
the linked load is experimentally demanding, such that these studies relied on the comparison of a
limited number of alleles (three S-alleles in each of the studies) and therefore each of them had
inherently low power. Second, the experimental procedures to reveal the load varied slightly.
Llaurens et al.[13] used CO_2 treatment to by-pass the SI system and obtain homozygous progenies from
crosses that would otherwise have been incompatible, whereas we used the "natural" masking by
dominant S-alleles to enable the obtention of recessive homozygous genotypes. Our approach is
experimentally simpler and avoids the possible contamination by offspring obtained by selfing,
which may combine the effect of the sheltered load with that of genome-wide inbreeding
depression (see Stift et al.[19] for a detailed discussion of this caveat). Third, a limitation of our
approach is that it is restricted to S-alleles that are recessive or intermediate along the dominance
hierarchy, and is thus not applicable to quantify the load associated with the most dominant S-alleles
under more intense balancing selection. It is therefore possible that the S-alleles we examined did
not exhibit sufficiently contrasted levels of dominance, in particular if only the most dominant ones
are generating a substantial load, as suggested for fully linked deleterious mutations. In addition,
since the homozygous S-allele genotypes we create correspond to different gene copies from the
population, they may carry distinct sets of linked deleterious variants, especially for the more
recessive alleles. The variation we observed in the phenotypic magnitude of the load among families
confirms that linked deleterious variants are unlikely to be fixed within all allele lineages. Finally, we
note that our genomic analysis of the genetic load shows that the dominant allele Ah15 previously
associated with reduced fitness in homozygotes[13], is indeed unusual in terms of the number of
mutations it carries. In fact, it is one of the most "loaded" alleles among all the dominant S-alleles
present in this population, possibly explaining why Llaurens et al.[13] observed a significant effect
despite the inherently limited experimental power of their analysis.

Our stochastic simulations and genomic analyses concur to the conclusion that the intensity of
balancing selection applied to each allele affects the genetic architecture of the linked load: a larger
proportion of putatively deleterious mutations are fixed among gene copies of the dominant as
compared to the recessive S-alleles, while gene copies of the recessive S-alleles tend to accumulate
more segregating deleterious variation. While these two processes eventually compensate one
another, they may have distinct consequences for the evolution of S-alleles. Uyenoyama[12] showed
that the existence of a sheltered load should influence the evolutionary dynamics of new S-alleles
through self-compatible intermediates. Specifically, antagonistic interactions are expected between
ancestral and derived functional specificities because they would initially share their linked
deleterious mutations, slowing down the establishment of new alleles. Our observation that partially
different sets of linked mutations are associated with S-alleles from the different populations raises

the question of whether the (short) time scale at which recombination decouples alleles from their
sets of linked mutation is sufficiently fast to impede such antagonistic interactions to take place. In
other words, the effect of the load on the diversification dynamics should be most important if the
two mutational steps required for the emergence of new alleles under this model take place within
local populations, rather than involving a metapopulation-scale process. As shown by Stetsenko et
al.[32], this is expected to occur under very low dispersal only. In addition, the observation that the
architecture of the sheltered load differs between dominant and recessive S-alleles suggests that
their diversification dynamics may also differ. Specifically, the self-compatible intermediates
required for the formation of new S-alleles[33,34] are expected to be capable of selfing as well as
forming homozygous genotypes that would otherwise be prevented. While the consequences of
selfing may be equivalent for all alleles (because the overall number of mutations to which they are
linked are equivalent), the consequences of the formation of homozygotes allowed by the crossing
of separate individuals sharing a given S-allele are expected to be more severe for dominant S-
alleles. The segregation of distinct deleterious variants linked to different gene copies of recessive S-
alleles implies that linked recessive deleterious mutations are likely to remain masked when two
distinct gene copies of a given recessive S-allele are brought together. Hence, our results lead to the
prediction that in natural populations self-compatible mutants may segregate more readily for the
more recessive than for the more dominant S-alleles, and more generally for allelic lineages under
lower intensity of balancing selection. Considering that self-compatible mutants are a necessary
intermediate stage in the formation of new S-alleles, one may predict that the diversification
dynamics should be more efficient for lineages of recessive than dominant S-alleles. This prediction
is in line with the deeper phylogenetic divergence among the most dominant S-alleles observed in
Arabidopsis[16]. Detailed quantification of the presence of self-compatible variants in natural
populations will now be necessary to test this hypothesis. At this stage, however, a proper model of
allelic diversification taking into account dominance interactions among S-alleles is still missing.

Our observation that the genetic load varies across balanced allelic lines is not unprecedented. The
classical case of Y or W sex chromosomes are indeed examples where one balanced line accumulates
a greater genetic load than the other (X or Z, respectively), eventually leading to substantial genetic
degeneration[35-37]. Another example is the supergene controlling variation in male plumage
phenotypes of the ruff, where the genetic load on the derived "Satellite" haplotype is higher than on
the ancestral "Independent" haplotype[3,38]. Similarly, in the butterfly *Heliconius numata*, the inverted
haplotypes conferring mimetic wing patterns tend to accumulate a greater load than the non-
inverted haplotypes[39]. Interestingly, in all these cases, the haplotypes with the greatest load also act
genetically in a dominant manner, establishing a clear parallel with our observations. An interesting
next step will be to determine whether similar asymmetries between the balanced allelic lines are
observed in the linked region for these other systems.

It is clear from our results that S-allele dominance can affect the linked load, through its effect on
the intensity of balancing selection, but in turn the differences in structure of the linked load may
affect the conditions under which dominance can evolve. The Brassicaceae S-locus is indeed an
interesting and rather unique system, where the mechanisms by which dominance is controlled and
evolves under the action of so-called "dominance modifiers" have been studied in detail[39,40]
(involving the interaction between small non-coding RNAs and their target sites[16, 40]). The presence
of deleterious mutations linked to S-alleles has been shown to affect the evolution of dominance

modifiers, favouring evolution towards greater dominance than towards greater recessivity[40]. This
asymmetry arises from the fact that *S*-alleles that become recessive (e.g. following acquisition of a
recessivity modifier such as a small RNA target) will start forming homozygous genotypes, leading to
expression of their linked load, while *S*-alleles that become dominant will not. Our observation that
many deleterious mutations linked to recessive *S*-alleles are indeed segregating, suggests that
expression of the load will be less severe for recessive than for dominant *S*-alleles, hence decreasing
this predicted asymmetry. It will now be essential to modify models for the evolution of dominance
to allow for such differential load among *S*-alleles.

Acknowledgements

This work was funded by the European Research Council (NOVEL project, grant #648321) and ANR
TE-MoMa (grant ANR-18-CE02-0020-01). AL's PhD thesis was funded by the ERC and the University
of Lille. The authors thank Barbara Mable for sharing seeds of *A. lyrata* and Camille Roux for
discussions. This work was performed using infrastructure and technical support of the Plateforme
Serre, cultures et terrains expérimentaux - Université de Lille for the greenhouse/field facilities. The
authors thank the UMR 8199 LIGAN-MP Genomics platform (Lille, France) which belongs to the
'Federation de Recherche' 3508 Labex EGID (European Genomics Institute for Diabetes; ANR-10-
LABX-46) and was supported by the ANR Equipex 2010 session (ANR-10-EQPX-07-01; 'LIGAN-MP').
The LIGAN-MP Genomics platform (Lille, France) is also supported by the FEDER and the Region des
Hauts-de-France. The authors thank the GenoScreen platform (Lille, France).

Author contributions

482 AL, ED, VC and XV developed and designed the experiments for the study. AL, CBL and CP performed
483 the experiments. AL and MG analysed and interpreted the data. AL, ED, VC and XV wrote the
484 manuscript. All authors edited the manuscript.

Declaration of interests

The authors declare no competing interests

Figures legend

Figure 1: Effect of homozygosity at the S-locus on 13 phenotypic traits compared to heterozygotes.
For each trait, the phenotypic values in homozygotes (in grey) were normalised relative to the mean
phenotypic values in heterozygotes (in black). The distributions were obtained by 10,000 random
permutations. SD: Standard Deviation.

Figure 2: Linkage to the S-locus locally distorts the phylogenetic relationships. *A. The two*
topologies of interest cluster haplotypes either by the S-allele to which they are linked (top) or by the
populations where they came (bottom). Different S-alleles are represented by symbols of different
colours, different populations of origin are represented by symbols of different shapes. B. Difference
in log likelihood of the two topologies of interest. Dots correspond to the difference in log likelihood
for overlapping series of 50 SNPs around the S-locus for A. halleri (top panel) and A. lyrata (bottom
panel). Positive values correspond to chromosomal positions where the topology by S-alleles explains
the phylogeny of haplotypes better than the topology by populations. The right panels show the
difference in log likelihood in the control regions. 2.5 and 97.5 percentiles of the distribution in the
control regions are indicated by dashed lines.

Figure 3: No overall effect of S-allele dominance on the total number of 0-fold degenerate
mutations (S_{0f}) in the linked genomic regions within 25kb. *Each dot represents the mean number of*
mutations observed among haplotypes linked to one S-allele in one population. The correlations
evaluated by a GLM are represented by lines, with confidence intervals represented in grey. A: A.
halleri. Black dots correspond to the Nivelle population, grey dots to the Mortagne population. The
red arrow points to the copy of Ah15, corresponding to the S-allele whose sheltered load was
phenotypically characterised by Llaurens et al.[13] in Nivelle. B: A. lyrata. Red dots correspond to the
IND population, black dots to the PIN population and blue dots to the TSS population.

Figure 4: The number of 0-fold degenerate mutations fixed in the 25kb regions flanking the S-locus
increases with dominance of the S-allele associated. *Each dot represents the value obtained for*
haplotypes linked to one S-allele in one population. The correlations evaluated by a GLM are
represented by lines, with confidence intervals represented in grey. A: A. halleri. Black dots
correspond to the Nivelle population, grey dots to the Mortagne population. B: A. lyrata. Red dots
correspond to the IND population, black dots to the PIN population, blue dots to the TSS population.

Figure 5: Stochastic simulations confirm the contrasted architecture of the load of deleterious
mutations linked to dominant vs. recessive S-alleles. *Number of fixed (A), segregating (B) and total*
(C) deleterious mutations linked to S-alleles at four different levels of dominance (I<II<III<IV). The
means (bold lines) were estimated per S-allele dominance classes over 100 replicate simulations after
discarding an initial burn-in of 100,000 generations. h=0. s=0.01.

Table 1: Proportion of S-locus homozygous offspring having reached the reproductive stage for
three different S-alleles. The test is performed relative to the expected proportion of homozygous
genotypes in the offspring (25% when both parents are heterozygous; 50% when one of the parents
is homozygous and the other heterozygous).

S-allele	Level of dominance	Number of seedlings having reached the reproductive stage	Observed proportion of homozygotes	Ratio of the observed to expected proportion of homozygous genotypes (p-values[a])	Observed proportion of heterozygotes with the S-allele (p-values compared to expected)
Ah01	I	35	0.29	1.14 (0.70; 0.76)	19 (0.40; 0.90)
Ah03	II	27	**0.074**	**0.3 (0.04; 0.02)**	17 (1; 0.69)
Ah04	III	96	0.479	0.96 (0.75; 0.39)	50

530   [a] *p-values were obtained by binomial test and after 10,000 random permutations, respectively.*
531   *Values departing from Mendelian expectations are figured in bold. For Ah04, the maternal parent*
532   *was Ah04/Ah04, so all heterozygous offspring carried the S-allele.*

Methods STAR

Source plant material

We worked on natural accessions from two closely related species, *A. halleri* and *A. lyrata*, represented by two population samples named Mortagne (50°47'N, 3°47'E, France, *n*=60) and Nivelle (50°47'N, 3°47'E, France, *n*=61) for *A. halleri*, and three highly outcrossing population samples from the North American Great Lakes, named IND (Indiana Dunes National Lakeshore in Michigan, *n*=9), PIN (Pinery Provincial Park in Ontario, *n*=11) and TSS (Tobermory Provincial Park in Ontario, *n*=8)[22] for *A. lyrata* (Fig. S9). The *A. lyrata* populations colonised North America from ancestral European populations about 20-30.000 years ago[29,30] and the *A. halleri* populations are peripheral and likely colonised the north of France during the last century from ancestral German populations[31].

We performed 92, 91, 40, 43 and 21 controlled crosses between randomly chosen individuals within the Nivelle, Mortagne, IND, PIN and TSS populations, respectively. We successfully obtained seeds from 60, 66, 21, 21 and 10 of these crosses, respectively. Because we were not interested in estimating population frequencies of *S*-alleles, we instead tried to maximise the number of reconstructed haplotypes and avoid over representing the most recessive *S*-allele (Ah01) that tends to segregate at very high frequencies in natural populations[15]. To do this, we performed PCR with *S*-allele-specific primers [15,18] to screen the parents of the crosses and we removed from the experiment offspring with two parents carrying allele Ah01. For *A. halleri*, we selected 19 individuals from the Nivelle population and 19 individuals from the Mortagne population, based on their genotype at the *S*-locus (Fig. S9; Table S7). We also selected one offspring of 9, 11, 5, 6 and 5 pairs of selected individuals in the Nivelle, Mortagne, IND, PIN and TSS populations respectively for the phasing of *S*-haplotypes (Table S8; Fig. S9). To increase sample size for the phenotypic measurements, we included offspring from five additional crosses from the Nivelle population (Table S8).

Library preparation, capture and sequencing

We used a previously developed sequence capture approach to specifically sequence genomic regions of interest[20]. Briefly, indexed genomic libraries were constructed for each individual and libraries were pooled in equimolar proportions. Fragments matching a series of regions of interest (including in particular the 75kb upstream and downstream of the non-recombining *S*-locus region as well as a series of 100 unlinked 25kb regions used as genomic controls[20]), were then enriched using synthetic 120bp RNA probes and sequenced by Illumina MiSeq (a total of 159 million paired-end reads).

For six individuals (Table S7, S8), we completed the sequencing with genome-wide resequencing (WGS) in order to distinguish the homozygous and heterozygous genotypes at the *S*-locus based on read depth[41], which is not possible using data from the capture protocol. The prepared libraries were sequenced by Illumina NovaSeq (2x 150pb, paired-end) from the GenoScreen platform (Lille, France).

Determination of the S-locus genotypes and dominance of S-alleles

We used a dedicated pipeline for genotyping the *S*-locus based on short reads sequencing[41] obtained from each individual (Table S7 and S8). The level of dominance of *S*-alleles found in our study was determined based on either previous assessment of dominance in *A. lyrata* and *A. halleri*[15,19,42-44] or indirectly inferred based on the observed association between the phylogeny of *S*-alleles and levels of dominance[45].

Read mapping and variant calling in A. halleri and A. lyrata populations

Raw reads were mapped on the complete *A. lyrata* reference genome V1.0.23[21] using Bowtie2
v2.4.1[46], as described in Le Veve et al[20]. File formats were then converted to BAM using samtools
v1.3.1[47] and duplicated reads were removed with the MarkDuplicates program of picard-tools
v1.119 (http://broadinstitute.github.io/picard). These steps were performed by the custom Python
script   sequencing_genome_vcf.py   available   at   https://github.com/leveveaudrey/analysis-of-
polymorphism-S-locus.

We obtained an average of 620 million properly mapped paired-end 300bp reads per population
sample. For consistency, we conserved only reads which mapped to the *S*-locus flanking or control
regions, even for samples sequenced by WGS, using the *targetintercept* option of bedtool v2.25.0[48].
We called all SNPs within the chromosomal segment comprising 50 kb upstream from the first base
of the gene *Ubox* in 3' and 50 kb downstream from the last base of the gene *ARK3* in 5' of the *S*-locus
using the Genome Analysis Toolkit v. 3.8 (GATK)[49] with the option GVCF and a quality score threshold
of 60 using vcftool v0.1.15[50]. This region contains 20 annotated protein-coding genes. In this study
we excluded the genes inside the *S*-locus itself (*SCR*, *SRK*). For each sample independently, we
computed the distribution of coverage depth across control regions using samtools depth[47]. We
excluded sites with either less than 15 reads aligned or coverage depth above the 97.5 % percentile,
as the latter are likely to correspond to repeated sequences (e.g. transposable elements or
paralogs). Finally, we removed SNPs fixed in each population using the script 1_fix_pos_vcf.py
(https://github.com/leveveaudrey/dominance_and_sheltered_load), thus retaining only nucleotide
sites that were variable in the population.

Quantifying the sheltered load of deleterious mutations

We examined the genetic load signatures based on the accumulation of mutations on 0-fold
degenerate sites, the vast majority of which are considered deleterious. The 0-fold and 4-fold
degenerate sites were identified and extracted from the reference genome and the gene annotation
using the script NewAnnotateRef.py[51]. We also examined the genetic load signatures based on the
accumulation of all non-synonymous mutations. We obtained a total of 2,441 and 2,435 variable
positions for the *A. halleri* samples from Nivelle and Mortagne respectively, and 2,360 variable
positions for the *A. lyrata* samples.

Phasing S-haplotypes

For each of the 9, 11, 5, 6 and 5 trios analysed in the Nivelle, Mortagne, IND, PIN and TSS
populations respectively, we phased mutations in the flanking regions, resulting in 130 phased
haplotypes (Fig. S9). Briefly, we used sites that were heterozygous in the offspring to resolve
parental haplotypes by assuming no recombination between parent and offspring, thus attributing
the allelic state that was shared between a parent and its offspring to their shared *S*-allele, and the
allelic state that was not shared to the other (untransmitted) haplotype of the parent. Twelve of the
parents had been used in more than one cross, and in these cases we phased their haplotypes only
once (Table S8). We implemented the phasing procedure in the script 3_phase_S_allele.py available
at https://github.com/leveveaudrey/dominance_and_sheltered_load.

Study of the structure of S-haplotypes

We first developed a new method to evaluate the distortion of the phylogenetic patterns caused by
linkage to *S*-alleles. To do this, we used phyml[52] v.3.3 to calculate the likelihood of two contrasted
topologies of interest: (1) the topology clustering haplotypes by the populations where they came
from vs. (2) the topology clustering them by the *S*-allele to which they are linked (Fig 2A). We used
sliding windows of sequences with 50 SNPs to obtain the variation of the difference in log-likelihood
between these two topologies along the chromosome. We then compared these values to their

distribution throughout the genome obtained by random draws of sequences with 50 SNPs from the
control regions. Second, we visualised the relationships among the phased haplotypes using
maximum likelihood phylogenies based on the Tamura-Nei model[53], with 1,000 replicates in MEGA
X[54]. Third, we followed Kamau et al.'s[24] approach and examined the variation of F_{ST} among
populations within each species (Nivelle and Mortagne for *A. halleri* and IND, PIN and TSS for *A.*
lyrata) along the flanking region in non-overlapping windows of 5kb. We also examined the variation
of F_{ST} along the flanking region obtained by grouping haplotypes by their linked *S*-allele rather than
by population of origin. Then, we compared these F_{ST} values computed in the *S*-locus flanking regions
with their genomic distribution as determined from the 100 control regions. The F_{ST} values were
estimated with the DNAsp 6 software[55]. Fourth, we performed a major component analysis (MCA)
based on SNPs in the first 5kb, SNPs between 5 and 25kb and SNPs between 25 and 50kb around the
S-locus, using the R packages 'ggplot2' (version 3.4.0), 'factoextra' (version 1.0.7) and 'FactoMiner'
(version 2.7). We compared the patterns obtained by these MCAs with those obtained from identical
numbers of SNP (+/- 1%) from the control regions. Finally, we analysed genetic association in each
population independently between each of the locally segregating variants and the *S*-alleles
considered as phenotypes, using STRAT[56] V1.1 combined with Structure[57] V2.3. We examined the
distribution of the top 0.1% most significant associations detected specifically for each *S*-allele in
each population.

Estimation of the number of fixed and segregating deleterious mutations within S-allele lineages

For each variable position considered in the phased haplotypes, we estimated the number of
mutations on 0-fold (S_{0f}) and 4-fold degenerate sites (S_{4f}) compared with the reference genome. We
distinguished SNPs that were fixed from those that were segregating within each of the allelic lines.
We used GLM with a Poisson distribution to test whether the number of fixed and segregating
mutations were associated with *S*-allele dominance, considering populations as random effects. We
reiterated the GLM analysis with the number of non-synonymous (S_{NS}) and synonymous (S_S)
mutations.

Estimation of the phenotypic impact of homozygosity at the S-locus for three S-alleles

To determine if the genetic sheltered load putatively linked to the *S*-locus has a detectable
phenotypic impact, we performed 45 crosses (Table S1) between offspring of the Nivelle individuals
that we chose so that they shared one *S*-allele (Fig. S9). Based on the dominance hierarchy in
pollen[16] (Table S1), these crosses should correspond to compatible partners. The general principle of
the experiment was to take advantage of the dominance hierarchy to mask recessive *S*-alleles and
generate full sibs that were either homozygous (because they inherited the *S*-allele that was shared
by their two parents) or heterozygous at the *S*-locus, and thus isolate the effect of homozygosity at
the *S*-locus. Note that all offspring in our experiments were thus "naturally" outcrossed, whereas
Llaurens et al.[13] based their comparisons on outcrossed progenies obtained by enforced
incompatible crosses and Stift et al.[19] based their comparisons on enforced selfed progenies. These
crosses generated 399 seeds overall, with homozygous genotypes expected for the *S*-alleles Ah01,
Ah03 and Ah04 forming the following dominance relationship: Ah01<Ah03<Ah04.

Seedlings were grown in a greenhouse between 14.5 and 23.1°C and a photoperiod of 16 hr day/8 hr
night. Offspring from the six families were placed on tables, and their position randomised every
three days. After three months of growing, all the germinated plants were vernalised under a
temperature between 6 and 8°C and a natural photoperiod for two months (January-February).
Then, all surviving plants began reproduction in a greenhouse under temperature between 10.6 and
25.3°C and a natural photoperiod. The genotypes at the *S*-locus were determined in surviving plants
by a PCR approach, using *S*-allele-specific primers for the pistil-expressed *SRK* gene. We assessed the
reproductive success of offspring from the different crosses on the basis of fourteen phenotypic
traits (detailed below) and computed the mean difference for the trait between homozygotes and

heterozygotes within each family. We also tested for departures from mendelian proportions of
each S-locus genotypic category in the family after the apparition of the first stem. Significant
departures were interpreted as reflecting differences in survival between homozygous and
heterozygous S-locus genotypes. We performed 10,000 replicate simulations of mendelian
segregation based on the S-locus genotype of the parents. We used GLM to test whether the
phenotypic impact of homozygosity at the S-locus increased with dominance of the S-alleles. The
models used for GLM depended on the type of trait analysed (poisson for the counts like the number
of leaves, flowers by stems or days; gaussian for continue traits like the lengths, widths and areas).

We measured the following fourteen phenotypic traits: the time (days) to the first leaf measured by
visual control every day during seven weeks after sowing the seeds, the number of leaves, the area
of the rosette (cm²), the mean length and width of leaves (cm), the standard deviation of length and
width of leaves (cm) and the mean area of leaves (cm²) measured by ImageJ[58] based on photographs
taken seven weeks (+/- five days) after the first leaf. At reproduction, we measured the time to the
first flower bud for the end of vernalisation (day), scored by visual control every three days during
nine weeks, the number of flower buds per flower stem produced during four week after the
appearance of the first bud, the number of flower stems, the length of the highest flower stem
produced four weeks after the appearance of the first bud (cm), and finally the total duration of
buds production (days), scored by visual control every three days during eleven weeks after the
appearance of the first bud. The last trait we measured was the proportion of homozygotes per
family that survived until reproduction assuming mendelian proportions in the seeds. During the
whole experiment, the presence of phytophagous insects, pathogens and stress markers were
scored as binary variables. The presence of phytophagous insects and pathogen attacks were
detected by the occurrence of gaps in leaves. Oxidative stress was scored qualitatively based on the
occurrence of purple leaves. We also controlled the effect of the family on the phenotypic trait.
These effects were controlled by redistributing 10,000 times the values observed in groups of the
same size observed for each effect (for example, presence or absence of pathogen attack) and
comparing the difference for the trait observed with the distribution of the differences obtained in
the permutations. We considered the impact of the effect on the trait if the observed difference
between groups was higher than the 95% percentile of the distribution obtained randomly (Table
S9). When the test was significant, the effect was implemented as a random effect in the GLM. We
used the same method to control for the family effect, which was included as a random effect in
GLM if necessary (Table S10). The general experimental procedure is summarised in Fig. S9 and all
data analyses were done in R ver. 3.1.2 (R Development Core Team 2014).

Simulations

Finally, we refined the model of Llaurens et al.,[13] in several ways. We simulated a panmictic
population of N diploid individuals with non-overlapping generations. Each individual was defined by
its genotype in a non-recombining genomic region. This region contains the S- locus, and a D locus
where deleterious mutations accumulated. For the S-locus, we used a simple model of sporophytic
SI, with four dominance classes, as observed in A. halleri[41] (only three classes were considered
before[13]), and fourteen S-alleles (eight alleles in the class IV, three in the class III, two in the class II
and one allele in the class I). This distribution mirrors that of the Nivelle population (Table S7), with
the exception that a class II allele has been added because its presence has been reported in
previous studies[15]. Alleles within classes were assumed to be codominant with each other, and
dominant over all alleles of the more recessive classes, with the following linear hierarchy between
classes : classI<classII<classIII<classIV). We also assumed that no new S-allele could appear by
mutation during the simulations. The population size was 10,000 diploid individuals, so as to be large
enough to avoid S-allele loss by drift during the simulations (previously it was 1,000). The "D locus"
comprised one hundred fully linked biallelic positions (versus a single one in Llaurens et al.[13]). Fully
recessive deleterious mutations were recurrently introduced (at a rate 10^{-4}), and reverse mutations
were possible (at a rate 10^{-5}). We ignored partially recessive deleterious mutations because these

mutations were predicted to be effectively eliminated by natural selection in Llaurens et al.[13]. The
survival probability p of a zygote depended on its genotype at the D locus: $p = (1 - s)^n$ with s the
selection coefficient and n the number of positions homozygous for the mutated allele. We explored
different values of the selection coefficient (0.1, 0.05, 0.03, 0.01 and 0.005). Under strong selection
(s=0.1, 0.05 and 0.03), the combined effect of multiple mutations led to low-fitness individuals,
eventually causing population extinction. Under weak selection, (s=0.005), we observed near fixation
of the deleterious mutations under the influence of asymmetrical mutation. Hence, we focused on
the intermediate value of the selection coefficient (s=0.01), where deleterious mutations segregated
stably in the simulations.

We first ran simulations without deleterious mutations until a deterministic equilibrium for S-allele
frequencies was reached, which was considered to be attained when allelic frequencies changed by
less than 10^{-3} between generations. Recessive deleterious mutations were then allowed to
accumulate at the positions within the D locus. Each simulation was performed with 100
independent replicates of 100,000 generations, and the frequency of the deleterious alleles was
recorded every 1,000 generations. At the end of the simulation runs, we estimated the number of
deleterious mutations found in each haplotype associated with each S-allele to determine the
expected patterns of association between the sheltered load and dominance at the S-locus.

Data Availability

755 All sequence data are available in the NCBI Short Read Archive (SRA;
756 https://www.ncbi.nlm.nih.gov/sra) with accession codes: PRJNA744343, PRJNA755829.

757 All scripts developed are available in Github
758 (https://github.com/leveveaudrey/dominance_and_sheltered_load
759 https://github.com/leveveaudrey/analysis-of-polymorphism-S-locus).

Supplementary data

Figures legend

Figure S1: Analysis of Major Components obtained for haplotypes of A. halleri of the Nivelle (black dots) and Mortagne (grey dots) populations) based on SNPs in the first 5kb, between 5 and 25kb, and between 25kb and 50kb away from the S-locus. *The S-allele to which the SNPs are linked are represented by different symbols. The right panels show the analysis on control regions, each time matching the number of SNPs with that of the corresponding linked regions (left panels).*

Figure S2: Analysis of major components (AMC) obtained for haplotypes of A. lyrata (of the PIN: grey dots, IND: red dots and TSS: blue dots) populations based on the SNPs in the first 5kb, between 5 and 25kb, and between 25kb and 50kb away from the S-locus. *The S-allele to which the SNPs are linked are represented by different symbols. The right panels show the analysis on control regions, each time matching the number of SNPs with that of the corresponding linked regions (left panels).*

Figure S3: Phylogenetic tree obtained by Maximum Likelihood for haplotypes of A. halleri (populations Nivelle and Mortagne) across the first 25kb flanking the S-locus. *The Tamura-Nei model was used and the percentage of trees in which the associated haplotypes clustered together is shown next to the branches. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. The black braces indicate haplotypes clustering by populations. The overall pattern shows a structure by S-alleles, with exceptions highlighted in yellow.*

Figure S4: Phylogenetic tree obtained by maximum likelihood for haplotypes of A. halleri (populations Nivelle and Mortagne) based on the nucleotide positions between 25kb and 50kb away from the S-locus. *The Tamura-Nei model was used and the percentage of trees in which the associated haplotypes clustered together is shown next to the branches. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. The black braces indicate haplotypes clustering by populations. The overall pattern shows a structure by S-alleles, with exceptions highlighted in yellow.*

Figure S5: Phylogenetic tree obtained by maximum likelihood for haplotypes of A. lyrata (populations PIN, IND, TSS) across the first 5kb flanking the S-locus. *The Tamura-Nei model was used and the percentage of trees in which the associated haplotypes clustered together is shown next to the branches. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. The black braces indicate haplotypes clustering by populations. The overall pattern shows a structure by S-alleles, with exceptions highlighted in yellow.*

Figure S6: Phylogenetic tree obtained by maximum likelihood for haplotypes of A. lyrata (populations PIN, IND, TSS) based on the nucleotide positions between 5kb and 10kb away from the S-locus. *The Tamura-Nei model was used and the percentage of trees in which the associated haplotypes clustered together is shown next to the branches. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. The black braces indicate haplotypes clustering by populations. The overall pattern shows a structure by S-alleles, with exceptions highlighted in yellow.*

Figure S7: The genetic structure of SNPs in the S-locus flanking regions in A. halleri and A. lyrata.
Left: Mean F_{ST} (lines) and F_{ST} by pair (point) analysed among S-alleles (grey) or among populations (black) in 5 kb windows in the S-locus flanking regions. Right: Distribution (count) of F_{ST} in the control regions analysed among S-alleles (grey bars) or among populations (black bars). The 95% percentiles of the distributions are represented by dotted lines and the medians by solid lines.

Figure S8: Patterns of genetic associations between S-alleles and SNPs across the genome. Each dot corresponds to a SNP showing statistically significant association (top 0.1%) with a given S-allele. The left panel confirms that SNPs physically linked to the S-locus (in red) are considerably more likely to show statistical associations with S-alleles. The chromosomes were signified by alternance of black and grey. The middle panel shows a zoom on the 50kb regions flanking the S-locus and shows that the statistical association extends over long distances, at least for some S-alleles. Each region of 25 kb was delimited by vertical lines. The right panel shows that the observed number of associated SNPs in the linked regions far exceeds that in regions of identical size from control regions. The histogram shows the distribution across control regions of the mean number of significantly associated SNPs per S-allele. The vertical lines correspond to the mean number of significantly associated SNPs in the first 25kb (solid lines) and the 25-50kb interval (dashed lines) away from S-locus.

Figure S9: Experimental protocol. A) We randomly crossed A. lyrata individuals from the PIN, TSS and IND populations in North America (left) and A. halleri from the Nivelle (middle) and Mortagne (right) populations. Individuals were sequenced by a capture protocol. Numbers between parentheses represent the number of individuals per dataset. B) One offspring from each cross was sequenced along with its two parents for trio haplotyping. Offspring from the Nivelle population (black circle) were conserved for the study of the impact of homozygosity at the S-locus on fitness (G1 population). C)Individuals sequenced in A and B were used to reconstruct haplotypes linked to each copy of S-allele, assuming no recombination between the S-locus and its flanking regions between parents and offspring. D) We used the dominance hierarchy between S-alleles expressed in pollen[22,23] to cross G1 individuals from theNivelle population and obtained six G2 families constituted of heterozygous and homozygous individuals for the alleles Ah01 , Ah03 and Ah04 . E) Description of the traits measured and the methods used to estimate the impact of homozygosity at the S-locus in homozygotes. Traits 1-8 are related to biomass and traits 10-14 are related to reproductive success.

Figure S1: Analysis of Major Components obtained for haplotypes of A. halleri of the Nivelle (black
dots) and Mortagne (grey dots) populations) based on SNPs in the first 5kb, between 5 and 25kb,
and between 25kb and 50kb away from the S-locus.

Figure S2: Analysis of major components (AMC) obtained for haplotypes of A. lyrata (of the PIN:
grey dots, IND: red dots and TSS: blue dots) populations based on the SNPs in the first 5kb,
between 5 and 25kb, and between 25kb and 50kb away from the S-locus.

Figure S3: Phylogenetic tree obtained by Maximum Likelihood for haplotypes of A. halleri

(populations Nivelle and Mortagne) across the first 25kb flanking the S-locus.

Figure S4: Phylogenetic tree obtained by maximum likelihood for haplotypes of A. halleri
(populations Nivelle and Mortagne) based on the nucleotide positions between 25kb and 50kb
away from the S-locus.

848 **Figure S5: Phylogenetic tree obtained by maximum likelihood for haplotypes of A. lyrata**
849 **(populations PIN, IND, TSS) across the first 5kb flanking the S-locus.**

Figure S6: Phylogenetic tree obtained by maximum likelihood for haplotypes of A. lyrata (populations PIN, IND, TSS) based on the nucleotide positions between 5kb and 10kb away from the S-locus.

856   *Figure S7: The genetic structure of SNPs in the S-locus flanking regions in A. halleri and A. lyrata.*

858   *Figure S8: Patterns of genetic associations between S-alleles and SNPs across the genome.*

860   *Figure S9: Experimental protocol.*

Supplemental items

Table S1: Crosses performed to obtain homozygotes for three S-alleles.

Table S2: Trait variation in S-locus homozygous individuals.

Table S3: Trait variation in homozygous at the S-locus for the S-alleles Ah01 and Ah04 between families.

Table S4: Effect of dominance on variation of phenotypic traits in S-locus homozygous individuals.

Table S5: Number of phased haplotypes linked to S-alleles in each sample.

Table S6: Effect of dominance on the accumulation of genetic load in S-flanking regions.

Table S7: S-locus genotypes of individuals sequenced using the capture protocol.

Table S8: S-locus genotypes of the offspring selected for haplotype phasing and the crosses for the study of phenotypic traits.

Table S9: Effect of phytopathogen, phytophagous attacks and oxidative stress on the phenotypic traits.

Table S10: Difference on the phenotypic traits variation between the two families for each allele tested.

885 **Table S1: Crosses performed to obtain homozygotes for three S-alleles.** *For each S-allele, we*
886 *obtained homozygous and heterozygous plants within two separate families (listed as separate*
887 *rows).*

Genotype pollen donor [a]	Genotype stigma recipient	Homozygote studied	Dominance level	Number of fruits obtained	Number of seeds
Ah20/Ah01	Ah12/Ah01	Ah01	1	5	48
Ah20/Ah01	Ah12/Ah01	Ah01	1	2	23
Ah02/Ah03	Ah03/Ah01	Ah03	2	6	69
Ah24/Ah03	Ah03/Ah01	Ah03	2	3	27
Ah20/Ah04	Ah04/Ah04	Ah04	3	21	167
Ah20/Ah04	Ah04/Ah04	Ah04	3	8	65

888 [a] *The S-alleles in bold represent the dominant S-allele expressed on pollen of each donor genotype.*
889 *Each line corresponds to one pair of individuals used as parents.*

893 **Table S2: Trait variation in S-locus homozygous individuals.**

Trait (unit)	Mean heterozygotes (SD)	Mean homozygotes (SD)	P value[a]
Time first leaf (day)	15.7 (1.04)	14.79 (1.19)	0.14
Number of leaves	13.22 (0.65)	13.46 (0.96)	0.35
Rosette area (cm²)	34.99 (5.11)	35.5 (7.65)	0.46
Mean leaf length (cm)	1.86 (0.15)	1.86 (0.21)	0.49
St dev leaf length (cm)	0.28 (0.02)	0.29 (0.04)	0.31
Mean leaf width (cm)	1.49 (0.11)	1.46 (0.14)	0.36
St dev leaf width (cm)	0.27 (0.02)	0.25 (0.03)	0.24
Mean leaf area (cm²)	3.16 (0.43)	3.11 (0.6)	0.45
Time first flower (day)	36.28 (1.14)	34.98 (1.25)	0.07
Number of flowering stems	12.9 (0.91)	13.3 (1.32)	0.31
Maximum flowering stem length (cm)	**56.19 (2.26)**	**64.55 (3.17)**	**$<1e^{-04}$**
Number of flowers by flowering stem	58.87 (5.96)	66.56 (10.71)	0.09
Flowering duration (day)	44.15 (1.7)	43.81 (1.88)	0.39

894 [a] The P value represents the proportions of the distribution with the expected difference of mean
895 between heterozygotes and homozygotes at the S-locus after 10,000 random resamples equal to or
896 less than the value observed. Significant values are represented in bold.

902 **Table S3: Trait variation in homozygous at the S-locus for the S-alleles Ah01 and Ah04 between**
903 **families.**

Allele	Trait (unit)	Mean first familly (SD)	Mean second familly (SD)	p value
Ah01[b]	**Time first leaf (day)**	**17.11 (2.26)**	**32 (NA)**	**<1.00^{e-04}**
	Number of leaves	14 (2.83)	9 (NA)	0.10
	Rosette area (cm²)	27.65 (18.22)	14.21 (NA)	0.50
	Mean leaf length (cm)	1.64 (0.68)	1.52 (NA)	0.49
	St dev leaf length (cm)	0.28 (0.11)	0.17 (NA)	0.30
	Mean leaf width (cm)	1.38 (0.43)	1.38 (NA)	0.49
	St dev leaf width (cm)	0.24 (0.07)	0.16 (NA)	0.30
	Mean leaf area (cm²)	2.52 (1.59)	2.1 (NA)	0.50
	Time first flower (day)	35.86 (3.13)	36 (NA)	0.25
	Number of flowering stems	12.29 (3.9)	7 (NA)	0.25
	Maximum flowering stem length (cm)	**43.43 (6.91)**	**62.4 (NA)**	**<1.00^{e-04}**
	Number of flowers by flowering stem	31.73 (24.16)	52.25 (NA)	0.12
	Flowering duration (day)	**44.71 (6.73)**	**52 (NA)**	**<1.00^{e-04}**
Ah04	Time first leaf (day)	13.15 (2.68)	14.88 (4.36)	0.11
	Number of leaves	13.43 (2.99)	14.04 (3.9)	0.24
	Rosette area (cm²)	36.63 (27.74)	45.86 (33.48)	0.08
	Mean leaf length (cm)	1.93 (0.81)	2.13 (0.89)	0.14
	St dev leaf length (cm)	0.29 (0.15)	0.3 (0.12)	0.33
	Mean leaf width (cm)	1.54 (0.57)	1.44 (0.51)	0.20
	St dev leaf width (cm)	**0.28 (0.13)**	**0.22 (0.09)**	**0.02**
	Mean leaf area (cm²)	3.4 (2.45)	3.46 (2.27)	0.45
	Time first flower (day)	35.57 (4.73)	33.2 (3.66)	0.06
	Number of flowering stems	**13.21 (4.25)**	**16.4 (5.25)**	**0.02**
	Maximum flowering stem length (cm)	64.98 (9.65)	62.41 (9.34)	0.13
	Number of flowers by flowering stem	**72.38 (33.07)**	**45.25 (15.45)**	**1.00^{e-04}**
	Flowering duration (day)	43.68 (6.96)	41.56 (7.38)	0.23

[a] The P value represents the proportions of the distribution with the expected difference of mean
between each family after 10,000 random resamples equal to or less than the observed value.
Significant values are represented in bold. The Ah03 S-allele is excluded because only two
homozygotes survived. [b] the standard deviation for the second family was not estimated because
only one homozygote survived.

Table S4: Effect of dominance on variation of phenotypic traits in S-locus homozygous individuals.

Trait (unit)	Random effect implemented in GLM model	Distribution of trait tested	Linear effect	p value
Time first leaf (days)	**Family**	**Poisson**	**-0.15**	**5.5^{e-04}**
Leaves (counts)	Oxydative stress	Poisson	0.006	0.91
Rosette area (cm²)	Oxydative stress + family	Gaussian	8.25	0.19
Mean leave length (cm)	Oxydative stress + family	Gaussian	0.17	0.28
St dev leave length (cm)	None	Gaussian	0.02	0.54
Mean leave width (cm)	Oxydative stress + phytophagous attack	Gaussian	0.05	0.83
St dev leave width (cm²)	Oxydative stress + phytophagous attack	Gaussian	0.01	0.91
Mean leave area (cm²)	Oxydative stress	Gaussian	0.39	0.36
Time first flower (days)	Phytopathogen + phytophagous attack + family	Poisson	-0.0002	1
Flowering stems (counts)	Family	Poisson	0.15	0.08
Maximum flowering stem length (cm)	Phytopathogen + family	Gaussian	7.6	0.06
Flowers by stem (counts)	Family	Poisson	0.15	0.44
Flowering duration (days)	Phytopathogen + family	Poisson	-0.05	0.1

910   *Dominance at the S-locus was implemented as a fixed effect.*

914 **Table S5: Number of phased haplotypes linked to S-alleles in each sample.**

Species	Population	Number of phased haplotypes	Number of S-alleles	Number of S-alleles with more than one copy	Mean number of gene copies per S-allele [a]
A. halleri	Nivelle	34	12	7	4.1
	Mortagne	38	11	9	4
A. lyrata	TSS	16	5	3	4.7
	IND	16	6	4	3.5
	PIN	22	7	3	6

915 [a] *based on S-alleles with more than one copy.*

917 *Table S6: Effect of dominance on the accumulation of genetic load in S-flanking regions.*

Type of mutations	Species	Genetic load estimated	Linear effect	p value
0fold	A. halleri	**Number of fixed mutations by S-allele**	**0.08**	**2.57^{e-04}**
		Number of segregated mutations by S-allele	**-0.49**	**2.21^{e-03}**
		Mean total number of mutation by S-allele (S_{0f})	0.01	0.54
	A. lyrata	**Number of fixed mutations by S-allele**	**0.08**	**4.50^{e-03}**
		Number of segregated mutations by S-allele	-0.44	0.09
		Mean total number of mutation by S-allele (S_{0f})	-0.07	0.07
4fold	A. halleri	**Number of fixed mutations by S-allele**	**0.23**	**8.95^{e-04}**
		Number of segregated mutations by S-allele	**-0.48**	**2.02^{e-10}**
		Mean total number of mutation by S-allele (S_{4f})	1.53^{e-03}	0.95
	A. lyrata	**Number of fixed mutations by S-allele**	**0.23**	**8.95^{e-04}**
		Number of segregated mutations by S-allele	**-0.46**	**0.04**
		Mean total number of mutation by S-allele (S_{4f})	-0.04	0.32
S_{0f}/S_{4f}	A. halleri	Ratio	0.02	0.42
	A. lyrata	Ratio	-0.11	0.11
Non-synonymous	A. halleri	**Number of fixed mutations by S-allele**	**0.67**	**1.86^{e-04}**
		Number of segregated mutations by S-allele	**-0.23**	**$<2^{e-16}$**
		Mean total number of mutation by S-allele (S_{NS})	0.02	0.12
	A. lyrata	**Number of fixed mutations by S-allele**	**0.24**	**1.74^{e-08}**
		Number of segregated mutations by S-allele	**-0.28**	**$<2^{e-16}$**
		Mean total number of mutation by S-allele (S_{NS})	7.99^{e-03}	0.65
Synonymous	A. halleri	**Number of fixed mutations by S-allele**	**0.08**	**4.55^{e-04}**
		Number of segregated mutations by S-allele	**-0.26**	**$<2^{e-16}$**
		Mean total number of mutation by S-allele (S_S)	0.02	0.15
	A. lyrata	**Number of fixed mutations by S-allele**	**0.27**	**2.3^{e-11}**
		Number of segregated mutations by S-allele	**-0.34**	**$<2^{e-16}$**
		Mean total number of mutation by S-allele (S_S)	-0.01	0.33
S_{NS}/S_S	A. halleri	Ratio	-5.82^{e-03}	0.81
	A. lyrata	Ratio	0.02	0.60

919   *Dominance at the S-locus was implemented as a fixed effect. Populations were implemented as random effects. The distribution of the genetic load*
920   *estimated followed a Poisson distribution, except the ratios that followed normal distributions.*

Table S7: S-locus genotypes of individuals sequenced using the capture protocol.

Individual	Allele 1[a]	Allele 2[a]	Population	Species	SRA
Mor_19_13	Ah04 (III)	Ah25 (III)	Mortagne	A. halleri	SAMN20088356
Mor_19_14	Ah04 (III)	Ah01 (I)	Mortagne	A. halleri	SAMN20844087
Mor_19_19	Ah12 (IV)	Ah36 (IV)	Mortagne	A. halleri	SAMN20088358
Mor_19_2	Ah03 (II)	Ah24 (IV)	Mortagne	A. halleri	SAMN20088359
Mor_19_23	Ah20 (IV)	Ah25 (III)	Mortagne	A. halleri	SAMN20844088
Mor_19_24	Ah12 (IV)	Ah24 (IV)	Mortagne	A. halleri	SAMN20088362
Mor_19_3	Ah03 (II)	Ah01 (I)	Mortagne	A. halleri	SAMN20844098
Mor_19_37	Ah59 (IV)	Ah20 (IV)	Mortagne	A. halleri	SAMN20088357
Mor_19_38	Ah36 (IV)	Ah01 (I)	Mortagne	A. halleri	SAMN20088371
Mor_19_4	Ah25 (III)	Ah12 (IV)	Mortagne	A. halleri	SAMN20088360
Mor_19_41	Ah25 (III)	Ah25 (III)	Mortagne	A. halleri	SAMN20088372
Mor_19_42	Ah12 (IV)	Ah04 (III)	Mortagne	A. halleri	SAMN20088373
Mor_19_45	Ah36 (IV)	Ah02 (III)	Mortagne	A. halleri	SAMN20088374
Mor_19_46	Ah03 (II)	Ah12 (IV)	Mortagne	A. halleri	SAMN20088375
Mor_19_51	Ah20 (IV)	Ah05 (IV)	Mortagne	A. halleri	SAMN20088361
Mor_19_53	Ah36 (IV)	Ah12 (IV)	Mortagne	A. halleri	SAMN20088365
Mor_19_54	Ah24 (IV)	Ah12 (IV)	Mortagne	A. halleri	SAMN20088366
Mor_19_55	Ah20 (IV)	Ah05 (IV)	Mortagne	A. halleri	SAMN20088377
Mor_19_56	Ah12 (IV)	Ah03 (II)	Mortagne	A. halleri	SAMN20088378
Niv_19_18	Ah15 (IV)	Ah03 (II)	Nivelle	A. halleri	SAMN20088349
Niv_19_19	Ah20 (IV)	Ah24 (IV)	Nivelle	A. halleri	SAMN20088350
Niv_19_22	Ah24 (IV)	Ah01 (I)	Nivelle	A. halleri	SAMN20088351
Niv_19_23	Ah20 (IV)	Ah12 (IV)	Nivelle	A. halleri	SAMN20088352
Niv_19_3	Ah04 (III)	Ah59 (IV)	Nivelle	A. halleri	SAMN20088336
Niv_19_31	Ah04 (III)	Ah24 (IV)	Nivelle	A. halleri	SAMN20088337

Niv_19_4	Ah01 (I)	Ah24 (IV)	Nivelle	A. halleri	SAMN20088382
Niv_19_42	Ah22 (IV)	Ah12 (IV)	Nivelle	A. halleri	SAMN20088341
Niv_19_45	Ah12 (IV)	Ah04 (III)	Nivelle	A. halleri	SAMN20088344
Niv_19_5	Ah25 (III)	Ah67 (IV)	Nivelle	A. halleri	SAMN20088340
Niv_19_52	Ah20 (IV)	Ah22 (IV)	Nivelle	A. halleri	SAMN20088331
Niv_19_53	Ah05 (IV)	Ah20 (IV)	Nivelle	A. halleri	SAMN20088339
Niv_19_54	Ah01 (I)	Ah12 (IV)	Nivelle	A. halleri	SAMN20844097
Niv_19_58	Ah02 (III)	Ah59 (IV)	Nivelle	A. halleri	SAMN20088343
Niv_19_59	Ah01 (I)	Ah67 (IV)	Nivelle	A. halleri	SAMN20844096
Niv_19_60	Ah04 (III)	Ah20 (IV)	Nivelle	A. halleri	SAMN20088332
Niv_19_7	Ah04 (III)	Ah67 (IV)	Nivelle	A. halleri	SAMN20088345
Niv_19_8	Ah04 (III)	Ah12 (IV)	Nivelle	A. halleri	SAMN20844090
Niv_19_9	Ah04 (III)	Ah67 (IV)	Nivelle	A. halleri	SAMN20088338
Pin_15_1	Ah03 (II)	Ah01 (I)	PIN	A. lyrata	SAMN20088324
Pin_16_1	Ah01 (I)	Ah46 (IV)	PIN	A. lyrata	SAMN20088318
Pin_4_24	Ah03 (II)	Ah29 (III)	PIN	A. lyrata	SAMN20088314
Pin_16_3	Ah18 (IV)	Ah29 (III)	PIN	A. lyrata	SAMN20088323
Pin_4_54	Ah03 (II)	Ah03 (II)	PIN	A. lyrata	SAMN20088319
Pin_5_1	Ah01 (I)	Ah01 (I)	PIN	A. lyrata	SAMN20844102
Pin_5_12	Ah29 (III)	Ah01 (I)	PIN	A. lyrata	SAMN20088325
Pin_5_2	Ah03 (II)	Ah01 (I)	PIN	A. lyrata	SAMN20088311
Pin_8_15	Ah01 (I)	Ah01 (I)	PIN	A. lyrata	SAMN20088328
Pin_8_2	Ah29 (III)	Ah63 (III)	PIN	A. lyrata	SAMN20088326
Pin_9_1	Ah42 (IV)	Ah29 (III)	PIN	A. lyrata	SAMN20088312
Tss_14_3	Ah01 (I)	Ah31 (IV)	TSS	A. lyrata	SAMN20088315
Tss_21_10	Ah01 (I)	Ah03 (II)	TSS	A. lyrata	SAMN20088329
Tss_22_24	Ah01 (I)	Ah31 (IV)	TSS	A. lyrata	SAMN20088306

Tss_22_7	*Ah01 (I)*	*Ah31 (IV)*	*TSS*	*A. lyrata*	*SAMN20088317*
Tss_3_10	*Ah31 (IV)*	*Ah01 (I)*	*TSS*	*A. lyrata*	*SAMN20088330*
Tss_23_2	*Ah03 (II)*	*Ah63 (III)*	*TSS*	*A. lyrata*	*SAMN20088313*
Tss_3_23	*Ah18 (IV)*	*Ah01 (I)*	*TSS*	*A. lyrata*	*SAMN20088307*
Tss_5_1	*Ah01 (I)*	*Ah31 (IV)*	*TSS*	*A. lyrata*	*SAMN20088316*
Ind_1_1	*Ah01 (I)*	*Ah01 (I)*	*IND*	*A. lyrata*	*SAMN20088320*
Ind_10_3	*Ah31 (IV)*	*Ah24 (IV)*	*IND*	*A. lyrata*	*SAMN20088327*
Ind_15_1	*Ah03 (II)*	*Ah24 (IV)*	*IND*	*A. lyrata*	*SAMN20088321*
Ind_15_2	*Ah63 (III)*	*Ah03 (II)*	*IND*	*A. lyrata*	*SAMN20088309*
Ind_9_3	*Ah01 (I)*	*Ah46 (IV)*	*IND*	*A. lyrata*	*SAMN20088322*
Ind_18_1	*Ah01 (I)*	*Ah01 (I)*	*IND*	*A. lyrata*	*SAMN20844103*
Ind_6_1	*Ah01 (I)*	*Ah63 (III)*	*IND*	*A. lyrata*	*SAMN20088308*
Ind_8_1	*Ah03 (II)*	*Ah63 (III)*	*IND*	*A. lyrata*	*SAMN20088305*

922 *[a] The individuals were mainly sequenced by capture approach. The genotypes of homozygotes were*
923 *confirmed after a whole genome sequencing. The class of dominance is specified in parenthesis.*

928 **Table S8: S-locus genotypes of the offspring selected for haplotype phasing and the crosses for the**
929 **study of phenotypic traits.**

Identity[a]	Allele 1	Allele 2	Pollen donor[f]	Stigma recipient[c]	SRA
d32	Ah03	Ah12	Mor_19_2	Mor_19_19	SAMN20844104
d33	Ah01	Ah12	Mor_19_3	Mor_19_4	SAMN20844105
d38[d]	Ah04	Ah04	Mor_19_13	Mor_19_14	SAMN20844106
d42	Ah20	Ah12	Mor_19_23	Mor_19_24	SAMN20844107
d72	Ah20	Ah36	Mor_19_37	Mor_19_38	SAMN20844108
d74	Ah25	Ah12	Mor_19_41	Mor_19_42	SAMN20844109
d76	Ah03	Ah36	Mor_19_46	Mor_19_45	SAMN20844110
d89	Ah03	Ah05	Mor_19_56	Mor_19_55	SAMN20844111
d217	Ah12	Ah59	Mor_19_19	Mor_19_37	SAMN20844112
d250	Ah12	Ah20	Mor_19_24	Mor_19_51	SAMN20844113
d265	Ah36	Ah24	Mor_19_53	Mor_19_54	SAMN20844114
d3.1	Ah24	Ah25	Niv_19_4	Niv_19_5	SAMN20844115
d10.1[f]	Ah03	Ah24	Niv_19_18	Niv_19_19	SAMN20844116
d12.1[f]	Ah01	Ah20	Niv_19_22	Niv_19_23	SAMN20844117
d30.1	Ah02	Ah01	Niv_19_58	Niv_19_59	SAMN20844118
d48.1	Ah20	Ah25	Niv_19_52	Niv_19_5	SAMN20844119
d50.1[e]	Ah12	Ah04	Niv_19_8	Niv_19_60	SAMN20844120
d51.1	Ah12	Ah04	Niv_19_54	Niv_19_7	SAMN20844121
d191.1[e,f]	Ah04	Ah04	Niv_19_3	Niv_19_31	SAMN20844122
d206.1	Ah22	Ah25	Niv_19_42	Niv_19_5	SAMN20844123
d205.1[f]	Ah20	Ah04	Niv_19_53	Niv_19_9	SAMN20844124
d208.1[f]	**Ah03**	**Ah02**	**Niv_19_58**	**Niv_19_18**	**SAMN20844125**
d17.1[f]	**Ah03**	**Ah01**	**Niv_19_33**	**Niv_19_32[b]**	**SAMN20844126**
d24.1[f]	**Ah20**	**Ah01**	**Niv_19_47[b]**	**Niv_19_46[b]**	**SAMN20844127**

d29.1[f]	*Ah12*	*Ah01*	*Niv_19_57[b]*	*Niv_19_56[b]*	*SAMN20844128*
d50.2[f]	*Ah04*	*Ah04*	*Niv_19_8*	*Niv_19_60*	*SAMN20844129*
d266.1	*Ah04*	*Ah12*	*Niv_19_7*	*Niv_19_45*	*SAMN20844130*
d122	*Ah01*	*Ah63*	*Pin_16_1*	*Pin_8_2*	*SAMN20844131*
d127	*Ah01*	*Ah03*	*Pin_5_1*	*Pin_4_54*	*SAMN20844132*
d173	*Ah03*	*Ah18*	*Pin_5_2*	*Pin_16_3*	*SAMN20844133*
d176	*Ah03*	*Ah42*	*Pin_15_1*	*Pin_9_1*	*SAMN20844134*
d177	*Ah01*	*Ah29*	*Pin_16_1*	*Pin_5_12*	*SAMN20844135*
d239	*Ah29*	*Ah01*	*Pin_4_24*	*Pin_8_15*	*SAMN20844136*
d113	*Ah18*	*Ah31*	*Tss_3_23*	*Tss_22_7*	*SAMN20844137*
d273	*Ah03*	*Ah01*	*Tss_23_2*	*Tss_22_24*	*SAMN20844138*
d275	*Ah63*	*Ah31*	*Tss_23_2*	*Tss_14_3*	*SAMN20844139*
d284	*Ah01*	*Ah31*	*Tss_21_10*	*Tss_5_1*	*SAMN20844140*
d285	*Ah01*	*Ah31*	*Tss_21_10*	*Tss_3_10*	*SAMN20844141*
d118	*Ah03*	*Ah01*	*Ind_9_3*	*Ind_8_1*	*SAMN20844142*
d163	*Ah03*	*Ah01*	*Ind_8_1*	*Ind_1_1*	*SAMN20844143*
d166	*Ah03*	*Ah01*	*Ind_6_1*	*Ind_15_1*	*SAMN20844144*
d170	*Ah01*	*Ah63*	*Ind_9_3*	*Ind_15_2*	*SAMN20844145*
d232	*Ah01*	*Ah31*	*Ind_18_1*	*Ind_10_3*	*SAMN20844146*

[a] *Offspring figured in bold were not used to reconstitute haplotype of parents.* [b] *Parents were not*
sequenced. [c] *Italics indicate parental haplotypes that were suppressed from the dataset because they*
had already been phased through another offspring . [d] *The homozygous genotype of d38 was*
confirmed by genome-wide sequencing. [e] *The homozygous genotypes of d191 and d50.1 were*
confirmed by the proportion of homozygous S-alleles in their offspring found by PCR after cross. [f] *The*
individuals used for crosses in the phenotypic experiments.

936
937
938

Table S9: Effect of phytopathogen, phytophagous attacks and oxidative stress on the phenotypic

traits.

Effect tested	Trait	Mean with attack (sd)	Mean without attack (sd)	p value [a]
	Time first leaf (day)	15.37 (4.84)	15.38 (5.24)	0.49
	Number of leaves	13.03 (2.9)	13.46 (3.71)	0.23
	Rosette area (cm²)	31.93 (27.19)	37.08 (27.14)	0.13
	Mean leave length (cm)	1.78 (0.78)	1.91 (0.8)	0.16
	St dev leave length (cm)	0.29 (0.14)	0.28 (0.13)	0.38
	Mean leave width (cm)	1.42 (0.56)	1.51 (0.53)	0.17
	St dev leave width (cm)	0.25 (0.11)	0.27 (0.12)	0.23
	Mean leave area (cm²)	2.94 (2.33)	3.25 (2.15)	0.2
	Time first flower (day)	**37.43 (4.56)**	**31.34 (5.55)**	**$2e^{-03}$**
	Number of flowering stems	12.05 (4.4)	11.84 (4.62)	0.05
	Maximum flowering stem length (cm)	56.78 (12.04)	52.55 (11.82)	0.1
	Number of flowers by flowering stem	56.56 (30.99)	56.22 (34.55)	0.11
Phytopathogen	**Flowering duration (day)**	**41.44 (8.13)**	**39.51 (7.32)**	**$9e^{-03}$**
	Time first leaf (day)	14.48 (4.4)	15.56 (5.2)	0.16
	Number of leaves	12.59 (3.76)	13.44 (3.35)	0.13
	Rosette area (cm²)	32.15 (32.24)	35.77 (26.16)	0.27
	Mean leave length (cm)	1.68 (0.83)	1.9 (0.78)	0.1
	St dev leave length (cm)	0.29 (0.1)	0.28 (0.14)	0.36
	Mean leave width (cm)	**1.29 (0.55)**	**1.51 (0.53)**	**0.03**
	St dev leave width (cm)	**0.22 (0.1)**	**0.27 (0.12)**	**0.03**
	Mean leave area (cm²)	2.59 (2.3)	3.25 (2.19)	0.08
	Time first flower (day)	**34.78 (6.93)**	**33.36 (4.92)**	**0.04**
Phytophagous	Number of flowering stems	10.78 (4.76)	12.15 (4.55)	0.34

	Maximum flowering stem length (cm)	**53.53 (9.81)**	**54.24 (12.2)**	**0.05**
	Number of flowers by flowering stem	47.46 (25.45)	58.12 (34.46)	0.17
	Flowering duration (day)	37.56 (8.1)	40.76 (7.76)	0.45
	Time first leaf (day)	14.92 (3.93)	15.46 (5.28)	0.34
	Number of leaves	**10.04 (3.88)**	**13.93 (2.96)**	**1e^{-04}**
	Rosette area (cm^2)	**11.79 (16.54)**	**39.64 (26.56)**	**1e^{-04}**
	Mean leave length (cm)	**1.05 (0.49)**	**2.02 (0.74)**	**1e^{-04}**
	St dev leave length (cm)	0.25 (0.1)	0.29 (0.14)	0.09
	Mean leave width (cm)	**0.88 (0.31)**	**1.59 (0.5)**	**1e^{-04}**
	St dev leave width (cm)	**0.23 (0.12)**	**0.27 (0.12)**	**0.04**
	Mean leave area (cm^2)	**1.05 (0.98)**	**3.54 (2.17)**	**1e^{-04}**
	Time first flower (day)	31.77 (5.6)	33.95 (5.25)	0.06
	Number of flowering stems	10.38 (4.85)	12.21 (4.54)	0.44
	Maximum flowering stem length (cm)	50.3 (13.1)	54.85 (11.71)	0.11
	Number of flowers by flowering stem	53.97 (55.98)	56.8 (28.05)	0.21
Oxidative stress	Flowering duration (day)	37.35 (6.5)	40.78 (7.94)	0.08

942 [a] *The P value represents the proportions of the distribution with the expected difference of mean*
943 *between individuals with and without attack after 10,000 random resamples equal to or less than the*
944 *value observed. Significant values are represented in bold.*

946 *Table S10: Difference on the phenotypic trait variation between the two families for each allele*
947 *tested.*

Allele	Family (Male/Female)	Trait (unit)	Mean (sd)	p value[a]
Ah01	*1 (d29.1/d24.1)*	**Time first leaf (day)**	**20.8 (5.43)**	**$2e^{-03}$**
		Number of leaves	11.9 (3.75)	0.09
		Rosette area (cm²)	24.9 (25.58)	0.11
		Mean leaf length (cm)	1.53 (0.79)	0.08
		St dev leaf length (cm)	0.21 (0.09)	0.03
		Mean leaf width (cm)	1.4 (0.58)	0.33
		St dev leaf width (cm)	0.24 (0.14)	0.32
		Mean leaf area (cm²)	2.47 (1.93)	0.17
		Time first flower (day)	28.8 (5.29)	0.44
		Number of flowering stems	**8 (4.07)**	**0.02**
		Maximum flowering stem length (cm)	**40.65 (6.76)**	**0.02**
		Number of flowers by flowering stem	**35.85 (10.72)**	**0.04**
		Flowering duration (day)	**40.35 (12.45)**	**$8e^{-03}$**
Ah01	*2 (d12.1/d29.1)*	**Time first leaf (day)**	**18.76 (5.57)**	**$1e^{-04}$**
		Number of leafs	13.45 (3.22)	0.42
		Rosette area (cm²)	31.99 (22.61)	0.25
		Mean leaf length (cm)	1.74 (0.71)	0.19
		St dev leaf length (cm)	0.29 (0.1)	0.42
		Mean leaf width (cm)	1.44 (0.51)	0.34
		St dev leaf width (cm)	0.29 (0.11)	0.11
		Mean leaf area (cm²)	2.84 (1.89)	0.21
		Time first flower (day)	**29.86 (5.43)**	**0.04**
		Number of flowering stems	9.59 (3.54)	0.32
		Maximum flowering stem length (cm)	**34.85 (9.35)**	**$1e^{-04}$**
		Number of flowers by flowering stem	**30.27 (25.45)**	**$1e^{-04}$**

		Flowering duration (day)	*31.62 (7.17)*	*0.06*
Ah03	*3 (d208.1/d17.1)*	*Time first leaf (day)*	*13.33 (5.27)*	*0.22*
		Number of leaves	*12.05 (4.25)*	*0.17*
		Rosette area (cm²)	*31.23 (25.7)*	*0.35*
		Mean leaf length (cm)	*1.67 (0.78)*	*0.32*
		St dev leaf length (cm)	*0.28 (0.15)*	*0.42*
		Mean leaf width (cm)	*1.39 (0.62)*	*0.5*
		St dev leaf width (cm)	*0.24 (0.13)*	*0.42*
		Mean leaf area (cm²)	*2.87 (2.24)*	*0.45*
		Time first flower (day)	***34.81 (7.87)***	***0.02***
		Number of flowering stems	***7.67 (3)***	***$3e^{-04}$***
		Maximum flowering stem length (cm)	***44.4 (13.87)***	***0.03***
		Number of flowers by flowering stem	*61.55 (46.06)*	*0.06*
		Flowering duration (day)	***39 (6.6)***	***$5e^{-03}$***
Ah03	*4 (d10.1/d17.1)*	Time first leaf (day)	18 (6.2)	0.12
		Number of leaves	12.33 (4.46)	0.22
		Rosette area (cm²)	**17.71 (11.23)**	**0.04**
		Mean leaf length (cm)	1.52 (0.53)	0.14
		St dev leaf length (cm)	0.23 (0.07)	0.14
		Mean leaf width (cm)	1.23 (0.35)	0.13
		St dev leaf width (cm)	0.23 (0.08)	0.23
		Mean leaf area (cm²)	2.01 (1.34)	0.1
		Time first flower (day)	33.33 (4.23)	0.09
		Number of flowering stems	11.67 (4.23)	0.22

		Maximum flowering stem length (cm)	57.25 (3.69)	0.34
		Number of flowers by flowering stem	75.05 (29.69)	0.12
		Flowering duration (day)	45.5 (8.67)	0.33
Ah04	5 (d205.1/d191.1)	**Time first leaf (day)**	**13.42 (3.28)**	**1e⁻⁰⁴**
		Number of leaves	13.45 (2.97)	0.32
		Rosette area (cm²)	36.29 (27.39)	0.31
		Mean leaf length (cm)	1.92 (0.8)	0.22
		St dev leaf length (cm)	0.29 (0.15)	0.34
		Mean leaf width (cm)	1.54 (0.56)	0.11
		St dev leaf width (cm)	0.27 (0.13)	0.15
		Mean leaf area (cm²)	3.36 (2.41)	0.14
		Time first flower (day)	35.1 (4.66)	0.33
		Number of flowering stems	12.87 (4.16)	0.32
		Maximum flowering stem length (cm)	**62.96 (9.48)**	**1e⁻⁰⁴**
		Number of flowers by flowering stem	**70.13 (32.5)**	**1e⁻⁰⁴**
		Flowering duration (day)	42.44 (7.05)	0.3
Ah04	6 (d205.1/d50.2)	Time first leaf (day)	14.31 (4.36)	0.30
		Number of leaves	13.5 (3.9)	0.13
		Rosette area (cm²)	**44.1 (33.48)**	**0.02**
		Mean leaf length (cm)	**2.04 (0.89)**	**0.04**
		St dev leaf length (cm)	0.29 (0.12)	0.21
		Mean leaf width (cm)	1.38 (0.51)	0.36
		St dev leaf width (cm)	**0.21 (0.09)**	**0.03**
		Mean leaf area (cm²)	3.33 (2.27)	0.21
		Time first flower (day)	**31.92 (3.66)**	**2e⁻⁰³**
		Number of flowering stems	**15.77 (5.25)**	**1e⁻⁰⁴**
		Maximum flowering stem length (cm)	60.01 (9.34)	0.07

		Maximum flowering stem length (cm)	57.25 (3.69)	0.34
		Number of flowers by flowering stem	75.05 (29.69)	0.12
		Flowering duration (day)	45.5 (8.67)	0.33
Ah04	5 (d205.1/d191.1)	**Time first leaf (day)**	**13.42 (3.28)**	$\mathbf{1e^{-04}}$
		Number of leaves	13.45 (2.97)	0.32
		Rosette area (cm²)	36.29 (27.39)	0.31
		Mean leaf length (cm)	1.92 (0.8)	0.22
		St dev leaf length (cm)	0.29 (0.15)	0.34
		Mean leaf width (cm)	1.54 (0.56)	0.11
		St dev leaf width (cm)	0.27 (0.13)	0.15
		Mean leaf area (cm²)	3.36 (2.41)	0.14
		Time first flower (day)	35.1 (4.66)	0.33
		Number of flowering stems	12.87 (4.16)	0.32
		Maximum flowering stem length (cm)	**62.96 (9.48)**	$\mathbf{1e^{-04}}$
		Number of flowers by flowering stem	**70.13 (32.5)**	$\mathbf{1e^{-04}}$
		Flowering duration (day)	42.44 (7.05)	0.3
Ah04	6 (d205.1/d50.2)	Time first leaf (day)	14.31 (4.36)	0.30
		Number of leaves	13.5 (3.9)	0.13
		Rosette area (cm²)	**44.1 (33.48)**	**0.02**
		Mean leaf length (cm)	**2.04 (0.89)**	**0.04**
		St dev leaf length (cm)	0.29 (0.12)	0.21
		Mean leaf width (cm)	1.38 (0.51)	0.36
		St dev leaf width (cm)	**0.21 (0.09)**	**0.03**
		Mean leaf area (cm²)	3.33 (2.27)	0.21
		Time first flower (day)	**31.92 (3.66)**	$\mathbf{2e^{-03}}$
		Number of flowering stems	**15.77 (5.25)**	$\mathbf{1e^{-04}}$
		Maximum flowering stem length (cm)	60.01 (9.34)	0.07

Number of flowers by flowering stem	*43.51 (15.45)*	*5e^{-04}*
Flowering duration (day)	*39.96 (7.38)*	*0.04*

948   *[a] The P value represents the proportions of the distribution with the expected difference of mean*
949   *between each family and the other individuals after 10,000 random resamples equal to or less than*
950   *the value observed. Significant values are represented in bold.*

References

1. Katsura, Y., Iwase, M., Satta, Y. (2012) Evolution of genomic structures on mammalian sex chromosomes. Current Genomics 13:115–123. https://doi.org/10.2174/138920212799860625.

2. Hartmann, F.E., Duhamel, M., Carpentier, F., et al. (2021) Recombination suppression and evolutionary strata around mating-type loci in fungi: documenting patterns and understanding evolutionary and mechanistic causes. New Phytologist 229:2470–2491. https://doi.org/10.1111/nph.17039.

3. Lamichhaney, S., Fan, G., Widemo, F., Gunnarsson, U., Thalmann, D.S., Hoeppner, M.P., Kerje, S., Gustafson, U., Shi, C., Zhang, H., Chen, W., Liang, X., Huang, L., Wang, J., Liang, E., Wu, Q., Lee, S.M.-Y., Xu, X., Höglund, J., Liu, X., and Andersson, L. (2016). Structural genomic changes underlie alternative reproductive strategies in the ruff (*Philomachus pugnax*). Nat Genet 48, 84–88. https://doi.org/10.1038/ng.3430.

4. Cocker, J.M., Wright, J., Li, J., Swarbreck, D., Dyer, S., Caccamo, M., and Gilmartin, P.M. (2018). *Primula vulgaris* (primrose) genome assembly, annotation and gene expression, with comparative genomics on the heterostyly supergene. Sci Rep 8, 17942. https://doi.org/10.1038/s41598-018-36304-4.

5. Uyenoyama, M.K. (1997). Genealogical structure among alleles regulating self-incompatibility in natural populations of flowering plants. Genetics 147, 1389–1400. https://doi.org/10.1093/genetics/147.3.1389.

6. Uyenoyama, M.K. (2005). Evolution under tight linkage to mating type. New Phytol. 165: 63–70. https://doi.org/10.1111/j.1469-8137.2004.01246.x.

7. Jay, P., Chouteau, M., Whibley, A., Bastide, H., Parrinello, H., Llaurens, V., and Joron, M. (2021). Mutation load at a mimicry supergene sheds new light on the evolution of inversion polymorphisms. Nat Genet 53, 288–293. https://doi.org/10.1038/s41588-020-00771-1.

8. Jay, P., Tezenas, E., Véber, A., and Giraud, T. (2022). Sheltering of deleterious mutations explains the stepwise extension of recombination suppression on sex chromosomes and other supergenes. PLOS Biology 20, e3001698. https://doi.org/10.1371/journal.pbio.3001698

9. Nettancourt, D. (2001). Incompatibility and incongruity in wild and cultivated plants. (2nd ed. Springer-Verlag, Berlin Heidelberg).

10. Schopfer, C.R., Nasrallah, M.E., and Nasrallah, J.B. (1999). The male determinant of self-incompatibility in Brassica. Science 286, 1697–1700. https://doi.org/10.1126/science.286.5445.1697.

11. Kusaba, M., Dwyer, K., Hendershot, J., Vrebalov, J., Nasrallah, J.B., and Nasrallah, M.E. (2001). Self-incompatibility in the genus Arabidopsis: characterization of the S locus in the outcrossing *A. lyrata* and its autogamous relative *A. thaliana*. Plant Cell 13, 627–643.

12. Uyenoyama, M.K. (2003). Genealogy-dependent variation in viability among self-incompatibility
genotypes. Theoretical Population Biology 63, 281–293. https://doi.org/10.1016/S0040-
5809(03)00020-0.

13. Llaurens, V., Gonthier, L., and Billiard, S. (2009). The sheltered genetic load linked to the S locus
in plants: new insights from theoretical and empirical approaches in sporophytic self-incompatibility.
Genetics 183, 1105–1118. https://doi.org/10.1534/genetics.109.102707.

14. Castric, V., and Vekemans, X. (2004). Plant self-incompatibility in natural populations: a critical
assessment of recent theoretical and empirical advances. Molecular Ecology 13, 2873–
2889.https://doi.org/10.1111/j.1365-294X.2004.02267.x.

15. Llaurens, V., Billiard, S., Leducq, J.-B., Castric, V., Klein, E.K., and Vekemans, X. (2008). Does
frequency-dependent selection with complex dominance interactions accurately predict allelic
frequencies at the self-incompatibility locus in *Arabidopsis halleri*? Evolution 62, 2545–2557.
https://doi.org/10.1111/j.1558-5646.2008.00469.x.

16. Durand, E., Méheust, R., Soucaze, M., Goubet, P.M., Gallina, S., Poux, C., Fobis-Loisy, I., Guillon,
E., Gaude, T., Sarazin, A., et al. (2014). Dominance hierarchy arising from the evolution of a complex
small RNA regulatory network. Science 346, 1200–1205. https://doi.org/10.1111/eva.12933.

17. Billiard, S., Castric, V., and Vekemans, X. (2006). A general model to explore complex dominance
patterns in plant sporophytic self-incompatibility systems. Genetics 175, 1351–1369.
https://doi.org/10.1016/j.tig.2011.06.005.

18. Goubet, P.M., Bergès, H., Bellec, A., Prat, E., Helmstetter, N., Mangenot, S., Gallina, S., Holl, A.-C.,
Fobis-Loisy, I., Vekemans, X., et al. (2012). Contrasted patterns of molecular evolution in dominant
and recessive self-incompatibility haplotypes in Arabidopsis. PLoS Genetics 8, e1002495.
https://doi.org/10.1371/journal.pgen.1002495.

19. Stift, M., Hunter, B.D., Shaw, B., Adam, A., Hoebe, P.N., and Mable, B.K. (2013). Inbreeding
depression in self-incompatible North-American *Arabidopsis lyrata*: disentangling genomic and S-
locus-specific genetic load. Heredity 110, 19–28. https://doi.org/10.1038/hdy.2012.49.

20. Le Veve, A., Burghgraeve, N., Genete, M., et al. (2023) Long-term balancing selection and the
genetic load linked to the self-incompatibility locus in *Arabidopsis halleri* and *A. lyrata*. Molecular
Biology and Evolution msad120. https://doi.org/10.1093/molbev/msad120.

21. Hu, T.T., Pattyn, P., Bakker, E.G.,, Cao, J., Cheng, J.-F., Clark, R.M., Fahlgren, N., Fawcett, J.A.,
Grimwood, J., Gundlach, H., et al. (2011). The *Arabidopsis lyrata* genome sequence and the basis of
rapid genome size change. Nat Genet 43, 476–481. https://doi.org/10.1038/ng.807.

22. Foxe, J.P., Stift, M., Tedder, A., Haudry, A., Wright, S.I., and Mable, B.K. (2010). Reconstructing
origins of loss of self-incompatibility and selfing in North american *Arabidopsis lyrata*: a population
genetic context. Evolution 64, 3495–3510. https://doi.org/10.1111/j.1558-5646.2010.01094.x.

23. Charlesworth, D. (2006). Balancing selection and its effects on sequences in nearby genome
regions. PLOS Genetics 2, e64. https://doi.org/10.1371/journal.pgen.0020064.

24. Kamau, E., Charlesworth, B., Charlesworth, D. (2007) Linkage Disequilibrium and Recombination
Rate Estimates in the Self-Incompatibility Region of Arabidopsis lyrata. Genetics 176, 2357–2369.
https://doi.org/10.1534/genetics.107.072231

25. Stone, J.L. (2004). Sheltered load associated with S-alleles in *Solanum carolinense*. Heredity 92,
335–342. https://doi.org/10.1038/sj.hdy.6800425.

26. Lane, M.D., and Lawrence, M.J. (1995). The population genetics of the self-incompatibility
polymorphism in *Papaver rhoeas*. X. An association between incompatibility genotype and seed
dormancy. Heredity 75, 92–97. https://doi.org/10.1038/hdy.1995.108.

27. Vieira, J., Pimenta, J., Gomes, A., Laia, J., Rocha, S., Heitzler, P., and Vieira, C.P. (2021). The
identification of the Rosa S-locus and implications on the evolution of the Rosaceae gametophytic
self-incompatibility systems. Sci Rep 11, 3710. https://doi.org/10.1038/s41598-021-83243-8.

28. Mena-Ali, J. I., Keser, L. H., Stephenson, A. G. (2009). The effect of sheltered load on
reproduction in *Solanum carolinense*, a species with variable self-incompatibility. Sex. Plant Reprod.
22 63–67. https://doi.org/10.1007/s00497-008-0092-x.

29. Clauss, M.J., and Mitchell-Olds, T. (2006). Population genetic structure of *Arabidopsis lyrata* in
Europe. Molecular Ecology 15, 2753–2766. Clauss, M.J., Mitchell-Olds, T., 2006. Population genetic
structure of *Arabidopsis lyrata* in Europe. Molecular Ecology 15, 2753–2766.
https://doi.org/10.1111/j.1365-294X.2006.02973.x.

30. Ross-Ibarra, J., Wright, S.I., Foxe, J.P., Kawabe, A., DeRose-Wilson, L., Gos, G., Charlesworth, D.,
and Gaut, B.S. (2008). Patterns of polymorphism and demographic history in natural populations of
Arabidopsis lyrata. PLOS ONE 3, e2411. https://doi.org/10.1371/journal.pone.0002411.

31. Pauwels, M., Saumitou-Laprade, P., Holl, A.C., Petit, D., and Bonnin, I. (2005). Multiple origin of
metallicolous populations of the pseudometallophyte *Arabidopsis halleri* (Brassicaceae) in central
Europe: the cpDNA testimony. Molecular Ecology 14, 4403–4414. https://doi.org/10.1111/j.1365-
294X.2005.02739.x.

32. Stetsenko, R., Brom, T., Castric, V., and Billiard, S. (2023). Balancing selection and the crossing of
fitness valleys in structured populations: diversification in the gametophytic self-incompatibility
system. Evolution 77:907–920. https://doi.org/10.1093/evolut/qpac065

33. Gervais, C.E., Castric, V., Ressayre, A., and Billiard, S. (2011). Origin and diversification dynamics
of self-Incompatibility haplotypes. Genetics 188, 625–636.
https://doi.org/10.1534/genetics.111.127399.

34. Bergero, R., and Charlesworth, D. (2009). The evolution of restricted recombination in sex
chromosomes. Trends in Ecology & Evolution 24, 94–102.
https://doi.org/10.1016/j.tree.2008.09.010.

35. Wright, A.E., Dean, R., Zimmer, F., and Mank, J.E. (2016). How to make a sex chromosome. Nat
Commun 7, 12087. https://doi.org/10.1038/ncomms12087.

36. Ponnikas, S., Sigeman, H., Abbott, J.K., and Hansson, B. (2018). Why do sex chromosomes stop
recombining? Trends Genet 34, 492–503. https://doi.org/10.1016/j.tig.2018.04.001.

37. Hill, J., Enbody, E., Bi, H., Lamichhaney, S., Schwochow, D., Younis, S., Widemo, F., and
Andersson, L. (2022). Low mutation load in a supergene underpinning alternative male mating
strategies in ruff. https://doi.org/10.1101/2022.04.27.489720.

38. Rosser, N., Edelman, N.B., Queste, L.M., et al. (2022) Complex basis of hybrid female sterility and
Haldane's rule in Heliconius butterflies: Z-linkage and epistasis. Molecular Ecology 31:959–977.
https://doi.org/10.1111/mec.16272.

39. Durand, E., Chantreau, M., Le Veve, A., et al. (2020). Evolution of self-incompatibility in the
Brassicaceae: Lessons from a textbook example of natural selection. Evolutionary Applications.
https://doi.org/10.1111/eva.12933.

40. Llaurens, V., Billiard, S., Castric, V., and Vekemans, X. (2009). Evolution of dominance in
sporophytic self-incompatibility systems: I. Genetic load and coevolution of levels of dominance in
pollen and pistil. Evolution 63, 2427–2437. https://doi.org/10.1111/j.1558-5646.2009.00709.x.

41. Genete, M., Castric, V., and Vekemans, X. (2020). Genotyping and *de novo* discovery of allelic
variants at the Brassicaceae self-incompatibility locus from short read sequencing data. Mol Biol
Evol. https://doi.org/10.1093/molbev/msz258.

42. Schierup, M.H., Mikkelsen, A.M., and Hein, J. (2001). Recombination, balancing selection and
phylogenies in *MHC* and self-incompatibility genes. Genetics 159, 1833–1844.
https://doi.org/10.1093/genetics/159.4.1833

43. Mable, B.K., Schierup, M.H., and Charlesworth, D. (2003). Estimating the number, frequency, and
dominance of S -alleles in a natural population of *Arabidopsis lyrata* (Brassicaceae) with sporophytic
control of self-incompatibility. Heredity 90, 422–431. https://doi.org/10.1038/sj.hdy.6800261.

44. Bechsgaard, J., Bataillon, T., and Schierup, M.H. (2004). Uneven segregation of sporophytic self-
incompatibility alleles in *Arabidopsis lyrata*. Journal of Evolutionary Biology 17, 554–561.
https://doi.org/10.1111/j.1420-9101.2004.00699.x.

45. Prigoda, N.L., Nassuth, A., and Mable, B.K. (2005). Phenotypic and genotypic expression of self-
incompatibility haplotypes in *Arabidopsis lyrata* suggests unique origin of alleles in different
dominance classes. Molecular Biology and Evolution 22, 1609–1620.
https://doi.org/10.1093/molbev/msi153.

46. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat Methods
9, 357–359. https://doi.org/10.1038/nmeth.1923.

47. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin,
R., and 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map
format and SAMtools. Bioinformatics 25, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352.

48. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics 26, 841–842. https://doi.org/10.1093/bioinformatics/btq033.

49. DePristo, M.A., Banks, E., Poplin, R.E., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis, A.A.,
del Angel, G., Rivas, M.A., Hanna, M., et al. (2011). A framework for variation discovery and
genotyping using next-generation DNA sequencing data. Nat Genet 43, 491–498.
https://doi.org/10.1038/ng.806.

50. Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E.,
Lunter, G., Marth, G.T., Sherry, S.T., et al. (2011). The variant call format and VCFtools.
Bioinformatics 27, 2156–2158. https://doi.org/10.1093/bioinformatics/btr330.

51. Williamson, R.J., Josephs, E.B., Platts, A.E., Hazzouri, K.M., Haudry, A., Blanchette, M., and
Wright, S.I. (2014). Evidence for widespread positive and negative selection in coding and conserved
noncoding regions of *Capsella grandiflora*. PLOS Genetics 10, e1004622.
https://doi.org/10.1371/journal.pgen.1004622.

52. Guindon, S., Dufayard, J-F., Lefort, V., et al (2010) New algorithms and methods to estimate
maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. Syst Biol 59, 307–321.
https://doi.org/10.1093/sysbio/syq010.

53. Tamura, K., and Nei, M. (1993). Estimation of the number of nucleotide substitutions in the
control region of mitochondrial DNA in humans and chimpanzees. Molecular Biology and Evolution
10, 512–526. https://doi.org/10.1093/oxfordjournals.molbev.a040023.

54. Kumar S, Stecher G, Li M, Knyaz C, and Tamura K (2018) MEGA X: Molecular evolutionary
genetics analysis across computing platforms. Molecular Biology and Evolution 35:1547-1549.
https://doi.org/10.1093/molbev/msy096.

55. Rozas, J., Ferrer-Mata, A., Sánchez-DelBarrio, J.C., Guirao-Rico, S., Librado, P., Ramos-Onsins, S.E.,
and Sánchez-Gracia, A. (2017). DnaSP 6: DNA sequence polymorphism analysis of large data sets.
Molecular Biology and Evolution 34, 3299–3302. https://doi.org/10.1093/molbev/msx248.

56. Pritchard, J. K., Stephens, M., Rosenberg, N. A., and Donnelly, P. (2000). Association mapping in
structured populations. The American Journal of Human Genetics 67:170–181.
https://doi.org/10.1086/302959.

57. Pritchard, J. K., Wen, X., & Falush, D. (2010). Documentation for structure software: Version 2.3.
University of Chicago, Chicago, IL, 1-37.

58. Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to ImageJ: 25 years of image
analysis. Nat Methods 9, 671–675. https://doi.org/10.1038/nmeth.2089.