Oral contraceptives reduce the avoidance bias to social threat signals

Jasmin Thurley[1], Macià Buades Rotger[2], Georg Serfling[3], Thessa Howaldt[1], Nicole Reisch[4],

Ulrike M. Krämer[1,5,6*]

[1] Dept. of Neurology, University of Lübeck, Germany

[2] Department of Clinical Psychology and Psychobiology, University of Barcelona, Barcelona,

Spain

[3] Medizinische Klinik I und Institut für Endokrinologie und Diabetes, Universitätsklinikum

Schleswig-Holstein, Campus Lübeck, Universität zu Lübeck, Lübeck, Germany

[4] Medizinische Klinik und Poliklinik IV Klinikum der Universität München, Munich, Germany

[5] Center of Brain, Behavior and Metabolism (CBBM), University of Lübeck, Lübeck,

Germany

[6] Department of Psychology, University of Lübeck, Lübeck, Germany

* Corresponding author:

Ulrike M. Krämer, Ph.D.

Department of Neurology, University of Lübeck

Ratzeburger Allee 160

23538 Lübeck, Germany

+49 451 31017421

ulrike.kraemer@uni-luebeck.de

29   **Abstract**

Recent research has increasingly acknowledged the impact of oral contraceptives on affective

behavior and stress responses; however, the underlying mechanisms are still not well

understood. Studies have previously shown that steroid hormones modulate automatic

approach and avoidance behavior. Here, we thus investigated the effects of oral

contraceptives on approach and avoidance behavior and whether these effects are modulated

by stress. The study comprised 130 female participants, half of whom were using oral

contraceptives, while the other half were not using any hormonal contraception (NC). The

participants completed the Approach Avoidance Task (AAT), which measures automatic

approach and avoidance behavior to socio-affective signals. The AAT was run once before

and once after a stress manipulation using the Socially Evaluated Cold Pressor Test. OC

users showed absent avoidance behavior to social threat signals and a stress-induced

increase in approach behavior to positive social signals. The latter was found in particular in

women taking androgenic acting OC, demonstrating that different OC preparations need to be

taken into account in research on OC effects. However, OC and NC group did not differ in

their cortisol stress response. Overall, the results suggest that OC usage impacts on approach

and avoidance behavior to social signals, which might also contribute to the development of

affective side effects.

Keywords: oral contraceptives, approach avoidance behavior, stress

1. Introduction

Oral contraceptives (OC) are amongst the most reliable methods to prevent undesired pregnancies and are the contraceptive of choice for many women worldwide (Christin-Maitre, 2013). Hormonal contraception, however, is associated with side effects, including changes in affective experience and a possibly increased risk for depression (Montoya & Bos, 2017). The impact of oral contraceptives on specific affective and cognitive mechanisms, which could contribute to these side effects, remains unclear. Altered approach and avoidance behavior may be a possible mechanism for several reasons: The steroid hormones, which are suppressed by OC, bind to receptors in brain regions implicated in approach and avoidance behavior and its regulation (Volman et al., 2013). OC users have been found to show functional and structural changes in these brain regions, namely the amygdala and prefrontal cortex (Petersen & Cahill, 2015; Sharma et al., 2020). Moreover, other steroid hormones, namely testosterone and cortisol, appear to have an impact on approach and avoidance behaviors (Radke et al., 2015; Roelofs et al., 2005). Lastly, considering that OC influence the stress axis and cortisol reactivity (Mordecai et al., 2017), which in turn affects approach and avoidance behavior (Roelofs et al., 2005), it is possible that OC influence affective behavior in conjunction with stress. Here, we thus tested whether women who are taking OC show altered approach and avoidance behavior, and whether this effect is modulated by stress using a cross-sectional design.

The evidence on the influence of oral contraceptives on affect is inconclusive. While some studies reported that only a small proportion of OC users experienced affective side-effects (Ernst et al., 2002), others, including a placebo-controlled longitudinal study, reported more depressive symptoms, mood swings and fatigue in OC users compared with the placebo group (Gingnell et al., 2013). Other recent studies have started to tap into the psychological and neural mechanisms responsible for these affective side-effects by investigating which specific cognitive, affective and neural processes are affected by OC usage (Brønnick et al., 2020; Lewis et al., 2019; Montoya & Bos, 2017).

One potential mechanism underlying affective symptoms might be altered approach

and avoidance behavior. Approach/avoidance behavior refers to the automatic tendency to

approach positive stimuli and avoid negative stimuli, which can be measured using the

Approach Avoidance task (AAT) (Chen & Bargh, 1999). In one commonly used variant of the

AAT (Kaldewaij et al., 2017), participants are asked to pull pictures of happy faces towards

them and push negative faces away as fast as possible using a joystick. Response times in

this congruent condition are compared with those of the reverse, incongruent condition (pull

angry faces and push happy faces away) (Beyer et al., 2017). Typically, participants are faster

in the congruent compared to the incongruent condition, referred to as congruency effect or

AAT bias. Previous research with the AAT reported that people with social anxiety show faster

avoidance (Roelofs et al., 2010), whereas psychopathic patients showed reduced avoidance

to angry faces (von Borries et al., 2012), speaking for the validity of the AAT.

A neurobiological model of the AAT (Volman et al., 2013) assumed that faces are first

processed in the fusiform face area (FFA) and subsequently projected to the amygdala, which

has a bidirectional connection to the anterior prefrontal cortex (aPFC). When automatic

affective behavior needs to be controlled in the incongruent condition, amygdala activity is

believed to be inhibited by the aPFC (Volman et al., 2013). Oral contraceptive use has

previously been found to be associated with changes in reactivity and activation of the PFC

and amygdala. OC women responded to negative emotional stimuli with reduced amygdala

reactivity and increased prefrontal activation (Petersen & Cahill, 2015; Sharma et al., 2020).

Furthermore, estradiol and progesterone receptors are expressed in the amygdala, whereby

their density may be influenced by oral contraceptives according to results from animal studies

(Brown, 2020; Österlund et al., 1998). Other steroid hormones whose receptors are also

located on the amygdala such as cortisol (Morimoto et al., 1996) and testosterone (Simerly et

al., 1990) are known to modulate AAT effects (Radke et al., 2015; Roelofs et al., 2005).

Moreover, Li and colleagues reported that approach and avoidance behavior was dependent

on menstrual cycle phase and estrogen and progesterone levels (Li et al., 2022). These data

suggest that OC women might differ from NC women in their behavior in the AAT. Since earlier

studies have found differences mainly for negatively arousing stimuli (Petersen & Cahill, 2015;

Sharma et al., 2020), we expect group differences especially for angry faces.

In addition to the contraceptive effect OC have on the HPG-axis (hypothalamic–

pituitary–gonadal axis), they also affect the HPA stress axis (hypothalamic-pituitary-adrenal

axis) (Hertel et al., 2017; Meulenberg & Hofman, 1990). While plasma cortisol is elevated

during OC intake, the stress-induced increase in free cortisol is reduced in OC users

(Kirschbaum et al., 1999). Although these changes in the HPA axis activity in OC users are

well documented, their behavioral implications are less clear. Given previous evidence that a

high cortisol increase due to social stress is associated with a reduced AAT congruency effect

(Roelofs et al., 2005), OC might also influence approach/avoidance behavior through this

pathway. In the current study, we therefore tested whether approach/avoidance behavior in

OC users interacts with stress exposure.

One challenge in research on OC is the large heterogeneity of preparations taken by

women (Tronson & Schuh, 2022). In our study, only women taking combined preparations

containing an estrogen and a progestin were included, but the preparations nevertheless

differed in the included estradiol dose and in the used anti- or androgenic acting progestin.

The different formulations could have an influence on AAT and stress effects via the different

efficacy profiles, as there is for example evidence that women with androgenic OC have a

more pronounced cardiovascular stress response (Straneva et al., 2000). Therefore, we

performed exploratory analyses on the subgroups.

2. Methods

2.1 Participants

We tested one hundred thirty women, of whom sixty-five had been using oral contraceptives

for at least three months (oral contraceptives, OC group) and 65 had not used any hormonal

contraception for at least six months (no hormonal contraception, NC group). The mean age

in the OC group was 21.9 years (SD = 2.13) and in the NC group 22.6 years (SD = 2.4). They

had all normal or corrected to normal vision (self-report). An analysis with G*Power 3.1.9.6

(Faul et al., 2007) suggested that at least 128 participants would be required to find an effect

of medium effect size for group differences in the AAT effect (f = 0.25) with a power of 0.8 and

an alpha error of 0.05. All participants were between 19 and 30 years old, had a BMI between

17.5 and 25 and no neurological, psychiatric, cardiovascular or endocrinological pre-existing

diseases.

Of the 130 participants, we had to exclude three for the following reasons: one due to

many errors in the AAT (mean error rate: 39.15%) and two due to reaction times in the AAT

that were more than 3 SD above the mean. Saliva samples from 20 participants could not be

used, resulting in a final sample of 106 participants (50=OC, 56=NC) for the cortisol analyses.

All of the women in the OC group were using combined oral contraceptives containing both

estradiol and progestin. The doses of ethinyl estradiol varied between 20 and 30 µg. Three

women where using OC with estetrol (14.2 mg) as estradiol and 4 women took OC containing

estradiol valerate (1-3 mg). The doses of progestin ranged between 100 and 3000 µg. Twelve

of the OC women reported affective side effects. The participants were recruited via mailing

lists of the University of Lübeck. For the verification of the inclusion and exclusion criteria, the

potential participants were subsequently called. The person who conducted the calls was not

involved in the measurements and the investigators were blinded to the group membership of

the respective participants. All participants were informed about the study objectives and

compensated for their participation with 10 Euros per hour or credit points. All participants

gave their written informed consent in accordance with the Declaration of Helsinki. The Ethics

Committee of the University of Lübeck approved this study protocol (22-044).

2.2 Design and Procedure

OC women were tested between the first and third day of pill intake, NC women

correspondingly on the eighth to tenth day of the menstrual cycle (follicular phase). The timing

was chosen because interpersonal variations in cycle length (in the NC group) and associated

differences in hormone concentrations are lowest at the beginning of the cycle (Hampson,

2020). Moreover, the low levels of estrogen and progesterone during this period are most

comparable to the levels of endogenous estradiol and progesterone that result from taking

oral contraceptives. Effects can therefore be attributed to the chronic intake of OC and not to

acute hormonal variations. All measurements took place in the afternoon between 2 and 7

p.m., since during this period salivary cortisol concentrations are relatively consistently low for

both women taking oral contraceptives and women with no hormonal contraception

(Meulenberg & Hofman, 1990).

After participants gave their informed consent, they were asked to fill out some

computerized questionnaires assessing emotion regulation, personality, and

approach/avoidance tendencies (see 2.6 Questionnaires and Demographic Data). In the next

step, they provided a saliva sample before performing the AAT on the computer (Figure 1).

After another saliva sample, the Socially evaluated cold pressor test (SECPT) (Schwabe &

Schächinger, 2018) was performed. After a break of approximately 12 minutes, the AAT was

performed a second time. After completion of the second AAT, another saliva sample was

collected and the participants filled in a questionnaire with demographic data and information

about the pill. Lastly, participants were debriefed about the research question and were

compensated. The testing lasted approximately one hour in total.

2.3 Approach Avoidance Task (AAT)

We used the Approach Avoidance Task (Chen & Bargh, 1999) in the version of Beyer et al.

(2017) with images of happy and angry faces from the Radboud Faces database (Langner et

al., 2010) as stimuli. Stimuli were presented using Presentation® (Neurobehavioral Systems).

In total, images of thirty individuals (15 male, 15 female) are shown during the AAT. There is

one picture of each person with an angry and a happy facial expression. For the practice

blocks, pictures of nine different people than in the main block are used. All pictures are cut

into an oval shape so that hair, neck and ears are not visible.

Participants used the joystick with their dominant hand. Before each stimulus, a small

black cross appeared on the white screen. As soon as the "shoot" button of the joystick was

pressed, the cross disappeared and a face appeared in the center of the screen. By pushing

the joystick away (avoidance response), the image got smaller step-by-step (seven steps in

total). Pulling the joystick towards the body (approach response) made the image larger. Two

conditions were run in two different blocks. In the congruent condition, participants were asked

to pull happy faces toward them as quickly as possible and to push angry faces away from

them. In the incongruent condition, the instruction was reversed. Here, happy faces were to

be pushed away from the body and angry ones were to be pulled in. Each block contained

thirty happy faces and thirty angry faces, using the same images in both blocks.

Before each block, the participants could familiarize themselves with the task in a

practice session. Here, the participants received direct feedback whether they had reacted

correctly (green tick for correct reaction and red cross for incorrect reaction). The first exercise

run consisted of twenty images. In the second practice run, 28 pictures were used because

the instruction that was internalized in the first block was reversed and task-switching effects

were to be avoided.

During each trial the reaction time was measured, which is determined as the interval

between the appearance of the stimulus and the start of the movement of the joystick.

2.4 Socially Evaluated Cold Pressor Test (SECPT)

The SECPT was performed as suggested by Schwabe and Schächinger (2018). The

participants were instructed to hold their dominant hand, including the wrist, in a bucket of ice

water (0-2 degrees Celsius) until the experimenter asked them to take the hand out again

(they were not told in advance how long they should keep the hand in the ice water). Only if

they could no longer stand the cold at all, they were allowed to take their hand out of the water

earlier. They were also told that they were videotaped during the experiment so that their facial

expressions can be analyzed later. Therefore, they were instructed to look into the camera

during the entire experiment and not to speak. The camera was placed about two meters in

front of the participants behind a screen facing them. The camera recorded the subject's face

and transferred it directly to the screen so that the participant could see herself. Next to the

camera, the experimenter stood at a distance that allowed the participant to simultaneously

look into the camera and see the experimenter out of the corner of her eye. The experimenter

made sure that the participant did not make a fist and that her hand was in ice water up to the

wrist. At the same time, she took notes and avoided giving confirming signals such as smiling

or nodding. After three minutes, the test ended and the participant was allowed to take her

hand out of the water. If she took her hand out before that, the experimenter responded by

asking the participant to put her hand back in the water. If this was not possible, the participant

was asked to remain standing and looking into the camera for the remainder of the three

minutes.

2.5 Saliva Sampling

Saliva samples were collected at three time points of the measurement. For this purpose, we

used Sarstedt cortisol Salivette® with a synthetic fiber roll. To prevent cortisol levels from

being influenced by factors other than the experimental factors, participants were instructed

not to exercise, eat anything, drink coffee, take medications, or smoke for one hour before the

appointment. Saliva samples were frozen and stored at -20°C until analysis. After thawing,

samples were centrifuged at 3,000 rpm for 5 min, which resulted in a clear supernatant of low

viscosity.   Salivary   concentrations   were   measured   using   commercially   available

chemiluminescence immunoassay with high sensitivity (Tecan - IBL International, Hamburg,

Germany; catalogue number R62111). The intra and interassay coefficients of variance were

below 9%.   Samples were considered empty if there was no saliva in the tube after

centrifugation.

2.6 Questionnaires and Demographic Data

We used the Behavioral Inhibition Scale (BIS) and the Behavioral Approach Scale (BAS)

designed by Carver and White (1994) to assess tendencies to exhibit avoidance or approach

behaviors. We also used the Emotion Regulation Questionnaire (ERQ) (Gross & John, 2003)

and the Big Five Inventory (BFI-10) (Rammstedt & John, 2007) to examine interpersonal

differences in emotion regulation and personality structure.

Participants also provided information on the name of the OC preparation taken,

previous duration of use and age at first use of oral contraceptives, time of daily use, reason

for use, and any side effects (OC group). The NC-women where asked if they had used

hormonal contraception in the past. If they had taken OC in the past, the duration and the time

since the last intake was recorded. The reason for taking the pill and the occurrence of any

side effects were asked as well. The weight, height and age of all test participants was

recorded.

Figure 1. A Procedure. At first, the participants completed the questionnaires ERQ, BFI-10, and BIS/BAS. Next,
they provided a saliva sample and performed the AAT (approach avoidance task). After another saliva sample,
they performed the Socially Evaluated Cold Pressor Test (SECPT) and another run of the AAT. 25 minutes after
the onset of the SECPT, the participants gave another saliva sample and finally provided demographic information.
B Approach Avoidance Task (AAT): Participants were asked to push away angry faces and pull happy faces as
fast as possible in the congruent condition and do it the other way round in the incongruent condition. Note that
drawings are shown for visualization only. In the experiment, we showed images of happy and angry faces, which
were taken from the Radboud Faces database (Langner et al., 2010). **C** Socially evaluated cold pressor test

(SECPT). Participants were required to keep their dominant hand in ice water for 3 minutes while being observed
by the experimenter and being recorded on video.
270

2.7 Data analysis

The main outcome of the AAT were the reaction times (RT) which were defined as the interval

between stimulus presentation and movement onset. Following previous work (Beyer et al.,

2017), we excluded trials with response latencies of less than 150 ms or more than 3 SD from

the individual participant's own mean. Moreover, we only considered correct trials. Incorrect

trials (errors), which also included trials with an initial movement into the wrong direction and

a following correction were excluded. We first log-transformed the reaction times (ms) before

computing mean response times for each condition (push, pull for angry and happy) and

participant. We also considered the error rates for determination of outliers. Participants were

excluded if reaction times (n=2) or error rates (n=1) in the AAT were more than 3 SD above

the mean across all participants (n = 2).

We computed RT means, outliers and error rates using MATLAB (MATLAB, 2021).

Statistical analyses were conducted in Jamovi (The jamovi project, 2021) and JASP (JASP

Team, 2023).

We performed a mixed effects ANOVA that included the factors emotion (happy,

angry), movement (push, pull), stress (within factors) and contraception (between factor) using

log-transformed reaction times. To parse interaction effects, we conducted further ANOVAs

to examine group differences between OC and NC women separately for happy and angry

faces and before and after stress. To determine the AAT effect, the reaction times of the

movement away from the body (push) were subtracted from the reaction times of the

movement towards the body (pull). A higher (more positive) score here reflects stronger

avoidance behavior and a lower (more negative) score reflects approach behavior. The AAT

effect was calculated as the difference between the bias score for angry faces and the bias

score for happy faces. Accordingly, the higher the bias scores, the larger the effect. The AAT

effect was reported only for visualization and to allow for comparability of results with previous

studies and is not interpreted on its own in this study. We conducted an additional mixed

effects ANOVA with factors stress and contraception to compare the error rates of OC and

NC women before and after stress.

To investigate the change in cortisol levels, we performed another mixed effects

ANOVA. Here, the within factor time (t_0, t_1, t_2) and the between factor contraception were

included. For the comparison of group differences in the questionnaires, we calculated two-

sided independent samples t-tests.

For further exploratory analyses of subgroups within the OC group, we ran additional

mixed effects ANOVAs (with the factors emotion, movement, stress (within factors) and the

respective between factors (androgenicity, EE-dose, onset, time of OC intake or

contraception). Moreover, we ran a correlation of duration of OC intake and the AAT effect

and a correlation of the AAT effect with the Emotion Regulation Questionnaire subscales

(across the whole sample).

All results with a p-value less than .05 were considered significant. For significant

ANOVA results, we used Bonferroni-Holm adjusted p-values for post-hoc tests. In order to

make informed statements regarding the evidence, we also provide the Bayes Factor, which

tells us how much more probable the observations are under the alternative hypothesis

compared to the null hypothesis. Our Bayesian analysis was conducted using the default

priors in JASP, and we present the findings of Bayesian model averaging (BF_{incl}) (Love et al.,

2019). To interpret the results, we adhere to established guidelines: BF values exceeding 3

indicate moderate support, while values above 10 indicate strong support for the alternative

hypothesis. Conversely, BF values below 0.1 strongly support the null hypothesis, and values

below 0.33 suggest moderate support (van Doorn et al., 2021). Because Bayesian analyses

are time-consuming and computationally expensive for large-sample multifactorial ANOVAs,

we report Bayes factors exclusively for specific lower-order effects.

3. Results

3.1 Approach Avoidance Behavior

The AAT showed the expected experimental effects, as participants showed avoidance of

angry faces and approach toward happy faces. Visual examination of the behavioral data

suggested, however, that OC women responded less avoidant overall and even showed a

slight tendency to approach angry faces before the SECPT (see Table 1 and Figure 2A&B).

Note that the reaction times were log-transformed to increase normality.

Results of the mixed ANOVA showed that, as expected, the responses to the emotions

in the congruent and incongruent condition differed in speed (see Figure 2A-D). Overall, it took

the participants longer to pull angry faces towards them than to push them away, whereas

they showed the opposite pattern for happy faces (emotion x movement: $F(1,125) = 66.733$,

$p < .001$, $\eta^2_p = 0.348$), reflecting the expected AAT effect. In general, the groups did not differ

in the expression of this pattern (emotion x movement x contraception: $F(1,125) = 2.482$, $p =$

$.118$, $\eta^2_p = 0.019$). However, there was a significant fourfold interaction of stress x emotion x

movement x contraception ($F(1,125) = 4.345$, $p = .039$, $\eta^2_p = 0.034$). Similar to previous studies

with the AAT (Radke et al., 2015; Volman et al., 2013), participants were faster in pull than

push movements (movement: $F(1,125) = 95.735$, $p < .001$, $\eta^2_p = 0.434$). Interestingly, OC

women responded overall faster than NC women (contraception: $F(1,125) = 6.35$, $p = .013$,

$\eta^2_p = 0.048$) (see Figure 2D). Women in the OC group were especially faster than NC women

in their pull movements, and less so in their push movements (interaction of movement x

contraception: $F(1,125) = 6.019$, $p = .016$, $\eta^2_p = 0.046$; pull: $t(125) = 2.827$, $p_{Holm} = .016$, $d =$

0.461; push: $t(125) = 2.166$, $p_{Holm} = .064$, $d = 0.353$).

To understand the four-way interaction of stress x emotion x movement x

contraception, we computed separate mixed effects ANOVAs for the two emotions (happy

and angry), since we expected differences especially for angry faces. In response to angry

faces, the groups indeed differed in their behavior, reflected in a significant interaction of

movement x contraception ($F(1,125) = 6.708$, $p = .011$, $\eta^2_p = 0.051$, $BF = 3.916$). The Bayes

factor indicated moderate support for this interaction. While NC women pushed away angry

faces faster than they pulled them closer ($p_{Holm} = 0.003$), showing the typical avoidance bias,

OC women responded equally fast when pushing and pulling (p_{Holm} = 0.937; see Figure 2C).

This pattern was not influenced by stress (stress x movement x contraception: $F(1,125)$ =

1.155, p = .285, η^2_p = 0.009, BF = 0.305). The groups did not differ in their general behavior

to happy faces (movement x contraception: $F(1,125)$ = 0.039, p = .843, η^2_p < 0.001, BF =

0.216), however, there was a threefold interaction of stress x movement x contraception for

happy faces ($F(1,125)$ = 6.596, p = .011, η^2_p = 0.050, BF = 3.073). To understand this

interaction, we computed separate mixed effects ANOVAs for the two groups (NC and OC).

While stress had no influence on approach behavior to happy faces in NC women (stress x

movement: $F(1,62)$ = 1.56, p = .216, η^2_p = 0.025, BF = 1.419), OC women differed in their

reactions towards happy faces before and after stress (stress x movement: $F(1,63)$ = 5.68, p

= .02, η^2_p = 0.083, BF = 9.59). This was due to more pronounced approach behavior after

stress than before stress (movement after stress: $F(1,63)$ = 87.2, p < .001, η^2_p = 0.580, BF =

6069.494; movement before stress: $F(1,63)$ = 27.1, p < .001, η^2_p = 0.301, BF = 2.379x10^{10})

(see Figure 2A&B). Note however, that OC women showed a clear approach bias to happy

faces both before and after stress.

Regarding error rates in the AAT, there was no group difference before or after stress

(mixed effects ANOVA with factors stress and contraception; contraception: $F(1,125)$ = 1.05,

p = 0.308, η^2_p = 0.008, BF = 0.257) (Figure 2E).

3.2 Stress effects on cortisol level

The stress manipulation generally succeeded, as cortisol levels were higher on average after

the SECPT than before. About 13% of the participants discontinued the SECPT early, which

is comparable to the data presented by Schwabe and Schächinger (2018). Since those

participants in previous studies did not differ in their cortisol response from those who

persisted for the three minutes in the stress situation (Schwabe & Schächinger, 2018), they

were not excluded from further analyses.

Cortisol values were log-transformed to obtain a normal distribution. Additionally, we

performed a Greenhouse Geisser correction due to the violation of the assumption of

sphericity. Cortisol levels differed significantly at the three measurement time points (two

before SECPT and one after) (time: $F(2,214) = 41.433$, $p < .001$, $\eta^2_p = 0.279$, $BF = 1.076 \times 10^{13}$).

As expected, we observed a strong increase in cortisol after the stress manipulation (t_2)

compared to directly before (t_1) ($t(107) = -8.452$, $p_{Holm} < .001$, $d = -0.647$, $BF = 3.983 \times 10^8$) and

compared to the first measurement (t_0) ($t(107) = -7.154$, $p_{Holm} < .001$, $d = -0.548$, $BF =$

256993.046) . However, there was no significant difference between the NC and OC women

in cortisol levels (contraception: $F(1,107) = 0.201$, $p = .655$, $\eta^2_p = 0.002$, $BF = 0.231$) and no

significant interaction of time and contraception ($F(2,214) = 0.547$, $p = .580$, $\eta^2_p = 0.005$, $BF =$

0.108) (see Figure 2F).

Figure 2. (A/B) Approach Avoidance Task (AAT) - Bias Scores (log-transformed) for angry and happy faces for
women with no hormonal contraception (NC; red) and women taking oral contraceptives (OC; blue) before stress
induction through Socially Evaluated Cold Pressor Test (SECPT) **(A)** and after stress exposure **(B)**. **(C)** AAT – Bias

Scores (log-transformed) for angry faces across both AAT-runs for NC (red) and OC (blue) women. **(D)** AAT -
Reaction times (log-transformed) for each emotion (happy or angry), movement (push or pull) and mean reaction
times over all trials before and after stress for NC and OC women. **(E)** Error rates of runs of the AAT for each
participant in percent before and after stress. **(F)** Cortisol levels (log-transformed) at three measuring points for NC
and OC women. Error bars indicate standard error, asterisks indicate significant differences (p < 0.05).
400

3.3 Exploratory Analyses

Composition of preparations. OC preparations differed in composition with regard to the type of progestin and the dose of ethinyl estradiol (EE) they contain. The partial effects of the progestin can be distinguished into androgenic and anti-androgenic effects. To compare OC women with androgenic (N=27, mean(SD) progestin dose: 122(100)) and anti-androgenic (N=37, mean(SD) progestin dose: 2108 (315)) preparations, we conducted a mixed effects ANOVA (within factor: stress, emotion, movement; between factor: androgenicity). The results showed a significant four-way interaction of stress x emotion x movement x androgenicity ($F(1,62) = 6.380$, $p = .014$, $\eta^2_p = 0.093$). Separate ANOVAs for the emotions happy and angry showed that only for happy faces there was a significant three-way interaction of movement x stress x androgenicity ($F(1,26) = 6.502$, $p = .013$, $\eta^2_p = 0.095$, $BF = 1.056$). Using further separate ANOVAs for androgenic and anti-androgenic groups with factors stress and movement, we observed that only for women taking androgenic acting OC there was a significant interaction of stress and movement ($F(1,26) = 19.162$, $p < .001$, $\eta^2_p = 0424$, $BF = 269.14$). Post hoc tests showed that women taking androgenic acting OC had a stronger approach bias for happy faces after stress than before (before stress: ($t(26) = 2.726$, $p_{Holm} = 0.022$, $d = 0.371$, $BF = 2.501$); after stress: ($t(26) = 6.199$, $p_{Holm} < 0.001$, $d = 0.843$, $BF = 96212.845$)). This suggests that the stress effect on the approach bias in OC reported above was due in particular to the behavior in the group of women taking androgenic acting OCs. The overall response times, however, did not differ between the androgenic and anti-androgenic groups ($F(1,62) = 1.126$, $p = .293$, $\eta^2_p = 0.018$, $BF = 0.179$) (Figure 3A&B).

Most OC preparations used in our sample contained 30µg (EE30) (N=41, one excluded in analysis) or 20µg (EE20) (N=17) of ethinyl estradiol. Note that the ethinyl estradiol

dosage was not independent of the progestin type: all EE20 preparations were combined with

an androgenic acting progestin. The EE30 preparations were combined with either androgenic

(N=10) and anti-androgenic (N=30) acting progestins. When conducting a mixed effects

ANOVA (within factors: stress, emotion, movement; between factor: EE-dose), we again found

a fourfold interaction of stress x emotion x movement x EE-dose (F(1,55) = 4.267, p = .044,

η^2_p = 0.072). Again, this reflected a stronger approach bias to happy faces after stress in

women with a low EE dosage (and androgenic preparation) (interaction of stress and

movement: F(1,16) = 14.52, p = .002, η^2_p = 0.476, BF = 56.229; movement effect before stress:

t(16) = 1.0, p_{Holm} = 0.399, d = 0.207, BF = 0.347; movement effect after stress: t(16) = 4.213,

p_{Holm} < 0.001, d = 0.871, BF = 262.054).

The women did not differ in their cortisol response as a function of either androgenicity

or EE dose (no significant interaction of time and androgenicity/ EE dose: F(2,98) = 0.566, p

= 0.596, η^2_p = 0.011 / F(2,88) = 0.434, p = 0.649, η^2_p = 0.01).

Conditions of OC intake. Previous research suggests that certain conditions of OC

use, such as age at first use or whether OC were taken before or after the measurement, may

be important (Anderl et al., 2022; Gravelsins et al., 2021). In our sample, participants did not

differ in their approach and avoidance behaviors depending on whether they started taking

OC as adolescents (aged younger than 18 years (N=42)) or as adults (aged older than 18

years (N=22)) (no interaction of stress x emotion x movement x onset: F(1,62) = 0.997, p =

.322, η^2_p = 0.016), nor did the timing of daily intake (before or after measurement) make a

difference (no interaction of stress x emotion x movement x time of OC intake: F(1,62) =

2.6674, p = .107, η^2_p = 0.041). We were also interested in whether the duration of intake

correlated with the expression of the AAT effects, which was not the case (r = -0.047, p =.710,

Fisher´s z = 0.128, *BF* = 0.167).

Previous OC intake. The work of Pletzer and Kerschbaum (2014) suggested that

psychological effects of OC use may not be fully reversible. As some women in the NC group

had previously taken oral contraceptives (N=41), we tested this for our data using another

mixed effects ANOVA (within factors: stress, emotion, movement; between factor:

contraception). The factor contraception had three levels: OC, NC - previously taken OC (NCp,

N=41) and NC who had never taken OC (NCn; N=22). We found a significant main effect of

contraception on response times (F(2,124) = 4.954, p = .009, η^2_p = 0.074). The Post Hoc

analysis shows that only the NCn and OC women differed significantly in their response times

in the AAT (t(124) = 3.117, p = .007, d = 0.699), while NCp and OC women did not (t(124) =

1.402, p = .164, d = 0.255). However, there was no significant interaction of emotion x

movement x contraception (F(2,124) = 1.397, p = .251, η^2_p = 0.022)) (see Figure 3C).

Figure 3. (A/B) Approach Avoidance Task (AAT) - Bias Scores (log-transformed) for angry and happy faces for
women taking oral contraceptives (OC) with androgenic progestin (light green) and those with anti-androgenic
progestin (light grey) before stress induction through Socially Evaluated Cold Pressor Test (SECPT) **(A)** and after
stress exposure **(B)**. **(C)** Bias Scores (log-transformed) for angry and happy faces for women who never took OC
(NCn, red), who took OC previously but discontinued (NCp, pink) and OC women (blue) over both AAT-trials. Error
bars indicate standard error.
470

472   **3.4 Questionnaires**

473   The groups did not differ significantly from each other in their scores in the BIS or BAS, BFI-

474   10, or ERQ scales (see Table 1), suggesting that there were no general personality differences

between NC women and OC users. The ERQ subscales did not correlate with the AAT bias scores (all p > 0,05).

Table 1. Descriptive statistics for computerized measures (mean ± standard error in log(ms)) and questionnaire scores (mean ± standard error) and results of independent t-test comparisons for questionnaires.

	NC (n=63)	OC (n=64)				
AAT-Effect (pre stress)	0.137 ± 0.024	0.059 ± 0.024				
Bias Score Angry (pre stress)	0.036 ± 0.015	-0.015 ± 0.013				
Bias Score Happy (pre stress)	-0.101 ± 0.014	-0.074 ± 0.014				
AAT-Effect (post stress)	0.125 ± 0.023	0.119 ± 0.020				
Bias Score Angry (post stress)	0.041 ± 0.013	0.013 ± 0.012				
Bias Score Happy (post stress)	-0.084 ± 0.012	-0.106 ± 0.011				
	NC (n=65)	OC (n=65)	t(128)	p	d	BF_{10}
BIS	2.09 ± 0.063	2.06 ± 0.055	0.368	.713	0.065	0.199
BAS	1.87 ± 0.039	1.91 ± 0.038	-0.836	.405	-0.147	0.258
BFI-10 (openness)	3.76 ± 0.125	3.59 ± 0.123	0.965	.336	0.169	0.286
BFI-10 (neuroticism)	3.22 ± 0.110	3.45 ± 0.114	-1.460	.148	-0.255	0.490
BFI-10 (extraversion)	3.62 ± 0.112	3.62 ± 0.118	0.047	.962	0.008	0.188
BFI-10 (conscientiousness)	3.89 ± 0.100	3.77 ± 0.094	0.897	.371	0.157	0.270
BFI-10 (agreeableness)	3.71 ± 0.109	3.49 ± 0.113	1.371	.173	0.240	0.440
ERQ (cognitive reappraisal)	4.91 ± 0.122	4.66 ± 0.107	1.522	.130	0.267	0.535
ERQ (expressive suppression)	3.17 ± 0.149	3.24 ± 0.133	-0.347	.729	-0.061	0.198

4. Discussion

Oral contraceptives have an effect on central nervous levels of hormones, which in turn are

known to modulate affective behavior. In the current study, we investigated whether OC usage

is related to altered approach and avoidance behavior in a between-subject design. The

results of this study suggest that women using oral contraceptives (OC) have significantly

reduced avoidance tendencies to social threat signals compared to women not using oral

contraceptives (NC). OC women were as quick to pull angry faces towards them as they were

to push them away and thus showed no avoidance tendencies. Socially evaluated physical

stress led to a similar rise in cortisol levels in both groups, but only in the OC group did stress

increase approach behavior to positive social signals. It was also noteworthy that women

taking OC responded overall faster than the NC women. Additional exploratory analyses

suggested that the composition of the preparations may have an influence on the AAT results.

Only the OC women with an androgenic preparation showed a significant stress effect on

approach behavior. The composition of preparations was not related to the decreased

avoidance effect to threat signals in OC users, however. In sum, our study clarifies how

contraceptives can impact affect in women, and points toward promising avenues for future

research on the topic.

4.1 OC- related effects on approach and avoidance behavior

Group differences in approach and avoidance behavior were particularly evident in the

response to angry faces. Here, OC women differed significantly from NC women in their

avoidance behavior. Specifically, OC women pulled angry faces significantly faster towards

them than NC women did, but were comparably fast at pushing them away. Thus, it appears

that OC primarily facilitate approach toward threat signals. However, the reaction to angry

faces was not influenced by stress. For happy faces, it was the reverse. Here, there were no

general differences between NC and OC women, but OC women responded with stronger

approach tendencies to happy faces after stress than before.

528   These results expand on findings of Hamstra et al. (2014), who reported that women

taking oral contraceptives detected fewer facial expressions of anger than NC women, but did

not differ in recognizing happy faces. Weaker recognition of angry faces might manifest

primarily in a higher error rate in the AAT with respect to angry faces, which is not what we

found. However, OC users also here showed behavioral changes in the emotional response

to angry faces. This might be related to the previously reported OC-related reduction in

amygdala activation to negative stimuli (Petersen & Cahill, 2015), but as we did not measure

neural activity, this remains speculative. The AAT effects we observed were independent of

the exact OC preparation. This, and the fact that we did not select for specific OC preparations,

makes conclusions regarding the relative role of estradiol and progesterone difficult. Previous

research has suggested that endogenous estradiol is associated with increased activity in

emotion-regulatory brain areas such as the dorsolateral prefrontal cortex or the anterior

cingulate cortex (Chung et al., 2019; Sharma et al., 2021). For progesterone it is known that

it impairs emotion processing and recognition (Derntl et al., 2008; Guapo et al., 2009; van

Wingen et al., 2007). Based on these previous findings, the observed AAT effects might rather

be attributed to the OC effects on estradiol levels, but this needs to be investigated in future

studies.

OC women were also generally faster than NC women. This is similar to results of

another study, who also reported faster response times in OC users, this time to visual stimuli

without any emotional component (Grikšienė & Rukšėnas, 2009). Older studies, however,

reported generally increased reaction times in OC users (Garrett & Elder, 1984; Wuttke et al.,

1975). However, it is important to note that other formulations of oral contraceptives were used

at the time the studies were conducted. In a more recent study presenting emotional stimuli,

women taking OC were also faster than NC women, but only in response to sad faces and not

for happy or angry faces (Hamstra et al., 2014). Moreover there is some evidence that the

menstrual cycle phase has an influence on the reaction times in various paradigms (Li et al.,

2022; Pletzer et al., 2014). This indicates that different hormone levels may have contributed

to the observed reaction times differences, but there may be other influencing factors. It

remains for future studies to examine how reliable these response time differences are and

how they relate to previously reported cortical changes such as increased prefrontal activity

in OC users (Sharma et al., 2020).

In our study, previous OC usage was not associated with an altered approach

avoidance pattern compared to women who had never taken OC. Still, contrary to women

who never took OC, women who previously had used oral contraceptives did not differ

significantly in their reaction times from those using OC currently. Put otherwise, women who

took OC previously ranged in between women who were taking OC currently and those who

had never taken OC. Pletzer and Kerschbaum (2015) suggested that effects of oral

contraceptives could persist beyond the duration of intake. While they observed

neurostructural changes after OC use (relative hippocampal gray matter volume was found to

be associated positively with the duration of previous OC use), we found an effect on a

behavioral level. Although the effect was of medium size, it should be treated with caution

because of the unequal group sizes and the exploratory nature of the analysis. Nevertheless,

the results show that, in future studies, a differentiation between current and previous OC

users and nonusers might be useful.

4.2 OC-related effects in their interaction with stress

As hypothesized, stress influenced approach/avoidance behavior of OC and NC women in

different ways. Only the OC women showed altered AAT behavior after stress compared to

before. They reacted after stress with stronger approach tendencies to happy faces and

unaltered to angry faces. The increased approach bias resulted primarily from faster pull

reactions to happy faces. This could be interpreted as kind a of tend-and-befriend response,

characterized by responding to stress by aligning with social groups to ensure security (Taylor

et al., 2000). The influence of OC use on tend-and-befriend behavior is not well documented

so far. In contrast, one study revealed no significant difference between OC and NC women

in prosocial behavior under stress (von Dawans et al., 2019). However, the study did not

distinguish between OC women using androgenic and anti-androgenic preparations. In our

study, we noticed that women taking androgenic OC reacted with heightened approach to

happy faces after stress. There is some evidence that women using androgenic OC showed

increased stress effects on vasoconstriction compared to women using anti-androgenic OC

(Straneva et al., 2000), but there is no previous data on cognitive and affective changes after

stress. The tend-and-befriend hypothesis states that the affiliating behavior after stress is

mediated by oxytocin, which is also supported by oxytocin intervention studies (Cardoso et

al., 2013), and moderated by sex hormones (Taylor et al., 2000). Oxytocin was also found to

influence the approach avoidance behavior by normalizing the reaction tendencies towards

angry faces in a sample that showed no avoidance bias (Schneider et al., 2020). Similarly,

oxytocin might as well have influenced AAT scores after stress exposure in women using

androgenic OC. However, since this was not our main question and we did not measure

oxytocin levels, further research would be needed to explore the possible interaction of

androgenicity, stress and oxytocin. Remarkably, neither NC and OC women nor women taking

androgenic and antiandrogenic OC differed significantly in their cortisol stress response.

However, since cortisol is only one of many stress markers, the groups might still have differed

in other stress responses.

Overall, it is conceivable that altered approach and avoidance behavior with OC use

could be linked to the development of affective side effects. However, most OC women in our

study reported no affective side effects. AAT effects were also found to be unrelated to self-

reported emotion regulation in our sample. It could be that these relatively subtle behavioral

changes in OC users only result in affective side effects in women with higher vulnerability or

predisposition for depressive symptoms in the first place. Studies measuring pre-existing

hormonal levels as well as potential genetic markers are therefore warranted.

4.4 Limitations and Future Directions

Like many other studies investigating the influence of oral contraceptives on the female

organism, our study is also limited by the "survivor effect" (Kutner & Brown, 1972). This means

that our study only comprised women who did not stop taking OC because of serious side

effects, whereas those who did develop side effects abandoned the treatment early. Future

studies should implement a longitudinal, within-subject design to avoid pre-selection of

participants. The cross-sectional study design also presents the challenge that the effects

found are not uniquely attributable to OC use but may be influenced by other factors which

caused women to decide for or against OC. Nevertheless, the lack of differences in personality

and emotion regulation between OC and NC women speaks against this possibility.

In our AAT paradigm, we only presented faces with positive and negative valence. It

would be beneficial to add a neutral condition in future studies, which would allow investigating

whether the shortened reaction times of OC women are indeed due to the affective

component. Repeating the experiment with fMRI measurements could also shed light on the

neural patterns associated with these effects. In particular, the amygdala and aPFC could be

examined more closely to understand the neural basis of the observed behavioral changes

(Kaldewaij et al., 2017). Other stress markers, as well as hormonal measurements, may also

be relevant to the interpretation of future studies.

In general, the results of this and previous studies show that OC intake can lead to

significantly different outcomes in established paradigms such as the AAT. Thus, in future

work, OC intake, even if not the focus, should at least be measured and/or considered as an

influencing variable. Our results also indicate, once again, that in future study designs, the NC

group should be subdivided according to whether OC have been taken previously.

4.5 Conclusion

This work stands in line with a series of previous studies showing that oral contraceptives can

influence cognitive and affective processes (Gingnell et al., 2016; Monciunskaite et al., 2019;

Petersen & Cahill, 2015; Sharma et al., 2020). Specifically, we showed that OC use is

associated with differences in approach and avoidance behaviors. The markedly reduced

avoidance of angry faces while taking oral contraceptives indicates that affective action control

is altered by the hormonal changes that OC use involves. This has relevance primarily in that

altered affective action control may be an underlying cause of the affect-related side effects

reported by some OC users. Since discontinuation of OC use is often not straightforward when

side effects occur, it is important to understand the mechanisms behind this to enable

successful coping, as well as to identify those at higher risk.

CRediT author statement:

Jasmin Thurley: Methodology, Software, Formal analysis, Investigation, Writing – Original

Draft, Visualization **Macía Buades Rotger:** Methodology, Writing – Review and Editing **Georg**

Serfling: Methodology, Writing – Review and Editing **Thessa Howaldt:** Investigation, Writing

– Review and Editing **Nicole Reisch:** Writing – Review and Editing **Ulrike Krämer:**

Conceptualization, Methodology, Software, Resources, Writing – Review and Editing,

Supervision, Project administration

Acknowledgements:

We thank Carla Bücker, Annika Kühle and Luisa Ackermann for help during data acquisition.

This research did not receive any specific grant from funding agencies in the public,

commercial, or not-for-profit sectors.

References

Anderl, C., Wit, A. E., Giltay, E. J., Oldehinkel, A. J., & Chen, F. S. (2022). Association between
adolescent oral contraceptive use and future major depressive disorder: a prospective cohort
study. *Journal of Child Psychology and Psychiatry*, *63*(3), 333-341.
https://doi.org/10.1111/jcpp.13476
Beyer, F., Buades-Rotger, M., Claes, M., & Krämer, U. M. (2017). Hit or Run: Exploring Aggressive and
Avoidant Reactions to Interpersonal Provocation Using a Novel Fight-or-Escape Paradigm
(FOE). *Front Behav Neurosci*, *11*, 190. https://doi.org/10.3389/fnbeh.2017.00190
Brønnick, M. K., Økland, I., Graugaard, C., & Brønnick, K. K. (2020). The Effects of Hormonal
Contraceptives on the Brain: A Systematic Review of Neuroimaging Studies. *Front Psychol*,
11, 2813. https://doi.org/10.3389/fpsyg.2020.556577
Brown, T. (2020). *Investigating Chronic Pain in a Work Setting Using Electroencephalography and an*
Approach Avoidance Task [PhD Thesis, Tarleton State University]. Google Scholar.
Cardoso, C., Ellenbogen, M. A., Serravalle, L., & Linnen, A.-M. (2013). Stress-induced negative mood
moderates the relation between oxytocin administration and trust: evidence for the tend-and-
befriend response to stress? *Psychoneuroendocrinology*, *38*(11), 2800-2804.
https://doi.org/10.1016/j.psyneuen.2013.05.006
Carver, C. S., & White, T. L. (1994). Behavioral inhibition, behavioral activation, and affective responses
to impending reward and punishment: The BIS/BAS Scales. *Journal of Personality and Social*
Psychology, *67*, 319-333. https://doi.org/10.1037/0022-3514.67.2.319

Chen, M., & Bargh, J. A. (1999). Consequences of Automatic Evaluation: Immediate Behavioral
679   Predispositions to Approach or Avoid the Stimulus. *Personality and Social Psychology Bulletin*,
680   *25*(2), 215-224. https://doi.org/10.1177/0146167299025002007
681   Christin-Maitre, S. (2013). History of oral contraceptive drugs and their use worldwide. *Best Practice &*
682   *Research   Clinical   Endocrinology   &   Metabolism*,   *27*(1),   3-12.
683   https://doi.org/10.1016/j.beem.2012.11.004
684   Chung, Y. S., Poppe, A., Novotny, S., Epperson, C. N., Kober, H., Granger, D. A., Blumberg, H. P.,
685   Ochsner, K., Gross, J. J., Pearlson, G., & Stevens, M. C. (2019). A preliminary study of
686   association between adolescent estradiol level and dorsolateral prefrontal cortex activity during
687   emotion   regulation.   *Psychoneuroendocrinology*,   *109*,   104398.
688   https://doi.org/10.1016/j.psyneuen.2019.104398
689   Derntl, B., Windischberger, C., Robinson, S., Lamplmayr, E., Kryspin-Exner, I., Gur, R. C., Moser, E.,
690   & Habel, U. (2008). Facial emotion recognition and amygdala activation are associated with
691   menstrual   cycle   phase.   *Psychoneuroendocrinology*,   *33*(8),   1031-1040.
692   https://doi.org/10.1016/j.psyneuen.2008.04.014
693   Ernst, U., Baumgartner, L., Bauer, U., & Janssen, G. (2002). Improvement of quality of life in women
694   using a low-dose desogestrel-containing contraceptive: results of an observational clinical
695   evaluation. *The European Journal of Contraception & Reproductive Health Care*, *7*(4), 238-
696   243. https://doi.org/10.1080/ejc.7.4.238.243
697   Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power
698   analysis program for the social, behavioral, and biomedical sciences. *Behavior Research*
699   *Methods*, *39*(2), 175-191. https://doi.org/10.3758/BF03193146
700   Garrett, K. F., & Elder, S. T. (1984). The menstrual cycle from a bio-behavioral approach: A comparison
701   of oral contraceptive and non-contraceptive users. *International Journal of Psychophysiology*,
702   *1*(2), 209-214. https://doi.org/10.1016/0167-8760(84)90039-4
703   Gingnell, M., Bannbers, E., Engman, J., Frick, A., Moby, L., Wikström, J., & Sundström-Poromaa, I.
704   (2016). The effect of combined hormonal contraceptives use on brain reactivity during response
705   inhibition. *The European Journal of Contraception & Reproductive Health Care*, *21*(2), 150-157.
706   https://doi.org/10.3109/13625187.2015.1077381
707   Gingnell, M., Engman, J., Frick, A., Moby, L., Wikström, J., Fredrikson, M., & Sundström-Poromaa, I.
708   (2013). Oral contraceptive use changes brain activity and mood in women with previous
709   negative affect on the pill—A double-blinded, placebo-controlled randomized trial of a
710   levonorgestrel-containing combined oral contraceptive. *Psychoneuroendocrinology*, *38*(7),
711   1133-1144. https://doi.org/10.1016/j.psyneuen.2012.11.006
712   Gravelsins, L., Duncan, K., & Einstein, G. (2021). Do oral contraceptives affect young women's
713   memory? Dopamine-dependent working memory is influenced by COMT genotype, but not time
714   of pill ingestion. *PLoS One*, *16*(6), e0252807. https://doi.org/10.1371/journal.pone.0252807
715   Grikšienė, R., & Rukšėnas, O. (2009). Cognitive effects of hormone-based contraception in young
716   healthy women. *Biologija*, *55*(3-4).
717   Gross, J. J., & John, O. P. (2003). Individual differences in two emotion regulation processes:
718   implications for affect, relationships, and well-being. *Journal of Personality and Social*
719   *Psychology*, *85*(2), 348-362. https://doi.org/10.1037/0022-3514.85.2.348
720   Guapo, V. G., Graeff, F. G., Zani, A. C., Labate, C. M., dos Reis, R. M., & Del-Ben, C. M. (2009). Effects
721   of sex hormonal levels and phases of the menstrual cycle in the processing of emotional faces.
722   *Psychoneuroendocrinology*,   *34*(7),   1087-1094.
723   https://doi.org/10.1016/j.psyneuen.2009.02.007
724   Hampson, E. (2020). A brief guide to the menstrual cycle and oral contraceptive use for researchers in
725   behavioral   endocrinology.   *Hormones   and   Behavior*,   *119*,   104655.
726   https://doi.org/10.1016/j.yhbeh.2019.104655
727   Hamstra, D. A., De Rover, M., De Rijk, R. H., & Van der Does, W. (2014). Oral contraceptives may alter
728   the detection of emotions in facial expressions. *European Neuropsychopharmacology*, *24*(11),
729   1855–1859.
730   Hertel, J., König, J., Homuth, G., Van der Auwera, S., Wittfeld, K., Pietzner, M., Kacprowski, T., Pfeiffer,
731   L., Kretschmer, A., Waldenberger, M., Kastenmüller, G., Artati, A., Suhre, K., Adamski, J.,
732   Langner, S., Völker, U., Völzke, H., Nauck, M., Friedrich, N., & Grabe, H. J. (2017). Evidence
733   for Stress-like Alterations in the HPA-Axis in Women Taking Oral Contraceptives. *Scientific*
734   *Reports*, *7*(1), 14111. https://doi.org/10.1038/s41598-017-13927-7
735   Kaldewaij, R., Koch, S. B., Volman, I., Toni, I., & Roelofs, K. (2017). On the Control of Social Approach-
736   Avoidance Behavior: Neural and Endocrine Mechanisms. *Curr Top Behav Neurosci*, *30*, 275-
737   293. https://doi.org/10.1007/7854_2016_446

Kirschbaum, C., Kudielka, B. M., Gaab, J., Schommer, N. C., & Hellhammer, D. H. (1999). Impact of
Gender, Menstrual Cycle Phase, and Oral Contraceptives on the Activity of the Hypothalamus-
Pituitary-Adrenal   Axis: .   *Psychosomatic   Medicine*,   *61*(2),   154-162.
https://doi.org/10.1097/00006842-199903000-00006
Kutner, S. J., & Brown, W. L. (1972). Types of oral contraceptives, depression, and premenstrual
symptoms. *J Nerv Ment Dis*, *155*(3), 153-162. https://doi.org/10.1097/00005053-197209000-
00001
Langner, O., Dotsch, R., Bijlstra, G., Wigboldus, D. H. J., Hawk, S. T., & van Knippenberg, A. (2010).
Presentation and validation of the Radboud Faces Database. *Cognition and Emotion*, *24*(8),
1377-1388. https://doi.org/10.1080/02699930903485076
Lewis, C. A., Kimmig, A.-C. S., Zsido, R. G., Jank, A., Derntl, B., & Sacher, J. (2019). Effects of
Hormonal Contraceptives on Mood: A Focus on Emotion Recognition and Reactivity, Reward
Processing,   and   Stress   Response.   *Current   Psychiatry   Reports*,   *21*(11),   115.
https://doi.org/10.1007/s11920-019-1095-z
Li, D., Zhang, L., & Wang, X. (2022). The Effect of Menstrual Cycle Phases on Approach–Avoidance
Behaviors in Women: Evidence from Conscious and Unconscious Processes. *Brain Sciences*,
12(10), 1417. https://doi.org/10.3390/brainsci12101417
Love, J., Selker, R., Marsman, M., Jamil, T., Dropmann, D., Verhagen, J., Ly, A., Gronau, Q. F., Šmíra,
M., Epskamp, S., Matzke, D., Wild, A., Knight, P., Rouder, J. N., Morey, R. D., & Wagenmakers,
E.-J. (2019). JASP: Graphical Statistical Software for Common Statistical Designs. *Journal of
Statistical Software*, *88*, 1-17. https://doi.org/10.18637/jss.v088.i02
Meulenberg, P. M. M., & Hofman, J. A. (1990). The effect of oral contraceptive use and pregnancy on
the daily rhythm of cortisol and cortisone. *Clinica Chimica Acta*, *190*(3), 211-221.
https://doi.org/10.1016/0009-8981(90)90175-R
Monciunskaite, R., Malden, L., Lukstaite, I., Ruksenas, O., & Griksiene, R. (2019). Do oral
contraceptives modulate an ERP response to affective pictures? *Biol Psychol*, *148*, 107767.
https://doi.org/10.1016/j.biopsycho.2019.107767
Montoya, E. R., & Bos, P. A. (2017). How oral contraceptives impact social-emotional behavior and
brain function. *Trends Cogn Sci*, *21*(2), 125–136.
Mordecai, K. L., Rubin, L. H., Eatough, E., Sundermann, E., Drogos, L., Savarese, A., & Maki, P. M.
(2017). Cortisol reactivity and emotional memory after psychosocial stress in oral contraceptive
users. *Journal of neuroscience research*, *95*(1-2), 126–135.
Morimoto, M., Morita, N., Ozawa, H., Yokoyama, K., & Kawata, M. (1996). Distribution of glucocorticoid
receptor immunoreactivity and mRNA in the rat brain: an immunohistochemical and in situ
hybridization   study.   *Neurosci   Res*,   *26*(3),   235-269.   https://doi.org/10.1016/s0168-
0102(96)01105-4
Österlund, M., Kuiper, G. G., Gustafsson, J.-Å., & Hurd, Y. L. (1998). Differential distribution and
regulation of estrogen receptor-α and-β mRNA within the female rat brain. *Molecular Brain
Research*, *54*(1), 175–180.
Petersen, N., & Cahill, L. (2015). Amygdala reactivity to negative stimuli is influenced by oral
contraceptive   use.   *Soc   Cogn   Affect   Neurosci*,   *10*(9),   1266-1272.
https://doi.org/10.1093/scan/nsv010
Pletzer, B., Kronbichler, M., & Kerschbaum, H. (2015). Differential effects of androgenic and anti-
androgenic progestins on fusiform and frontal gray matter volume and face recognition
performance. *Brain Res*, *1596*, 108–115.
Pletzer, B., Kronbichler, M., Nuerk, H.-C., & Kerschbaum, H. (2014). Hormonal contraceptives
masculinize brain activation patterns in the absence of behavioral changes in two numerical
tasks. *Brain Res*, *1543*, 128-142. https://doi.org/10.1016/j.brainres.2013.11.007
Radke, S., Volman, I., Mehta, P., van Son, V., Enter, D., Sanfey, A., Toni, I., de Bruijn, E. R., & Roelofs,
K. (2015). Testosterone biases the amygdala toward social threat approach. *Science
advances*, *1*(5), e1400074. https://doi.org/10.1126/sciadv.1400074
Rammstedt, B., & John, O. P. (2007). Measuring personality in one minute or less: A 10-item short
version of the Big Five Inventory in English and German. *Journal of Research in Personality*,
41, 203-212. https://doi.org/10.1016/j.jrp.2006.02.001
Roelofs, K., Elzinga, B. M., & Rotteveel, M. (2005). The effects of stress-induced cortisol responses on
approach–avoidance   behavior.   *Psychoneuroendocrinology*,   *30*(7),   665-677.
https://doi.org/10.1016/j.psyneuen.2005.02.008
Roelofs, K., Putman, P., Schouten, S., Lange, W. G., Volman, I., & Rinck, M. (2010). Gaze direction
differentially affects avoidance tendencies to happy and angry faces in socially anxious
individuals. *Behav Res Ther*, *48*(4), 290-294. https://doi.org/10.1016/j.brat.2009.11.008

Schneider, I., Boll, S., Volman, I., Roelofs, K., Spohn, A., Herpertz, S. C., & Bertsch, K. (2020). Oxytocin
Normalizes Approach–Avoidance Behavior in Women With Borderline Personality Disorder.
Frontiers in Psychiatry, *11*, 120. https://doi.org/10.3389/fpsyt.2020.00120
Schwabe, L., & Schächinger, H. (2018). Ten years of research with the Socially Evaluated Cold Pressor
Test: Data from the past and guidelines for the future. *Psychoneuroendocrinology*, *92*, 155–
161. https://doi.org/10.1016/j.psyneuen.2018.03.010
Sharma, R., Cameron, A., Fang, Z., Ismail, N., & Smith, A. (2021). The regulatory roles of progesterone
and estradiol on emotion processing in women. *Cogn Affect Behav Neurosci*, *21*(5), 1026-1038.
https://doi.org/10.3758/s13415-021-00908-7
Sharma, R., Smith, S. A., Boukina, N., Dordari, A., Mistry, A., Taylor, B. C., Felix, N., Cameron, A.,
Fang, Z., Smith, A., & Ismail, N. (2020). Use of the birth control pill affects stress reactivity and
brain   structure   and   function.   *Hormones   and   Behavior*,   *124*,   104783.
https://doi.org/10.1016/j.yhbeh.2020.104783
Simerly, R. B., Swanson, L. W., Chang, C., & Muramatsu, M. (1990). Distribution of androgen and
estrogen receptor mRNA-containing cells in the rat brain: An in situ hybridization study. *Journal
of Comparative Neurology*, *294*(1), 76-95. https://doi.org/10.1002/cne.902940107
Straneva, P., Hinderliter, A., Wells, E., Lenahan, H., & Girdler, S. (2000). Smoking, oral contraceptives,
and   cardiovascular   reactivity   to   stress.   *Obstetrics   &   Gynecology*,   *95*(1),   78-83.
https://doi.org/10.1016/S0029-7844(99)00497-4
Taylor, S. E., Klein, L. C., Lewis, B. P., Gruenewald, T. L., Gurung, R. A., & Updegraff, J. A. (2000).
Biobehavioral responses to stress in females: tend-and-befriend, not fight-or-flight. *Psychol
Rev*, *107*(3), 411-429. https://doi.org/10.1037/0033-295x.107.3.411
Tronson, N. C., & Schuh, K. M. (2022). Hormonal contraceptives, stress, and the brain: The critical
need   for   animal   models.   *Frontiers   in   Neuroendocrinology*,   *67*,   101035.
https://doi.org/10.1016/j.yfrne.2022.101035
van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz, A., Evans, N. J.,
Gronau, Q. F., Haaf, J. M., Hinne, M., Kucharský, Š., Ly, A., Marsman, M., Matzke, D., Gupta,
A. R. K. N., Sarafoglou, A., Stefan, A., Voelkel, J. G., & Wagenmakers, E.-J. (2021). The JASP
guidelines for conducting and reporting a Bayesian analysis. *Psychonomic Bulletin & Review*,
28(3), 813-826. https://doi.org/10.3758/s13423-020-01798-5
van Wingen, G., van Broekhoven, F., Verkes, R. J., Petersson, K. M., Backstrom, T., Buitelaar, J., &
Fernandez, G. (2007). How progesterone impairs memory for biologically salient stimuli in
healthy   young   women.   *J   Neurosci*,   *27*(42),   11416-11423.
https://doi.org/10.1523/JNEUROSCI.1715-07.2007
Volman, I., Verhagen, L., Den Ouden, H. E. M., Fernández, G., Rijpkema, M., Franke, B., Toni, I., &
Roelofs, K. (2013). Reduced Serotonin Transporter Availability Decreases Prefrontal Control
of   the   Amygdala.   *The   Journal   of   Neuroscience*,   *33*(21),   8974-8979.
https://doi.org/10.1523/JNEUROSCI.5518-12.2013
von Borries, L. A. K., Volman, I., de Bruijn, E. R., Bulten, B. H., Verkes, R. J., & Roelofs, K. (2012).
Psychopaths lack the automatic avoidance of social threat: relation to instrumental aggression.
Psychiatry Res, *200*(2-3), 761-766. https://doi.org/10.1016/j.psychres.2012.06.026
von Dawans, B., Ditzen, B., Trueg, A., Fischbacher, U., & Heinrichs, M. (2019). Effects of acute stress
on   social   behavior   in   women.   *Psychoneuroendocrinology*,   *99*,   137-144.
https://doi.org/10.1016/j.psyneuen.2018.08.031
Wuttke, W., Arnold, P., Becker, D., Creutzfeldt, O., Langenstein, S., & Tirsch, W. (1975). Circulating
hormones, EEG, and performance in psychological tests of women with and without oral
contraceptives. *Psychoneuroendocrinology*, *1*(2), 141-152. https://doi.org/10.1016/0306-
4530(75)90006-2
846