1   **Reduced threat avoidance but increased stress induced approach bias in women**

2   **taking oral contraceptives**

4   Jasmin Thurley[1], Macià Buades-Rotger[2], Georg Serfling[3], Thessa Howaldt[1], Nicole Reisch[4],

5   Ulrike M. Krämer[5*]

[1] Dept. of Neurology, University of Lübeck, Germany

[2] Department of Clinical Psychology and Psychobiology, University of Barcelona, Barcelona,

Spain

[3] Medizinische Klinik I und Institut für Endokrinologie und Diabetes, Universitätsklinikum

Schleswig-Holstein, Campus Lübeck, Universität zu Lübeck, Lübeck, Germany

[4] Medizinische Klinik und Poliklinik IV, LMU Klinikum München, Munich, Germany

[5] Institute of Medical Psychology, Center of Brain, Behavior and Metabolism (CBBM),

University of Lübeck, Lübeck, Germany

* Corresponding author:

Ulrike M. Krämer, Ph.D.

Institute of Medical Psychology, University of Lübeck

Ratzeburger Allee 160

23538 Lübeck, Germany

+49 451 31017421

ulrike.kraemer@uni-luebeck.de

28   **Abstract**

Recent research has increasingly acknowledged the impact of oral contraceptives on affective

behavior and stress responses; however, the underlying mechanisms are still not well

understood. Studies have previously shown that steroid hormones modulate automatic

approach and avoidance behavior. Here, we thus investigated the effects of oral

contraceptives on approach and avoidance behavior and whether these effects are modulated

by stress. The study comprised 130 female participants, half of whom were using oral

contraceptives, while the other half were not using any hormonal contraception (NC). The

participants completed the Approach Avoidance Task (AAT), which measures automatic

approach and avoidance behavior to socio-affective signals. The AAT was run once before

and once after a stress manipulation using the Socially Evaluated Cold Pressor Test. OC

users showed absent avoidance behavior to social threat signals and a stress-induced

increase in approach behavior to positive social signals. The latter was found in particular in

women taking androgenic acting OC, demonstrating that different OC preparations need to be

taken into account in research on OC effects. However, OC and NC group did not differ in

their cortisol stress response. Overall, the results suggest that OC usage impacts on approach

and avoidance behavior to social signals, which might also contribute to the development of

affective side effects.

Keywords: oral contraceptives, approach avoidance behavior, stress

1. Introduction

Oral contraceptives (OC) are amongst the most reliable methods to prevent undesired pregnancies and are the contraceptive of choice for many women worldwide (Christin-Maitre, 2013). Hormonal contraception, however, is associated with side effects, including changes in affective experience and a possibly increased risk for depression (Ernst et al., 2002; Gingnell et al., 2013; Montoya & Bos, 2017). Prevalence estimates of adverse effects on mood and affect are between 4 – 10% (Poromaa & Segebladh, 2012) with a higher risk among adolescents and women with previous mood-related side-effects of hormonal contraceptives (Lewis et al., 2019; Tronson & Schuh, 2022). Many OC users however also report improved mood and emotional well-being or no changes at all (Poromaa & Segebladh, 2012; Tronson & Schuh, 2022). The impact of oral contraceptives on specific neural and psychological mechanisms, which cause these changes in mood and affect, remains unclear. OCs can influence brain functions through multiple mechanisms: directly through binding of the synthetic estrogens and progestins to endocrine receptors, indirectly through a reduction of endogenous levels of estrogens and progesterone, and, finally, indirectly through effects on neurotransmitter systems or other endocrine receptors (androgen, glucocorticoid or mineralocorticoid receptors) (Tronson & Schuh, 2022).

Altered approach and avoidance behavior may be a possible psychological mechanism for several reasons: the brain regions which have been implicated in approach and avoidance behavior to socio-affective signals and its regulation, include especially the amygdala and anterior prefrontal cortex (Kaldewaij et al., 2017). OC users have been found to show functional and structural changes in these and neighboring brain regions (Brønnick et al., 2020; Gingnell et al., 2013; Petersen & Cahill, 2015; Sharma et al., 2020), which might contribute to altered approach/avoidance behavior. Converging evidence (Rehbein et al., 2021) suggests that amygdala activity to affective stimuli is modulated by estradiol (van Wingen et al., 2011; Zeidan et al., 2011), progesterone (van Wingen et al., 2007; van Wingen et al., 2008) and testosterone (Buades-Rotger et al., 2016; van Wingen et al., 2010). Moreover, the steroid hormones testosterone and cortisol, possibly modulated by OC use,

appear to have an impact on approach and avoidance behaviors (Enter et al., 2014; Radke et

al., 2015; Roelofs et al., 2005). Lastly, considering that OC influence the stress axis and

cortisol reactivity (Mordecai et al., 2017), which in turn affects approach and avoidance

behavior (Roelofs et al., 2005), it is possible that OC influence affective behavior in conjunction

with stress. Here, we thus tested whether women who are taking OC show altered approach

and avoidance behavior, and whether this effect is modulated by stress using a cross-

sectional design.

Approach/avoidance behavior refers to the automatic tendency to approach positive

stimuli and avoid negative stimuli, which can be measured using the Approach Avoidance

Task (AAT) (Chen & Bargh, 1999). In one commonly used variant of the AAT (Kaldewaij et

al., 2017), participants are asked to pull pictures of happy faces towards them and push

negative faces away as fast as possible using a joystick. Response times in this congruent

condition are compared with those of the reverse, incongruent condition (pull angry faces and

push happy faces away) (Beyer et al., 2017). Typically, participants are faster in the congruent

compared to the incongruent condition, referred to as congruency effect or AAT bias. Previous

research with the AAT reported that people with social anxiety show faster avoidance (Roelofs

et al., 2010), whereas psychopathic patients showed reduced avoidance to angry faces (von

Borries et al., 2012), speaking for the validity of the AAT. A neurobiological model of the AAT

(Volman et al., 2013) assumed that faces are first processed in the fusiform face area (FFA)

and subsequently projected to the amygdala, which has a bidirectional connection to the

anterior prefrontal cortex (aPFC). When automatic affective behavior needs to be controlled

in the incongruent condition, amygdala activity is believed to be inhibited by the aPFC (Volman

et al., 2013).

Oral contraceptive use has previously been found to be associated with changes in

reactivity and activation of the PFC and amygdala. Women using OCs responded to negative

emotional stimuli with reduced amygdala reactivity (Petersen & Cahill, 2015) and increased

prefrontal activation (Sharma et al., 2020). Rodent studies suggest that estradiol treatment

influences estrogen receptor density in the amygdala, whereby the direction of effects seems

to depend on the amygdala nucleus and estrogen receptor subtype (ERα or ERβ) (Österlund

et al., 1998). Estradiol has been shown to facilitate fear extinction learning through its effects

in the amygdala and prefrontal cortex (Zeidan et al., 2011). Reduced circulating estradiol

levels due to hormonal contraception in turn impaired extinction learning (Graham & Milad,

2013; Taxier et al., 2020). Although these findings cannot directly be translated to

approach/avoidance behavior to social signals, they speak for reduced reactivity to threat

signals in OC users. Moreover, Li and colleagues (Li et al., 2022) reported that in a sample of

women not taking hormonal contraceptives, progesterone and estradiol levels were correlated

with approach and avoidance behavior. Specifically, higher progesterone levels were related

to faster avoidance and higher estradiol levels were related to faster approach behavior (Li et

al., 2022). This would suggest reduced avoidance and approach behavior with the reduction

of endogenous progesterone and estradiol in women taking OCs.

In addition to the contraceptive effect OC have on the HPG-axis (hypothalamic–

pituitary–gonadal axis), they also affect the HPA stress axis (hypothalamic-pituitary-adrenal

axis). One often replicated effect of OC use (Lewis et al., 2019; Tronson & Schuh, 2022) is a

blunted cortisol stress response, which was first reported by Kirschbaum and colleagues

(Kirschbaum et al., 1999; Kirschbaum et al., 1995). Additionally, several studies reported an

increase of basal cortisol levels in OC users compared to women not taking hormonal

contraceptives (Tronson & Schuh, 2022). This pattern of elevated basal cortisol levels together

with a blunted cortisol stress response is also characteristic of depression as well as other

psychiatric disorders (Carroll et al., 2017). However, in a large-scale study on OC-related

changes in the stress axis, no association was found between basal plasma cortisol levels

and depressive symptoms in women using OCs (Hertel et al., 2017), calling basal cortisol

levels into question as a mediating factor between OC use and depression. Given the close

interrelation between stress and motivation (Slattery & Cryan, 2017), it is however likely that

a blunted stress response in OC users is associated with altered motivation and behavior.

There is only few research on stress effects on the AAT, but this work suggests that a high

cortisol increase due to social stress is associated with a reduced AAT congruency effect

(Roelofs et al., 2005). As OC might also influence approach/avoidance behavior through this

pathway, we tested whether approach/avoidance behavior in OC users interacts with stress

exposure.

One challenge in research on OC is the large heterogeneity of preparations taken by

women (Tronson & Schuh, 2022). In our study, only women taking combined preparations

containing an estrogen and a progestin were included, but the preparations nevertheless

differed in the included estradiol dose and in the used anti- or androgenic acting progestin.

The different formulations could have an influence on AAT and stress effects via the different

efficacy profiles, as there is for example evidence that women with androgenic OC have a

more pronounced cardiovascular stress response (Straneva et al., 2000). Therefore, we

performed exploratory analyses on the subgroups.

2. Methods

2.1 Participants

We tested one hundred thirty women, of whom sixty-five had been using oral contraceptives

for at least three months (oral contraceptives, OC group) and 65 had not used any hormonal

contraception for at least six months (no hormonal contraception, NC group). The mean age

in the OC group was 21.9 years (SD = 2.13) and in the NC group 22.6 years (SD = 2.4). They

had all normal or corrected to normal vision (self-report). An analysis with G*Power 3.1.9.6

(Faul et al., 2007) suggested that at least 128 participants would be required to find an effect

of medium effect size for group differences in the AAT effect (f = 0.25) with a power of 0.8 and

an alpha error of 0.05. All participants were between 19 and 30 years old, had a BMI between

17.5 and 25 and no neurological, psychiatric, cardiovascular or endocrinological pre-existing

diseases.

Of the 130 participants, we had to exclude three for the following reasons: one due to

many errors in the AAT (mean error rate: 39.15%) and two due to reaction times in the AAT

that were more than 3 SD above the mean. Saliva samples from 20 participants could not be

used, resulting in a final sample of 106 participants (50=OC, 56=NC) for the cortisol analyses.

All of the women in the OC group were using combined oral contraceptives containing both

estradiol and progestin. The doses of ethinyl estradiol varied between 20 and 30 µg. Three

women were using OC with estetrol (14.2 mg) as estradiol and 4 women took OC containing

estradiol valerate (1-3 mg). The doses of progestin ranged between 100 and 3000 µg (see

Supplementary Table 1). The participants were recruited via mailing lists of the University of

Lübeck. For the verification of the inclusion and exclusion criteria, the potential participants

were subsequently called. The person who conducted the calls was not involved in the

measurements and the investigators were blinded to the group membership of the respective

participants. All participants were informed about the study objectives and compensated for

their participation with 10 Euros per hour or credit points. All participants gave their written

informed consent in accordance with the Declaration of Helsinki. The Ethics Committee of the

University of Lübeck approved this study protocol (22-044).

2.2 Design and Procedure

OC women were tested between the first and third day of pill intake, NC women

correspondingly on the eighth to tenth day of the menstrual cycle (follicular phase). The timing

was chosen because interpersonal variations in cycle length (in the NC group) and associated

differences in hormone concentrations are lowest at the beginning of the cycle (Hampson,

2020) and to measure women taking OC already during the active phase of pill intake. All

measurements took place in the afternoon between 2 and 7 p.m., since during this period

salivary cortisol concentrations are relatively consistently low for both women taking oral

contraceptives and women with no hormonal contraception (Meulenberg & Hofman, 1990).

189   After participants gave their informed consent, they were asked to fill out some

computerized questionnaires assessing emotion regulation, personality, and

approach/avoidance tendencies (see 2.6 Questionnaires and Demographic Data). In the next

step, they provided a saliva sample (t_0) before performing the AAT on the computer (Figure

1). After another saliva sample (t_1), the Socially evaluated cold pressor test (SECPT)

(Schwabe & Schächinger, 2018) was performed. After a break of approximately 12 minutes,

the AAT was performed a second time. After completion of the second AAT, another saliva

sample (t_2) was collected and the participants filled in a questionnaire with demographic data

and information about the pill. Lastly, participants were debriefed about the research question

and were compensated. The testing lasted approximately one hour in total.

2.3 Approach Avoidance Task (AAT)

We used the Approach Avoidance Task (Chen & Bargh, 1999) in the version of Beyer et al.

(2017) with images of happy and angry faces from the Radboud Faces database (Langner et

al., 2010) as stimuli. Stimuli were presented using Presentation® (Neurobehavioral Systems).

In total, images of thirty individuals (15 male, 15 female) were shown during the AAT. There

was one picture of each person with an angry and a happy facial expression. For the practice

blocks, pictures of nine different people than in the main block were used. All pictures were

cut into an oval shape so that hair, neck and ears are not visible.

Participants used the joystick with their dominant hand. Before each stimulus, a small

black cross appeared on the white screen. As soon as the "shoot" button of the joystick was

pressed, the cross disappeared and a face appeared in the center of the screen. By pushing

the joystick away (avoidance response), the image got smaller step-by-step (seven steps in

total). Pulling the joystick towards the body (approach response) made the image larger. Two

conditions were run in two different blocks. In the congruent condition, participants were asked

to pull happy faces toward them as quickly as possible and to push angry faces away from

them. In the incongruent condition, the instruction was reversed. Block order was kept identical

across participants and stress conditions, starting always with the congruent condition. Here,

happy faces were to be pushed away from the body and angry ones were to be pulled in. Each

block contained thirty happy faces and thirty angry faces, using the same images in both

blocks.

Before each block, the participants could familiarize themselves with the task in a

practice session. Here, the participants received direct feedback whether they had reacted

correctly (green tick for correct reaction and red cross for incorrect reaction). The first exercise

run consisted of twenty images. In the second practice run, 28 pictures were used because

the instruction that was internalized in the first block was reversed and task-switching effects

were to be avoided.

During each trial the reaction time was measured, which is determined as the interval

between the appearance of the stimulus and the start of the movement of the joystick.

2.4 Socially Evaluated Cold Pressor Test (SECPT)

The SECPT was performed as suggested by Schwabe and Schächinger (2018). The

participants were instructed to hold their dominant hand, including the wrist, in a bucket of ice

water (0-2 degrees Celsius) until the experimenter asked them to take the hand out again

(they were not told in advance how long they should keep the hand in the ice water). Only if

they could no longer stand the cold at all, they were allowed to take their hand out of the water

earlier. They were also told that they were videotaped during the experiment so that their facial

expressions can be analyzed later. Therefore, they were instructed to look into the camera

during the entire experiment and not to speak. The camera was placed about two meters in

front of the participants behind a screen facing them. The camera recorded the subject's face

and transferred it directly to the screen so that the participant could see herself. Next to the

camera, the experimenter stood at a distance that allowed the participant to simultaneously

look into the camera and see the experimenter out of the corner of her eye. The experimenter

made sure that the participant did not make a fist and that her hand was in ice water up to the

wrist. At the same time, she took notes and avoided giving confirming signals such as smiling

or nodding. After three minutes, the test ended and the participant was allowed to take her

hand out of the water. If she took her hand out before that, the experimenter responded by

asking the participant to put her hand back in the water. If this was not possible, the participant

was asked to remain standing and looking into the camera for the remainder of the three

minutes.

2.5 Saliva Sampling

Saliva samples were collected at three time points of the measurement. For this purpose, we

used Sarstedt cortisol Salivette® with a synthetic fiber roll. To prevent cortisol levels from

being influenced by factors other than the experimental factors, participants were instructed

not to exercise, eat anything, drink coffee, take medications, or smoke for one hour before the

appointment (Roelofs et al., 2005). Saliva samples were frozen and stored at -20°C until

analysis. After thawing, samples were centrifuged at 3,000 rpm for 5 min, which resulted in a

clear supernatant of low viscosity. Salivary concentrations were measured using commercially

available chemiluminescence immunoassay with high sensitivity (Tecan - IBL International,

Hamburg, Germany; catalogue number R62111). The intra and interassay coefficients of

variance were below 9%. Samples were considered empty if there was no saliva in the tube

after centrifugation (samples from n=20 participants, resulting in a sample of 106 participants

(50=OC, 56=NC) for the cortisol analyses).

2.6 Questionnaires and Demographic Data

We used the Behavioral Inhibition Scale (BIS) and the Behavioral Approach Scale (BAS)

designed by Carver and White (1994) to assess tendencies to exhibit avoidance or approach

behaviors. We also used the Emotion Regulation Questionnaire (ERQ) (Gross & John, 2003)

and the Big Five Inventory (BFI-10) (Rammstedt & John, 2007) to examine interpersonal

differences in emotion regulation and personality structure.

270   Participants also provided information on the name of the OC preparation taken,

previous duration of use and age at first use of oral contraceptives, time of daily use, reason

for use, and any side effects (OC group). A summary of the frequency of reported side effects

in our sample is provided in Supplementary Table 2. The NC-women were asked if they had

used hormonal contraception in the past. If they had taken OC in the past, the duration and

the time since the last intake was recorded. The reason for taking the pill and the occurrence

of any side effects were asked as well. The weight, height and age of all test participants was

recorded.

A Procedure

B AAT   ## C SECPT

Figure 1. A Procedure. At first, the participants completed the questionnaires ERQ, BFI-10, and BIS/BAS. Next, they provided a saliva sample and performed the AAT (Approach Avoidance Task). After another saliva sample, they performed the Socially Evaluated Cold Pressor Test (SECPT) and another run of the AAT. 25 minutes after the onset of the SECPT, the participants gave another saliva sample and finally provided demographic information. **B** Approach Avoidance Task (AAT): Participants were asked to push away angry faces and pull happy faces as fast as possible in the congruent condition and do it the other way round in the incongruent condition. Note that drawings are shown for visualization only. In the experiment, we showed images of happy and angry faces, which were taken from the Radboud Faces database (Langner et al., 2010). **C** Socially evaluated cold pressor test (SECPT). Participants were required to keep their dominant hand in ice water for 3 minutes while being observed by the experimenter and being recorded on video.

2.7 Data analysis

The main outcome of the AAT were the reaction times (RT) which were defined as the interval between stimulus presentation and movement onset. Following previous work (Beyer et al., 2017), we excluded trials with response latencies of less than 150 ms or more than 3 SD from the individual participant's own mean. Moreover, we only considered correct trials. Incorrect trials (errors), which also included trials with an initial movement into the wrong direction and a following correction were excluded. We first log-transformed the reaction times (ms) before computing mean response times for each condition (push, pull for angry and happy) and

participant, as reaction times were not normally distributed. We also considered the error rates

for determination of outliers. Participants (total n=3) were excluded if reaction times (n=2) or

error rates (n=1) in the AAT were more than 3 SD above the mean across all participants.

We computed RT means, outliers and error rates using MATLAB (MATLAB, 2021).

Statistical analyses were conducted in Jamovi (The jamovi project, 2021) and JASP (JASP

Team, 2023).

We performed a mixed effects ANOVA that included the factors emotion (happy,

angry), movement (push, pull), stress (within-subject factors) and contraception (between-

subject factor) using log-transformed reaction times. To parse interaction effects, we

conducted further ANOVAs to examine group differences between OC and NC women

separately for happy and angry faces and before and after stress. To determine the AAT effect,

the reaction times of the movement away from the body (push) were subtracted from the

reaction times of the movement towards the body (pull). A higher (more positive) score here

reflects stronger avoidance behavior and a lower (more negative) score reflects approach

behavior. The AAT effect was calculated as the difference between the bias score for angry

faces and the bias score for happy faces. Accordingly, the higher the bias scores, the larger

the effect. The AAT effect was reported only for visualization and to allow for comparability of

results with previous studies and is not interpreted on its own in this study. We conducted an

additional mixed effects ANOVA with factors stress and contraception to compare the error

rates of OC and NC women before and after stress.

To investigate the change in cortisol levels, we performed another mixed effects

ANOVA. Here, the within-subject factor time (t_0, t_1, t_2) and the between-subject factor

contraception were included. For the comparison of group differences in the questionnaires,

we calculated two-sided independent samples t-tests.

For further exploratory analyses of subgroups within the OC group, we ran additional

mixed effects ANOVAs with the factors emotion, movement, stress (within-subject factors) and

the respective between-subject factors (androgenicity, EE-dose, onset, time of OC intake or

contraception). Moreover, we ran a correlation of duration of OC intake and the AAT effect

and a correlation of the AAT effect with the Emotion Regulation Questionnaire subscales

(across the whole sample).

All results with a p-value less than .05 were considered significant. For significant

ANOVA results, we used Bonferroni-Holm adjusted p-values for post-hoc tests. In order to

make informed statements regarding the evidence, we also provide the Bayes Factor, which

tells us how much more probable the observations are under the alternative hypothesis

compared to the null hypothesis. Our Bayesian analysis was conducted using the default

priors in JASP, and we present the findings of Bayesian model averaging (BF_{incl}) (Love et al.,

2019). To interpret the results, we adhere to established guidelines: BF values exceeding 3

indicate moderate support, while values above 10 indicate strong support for the alternative

hypothesis. Conversely, BF values below 0.1 strongly support the null hypothesis, and values

below 0.33 suggest moderate support (van Doorn et al., 2021). Because Bayesian analyses

are time-consuming and computationally expensive for large-sample multifactorial ANOVAs,

we report Bayes factors exclusively for specific lower-order effects.

Data is available at https://osf.io/kz4yp/?view_only=167cd3c80d064916ae0edcd82f02b5ac.

3. Results

3.1 Approach Avoidance Behavior

The AAT showed the expected experimental effects, as participants showed avoidance of

angry faces and approach toward happy faces. Visual examination of the behavioral data

suggested, however, that OC women responded less avoidant overall and even showed a

slight tendency to approach angry faces before the SECPT (see Table 1 and Figure 2A&B).

Note that the reaction times were log-transformed to increase normality.

Results of the mixed ANOVA showed that, as expected, the responses to the emotions

in the congruent and incongruent condition differed in speed (see Figure 2A-D). Overall, it took

the participants longer to pull angry faces towards them than to push them away, whereas

they showed the opposite pattern for happy faces (emotion x movement: $F(1,125) = 66.733$,

$p < .001$, $\eta^2_p = 0.348$), reflecting the expected AAT effect. In general, the groups did not differ

in the expression of this pattern (emotion x movement x contraception: $F(1,125) = 2.482$, $p =$

.118, $\eta^2_p = 0.019$). However, there was a significant fourfold interaction of stress x emotion x

movement x contraception ($F(1,125) = 4.345$, $p = .039$, $\eta^2_p = 0.034$). Similar to previous studies

with the AAT (Radke et al., 2015; Volman et al., 2013), participants were faster in pull than

push movements (movement: $F(1,125) = 95.735$, $p < .001$, $\eta^2_p = 0.434$). Interestingly, OC

women responded overall faster than NC women (contraception: $F(1,125) = 6.35$, $p = .013$,

$\eta^2_p = 0.048$) (see Figure 2D). Women in the OC group were especially faster than NC women

in their pull movements, and less so in their push movements (interaction of movement x

contraception: $F(1,125) = 6.019$, $p = .016$, $\eta^2_p = 0.046$; pull: $t(125) = 2.827$, $p_{Holm} = .016$, $d =$

0.461; push: $t(125) = 2.166$, $p_{Holm} = .064$, $d = 0.353$).

To understand the four-way interaction of stress x emotion x movement x

contraception, we computed separate mixed effects ANOVAs for the two emotions (happy

and angry), since we expected differences especially for angry faces. In response to angry

faces, the groups indeed differed in their behavior, reflected in a significant interaction of

movement x contraception ($F(1,125) = 6.708$, $p = .011$, $\eta^2_p = 0.051$, $BF = 3.916$). The Bayes

factor indicated moderate support for this interaction. While NC women pushed away angry

faces faster than they pulled them closer ($p_{Holm} = 0.003$), showing the typical avoidance bias,

OC women responded equally fast when pushing and pulling ($p_{Holm} = 0.937$; see Figure 2C).

This pattern was not influenced by stress (stress x movement x contraception: $F(1,125) =$

1.155, p = .285, $\eta^2_p = 0.009$, $BF = 0.305$). The groups did not differ in their general behavior

to happy faces (movement x contraception: $F(1,125) = 0.039$, $p = .843$, $\eta^2_p < 0.001$, $BF =$

0.216), however, there was a threefold interaction of stress x movement x contraception for

happy faces ($F(1,125) = 6.596$, $p = .011$, $\eta^2_p = 0.050$, $BF = 3.073$). To understand this

interaction, we computed separate mixed effects ANOVAs for the two groups (NC and OC).

While stress had no influence on approach behavior to happy faces in NC women (stress x

movement: $F(1,62) = 1.56$, $p = .216$, $\eta^2_p = 0.025$, $BF = 1.419$), OC women differed in their

reactions towards happy faces before and after stress (stress x movement: $F(1,63) = 5.68$, p

= .02, η^2_p = 0.083, *BF* = 9.59). This was due to more pronounced approach behavior after

stress than before stress (movement after stress: *F*(1,63) = 87.2, *p* < .001, η^2_p = 0.580, *BF* =

6069.494; movement before stress: *F*(1,63) = 27.1, *p* < .001, η^2_p = 0.301, *BF* = 2.379x10^{10})

(see Figure 2A&B). Note however, that OC women showed a clear approach bias to happy

faces both before and after stress.

Regarding error rates in the AAT, there was no group difference before or after stress

(mixed effects ANOVA with factors stress and contraception; contraception: *F*(1,125) = 1.05,

p = 0.308, η^2_p = 0.008, *BF* = 0.257) (Figure 2E).

3.2 Stress effects on cortisol level

The stress manipulation generally succeeded, as cortisol levels were higher on average after

the SECPT than before. About 13% of the participants discontinued the SECPT early, which

is comparable to the data presented by Schwabe and Schächinger (2018). Since those

participants in previous studies did not differ in their cortisol response from those who

persisted for the three minutes in the stress situation (Schwabe & Schächinger, 2018), they

were not excluded from further analyses.

Cortisol values were log-transformed to obtain a normal distribution. Additionally, we

performed a Greenhouse Geisser correction due to the violation of the assumption of

sphericity. Cortisol levels differed significantly at the three measurement time points (two

before SECPT and one after) (time: *F*(2,214) = 41.433, *p* < .001, η^2_p = 0.279, *BF* = 1.076x10^{13}).

As expected, we observed a strong increase in cortisol after the stress manipulation (t_2)

compared to directly before (t_1) (*t*(107) = -8.452, p_{Holm} < .001, *d* = -0.647, BF = 3.983x10^8) and

compared to the first measurement (t_0) (*t*(107) = -7.154, p_{Holm} < .001, *d* = -0.548, BF =

256993.046) . However, there was no significant difference between the NC and OC women

in cortisol levels (contraception: *F*(1,107) = 0.201, *p* = .655, η^2_p = 0.002, *BF* = 0.231) and no

significant interaction of time and contraception (*F*(2,214) = 0.547, *p* = .580, η^2_p = 0.005, *BF* =

0.108) (see Figure 2F). We also computed the area-under-the-curve values AUC-G and AUC-

I as suggested by Pruessner and colleagues (Pruessner et al., 2003), but these values did not

differ between the groups either (both t < 1 and p > 0.5).

Figure 2. (A/B) Approach Avoidance Task (AAT) - Bias Scores (log-transformed) for angry and happy faces for
women with no hormonal contraception (NC; red) and women taking oral contraceptives (OC; blue) before stress
induction through Socially Evaluated Cold Pressor Test (SECPT) **(A)** and after stress exposure **(B). (C)** AAT – Bias
Scores (log-transformed) for angry faces across both AAT-runs for NC (red) and OC (blue) women. **(D)** AAT -
Reaction times (log-transformed) for each emotion (happy or angry), movement (push or pull) and mean reaction
times over all trials before and after stress for NC and OC women. **(E)** Error rates of runs of the AAT for each
participant in percent before and after stress. **(F)** Cortisol levels (log-transformed) at three measuring points for NC
and OC women. Error bars indicate standard error, asterisks indicate significant differences (p < 0.05).

3.3 Exploratory Analyses

Composition of preparations. OC preparations differed in composition with regard to the type

of progestin and the dose of ethinyl estradiol (EE) they contain. The partial effects of the

progestin can be distinguished into androgenic and anti-androgenic effects. To compare OC

women with androgenic (N=27, mean(SD) progestin dose: 122(100)) and anti-androgenic

(N=37, mean(SD) progestin dose: 2108 (315)) preparations, we conducted a mixed effects

ANOVA (within factor: stress, emotion, movement; between factor: androgenicity). The results

showed a significant four-way interaction of stress x emotion x movement x androgenicity

(F(1,62) = 6.380, p = .014, η^2_p = 0.093). Separate ANOVAs for the emotions happy and angry

showed that only for happy faces there was a significant three-way interaction of movement x

stress x androgenicity (F(1,26) = 6.502, p = .013, η^2_p = 0.095, BF = 1.056). Using further

separate ANOVAs for androgenic and anti-androgenic groups with factors stress and

movement, we observed that only for women taking androgenic acting OC there was a

significant interaction of stress and movement (F(1,26) = 19.162, p < .001, η^2_p = 0424, BF =

269.14). Post hoc tests showed that women taking androgenic acting OC had a stronger

approach bias for happy faces after stress than before (before stress: (t(26) = 2.726, p_{Holm} =

0.022, d = 0.371, BF = 2.501); after stress: (t(26) = 6.199, p_{Holm} < 0.001, d = 0.843, BF =

96212.845)). This suggests that the stress effect on the approach bias in OC reported above

was due in particular to the behavior in the group of women taking androgenic acting OCs.

The overall response times, however, did not differ between the androgenic and anti-

androgenic groups (F(1,62) = 1.126, p = .293, η^2_p = 0.018, BF = 0.179) (Figure 3A&B).

Most OC preparations used in our sample contained 30µg (EE30) (N=41, one

excluded in analysis) or 20µg (EE20) (N=17) of ethinyl estradiol. Note that the ethinyl estradiol

dosage was not independent of the progestin type: all EE20 preparations were combined with

an androgenic acting progestin. The EE30 preparations were combined with either androgenic

(N=10) and anti-androgenic (N=30) acting progestins. When conducting a mixed effects

ANOVA (within factors: stress, emotion, movement; between factor: EE-dose), we again found

a fourfold interaction of stress x emotion x movement x EE-dose (F(1,55) = 4.267, p = .044,

η^2_p = 0.072). Again, this reflected a stronger approach bias to happy faces after stress in

women with a low EE dosage (and androgenic preparation) (interaction of stress and

movement: F(1,16) = 14.52, p = .002, η^2_p = 0.476, BF = 56.229; movement effect before stress:

$t(16) = 1.0$, $p_{Holm} = 0.399$, $d = 0.207$, $BF = 0.347$; movement effect after stress: $t(16) = 4.213$,

$p_{Holm} < 0.001$, $d = 0.871$, $BF = 262.054$).

The women did not differ in their cortisol response as a function of either androgenicity

or EE dose (no significant interaction of time and androgenicity/ EE dose: F(2,98) = 0.566, p

= 0.596, $\eta^2_p = 0.011$ / F(2,88) = 0.434, p = 0.649, $\eta^2_p = 0.01$).

Conditions of OC intake. Previous research suggests that certain conditions of OC

use, such as age at first use or whether OC were taken before or after the measurement, may

be important (Anderl et al., 2022; Gravelsins et al., 2021). In our sample, participants did not

differ in their approach and avoidance behaviors depending on whether they started taking

OC as adolescents (aged younger than 18 years (N=42)) or as adults (aged older than 18

years (N=22)) (no interaction of stress x emotion x movement x onset: $F(1,62) = 0.997$, $p =$

.322, $\eta^2_p = 0.016$), nor did the timing of daily intake (before or after measurement) make a

difference (no interaction of stress x emotion x movement x time of OC intake: $F(1,62) =$

2.6674, $p = .107$, $\eta^2_p = 0.041$). We were also interested in whether the duration of intake

correlated with the expression of the AAT effects, which was not the case ($r = -0.047$, $p = .710$,

Fisher´s z = 0.128, *BF* = 0.167).

Previous OC intake. The work of Pletzer and Kerschbaum (2014) suggested that

psychological effects of OC use may not be fully reversible. As some women in the NC group

had previously taken oral contraceptives (N=41), we tested this for our data using another

mixed effects ANOVA (within factors: stress, emotion, movement; between factor:

contraception). The factor contraception had three levels: OC, NC - previously taken OC (NCp,

N=41) and NC who had never taken OC (NCn; N=22). We found a significant main effect of

contraception on response times ($F(2,124) = 4.954$, $p = .009$, $\eta^2_p = 0.074$). The Post Hoc

analysis shows that only the NCn women were slower than OC women in the AAT ($t(124) =$

3.117, $p = .007$, $d = 0.699$), while NCp and OC women did not differ ($t(124) = 1.402$, $p = .164$,

$d = 0.255$). However, there was no significant interaction of emotion x movement x

contraception ($F(2,124) = 1.397$, p = .251, $\eta^2_p = 0.022$)) (see Figure 3C).

Figure 3. (A/B) Approach Avoidance Task (AAT) - Bias Scores (log-transformed) for angry and happy faces for
women taking oral contraceptives (OC) with androgenic progestin (light green) and those with anti-androgenic
progestin (light grey) before stress induction through Socially Evaluated Cold Pressor Test (SECPT) **(A)** and after
stress exposure **(B)**. **(C)** Bias Scores (log-transformed) for angry and happy faces for women who never took OC
(NCn, red), who took OC previously but discontinued (NCp, pink) and OC women (blue) over both AAT-trials. Error
bars indicate standard error.
490

3.4 Questionnaires

The groups did not differ significantly from each other in their scores in the BIS or BAS, BFI-

10, or ERQ scales (see Table 1), suggesting that there were no general personality differences

between NC women and OC users. The ERQ and the BIS and BAS subscales did not correlate

with the AAT bias scores (all p > 0,05), neither did the BFI-10 subscales.

Table 1. Sample descriptives

		NC (*n*=65)	OC (*n*=65)	*t*(128)	*p*	*d*	BF$_{10}$
500	Age	22.6 ± 2.40	21.9 ± 2.13	1.58	.116	.278	0.583
501	BMI	21.3 ± 1.72	21.1 ± 1.77	0.809	.420	.143	0.253
502							
503	*Questionnaires*						
504	BIS	2.09 ± 0.063	2.06 ± 0.055	0.368	.713	.065	0.199
505	BAS	1.87 ± 0.039	1.91 ± 0.038	-0.836	.405	-.147	0.258
506	BFI-10 (openness)	3.76 ± 0.125	3.59 ± 0.123	0.965	.336	.169	0.286
507	BFI-10 (neuroticism)	3.22 ± 0.110	3.45 ± 0.114	-1.460	.148	-.255	0.490
508	BFI-10 (extraversion)	3.62 ± 0.112	3.62 ± 0.118	0.047	.962	.008	0.188
509	BFI-10 (conscientiousness)	3.89 ± 0.100	3.77 ± 0.094	0.897	.371	.157	0.270
510	BFI-10 (agreeableness)	3.71 ± 0.109	3.49 ± 0.113	1.371	.173	.240	0.440
511	ERQ (cognitive reappraisal)	4.91 ± 0.122	4.66 ± 0.107	1.522	.130	.267	0.535
512	ERQ (expressive suppression)	3.17 ± 0.149	3.24 ± 0.133	-0.347	.729	-.061	0.198

513 Notes: BIS Behavioral Inhibition Scale, BAS Behavioral Approach Scale, BFI Big Five
514 Inventory, ERQ Emotion Regulation Questionnaire; values are mean ± standard deviation
515

4. Discussion

Oral contraceptives have an effect on central nervous levels of steroid hormones, which in

turn are known to modulate affective behavior. In the current study, we investigated whether

OC usage is related to altered approach and avoidance behavior in a between-subject design.

The results of this study suggest that women using oral contraceptives (OC) have significantly

reduced avoidance tendencies to social threat signals compared to women not using oral

contraceptives (NC). OC women were as quick to pull angry faces towards them as they were

to push them away and thus showed no avoidance tendencies. Socially evaluated physical

stress led to a similar rise in cortisol levels in both groups, but only in the OC group did stress

increase approach behavior to positive social signals. It was also noteworthy that women

taking OC responded overall faster than the NC women. Additional exploratory analyses

suggested that the composition of the preparations may have influenced the AAT results. Only

the OC women with an androgenic preparation showed a significant stress effect on approach

behavior. The composition of preparations was not related to the decreased avoidance effect

to threat signals in OC users, however. In sum, our study adds to the existing literature on

how contraceptives can impact affect in women, and points to altered approach/avoidance

tendencies as potential underlying mechanism of changes in affect and behavior.

4.1 OC-related differences in approach and avoidance behavior

We found that OC women pulled angry faces significantly faster towards them than NC women

did, but were comparably fast at pushing them away. This suggests that OC primarily facilitate

approach toward threat signals. Additionally, OC women responded with stronger approach

tendencies to happy faces after stress than before. These differences were found relative to

NC women who were in their early to mid follicular phase.

These results expand on recent studies on emotional reactivity changes in relation to

OC usage (Lewis et al., 2019; Montoya & Bos, 2017). For instance, Hamstra et al. (2014)

reported that women taking oral contraceptives detected fewer facial expressions of anger

than NC women, but did not differ in recognizing happy faces. Although we did not find a

reduced error rate, OC women did show a reduced avoidance response to angry faces. This

might be related to the previously reported OC-related reduction in amygdala activation to

negative stimuli (Petersen & Cahill, 2015) and reduced ERP responses to aversive stimuli

(Monciunskaite et al., 2019), but as we did not measure neural activity, this remains

speculative. Our results are in contrast to other studies, suggesting *enhanced* emotional

reactivity in OC users, especially to negative stimuli (Radke & Derntl, 2016; Spalek et al.,

2019). However, these studies measured explicit evaluations of the emotional stimuli, whereas

the AAT aims to measure automatic behavioral tendencies. Crucially, our data thus point to

motivational implications of OC usage, leading to altered affective action control and not mere

perceptual differences. Future studies should combine behavioral measures of automatic

affective behavior with neural measurements to tap into the underlying neural networks which

are affected by OCs.

The AAT effects we observed were independent of the exact OC preparation. As we

did not select for specific OC preparations and did not measure hormonal levels, conclusions

regarding the relative role of estradiol and progesterone are difficult. Previous research has

suggested that endogenous estradiol is associated with increased activity in emotion-

regulatory brain areas such as the dorsolateral prefrontal cortex or the anterior cingulate

cortex (Chung et al., 2019; Sharma et al., 2021). For progesterone it is known that it impairs

emotion processing and recognition (Derntl et al., 2008; Guapo et al., 2009; van Wingen et

al., 2007). Based on these previous findings, the observed AAT effects might rather be

attributed to the OC effects on estradiol levels, but this needs to be investigated in future

studies.

OC women were also generally faster than NC women. This is similar to results of

another study, who also reported faster response times in OC users, this time to visual stimuli

without any emotional component (Grikšienė & Rukšėnas, 2009). Older studies, however,

reported generally increased reaction times in OC users (Garrett & Elder, 1984; Wuttke et al.,

1975). However, it is important to note that other formulations of oral contraceptives were used

at the time the studies were conducted. In a more recent study presenting emotional stimuli,

women taking OC were also faster than NC women, but only in response to sad faces and not

for happy or angry faces (Hamstra et al., 2014). Moreover, there is some evidence that the

menstrual cycle phase has an influence on the reaction times in various paradigms (Li et al.,

2022; Pletzer et al., 2014). Group differences in response times independently of experimental

conditions are notoriously difficult to interpret as various factors might have contributed, such

as arousal, psychomotor processing speed or motivation. It remains for future studies to

examine how reliable these response time differences are.

In our study, women who had previously taken OC (NC_p) did not differ in their approach

avoidance pattern from women who had never taken OC (NC_n). Regarding general response

times however, women who had never taken OC were slowest, whereas women who had

taken OC previously were not slower than current OC users. Pletzer and Kerschbaum (2015)

suggested that effects of oral contraceptives could persist beyond the duration of intake. While

they observed neurostructural changes after OC use (relative hippocampal gray matter

volume was found to be associated positively with the duration of previous OC use), we found

an effect on a behavioral level. Although the effect was of medium size, it should be treated

with caution because of the unequal group sizes and the exploratory nature of the analysis.

Nevertheless, the results show that, in future studies, a differentiation between current and

previous OC users and nonusers might be useful.

4.2 OC-related effects in their interaction with stress

As hypothesized, stress influenced approach/avoidance behavior of OC and NC women in

different ways. Only the OC women showed altered AAT behavior after stress compared to

before. They reacted after stress with stronger approach tendencies to happy faces and

unaltered to angry faces. The increased approach bias resulted primarily from faster pull

reactions to happy faces. Our hypothesis was based on previous work by Roelofs and

colleagues (2005), who showed reduced approach and avoidance biases after stress in high

cortisol responders. Our results are only partly in line with these findings as the stress effects

in our sample were independent of the cortisol response and specific for approach behavior.

To our knowledge, no other study tested for stress effects on approach or avoidance biases

in women taking oral contraceptives. Our results thus remain to be replicated.

602   The stress induced increase in the approach bias in OC users could be interpreted as

kind a of tend-and-befriend response, characterized by responding to stress by aligning with

social groups to ensure security (Taylor et al., 2000). Although the tend-and-befriend theory

assumes that steroid hormones together with oxytocin drive the behavioral response to stress

(Taylor et al., 2000), there is to our knowledge little direct evidence for estradiol or

progesterone influencing social behavior after stress. In fact, one study revealed no significant

difference between OC and NC women in prosocial behavior under stress (von Dawans et al.,

2019). However, the study did not distinguish between OC women using androgenic and anti-

androgenic preparations. In our study, we noticed that women taking androgenic OC reacted

with heightened approach to happy faces after stress. There is some evidence that women

using androgenic OC showed increased stress effects on vasoconstriction compared to

women using anti-androgenic OC (Straneva et al., 2000). However, so far there is no study

directly comparing androgenic and anti-androgenic OC effects on cognitive and affective

changes after stress.

616   Indirect evidence for a role of steroid hormones on stress-induced changes in social

cognition and behavior comes from studies showing sex differences in the response to stress.

Women showed improved affective self-other distinction after stress (Tomova et al., 2014),

whereas men showed worse self-other distinction resulting in an increased emotional

egocentricity bias. In a female sample including both OC and NC women, Zhang et al. (2019)

found no stress effect on pro-social decision-making, whereas men showed more pro-social

behavior with higher cortisol stress responses. Both studies did not measure steroid hormone

levels however, making a direct comparison with our findings difficult. There is also extensive

literature about stress effects on empathy (Nitschke & Bartz, 2023) and some evidence for

sex differences in these effects. For instance, Nitschke and colleagues (2022) found no stress

effects on empathic accuracy in women, whereas men were more accurate after stress

compared to before stress. Interestingly, this study also reported generally reduced empathic

accuracy in OC users compared to NC women. Sex differences in stress effects on empathy

seem to be context dependent however, as others found a stress-induced reduction in

empathic accuracy in women compared to men (Crenshaw et al., 2019), when assessed in

romantic couples. Otherwise, social cognition even increased in women after stress, but only

in low cortisol responders (Smeets et al., 2009). In sum, these previous studies suggest

interactive effects of sex/gender and stress on social cognition and empathy, but the direction

of effects seems to depend on the experimental measure of social cognition and affect. It is

important to emphasize that our paradigm measures automatic emotional action control and

thus addresses very different affective and neural processes than paradigms of empathic

accuracy or social cognition. Clearly, more research is needed on how steroid hormones and

stress interact in their effects on various cognitive and affective processes and how this

ultimately affects social behavior.

Remarkably, neither NC and OC women nor women taking androgenic and

antiandrogenic OC differed significantly in their cortisol stress response. This is surprising as

a blunted cortisol stress response has been often replicated in previous studies (Lewis et al.,

2019; Tronson & Schuh, 2022). However, there are also other reports who did not find any

group differences in this stress marker (Lovallo et al., 2019; Sharma et al., 2020). At least

Lovallo and colleagues found the reduced cortisol stress response in OC users only when

tested in the morning, but not in the afternoon (Lovallo et al., 2019). As we tested participants

in the afternoon, this might explain the lack of group differences. Moreover, we instructed

participants not to exercise, eat anything, drink coffee, take medications, or smoke for one

hour before the appointment, following previous studies (Roelofs et al., 2005). The time

window of one hour might have been too short (Hansen et al., 2012), which might have

introduced variability within the groups that prevented us from detecting group differences.

Finally, we could not analyze cortisol levels in 20 women which reduced power to detect group

differences. Since cortisol is only one of many stress markers, the groups might still have

differed in other stress responses such as their subjective stress experience or other

physiological stress markers. Unfortunately, we did not measure any other stress responses

in our participants. Because we focused on the approach and avoidance behavior of OC users

and the potential modulating effect of stress-induced cortisol changes, we had no specific

research questions or hypotheses regarding the subjective experience of stress. Previous

work on OC effects on subjective stress experience or other stress markers is inconclusive

(Lewis et al., 2019), as some studies reported no OC related differences in subjective stress

responses (Kirschbaum et al., 1999; Merz, 2017; Mordecai et al., 2017; Sharma et al., 2020)

but others did not measure it. More research is clearly needed to understand the effects of

OC usage on stress responses on a subjective and behavioral level.

4.4 Limitations

Like many other studies investigating the influence of oral contraceptives on the female

organism, our study is also limited by the "survivor effect" (Kutner & Brown, 1972). This means

that our study only comprised women who did not stop taking OC because of serious side

effects, whereas those who did develop side effects abandoned the treatment early. Future

studies should implement a longitudinal, within-subject design to avoid pre-selection of

participants. To be able to directly relate behavioral effects in the experimental paradigms to

potential affective side effects, one should also systematically assess emotional well-being

and psychiatric symptoms. The cross-sectional study design also means that the effects are

not uniquely attributable to OC use but may be influenced by other factors which caused

women to decide for or against OC. Although the lack of group differences in personality and

emotion regulation questionnaires speaks against this argument, we cannot rule out the

possibility that there are influencing variables which differ between women taking oral

contraceptives or not.

In our AAT paradigm, we only presented faces with positive and negative valence. It

would be beneficial to add a neutral condition in future studies to clarify if the shortened

reaction times of OC women are due to the affective component. Combining the experiment

with fMRI measurements could also shed light on the neural mechanisms associated with

these effects. In particular, the amygdala and aPFC could be examined more closely to

understand the neural basis of the observed behavioral changes (Kaldewaij et al., 2017).

We did not measure steroid hormone levels in the participants, which limits the

conclusions we can derive from our observations. Steroid hormone measurements would

have allowed to go beyond a simple group comparison and to directly correlate behavioral

effects with hormonal levels. This could have helped to clarify the relative contribution of

estradiol and progesterone to the AAT effects (for comparison, see Strojny et al. (2021)).

Moreover, hormone measurements could have supported the presumed group differences in

steroid hormones. We decided to test participants on the 1st to 3rd day of the active pill phase

and, in case of women not taking hormonal contraception, on the 8th to 10th day of their

menstrual cycle. As the 8th to 10th day are not clearly early follicular phase anymore, this might

have led to more variability within the NC group already, as in some estradiol might have been

rising already again. Also, we did not track cycle length of the participants but only asked them

to come to the lab at the specified day after menses. However, such variability should rather

have led to smaller effects or null effects. Also with regard to the measurement time in the OC

group, there is debate on when to best perform measurements to study OC effects (Gravelsins

et al., 2023). Hormonal levels might vary depending on whether women are in their active or

inactive phase, when in the inactive phase and, depending on the pharmacokinetics, when

during the day (Gravelsins et al., 2023; Stanczyk et al., 2013). In our study, we decided to

keep the measurement time within the active phase constant, and we did not observe

differences between participants depending on pill intake. However, it should be noted that

hormone measurements could have clarified the hormonal status of participants and the lack

of hormone measurements presents a limitation of this study.

4.5 Conclusion

This work stands in line with a series of previous studies showing that oral contraceptives can

influence cognitive and affective processes (Gravelsins et al., 2023; Lewis et al., 2019;

Tronson & Schuh, 2022). Specifically, we showed that OC use is associated with a reduced

avoidance tendency to social threat signals and an increased stress-induced approach

tendency to positive social signals. This expands on other research demonstrating altered

reactivity to particularly negative emotional stimuli in OC users (Lewis et al., 2019) and

suggests that these differences are not limited to the perceptual level, but have motivational

implications. In addition to recent work speaking for a negative OC effect on affective empathy

and prosocial decision-making (Kimmig et al., 2022; Strojny et al., 2021), we show that women

taking oral contraceptives differ already on automatic motivated behaviors to affective signals.

Future studies will have to examine how these differences relate to neural mechanisms on the

one hand and presumed affective side-effects of OCs on the other hand.

CRediT author statement:

Jasmin Thurley: Methodology, Software, Formal analysis, Investigation, Writing – Original

Draft, Visualization **Macià Buades-Rotger:** Methodology, Writing – Review and Editing

Georg Serfling: Methodology, Writing – Review and Editing **Thessa Howaldt:** Investigation,

Writing – Review and Editing **Nicole Reisch:** Writing – Review and Editing **Ulrike Krämer:**

Conceptualization, Methodology, Software, Resources, Writing – Review and Editing,

Supervision, Project administration

Acknowledgements:

We thank Carla Bücker, Annika Kühle and Luisa Ackermann for help during data acquisition.

UMK and NR are supported by the German Research Foundation (DFG) as part of the CRC

1665.

References

References

Anderl, C., Wit, A. E., Giltay, E. J., Oldehinkel, A. J., & Chen, F. S. (2022). Association between
adolescent oral contraceptive use and future major depressive disorder: a prospective cohort
study. *Journal of Child Psychology and Psychiatry*, *63*(3), 333-341.
https://doi.org/10.1111/jcpp.13476
Beyer, F., Buades-Rotger, M., Claes, M., & Krämer, U. M. (2017). Hit or Run: Exploring Aggressive and
Avoidant Reactions to Interpersonal Provocation Using a Novel Fight-or-Escape Paradigm
(FOE). *Front Behav Neurosci*, *11*, 190. https://doi.org/10.3389/fnbeh.2017.00190
Brønnick, M. K., Økland, I., Graugaard, C., & Brønnick, K. K. (2020). The Effects of Hormonal
Contraceptives on the Brain: A Systematic Review of Neuroimaging Studies [Systematic
Review]. *Front Psychol*, *11*(2813). https://doi.org/10.3389/fpsyg.2020.556577
Buades-Rotger, M., Engelke, C., Beyer, F., Keevil, B. G., Brabant, G., & Krämer, U. M. (2016).
Endogenous testosterone is associated with lower amygdala reactivity to angry faces and
reduced aggressive behavior in healthy young women. *Scientific Reports*, *6*, 38538.
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5141420/pdf/srep38538.pdf
Carroll, D., Ginty, A. T., Whittaker, A. C., Lovallo, W. R., & de Rooij, S. R. (2017). The behavioural,
cognitive, and neural corollaries of blunted cardiovascular and cortisol reactions to acute
psychological stress. *Neuroscience & Biobehavioral Reviews*, *77*, 74-86.
https://doi.org/https://doi.org/10.1016/j.neubiorev.2017.02.025
Carver, C. S., & White, T. L. (1994). Behavioral inhibition, behavioral activation, and affective responses
to impending reward and punishment: The BIS/BAS Scales. *Journal of Personality and Social
Psychology*, *67*, 319-333. https://doi.org/10.1037/0022-3514.67.2.319
Chen, M., & Bargh, J. A. (1999). Consequences of Automatic Evaluation: Immediate Behavioral
Predispositions to Approach or Avoid the Stimulus. *Personality and Social Psychology Bulletin*,
25(2), 215-224. https://doi.org/10.1177/0146167299025002007
Christin-Maitre, S. (2013). History of oral contraceptive drugs and their use worldwide. *Best Practice &
Research Clinical Endocrinology & Metabolism*, *27*(1), 3-12.
https://doi.org/10.1016/j.beem.2012.11.004
Chung, Y. S., Poppe, A., Novotny, S., Epperson, C. N., Kober, H., Granger, D. A., Blumberg, H. P.,
Ochsner, K., Gross, J. J., Pearlson, G., & Stevens, M. C. (2019). A preliminary study of
association between adolescent estradiol level and dorsolateral prefrontal cortex activity during
emotion regulation. *Psychoneuroendocrinology*, *109*, 104398.
https://doi.org/10.1016/j.psyneuen.2019.104398
Crenshaw, A. O., Leo, K., & Baucom, B. R. W. (2019). The effect of stress on empathic accuracy in
romantic couples. *J Fam Psychol*, *33*(3), 327-337. https://doi.org/10.1037/fam0000508
Derntl, B., Windischberger, C., Robinson, S., Lamplmayr, E., Kryspin-Exner, I., Gur, R. C., Moser, E.,
& Habel, U. (2008). Facial emotion recognition and amygdala activation are associated with
menstrual cycle phase. *Psychoneuroendocrinology*, *33*(8), 1031-1040.
https://doi.org/10.1016/j.psyneuen.2008.04.014
Enter, D., Spinhoven, P., & Roelofs, K. (2014). Alleviating social avoidance: effects of single dose
testosterone administration on approach-avoidance action. *Hormones and Behavior*, *65*(4),
351-354. https://doi.org/10.1016/j.yhbeh.2014.02.001
Ernst, U., Baumgartner, L., Bauer, U., & Janssen, G. (2002). Improvement of quality of life in women
using a low-dose desogestrel-containing contraceptive: results of an observational clinical
evaluation. *The European Journal of Contraception & Reproductive Health Care*, *7*(4), 238-
243. https://doi.org/10.1080/ejc.7.4.238.243
Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. *Behavior Research
Methods*, *39*(2), 175-191. https://doi.org/10.3758/BF03193146
Garrett, K. F., & Elder, S. T. (1984). The menstrual cycle from a bio-behavioral approach: A comparison
of oral contraceptive and non-contraceptive users. *International Journal of Psychophysiology*,
1(2), 209-214. https://doi.org/10.1016/0167-8760(84)90039-4
Gingnell, M., Engman, J., Frick, A., Moby, L., Wikström, J., Fredrikson, M., & Sundström-Poromaa, I.
(2013). Oral contraceptive use changes brain activity and mood in women with previous
negative affect on the pill—A double-blinded, placebo-controlled randomized trial of a
levonorgestrel-containing combined oral contraceptive. *Psychoneuroendocrinology*, *38*(7),
1133-1144. https://doi.org/10.1016/j.psyneuen.2012.11.006

Graham, B. M., & Milad, M. R. (2013). Blockade of estrogen by hormonal contraceptives impairs fear
extinction   in   female   rats   and   women.   *Biol Psychiatry*,   *73*(4),   371-378.
https://doi.org/10.1016/j.biopsych.2012.09.018

Gravelsins, L., Duncan, K., & Einstein, G. (2021). Do oral contraceptives affect young women's
memory? Dopamine-dependent working memory is influenced by COMT genotype, but not time
of pill ingestion. *PLoS One*, *16*(6), e0252807. https://doi.org/10.1371/journal.pone.0252807

Gravelsins, L., Zhao, S., & Einstein, G. (2023). Hormonal contraception and cognition: Considering the
influence of endogenous ovarian hormones and genes for clinical translation. *Front
Neuroendocrinol*, *70*, 101067. https://doi.org/10.1016/j.yfrne.2023.101067

Grikšienė, R., & Rukšėnas, O. (2009). Cognitive effects of hormone-based contraception in young
healthy women. *Biologija*, *55*(3-4).

Gross, J. J., & John, O. P. (2003). Individual differences in two emotion regulation processes:
implications for affect, relationships, and well-being. *Journal of Personality and Social
Psychology*, *85*(2), 348-362. https://doi.org/10.1037/0022-3514.85.2.348

Guapo, V. G., Graeff, F. G., Zani, A. C., Labate, C. M., dos Reis, R. M., & Del-Ben, C. M. (2009). Effects
of sex hormonal levels and phases of the menstrual cycle in the processing of emotional faces.
Psychoneuroendocrinology,   *34*(7),   1087-1094.
https://doi.org/10.1016/j.psyneuen.2009.02.007

Hampson, E. (2020). A brief guide to the menstrual cycle and oral contraceptive use for researchers in
behavioral   endocrinology.   *Hormones   and   Behavior*,   *119*,   104655.
https://doi.org/10.1016/j.yhbeh.2019.104655

Hamstra, D. A., De Rover, M., De Rijk, R. H., & Van der Does, W. (2014). Oral contraceptives may alter
the detection of emotions in facial expressions. *European Neuropsychopharmacology*, *24*(11),
1855–1859.

Hansen, A. M., Gunnarsson, L., Harris, A., Eller, N., Garvin, P., & Garde, A. H. (2012). Biological
markers and salivary cortisol. *The Role of Saliva Cortisol Measurement in Health and Disease*.
Sharja: Bentham eBooks, 88–117.

Hertel, J., König, J., Homuth, G., Van der Auwera, S., Wittfeld, K., Pietzner, M., Kacprowski, T., Pfeiffer,
L., Kretschmer, A., Waldenberger, M., Kastenmüller, G., Artati, A., Suhre, K., Adamski, J.,
Langner, S., Völker, U., Völzke, H., Nauck, M., Friedrich, N., & Grabe, H. J. (2017). Evidence
for Stress-like Alterations in the HPA-Axis in Women Taking Oral Contraceptives. *Scientific
Reports*, *7*(1), 14111. https://doi.org/10.1038/s41598-017-13927-7

Kaldewaij, R., Koch, S. B., Volman, I., Toni, I., & Roelofs, K. (2017). On the Control of Social Approach-
Avoidance Behavior: Neural and Endocrine Mechanisms. *Curr Top Behav Neurosci*, *30*, 275-
293. https://doi.org/10.1007/7854_2016_446

Kimmig, A. S., Wildgruber, D., Gartner, A., Drotleff, B., Krylova, M., Lammerhofer, M., Sundstrom-
Poromaa, I., & Derntl, B. (2022). Lower affective empathy in oral contraceptive users: a cross-
sectional fMRI study. *Cereb Cortex*. https://doi.org/10.1093/cercor/bhac345

Kirschbaum, C., Kudielka, B. M., Gaab, J., Schommer, N. C., & Hellhammer, D. H. (1999). Impact of
Gender, Menstrual Cycle Phase, and Oral Contraceptives on the Activity of the Hypothalamus-
Pituitary-Adrenal   Axis: .   *Psychosomatic   Medicine*,   *61*(2),   154-162.
https://doi.org/10.1097/00006842-199903000-00006

Kirschbaum, C., Pirke, K.-m., & Hellhammer, D. H. (1995). Preliminary evidence for reduced cortisol
responsivity to psychological stress in women using oral contraceptive medication.
Psychoneuroendocrinology,   *20*(5),   509-514.   https://doi.org/https://doi.org/10.1016/0306-
4530(94)00078-O

Kutner, S. J., & Brown, W. L. (1972). Types of oral contraceptives, depression, and premenstrual
symptoms. *J Nerv Ment Dis*, *155*(3), 153-162. https://doi.org/10.1097/00005053-197209000-
00001

Langner, O., Dotsch, R., Bijlstra, G., Wigboldus, D. H. J., Hawk, S. T., & van Knippenberg, A. (2010).
Presentation and validation of the Radboud Faces Database. *Cognition and Emotion*, *24*(8),
1377-1388. https://doi.org/10.1080/02699930903485076

Lewis, C. A., Kimmig, A. S., Zsido, R. G., Jank, A., Derntl, B., & Sacher, J. (2019). Effects of Hormonal
Contraceptives on Mood: A Focus on Emotion Recognition and Reactivity, Reward Processing,
and Stress Response. *Curr Psychiatry Rep*, *21*(11), 115. https://doi.org/10.1007/s11920-019-
1095-z

Li, D., Zhang, L., & Wang, X. (2022). The Effect of Menstrual Cycle Phases on Approach–Avoidance
Behaviors in Women: Evidence from Conscious and Unconscious Processes. *Brain Sciences*,
12(10), 1417. https://doi.org/10.3390/brainsci12101417

Lovallo, W. R., Cohoon, A. J., Acheson, A., Vincent, A. S., & Sorocco, K. H. (2019). Cortisol stress reactivity in women, diurnal variations, and hormonal contraceptives: studies from the Family Health Patterns Project. *Stress*, *22*(4), 421-427. https://doi.org/10.1080/10253890.2019.1581760

Love, J., Selker, R., Marsman, M., Jamil, T., Dropmann, D., Verhagen, J., Ly, A., Gronau, Q. F., Šmíra, M., Epskamp, S., Matzke, D., Wild, A., Knight, P., Rouder, J. N., Morey, R. D., & Wagenmakers, E.-J. (2019). JASP: Graphical Statistical Software for Common Statistical Designs. *Journal of Statistical Software*, *88*, 1-17. https://doi.org/10.18637/jss.v088.i02

Merz, C. J. (2017). Contribution of stress and sex hormones to memory encoding. *Psychoneuroendocrinology*, *82*, 51-58. https://doi.org/10.1016/j.psyneuen.2017.05.002

Meulenberg, P. M. M., & Hofman, J. A. (1990). The effect of oral contraceptive use and pregnancy on the daily rhythm of cortisol and cortisone. *Clinica Chimica Acta*, *190*(3), 211-221. https://doi.org/10.1016/0009-8981(90)90175-R

Monciunskaite, R., Malden, L., Lukstaite, I., Ruksenas, O., & Griksiene, R. (2019). Do oral contraceptives modulate an ERP response to affective pictures? *Biol Psychol*, *148*, 107767. https://doi.org/10.1016/j.biopsycho.2019.107767

Montoya, E. R., & Bos, P. A. (2017). How Oral Contraceptives Impact Social-Emotional Behavior and Brain Function. *Trends Cogn Sci*, *21*(2), 125-136. https://doi.org/10.1016/j.tics.2016.11.005

Mordecai, K. L., Rubin, L. H., Eatough, E., Sundermann, E., Drogos, L., Savarese, A., & Maki, P. M. (2017). Cortisol reactivity and emotional memory after psychosocial stress in oral contraceptive users. *J Neurosci Res*, *95*(1-2), 126-135. https://doi.org/10.1002/jnr.23904

Nitschke, J. P., & Bartz, J. A. (2023). The association between acute stress & empathy: A systematic literature review. *Neurosci Biobehav Rev*, *144*, 105003. https://doi.org/10.1016/j.neubiorev.2022.105003

Nitschke, J. P., Pruessner, J. C., & Bartz, J. A. (2022). Stress and Stress-Induced Glucocorticoids Facilitate Empathic Accuracy in Men but Have No Effects for Women. *Psychol Sci*, *33*(10), 1783-1794. https://doi.org/10.1177/09567976221101315

Österlund, M., Kuiper, G. G., Gustafsson, J.-Å., & Hurd, Y. L. (1998). Differential distribution and regulation of estrogen receptor-α and-β mRNA within the female rat brain. *Molecular Brain Research*, *54*(1), 175–180.

Petersen, N., & Cahill, L. (2015). Amygdala reactivity to negative stimuli is influenced by oral contraceptive use. *Soc Cogn Affect Neurosci*, *10*(9), 1266-1272. https://doi.org/10.1093/scan/nsv010

Pletzer, B., Kronbichler, M., & Kerschbaum, H. (2015). Differential effects of androgenic and anti-androgenic progestins on fusiform and frontal gray matter volume and face recognition performance. *Brain Res*, *1596*, 108–115.

Pletzer, B., Kronbichler, M., Nuerk, H.-C., & Kerschbaum, H. (2014). Hormonal contraceptives masculinize brain activation patterns in the absence of behavioral changes in two numerical tasks. *Brain Res*, *1543*, 128-142. https://doi.org/10.1016/j.brainres.2013.11.007

Poromaa, I. S., & Segebladh, B. (2012). Adverse mood symptoms with oral contraceptives. *Acta Obstet Gynecol Scand*, *91*(4), 420-427. https://doi.org/10.1111/j.1600-0412.2011.01333.x

Pruessner, J. C., Kirschbaum, C., Meinlschmid, G., & Hellhammer, D. H. (2003). Two formulas for computation of the area under the curve represent measures of total hormone concentration versus time-dependent change. *Psychoneuroendocrinology*, *28*(7), 916-931. https://doi.org/10.1016/s0306-4530(02)00108-7

Radke, S., & Derntl, B. (2016). Affective responsiveness is influenced by intake of oral contraceptives. *Eur Neuropsychopharmacol*, *26*(6), 1014-1019. https://doi.org/10.1016/j.euroneuro.2016.03.004

Radke, S., Volman, I., Mehta, P., van Son, V., Enter, D., Sanfey, A., Toni, I., de Bruijn, E. R., & Roelofs, K. (2015). Testosterone biases the amygdala toward social threat approach. *Science advances*, *1*(5), e1400074. https://doi.org/10.1126/sciadv.1400074

Rammstedt, B., & John, O. P. (2007). Measuring personality in one minute or less: A 10-item short version of the Big Five Inventory in English and German. *Journal of Research in Personality*, *41*, 203-212. https://doi.org/10.1016/j.jrp.2006.02.001

Rehbein, E., Hornung, J., Sundstrom Poromaa, I., & Derntl, B. (2021). Shaping of the Female Human Brain by Sex Hormones: A Review. *Neuroendocrinology*, *111*(3), 183-206. https://doi.org/10.1159/000507083

Roelofs, K., Elzinga, B. M., & Rotteveel, M. (2005). The effects of stress-induced cortisol responses on approach–avoidance behavior. *Psychoneuroendocrinology*, *30*(7), 665-677. https://doi.org/10.1016/j.psyneuen.2005.02.008

Roelofs, K., Putman, P., Schouten, S., Lange, W. G., Volman, I., & Rinck, M. (2010). Gaze direction differentially affects avoidance tendencies to happy and angry faces in socially anxious individuals. *Behav Res Ther*, *48*(4), 290-294. https://doi.org/10.1016/j.brat.2009.11.008

Schwabe, L., & Schächinger, H. (2018). Ten years of research with the Socially Evaluated Cold Pressor Test: Data from the past and guidelines for the future. *Psychoneuroendocrinology*, *92*, 155–161. https://doi.org/10.1016/j.psyneuen.2018.03.010

Sharma, R., Cameron, A., Fang, Z., Ismail, N., & Smith, A. (2021). The regulatory roles of progesterone and estradiol on emotion processing in women. *Cogn Affect Behav Neurosci*, *21*(5), 1026-1038. https://doi.org/10.3758/s13415-021-00908-7

Sharma, R., Smith, S. A., Boukina, N., Dordari, A., Mistry, A., Taylor, B. C., Felix, N., Cameron, A., Fang, Z., Smith, A., & Ismail, N. (2020). Use of the birth control pill affects stress reactivity and brain structure and function. *Hormones and Behavior*, *124*, 104783. https://doi.org/10.1016/j.yhbeh.2020.104783

Slattery, D. A., & Cryan, J. F. (2017). Modelling depression in animals: at the interface of reward and stress pathways. *Psychopharmacology (Berl)*, *234*(9-10), 1451-1465. https://doi.org/10.1007/s00213-017-4552-6

Smeets, T., Dziobek, I., & Wolf, O. T. (2009). Social cognition under stress: differential effects of stress-induced cortisol elevations in healthy young men and women. *Hormones and Behavior*, *55*(4), 507-513. https://doi.org/10.1016/j.yhbeh.2009.01.011

Spalek, K., Loos, E., Schicktanz, N., Hartmann, F., de Quervain, D., Stier, C., & Milnik, A. (2019). Women using hormonal contraceptives show increased valence ratings and memory performance for emotional information. *Neuropsychopharmacology*, *44*(7), 1258-1264. https://doi.org/10.1038/s41386-019-0362-3

Stanczyk, F. Z., Archer, D. F., & Bhavnani, B. R. (2013). Ethinyl estradiol and 17beta-estradiol in combined oral contraceptives: pharmacokinetics, pharmacodynamics and risk assessment. *Contraception*, *87*(6), 706-727. https://doi.org/10.1016/j.contraception.2012.12.011

Straneva, P., Hinderliter, A., Wells, E., Lenahan, H., & Girdler, S. (2000). Smoking, oral contraceptives, and cardiovascular reactivity to stress. *Obstetrics & Gynecology*, *95*(1), 78-83. https://doi.org/10.1016/S0029-7844(99)00497-4

Strojny, J., Domes, G., Fischbacher, U., & von Dawans, B. (2021). The modulation of social behavior and empathy via oral contraceptives and female sex hormones. *Psychoneuroendocrinology*, *131*, 105250. https://doi.org/10.1016/j.psyneuen.2021.105250

Taxier, L. R., Gross, K. S., & Frick, K. M. (2020). Oestradiol as a neuromodulator of learning and memory. *Nature Reviews Neuroscience*, *21*(10), 535-550. https://doi.org/10.1038/s41583-020-0362-7

Taylor, S. E., Klein, L. C., Lewis, B. P., Gruenewald, T. L., Gurung, R. A., & Updegraff, J. A. (2000). Biobehavioral responses to stress in females: tend-and-befriend, not fight-or-flight. *Psychol Rev*, *107*(3), 411-429. https://doi.org/10.1037/0033-295x.107.3.411

Tomova, L., von Dawans, B., Heinrichs, M., Silani, G., & Lamm, C. (2014). Is stress affecting our ability to tune into others? Evidence for gender differences in the effects of stress on self-other distinction. *Psychoneuroendocrinology*, *43*, 95-104. https://doi.org/10.1016/j.psyneuen.2014.02.006

Tronson, N. C., & Schuh, K. M. (2022). Hormonal contraceptives, stress, and the brain: The critical need for animal models. *Frontiers in Neuroendocrinology*, *67*, 101035. https://doi.org/10.1016/j.yfrne.2022.101035

van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz, A., Evans, N. J., Gronau, Q. F., Haaf, J. M., Hinne, M., Kucharský, Š., Ly, A., Marsman, M., Matzke, D., Gupta, A. R. K. N., Sarafoglou, A., Stefan, A., Voelkel, J. G., & Wagenmakers, E.-J. (2021). The JASP guidelines for conducting and reporting a Bayesian analysis. *Psychonomic Bulletin & Review*, *28*(3), 813-826. https://doi.org/10.3758/s13423-020-01798-5

van Wingen, G., Mattern, C., Verkes, R. J., Buitelaar, J., & Fernandez, G. (2010). Testosterone reduces amygdala-orbitofrontal cortex coupling. *Psychoneuroendocrinology*, *35*(1), 105-113. https://doi.org/10.1016/j.psyneuen.2009.09.007

van Wingen, G., van Broekhoven, F., Verkes, R. J., Petersson, K. M., Backstrom, T., Buitelaar, J., & Fernandez, G. (2007). How progesterone impairs memory for biologically salient stimuli in healthy young women. *J Neurosci*, *27*(42), 11416-11423. https://doi.org/10.1523/JNEUROSCI.1715-07.2007

van Wingen, G. A., Ossewaarde, L., Backstrom, T., Hermans, E. J., & Fernandez, G. (2011). Gonadal hormone regulation of the emotion circuitry in humans. *Neuroscience*, *191*, 38-45. https://doi.org/10.1016/j.neuroscience.2011.04.042

van Wingen, G. A., van Broekhoven, F., Verkes, R. J., Petersson, K. M., Bäckström, T., Buitelaar, J.
K., & Fernández, G. (2008). Progesterone selectively increases amygdala reactivity in women.
Molecular Psychiatry, *13*(3), 325-333. https://doi.org/10.1038/sj.mp.4002030
Volman, I., Verhagen, L., Den Ouden, H. E. M., Fernández, G., Rijpkema, M., Franke, B., Toni, I., &
Roelofs, K. (2013). Reduced Serotonin Transporter Availability Decreases Prefrontal Control
of the Amygdala. *The Journal of Neuroscience*, *33*(21), 8974-8979.
https://doi.org/10.1523/JNEUROSCI.5518-12.2013
von Borries, L. A. K., Volman, I., de Bruijn, E. R., Bulten, B. H., Verkes, R. J., & Roelofs, K. (2012).
Psychopaths lack the automatic avoidance of social threat: relation to instrumental aggression.
Psychiatry Res, *200*(2-3), 761-766. https://doi.org/10.1016/j.psychres.2012.06.026
von Dawans, B., Ditzen, B., Trueg, A., Fischbacher, U., & Heinrichs, M. (2019). Effects of acute stress
on social behavior in women. *Psychoneuroendocrinology*, *99*, 137-144.
https://doi.org/10.1016/j.psyneuen.2018.08.031
Wuttke, W., Arnold, P., Becker, D., Creutzfeldt, O., Langenstein, S., & Tirsch, W. (1975). Circulating
hormones, EEG, and performance in psychological tests of women with and without oral
contraceptives. *Psychoneuroendocrinology*, *1*(2), 141-152. https://doi.org/10.1016/0306-
4530(75)90006-2
Zeidan, M. A., Igoe, S. A., Linnman, C., Vitalo, A., Levine, J. B., Klibanski, A., Goldstein, J. M., & Milad,
M. R. (2011). Estradiol modulates medial prefrontal cortex and amygdala activity during fear
extinction in women and female rats. *Biol Psychiatry*, *70*(10), 920-927.
https://doi.org/10.1016/j.biopsych.2011.05.016
Zhang, Q., Ma, J., & Nater, U. M. (2019). How Cortisol Reactivity Influences Prosocial Decision-Making:
The Moderating Role of Sex and Empathic Concern. *Front Hum Neurosci*, *13*, 415.
https://doi.org/10.3389/fnhum.2019.00415
995