Maturational Differences in Affective Behaviors Involves Changes in Frontal Cortical-Hippocampal Functional Connectivity and Metabolomic Profiles

[1,3,4]Marcelo Febo, [5]Rohit Mahar, [1]Nicholas A. Rodriguez, [1]Joy Buraima, [1]Marjory Pompilus, [1]Aeja M. Pinto, [1]Matteo Grudny, [1,4]Adriaan W. Bruijnzeel, [2,3]Matthew E. Merritt

[1]Departments of Psychiatry, [2]Biochemistry and Molecular Biology, [3]Advanced Magnetic Resonance Imaging and Spectroscopy (AMRIS) Facility, [4]McKnight Brain Institute, College of Medicine, University of Florida, Gainesville, Florida. [5]Department of Chemistry, Hemvati Nandan Bahuguna Garhwal University (A Central University), Srinagar Garhwal, Uttarakhand, India

Short title: Maturational changes in functional connectivity and affective behaviors

Corresponding author: Marcelo Febo, PhD

Contact information: Department of Psychiatry, P.O. Box 100256, Gainesville, Florida 32611; Office: (352) 294 4911; Email: febo@ufl.edu

Acknowledgements: This study was funded by a National Institute on Aging grant (AG065819). Joy Buraima was supported by the Summer Neuroscience Internship Program (SNIP) of the McKnight Brain Institute. The contents of this manuscript are solely the responsibility of the authors and do not necessarily represent the official views of the funding agencies. This work was performed in the McKnight Brain Institute at the National High Magnetic Field Laboratory's AMRIS Facility, which is supported by National Science Foundation Cooperative Agreement No. DMR-1644779 and the State of Florida.

Abstract

Differences in affective behavioral expression from early to late adulthood is thought to

involve changes in frontal cortical responsiveness to negative valence stimuli. In mice, similar

maturational changes in affective behaviors have also been reported but the functional neural

circuitry remains unclear. In the present study we investigated age variations in affective

behaviors and functional connectivity in male and female C57BL6/J mice. Mice aged 10, 30 and

60 weeks (wo) were tested over 8 weeks for open field activity, sucrose reward preference,

social interactions and fear conditioning and functional neuroimaging. Frontal cortical and

hippocampal tissues were excised for metabolomics analysis. Our results indicate that young

10wo mice display greater levels of anxiety-like locomotor behavior and develop robust fear

conditioning compared to older adult and late middle-aged mice (30-60wo). This was

accompanied by greater functional connectivity between a temporal cortical/auditory cortex

network and subregions of the anterior cingulate cortex and ventral hippocampus, and a greater

network modularity and assortative mixing of nodes in young versus older adult mice.

Metabolome analyses identified differences in several essential amino acids between 10wo

mice and the other age groups. The results support high 'emotionality' in younger versus older

adult mice involving greater prefrontal-hippocampal connectivity.

Introduction

The U.S. is seeing a rise in affective conditions among young adults (Kessler et al.,

2012; Terlizzi and Villarroel, 2020). The most recent national mental health statistics show that

young adults (ages 18-25 years) have a greater prevalence of anxiety and mood conditions but

seek less treatment than older and middle-aged adults (ages 50 and older) (SAMHSA, 2022).

Prior studies have revealed maturational changes in 'emotionality' and emotion processing that

may contribute to a differential vulnerability of young individuals to experience an affective

condition. Compared to older individuals, young adults cite greater negative emotional

experiences, show less emotional control (Gross et al., 1997) and stability (Carstensen et al.,

2011), have a greater tendency to report negative mood states (Machado et al., 2019), more

worries (Brenes, 2006), and a greater recall of negative memories (Charles et al., 2003).

Neuroimaging studies have determined the major components of the neural circuitry involved in

the differential processing of negative valence stimuli between young and older adults. The

dorsal and rostral subregions of anterior cingulate cortex (ACg) show weaker responses to

negative stimuli in old versus young adults (Chaudhary et al., 2023). Functional magnetic

resonance imaging (fMRI) studies employing face perception tasks indicate that the amygdala

has greater blood oxygenation level dependent (BOLD) response to negative emotion stimuli in

young compared older adults (Fischer et al., 2005; Iidaka et al., 2002; Tessitore et al., 2005).

Other studies, however, observed either no age associated differences in the amygdala

response to negative stimuli (Wright et al., 2006) or report an increased amygdala functional

connectivity from adolescence to adulthood (Ravindranath et al., 2020). Recent evidence

suggests that the neurobiological adaptations associated with adulthood variations in affective

behaviors likely involves a larger number of brain areas outside the amygdala and changes in

neural interactions at a network level (Malezieux et al., 2023; Wang et al., 2023).

Maturational changes in affective behaviors in early versus late adulthood have been

reported in rodents (Baumgartner et al., 2023; Flurkey et al., 2007; Shoji and Miyakawa, 2019).

Young (2–3-month-old; mo) C57BL/6J male mice display greater anxiety-like behaviors

compared to older adult, late middle age and aged mice (Ammassari-Teule et al., 1994; Shoji

and Miyakawa, 2019). Although this remains unclear, since other studies provide evidence of

greater anxiety-like behaviors in aged compared to young adult mice (Baumgartner et al., 2023;

Shoji and Miyakawa, 2019). Nevertheless, the expression of fear-related associative memory

(Evans et al., 2021; Ferreira et al., 2023; Moyer and Brown, 2006; von Bohlen und Halbach et

al., 2006), reward sensitivity (Ammassari-Teule et al., 1994; Malatynska et al., 2012), sociability

and social recognition (Shoji and Miyakawa, 2019), and learned helplessness on a Porsolt test

(Ferreira et al., 2023) are all observed to vary in young versus older adulthood mice. In parallel

with these affective behavior findings, there is recent evidence of brain structural differences

between early and late adulthood (Clifford et al., 2023). In an analysis of structural covariance

network data, it was shown that young mice have a higher modularity with greater node degree

in anterior cingulate cortex and lower in hippocampal formation than older adult mice (Clifford et

al., 2023). This age-related difference in structural network modularity was linked to cognitive

differences (Clifford et al., 2023). It remains unclear whether functional network topology also

differs in a similar manner to the reported structural changes in young versus older adult mice.

Moreover, the relationship between functional network activity as measured by functional

magnetic resonance imaging (fMRI) and affective behaviors in early and late adulthood warrants

investigation. In the present study, we measured affective behaviors involving anxiety-like open

field activity, reward sensitivity, sociability, and fear memory in young adult, older adult, and late

middle aged C57BL/6J male and female mice. We also evaluated the relationship between

behavioral measures and fMRI markers of network connectivity. Finally, we conducted

hippocampal and frontal cortical metabolomic analysis and observed several novel protein-

related substrates differing between young, older adult and late middle-aged mice.

                                     Materials and Methods

Subjects

        Male and female C57BL/6J (B6) mice aged 10, 30, or 60 weeks old (wo) at the time of

arrival were allowed a week of acclimation to the vivarium prior to the start of experiments

(N=30, total; n=10 per age group, 50% females; The Jackson Laboratory, Bar Harbor, ME).

Female mice initially weighed (in grams; mean ± standard deviation) 19±0.93, 32±4.6, 29±2.8

and males 26.7±2, 30±2.5, 46.8±5.6 (for 10, 30, 60wo, respectively). The selected age groups

represent 3 stages of adulthood. B6 mice at 3-6MO (12-24wo) are considered mature adults

and at 10-14MO (40-56wo) are considered middle age (Flurkey et al., 2007). The youngest

10wo group (2.5MO) was tested just before mature adulthood and the 60wo group was just at

the end of middle age (15MO). At 30wo (7.5MO), mice start to show early signs of senescence

(such as retirement from breeding at 8MO) and were included here as reference to compare

early adulthood and late middle age changes (Flurkey et al., 2007). Mice were housed in age-

and sex-matched groups of 3-4 in a temperature- and humidity-controlled room, inside

conventional air filtered cages (dimensions: 29 x 18 x 13 cm) with food and water available *ad*

libitum (vivarium lights on from 07:00-19:00 hours). All procedures received prior approval from

the Institutional Animal Care and Use Committee of the University of Florida and follow all

applicable NIH guidelines.

Experimental design

        Figure 1 shows the experimental timeline. All behavioral tests and functional

neuroimaging were conducted over an 8-week period. Behavioral tests included locomotor

activity in a novel environment, sociability and social recognition tests, sucrose preference tests,

and cued fear conditioning (**Figure 1**). Functional magnetic resonance imaging (fMRI) was

conducted a week after sucrose preference testing and two weeks before the final behavioral

test (fear conditioning). At the end of the experiments, mice were deeply anesthetized, and brain

tissue harvested for the metabolomic assay of excised frontal cortex and hippocampus tissue.

Imaging was thus carried out when mice were 14, 34 and 63wo and these were euthanized 8

weeks after arrival, at 18, 38 and 68wo. However, to simplify presentation of the results, we

keep group names in figures as 10wo, 30wo and 60wo.

Locomotor activity in a novel environment

An automated rodent activity cage system (Accuscan Instruments, Columbus, OH) was

used to measure spontaneous locomotor activity in a novel environment. The day before

locomotor measurements, mice were acclimatized for 1 hour to the testing room. The activity

monitors are made of clear Plexiglas (40 cm^3) with an outer steel frame containing infrared

emitting and receiving panels. Each panel has 16 equally spaced (2.5 cm) infrared beams

crossing the length and width of the cages at a height of 2 cm (horizontal beams). Several

patterns of sequential and repetitive infrared beam interruptions were registered and sent to a

programmable counter, which sorted beam counts and sent sorted data to a PC running

Versamax™ software. The sorted activity counts were classified as horizontal (forward

locomotion), repetitive (stereotyped activity), total, center and margin distances travelled (in cm).

Activity was measured for 30 minutes divided into six bins of 5 minutes each.

Sucrose preference test

A decrease in preference to consume a 2% sucrose solution over water suggests

anhedonia (D'Aquila et al., 1997). 'Leakproof' bottles (100 ml, graduated by 1 ml) with stoppers

and sipper tubes were purchased (Braintree Scientific Inc, Braintree, MA; model no. WTRBTL).

Filtered home cage lids with two bottle openings (Allentown, Inc., Allentown, NJ), were used for

two bottle fluid delivery in mouse home cages. Mice were temporarily single housed for testing

and once testing concluded they were returned to their original group cages. First, mice were

acclimatized to bottles filled with 2% sucrose solution for 3 days (96 h). On each of the 3 days,

bottles were weighed (in grams) to monitor fluid intake and replaced with a full bottle. The 96 h

acclimation period was followed by a single 24 h two-bottle choice test in which one bottle was

filled with 2% sucrose solution and the other with water. The side of the cage lid where the

sucrose and water filled bottles were located was counterbalanced across animals. Water and

sucrose consumed was measured after a 24 h period (McCall et al., 2013). Bottles were

weighted and values converted from grams to ml. An index (S_I) of how much sucrose (s)

consumed versus water (w) was calculated as: $S_i = \frac{s-w}{s+w}$.

Sociability and social recognition test

We used a modified version of a previously reported sociability and social recognition

test protocol (Tan et al., 2019). Test arenas were made of clear Plexiglas, divided by walls into 3

equal sized quadrants (arena total length x width x height dimensions: 60 x 45 x 20 cm). The

middle quadrant is the neutral zone, and the side quadrants (left and right) was where 'bait'

mice are placed for sociability and recognition testing. Each side quadrant can be entered

through a central door in the dividing walls. A pair of clear plastic 'bait' cups covered with

equally sized and equidistant perforations for air flow were inverted over the 'bait' mouse for

sociability testing. Test mice were acclimatized to the room and the neutral quadrant of the test

arena on 3 days, 15 minutes a day. During sociability testing, a novel (unfamiliar) mouse (an

adult mouse of the same strain and sex) is placed under a cup and the entrance to the 'bait'

side of the arena is opened for exploration by the test mouse while the other side is kept sealed.

In preliminary tests, this encouraged mice to explore the unfamiliar 'bait' mouse more vigorously

during a first encounter. Mice were tested for 10 minutes for time spent exploring the unfamiliar

mouse. Upon completion of the first test, and after a 5-minute break to clean the arena, mice

were placed back into the arena in the presence of a novel mouse and the previously

encountered (now familiar) mouse and re-tested for 10 more minutes with both sides opened for

exploratoration. In the first test, the degree of exploration of an unfamiliar mouse ("novel 1") is

considered here as a measure of the degree of sociability. In the second test, less exploration of

the familiar mouse ("familiar 1") relative to the new unfamiliar mouse ("novel 2") is a sign of

social recognition (Crawley et al., 2007). Behaviors were recorded using a digital camera

directly connected via FireWire to a Linux computer. We used the '*dvgrab*' tool from the Unix

command terminal to capture videos remotely via Firewire and program video capture settings.

Behavioral analyses were carried out using DeepLabCut (DLC, version 2.2.1) (Mathis et

al., 2018; Nath et al., 2019). We labeled 120 frames of mouse trajectories within the test cages

taken from 6 different videos and 95% of those labels were used to train a machine learning

algorithm. A ResNet-50-based neural network with default parameters for 2 training iterations

was used for training. Data (digital videos) were split into training and test datasets. The training

dataset was used to train a DLC model. Conversely, the test dataset was used to evaluate the

performance of the model. We validated the DLC model with 7 train-test-split shuffles. The

model with the smallest mean average Euclidean error (MAE) was selected. Before model

selection, we applied a p-value cutoff of 0.6 to exclude model predictions with a likelihood under

0.6. The final selected model had a MAE of 2.9 pixels. This neural network was used to analyze

videos from similar experimental settings to generate trajectory data. Using the R packages

'trajr' (McLean and Volponi, 2018) and 'DLCanalyzer' (Sturman et al., 2020), spurious

trajectories were removed before analyses. The videos were analyzed in two parts, the first

sociability test (T1) and the second social recognition test (T2). The immediate area surrounding

and including the mouse 'bait' cup on one side of the arena was designated 'ROI A' and the

other 'ROI B' (region of interest A and B, respectively). A python script was compiled to

generate the number of entrances, the velocity (in pixels per frame or ppf), and the duration of

initial approach and time spent within the ROIs.

Classical fear conditioning

Fear conditioning was conducted two weeks following the fMRI studies (**Figure 1**). The

fear conditioning cages were housed in sound attenuation chambers. The chambers had

internal computer-controlled lighting, white noise generator, a fan, and control units for distinct

aspects of the fear conditioning assay. These included audio speaker controls for tone

generation, shock controllers for the internal cage floors, and infrared and white lighting

controllers. The internal operant cage was made of translucent Plexiglas with a steel frame and

steel shock grid floor connected to a computer controlled current stimulator (Med-Associates,

Inc. St. Albans, VT). Activity within the fear conditioning cages was captured by a monochrome

video camera and video recordings were optimized via externally controlled near infrared (NIR),

white light and sound controllers. All cage accessories were controlled by PC running Video

Freeze Software^{TM}. Camera lens brightness, gain, and shutter settings were verified and

adjusted before collecting data and were kept constant in all mice. For each session, the

detected motion-sensitive signals were calibrated with the mouse inside the cage and NIR lights

on. The estimated motion index was set to the same threshold level (10 a.u.) in all mice prior to

data exporting or analyses. Videos were acquired at 30 frames per second (fps) and the

minimum freeze duration was 30 frames. For training on the first day, a 180 second baseline

was followed by four presentations of a 20 second tone (90 dB) ending with the presentation of

moderate level current to the entire grid floor for 1 second (0.9mA) (inescapable shock). Each of

the four tone-shock presentations were separated by 190 seconds and the entire session lasted

17 minutes. On days 2 and 3 (recall tests at 24 and 48 h of training day 1, respectively), the

same protocol was carried out but in the absence of a shock. On day 3 (modified context test),

the cage environment was modified by placing a plastic smooth white floor and diagonal walls,

which covered the grid floor, light, and speaker. For each of the 4 tone-shock pairings, a 20

second pre-shock tone interval and a 40 second post-shock interval were used to quantify

number of freezing bouts (freezing counts) in a total 60 second epoch (per day there were four

60 second epochs summed to provide total freezing counts per day. Motion-index was also

used as a secondary measure of general activity inside the cage during the same 60 second

epochs/day.

Magnetic resonance imaging instrumentation

Images were collected on a magnetic resonance spectrometer tuned to 200.6 MHz

proton resonant frequency (4.7 Tesla/33 cm actively shielded magnet). The MR system was

equipped with Resonance Research Inc. spatial encoding gradients (RRI BFG-200/115-S14,

maximum gradient strength of 670 mT/m at 300 Amps and a 120 µs risetime; RRI, Billerica, MA)

and controlled by a VnmrJ 3.1 console (Agilent, Palo Alto, CA). A custom-built 8.8 cm x 11.6 cm

radiofrequency (RF) transmit coil used in combination with a 3 cm x 3.5 cm actively

decoupled/tunable RF receive quadrature surface coil (RF engineering lab, Advanced Magnetic

Resonance Imaging and Spectroscopy Facility, Gainesville, FL).

Mouse imaging setup

Mice were scanned sedated under a continuous paranasal flow of 0.25 % isoflurane gas

(delivered at 0.04L/min mixed with medical grade air containing 70% N_2 and 30% O_2, Airgas,

Inc.) and a continuous subcutaneous infusion of dexmedetomidine. Prior to setup, mice were

anesthetized under 2% isoflurane and administered an intraperitoneal (i.p.) injection of 0.1mg/kg

dexmedetomidine (at a 1ml/kg volume). Isoflurane was reduced to 0.25% throughout the

remaining imaging session. An infusion line was setup for subcutaneous delivery of

dexmedetomidine over the course of scanning (0.1mg/kg/ml at an infusion rate of 25 µl/h using

a PHD-Ultra microinfusion pump, Harvard Apparatus, Holliston, MA). Functional MRI scans

were collected at least 50 minutes after the i.p. injection. Respiratory rates were monitored

continuously, and body temperature was maintained at 36-37°C using a warm water

recirculation system (SA Instruments, Inc., New York).

Functional MRI acquisition protocol

We acquired one T2 weighted anatomical and an fMRI scan per mouse. The T2-weighted

fast spin echo (FSE) sequence was acquired with the following parameters: effective echo time

= 48 ms, repetition time (TR) = 2 seconds, echo train length = 8, number of averages = 12, field

of view (FOV) of 19.2 mm x 19.2 mm and 0.8 mm thick slice, and a data matrix of 192 x 192

(0.1 μm^2 in plane) and 12 interleaved ascending coronal (axial) slices covering the entire brain

from the rostral-most extent of the anterior frontal cortical surface, caudally towards the upper

brainstem and cerebellum (acquisition time 9 minutes 36 seconds). Functional images were

collected using a single-shot gradient echo planar imaging sequence with the following

parameters: TE = 22 ms, flip angle = 90˚, TR = 1.2 seconds, 600 repetitions, FOV = 19.2 x 19.2

mm and 0.8 mm thick slice, and a data matrix of 48 x 48 (0.4 μm^2 in plane) with 12 interleaved

ascending coronal slices in the same position as the anatomical scan. Ten dummy EPI scans

were run prior to acquiring data under steady state conditions. Respiratory rates, isoflurane and

dexmedetomidine delivery, body temperature, lighting, and room conditions were kept constant

across subjects.

Image pre-processing

The image processing workflow is illustrated in **Figure 2** (Colon-Perez et al., 2019;

Kotlarz et al., 2022; Sadaka et al., 2023; Sakthivel et al., 2023). Resting state processing was

carried out using Analysis of Functional NeuroImages (AFNI) (Cox, 1996), FMRIB Software

Library (FSL) (Jenkinson et al., 2002), and Advanced Normalization Tools (ANTs) (Klein et al.,

2009). Binary masks for anatomical and functional scans were created using ITKSNAP

(Yushkevich et al., 2006). The brain binary masks were used for brain extraction prior to

registration steps. Times series spikes were removed (3dDespike, AFNI), image repetition

frames aligned to the first time series volume (3dvolreg, AFNI), and detrended (high pass filter

<0.009 Hz using AFNI 3dTproject). Independent component analysis (ICA) using Multivariate

Exploratory Optimized Decomposition into Independent Components (FSL MELODIC version

3.0) was used to assess structured 'noise' or artefact components in each subject scan, in their

native space. Most, if not all ICA components in this first stage contained artefact signal voxels

along brain edges, in ventricles, and large vessel regions. These components were suppressed

using a soft ('non-aggressive') regression approach, as implemented in FMRIB Software Library

(FSL 6.0.3) using fsl_regfilt (Jenkinson et al., 2002). A low-pass filter (>0.12Hz) and spatial

smoothing (0.4mm FWHM) was next applied to the fMRI scans prior to registration steps. Post-

regression ICA was carried out to verify removal of artefact components and preliminary

assessment of putative ICA networks in individual scans.

Image template construction, subject normalization, and resting state signal extraction

Anatomical scans were cropped and bias field corrected (N4BiasFieldCorrection, ANTs).

Functional scans were cropped, and a temporal mean image was registered to the anatomical.

The functional-to-anatomical registration matrix was applied to all fMRI volumes. A multi-T2

anatomical template was generated using ANTs 'antsMultivariateTemplateConstruction2.sh'

script (cross correlation used as a similarity metric and a greedy SyN as transformation model).

Individual subject-to-template transformation matrices were applied to fMRI datasets. A group

ICA was applied to template-registered concatenated fMRI scans to generate a set of 60 nodes

for ROI-based intrinsic fMRI signal extraction, pairwise cross-correlation statistics, and

adjacency matrix construction. The ICA was set to an upper limit 60 component signal

decomposition and a mixture model threshold of 0.7. The resulting 60 ICA components included

peak signal (peak T statistic) nodes in midline frontal cortex (anterior cingulate), somatosensory

regions, temporal, visual, auditory cortices, entorhinal cortex, several hippocampal nodes,

subiculum, amygdala, thalamic nuclei, and striatum. The generated 60 nodes were each

binarized and assessed to ensure non-overlapping coverage of brain areas on the template and

registered scans (**Figure 2**). Both the template and ICA nodes were normalized to the Allen

Brain Institutes common coordinate framework version 3 (CCFv3) atlas of the B6 mouse brain

for 3D visualizations (Lein et al., 2007). Signal timeseries were extracted from preprocessed

fMRI scans with the assistance of 60 ROI mask overlays. This generated 60 individual ROI text

files per subject that contained L2-normalized resting state signals as a vector of 600 data

points.

The timeseries files were used in cross-correlations and in calculations of Pearson r

coefficients for every pairwise combinations of ROIs (using *corrcoef* function in MATLAB). The

resulting number of pairwise correlations was 1,710 per subject (after removing 60 self-

correlations). Correlation coefficients were Fisher's transform to ensure normality prior to

statistical analyses. The center voxel coordinates for the ICA-based nodes normalized to the

CCFv3 were used in 3D network visualizations using BrainNet viewer (Xia et al., 2013).

Network analysis

Weighted undirected matrices were analyzed using Brain Connectivity Toolbox (Rubinov

and Sporns, 2010), igraph (version 1.5.1.9005) (G. and Nepusz, 2006) in R, and tools in

MATLAB (Mathworks, Natick, MA). Global graph metrics were calculated for edge density

thresholds ranging from 6-32%. Global network measures for this density range were converted

to area under the curve (AUC) values prior to multivariate statistical assessments. Node-specific

(ROI) network measures were calculated at a 16% threshold, which in the present dataset

corresponds to a Pearson r of approximately 0.2 or more, an average degree of 9.1, and

network strength of 2.5-5.0 (**Supplemental Figure 1**). As a comparison, graph density based on

bootstrapped calculations of confidence intervals for significant Pearson correlations

corresponds to a significant r of ≥ 0.06, 40% density ($\delta \geq 0.4$), degree(k) ≥ 20, and strength ≥ 4

(**Supplemental Figure 1**).

A major objective was to assess maturational and aging changes in functional network

topology and determine links between network measures and spontaneous 'affective' behaviors.

To this end, we assessed several graph measures of network integration and communication

efficiency. These included node clustering coefficient (CC; number of interconnected node pairs

connected to a node) (Onnela et al., 2005), node strength (sum of edge weights/node) and

degree (sum of edges/node) and global measures such as transitivity (related to CC; number of

triad groups normalized by all possible triad nodes in a network), mean distance or

characteristic path length (CPL; the average edges or edge weights between node pairs), the

small world coefficient (SWC) (Humphries and Gurney, 2008; Humphries et al., 2006) and

small world measure (SWM) (Telesford et al., 2011). Global efficiency is the inverse of CPL. For

local node efficiency, a length matrix was used to calculate inverse values in vicinity of a node,

with added weights used to emphasize the highest efficiency node paths (Rubinov and Sporns,

2010; Wang et al., 2017). To corroborate results relative to random networks, all network

measures were calculated on original and randomized versions of the functional connectivity

matrices. Positive and negative edges were randomized by ~5 binary swaps with edge weights

re-sorted at each step. The approach preserves weights, degree, and strength distributions

(Rubinov and Sporns, 2010). We confirmed that the randomization approach preserved weights

per nodes, node degree and strength distributions (**Supplemental Figure 2**). Node strength and

degree distributions were similar between random and original networks, with distributions

having long tail at low density threshold and near Gaussian at high density (**Supplemental

Figure 2**). Degree and strength were approximately preserved across nodes, although the

specific organization of edge weights differed when viewed on 3D brains (**Supplemental Figure

2**). The positive and negative matrix strength sequence accuracies for input and output matrices

were r = 0.99 and r = 0.98. SWC was calculated as the ratio of gamma to lambda coefficients,

where gamma is the ratio of real to random network CC, and lambda the ratio of real to random

network CPL (Humphries and Gurney, 2008). SWM was calculated with an artificial non-ring

lattice network as reference, as previously reported (Telesford et al., 2011).

A probabilistic approach for community detection was used to calculate a modularity

statistic (Q), which indexes the rate of intra-group connections versus connections due to

chance (Blondel et al., 2008b). The procedure starts with a random grouping of nodes and

iteratively moving nodes into groups which maximize the value of Q. The final number of

modules and node assignments to each group (e.g., community affiliation assignments) was

taken as the median of 100 iterations of the modularity maximization procedure (Pompilus et al.,

2020). The number of communities per mouse brain functional network and population size of

each community were also assessed. We evaluated other community detection routines in

igraph as a comparison to results obtained with the implementation of Louvain's algorithm in

Brain Connectivity Toolbox (results in **Table 1**). The role each node plays within their assigned

module can vary according to their participation coefficient. The participation coefficient is

calculated as: $P = 1 - \sum_{m_i}^{m_n} \left(\frac{d_{im}}{d_i}\right)^2$, where d_{im} is the within-module node degree and d_i is

the overall network degree for a node.

We analyzed the tendency of assortative vs dissortative mixing of nodes (Newman,

2002). The assortativity index is a correlation coefficient comparing node strength values

between pairs of edge-connected nodes. Positive r values indicate connectivity between pairs of

nodes with similar strengths (e.g., high strength nodes pairs with high and low with low), while

negative r values indicate cross-pairings between low and high strength nodes. We generated

and analyzed weighted rich club coefficient curves (Φ) at an edge density of 16% (Colizza et al.,

2006). Nodes belonging to a rich club subnetwork have an above-chance tendency to tightly

connect with each other and share the bulk of connections within the network (Colizza et al.,

2006; van den Heuvel and Sporns, 2011). The approach creates subgraphs containing nodes

with strength values at a predetermined degree value, *k*. For each k-level subgraph, Φ_w is

calculated as the ratio of the sum of edge weights that connect nodes of degree k or higher

($W_{>k}$) to the total possible number of summed edge-weights (van den Heuvel and Sporns,

2011). Rich club coefficient values were calculated for original and randomized networks and a

normalized rich club index analyzed (i.e., ($\Phi norm = \Phi orig/\Phi rand$). Functional connectivity

networks were visualized in BrainNet viewer (Xia et al., 2013). Center coordinates for each node

were based on CCFv3 parcellations, as indicated above. Several 3D mouse brain connectome

maps were generated in which size of nodes (spheres) was scaled according to AUC values for

several node-centric measures at various thresholds. Additional 3D maps were generated with

node colors representing the module assignment (e.g., community affiliation vector) and node

size weighted by modularity index.

Sample Preparation for Gas Chromatography-Mass Spectrometry (GC-MS)

Fresh brain tissue was extracted, and two isolated regions were weighed, and flash

frozen for metabolomics (**Supplemental Figure 3**). A 1mm thick slice of the prefrontal cortex

was manually dissected from approximately Begma anterior-posterior -2.96mm to -1.94mm

(Paxinos and Franklin, 2001), and a 2mm wide midline cut including from the median -/+ 1mm

medial lateral structures -3 to -3.5 mm deep. The dissected regions thus included areas of the

infralimbic, prelimbic, medial orbital, anterior cingulate and midline primary motor cortices. The

dorsal hippocampus was dissected at -1.46 mm rostrally to about -2.46 mm caudally (Bregma

coordinates) and stripped of its overlying cortex and white matter.

Mice brain tissue (20-25 mg) was extracted with 1 mL of Acetonitrile:Isopropanol:Water

(3:3:2, $v.v.v$), centrifuged for 25 min at 10,000× g. The supernatant was transferred into a new

tube and dried in a speedvac. The dried extract was resuspended in 0.5 mL of

Acetonitrile:Water (1:1, $v.v$) and centrifuged for 15 min and the supernatant was transferred into

reaction vial and dried down under nitrogen. Fifty microliters of 2% MOX reagent were added to

the vial and incubated at 30 °C for 90 min followed by addition of 50μL of MTBSTFA +TBDMS

and incubation at 60 °C for 1 h. Finally, reaction solution was transferred into GC vial for GC-MS

analysis (Mahar et al., 2021).

GC-MS analysis

396   GC-MS data was recorded with Single Quadrupole Mass Spectrometer and GC (Trace

397 1310) (Thermo Fisher Scientific, USA). The column was 30 m long dimethyl (95%)/diphenyl

398 polysiloxane (5%) RTX-5MS with 0.25 mm ID, and guard column (10 m) (Restek). The initial

399 temperature of GC oven was 60 °C for 60 seconds and the temperature increased to 325 °C

400 (@10 °C/min). Metabolites were identified using NIST library with Xcalibur software.

Statistical analysis

Statistical analyses used MATLAB or the R stats package. Unless otherwise noted, all

analyses involved either a two-way full factorial analysis of variance (ANOVA) without or with

repeated measures, depending on the specific test (critical $\alpha < 0.05$). Post hoc tests used

Tukey's honest significant difference (HSD) procedure. Where appropriate, false discovery rate

(FDR) correction was used. The resultant statistical test parameters are indicated in figure

legends or in the results section. Initial analyses revealed only trends for sex. Male and female

data were therefore pooled to increase statistical power. The dependent variables in locomotor

tests were horizontal activity, stereotyped activity, total, margin, and center distances.

Locomotor activity data collected in 5-minute bins over 30 minutes were summed prior to

analysis. Dependent variables for sucrose preference tests are fluid intake consumed in 24 h

(ml) and sucrose preference index. For social interaction, the dependent variables are number

of ROI entrances, average speed in ROI (in pixels per frame, or ppf), time spent in ROI, and

time of initial approach towards ROI. For cued fear conditioning, the dependent variable is

freezing counts summed over four 60 second trial bins per day.

Global network measures (i.e., SW, modularity, CPL, efficiency, transitivity, assortativity)

were analyzed first as AUC values. Measures with significant differences were again analyzed

across density thresholds to determine the threshold level where significant differences are

observed. Upon determining the significance of modularity and assortativity, we analyzed the

number of modules per group, number of nodes per module, and the normalized rich club index.

All single threshold nodal analyses were carried out for a network density threshold of 16%.

These included node degree, node strength, participation coefficient, clustering coefficient, and

local efficiency. These were corrected for multiple comparisons across 60 nodes using the

Benjamini-Hochberg FDR procedure.

Additional exploratory analyses were carried out on several variables to assess novel links

between network measures and affective behavioral measures. For this, we used 9 variables,

including AUC values for SW, modularity, CPL, transitivity, assortativity, freezing counts on days

1-3 and margin distance. This data set provided 9 variables with 30 observations each. Factor

analysis was first used to explore associations between variables of most importance in the data

set (i.e., evaluate variable loadings for the first two factors contributing most to the variance in

the data). Projections of variables and data points onto the first 2 dimensions was carried out

with principal components analysis (PCA) and cluster analysis using FactoMineR (Lê et al.,

2008). A multivariate ANOVA (MANOVA) was used to examine overall differences and report

specific univariate associations of potential future interest ($\alpha < 0.05$).

Further analysis of fMRI scans was carried out using probabilistic ICA (Beckmann and

Smith, 2004) as implemented in MELODIC Version 3.15, part of FSL. The following data pre-

processing was applied to the input data: masking of non-brain voxels; voxel-wise de-meaning

of the data; normalization of the voxel-wise variance. Pre-processed data were whitened and

projected into a 20-dimensional subspace using PCA. The whitened observations were

decomposed into sets of vectors which describe signal variation across the temporal domain

(time-courses), the session/subject domain and across the spatial domain (maps) by optimizing

for non-Gaussian spatial source distributions using a fixed-point iteration technique (Hyvarinen,

1999). Estimated component maps were divided by the standard deviation of the residual noise

and thresholded (0.6) by fitting a mixture model to the histogram of intensity values (Beckmann

and Smith, 2005). The resulting components were overlaid on the multi-subject template and

classified according to peak z score anatomical location. A series of two-stage multiple linear

regressions were used to back-project the group ICA components to subject-specific time

courses (using spatial regressions) and spatial components (using temporal regressions). The

subject-specific spatial component maps were then used in statistical comparisons for each

component in the FSL randomise tool. The statistical design matrix was generated using the

FSL Glm tool. We used a one-way ANOVA general linear model design. Randomization tests

were carried out with 500 permutations (corrected p-level for significance is 0.05 and statistical

thresholding of maps by threshold free cluster enhancement).

For metabolomics data, the intensity of metabolites was imported to the web-based

MetaboAnalyst 5.0 platform (https://www.metaboanalyst.ca/) and normalized by the sum of

intensities followed by generalized log10 transformation and pareto scaling before multivariate

analysis. Further, principal component analysis (PCA), partial least square discriminant analysis

(PLS-DA), and Hierarchical Clustering analysis were employed on the dataset (Pang et al.,

2021). The Q^2 and R^2 values were calculated to assess supervised PLS-DA model. The

Variable Importance in Projection (VIP) derived from PLS-DA was used to extract the

importance of metabolites to PLS-DA model. Metabolites with a VIP score>1 is weighed to be

influential in PLS-DA.

 Results

 Young 10wo mice display greater anxiety-like activity in a novel open field environment

 compared to adult 30wo and older middle aged 60wo mice.

Mice were tested in a novel open field environment one week after arrival. The ages at

the time of testing were 11, 31, and 61wo. The 10wo group showed a significantly greater total

and margin distance than 30 and 60wo mice (**Figure 3**). The margin distance result is indicative

of high levels of movement along the walls of the test cage. No significant age or sex

differences were observed in other locomotor activity measures, such as horizontal, stereotyped

activities and center distance.

No age differences in sociability, recognition, and reward sensitivity

Sociability tests were carried out when mice were 12, 32, and 62wo. No age differences

in the number of ROI entrances and amount of time spent near an unfamiliar mouse were

observed in the first test for sociability (T1) (**Supplemental Figure 4**). All animals showed a

reduced number of entrances and time spent approaching the familiar mouse in test 2 (T2), and

they showed an increase in these measures towards a novel unfamiliar mouse. No significant

differences were observed between age groups. Sucrose preference testing was conducted a

week later when mice were 13, 33, and 63wo. All mice consumed more sucrose solution than

water. No age-related differences in amount of sucrose and water consumed nor differences in

the sucrose preference index were found (**Supplemental Figure 5**). No sex differences were

observed in social interactions and sucrose preference. We also analyzed sucrose consumption

during the 3 days of acclimation and observed no differences between the groups

(**Supplemental Figure 5**).

Young adult mice develop stronger fear memory than older adult mice

Fear conditioning was conducted a week after completing fMRI scans when mice were

16, 36, and 66wo. On day 1 of fear conditioning, 10wo and 30wo mice progressively developed

unconditioned-conditioned stimulus (US+/CS+) associations (**Figure 4**). This is observed as a

gradual rise in freezing counts over the course of each tone-shock exposure across 4 trials

(**Figure 4A**). 60wo mice showed highly variable freezing responses, with high amounts of

freezing counts across all 4 cue-shock trials. Data were next analyzed as average freezing

counts per day (**Figure 4B**). When compared across days, 10wo mice had a robust recall of

fear conditioning, observed as high levels of freezing in response to the tone alone, at 24 and

48 h. At 48 h, the cages were modified such that the tone was delivered in a context with

different sensory features (walls and floors were smooth white plastic in a triangular

configuration, as reported in (Anagnostaras et al., 2010). Still, 10wo mice maintained high levels

of freezing in response to the conditioned cue. The 30wo mice showed a trend towards

increased fear recall at 24 h, which was significantly higher on day 3 vs day 1 under modified

context conditions. The 60wo group showed no evidence of fear conditioning nor recall on any

of the test days (**Figure 4B**). When compared across groups, 10wo mice showed higher

freezing levels during recall on days 2 and 3 compared to the two older age groups. We

analyzed general movement activity using an infrared motion sensitive camera inside the test

cages. Consistent with results in **Figure 3**, 10wo mice showed higher motor activity in the novel

test cage on day 1 of testing than the other two age groups (**Figure 4C**). Activity levels dropped

significantly in all groups by days 2 and 3 of testing. As with the freezing levels during recall, the

activity levels of 60wo mice were highly variable on day 1 of testing. Activity in 60wo mice was

significantly less on day 1 than 10wo mice.

Young adult mice had a more robust network modularity compared to older adult mice

Functional MRI scanning was carried out over a two-week period following sucrose

preference testing (**Figure 1**). Mice were 14-15, 34-35, and 64-65wo at the time of scanning.

Analysis of network measures obtained from functional connectivity matrices indicated

significant main effects of age, but not sex, for SWC, modularity, and assortativity (**Figure 5**).

Tukey's HSD post hoc comparisons further revealed that SWI was highest in 30wo mice relative

to 10wo mice and both modularity and assortativity were highest in 10wo mice relative to 30 and

60wo mice (**Figure 5**). Significant pairwise differences were observed across several edge

density thresholds (see **Figure 5C-5E**; details in figure legend). We analyzed the number of

modules (at 16% edge density) per age group and observed a significantly greater number of

modules in 30 vs 10wo mice (**Figure 5A** and **5F**). Network randomization removed these

differences (**Figure 5B**). The number of nodes per module did not differ between groups

(**Figure 5G**). To confirm the significant maturational changes in functional network modularity,

we further investigated 5 additional community detection routines available in igraph, including

greedy optimization (Clauset et al., 2004), edge betweenness (Newman and Girvan, 2004),

random walk (Pons and Latapy, 2005), lead Eigevector (Newman, 2006), and a different

implementation of Louvain's algorithm (Blondel et al., 2008a) (**Table 1**). These were all

estimated at 16% edge density. Across most of these algorithms, the number of modules

detected was lowest in 10wo mice compared to the other two age groups. In addition, the

modularity index was highest in 10wo mice compared to the two older age groups (**Table 1**).

At 16% edge density, we also observed significant main effects of age on transitivity and

assortativity (**Table 1**). Transitivity was lowest in 60wo vs 30 and 10 wo mice. Assortativity was

significantly greater in 10wo vs 30wo mice. This was observed using Brain Connectivity

Toolbox's routine for calculating network Assortativity. Although a similar trend was observed

using igraph, it did not reach statistical significance (**Table 1**).

Young adult mice have robust functional connectivity between affective brain regions in

auditory/temporal cortical and striatal networks

Group ICA identified 9 of 20 functional network components similar to previously

identified in mice (Bukhari et al., 2017; Grandjean et al., 2014a; Jonckers et al., 2011; Sforazzini

et al., 2014). The ICA components included anterior and posterior cingulate, somatosensory,

visual, auditory/temporal, piriform/amygdala, hypothalamic and dorsal striatal networks

(**Supplemental Figure 6**). **Figure 6** shows the results for statistical comparisons between age

groups. The 10wo mice had greater functional connectivity in the auditory/temporal network

than 60wo mice. The areas showing greater connectivity within the right hemisphere of this

network in 10wo mice included anterior cingulate, auditory/temporal subregions, ventral

hippocampus, dorsal striatum, midbrain reticular nucleus, bed nucleus of stria terminalis,

posterior thalamic areas, and superior colliculus. The left hemisphere had less differences and

included primary and secondary somatosensory areas, striatum, visual cortex, hippocampus,

and thalamic subregions (**Figure 6**). The left hemisphere striatal network had greater functional

connectivity in 10wo compared to 60wo mice. The greater functional connectivity in the left

striatal network of 10wo mice included piriform, hippocampus and superior colliculus. We also

observed greater functional connectivity in a left hemisphere striatal network in 30wo compared

to 60wo mice and in a right hemisphere somatosensory network in 10wo compared to 30wo

mice.

Significant statistical relationship between high fear and anxiety trait with high indices of

network communication in 10mo mice

MANOVA revealed a significant effect of age across the full range of tested connectomic

and affective behavior variables ($F_{9,20}$=4.6, p=0.002). Principal components analysis (PCA)

revealed a relatively strong relationship between several of these variables. This is observed by

contributions to the first 2 dimensions, which jointly explain 56.08% of variance in the data. To

establish significance, PCA results for the first 2 dimensions (56.08%) were compared against a

reference value (43.65%) representing 95% quartiles of distribution of percent explained

variances (inertia) of simulated data tables of equivalent size and normal distribution (**Figure 7**).

The positive direction along dimension 1 opposes 90% of mice in the youngest group (1-8 and

10 are in the youngest 10mo group) from many of the mice in the other two age groups (**Figure**

5A). This positive direction is characterized by high values freezing counts on days 2-3 of fear

conditioning and high margin distance activity in a novel open field environment and high values

for small world coefficient, transitivity, and modularity in 70% of the 10mo mice (**Figure 7B**).

Cluster dendrogram in **Figure 7C** classified 3 clusters. A majority of young 10mo mice were

classified in cluster 3 (green). This group is again characterized by high values for modularity,

transitivity, small world coefficient, margin distance activity and days 2-3 of fear conditioning.

Metabolomic profiles in frontal cortex and hippocampus differ between 10, 30 and 60wo

mice

Results for PCA and partial least squares discriminant analysis (PLS-DA) for GC-MS

data obtained from the mouse hippocampus are shown in **Supplemental Figure 7** and **Figure**

8 (panels A and B). Variable importance in projection (VIP) scores plot of the PLS-DA model for

mouse hippocampus indicated differences in the top 25 metabolites between 2.5, 7.5, and 15

mo mice. Among the top 25 listed in **Figure 8** (panel A), methionine and cysteine had the

highest VIP scores. The top 7 had a bidirectional transition in relative levels when youngest

mice having lowest or highest levels and late middle-aged mice having diametrically distinct

levels. This relationship included methionine, cysteine, malate, fumarate, succinate, N-

acetylglycine, alanine, and taurine. The differential prominence of these metabolites across age

groups is supported by clustering in heatmap shown in **Figure 8** (panel B). Results for PCA and

PLS-DA for GC-MS data obtained from the mouse midline frontal cortex are shown in

Supplemental Figure 8 and **Figure 8** (panels C and D). VIP scores plot of the PLS-DA model

for mouse frontal indicated differences in the top 25 metabolites between 2.5, 7.5, and 15 mo

mice. Among the top 25 listed in **Figure 8** (panel C), proline, taurine, and methionine had the

highest VIP scores. The differential prominence of these metabolites across age groups is

supported by clustering in the heatmap in **Figure 8** (panel D).

592   **Discussion**

Our data indicates that young 10wo mice display greater levels of anxiety-like locomotor

behavior and develop robust fear conditioning compared to older adult and late middle-aged

mice (30-60wo). This was accompanied by greater functional connectivity between a temporal

cortical/auditory cortex network and subregions of the anterior cingulate cortex and ventral

hippocampus, among other regions, and a greater network modularity and assortative mixing of

nodes in young versus older adult mice. We observed a close correspondence between

behavioral measures of fear conditioning and anxiety-like locomotor activity and measures of

network efficiency and segregation in young mice, which was no longer apparent in older adult

mice. Finally, metabolome analyses indicated significantly different neurochemical profiles in

frontal cortex and hippocampus between 10wo mice and the other age groups, with several

essential amino acids differing the most between the groups (e.g., methionine, cysteine, proline,

and taurine). The results collectively reinforce the notion of high 'emotionality' in younger versus

older adults, an affective behavioral trait in mice that can be further interrogated in future studies

to determine neurobiological substrates conferring 'risk or resilience' to affective disorders.

In mice, conditioned fear memory varies with age in both context (hippocampal-

dependent) and cued (amygdala-dependent) paradigms. Compared to adult 6mo mice, late

middle age 17mo C57BL/6 mice show deficits in renewal of extinguished freezing in a modified

context (Sanders, 2011). Eighteen-month-old mice showed greater freezing in response to

context but lower freezing in response to tone than 3mo mice (Evans et al., 2021). These results

suggest dissociable aging changes in conditioned fear memory, perhaps involving distinct

neural circuits involving amygdala and hippocampus (Evans et al., 2021). Twenty three month

old aged mice showed higher cued fear compared to 2mo mice but impaired renewal of

conditioned freezing following extinction than 12mo mice (Oh et al., 2018). This was associated

with altered stress reactivity (Oh et al., 2018). Using a trace eye blinking conditioning protocol,

Kishimoto showed significant impairments in conditioned responses in aged 20mo compared to

12 and 2mo mice (Kishimoto et al., 2001). The above studies support differences in conditioned

fear in aged versus young mice (Moyer and Brown, 2006). Age differences in

hippocampal/prefrontal structure and function at the neuronal and synaptic level, stress

coping/stress molecular signaling changes and changes in neuroimmune activity are thought to

account for the observed age differences in conditioned freezing (Evans et al., 2021; Oh et al.,

2018; von Bohlen und Halbach et al., 2006). Viewed from this perspective, human studies

reporting lower amygdala or cingulate activation in response to negative valence stimuli in aged

versus young adults may be accounted for by deficits in these brain areas as well (as opposed

to a greater reactivity in young adults) (Chaudhary et al., 2023). Our present data reveals a

close correspondence between conditioned fear recall and modularity, transitivity, and small

world coefficient. This implies broader functional network involvement in age related changes in

fear conditioning. It is possible that functional interactions between ventral hippocampal and

frontal cortical regions, particularly anterior cingulate and/or prelimbic cortices, drive changes in

the network configurations observed in the present study.

Young 10wo mice displayed greater locomotor activity than 30 and 60wo mice,

consistent with evidence of decreasing locomotion with age (Elias et al., 1975). Male B6 mice

are generally more active at 1-3mo than at 7 or 13mo (Lhotellier and Cohen-Salmon, 1989;

Serradj and Jamon, 2007), although others have reported greater locomotor activity in 17mo vs

5 and 25 mo B6NIA mice (Frick et al., 2000), greater locomotor activity in 3-8mo versus 28 mo

B6NJ (Fahlstrom et al., 2011), or no difference between 2 and 18mo B6 mice (Barreto et al.,

2010). In our present observations, locomotor activity in the younger B6 group was largely along

the margins of a novel test cage. This suggests exploratory activity along the safer walls of the

cages, possibly due to high anxiety levels in 10wo mice. Age differences in spontaneous (non-

learned) anxiety-like behaviors in open field tests have been previously reported and supported

by additional experimental tests such as the EPM, light-dark box, Porsolt swim test, and others.

Number of open arm entries and exploratory activity in an open field are greater in 28mo

compared to 3mo mice (Fahlstrom et al., 2011). Under novel exploratory conditions, however,

24mo B6 mice were observed to display greater exploratory activity than 2mo mice (Ammassari-

Teule et al., 1994). In a comprehensive analysis of B6 motor and several affective behaviors,

Shoji and Miyakawa reported greater open field activity, lower light-dark transitions, and lower

total number of arm entries in 23mo versus 2mo mice (Shoji and Miyakawa, 2019). In sum, the

above results suggest differences in affective behaviors in early versus late adulthood. These

are observed as age differences in open field behaviors, performance on other assays such as

EPM and light-dark shuttling, and in fear conditioning. In most cases younger aged adult mice

develop stronger fear memories and also display anxiety like behaviors relative to older adult

mice. Open field activity in our present work was greatest in 10mo mice and specifically female

mice. Thus, sex differences also should be considered when assessing affective behaviors.

Our results showed no differences in sucrose preference between the studied adult age

groups. All mice consumed preferred sucrose solution over water to the same degree. This

result contrasts with previous reports. Eighteen-month-old (~72wo) C57BL/6N mice reportedly

show less sucrose reward preference than 3mo mice (Malatynska et al., 2012). This was

correlated with other signs of depressive-like behaviors in aged versus young mice (Malatynska

et al., 2012). Similar reductions in sucrose and milk reward preference in 13-15mo versus 3mo

C57BL/6 were reported (Harb et al., 2014). Conversely, aged 24mo Balb/c mice did not show

differences in sucrose preference, locomotor activity, immobility in a tail suspension test

compared to 4-6mo mice (Kelley et al., 2013). Discrepancies between studies may arise from

differences in the sucrose preference protocol, differences in mouse strains, and age groups.

We also examined social interactions in mice. No age-related differences were observed. This is

consistent with results from Shoji and Miyakawa, who reported similar levels of social

interactions in B6 mice ages 2-23mo. However, they reported reduced preference for novel

social interactions in 17 and 23mo relative to 11mo old mice. Although we observed no such

preferences using a supervised learning approach to assess social interactions, 60wo mice did

display slightly lower novelty preferences during a second novel social encounter relative to a

first novel encounter (Supplemental Figure 4A). This suggests habituation effects of multiple

social encounters in 60wo mice.

In addition to examining affective behavioral differences in young and old adult mice, an

objective of the present study was to assess resting state networks across the age groups. We

observed differences in functional connectivity within a temporal/auditory cortex network.

Specifically, anterior cingulate cortex and ventral hippocampal connectivity within this network

was strongest in 10wo compared to 60wo mice. Our results using functional MRI agree with

previous structural neuroimaging of mice along the same age range, which identified

maturational changes from young to middle age in auditory, cingulate, hippocampal formation,

and somatosensory (parietal) cortices (Clifford et al., 2023). The results in our present study

showing stronger temporal/auditory connectivity with the ACg and vHPC in the young age group

is consistent with the robust cued fear conditioning these mice developed compared to older

age groups. These structures have been studied for their roles in generalized fear and anxiety in

murine models (Ortiz et al., 2019). Increased synaptic activity in projections from ACg to vHPC

regulate the long-term generalization of fear memories (Bian et al., 2019). The vHPC in mice

contains neurons that show higher levels of neuronal activity and conditioned freezing behavior

in response to threat cues than safety or combined threat/safety cues (compound cues) (Meyer

et al., 2019). Conversely, a subset of prelimbic cortical projecting vHPC neurons show

increased neuronal activity with safety cues and this was accompanied by lower levels of

freezing relative to their response to threat cues (Meyer et al., 2019). Thus, the stronger

functional connectivity observed in the present study in young adult mice may be associated

with the fear and anxiety related functions of this vHPC-PFC circuit. Importantly, we observed

the higher functional connectivity between temporal/auditory and vHPC-ACg regions in young

versus older mice prior to their exposure to fear conditioning, thus averting effects introduced by

learned fear on functional connectivity. We should point out that the age differences in

functional connectivity between ACg, vHPC and other regions within the temporal/auditory

network (Figure 6) were lateralized, with a much greater functional connectivity observed in the

right versus left temporal cortical network. Fear-related anterior cingulate activity has been

previously reported to be lateralized in mice (Kim et al., 2012). Inhibition of the right but not left

ACg resulted in reduced observational fear learning relative to control mice (Kim et al., 2012).

Our results are consistent with this functional lateralization of ACg in fear related behaviors.

We also observed a greater functional connectivity within a left dorsal striatal network in

10wo and 30wo compared to 60wo mice. The regions observed to show significantly greater

functional connectivity with the dorsal striatum included a subregion of the HPC, superior

colliculus, bed nucleus of stria terminalis. The role of the dorsal striatum in regards to fear and

anxiety are supported by recent work indicating its synaptic connectivity arising from apical

intercalated cells in the amygdala, which regulate fear conditioning in mice via inputs from

thalamic, temporal association cortex and basolateral amygdala nuclei (Asede et al., 2021).

Finally, we should note that age differences between young (1-2mo) and older adult (5-19mo)

wildtype littermates of the transgenic Arcβ familial amyloidosis mouse have been reported

(Grandjean et al., 2014b). Young adult mice around 2 months of age had weaker functional

connectivity within subregions of the parietal, cingulate and dorsal striatal areas than older mice

at 5mo and older (Grandjean et al., 2014b). These results identify similar regions to the ones

reported here. However, whereas we observed decreases with age, Grandjean and colleagues

(Grandjean et al., 2014b) observed increases in control aging mice. While this may arguably be

due to differences in anesthesia protocol (isoflurane 1.4% in Grandjean *et al.* and 0.25%

isoflurane and dexmedetomidine used here), it is more likely that differences are associated

with the background strain of mice. Mouse strain-dependent genetics likely weighs heavily on

brain microstructure and function (Ashbrook et al., 2021; Murdy et al., 2022; Neuner et al.,

2019).

We applied graph theory derived network measures to evaluate age differences in

functional connectome topology. Our results indicate a higher functional network modularity and

assortative mixing of nodes of young versus older adult mice. Although the relevance to

underlying neurophysiology for such graph indices remains unclear, it reveals a segregated

pattern of connectivity of groups of nodes with high degree (high 'connectivity') nodes

connected to other similarly high degree nodes and low degree with low degree. Maturation of

the modularity of functional networks implies a 'diversification' of nodal interactions and this is

further supported by reductions in assortative mixing. The modularity changes we observed in

young versus older adult mice were of particular interest since it persisted across several

different modularity algorithms. Graph analysis of brain structural covariance matrices of 2, 10,

and 22mo mice showed reductions in node degree values from young to middle age in ACg,

auditory and somatosensory cortices and an increase in hippocampus (Clifford et al., 2023).

They also reported reduced modularity from 2 to 10mo (Clifford et al., 2023). In human subjects,

young and early middle age groups have higher local nodal efficiency, clustering coefficients,

and path lengths than late middle age and elderly groups (Wang et al., 2023). These graph

measures, which suggest changes in network efficiency from young adulthood through aging,

were considered in the context of cognitive changes and not affective behaviors. In the present

work, we observed statistically meaningful associations between the multiple affective

behavioral and connectomic variables. Although this does not confirm causation or even

correlation between the variables, the results do point to interactions of interest that should be

considered with other circuit specific approaches to determine the relationship between network

configurations and affective behaviors.

Finally, we observed age related changes in several amino acid substrates in frontal and

hippocampal tissue, which supports biochemical changes in these regions that might be linked

to the observed maturational changes in brain function and behavior. Each of the substrates

that differed the most between the groups play important roles in neuronal function and

development. Proline is important for glutamate homeostasis following methamphetamine

challenge (Jones et al., 2021). Taurine is involved in developmental changes to neurons in the

hippocampus, cerebellum, hypothalamus, and can act via GABA$_A$ receptors to modulate firing of

GABA neurons (Curran and Marczinski, 2017). Methionine is associated with a host of distinct

functions related to the control of gene expression in neurons and high methionine diets have

been linked to neuronal damage and accelerated signs of senescence in animal models (Pi et

al., 2021). The D enantiomer of cysteine is found in neuroprogenitor cells and is critical to early

brain development (Semenza et al., 2021). Taken together, these results suggest ongoing

maturational developmental changes in frontal cortex and hippocampus in young adult mice as

well as senescence-related neurochemical changes in late middle age mice.

761 **Figures with captions**

762 **Figure 1.** Experimental timeline. See text for details.

Figure 2. Image processing workflow. Anatomical scans from all subjects were used to generate a group-wise template for anatomical and functional image registrations and individual and group ICA-based signal processing steps. The decomposition of whole brain signals into 60 components was used to generate region of interest masks for signal extraction and pairwise correlations. The study specific node-set were normalized to the common coordinate framework version 3 mouse atlas for 3D visualizations of nodes and edges. See text for details.

Figure 3. Young 10wo mice showed greater activity in a novel test environment compared to 30

and 60wo mice. A-B) No differences observed for horizontal (forward locomotion) and

stereotyped (repetitive) activities between the age groups. C) Total distance (in cm) differed

between the age groups ($F_{2,29}$=5.1, p=0.01), with 10wo mice showing greater movement

distance relative to both 30 and 60wo mice (*p<0.05, Tukey's HSD test). D) Movement along

the margins of the novel test cages different between the age groups ($F_{2,29}$=7.6, p=0.003), with

10wo mice showing greater movement relative to both 30 and 60wo mice (*p<0.05, Tukey's

HSD test). The greater movement in 10wo mice is largely driven by female mice (sex main

effect: $F_{1,29}$=5.9, p=0.02; Tukey's HSD: p<0.05 for 10wo female compared to 30 and 60wo

female mice).

Figure 4. Young 10wo mice develop robust fear conditioning compared to 30 and 60wo mice.

A) Although all mice develop conditioned freezing by trials 3 and 4, the oldest age group

showed high levels of freezing in the first two conditioning trials. This is observed in 4 of 10 mice

(2 females and 2 males) with unusually high freezing counts in the first 2 trials (trial by age

interaction $F_{6,81}$ = 4.6 p = 0.0005; age effect $F_{2,27}$ = 3.2 p = 0.054; Tukey's HSD p<0.05

comparing 60wo to 10* and 30wo** mice on trials 1 and 2). B) Total freezing counts over 3 days

of testing (conditioning, recall, recall in modified context). A 3-way ANOVA with repeated

measures indicated that young 10wo mice had higher freezing counts during recall and in a

modified context than older adult 30 and 60wo mice (main effect of age $F_{4,48}$ = 5.5 p =0.0009,

Tukey's HSD *p < 0.05 10wo vs 30 and 60wo groups; main effect test day $F_{2,48}$ = 25.1 p <

0.00001, Tukey's HSD **p < 0.05 day 2 and 3 comparing 10wo vs 30 and 60wo groups). C)

Motion index (arbitrary units, a.u.) over 3 days of testing. A 3-way ANOVA with repeated

measures indicated that young 10wo mice had higher motion index values on day 1 than the

oldest age group (days by age interaction $F_{4,54}$ = 2.8 p =0.04, Tukey's HSD *p = 0.007 day 1

comparing 10wo vs the 60wo group; main effect test day $F_{2,54}$ = 84.9 p < 0.00001, Tukey's HSD

**p < 0.05 comparing days 1 vs 2 and 3 for all age groups).

Figure 5. Young 10wo mice had greater modularity and assortativity of functional connectivity

networks than 30 and 60wo mice. A) 3D functional connectome maps illustrating mean

modularity value (node size) and node community affiliation (node and edge color) for 10, 30,

and 60wo mice (density threshold = 16% edges retained). The bar below maps indicate number

of identified modules. B) Randomized 3D connectome maps for the same data in A. C) Small

world coefficient was lowest in 10wo mice (age by threshold interaction $F_{26,312}$ = 2.4 p = 0.0002,

Tukey's HSD * and *** p<0.05 comparing 10wo and 30wo to 60 weeks, respectively). D)

Modularity was highest in 10wo mice (age by threshold interaction $F_{26,312}$ = 1.79 p = 0.01,

Tukey's HSD * and ** p<0.05 comparing 10wo to 30wo and 60 weeks, respectively). E)

Assortativity was highest in 10wo mice (age by threshold interaction $F_{26,312}$ = 2.6 p = 0.00006,

Tukey's HSD * and ** p<0.05 comparing 10wo to 30wo and 60 weeks, respectively). F) Number

of modules identified at density-16% was higher in 30wo compared to 10wo mice (Kruskall-

Wallis test p = 0.02). G) No difference in number of nodes per module was observed. H) No

difference in the normalized rich club coefficient was observed.

Figure 6. Age associated reductions in functional connectivity in primary sensory and association cortical areas and striatum. Lines indicate regions of the brain with statistically significant differences in connectivity with the network component (indicated above maps). Statistical analysis used a one-way ANOVA with mixed effects (p<0.05, cluster corrected; higher level permutation tests with 5000 permutations). Maps thresholded at t >2.3 (bar graph below maps) and overlaid onto multi-subject T2 anatomical template. Titles over the maps indicate the original network component identified by group pICA (see methods for details). Column maps on the left correspond to CCFv3 atlas of the mouse brain. Abbreviations: ACg, anterior cingulate; AMYG, amygdala; APN, anteroposterior nucleus of thalamus; AUD, auditory cortex; BNST, bed nucleus of stria terminalis; CA3, 'Ammons Cornus' 3 subfield of hippocampus; DStr, dorsal striatum; HPC, hippocampus; LDT, laterodorsal thalamus; MDT, mediodorsal thalamus;

MRN, median Raphe nucleus; NAc, nucleus accumbens; PHA, posterior hypothalamic area; Pir,

piriform cortex; PfT, parafascicular thalamus; S2, secondary somatosensory cortex; Sept,

septum SCol, superior colliculus; SI, substantia innominate; SSC1, primary somatosensory

cortex; V1, primary visual cortex; VALT, ventral anterolateral thalamus; vHPC, ventral

hippocampus; VMT, ventromedial thalamus; VPMT, ventral posteromedial thalamus; vSub,

ventral subiculum.

Figure 7. Exploratory multivariate analyses reveal significant associations of interest between

functional network measures and affective behavior variables. The first two dimensions of the

PCA explain 56.08%, which is above 95%-quartiles of random distributions predicting 43.62% of

variance explain with equal size data tables. A) Individual factor map with labeled individuals

contributing to a high degree to the first two dimensions. B) Variables factor map with labeled

variables projected best on the first two dimensions. Data points along the positive dimension 1

axis have high values for modularity and marginal distance activity. Data points in dimension 2

with high CPL values have low values for fear conditioning recall. C) Cluster tree partitioning

data by ascending hierarchical classification of principal components. Cluster 3, made mostly of

individuals in the youngest age group, is characterized by high values for modularity, transitivity,

small worldness, marginal distance activity and fear recall.

Figure 8. Mouse brain metabolomics identifies age variations in methionine, cysteine, and proline in frontal cortical and hippocampal regions. A-B) Variable importance in projection (VIP) scores plot of the PLS-DA model for mouse hippocampus indicated differences in the top 25 metabolites between 10, 30, and 60wo mice). The differential prominence of these metabolites across age groups is supported by clustering in heatmap. C-D) VIP scores plot of the PLS-DA model for mouse frontal indicated differences in the top 25 metabolites between 10, 30, and

921   60wo mice. D) The differential prominence of these metabolites across age groups is supported

922   by clustering in the heatmap

Supplemental figures and legends

Supplemental Figure 1. Thresholding graph density at 16% retains significant edge weights

(Pearson r correlation coefficients). A) A graph density of 16% (δ=0.16) corresponds to a mean

node degree of 9.1, mean node strength >2.4 and <4.6, and a r ~0.15-0.42. B) Thresholding

graphs based on statistically significant Pearson r values (within 95% confidence bands) results

in r values between 0.06-0.07 (excluding 2 outliers in 7.5 MO group), density between 40%-

95%, node degree between 20-60 and node strength between 4-18.

Supplemental Figure 2. Randomization of functional connectivity networks results in

differential organization of connectivity between original and random graphs, while preserving

node weights, degree, and strength distributions. A-F) Histograms of node strength and degree

distributions at 3 density thresholds (δ=0.08, 0.16, 0.32). Relaxation of graph densities (δ=0.32)

adds 'spurious' correlations shaping 'random-like' distributions in histograms. G-H) Node

strength and degree across 60 nodes (at δ=0.16) shows preservation of values between original

and randomized networks. I) High correlation between input and output strength sequence

accuracies with the randomization of signed graphs (i.e., positive, and negative weights). J-K)

Node strength values (sphere sizes at each anatomical location) were preserved while

organization of edges in 3D maps in J and K differed. See text for further details.

Supplemental Figure 3. Schematic of GC-MS workflow.

Supplemental Figure 4. No differences in sociability and social recognition between age

groups. A) All mice showed similar number of entrances in the novel mouse ROI in the first test

(T1), reduced entrances during second exposure to the familiar mouse in a second test (T2) and

showed high levels of entrances in a second novel mouse ROI during the second test (T2)

(significant effect of time point, $F_{2,40}$=52.7 p<0.00001, but not age or sex). Guided asterisks

indicate significant differences at *p<0.05, ***p<0.001, and ****p<0.0001 over time points. B) No

significant differences observed in average speed inside the ROI. C) The 10wo mice showed a

significant effect of time ($F_{2,40}$=4.6 p=0.05, **p<0.01 Tukey's HSD) but no significant differences

with age or sex. D) No significant differences in time to initial approach. E-F) Video frames

capturing exploratory activity by the experimental mouse and novel and familiar mice under

inverted clear plastic 'bait' cups.

1000

1001

1002

1003

1004

1005

1006

Supplemental Figure 5. No difference in reward sensitivity between age groups. A) Sucrose

intake from the left-side cage bottle during the 3 days of acclimatization. B) Sucrose intake from

the right-side cage bottle during the 3 days of acclimatization. C) No effect of age or sex was

observed on amount of water and sucrose consumed in 24 h. A main effect of type of fluid

(sucrose vs water) was observed, with all mice consuming more sucrose solution than water

($F_{1,23}$ = 120 p< 0.0001; *p<0.05 Tukey's HSD). D) No difference in sucrose preference index

was observed.

Supplemental Figure 6. Identified functional connectivity networks from pICA (see methods for

details). Networks are qualitatively classified based on anatomical location of peak t statistic

voxels (threshold free cluster enhancement corrected). Classification names and component

order number are indicated overlying the statistical maps. Unilateral maps that are matched by a

separate, but mirror image contralateral component is merged into a single map and both

component numbers shown above maps. The right and left hemisphere sides are indicated at

1051   the bottom of the figure. Statistical maps overlaid onto multi-subject T2 anatomical template

1052   (t>2.3 shows significant voxels)

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

Supplemental Figure 7. 2D scores plots of (**A**) principal component analysis (PCA) and (**B**)

partial least squares discriminant analysis (PLS-DA) from GC-MS data of hippocampus region

of the mouse brain. The variation is displayed in parentheses on each axis of PCA and PLS-DA.

The shaded areas indicate 95% confidence regions. PLS-DA maximizes the separation among

groups using the group label and the optimal number of components required to build the PLS-

DA model was determined by cross validation. The robustness of the mathematical model was

determined by sum of squares captured (R^2) and the cross-validated R^2 (Q^2) parameters.

1079

1080

1081

1082

1083

1084

1085

Supplemental Figure 8. 2D scores plots of (**A**) principal component analysis (PCA) and (**B**)

partial least squares discriminant analysis (PLS-DA) from GC-MS data of prefrontal cortex

(PFC) region of the mouse brain. The variation is displayed in parentheses on each axis of PCA

and PLS-DA. The shaded areas indicate 95% confidence regions. PLS-DA maximizes the

separation among groups using the group label and the optimal number of components required

to build the PLS-DA model was determined by cross validation. The robustness of the

mathematical model was determined by sum of squares captured (R^2) and the cross-validated

R^2 (Q^2) parameters.

Bibliography

Ammassari-Teule, M., Fagioli, S., Rossi-Arnaud, C., 1994. Radial maze performance and open-
field behaviours in aged C57BL/6 mice: further evidence for preserved cognitive abilities during
senescence. Physiol Behav 55(2), 341-345.

Anagnostaras, S.G., Wood, S.C., Shuman, T., Cai, D.J., Leduc, A.D., Zurn, K.R., Zurn, J.B.,
Sage, J.R., Herrera, G.M., 2010. Automated assessment of pavlovian conditioned freezing and
shock reactivity in mice using the video freeze system. Frontiers in behavioral neuroscience 4.

Asede, D., Doddapaneni, D., Chavez, A., Okoh, J., Ali, S., Von-Walter, C., Bolton, M.M., 2021.
Apical intercalated cell cluster: A distinct sensory regulator in the amygdala. Cell Rep 35(7),
109151.

Ashbrook, D.G., Arends, D., Prins, P., Mulligan, M.K., Roy, S., Williams, E.G., Lutz, C.M.,
Valenzuela, A., Bohl, C.J., Ingels, J.F., McCarty, M.S., Centeno, A.G., Hager, R., Auwerx, J.,
Lu, L., Williams, R.W., 2021. A platform for experimental precision medicine: The extended BXD
mouse family. Cell Syst 12(3), 235-247 e239.

Barreto, G., Huang, T.T., Giffard, R.G., 2010. Age-related defects in sensorimotor activity,
spatial learning, and memory in C57BL/6 mice. J Neurosurg Anesthesiol 22(3), 214-219.

Baumgartner, N.E., Biraud, M.C., Lucas, E.K., 2023. Sex differences in socioemotional behavior
and changes in ventral hippocampal transcription across aging in C57Bl/6J mice. Neurobiol
Aging 130, 141-153.

Beckmann, C.F., Smith, S.M., 2004. Probabilistic independent component analysis for functional
magnetic resonance imaging. IEEE Trans Med Imaging 23(2), 137-152.

Beckmann, C.F., Smith, S.M., 2005. Tensorial extensions of independent component analysis
for multisubject FMRI analysis. NeuroImage 25(1), 294-311.

Bian, X.L., Qin, C., Cai, C.Y., Zhou, Y., Tao, Y., Lin, Y.H., Wu, H.Y., Chang, L., Luo, C.X., Zhu,
D.Y., 2019. Anterior Cingulate Cortex to Ventral Hippocampus Circuit Mediates Contextual Fear
Generalization. J Neurosci 39(29), 5728-5739.

Blondel, V.D., Guillaume, J.-P., Lambiotte, R., Lefebvre, E., 2008a. Fast unfolding of
communities in large networks. Journal of Statistical Mechanics: Theory and Experiment 2008,
2-12.

Blondel, V.D., Guillaume, J.L., Lambiotte, R., Lefebvre, E., 2008b. Fast unfolding of
communities in large networks. Journal of Statistical Mechanics: Theory and Experiment 1-12.

Brenes, G.A., 2006. Age differences in the presentation of anxiety. Aging Ment Health 10(3),
298-302.

Bukhari, Q., Schroeter, A., Cole, D.M., Rudin, M., 2017. Resting State fMRI in Mice Reveals
Anesthesia Specific Signatures of Brain Functional Networks and Their Interactions. Front
Neural Circuits 11, 5.

Carstensen, L.L., Turan, B., Scheibe, S., Ram, N., Ersner-Hershfield, H., Samanez-Larkin, G.R.,
Brooks, K.P., Nesselroade, J.R., 2011. Emotional experience improves with age: evidence
based on over 10 years of experience sampling. Psychol Aging 26(1), 21-33.

Charles, S.T., Mather, M., Carstensen, L.L., 2003. Aging and emotional memory: the forgettable
nature of negative images for older adults. J Exp Psychol Gen 132(2), 310-324.

Chaudhary, S., Zhang, S., Zhornitsky, S., Chen, Y., Chao, H.H., Li, C.R., 2023. Age-related
reduction in trait anxiety: Behavioral and neural evidence of automaticity in negative facial
emotion processing. Neuroimage 276, 120207.

Clauset, A., Newman, M.E., Moore, C., 2004. Finding community structure in very large
networks. Phys Rev E Stat Nonlin Soft Matter Phys 70(6 Pt 2), 066111.

Clifford, K.P., Miles, A.E., Prevot, T.D., Misquitta, K.A., Ellegood, J., Lerch, J.P., Sibille, E.,
Nikolova, Y.S., Banasr, M., 2023. Brain structure and working memory adaptations associated
with maturation and aging in mice. Front Aging Neurosci 15, 1195748.

Colizza, V., Flammini, A., Serrano, M.A., Vespignani, A., 2006. Detecting rich-club ordering in
complex networks. Nature Physics 2, 110-115.

Colon-Perez, L.M., Ibanez, K.R., Suarez, M., Torroella, K., Acuna, K., Ofori, E., Levites, Y.,

Vaillancourt, D.E., Golde, T.E., Chakrabarty, P., Febo, M., 2019. Neurite orientation dispersion

and density imaging reveals white matter and hippocampal microstructure changes produced by

Interleukin-6 in the TgCRND8 mouse model of amyloidosis. NeuroImage 202, 116138.

Cox, R.W., 1996. AFNI: software for analysis and visualization of functional magnetic resonance

neuroimages. Comput Biomed Res 29(3), 162-173.

Crawley, J.N., Chen, T., Puri, A., Washburn, R., Sullivan, T.L., Hill, J.M., Young, N.B., Nadler,

J.J., Moy, S.S., Young, L.J., Caldwell, H.K., Young, W.S., 2007. Social approach behaviors in

oxytocin knockout mice: comparison of two independent lines tested in different laboratory

environments. Neuropeptides 41(3), 145-163.

Curran, C.P., Marczinski, C.A., 2017. Taurine, caffeine, and energy drinks: Reviewing the risks

to the adolescent brain. Birth Defects Res 109(20), 1640-1648.

D'Aquila, P.S., Newton, J., Willner, P., 1997. Diurnal variation in the effect of chronic mild stress

on sucrose intake and preference. Physiol Behav 62(2), 421-426.

Elias, P.K., Elias, M.F., Eleftheriou, B.E., 1975. Emotionality, exploratory behavior, and

locomotion in aging inbred strains of mice. Gerontologia 21(1), 46-55.

Evans, A.K., Park, H.H., Saw, N.L., Singhal, K., Ogawa, G., Leib, R.D., Shamloo, M., 2021.

Age-related neuroinflammation and pathology in the locus coeruleus and hippocampus: beta-

adrenergic antagonists exacerbate impairment of learning and memory in aged mice. Neurobiol

Aging 106, 241-256.

Fahlstrom, A., Yu, Q., Ulfhake, B., 2011. Behavioral changes in aging female C57BL/6 mice.

Neurobiol Aging 32(10), 1868-1880.

Ferreira, A.C., Sousa, N., Sousa, J.C., Marques, F., 2023. Age-related changes in mice

behavior and the contribution of lipocalin-2. Front Aging Neurosci 15, 1179302.

Fischer, H., Sandblom, J., Gavazzeni, J., Fransson, P., Wright, C.I., Backman, L., 2005. Age-
differential patterns of brain activation during perception of angry faces. Neurosci Lett 386(2),
99-104.

Flurkey, K., Currer, J.M., Harrison, D.E., 2007. Mouse models in aging research, in: Fox J. G.,
Q.F.W., Newcomer C. E., Davisson M. T., Barthold S. W., Smith A. L. (Ed.) The mouse in
biomedical research: normative biology, husbandry, and models. Academic Press, pp. 637-672.

Frick, K.M., Burlingame, L.A., Arters, J.A., Berger-Sweeney, J., 2000. Reference memory,
anxiety and estrous cyclicity in C57BL/6NIA mice are affected by age and sex. Neuroscience
95(1), 293-307.

G., C., Nepusz, T., 2006. The igraph software package for complex network research. Complex
Systems, 1695.

Grandjean, J., Schroeter, A., Batata, I., Rudin, M., 2014a. Optimization of anesthesia protocol
for resting-state fMRI in mice based on differential effects of anesthetics on functional
connectivity patterns. Neuroimage 102 Pt 2, 838-847.

Grandjean, J., Schroeter, A., He, P., Tanadini, M., Keist, R., Krstic, D., Konietzko, U., Klohs, J.,
Nitsch, R.M., Rudin, M., 2014b. Early alterations in functional connectivity and white matter
structure in a transgenic mouse model of cerebral amyloidosis. J Neurosci 34(41), 13780-
13789.

Gross, J.J., Carstensen, L.L., Pasupathi, M., Tsai, J., Skorpen, C.G., Hsu, A.Y., 1997. Emotion
and aging: experience, expression, and control. Psychol Aging 12(4), 590-599.

Harb, M.R., Sousa, N., Zihl, J., Almeida, O.F., 2014. Reward components of feeding behavior
are preserved during mouse aging. Front Aging Neurosci 6, 242.

Humphries, M.D., Gurney, K., 2008. Network 'small-world-ness': a quantitative method for
determining canonical network equivalence. PLoS One 3(4), e0002051.

Humphries, M.D., Gurney, K., Prescott, T.J., 2006. The brainstem reticular formation is a small-
world, not scale-free, network. Proc Biol Sci 273(1585), 503-511.

Hyvarinen, A., 1999. Fast and robust fixed-point algorithms for independent component
analysis. IEEE Trans Neural Netw 10(3), 626-634.

Iidaka, T., Okada, T., Murata, T., Omori, M., Kosaka, H., Sadato, N., Yonekura, Y., 2002. Age-
related differences in the medial temporal lobe responses to emotional faces as revealed by
fMRI. Hippocampus 12(3), 352-362.

Jenkinson, M., Bannister, P., Brady, M., Smith, S., 2002. Improved optimization for the robust
and accurate linear registration and motion correction of brain images. Neuroimage 17(2), 825-
841.

Jonckers, E., Van Audekerke, J., De Visscher, G., Van der Linden, A., Verhoye, M., 2011.
Functional connectivity fMRI of the rodent brain: comparison of functional connectivity networks
in rat and mouse. PLoS One 6(4), e18876.

Jones, B., Balasubramaniam, M., Lebowitz, J.J., Taylor, A., Villalta, F., Khoshbouei, H., Grueter,
C., Grueter, B., Dash, C., Pandhare, J., 2021. Activation of proline biosynthesis is critical to
maintain glutamate homeostasis during acute methamphetamine exposure. Sci Rep 11(1),
1422.

Kelley, K.W., O'Connor, J.C., Lawson, M.A., Dantzer, R., Rodriguez-Zas, S.L., McCusker, R.H.,
2013. Aging leads to prolonged duration of inflammation-induced depression-like behavior
caused by Bacillus Calmette-Guerin. Brain Behav Immun 32, 63-69.

Kessler, R.C., Petukhova, M., Sampson, N.A., Zaslavsky, A.M., Wittchen, H.U., 2012. Twelve-
month and lifetime prevalence and lifetime morbid risk of anxiety and mood disorders in the
United States. Int J Methods Psychiatr Res 21(3), 169-184.

Kim, S., Matyas, F., Lee, S., Acsady, L., Shin, H.S., 2012. Lateralization of observational fear
learning at the cortical but not thalamic level in mice. Proc Natl Acad Sci U S A 109(38), 15497-
15501.

Kishimoto, Y., Suzuki, M., Kawahara, S., Kirino, Y., 2001. Age-dependent impairment of delay
and trace eyeblink conditioning in mice. Neuroreport 12(15), 3349-3352.

Klein, A., Andersson, J., Ardekani, B.A., Ashburner, J., Avants, B., Chiang, M.C., Christensen,
G.E., Collins, D.L., Gee, J., Hellier, P., Song, J.H., Jenkinson, M., Lepage, C., Rueckert, D.,
Thompson, P., Vercauteren, T., Woods, R.P., Mann, J.J., Parsey, R.V., 2009. Evaluation of 14
nonlinear deformation algorithms applied to human brain MRI registration. Neuroimage 46(3),
786-802.

Kotlarz, P., Nino, J.C., Febo, M., 2022. Connectomic analysis of Alzheimer's disease using
percolation theory. Network Neuroscience 6(1), 213–233.

Lê, S., Josse, J., Husson, F., 2008. FactoMineR: An R Package for Multivariate Analysis.
Journal of Statistical Software 25(1), 1 - 18.

Lein, E.S., Hawrylycz, M.J., Ao, N., Ayres, M., Bensinger, A., Bernard, A., Boe, A.F., Boguski,
M.S., Brockway, K.S., Byrnes, E.J., Chen, L., Chen, L., Chen, T.M., Chin, M.C., Chong, J.,
Crook, B.E., Czaplinska, A., Dang, C.N., Datta, S., Dee, N.R., Desaki, A.L., Desta, T., Diep, E.,
Dolbeare, T.A., Donelan, M.J., Dong, H.W., Dougherty, J.G., Duncan, B.J., Ebbert, A.J.,
Eichele, G., Estin, L.K., Faber, C., Facer, B.A., Fields, R., Fischer, S.R., Fliss, T.P., Frensley,
C., Gates, S.N., Glattfelder, K.J., Halverson, K.R., Hart, M.R., Hohmann, J.G., Howell, M.P.,
Jeung, D.P., Johnson, R.A., Karr, P.T., Kawal, R., Kidney, J.M., Knapik, R.H., Kuan, C.L., Lake,
J.H., Laramee, A.R., Larsen, K.D., Lau, C., Lemon, T.A., Liang, A.J., Liu, Y., Luong, L.T.,
Michaels, J., Morgan, J.J., Morgan, R.J., Mortrud, M.T., Mosqueda, N.F., Ng, L.L., Ng, R., Orta,
G.J., Overly, C.C., Pak, T.H., Parry, S.E., Pathak, S.D., Pearson, O.C., Puchalski, R.B., Riley,
Z.L., Rockett, H.R., Rowland, S.A., Royall, J.J., Ruiz, M.J., Sarno, N.R., Schaffnit, K.,
Shapovalova, N.V., Sivisay, T., Slaughterbeck, C.R., Smith, S.C., Smith, K.A., Smith, B.I., Sodt,
A.J., Stewart, N.N., Stumpf, K.R., Sunkin, S.M., Sutram, M., Tam, A., Teemer, C.D., Thaller, C.,
Thompson, C.L., Varnam, L.R., Visel, A., Whitlock, R.M., Wohnoutka, P.E., Wolkey, C.K.,
Wong, V.Y., Wood, M., Yaylaoglu, M.B., Young, R.C., Youngstrom, B.L., Yuan, X.F., Zhang, B.,
Zwingman, T.A., Jones, A.R., 2007. Genome-wide atlas of gene expression in the adult mouse
brain. Nature 445(7124), 168-176.

Lhotellier, L., Cohen-Salmon, C., 1989. Genetics and senescence. I. Age-related changes in

activity and exploration in three inbred strains of mice. Physiol Behav 45(3), 491-493.

Machado, L., Thompson, L.M., Brett, C.H.R., 2019. Visual analogue mood scale scores in

healthy young versus older adults. Int Psychogeriatr 31(3), 417-424.

Mahar, R., Chang, M.C., Merritt, M.E., 2021. Measuring NQO1 Bioactivation Using

[(2)H7]Glucose. Cancers (Basel) 13(16).

Malatynska, E., Steinbusch, H.W., Redkozubova, O., Bolkunov, A., Kubatiev, A., Yeritsyan,

N.B., Vignisse, J., Bachurin, S., Strekalova, T., 2012. Anhedonic-like traits and lack of affective

deficits in 18-month-old C57BL/6 mice: Implications for modeling elderly depression. Exp

Gerontol 47(8), 552-564.

Malezieux, M., Klein, A.S., Gogolla, N., 2023. Neural Circuits for Emotion. Annual review of

neuroscience 46, 211-231.

Mathis, A., Mamidanna, P., Cury, K.M., Abe, T., Murthy, V.N., Mathis, M.W., Bethge, M., 2018.

DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. Nat

Neurosci 21(9), 1281-1289.

McCall, N.M., Sprow, G.M., Delpire, E., Thiele, T.E., Kash, T.L., Pleil, K.E., 2013. Effects of sex

and deletion of neuropeptide Y2 receptors from GABAergic neurons on affective and alcohol

drinking behaviors in mice. Front Integr Neurosci 7, 100.

McLean, D.J., Volponi, M.A.S., 2018. trajr: An R package for characterisation of animal

trajectories. Ethology 124(6), 440-448.

Meyer, H.C., Odriozola, P., Cohodes, E.M., Mandell, J.D., Li, A., Yang, R., Hall, B.S.,

Haberman, J.T., Zacharek, S.J., Liston, C., Lee, F.S., Gee, D.G., 2019. Ventral hippocampus

interacts with prelimbic cortex during inhibition of threat response via learned safety in both mice

and humans. Proc Natl Acad Sci U S A 116(52), 26970-26979.

Moyer, J.R., Jr., Brown, T.H., 2006. Impaired trace and contextual fear conditioning in aged rats.
Behavioral neuroscience 120(3), 612-624.

Murdy, T.J., Dunn, A.R., Singh, S., Telpoukhovskaia, M.A., Zhang, S., White, J.K., Kahn, I.,
Febo, M., Kaczorowski, C.C., 2022. Leveraging genetic diversity in mice to inform individual
differences in brain microstructure and memory. Frontiers in behavioral neuroscience 16,
1033975.

Nath, T., Mathis, A., Chen, A.C., Patel, A., Bethge, M., Mathis, M.W., 2019. Using DeepLabCut
for 3D markerless pose estimation across species and behaviors. Nat Protoc 14(7), 2152-2176.

Neuner, S.M., Heuer, S.E., Huentelman, M.J., O'Connell, K.M.S., Kaczorowski, C.C., 2019.
Harnessing Genetic Complexity to Enhance Translatability of Alzheimer's Disease Mouse
Models: A Path toward Precision Medicine. Neuron 101(3), 399-411 e395.

Newman, M.E., 2002. Assortative mixing in networks. Physical review letters 89(20), 208701.

Newman, M.E., 2006. Finding community structure in networks using the eigenvectors of
matrices. Phys Rev E Stat Nonlin Soft Matter Phys 74(3 Pt 2), 036104.

Newman, M.E., Girvan, M., 2004. Finding and evaluating community structure in networks. Phys
Rev E 69(2), 026113.

Oh, H.J., Song, M., Kim, Y.K., Bae, J.R., Cha, S.Y., Bae, J.Y., Kim, Y., You, M., Lee, Y., Shim,
J., Maeng, S., 2018. Age-Related Decrease in Stress Responsiveness and Proactive Coping in
Male Mice. Front Aging Neurosci 10, 128.

Onnela, J.P., Saramaki, J., Kertesz, J., Kaski, K., 2005. Intensity and coherence of motifs in
weighted complex networks. Phys Rev E Stat Nonlin Soft Matter Phys 71(6 Pt 2), 065103.

Ortiz, S., Latsko, M.S., Fouty, J.L., Dutta, S., Adkins, J.M., Jasnow, A.M., 2019. Anterior
Cingulate Cortex and Ventral Hippocampal Inputs to the Basolateral Amygdala Selectively
Control Generalized Fear. J Neurosci 39(33), 6526-6539.

Pang, Z., Chong, J., Zhou, G., de Lima Morais, D.A., Chang, L., Barrette, M., Gauthier, C.,

Jacques, P.E., Li, S., Xia, J., 2021. MetaboAnalyst 5.0: narrowing the gap between raw spectra

and functional insights. Nucleic Acids Res 49(W1), W388-W396.

Paxinos, G., Franklin, K.B.J., 2001. The mouse brain in stereotaxic coordinates, 2nd ed.

Academic Press, San Diego.

Pi, T., Wei, S., Jiang, Y., Shi, J.S., 2021. High Methionine Diet-Induced Alzheimer's Disease like

Symptoms Are Accompanied by 5-Methylcytosine Elevated Levels in the Brain. Behav Neurol

2021, 6683318.

Pompilus, M., Colon-Perez, L.M., Grudny, M.M., Febo, M., 2020. Contextual experience

modifies functional connectome indices of topological strength and efficiency. Sci Rep 10(1),

19843.

Pons, P., Latapy, M., 2005. Computing Communities in Large Networks Using Random Walks.

Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 284-293.

Ravindranath, O., Ordaz, S.J., Padmanabhan, A., Foran, W., Jalbrzikowski, M., Calabro, F.J.,

Luna, B., 2020. Influences of affective context on amygdala functional connectivity during

cognitive control from adolescence through adulthood. Dev Cogn Neurosci 45, 100836.

Rubinov, M., Sporns, O., 2010. Complex network measures of brain connectivity: uses and

interpretations. Neuroimage 52(3), 1059-1069.

Sadaka, A.H., Canuel, J., Febo, M., Johnson, C.T., Bradshaw, H.B., Ortiz, R., Ciumo, F.,

Kulkarni, P., Gitcho, M.A., Ferris, C.F., 2023. Effects of inhaled cannabis high in Delta9-THC or

CBD on the aging brain: A translational MRI and behavioral study. Front Aging Neurosci 15,

1055433.

Sakthivel, R., Criado-Marrero, M., Barroso, D., Braga, I.M., Bolen, M., Rubinovich, U., Hery,

G.P., Grudny, M.M., Koren, J., 3rd, Prokop, S., Febo, M., Abisambra, J.F., 2023. Fixed Time-

Point Analysis Reveals Repetitive Mild Traumatic Brain Injury Effects on Resting State

Functional Magnetic Resonance Imaging Connectivity and Neuro-Spatial Protein Profiles. J
Neurotrauma.

SAMHSA, 2022. Key substance use and mental health indicators in the United States: Results
from the 2021 National Survey on Drug Use and Health. Center for Behavioral Health Statistics
and Quality, Substance Abuse and Mental Health Services Administration.

Sanders, M.J., 2011. Context processing in aging: older mice are impaired in renewal of
extinguished fear. Exp Aging Res 37(5), 572-594.

Semenza, E.R., Harraz, M.M., Abramson, E., Malla, A.P., Vasavda, C., Gadalla, M.M.,
Kornberg, M.D., Snyder, S.H., Roychaudhuri, R., 2021. D-cysteine is an endogenous regulator
of neural progenitor cell dynamics in the mammalian brain. Proc Natl Acad Sci U S A 118(39).

Serradj, N., Jamon, M., 2007. Age-related changes in the motricity of the inbred mice strains
129/sv and C57BL/6j. Behav Brain Res 177(1), 80-89.

Sforazzini, F., Schwarz, A.J., Galbusera, A., Bifone, A., Gozzi, A., 2014. Distributed BOLD and
CBV-weighted resting-state networks in the mouse brain. Neuroimage 87, 403-415.

Shoji, H., Miyakawa, T., 2019. Age-related behavioral changes from young to old age in male
mice of a C57BL/6J strain maintained under a genetic stability program. Neuropsychopharmacol
Rep 39(2), 100-118.

Sturman, O., von Ziegler, L., Schlappi, C., Akyol, F., Privitera, M., Slominski, D., Grimm, C.,
Thieren, L., Zerbi, V., Grewe, B., Bohacek, J., 2020. Deep learning-based behavioral analysis
reaches human accuracy and is capable of outperforming commercial solutions.
Neuropsychopharmacology 45(11), 1942-1952.

Tan, Y., Singhal, S.M., Harden, S.W., Cahill, K.M., Nguyen, D.M., Colon-Perez, L.M., Sahagian,
T.J., Thinschmidt, J.S., de Kloet, A.D., Febo, M., Frazier, C.J., Krause, E.G., 2019. Oxytocin
Receptors Are Expressed by Glutamatergic Prefrontal Cortical Neurons That Selectively
Modulate Social Recognition. J Neurosci 39(17), 3249-3263.

Telesford, Q.K., Joyce, K.E., Hayasaka, S., Burdette, J.H., Laurienti, P.J., 2011. The ubiquity of
small-world networks. Brain connectivity 1(5), 367-375.

Terlizzi, E.P., Villarroel, M.A., 2020. Symptoms of generalized anxiety disorder among adults:
United States, 2019, NCHS Data Brief. National Center for Health Statistics, Hyattsville, MD.

Tessitore, A., Hariri, A.R., Fera, F., Smith, W.G., Das, S., Weinberger, D.R., Mattay, V.S., 2005.
Functional changes in the activity of brain regions underlying emotion processing in the elderly.
Psychiatry Res 139(1), 9-18.

van den Heuvel, M.P., Sporns, O., 2011. Rich-club organization of the human connectome. J
Neurosci 31(44), 15775-15786.

von Bohlen und Halbach, O., Zacher, C., Gass, P., Unsicker, K., 2006. Age-related alterations in
hippocampal spines and deficiencies in spatial memory in mice. J Neurosci Res 83(4), 525-531.

Wang, Q., Qi, L., He, C., Feng, H., Xie, C., Depression Imaging, R.C., 2023. Age- and gender-
related dispersion of brain networks across the lifespan. Geroscience.

Wang, Y., Ghumare, E., Vandenberghe, R., Dupont, P., 2017. Comparison of Different
Generalizations of Clustering Coefficient and Local Efficiency for Weighted Undirected Graphs.
Neural computation 29(2), 313-331.

Wright, C.I., Wedig, M.M., Williams, D., Rauch, S.L., Albert, M.S., 2006. Novel fearful faces
activate the amygdala in healthy young and elderly adults. Neurobiol Aging 27(2), 361-374.

Xia, M., Wang, J., He, Y., 2013. BrainNet Viewer: a network visualization tool for human brain
connectomics. PLoS One 8(7), e68910.

Yushkevich, P.A., Piven, J., Hazlett, H.C., Smith, R.G., Ho, S., Gee, J.C., Gerig, G., 2006. User-
guided 3D active contour segmentation of anatomical structures: significantly improved
efficiency and reliability. NeuroImage 31(3), 1116-1128.

1374

Table 1. Age differences in modularity persist across different cluster detection methods. All network measures calculated on mouse functional connectivity graphs with an edge density threshold preserving the top 16% Pearson correlations. Randomization of original networks preserved weights, node degree and strength sequences. Synthetic lattice structure used in small world index calculation was generated with the same number of nodes and approximately similar edge density.

Source	Network routine	Original			Random			Lattice
		2.5MO	7.5MO	15MO	2.5MO	7.5MO	15MO	
igraph	Assortativity	0.11±0.07	0.06±0.03	0.07±0.06	0.00±0.08	-0.02±0.05	-0.04±0.08	0.43
	Mean Distance	1.03±0.18	1.07±0.20	0.99±0.13	2.03±0.02	2.03±0.01	2.02±0.01	7.11
	Transitivity[1]	0.38±0.04	0.38±0.06	0.32±0.03*, ***	0.16±0.02	0.16±0.01	0.16±0.01	0.61
	Lambda	0.51±0.09	0.52±0.10	0.49±0.06	-	-	-	2.83
	Gamma[2]	1.35±0.12	1.39±0.12	1.54±0.20*	-	-	-	4.88
	Small world coefficient	4.68±0.76	4.64±0.51	4.09±0.55	-	-	-	1.73
	Small world index	1.39±0.40	1.33±0.45	1.54±0.20	-	-	-	0.15
	Modularity algorithm (cluster detection)							
	Fast greedy	4.50±0.71	5.70±1.70	5.00±0.67	5.40±1.07	4.70±0.68	4.90±0.74	3
	Edge betweenness[3]	15.8±6.44 *, **	27.8±5.1	23.8±8.32	26.4±6.60	26.2±4.87	24.3±6.60	4
	Lead eigenvector	4.20±1.32	5.50±1.84	4.50±1.18	4.50±1.18	4.70±1.49	4.70±1.34	6
	Walk trap[4]	6.10±2.64 **	9.90±4.36	7.90±1.29	5.60±1.65	5.90±1.60	6.40±1.96	6
	Louvain's algorithm	4.80±0.92	6.00±1.56	5.50±0.85	5.40±0.52	5.70±0.68	5.90±0.32	5
	Modularity Q value							
	Fast greedy[5]	0.34±0.04 **	0.30±0.02	0.31±0.03	0.24±0.04	0.23±0.01	0.23±0.01	0.60
	Edge betweenness[6]	0.25±0.07 *, **	0.12±0.04	0.16±0.08	0.13±0.04	0.13±0.02	0.15±0.03	0.65
	Lead eigenvector[7]	0.32±0.04 *, **	0.27±0.02	0.28±0.03	0.22±0.02	0.21±0.02	0.22±0.01	0.65
	Walk trap[8]	0.32±0.04 **	0.26±0.04	0.28±0.04	0.22±0.01	0.21±0.02	0.21±0.01	0.65
	Louvain's[9]	0.35±0.03 *, **	0.31±0.03	0.31±0.02	0.25±0.01	0.24±0.01	0.24±0.01	0.66

Brain connectivity toolbox							
Assortativity[10]	0.13±0.06 **	0.06±0.04	0.08±0.06	0.02±0.10	-0.05±0.11	-0.08±0.10	0.11
Characteristic path length	4.28±0.54	4.16±0.45	4.17±0.48	3.94±0.49	3.85±0.56	3.90±0.46	8.52
Transitivity	0.20±0.05	0.21±0.06	0.17±0.03	0.06±0.03	0.07±0.03	0.06±0.02	0.59
Gamma	1.77±0.35	1.57±0.28	1.49±0.29	-	-	-	9.83
Lambda	1.08±0.03	1.09±0.07	1.07±0.02	-	-	-	3.09
Small world coefficient	1.63±0.31	1.46±0.31	1.39±0.27	-	-	-	3.18
Small world index	0.58±0.08	0.57±0.15	0.65±0.05	-	-	-	0.22
Louvain's groups[11]	4.30±0.68 **	5.80±1.62	5.10±0.74	4.30±0.68	5.80±1.62	5.10±0.74	6
Louvain's Q[12]	0.35±0.04 *, **	0.30±0.03	0.31±0.03	0.24±0.01	0.22±0.02	0.24±0.02	0.70

Multivariate analysis of variance (MANOVA) was used to determine differences between groups (Age x Sex main effects and interactions, $\alpha<0.05$). Superscript numbers next to network routines indicate the following main effects of Age: (1) $F_{2, 24} = 4.3$, $p = 0.03$; (2) $F_{2, 24} = 3.8$, $p = 0.04$; (3) $F_{2, 24} = 7.6$, $p = 0.003$; (4) $F_{2, 24} = 3.6$, $p = 0.04$; (5) $F_{2, 24} = 3.9$, $p = 0.03$; (6) $F_{2, 24} = 8.6$, $p = 0.001$; (7) $F_{2, 24} = 5.7$, $p = 0.009$; (8) $F_{2, 24} = 4.5$, $p = 0.02$; (9) $F_{2, 24} = 6.1$, $p = 0.007$; (10) $F_{2, 24}= 3.5$, $p = 0.04$; (11) $F_{2, 24} = 4.7$, $p = 0.02$; (12) $F_{2, 24} = 5.1$, $p = 0.01$. Significant differences for post hoc multiple comparison Tukey's honest significant difference test indicated by *, **, *** for the following comparisons: 2.5MO vs 15MO, 2.5MO vs 7.5MO, and 2.5MO vs 15MO, respectively. Network measure values are presented as mean ± standard deviation. Values for randomized versions of the original functional MRI networks are shown as a comparison. Calculations were made on a non-toroidal lattice structure with approximately 300 weighted edges per 60 nodes (no on-diagonal connections; igraph version was undirected and BCT version directed).