# *In-vivo* multicellular control of gene expression in microbial consortia

Davide Salzano[a,b], Barbara Shannon[b], Claire Grierson[c,d], Lucia Marucci[d,e,f*], Nigel J Savery[b,d*],

Mario di Bernardo [a,g,*]

[a]*Scuola Superiore Meridionale, Naples, Italy*

[b] *School of Biochemistry, University of Bristol, United Kingdom*

[c] *School of Biological Sciences, University of Bristol, United Kingdom*

[d] *Bristol BioDesign Institute, University of Bristol, United Kingdom*

[e] *School of Engineering Mathematics and Technology, University of Bristol, United Kingdom*

[f] *School of Cellular and Molecular Medicine, University of Bristol, United Kingdom*

[g] *Department of Electrical Engineering and Information Technology, University of Naples Federico II,*
*Italy*

**\*** *These authors contributed equally*

[#] Correspondence should be addressed to: mario.di.bernardo@unina.it

**Keywords**: Cybergenetics, synthetic biology, control engineering, multicellular control, microbial consortia, gene regulation, Escherichia coli, modularity, biomedicine, industrial bioproduction.

**Abstract**

Cybergenetics, an emerging discipline straddling synthetic biology and control engineering,

seeks to harness feedback control mechanisms for engineering reliable synthetic biological

circuits that dynamically regulate target gene expression within host organisms. Current

approaches, including embedded control strategies in both Escherichia coli and mammalian

cells, suffer from limited modularity, requiring extensive re-engineering for controller design

modifications. Furthermore, the challenge of engineering dependable microbial consortia

remains a significant hurdle in synthetic biology and biotechnology. To address these

limitations, we propose the concept of multicellular control, where distinct populations within

a consortium perform specific tasks. In this paradigm, a controller population senses and

computes real-time control actions to regulate processes within a second population, denoted

as targets. Despite being theoretically proposed, in-vivo feasibility of multicellular control had

remained unverified until now.

In this study, we propose for the first time a biological implementation and the experimental

validation of multicellular control in a consortium of two Escherichia coli populations.

Through comprehensive characterization, we demonstrate effective inter-population

communication and behavioural influence, allowing the targets to achieve predetermined

outcomes set by the controllers. Importantly, our approach enables tunable regulation of

desired genes within the targets, showcasing robustness against perturbations, notably

imbalances in population sizes.

Our innovative architecture not only establishes efficient cross-population communication

but also offers adaptable gene regulation, even in the face of population imbalances. This

advancement opens avenues for robust module development, with potential applications in

biomedicine and industrial bioproduction, marking a significant stride toward overcoming

the challenges faced in the realm of synthetic biology and biotechnology.

## Introduction

Synthetic biology aims to engineer biomolecular systems to achieve new functions [1]; potential applications range from designing bacteria that can produce biofuels or sense and degrade pollutants in the environment (such as hydrocarbons and plastics), to immune cells that can track and kill cancer cells, or that can release drugs at specific points in desired conditions to avoid side effects (see [1] for an extensive review). Biomolecular feedback controllers have been highlighted as a crucial ingredient for reliable and robust synthetic biology devices [2]. Specifically, by applying concepts from control theory to guarantee desired static and dynamic performance of such devices, it is possible to drastically decrease the development time of new synthetic systems and to increase their flexibility and modularity [3, 4, 5].

Different designs have been proposed to implement biomolecular feedback control strategies both *in-silico* and *in-vivo*, ranging from simple feedback and feedforward loops to PID (Proportional Integral Derivative) controllers (see [3] and references therein). In particular, the introduction of an integral control action via an antithetic feedback motif has been highlighted as a fundamental ingredient for a synthetic circuit able to achieve and maintain a desired output even in the presence of disturbances [6], a property that has been termed *robust perfect adaptation* [5]. Integral biomolecular controllers have been thoroughly studied both theoretically [7] and experimentally, and different implementations have been proposed in bacteria [5] and mammalian cells [8]. For a review of the recent advancements in the implementations of biomolecular controllers guaranteeing robust perfect adaptation see [2].

However, existing robust controllers predominantly function at the single-cell level, posing inherent limitations on circuit complexity due to metabolic burdens and hindering their application as flexible plug-and-play modules across diverse synthetic and endogenous systems [6]. To overcome these constraints, a multicellular control strategy was conceived and computationally investigated [9]. In this approach, functionalities are distributed across distinct cell populations within a microbial consortium. Each cell strain encapsulates specific engineered gene networks, forming the foundation for a *distributed* feedback control loop [10] involving two populations: a controller population and a target cell population. The former

senses the targets' output, compares it with a reference signal encoding the desired expression

level, and provides the latter with stimuli based on antithetic control logic [11, 12, 13, 14].

Following this work, in [15, 16] a biological implementation of a comparator, the logic core of

the controller population, was designed and characterized both statically and dynamically but

never tested *in vivo* with a target population. Further multicellular controllers were proposed

(see [17] [18] and references within) but, to the best of our knowledge, the in-vivo

implementation and validation of a distributed multicellular feedback control loop in a

microbial consortium remains unexplored.

In this paper, we address this fundamental gap in the literature by developing and validating

*in-vivo* a feedback control loop distributed across a bacterial consortium. We demonstrate its

effectiveness in regulating a phenotype of interest in the target population even in the

presence of variations in the consortium composition. This is achieved by designing two

*Escherichia coli* strains embedding all the modules required to implement the three

fundamental functions needed for control, namely computation, sensing and actuation [19].

We show that the designed cellular populations respond to their respective inputs and that

the two populations can exchange information through two rationally designed

communication channels, implemented using orthogonal quorum sensing signalling

molecules. To demonstrate the validity of the proposed strategy, we carry out *in-vivo*

experiments comparing the closed loop configuration, where the controller cells can sense the

signalling molecules from the targets, with an open-loop implementation, where they cannot.

Our results show that the proposed multicellular control approach can successfully be used

to achieve reliable and robust regulation of the expression level of a gene of interest in the

targets while guaranteeing robustness to composition imbalances in the consortium, a

particularly important property for practical implementations.

**Results**

*A multicellular distributed control architecture*

The implementation of a multicellular control architecture requires the rational design of two

populations, namely the *controllers* and the *targets* (see Figure 1a). The controllers receive

information about the targets' output ($y$) which is the expression level of some gene of interest

therein. This value is compared to that of some reference signal (*ref*) encoding the desired

expression level and a control input ($u$) is then generated via the designed control logic. This

signal is transmitted to the targets in order to regulate the level of their output to that encoded

by the reference. The communication between controllers and targets is implemented via

orthogonal Quorum Sensing molecules, where each cell population produces signalling

molecules that can diffuse through the growth environment and into the other population.

The control logic, embedded in the controllers, is an antithetic feedback control strategy [5],

based on the *in-silico* analysis presented in [9] and derived from the error computation module

(comparator) implemented *in-vivo* using molecular titration in [15] and characterized

dynamically in [16] . The controller network is a multi-input, single-output device that uses

two independent signals to regulate the expression level of its output protein (see Figure 1b).

It is constructed around a $\sigma$ factor [20], which is a protein that recruits RNA polymerase

(RNAP) to target promotors by binding to DNA sequences in the -10 and -35 regions of those

promoters , and their cognate anti-$\sigma$, which blocks the interaction between RNAP and the $\sigma$

factor by sequestering the $\sigma$ factor. The production of $\sigma$ is controlled by IPTG (the reference

signal), which induces expression from the promoter *plac-UV5*. The production of anti-$\sigma$ is

regulated by the *plux* promoter, which is activated by the complex formed by LuxR, produced

constitutively, and 3-O-C6-HSL, produced by the targets. The output signal of the controllers

(3-O-C12-HSL) is produced by the product of the *lasI* gene,  which is regulated by the *p20 992*

promoter. This promoter, placed upstream of *lasI*, is induced by the presence of the orthogonal

$\sigma$ factor. The $\sigma$ and anti-$\sigma$ factors interact by binding and forming a complex that is unable to

recruit RNAP to *p20_992*. All proteins are fused to a degradation tag (ssrA tag) [21].

The targets can express a green fluorescent protein (GFP), whose expression level we aim to

regulate. The *gfp* gene is transcribed together with *luxI*, whose product is responsible for the

production of the molecule 3-O-C-6-HSL. This small molecule operates as a proxy of the *gfp*

expression levels by diffusing in the growth environment and activating the production of

anti-$\sigma$ in the controllers. The production of LuxI and GFP is regulated by *plas*. This promoter

is activated by the complex formed by LasR, which is constitutively expressed, and 3-O-C12-

HSL, which is produced by the controller population. In our implementation, LuxI was fused

to a degradation tag (ssrA tag) to ensure its fast degradation. All the details of the plasmids

and strains used in this work can be found in the *Strains and constructs* section of the Methods.

The goal of the design is to make the two engineered populations cooperate to steer and

maintain *gfp* expression in the targets at a desired setpoint level, which can be changed by

modifying the concentration of IPTG. Specifically, when *gfp* is overexpressed, the excessive

production of 3-O-C6-HSL upregulates the production of anti-$\sigma$. In turn, a reduction of the

free available $\sigma$, downregulates the production of LasI and, in turn, of the signalling molecule

3-O-C12-HSL. As a consequence, the activity of *pLas* in the targets should be reduced,

decreasing GFP expression levels. Similar mechanisms lead to an increase in GFP when GFP

falls below the desired level.

*Gene expression can be regulated via a multicellular architecture*

Prior to mixing the two populations in the consortium, we tested the functional modules

embedded in each population using flow cytometry [22]. Details on the machines and the

settings used can be found in section *Samples preparation and analysis using flow cytometry* of the

Methods.

We started by characterizing the response of the controller population to IPTG and 3-O-C6-

HSL, which are used in the architecture as a reference signal and a proxy for the targets state,

respectively. Note that, to measure the output level of the controllers, we substituted *lasI* with

a *gfp* fused with an ssrA tag (see Figure S1). We grew the controllers at 37 °C for 3h adding

different concentrations of IPTG and 3-O-C6-HSL to the growth media to test their response

to their inputs. The fluorescence levels of the population was then analysed by Flow

Cytometry (for more details see section *Induction protocol* of the Methods). The addition of

IPTG promoted *plac-UV5* activity by inhibiting the repression exerted by LacI, thus inducing

up to 300 fold increase in the steady state fluorescence when 100 $\mu M$ IPTG was added to the

media (Figure 2a). Conversely, the addition of 3-O-C6-HSL increased the anti-$\sigma$ level of

transcription by activating *plux;* as a consequence, the fluorescence levels consistently

decreased, resulting in the inhibition of *gfp* expression levels up to 100 folds when 100 nM 3-

O-C6-HSL was added to the growth media (Figure 2a). We used the same procedure to assess

the response of the targets to 3-O-C12-HSL; as expected, addition of 10 $\mu M$ 3-O-C12 in the

growth media promoted *plas* activity and a 14 fold increase was detected in average GFP

fluorescence (Figure 2b). These assays confirmed that the controllers were correctly

responding to changes either in the reference signal or in the output of the targets, and,

confirmed that the targets were able to adjust GFP production according to the concentration

of 3-O-C12-HSL. For further details on the standard deviation of the response of targets and

controllers see Figure S2.

We then assayed the ability of the two populations to communicate with each other. To test

the ability of the controllers to stimulate the targets in the absence of any feedback from the

latter, we implemented an open loop configuration of the consortium where the output

sensing module was removed from the controller cells (see Figure 2c, right panel). This made

them unable to receive feedback sensing molecules from the targets. We will refer to this

population as *open loop controllers*.

We mixed the open loop controllers with the targets every hour, grew the consortium at 37 ᵒC

for 6h and analysed the targets fluorescence via flow cytometry (for further details on the

experimental protocol see section *Time-course assays* of the methods). This time course,

reported in Figure 2d (light blue trajectory), shows that the open loop controllers are able to

communicate with the targets, i.e. that the 3-O-C12-HSL is successfully produced by the open

loop controllers and sensed by the targets. Indeed, the controllers were able to induce a 15

fold increase in the average fluorescence of the targets after 6h. However, over this time period

the open loop controllers were not able to stabilize the *gfp* expression levels in the targets

confirming that, as expected, they were unable to receive any feedback signal sent from them.

We performed a similar experiments mixing the controllers with the targets. In closed loop,

where  controllers possess the module to sense the expression level of the controlled gene in

the targets, stabilization of the targets fluorescence was observed after 3h (Figure 2d, yellow

trajectory). This confirms that the targets were able to successfully produce 3-O-C6-HSL

molecules which were correctly received by the controller population.

We then tested the ability of the controllers in the closed loop configuration to regulate *gfp*

expression levels in the targets to different values by modulating the concentration of IPTG in

the culture medium. Specifically, we grew controllers and targets in coculture at 37 ᵒC for 6h,

both with and without IPTG in the growth medium and then analysed the fluorescence level

of the targets using flow cytometry. When the consortium was induced with 50 $\mu$M IPTG, the

relative increase in fluorescence was approximately 50% (Figure 2e). This increase was
surprisingly small compared to the expected increase in the output observed during the
preliminary characterisation of the populations. Comparing the 6h timepoint of the closed
loop time course without IPTG shown in Figure 2d with the uninduced targets response
shown in Figure 2b, we observed that the targets were expressing high levels of fluorescence
when mixed with the controllers, even when no IPTG was present in the culture media (Figure
S3 shows a comparison of the distribution of fluorescence of the targets in the two scenarios).
This suggests that there was an unexpected high basal production of 3-O-C12-HSL, which was
not revealed during the initial characterisation of the populations shown in Figure 2a. The
inability to reveal the base expression level in the initial characterisation highlights the
difference between the expression level needed to detect any GFP signal and the levels of
enzyme needed to produce enough signalling molecules to induce the targets.

We addressed this problem by reducing the synthesis rate of $\sigma$ in the absence of IPTG, aiming
to increase the range of GFP fluorescence levels we could regulate the targets to. Specifically,
we increased the repression efficiency of *plac-UV5* by LacI by adding an extra lac operator
upstream of the promoter. This, as shown in [23], reduces the basal expression of the gene
controlled by LacI enabling the LacI tetramer to bind two operators simultaneously, thus
increasing the apparent affinity of the interaction (see *strains and constructs* section of the
methods for more details on the construction of the plasmid). We will henceforth denote the
modified controllers/open loop controllers "controllers/open loop controllers 2.0". After
adding the extra operator, the steady state fluorescence of the targets in closed loop and open
loop was reduced to levels that were lower than the ones reached by uninduced target
monocultures after 6h of growth (see Figure 3a,b). This indicated that the addition of the extra
operator reduced dramatically the leakiness of the *plac-UV5* promoter. We then repeated the
dynamic characterisation of the open and closed loop configurations of the consortium using
the new version of both controller populations. In open loop, the open loop controllers 2.0
were able to stabilize *gfp* expression levels in the targets after 6h. In addition, by increasing
the IPTG concentration in the culture media, we could achieve up to a 10-fold increase in GFP
expression levels. However, the regulation was highly variable across the different biological
replicates, especially at $0\ \mu M$ and $3\ \mu M$ IPTG (see Figure 3a). By closing the feedback loop with
the addition of the output sensing module in the controllers, we were able to increase the

reliability of the regulation, reducing the standard deviation at the 6h timepoint 5-fold and
29-fold, when we used 0 $\mu$M and 3 $\mu$M IPTG, respectively. In addition, the settling time was
comparable to the open loop configuration (approximately 3h), and we were able to induce
up to a 5-fold increase when 7 $\mu$M IPTG was added to the culture medium (see Figure 3b).

We further investigated the tunability of the regulation in both configurations by analysing
the steady state response with increasing levels of IPTG (details of the protocol used can be
found in the *Titration assays* section of the Methods). Specifically, we mixed the
controllers/open loop controllers 2.0 with the targets and grew the consortium for 6h at 37 ℃.
Figure 3c shows that both consortia were able to regulate the *gfp* gene to different expression
levels. Both architectures showed saturation when IPTG $\in [0\ \mu M; 2\ \mu M]$. Above that
concentration, increases in IPTG concentrations drove an increase in the targets' fluorescence
levels. In addition, both consortia underwent an abrupt increase in the targets' average
fluorescence and in its variance when we used IPTG at concentrations above 8$\mu M$. Using the
remaining data (IPTG $\in [2\ \mu M,\ 8\ \mu M]$), where the response of both systems was mostly linear,
we observed some differences in  the regulation patterns between the two configurations. The
closed loop system showed an almost linear increase in GFP fluorescence, while in open loop
we noticed a sharp increase in fluorescence levels between 2$\mu$M and 3$\mu$M IPTG, followed by
much flatter increase in fluorescence when IPTG was present at concentrations above 5 $\mu M$.
We quantified the linearity of the input-output response of both configurations to IPTG by
fitting the data with linear least square estimators and comparing the $R^2$ values. Using the
data where IPTG $\in [2\ \mu M, 8\ \mu M]$, the estimator fitting the closed loop data (yellow dashed line
in Figure 3c) predicts almost perfectly the variance coming from the data ($R^2$ = 0.927). In
contrast, the estimator that fits the data obtained using the open loop configuration of the
consortium explained just above half of such variance ($R^2$ = 0.624).

In our experiments, we established the effectiveness of a multicellular control architecture in
regulating the expression of a desired gene within the target population. Our findings
underscore the significant impact of feedback incorporation, leading to a substantial reduction
in response variability and enhanced linearity in the input-output relationship of the
architecture.

*Closed loop control enhances robustness to changes in the consortium composition*

A crucial problem when multicellular strategies are deployed in a consortium across multiple

cellular populations [24] is that different populations may grow at different rates because of

differences in their metabolic loads. This may hinder the ability of the controllers to steer the

dynamics of the targets as it might cause fluctuations in the production rate of the sensing (3-

O-C6-HSL) and actuation (3-O-C12-HSL) molecules. We therefore analysed the robustness of

both the closed loop and the open loop configurations to changes in the consortium

composition (see Figure 4).

We produced consortia of different compositions by inoculating targets and controllers/open

loop controllers 2.0. in different ratios, then measuring the average fluorescence of the targets

after 6h (for more detail see section *Consortium composition assay* of the Methods). Figure 4 a-d

shows that varying the proportion of controllers2.0 and targets had no detectable effect on the

reliability of the regulation of the closed loop configuration where feedback between the two

populations is present. Conversely, in the absence of feedback (in the open-loop

configuration), we noted a declining trend in the fluorescence levels of the targets as the

percentage of targets increased, especially when 0 $\mu$M and 3 $\mu$M IPTG concentrations were

employed. This trend was not seen when 5 $\mu M$ and 7 $\mu$M IPTG were present in the culture

medium. We quantified the difference between the fluorescence of the targets depending on

their percentage in the consortium by fitting linear least square estimators to the data (dashed

lines in Figure 4 a-d). We then compared the slopes of the lines fitting the open and closed

loop datapoints at different IPTG concentrations (see Figure 4 e). Using analysis of covariance

and a multiple comparison test, we found that when 0$\mu$M and 3$\mu$M IPTG were used, the

slopes of the linear models fitting the closed and open loop datapoints were statistically

different with *p*-values equal to 0.01527 and 1.13 $10^{-6}$, respectively. In contrast, when 5$\mu$M

and 7$\mu$M IPTG were present in the culture media, the changes were not significant. This could

be attributed to the saturation of the *plas* promoter activity at higher IPTG concentrations,

which reduced the sensitivity of the targets to variations in the 3-O-C12-HSL concentration

(see Supplementary Figure S4 for more details and section S1 of the Supplementary

Information for the details on the protocol used).

The analysis was finalized by conducting a statistical comparison of the slopes of the curves

fitted to the closed-loop data points across all IPTG concentrations. Employing the same

methodology, we established that there were no significant differences in slopes within the
closed-loop system (detailed p-values provided in Table S3). This confirmed the consistent
robustness of the multicellular feedback control strategy in maintaining stability, even in the
presence of imbalances between the controllers' and targets' populations.

## Discussion

Our study presents for the first time the *in-vivo* biological implementation of synthetic
feedback controllers within cellular populations, using a multicellular control architecture
where, withing a consortium, a specific population of controller cells possesses the ability to
sense and regulate targeted phenotypes within another population of cells. This strategic
distribution of control functionalities across members of a microbial consortium alleviates the
metabolic burden that would otherwise be imposed if all functions were concentrated within
a single cell, as noted in prior research [5]. Moreover, this approach enhances the consortium's
robustness, enabling it to adapt effectively to changes in its composition.

Following the validation of individual components within our microbial community, we
meticulously assessed the architecture's performance and robustness using flow cytometry.
Our results unequivocally demonstrate the distributed feedback controller's capability to
effectively regulate gene expression within target cells, maintaining the desired levels even in
the face of alterations in the consortium's composition. To substantiate the effectiveness of our
control architecture, we conducted a comparative analysis between open and closed loop
implementations. This examination clearly underscores the pivotal role of feedback
mechanisms, revealing that their incorporation significantly enhances both the reliability and
robustness of gene regulation within the system.

The development of distributed feedback controllers across microbial communities paves the
way for the construction of complex interacting communities, where each population carries
out a specific function [25]. Implementation of our distributed consortia within
microfluidics/microscopy platforms would allow the realisation of more complex feedback
control tasks in space and time [26] [27] [16].

Such microbes could be used in applications ranging from the integration of bacteria in the

human gut microbiome to treat a class of diseases [28] to engineering the soil microbiome to

increase plant growth and health [29], and bacteria-based production of fuels, which presents

a promising source for sustainable energy [30].

Current theoretical work is aimed at improving the performance and robustness of the

proposed design by expanding the consortium to include additional controller populations,

adding, for example, populations able to implement proportional or derivative control actions

in order to achieve a multicellular implementation of biomolecular PID controllers recently

proposed in [31], [32].

**Methods**

*Strains and constructs*

*Escherichia coli* strain TOP10 was used for all the cloning manipulations of this work.

*Escherichia coli* strain MG1655 $\lambda$-, rph-1 [33] was used for all the assays in this work and

transformed with the plasmids constituting either the controllers or the targets. For all the

sequences of the primers and the synthesised genes used in this study see Table S1, Table S2.

In addition, all the maps of the plasmids are reported In Figure S5.

The controllers contain three plasmids: pVRa20_992_BS2 (medium copy number),

pVRc20_992_BS1 (medium copy number) and pVRb_LasI (low copy number). The genes

constituting the essential core of the computation module ($\sigma$, anti-$\sigma$ and lasI) are ssrA tagged

(AANDENYALAA) to ensure fast dynamics of expression and degradation [21].

pVRb_lasI encodes the *lasI* gene, under the control of the *p_20992* promoter, and was built in

[15], together with pVRb_ssrA, where the lasI gene was swapped with a superfolder *gfp*.

pVRb_lasI was present in the controllers in each of the assays where they were mixed with

the targets, whereas pVRb_ssra was present in the controllers for their input output

characterisation.

Vector backbones for pVRa20_992_BS2 and pVRc20_992_BS1 were derived from plasmids

pVRa20_992 and pVRc20_992 [20]. pVRa20_992_BS2 produces a $\sigma$ factor under the control of

a promoter that responds to IPTG.. To construct this plasmid, first the pVRa20_992_BS1

plasmid was obtained by standard restriction digestion and ligation procedure, by amplifying

the *σ* gene from the pLusB plasmid [15] with primers Sigma_plus_tag_F and Sigma_plus_tag_R and cloning the PCR product in the pVRa_20_992 plasmid after NcoI/BamHI digestion. Then, the *plac* promoter was substituted with *plac-UV5* to obtain the pVRa20_992_BS2 plasmid. The construct was built by standard restriction digestion and ligation procedure, by amplifying *plac-UV5* from the pVRa_placASb_Flag plasmid [15] with primers PlacUV5_F and PlacUV5_R and cloning the PCR product in pVRa_20_992_BS1 plasmid after XbaI digestion. Finally, we modified pVRa_20_992_BS2 by introducing a second lac operator upstream of *plac_UV5*. This construct, denoted as PVRa_20_992_DS, was constructed by Gibson assembly after PCR amplification of pVRa20_992_BS2 using primers Back_extralac_for and Back_extralac_rev, and of the pVRbLacO1O1Del from [34] using the Ins_extralac_for and Ins_extralac_rev primers. pVRc20_992_BS1 produces the anti-*σ* factor under the induction of 3-O-C6-HSL. The plasmid was constructed by standard restriction digestion and ligation procedure, by amplifying anti-*σ* from pVRa_placASb_Flag [15] with primers Anti_Sigma_plus_tag_F and Anti_Sigma_plus_tag_R and cloning the PCR product in pVRc_20_992 plasmid after BamHI/PstI digestion. Finally, the controllers expressing a Red Fluorescent Protein were implemented by substituting the pVRb_lasI plasmid with the pVRb_lasI_RFP plasmid. pVRb_lasI_RFP was assembled using standard restriction digestion and ligation procedure, by amplifying a synthetically generated RFP with primers RFP_Afe_for and RFP_Nde_rev and cloning the PCR product in the pVRb_lasI plasmid after AfeI/NdeI digestion.

The targets hosted p_Las_Lux_GFP_3.0, a plasmid that embeds the 3-O-C12-HSL inducible promoter *plas*, driving the expression of a mut3 green fluorescent protein and of the luxI gene (fused with and ssrA tag). This plasmid was assembled by Gibson assembly after PCR amplification of the las_composite_device plasmid from [35] with p_Las_Lux_GFP_3_0_fwd_20 and p_Las_Lux_GFP_3_0_rev_20 primers, and of the luxI from pTeLu_GFP constructed in [15] with the luxI_fwd and luxI_rev primers.

*Chemicals*

For all experiments, kanamycin (50 $\mu$g/mL), ampicillin (100 $\mu$g/mL) and chloramphenicol (25 $\mu$g/mL) were added to the growth media, depending on the bacteria selective resistance. These antibiotics were supplied by Sigma-Aldrich. pVRa_20992_BS2 and pVRa_20_992_DS have an

Ampicillin resistant gene, pVRb_lasI, pVRB_lasI_RFP and pVRb_ssrA have a gene conferring

Kanamycin resistantce, and pVRc_20_992_BS1 and p_Las_Lux_GFP_3.0 have a gene

guaranteeing Chroranmphenicol resistance. When multiple populations were mixed, the

antibiotics to which both populations were resistant were used. If no resistance was in

common, no antibiotic was used.

3-O-C6-HSL (N-(B-Ketocaproyl)-L-Homoserine Lactone from Sigma-Aldrich, cat # K3007)

and 3-O-C12-HSL (N-(3-Oxododecanoyl)-L-homoserine lactone from Sigma-Aldrich, cat #

O9139) were dissolved in DMSO, filter-sterilized and added to the LB growth medium at the

indicated concentrations. IPTG (Isopropyl $\beta$-D-1-thiogalactopyranoside, supplied by Sigma-

Aldrich, cat # I5502) was dissolved in water, filter-sterilized and added to LB medium at the

indicated concentrations.

*Samples preparation and analysis using flow cytometry*

Each sample (0.5 mL volume) was resuspended in 1xPBS and diluted to achieve a final $OD_{600}$

for the culture of 0.05. Data were acquired using either BD LSRFortessa X-20 or Acea Novocyte

flow cytometer. The results analysed using FlowJo.

The events recorded by a flow cytometer were first gated in the forward scatter (FSC-H, proxy

of cellular size), side scatter (SSC-H, proxy of cellular complexity) plane. Here, the events were

divided in three subpopulations: healthy cells where size and complexity were in

physiological ranges [36], stressed cells where the bacteria had increased size and complexity

due to filamentation, and cellular debris corresponding to events with high size with respect

to their complexity (Figure S6 a). Subsequently, healthy cells were gated in the SSC-H, SSC-A

plane to select for single cells. Cells with comparable signals in the two channels were selected

as single cells, whereas events with much larger SSC-A were classified as cellular

conglomerates and excluded from the analysis (Figure S6b). Finally, single cells were gated

using the FITC-H (Novocyte machine)/FITC-A (Fortessa machine) (490 nm excitation, 530/30

nm emission) channel to discriminate targets and controllers based on their fluorescence

levels. Specifically, cells with low FITC-H signal were classified as controllers and cells having

high FITC-H fluorescence were selected to be targets (Figure S6c). Alternatively, when

controllers constitutively expressed RFP, events with a high signal in the PE-CF594 (560 nm

excitation, 610/20 nm emission filter, Red) channel were distinguished as controllers while

cells with low signal were designated as targets.

*Induction protocol*

387  Cells were cultured in 10 mL LB broth in 50 mL Falcon tubes. $20\mu L$ of overnight cultures were

388  added to the media, which was supplemented with the inducer(s) at the specified

389  concentration. The culture was then incubated at 37 $^{\circ}C$, shaking at 250rpm for 3h. Then, the

390  samples were prepared and analysed by flow cytometry. This protocol was used to obtain

391  Figures 2 a,b.

*Time-course assays*

35 $\mu L$ of overnight cultures of controllers and 140 $\mu L$ of overnight cultures of targets were

separately diluted each in 35 mL LB added to a 250mL sterile flask. Each culture was

supplemented with the specified concentration of IPTG. Both cultures were incubated at 37

$^{\circ}C$, shaking at 250rpm for 6h. Each hour, 5 mL of both cultures were mixed in sterile 50 mL

tubes, which were then incubated alongside the single strain cultures. After 6h, 0.5 mL of all

mixed cultures were sampled and analysed via flow cytometry. Note that the hours in Figure

2c and Figure 3a,b refer to the time targets and controllers have been mixed for. Figure 2c

Figure 3a,b were obtained using this protocol.

*Titration assay*

402  For each condition, $8\mu L$ of overnight culture of targets and $2\mu L$ of overnight culture of

403  controllers were mixed in 10 mL LB broth in 50 mL Falcon tubes. Each culture was

404  supplemented with IPTG at the specified concentration. The cultures were then incubated at

405  37 $^{\circ}C$, shaking at 250rpm for 6h. Then, the samples were prepared and analysed by flow

406  cytometry. This protocol was used to obtain Figure 3 c.

*Consortium composition assay*

408  Overnight cultures of targets and controllers were diluted in 10 mL LB using different

409  targets:controllers ratios. Specifically, 15:1, 8:1, 4:1, 2:1, 1:1 ratios were created by adding $10\mu L$

410  of controllers and 150 $\mu L$, 80 $\mu L$, 40 $\mu L$, 20 $\mu L$, 10 $\mu L$ of targets, respectively. All cultures were

411　incubated at 37 °C shaking at 250rpm for 6h. Then, 0.5 mL of each culture was sampled and

412　analysed by flow cytometry. This protocol was used in Figure 2d, Figure 4a-d.

**Data availability**

415　The authors declare that the data supporting the findings of this study are available within

416　the paper and its Supplementary Information files or from the corresponding author on

417　reasonable request. The source data underlying all figures in the main text and supplementary

418　information are provided as a Source Data file.

**Acknowledgements**

The authors wish to acknowledge Gwen Brouwer and Abigail Smith for their support with

the cloning of the constructs needed for this work. Also, the authors also wish to

acknowledge the FACS facility hosted at the University of Bristol for their support with the

use of the Flow Cytometers.　M.d.B., L.M. wish to acknowledge support from the European

Project FET-OPEN Cosy-Bio (Grant Agreement number 766840). M.d.B. also wishes to

acknowledge support from the PNRR and PNC projects of the Italian Government. L.M.

acknowledges funding from the Engineering and Physical Sciences Research Council

(EPSRC, Fellowship EP/S01876X/1) and the Biotechnology and Biological Sciences Research

Council (BrisEngBio grant, BB/W013959/1). Finally, L.M., N.J.S., C.G. and M.d.B. wish to

acknowledge support form BrisSynBio, a BBSRC/EPSRC Synthetic Biology Research Centre

(BB/L01386X/1).

**Author Contributions**

435　M.d.B, L.M., N.J.S. and C.G. designed the research; D.S. designed the experiments; D.S. and

436　B.Scarried out the experiments; D.S. with support from M.d.B., L.M., N.J.S. and C.G. analyzed

437　the data; D.S. and M.d.B. wrote the manuscript with inputs from L.M., N.J.S. and C.G.

**Competing Interests**

441 The authors declare no competing interests.

443 References

[1] D. D. Vecchio, Y. Qian, R. M. Murray and E. D. Sontag, "Future systems and control research in synthetic biology," Annual Reviews in Control, vol. 45, p. 5–17, 2018.

[2] M. Filo, C.-H. Chang and M. Khammash, "Biomolecular feedback controllers: from theory to applications," Current Opinion in Biotechnology, vol. 79, p. 102882, February 2023.

[3] M. Khammash, M. D. Bernardo and D. D. Bernardo, "Cybergenetics: Theory and Methods for Genetic Control System," in 2019 IEEE 58th Conference on Decision and Control (CDC), 2019.

[4] M. Chevalier, M. Gómez-Schiavon, A. H. Ng and H. El-Samad, "Design and Analysis of a Proportional-Integral-Derivative Controller with Biological Molecules," Cell Systems, vol. 9, p. 338–353.e10, October 2019.

[5] S. K. Aoki, G. Lillacci, A. Gupta, A. Baumschlager, D. Schweingruber and M. Khammash, "A universal biomolecular integral feedback controller for robust perfect adaptation," Nature, vol. 570, p. 533–537, June 2019.

[6] I. Ruolo, S. Napolitano, D. Salzano, M. di Bernardo and D. di Bernardo, "Control engineering meets synthetic biology: Foundations and applications," Current Opinion in Systems Biology, vol. 28, p. 100397, December 2021.

[7] C. Briat, C. Zechner and M. Khammash, "Design of a Synthetic Integral Feedback Circuit: Dynamic Analysis and DNA Implementation," ACS Synthetic Biology, vol. 5, p. 1108–1116, July 2016.

[8] T. Frei, C.-H. Chang, M. Filo, A. Arampatzis and M. Khammash, "A genetic mammalian proportional–integral feedback control circuit for robust and precise gene regulation," Proceedings of the National Academy of Sciences, vol. 119, June 2022.

[9] G. Fiore, A. Matyjaszkiewicz, F. Annunziata, C. Grierson, N. J. Savery, L. Marucci and M. di Bernardo, "In-silico Analysis and Implementation of a Multicellular Feedback Control Strategy in a Synthetic Bacterial Consortium," ACS Synthetic Biology, vol. 6, p. 507–517, December 2016.

[10] F. Bullo, J. Cortés and S. Martinez , Distributed control of robotic networks: a mathematical approach to motion coordination algorithms., Princeton University Press, 2009.

[11] P. Bittihn, M. O. Din, L. S. Tsimring and J. Hasty, "Rational engineering of synthetic microbial systems: from single cells to consortia," Current Opinion in Microbiology, vol. 45, p. 92–99, October 2018.

[12] A. Guarino, D. Fiore, D. Salzano and M. di Bernardo, "Balancing Cell Populations Endowed with a Synthetic Toggle Switch via Adaptive Pulsatile Feedback Control," ACS Synthetic Biology, vol. 9, p. 793–803, March 2020.

[13] D. Fiore, D. Salzano, E. Cristobal-Coppulo, J. M. Olm and M. di Bernardo, "Multicellular Feedback Control of a Genetic Toggle-Switch in Microbial Consortia," IEEE Control Systems Letters, vol. 5, p. 151–156, January 2021.

[14] E. Weill, V. Andreani, C. Aditya, P. Martinon, J. Ruess, G. Batt and F. Bonnans, "Optimal control of an artificial microbial differentiation system for protein bioproduction," in 2019 18th European Control Conference (ECC), 2019.

[15] F. Annunziata, A. Matyjaszkiewicz, G. Fiore, C. S. Grierson, L. Marucci, M. di Bernardo and N. J. Savery, "An Orthogonal Multi-input Integration System to Control Gene Expression in Escherichia coli," ACS Synthetic Biology, vol. 6, p. 1816–1824, July 2017.

[16] B. Shannon, C. G. Zamora-Chimal, L. Postiglione, D. Salzano, C. S. Grierson, L. Marucci, N. J. Savery and M. di Bernardo, "In Vivo Feedback Control of an Antithetic Molecular-Titration Motif in Escherichia coli Using Microfluidics," ACS Synthetic Biology, vol. 9, p. 2617–2624, September 2020.

[17] G. Perrino, A. Hadjimitsis, R. Ledesma-Amaro and G.-B. Stan, "Control engineering and synthetic biology: working in synergy for the analysis and control of microbial systems," Current Opinion in Microbiology, vol. 62, p. 68–75, August 2021.

[18] N. S. McCarty and R. Ledesma-Amaro, "Synthetic Biology Tools to Engineer Microbial Communities for Biotechnology," Trends in Biotechnology, vol. 37, p. 181–197, February 2019.

[19] K. J. Åström and R. M. Murray, Feedback Systems An Introduction for Scientists and Engineers, Second Edition, Princeton University Press, 2021, p. 528.

[20] V. A. Rhodius, T. H. Segall-Shapiro, B. D. Sharon, A. Ghodasara, E. Orlova, H. Tabakh, D. H. Burkhardt, K. Clancy, T. C. Peterson, C. A. Gross and C. A. Voigt, "Design of orthogonal genetic switches based on a crosstalk map of σ anti- σ and promoters," Molecular Systems Biology, vol. 9, p. 702, January 2013.

[21] K. E. McGinness, T. A. Baker and R. T. Sauer, "Engineering Controllable Protein Degradation," Molecular Cell, vol. 22, p. 701–707, June 2006.

[22] K. M. McKinnon, "Flow Cytometry: An Overview," Current Protocols in Immunology, vol. 120, January 2018.

[23] S. Oehler, E. R. Eismann, H. Krämer and B. Müller-Hill, "The three operators of the lac operon cooperate in repression.," The EMBO Journal, vol. 9, p. 973–979, April 1990.

[24] D. Salzano, D. Fiore and M. di Bernardo, "Ratiometric control of cell phenotypes in monostrain microbial consortia," Journal of The Royal Society Interface, vol. 19, July 2022.

[25] N. E. Grandel, K. R. Gamas and M. R. Bennett, "Control of synthetic microbial consortia in time, space, and composition," Trends in Microbiology, vol. 29, p. 1095–1105, December 2021.

[26] I. de Cesare, D. Salzano, M. di Bernardo, L. Renson and L. Marucci, "Control-Based Continuation: A New Approach to Prototype Synthetic Gene Networks," ACS Synthetic Biology, vol. 11, p. 2300–2313, June 2022.

[27] E. Pedone, I. de Cesare, C. G. Zamora-Chimal, D. Haener, L. Postiglione, A. L. Regina, B. Shannon, N. J. Savery, C. S. Grierson, M. di Bernardo, T. E. Gorochowski and L. Marucci, "Cheetah: A Computational Toolkit for Cybergenetic Control," ACS Synthetic Biology, vol. 10, p. 979–989, April 2021.

[28] I. Y. Hwang and M. W. Chang, "Engineering commensal bacteria to rewire host–microbiome interactions," Current Opinion in Biotechnology, vol. 62, p. 116–122, April 2020.

[29] Y. Arif, P. Singh, H. Siddiqui, A. Bajguz and S. Hayat, "Salinity induced physiological and biochemical changes in plants: An omic approach towards salt stress tolerance," Plant Physiology and Biochemistry, vol. 156, p. 64–77, November 2020.

[30] T. S. Bayer, D. M. Widmaier, K. Temme, E. A. Mirsky, D. V. Santi and C. A. Voigt, "Synthesis of Methyl Halides from Biomass Using Engineered Microbes," Journal of the American Chemical Society, vol. 131, p. 6508–6515, April 2009.

[31] M. Filo, S. Kumar and M. Khammash, "A hierarchy of biomolecular proportional-integral-derivative feedback controllers for robust perfect adaptation and dynamic performance," Nature Communications, vol. 13, April 2022.

[32] V. Martinelli, D. Salzano, D. Fiore and M. di Bernardo, "Multicellular PI Control for Gene Regulation in Microbial Consortia," IEEE Control Systems Letters, vol. 6, p. 3373–3378, 2022.

[33] M. S. Guyer, R. R. Reed, J. A. Steitz and K. B. Low, "Identification of a Sex-factor-affinity Site in E. coli as," Cold Spring Harbor Symposia on Quantitative Biology, vol. 45, p. 135–140, January 1981.

[34] A. J. Smith, F. Thomas, D. Shoemark, D. N. Woolfson and N. J. Savery, "Guiding Biomolecular Interactions in Cells Using de Novo Protein-Protein Interfaces," ACS Synthetic Biology, vol. 8, p. 1284–1293, May 2019.

[35] N. Kylilis, Z. A. Tuza, G.-B. Stan and K. M. Polizzi, "Tools for engineering coordinated system behaviour in synthetic microbial consortia," Nature Communications, vol. 9, July 2018.

[36] D. F. Ackerley, Y. Barak, S. V. Lynch, J. Curtin and A. Matin, "Effect of Chromate Stress on Escherichia coli K-12," Journal of Bacteriology, vol. 188, p. 3371–3381, May 2006.

Figure 1 **Multicellular control schematic. A** Microbial consortium implementing a

multicellular control architecture. The two populations, i.e. controllers and targets, share

information using orthogonal quorum sensing channels (**u** and **y**), playing the role of the

control input and the output of the process, respectively. In addition, the controllers sense

the external stimulus labeled as **ref,** used to set up the reference of the feedback loop. **b.**

Biological implementation of the multicellular control architecture. The Synthetic Biology

Open Language (SBOL) notation is used to denote promoters, genes, promotion/inhibition

relationships and quorum sensing molecules. The shaded areas identify different functional

modules within each population.

Figure 2 **Characterization of the controllers and targets cell populations. a:** Average fluorescence of the controller population induced using different concentrations of 3-O-C6-HSL and IPTG. The data were collected after 3h from the initial inoculation. **b:** Average fluorescence levels of the target population induced using 3-O-C12-HSL. The data were collected after 3h from the initial inoculation. **c:** Schematic representation of the closed loop (left) and the open loop (right) configurations. **d:** Average fluorescence of the targets over a 6h time course in open (blue) and closed (yellow) loop. The solid dots and vertical bars represent the average and standard deviation over n=3 biological replicates, respectively. **e:** Normalized average fluorescence of the targets mixed with the controllers in closed loop when 0 $\mu$M and 50 $\mu$M IPTG were present in the growth media. The data were collected 6h after the initial inoculation. The fluorescence was normalized to the average fluorescence reached when no IPTG was added in the media. All data are averaged over n=3 biological replicates.

Figure 3 **Regulation of the targets fluorescence using open and closed loop control using**

**controllers2.0/open loop controllers 2.0. a.** Average fluorescence of the targets over a 6h

time course, using an open loop control architecture. The solid dots represent the average

and the vertical bars the standard deviation of the data over n=3 biological replicates. Each

trajectory is color coded and labelled with respect to the concentration of IPTG used in the

experiment. **b.** Average fluorescence of the targets over a 6h time course, using a closed loop

control architecture. The solid dots represent the average and the vertical bars the standard

deviation of the data over n=3 biological replicates. Each trajectory is color coded and

labelled with respect to the concentration of IPTG used in the experiment.  **c.** Average steady

state fluorescence of the targets over different IPTG concentrations. The blue (open loop)

and yellow (closed loop) vertical bars are centered on the average values and their

amplitude represented the standard deviation of the data over n=3 biological replicates. The

dashed lines represent the linear interpolation of the data obtained with IPTG concentrations

in the range [2 $\mu M$, 8 $\mu M$]. The shaded areas represent the confidence intervals of the linear

model prediction. All steady states are collected after 6h from the initial inoculation.

Figure 4 **Closed loop control robustness to changes consortium composition. a-d.**

Normalized average fluorescence in open (blue) and closed (yellow) loop across different

consortium compositions. The solid dots are the data collected over n=3 biological replicates.

Dots of the same size belong to the same replica    te. The data were fitted with a first order

polynomial. The fitting and its confidence intervals are represented with the solid lines and

the shaded areas, respectively. For each replicate, the data are normalized to the average

fluorescence reached in closed loop across all compositions. **e.** Comparison of the estimates

of the slopes of the linear fittings of data in panels **a-d** when a closed (yellow) or open (blue)

loop architecture was used. The solid dots represent the slope estimates and the horizontal

bars the 90% confidence interval on the estimates.