Stereotyped goal-directed manifold dynamics in the insular cortex

Itay Talpir[1] and Yoav Livneh[1*#]

[1]Department of Brain Sciences, Weizmann Institute of Science, Rehovot 76100, Israel

*Correspondence: yoav.livneh@weizmann.ac.il

#Lead contact: : yoav.livneh@weizmann.ac.il

Summary

The insular cortex is involved in diverse processes including bodily homeostasis, emotions, and

cognition. Yet we lack a comprehensive understanding of how it processes information at the

level of neuronal populations. We leveraged recent advances in unsupervised machine learning

to study insular cortex population activity patterns (i.e., neuronal manifold) in mice performing

goal-directed behaviors. We find that the insular cortex activity manifold is remarkably

consistent across different animals and under different motivational states. Activity dynamics

within the neuronal manifold are highly stereotyped during rewarded trials, enabling robust

prediction of single-trial outcomes across different mice, and across various natural and artificial

motivational states. Comparing goal-directed behavior with self-paced free consumption, we find

that the stereotyped activity patterns reflect task-dependent goal-directed reward anticipation,

and not licking, taste, or positive valence. These findings reveal a core computation in insular

cortex that could explain its involvement in pathologies involving aberrant motivations.

Introduction

The insular cortex (InsCtx) is involved in a variety of processes, ranging from basic functions including bodily homeostasis, emotions, motivation, decision making, social interactions, and sensory processing, to maladaptive conditions including drug addiction, chronic pain, anxiety, obesity, and eating disorders[1–8]. Yet we lack a comprehensive mechanistic understanding of the computations in InsCtx that underlie its contribution to so many diverse processes.

A central path to understanding the computations within and across brain regions lies in measuring and deciphering neural activity of large populations of neurons in different behavioral contexts[9–11]. Indeed, there has been a recent explosion of artificial intelligence and machine learning approaches to decipher neural activity at the level of large neuronal populations[12–14] . These methods involve exploration of the multi-dimensional structure of activity patterns, also known as the activity manifold. By doing so, these approaches have begun to provide insights into the core computations of neural circuits, such as motor cortex control of limb movement, thalamic representation of head direction, and entorhinal cortex representation of spatial position[15–19]. Most previous animal model studies of InsCtx activity in behaving animals have used various recording and analysis techniques, including fiber photometry[20–23], single neuron electrophysiology[24–31], and ensemble analyses of electrophysiological recordings from relatively small populations (~5-10 neurons)[32–37]. While these studies have substantially advanced our understanding of InsCtx, they have not fully leveraged these recent artificial intelligence and machine learning approaches. Therefore, there is potentially much to be gained from comprehensive unbiased investigations of activity patterns of large InsCtx populations, using these recent developments in neuroscience analytical tools.

Experimental investigation of the population activity manifold can be used to test predictions from theoretical studies making explicit assumptions regarding the precise computations in the network (e.g., aforementioned studies of head direction and spatial position[15–17,38–41]). Alternatively, unbiased analyses of the manifold structure using, e.g., unsupervised machine learning, can aim to infer computations from the revealed manifold structure and dynamics within. Our study here falls within the latter category.

Previous work has suggested that InsCtx activity on short-timescales (~seconds) represents multi-modal gustatory and interoceptive sensory stimuli[26,42–46], as well as multi-modal salient

external cues[24,25,31,36,44,47–50]. Further work has suggested that such activity may also convey emotional, motivational and valence information[20–22,51–53]. Longer timescale changes in InsCtx activity (minutes to hours) have been suggested to represent slow changes in the physiological state of the body[22,47,54–57], as well as specific positive and negative emotional states[20–22,51]. However, seeing as InsCtx integrates many different inputs and is involved in many diverse processes, understanding its function at the neuronal population level requires comprehensive analyses of the repertoire of its activity patterns (i.e., the neuronal manifold) and their dynamics. Doing so would help answer fundamental questions that remain unanswered. For example, how constrained are InsCtx population activity patterns by intrinsic factors vs. extrinsic factors, such as the current behavioral context? How are different external and internal behavioral variables represented concomitantly in InsCtx population activity patterns? Are there core computations that InsCtx performs across different contexts? If so, will they have distinct signatures in population activity space? Here, we leverage recent advances in unsupervised machine learning and topological data analysis to perform an unbiased investigation of InsCtx population activity during goal-directed behaviors.

Results

Insular cortex population activity manifold structure is stereotyped across different mice and motivational states

To investigate how neuronal activity patterns are structured in InsCtx, we analyzed two-photon calcium imaging data from previously published and unpublished datasets from layer 2/3 of mid-InsCtx (**Table S1**). We started with data from thirsty (water-restricted) mice performing an operant go/no-go visual discrimination task in which licking following three initially arbitrary visual cues (drifting gratings) leads to rewarding (water), aversive (1M NaCl), or neutral outcomes[55] (**Figure 1A-B**). We previously showed that most InsCtx neurons respond to aspects of this behavioral task with either increase or decrease in activity (e.g., visual cues, licking, reward consumption)[55]. Therefore, to detect any underlying manifold structure within the observed activity patterns, we compared the temporal similarities across all simultaneously recorded neurons. Importantly, points in the manifold represent population activity patterns at specific times (**Figure 1C-D**).

Previous studies have shown activity manifolds to be low dimensional during simple behavioral tasks, i.e., they can be described by relatively few variables as compared to the number of recorded neurons[10,13,15,16,41,58–60]. However, because InsCtx integrates multimodal information from somatosensory, auditory, visceral, gustatory, and other limbic regions[61,62], we wondered whether InsCtx activity would be similarly low dimensional. Importantly, in all our analyses we analyzed each dataset in its entirety, including all time-point from all trials, starting from a thirsty (or hungry) state, and ending in a quenched (or sated) state. We estimated the dimensionality of InsCtx population data[58,63] (**Figure S1A-B**), and found that across all datasets activity manifolds consistently had 6±1 dimensions (mean ± SD). This was independent of the specific parameters used to assess dimensionality and of the number of recorded neurons (**Figure 1E, S1C-E**). This consistently low intrinsic dimensionality, together with the diverse multi-modal inputs InsCtx receives, suggests substantial convergence and integration of information in InsCtx during goal-directed behavior.

Building on this finding, we used non-linear dimensionality reduction to unveil the low dimensional manifold structure of InsCtx activity[12,13,15–17,41,60,63]. Specifically, we used Laplacian Eigenmaps (LEM), due to its capability to effectively examine the local geometry of high-dimensional data[41,64,65]. This provided a finer-grain approximation, particularly beneficial when

studying dynamics. Using LEM, we reduced dimensionality to the mean estimated intrinsic
dimensionality across all datasets (6 dimensions). This uncovered a remarkably consistent
activity manifold structure across datasets and different mice (**Figure 1D, S1F**). To quantify
structure similarity, we used Topological Data Analysis, an emerging approach for quantitatively
describing complex data structures[12,15,66–68]. We measured structure similarity by assessing the
difference (i.e., divergence) between the distributions of topological features across datasets (see
procedure illustration in **Figure S2A**). This revealed that the low-dimensional manifold structure
remained consistent across different days and mice. Structural similarity was significantly lower
for shuffled datasets (**Figure 1F**). We further validated these findings using pairwise permutation
tests[69] (**Figure S2B-C**).

We next assessed whether InsCtx dimensionality and manifold structure are conserved across
different motivations. We compared data from thirsty (water-restricted) mice working for water
rewards and hungry (food-restricted) mice working for liquid food rewards within the same
behavioral task structure[50] (**Figure 1G**). Activity manifold dimensionality in hungry and thirsty
mice were quite similar (5.7±1 vs. 6±1; mean ± SD; **Figure 1H**). Moreover, activity manifold
structure in hungry and thirsty mice were similar (**Figure 1I**).

We wondered whether the consistent dimensionality and topological similarity we observed in
InsCtx across similar behaviors during different motivations merely reflect behavioral constraints
on brain-wide activity[70–72]. We compared the InsCtx activity manifold with the activity manifold
of visual cortical areas of mice performing the same behavioral task during the same hunger
motivation (primary visual cortex and postrhinal cortex[73]). We found that the dimensionality and
structure of the activity manifold in visual areas were both significantly higher and less
consistent than those in InsCtx (**Figure 1J-K, S3**), and this did not depend on the number of
sampled neurons (see Methods). We further confirmed this by examining the distribution of P-
values from pairwise comparisons of manifold structure similarity between datasets within each
group (**Figure S2D**). Notably, dimensionality and structure (topology) can be independent (e.g.,
two-dimensional datasets could exhibit either ring or sheet structures). As such, the fact that we
observed different structures (topology) between InsCtx and visual areas, cannot be explained by
the higher variability in the estimated dimensionality of the visual areas datasets.

Collectively, our findings show that the activity manifold in InsCtx (but not visual areas) is low-
dimensional and structurally similar across mice engaged in similar goal-directed behaviors. The

consistent InsCtx manifold structure across animals may indicate common encoded information.

Moreover, these characteristics likely do not result from brain-wide constraints on behavior, and

could thus be more specific to a subset of brain regions, including InsCtx

Insular cortex activity dynamics within the manifold are stereotyped across different mice

Based on previous work[22,24,50,52,55], we expected InsCx activity patterns on both short and long

time-scales to correlate with external and internal behavioral variables (e.g., rewarding/aversive

outcomes, satiety, arousal, etc.). We color-coded the manifold based on different experimental

parameters (**Figure 2A**, left and middle; **S4A**) and found that some time-points associated with a

given variable aggregated spatially within the activity manifold, while others were more

uniformly distributed. We developed an automated method to cluster time-points within the

manifold (termed here as "clustered structure"). This method clustered time-points that diverged

from the center of the manifold ('manifold clusters'), which was not assigned to any cluster

('central point cloud'; **Figure 2A**, right; **S4B-C**). Notably, we consistently found 7 manifold

clusters in 46/47 of analyzed datasets. Note that different cluster indices (represented by different

colors; **Figure 2A**) were randomly assigned during clustering, and therefore do not necessarily

map onto the same behavioral variables across datasets (see more below).

Using the clustered structure, we quantitatively assessed the relationship between the activity

manifold and external/internal behavioral variables by computing their mutual information

(**Figure S4D-E**; Methods). Three variables had higher mutual information with the activity

manifold than others: (1) pupil size in inter-trial intervals (proxy for ongoing arousal levels[74]),

(2) cumulative water rewards (proxy for slow changes in physiological state, i.e., water satiety[55]),

and (3) trials in which the mice correctly responded to reward predicting cues to receive water

rewards (referred to hereafter as "rewarded trials"; **Figure 2B**). Moreover, the mutual

information for these three variables was significantly higher than that of shuffled data (**Figure

2C**). These results are consistent with previous work showing changes in InsCtx activity that are

associated with rewards, arousal, and physiological state[20–22,50–52,55]. While we have previously

shown that water rewards shift population activity along a linear thirsty-quenched axis[55],

manifold clusters associated with the quenched state were different from those associated with

rewarded trials (**Figure S5A-B**). This suggests that there are also distinct representations of

water satiety and reward in InsCtx population activity.

We next investigated activity manifold dynamics associated with behavioral variables. We

represented each activity pattern by its manifold cluster, and then examined the temporal

sequence of clusters during different trials (**Figure 2D**). InsCtx neuronal populations displayed

somewhat stereotyped patterns of cluster transitions during rewarded trials, characterized by

movement from the central point cloud to a sequence of transitions between the same 2-3

manifold clusters (**Videos S1, S2**). To evaluate the consistency of these dynamics, we calculated

the pairwise similarity of cluster sequences across all trial types. Cluster sequences were highly

similar across rewarded trials, substantially more so than all other trial types (**Figure 2E**). We

confirmed these results using a different dimensionality reduction method (Isomap[75]; **Figure**

S6,S1F). We also directly compared this cluster sequence method to our previous method of

projecting activity onto linear thirsty-quenched axes[55]. As expected, the linear projection

captures less of the variance of activity within the manifold and is less consistent across trials

(**Figure S5C-D**).

Could this stereotyped sequence of activity patterns be used to decode trial outcomes across days

and mice? The challenge is that our clustering technique does not yield functionally consistent

cluster indices across mice. Thus, clusters associated with the same behavioral variable would

not have the same label across mice. We therefore developed a method for translating cluster

labels across dataset pairs (different days, different mice) to align activity dynamics between

them for decoding (see Methods). We first determined the probability of transitions between

different clusters in each dataset (**Figure 2F**, Transition matrix). We then translated the cluster

labels from one "reference dataset" to relabel the clusters transition probabilities of the other

dataset (**Figure 2F**, "Sorted transition matrix"). Finally, we used the translated cluster labels to

train the decoder and tested it on the other "reference dataset" (**Figure 2F**). Using this approach,

we could classify rewarded trials across different days (neurons were not aligned across days),

and even across different mice, with high accuracy. We achieved an average accuracy of

$75\pm13\%$ and $65\pm13\%$ (mean $\pm$ SD), respectively, which was ~5 times higher than chance (~12-

15%). Decoding accuracy for other trial types did not significantly surpass that of shuffled trial

labels (**Figure 2G**). We further validated these results using P-values derived from the shuffle

distributions for each pairwise decoding (**Figure S7**). These results underscore that rewarded

trials were consistently characterized by movement from the central point cloud to a sequence of

transitions between the same 2-3 manifold clusters (**Videos S1, S2**).

Importantly, decoding trial types on single trials using this method was substantially more
accurate than using average activity levels across all neurons, or using our previous approach of
projecting activity on linear axes[55] (**Figure S7**). Therefore, decoding accuracy for rewarded trials
results from the stereotyped *patterns* of activity and does not trivially reflect global changes in
activity levels.

198 Insular cortex activity dynamics within the manifold are stereotyped across different
199 motivational states and rewards

Given the remarkably stereotyped activity patterns we observed for rewarded trials in thirsty
mice, and the consistent InsCtx manifold structure during both hunger and thirst, we next asked
whether hungry mice would exhibit the similar dynamics with food rewards. We first visualized
the temporal sequence of InsCtx manifold clusters in hungry mice and found that it displayed
similar stereotyped sequences as in thirsty mice (**Figure 3A**). In contrast, activity sequences in
visual areas were very different, primarily tracking visual cues, independent of rewards (see also
Figure S3). We then tested whether we could decode trial outcome across different thirsty and
hungry mice, receiving water and food rewards, respectively. Decoding accuracy in *hungry* mice
was high when training the decoder on InsCtx activity from *thirsty* mice (67±23%, mean ± SD,
vs. ~12-15% chance). However, decoding accuracy was near chance when training the decoder
on activity from visual areas (11±12%; **Figure 3B**). Notably, this differential decoding was not
due to the clustering procedure as InsCtx and visual areas datasets both had 7 clusters. The
consistent activity dynamics in InsCtx across hunger and thirst suggest that these activity
patterns likely do not reflect salient sensory attributes of the reward that are different in water vs.
Ensure (e.g., taste, viscosity), or the underlying physiological need. Nevertheless, we cannot rule
out that other sensory features, common to Ensure and water rewards, could be represented in
these activity patterns.

We further tested the extent to which these activity patterns remain stereotyped in mice
performing similar behavioral tasks across additional natural and artificial motivational states.
Specifically, we examined InsCtx activity from mice in which artificial hunger or thirst were
induced by chemogenetic activation of hypothalamic AgRP "hunger neurons" or SFO[GLUT] "thirst
neurons" – the primary sensors of physiological deficits and actuators of relevant behavioral and
physiological responses[76–79]. We also analyzed InsCtx activity from mice in which we induced

hyper-osmotic thirst by administration of hypertonic saline[80]. These datasets gave us the unique

opportunity to compare, within each mouse, the activity during the natural motivation driven by

physiological need, with the artificially induced motivation.

We first visualized the sequences of manifold clusters and compared them during the natural

motivational state to after induction of the artificial motivational state. Sequence transitions

during rewarded trials were remarkably similar in both states (**Figure 3C,F,I**). We quantified this

using pairwise trial similarity before and after induction of the artificial motivation. For all

motivational states, the pairwise similarities remained similar between the natural and

artificially-induced states, with no significant changes (**Figure 3D,G,J**). We therefore trained a

decoder on datasets from hungry and thirsty mice, which were not part of the artificial

motivation experiments, and used these to test decoding before and after the artificial induction

of motivational states. Decoding accuracy for rewarded trials remained the same for both states,

and was significantly higher than chance (**Figure 3E,H,K**). These findings demonstrate that the

stereotyped activity patterns in InsCtx are not specific to a particular reward (including its

sensory properties), nor to the specific underlying motivational state. Importantly, closer

examination of activity patterns and of individual neurons, will likely reveal distinct

representations of different rewards and motivational states. Nevertheless, our analyses reveal a

common activity pattern, which is so robust as to enable single-trial decoding across individuals

and motivational states.

Insular cortex stereotyped activity dynamics reflect goal-directed reward anticipation, but

not licking, taste, or positive valence

The stereotyped activity patterns we observed in InsCtx across rewarded trials with different

rewards and motivations could reflect goal-directed behavior, licking, positive valence, or need

fulfilment. We next tested these different options.

In hungry and thirsty mice, we observed that the stereotyped cluster sequence during rewarded

trials appeared linked with anticipatory licking, as evidenced by sorting trials based on the onset

of anticipatory licking (**Figure 4A**). We thus quantified the correlation between the onsets of

sequential activity and anticipatory licking in individual trials (**Figure 4B**). These correlations

were notably high and statistically significant in hungry mice (0.5±0.13; mean ± SD) and thirsty

mice (0.47±0.15; **Figure 4C**). Additionally, the onset of anticipatory licking consistently

preceded the onset of sequential cluster activity, and importantly, occurred independently of

reward delivery (**Figure 4D-E**). Together with similar activity dynamics across food and water

rewards (**Figure 3**), this supports a dissociation from the sensory aspect of the reward. Thus,

these stereotyped patterns could potentially reflect licking per se, positive valence (independent

of the reward's sensory properties), or cue-driven goal-directed behavior.

To distinguish between these possibilities, we examined trials in which mice incorrectly licked to

cues that do not predict reward ("false-alarm" trials). Cluster sequences in false-alarm trials

resembled rewarded trials more than other trial types (**Figure 4F**). To further investigate this, we

directly compared cluster sequences for false-alarm and rewarded trials. We sorted false-alarm

trials based on the similarity of *licking* patterns to rewarded trial licking (**Figure 4G**). False-

alarm trials with similar licking patterns to rewarded trials also displayed similar cluster

sequence dynamics. We quantified this by assessing correlations between licking similarity and

cluster sequence similarity in false-alarms vs. rewarded trials, finding positive correlations in

9/11 datasets (**Figure 4H**). As mice are not actually rewarded in false-alarm trials, these results

support the conclusion that InsCtx stereotyped activity patterns do not reflect reward or positive

valence of cues or outcomes. Nevertheless, they could still trivially reflect licking, or cue-

associated reward expectation.

To differentiate between these possibilities, we compared cluster sequences in hungry mice

during reward consumption in the operant discrimination task vs. during free consumption in the

same context without predictive cues or task structure (head-fixed, visual cue monitor on without

cues). Notably, in both conditions mice had a similar physiological need, fulfilled using the same

action (licking), to obtain the same reward. We constructed 'pseudo-trials' during the free

consumption epochs by identifying significant breaks between lick bouts and analyzed the

corresponding sequences of manifold clusters. As expected, average population activity

increased during reward consumption in both task-structured trials and self-paced consumption

(**Figure S8**). However, activity dynamics during task engagement and free reward consumption

were distinctly different (**Figure 4I**). Quantitatively, pairwise trial similarity between operant

trials and free consumption trials was significantly lower than the pairwise similarity within

operant trials. (**Figure 4J, S8**). We verified that these results could not be explained by

differences in licking patterns (**Figure S8**). Specifically, we compared licking patterns across

task performance and free consumption, that are as similar as possible (and statistically indistinguishable). We still found the same dramatic differences in activity dynamics (**Figure S8**). However, we cannot rule out the existence of subtle differences in licking patterns between these two conditions. It seems unlikely that such subtle differences, rather than behavioral context, account for such dramatic differences in InsCtx population activity patterns. Future work could directly test this.

In summary, by comparing highly similar licking patterns across different behavioral contexts (rewarded trials, false-alarm trials, free reward consumption), we show that: (1) licking per se cannot explain the stereotyped activity patterns we observed (comparing rewarded trials vs. free reward consumption); and that (2) reward expectation regardless of reward receipt does explain the stereotyped activity patterns we observed (comparing rewarded trials vs. non-rewarded false-alarm trials). Taken together, these results suggest that the stereotyped sequential activity patterns we observed in InsCtx are closely related to learned anticipation of reward during goal-directed behavior, rather than licking, the sensory aspects of the reward, positive valence, or specific motivational states.

Discussion

We used unbiased unsupervised machine learning and topological approaches to discover that InsCtx population activity had consistent dimensionality, manifold structure, and dynamics across days, across different mice, and across different motivational states.

Most cellular level InsCtx neural recordings in behaving animals have focused on taste processing[26,43,44,46]. These studies revealed sequential temporal coding of tactile, chemosensory, and palatability information in individual neurons on the order of hundreds of milliseconds[26]. Small populations of 5-10 InsCtx neurons have been shown to exhibit metastable dynamics, transitioning between different states that can reflect different behavioral variables, each lasting tens to hundreds of milliseconds[33–35]. Other work using activity manipulations and bulk activity recordings has shown that InsCtx activity is related to taste, palatability, motivation, valence, and anxiety[20,22,32,36,52,53,81]. Future experiments using manifold analyses of large population activity should be designed specifically to capture these variables, whether on slow or fast time-scales. The faster stereotyped changes we discovered appear not to be related to licking, taste, positive valence, or general reward. These observations (**Figure 4**) may explain previous manipulation experiments, in which inhibiting InsCtx activity reduces task-related behaviors, while not affecting free consumption behavior[50,82]. Future work could test this interpretation.

A recent study found that activity of anterior InsCtx (deep layer) projections to the brainstem encodes motivational vigor[52]. Notably, the stereotyped activity patterns we described here in superficial layers 2/3 of mid InsCtx likely do not reflect motivational vigor per se, as they depended on behavioral task structure and did not change with partial satiation, which reduces motivational vigor[52]. Nevertheless, it will be important to understand how InsCtx can simultaneously represent different types of information, and to what extent they are eventually integrated before being relayed to downstream brain regions to affect behavior and bodily physiology. Indeed, different InsCtx projection populations (e.g., projecting to striatum, thalamus, or brainstem[8,61,62]), different InsCtx layers (e.g., superficial vs. deep), and different InsCtx subregions (e.g., granular/dysgranular/agranular, and anterior/mid/posterior), may convey different types of information to downstream regions.

We previously used linear projections in activity space to suggest that reward consumption

transiently shifts activity towards a future satiety state[55]. Here, using analyses of the entire

activity manifold we find stereotyped goal-directed reward anticipation activity patterns, which

are mostly independent of changes in ongoing activity that reflect physiological state. Thus,

while our previous hypothesis-driven work provided support to a dominant model of

interoception[5,7], our unbiased analyses here highlight a general computation InsCtx performs,

which could be important for many behaviors that do not directly involve interoception.

Activity dynamics on the manifold were remarkably consistent across individuals and

motivations, suggesting that these dynamics may reflect an important fundamental aspect of

InsCtx function. We speculate that this is related to the consistent involvement of InsCtx in

cravings during pathological conditions associated with aberrant reward processing, including

obesity, binge eating disorder, and drug addiction[1–4,8]. InsCtx has been suggested to encode the

anticipated interoceptive value, which when goes awry can lead to aberrant food or drug

cravings[1]. We thus speculate that the stereotyped goal-directed reward anticipation patterns we

discovered reflect this fundamental computation in InsCtx. As such, future work using the same

analytical framework in mouse models of pathological conditions, could test this idea to form a

deeper understanding of the role of InsCtx in pathological conditions that involve aberrant

motivations.

Limitations of the study

We developed and used one manifold analyses approach. Future studies should compare our

methods to other recent powerful methods[12]. Additionally, our analyses were limited by the

behavioral variables that were measured in our studies, and the specific behavioral context of

cue-driven reward consumption. Other parts of the manifold could be determined by the intrinsic

connectivity of the network, as has been previously suggested[35], and/or by other variables we did

not measure in these experiments[21,42]. Furthermore, it will be important to compare InsCtx

activity across different behavioral contexts, including those that do not involve reward

consumption, to assess its intrinsic dimensionality and structure of the manifold.

Acknowledgements

We thank Mark Andermann, Alon Rubin, Yarden Cohen, Sasha Devore, Stav Shtiglitz, Ayal Lavi, Yael Prilutski, Einav Litvak, Omer Izhaki, Inbar Perets, Daniel Deitch, Tom Talpir, Omer Richmond, Gal Elyasaf, Birgit Jickeli, and Daphna Nachmani for fruitful discussions and helpful comments on the manuscript. YL is supported by research grants from the Israel Science Foundation (ISF #860/21), the European Research Council (ERC StG #101039145), and the Center for New Scientists at the Weizmann Institute of Science.

Author contributions

IT and YL conceived, designed, and executed the study. IT analyzed the data. IT and YL wrote the manuscript.

Declaration of Interests

The authors declare no competing interests.

Figure 1 – InsCtx activity manifold structure is stereotyped across different mice and

motivational states

A. Schematic of the operant visual discrimination task.

B. Two-photon calcium imaging of InsCtx through a microprism: schematic coronal brain

section illustrating the approach, and example field-of-view.

C. Illustration of the approach to study InsCtx population activity.

D. Three representative planes of dimensionality-reduced neuronal activity over time, from three

different thirsty mice. Each point represents the population activity pattern in a 0.5 sec time bin.

Note the high qualitative similarity between different mice.

E. Estimation of intrinsic dimensionality for each individual dataset vs. shuffled data. The

estimated dimensionality was significantly lower than chance (P<$7 * 10^{-5}$, one-tailed Wilcoxon

signed-rank test).

F. Quantification of topological dissimilarity ($\beta 0$ features) between all datasets (see also **Figure**

S2). Notably, there was a significant difference when comparing either across days or across

mice to chance levels (P<0.0003, one-tailed Wilcoxon signed-rank test), while comparisons of

the topological dissimilarity across days and across mice yielded no significant differences

(P>0.1, two-tailed Wilcoxon rank sum test).

G. Same as in '**D**', for hungry mice. Different mice than in '**D**'.

H. Similar estimated intrinsic dimensionality across hungry and thirsty mice (P>0.4, two-tailed

Wilcoxon rank-sum test).

I. Similar activity manifold shape (topological dissimilarity, $\beta 0$ features) across hungry and

thirsty mice (P>0.25, two-tailed Wilcoxon rank-sum test).

J. Comparison of estimated intrinsic dimensionality in InsCtx and cortical visual areas (POR –

dark shading, V1 – light shading) in hungry mice performing the same behavioral task.

*P<0.015, one-tailed Wilcoxon rank-sum test.

K. Comparison of activity manifold shape (topological dissimilarity, $\beta 0$ features) between

InsCtx and visual areas in hungry mice performing the same behavioral task. **P<0.0035, one-

tailed Wilcoxon rank-sum test.

Figure 2 – InsCtx activity dynamics within the manifold are stereotyped across different mice

A. Example visualization of time-points on the activity manifold that are associated with certain behavioral variables (red points). Different colors in the clustered structure reflect assignment to manifold clusters (**Figure S4**). Each row depicts a different plane of the activity manifold.

B. Mutual Information between behavioral variables and the activity manifold. ***P<0.0004, NS: P<0.02, one-tailed Wilcoxon signed-rank test with Bonferroni correction.

C. The most informative variables in '**B**' exceeded chance levels, i.e., randomly shuffled labels (P<0.00095, one-tailed Wilcoxon signed-rank test).

D. Example cluster sequence trial dynamics in two different mice. Left: two representative planes of the clustered structure (mouse 1 appears in '**A**'). Right: example of the cluster sequence dynamics for different trial types and outcomes (same cluster colors as the structures on the left). Note that for each mouse, cluster indices (colors) are randomly selected during the clustering procedure.

E. Distribution of trial similarity (cosine similarity). ***P$<5 * 10^{-10}$, two-sample Kolmogorov-Smirnov test with Bonferroni correction.

F. Decoding procedure across datasets. Cluster labels are translated across datasets by aligning the order of the transition probability matrix between the different neuronal clusters. The translated dataset is then used for training to decode the different trial outcomes of the test dataset.

G. Decoding accuracy for different trial types. ***P<0.00075 for across days and across mice versus shuffle, NS: P>0.2, one-and two-tailed Wilcoxon rank-sum test with Bonferroni correction, respectively.

A-G. N=14 datasets from 5 mice

Figure 3 – InsCtx activity dynamics within the manifold are stereotyped across different motivational states and rewards

A. Visualization of reward cue trial dynamics in two hungry mice from InsCtx and visual areas.

B. Decoding accuracy of rewarded trials from hungry mice. The decoder was trained using datasets from InsCtx in thirsty mice or from visual areas in hungry mice. ***P<0.0002, **P<0.0065, NS: P>0.35. Dunn test with Bonferroni correction. N=9 datasets from 6 mice.

C. Left: Example activity dynamics of reward cue trials before and after artificial induction of hunger using chemogenetic activation of AgRP "hunger neurons". Red arrow: first responded reward cue trial following induction.

D. Distribution of pairwise trial similarity (cosine similarity) among rewarded trials across natural and artificially induced hunger. Left: distributions pooled from all datasets. Right: paired comparisons of the mean pairwise trial similarity for each dataset trials. NS: P>0.15, two-tailed paired t-test.

E. Comparison of the decoding accuracy of rewarded trials across natural and artificially induced hunger. Comparisons between natural and artificial data across all states: P>0.3. Comparisons between natural and artificial shuffles across all states: P>0.08. **P<0.008. One-tailed Wilcoxon signed-rank test with Bonferroni correction.

F-H. Same as in 'C'-'E' before and after artificial induction of thirst using chemogenetic activation of SFO "thirst neurons".

I-K. Same as in 'C'-'E' with before and after induction of hypertonic thirst using injection of hypertonic saline.

A-K. N=10 hungry datasets from 7 mice, N=14 thirsty datasets from 5 mice, N=7 visual datasets from 5 mice, N=4 AgRP activation datasets from 4 mice, N=4 SFO activation datasets from 4 mice, N=3 hypertonic saline injection datasets from 3 mice.

Figure 4 – Stereotyped activity dynamics reflect goal-directed reward anticipation, but not licking, taste, or valence

A. Example neuronal dynamics during rewarded trials and task-related licking behavior.

B. Example correlations between anticipatory licking onsets and rewarded trials cluster onsets in 6 datasets from hungry (top) and thirsty mice (bottom).

C. Left: Distribution of all correlations across all hungry (N=10) and thirsty (N=11) datasets. Right: Scatterplot of correlation levels vs. p-values. Dashed vertical line: P=0.05. P<0.04, for all datasets, Pearson's product-moment correlation.

D. Example visualization of reward onset, anticipatory licking onset, and activity cluster onset. Left: Cluster dynamics of all rewarded trials. Right: Onsets of anticipatory licking, reward delivery, and cluster dynamics per trial. Note that the onset of the sequential cluster pattern consistently follows the onset of anticipatory licking and appears unrelated to the onset of reward delivery.

E. Distribution of the time difference between rewarded trial cluster onset and anticipatory licking onset. Across all trials, cluster onset occurs approximately 1 second *after* lick onset.

F. Distribution of trial similarity between incorrectly responded aversive and neutral trials ("false-alarms") and rewarded trials (purple), compared to the similarity between correctly non-responded aversive and neutral trials ("correct-rejections") to rewarded trials (red). N=14 datasets from 5 thirsty mice.

G. Two examples of cluster sequence trial dynamics during false-alarm trials and rewarded trials. False alarm trials were sorted based on their *licking* pattern similarity to the *licking* patterns in rewarded trials. Note that as licking pattern becomes more similar, false-alarm cluster sequence trials become more similar to rewarded trials.

H. Distribution of all correlations between licking similarity and trial similarity in false-alarm trials and rewarded trials, as a function of p-value. 9/11 had positive correlations, 7/9 were statistically significant (P<0.05, Pearson's product-moment correlation).

I. Example cluster sequence dynamics and licking behavior in two mice during reward consumption in the operant task and during self-paced free consumption. Dynamics were aligned based on the onset of licking.

J. Left: Distribution of all pairwise comparisons between reward consumption trials during task engagement and free consumption behavior (N=5 datasets from 5 mice). Right: Comparison of average trial similarity for each dataset individually. **P<0.009, one-tailed paired t-test.

492   **STAR Methods**

493   **Key resources table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and Virus Strains		
AAV1-hSyn-GCaMP6f	Penn Vector Core	Addgene 100837
AAV8-hSyn-DIO-hM3Dq-mCherry	UNC Vector Core	Addgene 44361
AAV8-CaMKII-hM3Dq-mCherry	Addgene	Addgene 50476
Chemicals, Peptides, and Recombinant Proteins		
Clozapine-N-Oxide (CNO)	NIMH Drug Supply	Program Cat# C-929
Compound 21 (C21)	Tocris	6422
Experimental Models: Organisms/Strains		
Mouse: C57BL/6J	The Jackson Laboratory	RRID: IMSR_JAX:000664
Software and Algorithms		
R	RStudio	RRID: SCR_000432
MATLAB R2015b, 2019a	Mathworks	https://www.mathworks.com/products/matlab.html; RRID: SCR_001622
MonkeyLogic	Asaad and Eskandar, 2008	http://www.brown.edu/Research/monkeylogic/
MonkeyLogic2	NIMH	https://monkeylogic.nimh.nih.gov/
Scanbox	Neurolabware	https://scanbox.org/
PCA/ICA neuron identification	Mukamel et al., 2009	https://github.com/mukamel-lab/CellSort
TDAStats	Wadha RR et al, 2018	https://github.com/rrrlw/TDAstats
dimRed	Kraemer et al, 2018	https://github.com/gdkrmr/dimRed
intRinsic	Denti Francesco, 2023	https://github.com/Fradenti/intRinsic
Custom code in R	This manuscript	https://github.com/livnehlab/InsCtxManifolds ; https://doi.org/10.5281/zenodo.10632243

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to the lead

contact, Yoav Livneh (yoav.livneh@weizmann.ac.il).

Material availability

This study did not generate new unique reagents.

Data and code availability

The code used for the analyses in this work is available at the LivnehLab github
(https://github.com/livnehlab/InsCtxManifolds; https://doi.org/10.5281/zenodo.10632243).

Any additional information required to reanalyze the data reported in this paper is available from
the lead contact upon request.

All data reported in this paper will be shared by the lead contact upon request.

Experimental model and study participant details

509 All animal care and experimental procedures were approved by the relevant Institutional Animal
510 Care and Use Committee. Mice used for *in vivo* two-photon imaging (males, age at surgery: 9-15
511 weeks) were instrumented with a headpost and a 2 mm microprism, centered over the mid InsCtx
512 (see details below).

Method details

Datasets analyzed for this paper were both previously published data[50,55,73], as well as
unpublished data. Unpublished datasets included experiments similar to those described in the
published datasets (e.g., in water or food restricted mice performing an operant visual
discrimination task for water or Ensure rewards), as well as imaging during head-fixed satiation
by Ensure consumption and induction of hyperosmotic thirst by intraperitoneal injection of
hypertonic saline. We briefly describe the experiments below, see also refs[50,55] for more detailed
descriptions.

Behavioral task

To perform the behavioral task, mice were water restricted to ~80% of their pre-restriction body
weight or food-restricted to ~85% of their free-feeding body weight. We trained mice to
discriminate between square-wave drifting gratings differing in orientation presented on an LCD
screen (2 Hz and 0.04 cycles/degree, full-field square wave drifting gratings, 80% contrast; food
cue: 0°, aversive cue: 270°, neutral cue: 135°[50,55]. All drifting gratings were presented for 2 sec,
after which the mouse had a 2 sec window to respond with a lick. Licking during the visual cue
was not punished, but also did not trigger delivery of the water/Ensure/salt-water/quinine. Only
the first lick (if any) occurring during the response window triggered delivery of
water/Ensure/salt-water/quinine. The lickspout was designed with two adjacent lick tubes (one
for each outcome), such that the tongue contacted both tubes on each lick, which served as an

effective deterrent for lick responses following aversive cues. Well-trained mice had a high rate

of correct water/food cue licking responses (criterion: >80% of trials, usually ~90-95%), and a

low rate of licking following aversive cue presentations (criterion: <50%, usually ~20-30%).

Each water reward was a ~2-3 µL drop, and Ensure reward was a ~5 µL drop (0.0075 calories).

Behavioral training was performed using MonkeyLogic[83] and MonkeyLogic2

(https://monkeylogic.nimh.nih.gov/).

Surgical procedures

539   Stereotaxic injections AAV8-CaMKII-hM3Dq-mCherry, AAV8-hSyn-DIO-hM3Dq-mCherry,

540   and AAV1-hSyn-GCaMP6f, as well as and implantation of microprisms (2 mm prisms; #MCPH-

541   1.0; Tower Optical; coated with aluminum along their hypotenuse) were performed as previously

542   described[50,55].

Two-photon imaging across different natural and artificial states

Two-photon imaging of GCaMP6f was performed using a resonant-scanning two-photon

microscope with tiltable scanhead (Neurolabware; 31 frames/second; 1154x512 pixels). All

imaging was performed with a 20x 0.45 NA air objective (Olympus) with a 540 x 360 µm2 field

of view. All imaged fields of view (FOV) were at a depth of 90-200 µm below the pial surface,

using a Mai Tai DeepSee laser (Newport Corp.) with laser power at 920-960 nm of 35-80 mW at

the front aperture of the objective (power at the sample was likely substantially less due to partial

transmission via the microprism). Imaging depth was adjusted in between runs (every 30 min) to

account for slow drift in the z plane (< 7 µm). Recording locations were approximately +0.5 mm

to -1.0 mm to the middle cerebral artery (see[50,55] for further details).

Imaging across thirsty and quenched states

554   We imaged mice during gradual water satiation during consecutive 30-minute runs until the mice

555   voluntarily stopped performing the task. We then performed one more imaging run, the

556   'quenched' run, in which mice did not respond to the water cue.

Imaging across hungry, satiation, and sated states

558   We imaged mice in two blocks of trials within a session, one block during food restriction and a

559   subsequent block immediately following re-feeding. At the start of each imaging session, food-

560   restricted mice performed the visual cue discrimination task. After ~180 trials (30-min imaging

run), we provided the mouse with ad libitum access to Ensure until voluntary cessation of consumption. Ensure consumption lasted 45-75 minutes. We triggered delivery of Ensure with every lick, but with a minimum inter-trial interval of 2.5 s between Ensure deliveries. During this period of time, mice consumed ~3-5 mL of Ensure and then voluntarily stopped licking for rewards. We then imaged additional ~180 trials (30 min imaging run) while mice were satiated (operationally defined as the absence of voluntary licking).

Imaging during chemogenetic activation of SFO^{GLUT} neurons

Following imaging during the quenched state (see above), we injected 150 µl 0.9% of either saline or CNO (5 mg/kg), waited 10-15 min and started another imaging run (~180 trials, 30 min). For every mouse used for these experiments, we used postmortem histology and immunohistochemistry to verify hM3Dq-mCherry expression in the SFO.

Imaging during following injection of hypertonic saline to induce hyperosmotic thirst

Following imaging during the quenched state, we injected ~200 µL of 2M NaCl, waited 5-10min and started another imaging run (~180 trials, 30 min). Mice usually re-engaged in the behavioral task immediately or after 2-3 minutes.

Imaging during chemogenetic activation of AgRP neurons

Following imaging during hunger and satiety, mice were returned to their home-cage with ad libitum access to regular chow. The next morning, we imaged the same InsCtx field of view in this satiety state (100-110% of normal body weight) during ~180 trials (30 min.). We then injected CNO (1-3 mg/kg). Ten minutes later, we initiated an additional imaging run of ~180 trials. For every mouse used for these experiments, postmortem histology and immunohistochemistry confirmed hM3Dq-mCherry expression in the hypothalamic arcuate nucleus.

Pupil videography during two-photon imaging

We acquired data using a GigE Vision camera (Dalsa) with a 60 mm lens (Nikon MicroNikkor) from a pre-selected region of interest around the eye ipsilateral to the LCD monitor used to present visual cues (contralateral to the InsCtx microprism). Acquisition of each frame (frame rate of 15.5 Hz) was triggered on every other frame of two-photon acquisition (acquired at 31 Hz) using Scanbox software (Neurolabware). The pupil was backlit with illumination originating

590 from diffusion within the brain of the IR light used for two-photon excitation during imaging.

591 See below for details of data analysis.

Image registration and time course extraction

First, each acquired image was spatially down sampled by 2X. To correct for motion along the

imaged plane (x-y motion), each frame was registered to an average field-of-view using efficient

subpixel registration methods[84] Within each imaging session, each run (2-8 runs/session) was

registered to the first run of the day. Image stacks were de-noised using principal components

analyses (PCA) of every pixel across time, and by user identification and removal of noise

principal components (low eigenvalues; based on[85]). Cell masks and calcium activity time

courses ('F(t)') were extracted using custom implementation of common methods[85]. To avoid

use of cell masks with overlapping pixels, we only included the top 75% of pixel weights for a

given mask, but users screened each prospective ROI and could edit the size of the mask,

selectively removing the lowest probability pixels. We then excluded any remaining pixels

identified in multiple masks. We manually verified that all cell masks had typical cell body

morphology and size.

Fluorescence time courses were extracted by averaging the pixels within each region-of-interest

('ROI') mask. Fluorescence time courses for neuropil within a 25 µm annulus surrounding each

ROI (but excluding adjacent ROIs and a protected ring surrounding each ROI) were also

extracted ($F_{neuropil}(t)$: median value from the neuropil ring on each frame). Fluorescence

timecourses were calculated as $F_{neuropil\_corrected}(t) = F_{ROI}(t) - F_{neuropil}(t)$. The change in fluorescence

was calculated by subtracting a running estimate of baseline fluorescence ($F_0(t)$) from

$F_{neuropil\_corrected}(t)$, then dividing by $F_0(t)$: $\Delta F/F(t) = (F_{neuropil\_corrected}(t) - F_0(t))/ F_0(t)$. $F_0(t)$ was

estimated as the 10th percentile of a 32 sec sliding window[50,55,73].

Preprocessing of neuronal data

For each collected dataset, we down-sampled $\Delta F/F$ traces by averaging every consecutive 15

frames (acquired at 31 Hz), resulting in ~0.5 seconds time-bins. Subsequently, we excluded

outlier neurons with abnormally high $\Delta F/F$ values, calibrated separately for two different

microscopes (2000% and 5000%). Following this, we Z-scored the activity time course for each

individual neuron. Each dataset consisted of multiple 30-minute runs, acquired within the same

day. For certain datasets, we applied separate Z-scoring to each run if substantial extended
breaks occurred between runs or if there was a z-plane shift between the runs.

Intrinsic dimensionality estimation

To estimate the intrinsic dimensionality of the neuronal activity data, we employed the minimal

neighborhood information technique as an estimator[86]. Briefly, the intrinsic dimension is

estimated using the ratio formula:

$$intrinsic\ dimension = -\frac{\log(1 - F(\mu))}{\log(\mu)}$$

Here, μ_i was determined for each data point i, where $\mu_i = \frac{r_{1,i}}{r_{2,i}}$, with $r_{1,i}$ and $r_{2,i}$ representing the

Euclidean distances of the nearest and second-nearest data points to point i, respectively. $F(\mu_i)$

indicates the percentile of μ_i with respect to all other data points. Subsequently, μ_i was computed

for each data point, and the highest 4% of points were excluded. The intrinsic dimension was

defined as the slope calculated from a linear regression between $\log(1 - F(\mu))$ and $\log(\mu)$

across all data points. To better estimate the intrinsic dimensionality of the neuronal activity data,

we used dimensionality reduction for denoising prior to dimensionality estimation. In essence,

dimensionality estimation remains invariant to the embedding dimension. To ensure this, we

iteratively assessed the intrinsic dimension across intermediate dimensionality reductions (see

Figure S1B-E). We further verified this in certain datasets where the number of neurons was

lower (specifically in visual areas from hungry mice; see **Table S1**). In visual areas datasets, we

compared the estimated dimensionality with InsCtx hungry-sated datasets (see **Figure 1**). To

ensure that dimensionality was unaffected by the number of neurons, we repeatedly randomly

subsampled to the lowest number of neurons (see **Table S1**). This confirmed that the estimated

dimensionality was consistently lower in InsCtx vs. visual areas. Specifically, for visual areas,

the average estimated dimension was 11.7±3 (mean ± SD) for both subsampled and full data. In

contrast, hungry-sated datasets, this resulted in 7.7±2 and 8.7±2 for subsampled and full data,

respectively.

Non-linear dimensionality reduction

To reduce the dimensionality of the neuronal data matrix X, we applied Laplacian eigenmaps[64]

implemented by a pre-established pipeline[87]. Laplacian eigenmaps, a spectral non-linear

dimensionality reduction technique, operates on a data matrix X with dimensions $N \times T$, by

generating a weighted adjacency graph W for the T high-dimensional data points such as each

time point $t \in \mathbb{R}^N$. This adjacency graph is computed for all pairs of data points, where the

weight between a pair of data points x_i, $x_j \in X$ is assigned as $W_{ij} = 1$ if the data points are

connected and $W_{ij} = 0$ otherwise. Data points are considered connected if they fall within the K

nearest neighbors of each other, where K is a user-defined parameter (see below). Following the

computation of W, a diagonal weight matrix D is constructed with $D_{ii} = \sum_j W_{ji}$, which is

equivalent to aggregating the rows of W. Notably, the Laplacian of the original data matrix is

$L = D - W$. The process of obtaining the low-dimensional embedding is achieved by calculating

the eigenvalues and eigenvectors of $L\boldsymbol{f} = \lambda D\boldsymbol{f}$. After excluding the leading eigenvector[64], the

following m eigenvectors are utilized to obtain a new low-dimensional mapping for each data

point x_i, which is given by[17]

$$x_i \rightarrow (f_1(i), \dots, f_m(i))$$

with m representing the desired output dimensionality. For every dataset X, we conducted two

rounds[41] of dimensionality reduction using the algorithm described above. In the initial iteration,

a 20-dimensional matrix was generated, with a parameter K set to 7.5% of the time points in

matrix X. The second iteration was performed on the 20-dimensional matrix obtained from the

preceding step. This culminated in the matrix reaching its ultimate dimension – the mean

estimated intrinsic dimension of 6 – with a parameter K set to 2.5% of the time points in matrix

X. The mean estimated intrinsic dimension of 6 was derived from the process described above.

Topological features analysis and topological similarity

To assess the topological features of the data, we quantified for each dataset its components (β0)

and holes (β1) as a function of the radius threshold, using a pre-established algorithm[67]

(TDAStats: R pipeline for computing persistent homology in topological data analysis,

https://github.com/rrrlw/TDAstats). Due to computational considerations, we determined the

topological characteristics of clustered timepoints within the reduced data matrix, acquired

through K-means clustering with a chosen K value of 80. Given our utilization of a stochastic

variant of K-means, this process was repeated 20 times for each dataset. For every topological

feature extracted from the clustered data matrix, we quantified its lifespan and derived the

lifespan distribution for all topological features. The topological resemblance between a pair of

datasets was defined as the mean Wasserstein distance[88] between the lifespan distribution of
features from both datasets, covering all 20 iterations of K-means clustering. To ascertain the
significance of differences between dataset pairs, permutation tests were conducted, following a
methodology outlined in prior work[69]. The lifespan distributions of a pair of datasets were pooled
and then systematically rearranged to generate two separate subsamples, adhering to the original
sizes of each distribution. The Wasserstein distance was computed between these subsamples.
This procedure was repeated n=200 times, and a P-value was established as the percentage of
permutations yielding a Wasserstein distance less than the original distance between the dataset
pairs. The ultimate P-value for every pair of datasets was determined as the average of the P-
values derived from this process across all 20 K-means clustering iterations.

Clustering the activity manifold

To evaluate the activity manifold, our objective was to create a standardized comparison across
datasets through an automated clustering approach. To achieve this, we devised an unsupervised
clustering strategy that recognizes data points that deviate from the central point cloud, and thus
classifies them as distinct neuronal clusters. To do so we systematically excluded time points that
were closest to the centroid of the reduced data matrix. Starting from 40 times points and
progressing to the entire dimensionality of the data matrix. For each set of excluded timepoints,
we conducted 20 iterations of K-means clustering with K values spanning from 2 to 20. The
mean squared error (MSE) was computed between the resulting clusters and their centroids
across all iterations and for each K value. This procedure produced a two-dimensional grid with
its two dimensions being the fraction of excluded timepoints and the chosen K values for K-
means clustering. The cells of the grid contained MSE values over the 20 clustering repetitions.
To identify the optimal configuration, we scaled both grid dimensions to 1 and determined the
grid cell closest to the origin of a three-dimensional Cartesian system formed by the grid, where
the third dimension was the MSE value within each cell. This process yielded, for each dataset
an optimal configuration consisting of 2 parameters: a K value, and the number of datapoints to
be considered as the central point cloud. Across virtually all utilized datasets the identified
optimal K value was 8, except one dataset in which it was 7. After obtaining the optimal
configuration, our aim was to derive deterministic clustering labels for each dataset. This was
accomplished by implementing the optimal configuration, isolating timepoints deemed part of
the central point cloud, and executing 500 iterations of K-means clustering with the optimal K

value for the remaining timepoints. The resulting clustering labels were aggregated into a
$500 \times T$ matrix. This matrix was then subjected to an additional clustering step using the ward
agglomerative complete linkage hierarchical clustering method[89]. This yielded the final labels of
the clustered activity manifold.

Activity manifold parameterization

To evaluate the contribution of behavioral variables to the activity manifold, we devised labeling
for the distinct behavioral variables, considering the total timepoints under analysis. These
variables were categorized into two groups: discrete (binary) and continuous. Discrete binary
variables encompassed the potential classes of trial outcomes within the behavioral task (e.g.,
response vs. no response for a given trial type). For each of the six trial outcome classes, a
corresponding label vector was generated, spanning T timepoints. A value of
$Trial\ outcome\ Label = 1$ was assigned for timepoints ranging from 3 to 6 seconds after the
cue onset for each class, while a value of $Trial\ outcome\ Label = 0$ was allocated to the
remaining timepoints. This enabled the isolation of timepoints exclusively linked to that specific
trial outcome, excluding those related to visual cues. As for continuous variables, namely Pupil
Size and Cumulative Consumed Rewards, their magnitude across the entire experiment was
quantified. Pupil size was quantified using code from previously published work[90]. Two label
vectors, each with a length of T, were established for these variables. To ensure equitable
comparison with discrete variables, both Pupil Size and Cumulative Consumed Rewards were
divided into eight bins.

Mutual information between behavioral variables and the activity manifold

To determine the mutual information between behavioral variables and the activity manifold, we
constructed a joint contingency table that related the two label vectors: $Cluster$ and $Behavior$.
Here, $Cluster$ represents a vector of size T containing different neuronal cluster labels
(including the central point cloud), while Behavior was a T-sized vector representing behavioral
variables (ranging from 0 to 1 for discrete variables and 1 to 8 for continuous variables).
Following this, we utilized the contingency table to formulate a joint probability space, denoted
as $P_{Cluster \times Behavior}$. The calculation of mutual information was carried out according to the
formula:

Where $P_{Cluster}$ and $P_{Behavioor}$ are the respective marginal probability distributions. To reduce

potential variations caused by different behavioral factors and ensure an equitable comparison

with discrete variables, for the continuous variables we exclusively included time points

unaffected by behavior, following established procedures from our previous work[55]. These time

points corresponded to the final 3 seconds of inter-trial intervals following the correct rejection

of aversive and neutral trials (referred to as non-responded trials).

Trial structure manifold dynamics

To evaluate manifold dynamics in relation to the trial structure, we compiled the sequential

neuronal cluster label (including the central point cloud) time course for every trial. This was

achieved by capturing a 10-second interval from the onset of the cue for each trial class. They

yielded a sequence of 20 neuronal cluster labels, subsequently combined into an $M \times 20$ matrix.

Here, M denotes the trial count, and each row signifies the sequence of visited clusters during the

respective trial.

Trial similarity

To assess trial dynamics similarity, our focus was on highlighting the sequence of neuronal

clusters. To achieve this, we designated the label -1 to each timepoint associated with the central

point cloud. Meanwhile, other timepoints were allocated labels ranging from 1 to 7,

corresponding to the respective neuronal cluster assignments. Subsequently, we created a trial

structure matrix as previously outlined and computed pairwise trial similarity using cosine

distance.

$$Trial\ Similarity(Trial_i, Trial_j) = \frac{Trial_i \cdot Trial_j}{\|Trial_i\| \, \|Trial_j\|}$$

Here, $Trial_i$ represents the timepoint sequence of labels in the ith trial. This procedure resulted

in a $M \times M$ for each dataset. Following that, we concentrated on the lower triangle of the matrix

and pooled all values across all datasets.

Decoding trial outcome across datasets

To enable across-datasets decoding, we generated a trial dynamics matrix using the same

labeling approach employed for assessing trial similarity. Our goal was to enable decoding

between datasets, which necessitated an alignment process. For each dataset, we initiated

alignment by constructing a transition probability matrix between all neuronal clusters and the

central point cloud. This process yielded a 8×8 probability matrix for each dataset, except for

one dataset with 7 clusters that was excluded from this analysis. Following this, the probabilities

underwent clustering through the ward agglomerative complete linkage hierarchical clustering

method[89]. Using the hierarchical clustering structure, we derived a cluster order based on the

similarity of their probabilities. This order served as a mask across datasets to allow translation

of clustering labels among dataset pairs. Essentially, this step aimed to address the question:

Given transition probabilities, which cluster in dataset A most closely aligns with a cluster in

dataset B. With the labels clusters of dataset B aligned to dataset A, we proceeded to compute the

pairwise trial similarity between each trial pair in both datasets. For each trial a in dataset A, the

decoded class of trial outcome was determined by maximizing the sum of Trial Similarity values

over trials b in dataset B:

$$argmax_{class} \sum_{b \in B} Trial\ Similarity(Trial_{b(class)}, Trial_a)$$

This process is analogous to identifying, on average after translation, which class of trials in

dataset B most closely resembles an individual trial in dataset A. To further validate the decoding

accuracy, we established a surrogate shuffle distribution by randomly shuffling trial classes in

both datasets 200 times. We then calculated the decoding accuracy for each shuffle and

computed the P-value by determining the percentile of shuffles that exhibited a higher decoding

accuracy compared to the measured decoding accuracy. This entire process was repeated for

every pairwise combination of datasets.

Decoding trial outcome across datasets based on overall activity levels and thirsty-
quenched linear axis

To decode trial outcomes across dataset based on overall activity levels, we calculated the

overall activity levels within each trial by averaging the collective population activity across

neurons. To decode trial outcomes across dataset based on the thirsty-quenched linear axis, we

defined this axis for each mouse by $\bar{x}_{Thirsty} - \bar{x}_{Quenched}$, where $\bar{x}$ is the population vector of

mean ongoing activity in each state. We projected peri-cue activity onto this axis by calculating

the dot product of this vector with the time-varying pattern of InsCtx population activity, x(t).

We then scaled values along this axis per mouse, ascribing a value of 1 when x(t) = $\bar{x}_{Thirsty}$, a

value of 0 when x(t) = $\bar{x}_{Quenched}$ as 0, and intermediate values for patterns that fall between

$\bar{x}_{Thirsty}$ and $\bar{x}_{Quenched}$ projected this data onto a thirsty-quenched linear axis, as outlined in

reference[55]. Second, we utilized a similar decoding as explained above. Instead of training it with

aligned trial labels for pairs of trials, we evaluated used the overall activity levels within each

trial or the projection of the trial onto the thirsty-quenched axis as a training dataset.

Quantification of anticipatory licking onset

796   The onset of anticipatory licking during a trial was defined as the time point when the first

797   licking bout took place within the response window after cue presentation. Licking bouts were

798   considered as instances where mice notably increased their licking rate. To detect these bouts, for

799   each trial, we calculated the time intervals between consecutive licks occurring after cue

800   presentation. We then identified the pair of licks with intervals shorter than a specific threshold:

$$\frac{1}{\mu_{Cue\ licks} + C * \sigma_{Cue\ licks}}$$

801   Where $\mu_{Cue\ licks}, \sigma_{Cue\ licks}$ were determined by calculating the average and standard deviation of

802   licks based on the first 15 frames (0.5 sec) of cue presentation across all trials. The constant C

803   varied; it was set as 0.1 for datasets involving water rewards and 0.35 for datasets involving food

804   rewards. The time point at which the first pair of licks exceeded this threshold was considered

805   the onset of anticipatory licking.

Quantification of cluster onset

To identify the onset of the cluster sequence, we first determined the two neuronal clusters with

the highest occupancy during the response window for each dataset. Following this, we

identified the initial time point within each trial where a transition occurred from the central

point cloud to each of the identified clusters. The onset was defined as the first time point in each

trial where the transition occurred with the cluster that exhibited the highest onset correlations

with the lick onsets across all trials.

Lick similarity between pairs of trials

814   To define the lick similarity between pairs of trials, for each trial, we smoothed the binary licks

815   vector $Licks_i$. To do so we applied a sliding window average of 100 frames (~3.3 sec). The

816   smoothed lick pattern for the ith trial was calculated as follows:

$$SL_{i,a} = \frac{1}{100} \sum_{a}^{a+100} Licks_{i,a}$$

817   Where a is the ath licking frame in the original binary vector and SL_i is the overall smoothed

818   licking pattern. Lick similarity between the ith and jth trials was determined using the formula:

$$Lick\ Similarity(Licks_i, Licks_j)\ \frac{1}{\left\| SL_i - SL_j \right\|^2}$$

Construction pseudo-trials in freely consuming mice

To construct pseudo-trials of consumption in freely consuming mice, we identified time points in

which mice initiated a licking bout. To do so we identified all the time points within the free

consumption epoch of the experiment in which the interval between two consecutive licks was

above 1 second, unless this did not yield enough pseudo-trials, in which case we used 0.5 sec as

the threshold (2/5 datasets). Every pair of licks that passed these criteria could be used to define

the onset of a pseudo-trial. We then took all the time points that occurred two seconds before and

four seconds after the second lick in that interval. We used these time points to identify the

neuronal cluster dynamics and licking pattern that took place in the experiment and constructed a

pseudo-trial matrix. We then removed pseudo-trials that overlapped with each other to avoid

inflation of the matrix. To control for behavioral differences between operant task licking and

free consumption licking, we repeated the analyses in (Fig. 4j-i) while selecting only pseudo-

trials of free consumption in which the average licking rate in the first two seconds of the lick

bout were within the mean ± standard deviation of the average licking rate in the first two

seconds of operant task licking (Supp. Fig. 8c-e).

Quantification and statistical analysis

All statistical details, including the specific statistical tests, are outlined in the corresponding

figure legends. For unrelated samples from two different groups, we conducted a Wilcoxon rank

sum test. In the case of matched-pairs related samples, we performed a Wilcoxon signed rank

test if the number of samples exceeded five; otherwise, we used a matched pairs t-test. Dunn test

was employed for multiple samples, encompassing all possible combinations of comparisons.
When comparing distributions, such as the distribution of pairwise trial similarity, we employed
a two-sample Kolmogorov-Smirnov test. A one-sided Pearson correlation moment test was
carried out to determine the significance levels of sample correlations. To assess the significance
levels of pairwise decoding, p-values were computed by extracting the percentile of the original
decoding accuracy within the distribution of accuracies obtained by shuffling labels. For
evaluating significance levels between pairs of manifolds, permutation tests were performed. The
significance threshold was set at alpha = 0.05, and we corrected for multiple comparisons using
Bonferroni-Holm correction.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the authors used Grammarly and ChatGPT in order to detect
grammatical errors. After using these tools, the authors reviewed and edited the content as
needed and take full responsibility for the content of the publication.

<u>Video S1 – Stereotyped activity dynamics during rewarded trials on a single plane (mouse 1)</u>

Right: Single plane of an example clustered structure. *Left*: Cluster dynamics heatmap in
rewarded trials. Highlighted row on the cluster dynamics heatmap indicates the current rewarded
trial, red dot on the clustered structure indicates current timepoint within the highlighted trial.
Each timepoint is a 0.5 sec time-bin.

<u>Video S2 – Stereotyped activity dynamics during rewarded trials on all planes (mouse 2)</u>

Right: All planes of an example structure. *Middle*: All planes of the same clustered structure.
Left: Cluster dynamics heatmap in rewarded trials. Highlighted row on the cluster dynamics
heatmap indicates the current rewarded trial, red dots on the structure planes and clustered
structure planes indicate current timepoint within the highlighted trial. Each timepoint is a 0.5
sec time-bin.

References

1. Naqvi, N.H., Gaznick, N., Tranel, D., and Bechara, A. (2014). The insula: A critical neural substrate
for craving and drug seeking under conflict and risk. Ann N Y Acad Sci *1316*, 53–70.
10.1111/NYAS.12415.

2. Uddin, L.Q. (2015). Salience processing and insular cortical function and dysfunction. Preprint at
Nature Publishing Group, 10.1038/nrn3857 10.1038/nrn3857.

3. Craig, A.D. (2003). Interoception: the sense of the physiological condition of the body. Curr Opin
Neurobiol *13*, 500–505. 10.1016/S0959-4388(03)00090-4.

4. Gogolla, N. (2017). The insular cortex. Current Biology, 573–591.

5. Owens, A.P., Allen, M., Ondobaka, S., and Friston, K.J. (2018). Interoceptive inference: From
computational neuroscience to clinic. Neurosci Biobehav Rev *90*, 174–183.
10.1016/J.NEUBIOREV.2018.04.017.

6. Quadt, L., Critchley, H.D., and Garfinkel, S.N. (2018). The neurobiology of interoception in health
and disease. Ann N Y Acad Sci *1428*, 112–128. 10.1111/NYAS.13915.

7. Barrett, L.F., and Simmons, W.K. (2015). Interoceptive predictions in the brain. Preprint at Nature
Publishing Group, 10.1038/nrn3950 10.1038/nrn3950.

8. Livneh, Y., and Andermann, M.L. (2021). Cellular activity in insular cortex across seconds to
hours: Sensations and predictions of bodily states. Preprint at Cell Press,
10.1016/j.neuron.2021.08.036 10.1016/j.neuron.2021.08.036.

9. Vyas, S., Golub, M.D., Sussillo, D., and Shenoy, K. V (2020). Computation Through Neural
Population Dynamics. Annu Rev Neurosci, 249–275. 10.1146/annurev-neuro-092619.

10. Jazayeri, M., and Afraz, A. (2017). Navigating the Neural Space in Search of the Neural Code.
Preprint at Cell Press, 10.1016/j.neuron.2017.02.019 10.1016/j.neuron.2017.02.019.

11. Saxena, S., and Cunningham, J.P. (2019). Towards the neural population doctrine. Preprint at
Elsevier Ltd, 10.1016/j.conb.2019.02.002 10.1016/j.conb.2019.02.002.

12. Schneider, S., Lee, J.H., and Mathis, M.W. (2023). Learnable latent embeddings for joint
behavioural and neural analysis. Nature *617*, 360–368. 10.1038/s41586-023-06031-6.

13. Mitchell-Heggs, R., Prado, S., Gava, G.P., Go, M.A., and Schultz, S.R. (2023). Neural manifold
analysis of brain circuit dynamics in health and disease. Preprint at Springer, 10.1007/s10827-
022-00839-3 10.1007/s10827-022-00839-3.

14. Langdon, C., Genkin, M., and Engel, T.A. (2023). A unifying perspective on neural manifolds and
circuits for cognition. Nat Rev Neurosci *24*, 363–377. 10.1038/s41583-023-00693-x.

15. Gardner, R.J., Hermansen, E., Pachitariu, M., Burak, Y., Baas, N.A., Dunn, B.A., Moser, M.B., and
Moser, E.I. (2022). Toroidal topology of population activity in grid cells. Nature *602*, 123–128.
10.1038/s41586-021-04268-7.

16. Chaudhuri, R., Gerçek, B., Pandey, B., Peyrache, A., and Fiete, I. (2019). The intrinsic attractor
manifold and population dynamics of a canonical cognitive circuit across waking and sleep. Nat
Neurosci *22*, 1512–1520. 10.1038/s41593-019-0460-x.

17. Ajabi, Z., Keinath, A.T., Wei, X.-X., and Brandon, M.P. (2023). Population dynamics of head-
direction neurons during drift and reorientation. Nature. 10.1038/s41586-023-05813-2.

18. Gallego, J.A., Perich, M.G., Chowdhury, R.H., Solla, S.A., and Miller, L.E. (2020). Long-term
stability of cortical population dynamics underlying consistent behavior. Nat Neurosci *23*, 260–
270. 10.1038/s41593-019-0555-4.

19. Churchland, M.M., Cunningham, J.P., Kaufman, M.T., Foster, J.D., Nuyujukian, P., Ryu, S.I.,
Shenoy, K. V., and Shenoy, K. V. (2012). Neural population dynamics during reaching. Nature *487*,
51–56. 10.1038/nature11129.

20. Nicolas, C., Ju, A., Wu, Y., Eldirdiri, H., Delcasso, S., Couderc, Y., Fornari, C., Mitra, A., Supiot, L.,
Vérité, A., et al. (2023). Linking emotional valence and anxiety in a mouse insula-amygdala circuit.
Nat Commun *14*, 5073. 10.1038/s41467-023-40517-1.

21. Klein, A.S., Dolensek, N., Weiand, C., and Gogolla, N. (2021). Fear balance is maintained by bodily
feedback to the insular cortex in mice. Science (1979) *374*, 1010–1015.

22. Gehrlach, D.A., Dolensek, N., Klein, A.S., Roy Chowdhury, R., Matthys, A., Junghänel, M.,
Gaitanos, T.N., Podgornik, A., Black, T.D., Reddy Vaka, N., et al. (2019). Aversive state processing
in the posterior insular cortex. Nat Neurosci *22*, 1424–1437. 10.1038/s41593-019-0469-1.

23. Juen, Z., Villavicencio, M., and Zuker, C.S. (2024). A neural substrate for short-term taste
memories. Neuron *112*, 277-287.e4. 10.1016/J.NEURON.2023.10.009.

24. Gardner, M.P.H., Fontanini, A., Gardner, M.P.H., and Fontanini, A. (2014). Encoding and Tracking
of Outcome-Specific Expectancy in the Gustatory Cortex of Alert Rats. Journal of Neuroscience
34, 13000–13017. 10.1523/JNEUROSCI.1820-14.2014.

25. Vincis, R., and Fontanini, A. (2016). Associative learning changes cross-modal representations in
the gustatory cortex. Elife *5*. 10.7554/ELIFE.16420.

26. Katz, D.B., Simon, S.A., and Nicolelis, M.A.L. (2001). Dynamic and Multimodal Responses of
Gustatory Cortical Neurons in Awake Rats.

27. Fonseca, E., De Lafuente, V., Simon, S.A., and Gutierrez, R. (2018). Sucrose intensity coding and
decision-making in rat gustatory cortices. Elife *7*. 10.7554/eLife.41152.001.

28. Fontanini, A., and Katz, D.B. (2006). State-dependent modulation of time-varying gustatory
responses. J Neurophysiol *96*, 3183–3193.
10.1152/JN.00804.2006/ASSET/IMAGES/LARGE/Z9K0120677960006.JPEG.

29. Kusumoto-Yoshida, I., Liu, H., Chen, B.T., Fontanini, A., and Bonci, A. (2015). Central role for the
insular cortex in mediating conditioned responses to anticipatory cues. Proc Natl Acad Sci U S A
112, 1190–1195. 10.1073/PNAS.1416573112/SUPPL_FILE/PNAS.201416573SI.PDF.

30.   Maier, J.X., and Katz, D.B. (2013). Neural dynamics in response to binary taste mixtures. J
Neurophysiol *109*, 2108–2117.
10.1152/JN.00917.2012/ASSET/IMAGES/LARGE/Z9K0081318880009.JPEG.

31.   Stapleton, J.R., Lavine, M.L., Wolpert, R.L., Nicolelis, M.A.L., and Simon, S.A. (2006). Rapid Taste
Responses in the Gustatory Cortex during Licking. Journal of Neuroscience *26*, 4126–4138.
10.1523/JNEUROSCI.0092-06.2006.

32.   Mukherjee, N., Wachutka, J., and Katz, D.B. (2019). Impact of precisely-timed inhibition of
gustatory cortex on taste behavior depends on single-trial ensemble dynamics. Elife *8*.
10.7554/ELIFE.45968.

33.   Jones, L., Fontanini, A., Sadacca, B., Miller, P., and Katz, D. (2007). Natural stimuli evoke dynamic
sequences of statesin sensory cortical ensembles. Proc Natl Acad Sci U S A *104*, 18772–18777.

34.   Lang, L., Camera, G. La, and Fontanini, A. (2023). Temporal progression along discrete coding
states during decision-making in the mouse gustatory cortex. PLoS Comput Biol *19*.
10.1371/journal.pcbi.1010865.

35.   Mazzucato, L., La Camera, G., and Fontanini, A. (2019). Expectation-induced modulation of
metastable activity underlies faster coding of sensory stimuli. Nat Neurosci *22*, 787–796.
10.1038/s41593-019-0364-9.

36.   Samuelsen, C.L., Gardner, M.P.H., and Fontanini, A. (2012). Effects of Cue-Triggered Expectation
on Cortical Processing of Taste. Neuron *74*, 410–422. 10.1016/J.NEURON.2012.02.031.

37.   Moran, A., and Katz, D.B. (2014). Sensory Cortical Population Dynamics Uniquely Track Behavior
across Learning and Extinction. Journal of Neuroscience *34*, 1248–1257.
10.1523/JNEUROSCI.3331-13.2014.

38.   Skaggs, W.E., Knierim, J.J., Kudrimoti, H.S., and Mcnaughton, B.L. (1994). A Model of the Neural
Basis of the Rat's Sense of Direction. Advances in Neural Information Processing System *7*.

39.   Guanella, A., Kiper, D., and Verschure, P. (2011). A MODEL OF GRID CELLS BASED ON A TWISTED
TORUS TOPOLOGY. https://doi.org/10.1142/S0129065707001093 *17*, 231–240.
10.1142/S0129065707001093.

40.   Burak, Y., and Fiete, I.R. (2009). Accurate path integration in continuous attractor network
models of grid cells. PLoS Comput Biol *5*. 10.1371/journal.pcbi.1000291.

41.   Rubin, A., Sheintuch, L., Brande-Eilat, N., Pinchasof, O., Rechavi, Y., Geva, N., and Ziv, Y. (2019).
Revealing neural correlates of behavior without behavioral measurements. Nat Commun *10*.
10.1038/s41467-019-12724-2.

42.   Cechetto, D.F., and Saper, C.B. (1987). Evidence for a Viscerotopic Sensory Representation in the
Cortex and Thalamus in the Rat.

43.   Jones, L.M., Fontanini, A., and Katz, D.B. (2006). Gustatory processing: a dynamic systems
approach. Curr Opin Neurobiol *16*, 420–428. 10.1016/J.CONB.2006.06.011.

44.   Kadohisa, M., Rolls, E.T., and Verhagen, J. V. (2005). Neuronal Representations of Stimuli in the
976   Mouth: The Primate Insular Taste Cortex, Orbitofrontal Cortex and Amygdala. Chem Senses *30*,
977   401–419. 10.1093/CHEMSE/BJI036.

978   45.   Small, D.M. (2012). Flavor is in the brain. Physiol Behav *107*, 540–552.
979   10.1016/J.PHYSBEH.2012.04.011.

980   46.   De Araujo, I.E., and Simon, S.A. (2009). The gustatory cortex and multisensory integration.
981   International Journal of Obesity 2009 33:2 *33*, S34–S43. 10.1038/ijo.2009.70.

982   47.   de Araujo, I.E., Gutierrez, R., Oliveira-Maia, A.J., Pereira, A., Nicolelis, M.A.L., and Simon, S.A.
983   (2006). Neural Ensemble Coding of Satiety States. Neuron *51*, 483–494.
984   10.1016/J.NEURON.2006.07.009.

985   48.   Rodgers, K.M., Benison, A.M., Klein, A., and Barth, D.S. (2008). Auditory, Somatosensory, and
986   Multisensory Insular Cortex in the Rat. Cerebral Cortex *18*, 2941–2951.
987   10.1093/CERCOR/BHN054.

988   49.   Yamamoto, T., Matsuo, R., Kiyomitsu, Y., and Kitamura, R. (1988). Sensory inputs from the oral
989   region to the cerebral cortex in behaving rats: an analysis of unit responses in cortical
990   somatosensory and taste areas during ingestive behavior.
991   https://doi.org/10.1152/jn.1988.60.4.1303 *60*, 1303–1321. 10.1152/JN.1988.60.4.1303.

992   50.   Livneh, Y., Ramesh, R.N., Burgess, C.R., Levandowski, K.M., Madara, J.C., Fenselau, H., Goldey,
993   G.J., Diaz, V.E., Jikomes, N., Resch, J.M., et al. (2017). Homeostatic circuits selectively gate food
994   cue responses in insular cortex. Nature *546*, 611–616. 10.1038/nature22375.

995   51.   Dolensek, N., Gehrlach, D.A., Klein, A.S., and Gogolla, N. (2020). Facial expressions of emotion
996   states and their neuronal correlates in mice. Science (1979) *368*.
997   10.1126/SCIENCE.AAZ9468/SUPPL_FILE/AAZ9468S3.MP4.

998   52.   Deng, H., Xiao, X., Yang, T., Ritola, K., Hantman, A., Li, Y., Huang, Z.J., and Li, B. (2021). A
999   genetically defined insula-brainstem circuit selectively controls motivational vigor. Cell *184*, 6344-
6360.e18. 10.1016/J.CELL.2021.11.019.

53.   Wang, L., Gillis-Smith, S., Peng, Y., Zhang, J., Chen, X., Salzman, C., Ryba, N., and Zuker, C. (2018).
The coding of valence and identity in the mammalian taste system. Nature *558*, 127–131.

54.   Egan, G., Silk, T., Zamarripa, F., Williams, J., Federico, P., Cunnington, R., Carabott, L., Blair-West,
J., Shade, R., McKinley, M., et al. (2003). Neural correlates of the emergence of consciousness of
thirst. Proc Natl Acad Sci U S A *100*, 15241–15246. 10.1073/PNAS.2136650100.

55.   Livneh, Y., Sugden, A.U., Madara, J.C., Essner, R.A., Flores, V.I., Sugden, L.A., Resch, J.M., Lowell,
B.B., and Andermann, M.L. (2020). Estimation of Current and Future Physiological States in
Insular Cortex. Neuron *105*, 1094-1111.e10. 10.1016/j.neuron.2019.12.027.

56.   Meier, L., Federspiel, A., Jann, K., Wiest, R., Neuroscience, W.S.-, and 2018, undefined Thirst-
dependent activity of the insular cortex reflects its emotion-related subdivision: a cerebral blood
flow study. Elsevier.

57.   Tataranni, P.A., Gautier, J.F., Chen, K., Uecker, A., Bandy, D., Salbe, A.D., Pratley, R.E., Lawson, M.,
Reiman, E.M., and Ravussin, E. (1999). Neuroanatomical correlates of hunger and satiation in

humans using positron emission tomography. Proc Natl Acad Sci U S A *96*, 4569–4574.
10.1073/PNAS.96.8.4569.

58. Jazayeri, M., and Ostojic, S. (2021). Interpreting neural computations by examining intrinsic and
embedding dimensionality of neural activity. Preprint at Elsevier Ltd, 10.1016/j.conb.2021.08.002
10.1016/j.conb.2021.08.002.

59. Nieh, E.H., Schottdorf, M., Freeman, N.W., Low, R.J., Lewallen, S., Koay, S.A., Pinto, L., Gauthier,
J.L., Brody, C.D., and Tank, D.W. (2021). Geometry of abstract learned knowledge in the
hippocampus. Nature *595*, 80–84. 10.1038/s41586-021-03652-7.

60. Humphries, M.D. (2021). Strong and weak principles of neural dimension reduction. Neuron
Behav Data Anal Theory *5*, 1–28. 10.51628/001c.24619.

61. Gehrlach, D.A., Weiand, C., Gaitanos, T.N., Cho, E., Klein, A.S., Hennrich, A.A., Conzelmann, K.K.,
and Gogolla, N. (2020). A whole-brain connectivity map of mouse insular cortex. Elife *9*, 1–78.
10.7554/ELIFE.55585.

62. Saper, C.B. (2002). The central autonomic nervous system: Conscious visceral perception and
autonomic pattern generation. Preprint, 10.1146/annurev.neuro.25.032502.111311
10.1146/annurev.neuro.25.032502.111311.

63. Deitch, D., Rubin, A., and Ziv, Y. (2021). Representational drift in the mouse visual cortex. Current
Biology *31*, 4327-4339.e6. 10.1016/j.cub.2021.07.062.

64. Belkin, M., and Niyogi, P. (2003). Laplacian Eigenmaps for Dimensionality Reduction and Data
Representation. Neural Comput *15*, 1373–1396.

65. Wang, J. (2012). Laplacian Eigenmaps. In Geometric Structure of High-Dimensional Data and
Dimensionality Reduction (Springer), pp. 51–77.

66. Rybakken, E., Baas, N., and Dunn, B. (2019). Decoding of neural data using cohomological feature
extraction. Preprint at MIT Press Journals, 10.1162/neco_a_01150 10.1162/neco_a_01150.

67. R. Wadhwa, R., F.K. Williamson, D., Dhawan, A., and G. Scott, J. (2018). TDAstats: R pipeline for
computing persistent homology in topological data analysis. J Open Source Softw *3*, 860.
10.21105/joss.00860.

68. Rouse, T.C., Ni ID, A.M., Huang, C., and Cohen, M.R. (2023). Topological insights into the neural
basis of flexible behavior. Proceedings of the National Academy of Sciences *120*. 10.1073/pnas.

69. Robinson, A., and Turner, K. (2017). Hypothesis testing for topological data analysis. J Appl
Comput Topol *1*, 241–261. 10.1007/s41468-017-0008-7.

70. Musall, S., Kaufman, M., Juavinett, A., Gluf, S., and Churchland, A. (2019). Single-trial neural
dynamics are dominated by richly varied movements. Nat Neurosci *22*, 1677–1686.

71. Salkoff, D., Zagha, E., … E.M.-C., and 2020, undefined Movement and performance explain
widespread cortical activity in a visual detection task. academic.oup.comDB Salkoff, E Zagha, E
McCarthy, DA McCormickCerebral Cortex, 2020•academic.oup.com.

72.   Stringer, C., Pachitariu, M., Steinmetz, N., Reddy, C.B., Carandini, M., and Harris, K.D. (2019).
1051   Spontaneous behaviors drive multidimensional, brainwide activity. Science (1979) *364*.
1052   10.1126/SCIENCE.AAV7893.

1053   73.   Burgess, C.R., Ramesh, R.N., Sugden, A.U., Levandowski, K.M., Minnig, M.A., Fenselau, H., Lowell,
1054   B.B., and Andermann, M.L. (2016). Hunger-Dependent Enhancement of Food Cue Responses in
1055   Mouse Postrhinal Cortex and Lateral Amygdala. Neuron *91*, 1154–1169.
1056   10.1016/j.neuron.2016.07.032.

1057   74.   McGinley, M.J., Vinck, M., Reimer, J., Batista-Brito, R., Zagha, E., Cadwell, C.R., Tolias, A.S., Cardin,
1058   J.A., and McCormick, D.A. (2015). Waking State: Rapid Variations Modulate Neural and
1059   Behavioral Responses. Neuron *87*, 1143–1161. 10.1016/J.NEURON.2015.09.012.

1060   75.   Tenenbaum, J.B., de Silva, V., and Langford, J.C. (2000). A Global Geometric Framework for
1061   Nonlinear Dimensionality Reduction. Science (1979) *290*, 2319–2322.

1062   76.   Augustine, V., Gokce, S.K., and Oka, Y. (2018). Peripheral and Central Nutrient Sensing Underlying
1063   Appetite Regulation. Trends Neurosci *41*, 526–539. 10.1016/J.TINS.2018.05.003.

1064   77.   Andermann, M.L., and Lowell, B.B. (2017). Toward a Wiring Diagram Understanding of Appetite
1065   Control. Neuron *95*, 757–778. 10.1016/J.NEURON.2017.06.014.

1066   78.   Zimmerman, C.A., and Knight, Z.A. (2020). Layers of signals that regulate appetite. Curr Opin
1067   Neurobiol *64*, 79–88. 10.1016/J.CONB.2020.03.007.

1068   79.   Sternson, S.M., and Eiselt, A.K. (2017). Three Pillars for the Neural Control of Appetite.
1069   https://doi.org/10.1146/annurev-physiol-021115-104948 *79*, 401–423. 10.1146/ANNUREV-
1070   PHYSIOL-021115-104948.

1071   80.   Zimmerman, C.A., Lin, Y.C., Leib, D.E., Guo, L., Huey, E.L., Daly, G.E., Chen, Y., and Knight, Z.A.
1072   (2016). Thirst neurons anticipate the homeostatic consequences of eating and drinking. Nature
1073   *537*, 680–684. 10.1038/nature18950.

1074   81.   Yiannakas, A., Kolatt Chandran, S., Kayyal, H., Gould, N., Khamaisy, M., and Rosenblum, K. (2021).
1075   Parvalbumin interneuron inhibition onto anterior insula neurons projecting to the basolateral
1076   amygdala drives aversive taste memory retrieval. Curr Biol *31*, 2770-2784.e6.
1077   10.1016/J.CUB.2021.04.010.

1078   82.   Stern, S.A., Azevedo, E.P., Pomeranz, L.E., Doerig, K.R., Ivan, V.J., and Friedman, J.M. (2021). Top-
1079   down control of conditioned overconsumption is mediated by insular cortex Nos1 neurons. Cell
1080   Metab *33*, 1418-1432.e6. 10.1016/j.cmet.2021.03.001.

1081   83.   Asaad, W.F., and Eskandar, E.N. (2008). A flexible software tool for temporally-precise behavioral
1082   control in Matlab. J Neurosci Methods *174*, 245–258. 10.1016/j.jneumeth.2008.07.014.

1083   84.   Bonin, V., Histed, M.H., Yurgenson, S., and Clay Reid, R. (2011). Local diversity and fine-scale
1084   organization of receptive fields in mouse visual cortex. Journal of Neuroscience *31*, 18506–18521.
1085   10.1523/JNEUROSCI.2974-11.2011.

1086   85.   Mukamel, E.A., Nimmerjahn, A., and Schnitzer, M.J. (2009). Automated Analysis of Cellular
1087   Signals from Large-Scale Calcium Imaging Data. Neuron *63*, 747–760.
1088   10.1016/j.neuron.2009.08.009.

86. Facco, E., D'Errico, M., Rodriguez, A., and Laio, A. (2017). Estimating the intrinsic dimension of
datasets by a minimal neighborhood information. Sci Rep 7. 10.1038/s41598-017-11873-y.

87. Kraemer, G., Reichstein, M., and Mahecha, M.D. (2018). dimRed and coRanking-Unifying
Dimensionality Reduction in R. R J 10, 342–358.

88. Ramdas, A., Garcia, N., and Cuturi, M. (2015). On Wasserstein Two Sample Testing and Related
Families of Nonparametric Tests. ArXiv.

89. Murtagh, F., and Legendre, P. (2014). Ward's Hierarchical Agglomerative Clustering Method:
Which Algorithms Implement Ward's Criterion? J Classif 31, 274–295. 10.1007/s00357-014-9161-
z.

90. Winkelmeier, L., Filosa, C., Hartig, R., Scheller, M., Sack, M., Reinwald, J.R., Becker, R., Wolf, D.,
Gerchen, M.F., Sartorius, A., et al. (2022). Striatal hub of dynamic and stabilized prediction coding
in forebrain networks for olfactory reinforcement learning. Nat Commun 13. 10.1038/s41467-
022-30978-1.

1102

A

Cue   Action   Outcome

Reward cue → Lick → Reward
        → No lick → Nothing

Aversive cue → Lick → Salt
        → No lick → Nothing

Neutral cue → Lick → Nothing
        → No lick → Nothing

Visual cue (2s) | Response window (2s) | inter-trial Interval (6s)

B Two-photon imaging of insular cortex

10 μm

C Neuronal activity state space

Neurons

n_1
n_2
n_3
n_4
n_5
n_k

dF/F

t_1 .... t_n

Time

Bin, Normalize →

n_2
t_1
t_n
n_3
n_4
n_5
n_k
n_1

D Insular Cortex (Thirsty mice)

Mouse 1 — Dim 2 / Dim 1 — Dim 4 / Dim 3 — Dim 5 / Dim 1

Mouse 2 — Dim 2 / Dim 1 — Dim 2 / Dim 3 — Dim 5 / Dim 4

Mouse 3 — Dim 2 / Dim 1 — Dim 3 / Dim 2 — Dim 5 / Dim 4

E

Estimated dimensionality

Data   Shuffle

****

F

☐ Data ☐ Shuffle

Structure dissimilarity (log2 divergence)

***   ***

NS

Across days   Across mice

G Insular Cortex (Hungry mice)

Mouse 1 — Dim 2 / Dim 1 — Dim 2 / Dim 3 — Dim 5 / Dim 4

Mouse 2 — Dim 2 / Dim 1 — Dim 5 / Dim 4 — Dim 2 / Dim 3

Mouse 3 — Dim 2 / Dim 1 — Dim 4 / Dim 3 — Dim 5 / Dim 1

H

Estimated dimensionality

NS

Hungry   Thirsty

I

Structure dissimilarity (log2 divergence)

NS

Hungry   Thirsty

J

Estimated dimensionality

*

InsCtx (Hungry)   Visual (Hungry)

K

Structure dissimilarity (log2 divergence)

**

InsCtx (Hungry)   Visual (Hungry)

A Mouse 1

Neutral cue trial (Response) | Reward cue trial (Response) | Clustered structure

B Mutual Information with Clustered structure (log2)

Pupil | Cuml. reward | Reward (R) | Reward (NR) | Aversive (R) | Aversive (NR) | Neutral (R) | Neutral (NR)

*** * *** NS

C Mutual Information with Clustered structure (log2)

Pupil | Cuml. reward | Reward (R)

*** *** ****

Data Shuffle

D Mouse 1 | Mouse 2 | **E** All mice

Reward | Aversive | Neutral

Time (s) 0   10

Fraction

Trial similarity (cosine) −1   0   1

****

Trial outcome
Response   No Response

Central point cloud
−1

Neuronal   Cluster
1 2 3 4 5 6 7

Reward (R)   Aversive (R)   Neutral (R)
Reward (NR)   Aversive (NR)   Neutral(NR)

F Trial dynamics | Transition matrix | Sorted transition matrix | Aligned trials mouse 1 → mouse 2 | Decoded trials mouse 2

Mouse 1

Mouse 2

Transition probabilities

Hierarchial clustering

Align by sorting

Decode

Time

Neuronal cluster

0   .05
Probability

Time   Time

G Decoding accuracy (%)

Reward cue trial | Aversive cue trial | Neutral cue trial

Response | No Response

Across days | Across mice

Data Shuffle

*** *** NS NS NS NS NS NS NS NS NS NS

A

InsCtx (Hungry)

Mouse 1   Mouse 2

Visual (Hungry)

Mouse 3   Mouse 4

0   Time (s)   10

Trial outcome (Reward) Central point cloud Neuronal Cluster

Response No Response   -1   1 2 3 4 5 6 7

B Responded reward trials

Decoding accuracy (%)

Data Shuffle

***

**

NS

Thirsty   Visual (hungry)
↓   ↓
Hungry   InsCtx (hungry)

C

Natural / artifical hunger

Experiment progression

Natural

Artificial

0   Time (s)   10

D

Natural vs Natural   Natural vs Artificial

Fraction

Trial similarity (cosine)

Trial similarity

NS

E

Decoding accuracy (%)

Data Shuffle

NS   **

NS

Hungry → Natural   Hungry → Artificial   Hungry → Natural   Hungry → Artificial

F

Natural / artifical thirst

Experiment progression

Natural

Artificial

0   Time (s)   10

G

Natural vs Natural   Natural vs Artificial

Fraction

Trial similarity (cosine)

Trial similarity

NS

H

Decoding accuracy (%)

Data Shuffle

NS   ***

NS

Thirsty → Natural   Thirsty → Artificial   Thirsty → Natural   Thirsty → Artificial

I

Natural / hyperosmotic thirst

Experiment progression

Natural

Hyperos.

0   Time (s)   10

J

Natural vs Natural   Natural vs Hyperos.

Fraction

Trial similarity (cosine)

Trial similarity

NS

K

Decoding accuracy (%)

Data Shuffle

NS   ***

NS

Thirsty → Natural   Thirsty → Hyperos.   Thirsty → Natural   Thirsty → Hyperos.