Stereotyped goal-directed activity patterns in the insular cortex

Itay Talpir[1] and Yoav Livneh[1*]

[1]Department of Brain Sciences, Weizmann Institute of Science, Rehovot 76100, Israel

*Correspondence: yoav.livneh@weizmann.ac.il

- InsCtx activity manifold structure is stereotyped across different mice and motivations.
- InsCtx activity dynamics within the manifold are stereotyped across different mice and motivations.
- Stereotyped activity dynamics reflect goal-directed reward anticipation.
- Stereotyped activity dynamics do not reflect licking, taste, or valence.

Abstract

The insular cortex is involved in diverse processes including bodily homeostasis,
emotions, and cognition. Yet little is known about how it processes information at the level
of neuronal populations, precluding an understanding of the computations it performs.
We leveraged recent advances in machine learning to study insular cortex population
activity patterns (i.e., neuronal manifold) in mice performing goal-directed behaviors. We
find that the insular cortex activity manifold is remarkably consistent across different
animals and under different motivational states. Activity dynamics within the neuronal
manifold are highly stereotyped during rewarded trials, enabling robust prediction of trial
outcomes across different mice, and across various natural and artificial motivational
states. Comparing goal-directed behavior with self-paced free consumption, we find that
the stereotyped activity patterns reflect goal-directed reward anticipation, and not licking,
taste, or valence. These findings reveal a core computation in insular cortex that could
explain its involvement in pathologies involving aberrant motivations.

Introduction

The insular cortex (or insula) is involved in a variety of processes, ranging from basic functions including bodily homeostasis, emotions, motivation, decision making, social interactions, and sensory processing, to maladaptive conditions including drug addiction, chronic pain, anxiety, obesity, and eating disorders[1–8 5–8]. Yet we lack a mechanistic understanding of the core computations in InsCtx that underlie its contribution to so many diverse processes.

A central path to understanding the core computations within and across brain regions lies in measuring and making sense of neural activity of large populations of neurons in different behavioral contexts[1–3]. Indeed, there has been a recent explosion of artificial intelligence and machine learning approaches to deciphering neural activity at the level of large neuronal populations[4–6]. These approaches have begun to provide insights into the core computations of neural circuits, such as motor cortex control of limb movement, thalamic representation of head direction, and entorhinal cortex representation of spatial position[7–11]. And yet, most animal model studies of InsCtx have used either bulk signal recordings like fiber photometry, or electrical recordings from relatively small populations, not yet leveraging these new approaches to better understand InsCtx computations. These studies have indeed substantially advanced our understanding of the bulk activity of InsCtx, of the activity of individual InsCtx neurons, and also of activity patterns in small populations. Nevertheless, a comprehensive unbiased investigation of activity patterns of large InsCtx populations, taking advantage of recent developments in machine learning tools, has yet to be performed.

Previous work has suggested that InsCtx activity on short-timescales (~seconds) represents multi-modal gustatory and interoceptive sensory stimuli[12–17], as well as multi-modal salient external cues[15,18–25]. Further work has suggested that such activity may also convey emotional, motivational and valence related information[26–31]. Longer timescale changes in InsCtx activity (minutes to hours) have been suggested to represent slow changes in the physiological state of the body[20,26,32–35], as well as specific positive and negative emotional states[26–29]. However, seeing as InsCtx integrates many different inputs and is involved in many diverse processes, understanding its function at the neuronal population level requires comprehensive analyses of the repertoire of its activity

patterns (i.e., the neuronal manifold) and their dynamics. Doing so would help answer

fundamental questions that remain unanswered. For example, how constrained are

InsCtx population activity patterns by intrinsic factors vs. extrinsic factors such as the

current behavioral context? How are different external and internal behavioral variables

represented concomitantly in InsCtx population activity patterns? Are there core

computations that InsCtx performs across different contexts? If so, will they have distinct

signatures in population activity space?

Here, we leverage recent advances in machine learning and topological data analysis to

perform an unbiased investigation of InsCtx population activity during goal-directed

behaviors. We use these approaches to analyze new as well as previously published data

[25,33]. We find that the repertoire of InsCtx population activity patterns (i.e., neuronal

manifold) has a remarkably consistent structure across different animals, physiological

needs, and behavioral contexts. The InsCtx activity manifold can be partially explained

by experimentally measured external and internal behavioral variables. Moreover, by

tracking the dynamics of activity within the manifold, we uncover previously unknown

stereotyped patterns of activity. These stereotyped activity dynamics were highly

conserved and allowed decoding across individuals, across different physiological needs,

different motivational states, and different available rewards. A comparison of population

dynamics during task-structured goal-directed behavior vs. self-paced, free consumption

behavior revealed that the stereotyped activity patterns reflect task-dependent reward

anticipation, as opposed to specific behavioral actions, taste, or valence. Our findings

demonstrate the power of comprehensive population level analyses to reveal core

computations of insular cortex.

Results

Insular cortex population activity manifold structure is stereotyped across different mice and motivational states

To investigate how neuronal activity patterns are structured in the insular cortex (InsCtx), we analyzed two-photon calcium imaging data from new datasets, as well as previously published ones[33]. These data were collected while thirsty (water restricted) mice performed an operant go/no go visual discrimination task in which licking following presentation of three initially arbitrary visual cues (drifting gratings) leads to rewarding (water), aversive (1 M NaCl), or neutral outcomes (Figure 1a-b)[33]. We have previously shown that most InsCtx neurons respond to some aspect of this behavioral task with either increases or decrease in activity (e.g., visual cues, licking, reward consumption)[33]. Therefore, in order to detect any underlying structure within the observed activity patterns (i.e., the "activity manifold"), we aimed to capture the temporal similarities across the entire neuronal population (Figure 1c-d).

Previous studies have shown that activity manifolds are of low dimensionality during relatively simple behavioral tasks and are thus determined by relatively few latent variables, as compared to the number of observed neurons[2,5,7,8,36-39]. However, because InsCtx is a central hub that integrates multimodal information from somatosensory, auditory, visceral, gustatory, and other limbic regions[40,41], we wondered to what extent InsCtx activity would be similarly low dimensional. Our initial goal was to determine the intrinsic dimensionality of the InsCtx population data, i.e., the minimal number of continuous variables needed to parametrize the activity manifold[36,42] (Supp. Fig. 1a-b[42]). We found that across all of our datasets, the activity exhibited a consistent intrinsic dimensionality of 6 ± 1 dimensions (mean ± SD), independent of the specific parameters used to assess dimensionality and of the number of observed neurons (Figure 1e; Supp. Fig. 1c-e). This consistently low intrinsic dimensionality, together with the diverse array of multi-modal inputs that InsCtx receives, suggest a large degree of convergence and integration of information in InsCtx during goal-directed behavior (but not in other cortical areas, see below).

Building on this finding, we used non-linear dimensionality reduction techniques to unveil the low dimensional structure of InsCtx activity patterns[4,5,7-9,38,39,43]. We selected the

target dimensionality based on the mean estimated intrinsic dimensionality across all
datasets. This analysis revealed remarkably consistent structure of neuronal activity
across datasets and across different mice (Figure 1d). To assess this structure similarity
quantitatively, we used topological data analysis, which is emerging as an important
approach for quantitative description of the structure of complex data[4,7,44–46]. We
measured the similarity by assessing the difference (i.e., divergence) between the
distributions of topological features across datasets (see detailed illustration of the
procedure in Supp. Fig. 2a). Notably, this analysis revealed that the structure of low-
dimensional activity for a dataset remained consistent across different mice and days.
Structural similarity was significantly lower for shuffled datasets (Figure 1f). We further
validated these findings using pairwise permutation tests[47] (Supp. Fig. 2b-c).

We next assessed whether InsCtx dimensionality and manifold structure are conserved
across different motivations. To do this, we compared data from thirsty (water-restricted)
mice engaged in a behavioral task to obtain water rewards, to data from hungry (food-
restricted) mice engaged in the same task but to obtain liquid food rewards[25] (Figure 1g)
The intrinsic dimensionality of the activity manifold in hungry mice and thirsty were quite
similar (5.7 ± 1 vs. 6 ± 1; mean ± SD; Figure 1h). Moreover, the structure of the activity
manifold in hungry mice was similar to that observed in thirsty mice (Figure 1i).

We next asked whether the consistent dimensionality and topological similarity we
observed in InsCtx reflect behavioral constraints on brain-wide activity, i.e., whether they
are unique characteristics of InsCtx. We compared the InsCtx activity manifold with the
activity manifold of visual cortical areas of food-deprived mice performing the exact same
behavioral task (primary visual cortex and postrhinal cortex[48]). Of note, previous work has
shown that population activity patterns in visual areas, including primary visual cortex, are
heavily influenced by ongoing task-related and "spontaneous" behaviors[49–51]. We found
that the dimensionality and structure of the activity manifold in visual areas were both
significantly higher and less consistent than those in InsCtx (Figure 1j-k). This was further
confirmed by examining the distribution of P-values resulting from pairwise comparisons
of the structure of activity manifolds between datasets within each group (Supp. Fig. 2d).
These results suggest that the consistent dimensionality and topological similarity in

InsCtx are not a result of brain-wide constraints on behavior, and could be more specific
to a subset of brain regions that include InsCtx.

Collectively, our findings indicate that the neuronal activity manifold within InsCtx is low
dimensional and is structurally similar across mice engaged in goal-directed behaviors.
The consistent manifold structure across animals, suggests common activity patterns,
and therefore perhaps also common encoded information. We next sought to investigate
the dynamics of activity within this stereotyped manifold and how they relate to ongoing
behavior.

154 Insular cortex activity dynamics within the manifold are stereotyped across
155 different mice

Based on previous work[18,25,26,30,33], we expected InsCx activity patterns on both short and
long time-scales to correlate with external and internal behavioral variables (e.g., different
trial types, rewarding or aversive outcomes, changing satiety levels, arousal levels, etc.).
We color-coded the manifold based on different experimental parameters (Figure 2a, left
and middle; Supp. Fig. 3a) and found that some time-points that were associated with a
given variable were spatially clustered within the neuronal activity manifold, while others
were more uniformly distributed. We developed an automated method to cluster the
activity patterns within the manifold (termed here as "Clustered structure"). This method
identified different clusters of activity patterns that diverged from the center of the
manifold. We refer here to these clusters as 'manifold clusters' and refer to the center that
was not part of any distinct cluster as the 'central point cloud'. (Figure 2a, right; Supp. Fig.
3b-c). Note that different cluster indices (represented by different colors in the figure) are
randomly assigned during the clustering procedure, and therefore do not necessarily map
to the same behavioral variable across different mice (see more below).

Using the clustered structure approach, we quantitatively assessed the relationship
between the activity manifold and external/internal behavioral variables by computing the
mutual information between them (Supp. Fig. 3d-e; see Methods). Three variables had
higher mutual information with the activity manifold than others: (1) pupil size in inter-trial
intervals (as a proxy for ongoing arousal levels in between epochs of overt behavior[52]),

(2) cumulative water rewards (as a proxy for slow changes in physiological state, i.e.,

water satiety[33]), and (3) trials in which the mice correctly responded to reward predicting

cues to receive water rewards (referred to hereafter as "rewarded trials"; Figure 2b).

Moreover, the mutual information for these three variables was significantly higher than

what was obtained for shuffled data (Figure 2c). These results are consistent with

previous work showing changes in InsCtx activity that are associated with rewards,

arousal, and physiological state[25–30,33]. We have previously shown that water rewards

shift population activity along a linear projection on a thirsty-quenched axis[33]. However,

manifold clusters associated with the quenched state were different from those

associated with rewarded trials (Supp. Fig. 4a-b), suggesting that there are also distinct

representations of water satiety and reward in InsCtx population activity.

We next sought to understand the dynamics associated with behavioral variables within

the activity manifold. We represented each activity pattern by its manifold cluster, and

then examined the temporal sequence of clusters during different trial types (Figure 2d).

InsCtx neuronal populations displayed somewhat stereotyped patterns of cluster

transitions during rewarded trials, characterized by movement from the central point cloud

to a sequence of transitions between the same 2-3 manifold clusters. To evaluate the

consistency of these dynamics, we calculated the pairwise similarity of cluster sequences

across all trial types. Cluster sequences were highly similar across different rewarded

trials, and substantially more so than all other trial types (Figure 2e). We confirmed these

results using a different dimensionality reduction method (Isomap[53]; Supp. Fig. 5a-b). We

also directly compared the new cluster sequence method to our previous method of

projecting the activity onto empirically defined linear thirsty-quenched axes in population

activity space[33]. As expected, we found that the linear projection naturally captures less

of the variance of activity within the manifold and is much less consistent across trials

(Supp. Fig. 4c-d).

We next examined to what extent trial outcomes could be decoded from the stereotyped

sequence of activity patterns across days and mice. The challenge in decoding across

mice is that our clustering technique does not yield functionally consistent cluster indices

across mice. That is, clusters associated with the same behavioral variable would not

have the same label across mice. We therefore developed a method for translating cluster

labels between data sets in order to properly align the activity dynamics between different
datasets (different days, different mice) for decoding. We first determined the probability
of transitions between different clusters in each individual dataset (Figure 2f, Transition
matrix). We then translated the cluster labels from one "reference dataset" to relabel the
clusters transition probabilities of the other dataset (Figure 2f, Sorted transition matrix).
Finally, we used the translated cluster labels to train the decoder and tested it on the other
"reference dataset" (Figure 2f). Using this approach we could classify rewarded trials
across different days, and even across different mice, with high accuracy. We achieved
an average accuracy of 75% ± 13% and 65% ± 13% (mean ± SD), respectively, which
was 4-5 times higher than chance (~12-15%). The decoding accuracy for other trial types
did not significantly surpass that obtained from randomly shuffling trial labels (Figure 2g).
We further validated these results using the P-values derived from the shuffle distributions
for each pairwise decoding (Supp. Fig. 6).

Importantly, decoding trial types on single trials across mice using this method was
substantially more accurate than using average activity levels across all neurons, or using
our previous approach of projecting activity on linear axes[33] (Supp. Fig. 6). These
analyses suggest that decoding accuracy for rewarded trials results from the stereotyped
patterns of activity and does not trivially reflect global changes in activity. These findings
demonstrate that there are highly stereotyped activity patterns within the InsCtx activity
manifold during rewarded trials, both across days and across mice.

**Insular cortex activity dynamics within the manifold are stereotyped across
different motivational states and rewards**

Given the remarkably stereotyped activity patterns we observed for rewarded trials in
thirsty mice (motivated by water rewards), and the consistent InsCtx manifold structure
during both hunger and thirst, we next tested whether similar dynamics were also evident
in mice motivated by food rewards. We first visualized the temporal sequence of manifold
clusters of rewarded trials in hungry mice in the InsCtx and found that it displayed similar
stereotyped cluster sequences as in thirsty mice (Figure 3a). In contrast, activity
sequences in visual cortical areas were very different and, as expected, primarily showed
activity during the presentation of the visual cue (0-2.5 seconds into the trial), independent

of reward. We then tested whether we could decode trial outcome across thirsty and

hungry mice, receiving water and food rewards, respectively. The trial outcome in hungry

mice could be decoded with high accuracy of $67 \pm 23\%$ (mean $\pm$ SD; vs. ~12-15% chance,

see Methods) when we trained the decoder on InsCtx activity dynamics from thirsty mice.

However, the decoding accuracy was similar to chance ($11 \pm 12\%$) when the decoder

was trained on activity from visual areas (Figure 3b). The consistent activity dynamics in

InsCtx across hunger and thirst suggests that these activity patterns do not reflect the

pure sensory attributes of the reward or the underlying physiological need.

We performed further analyses to test the extent to which these activity patterns remain

stereotyped in mice performing similar behavioral tasks across additional different natural

and artificial motivational states. Specifically, we examined InsCtx activity from mice in

which artificial hunger or thirst states were induced by chemogenetic activation of

hypothalamic AgRP "hunger neurons" or SFO^{GLUT} "thirst neurons", which are the primary

sensors of physiological imbalances and actuators of relevant behavioral and

physiological responses[54–57]. We also analyzed InsCtx activity from mice in which we

induced hyper-osmotic thirst by administration of hypertonic saline[58]. These 3 datasets

gave us the unique opportunity to compare, within each mouse, the activity during the

natural motivation driven by physiological need, with the artificially induced motivation.

We first visualized the sequences of manifold clusters in these mice and compared them

during the natural motivational state to those after induction of the artificial motivational

state. Sequence transitions during rewarded trials appeared remarkably similar in both

the natural and artificially induced motivational states (Figure 3c,f,i). We quantified this

using the pairwise trial similarity before and after the induction of the artificial motivation.

For all motivational states, the pairwise trial similarities remained very similar between the

natural and artificially induced motivational states, with no significant changes observed

(Figure 3d,g,j). We therefore trained the decoder on datasets from hungry and thirsty mice

that were not part of the artificial motivation experiments and then used these to test

decoding in the same mice before and after artificial induction of a motivational state.

Decoding accuracy for rewarded trials in artificial motivational states was similar to the

natural motivational states, and both were significantly higher than chance (Figure

3e,h,k). These findings collectively demonstrate that the stereotyped activity patterns in

InsCtx are not specific to a particular reward (including its sensory properties), nor to the
specific underlying motivational state.

Insular cortex stereotyped activity dynamics reflect goal-directed reward
anticipation, but not licking, taste, or valence

The stereotyped activity patterns we observed in InsCtx across rewarded trials with
different rewards and motivations could reflect goal-directed behavior, or it could simply
reflect licking, valence, or fulfilment of a need. We next tested these different options.

In both hungry and thirsty mice, we observed that the onset of the stereotyped sequence
of activity clusters during rewarded trials appeared to be linked with anticipatory licking
during the trial, as evidenced by sorting trials based on the onset of anticipatory licking
(Figure 4a). We thus quantified the correlation between the onsets of sequential activity
and anticipatory licking in individual trials (Figure 4b). These correlations were notably
high in hungry mice (0.5 ± 0.13; mean ± SD) and in thirsty mice (0.47 ± 0.15), with all
correlations being statistically significant (Figure 4c). Additionally, we observed that the
onset of anticipatory licking consistently preceded the onset of sequential cluster activity,
and occurred independently of when the reward was delivered (Figure 4d-e). Together
with similar activity dynamics across food and water rewards (Figure 3), this further
supports a dissociation from the sensory aspect of the reward. Thus, these stereotyped
activity patterns could potentially reflect licking per se, valence (independent of the
reward's sensory properties), or cue-driven goal-directed behavior.

To distinguish between these possibilities, we examined trials in which mice incorrectly
licked to cues that do not predict reward ("false alarm" trials). We found that cluster
sequences in false alarm trials were more similar to rewarded trials than other trial types
(Figure 4f). To further investigate this, we compared cluster sequence dynamics for false
alarm trials and rewarded trials. We sorted false alarm trials based on the similarity of
licking pattern to rewarded trial licking (Figure 4g). Indeed, false alarm trials with similar
licking patterns to rewarded trials also displayed similar cluster sequence dynamics. To
quantify this, we assessed the correlations between licking similarity and cluster

sequence similarity in false alarms and rewarded trials within individual datasets. We
observed positive correlations in 9/11 datasets (Figure 4h). Because mice do not actually
receive reward in "false alarm" trials, these results further support the conclusion that
InsCtx stereotyped activity patterns do not reflect reward or valence. Nevertheless, they
could still potentially reflect licking behavior, or cue-associated goal-directed reward
expectation.

To further distinguish between these possibilities, we compared cluster sequences when
hungry mice consumed rewards while actively engaged in the operant visual
discrimination task, and when they did so while freely licking in the same physical context
(head-fixed, visual cue monitor is on but does not present cues) to receive rewards in the
absence of predictive cues or any task structure. Notably, in both conditions mice had a
similar physiological need, fulfilled using the same action (licking), to obtain the same
reward. We constructed 'pseudo-trials' during the free reward consumption epochs by
identifying significant breaks between lick bouts and analyzed the corresponding
sequences of manifold clusters. As expected, average population activity increased
during reward consumption in both task-structured trials and self-paced consumption
(Supp. Fig. 7). However, activity dynamics during task engagement and free reward
consumption were distinctly different (Figure 4i). Quantitatively, the pairwise trial similarity
between operant trials and free reward consumption trials was significantly lower than the
pairwise similarity within operant trials. (Figure 4j, Supp. Fig. 7). These results could be
explained by differences in patterns of licking (Supp. Fig. 7). Taken together, these results
suggest that the stereotyped sequential activity pattern we observed in InsCtx is closely
related to learned anticipation of reward during goal-directed behavior, rather than licking,
the sensory aspects of the reward, general reward, valence, or specific motivational
states.

Discussion

We used unbiased topological and machine learning approaches to investigate population activity in InsCtx. We discovered that InsCtx population activity had consistent dimensionality and manifold structure across days, across different mice, and across different motivational states. Dynamics within the manifold were remarkably stereotyped, allowing us to decode trial outcomes with high accuracy across different animals, motivations, and rewards. These stereotyped dynamics reflected goal-directed reward anticipation, rather than general reward, valence, taste, or licking. We propose that such stereotyped activity patterns could reflect a core computational function of InsCtx across the many processes it is thought to be involved in.

We found that the InsCtx activity manifold (i.e., the repertoire of activity patterns), is constrained by the current behavioral context. Importantly however, we found that the same behavioral context does not yield the same manifold structure in other cortical regions involved in the behavior (e.g., visual cortex), suggesting a conjunction of two constraints that determine manifold structure – behavior and brain region. Several groups have recently reported that activity related to movement and behavior is prevalent and dominant across many brain regions[49–51,59]. Because we found different manifold structure in InsCtx and visual cortex during the same behavior, our results suggest that while behavior does constrain manifold structure, it does so in conjunction with more localized and potentially unique functions of distinct subnetworks of different brain regions.

The InsCtx activity manifold was consistent across individuals and was partially correlated with external and internal behavioral variables, which included pupil size (as a proxy for arousal behavior[52]), cumulative rewards (as a proxy for satiation[33]), and reward consumption. Our analyses were limited by the behavioral variables that were measured in our studies. It remains to be tested whether other parts of the manifold could be determined by the intrinsic connectivity of the network, as has been previously suggested[60], and/or by other variables we did not measure in these experiments[5]. These variables could include interoceptive changes, such as changes in blood pressure, gastric stretch, and blood oxygenation, all previously shown to be represented by single InsCtx

neurons or bulk activity[12,27]. Future studies comparing InsCtx activity across different

behavioral contexts could assess to what extent the intrinsic dimensionality and structure

of the manifold depend on specific ongoing behaviors and sensory inputs.

Most of the cellular level neural activity recordings in the InsCtx of behaving animals have

focused on taste processing[13–15,17]. These studies investigated relatively fast activity

changes on the order of hundreds of milliseconds, revealing sequential temporal coding

of tactile, chemosensory, and palatability information in individual neurons[13]. At the level

of small populations of 5-10 neurons, InsCtx activity has been shown to exhibit metastable

dynamics, transitioning between different states that can reflect different behavioral

variables, each lasting tens to hundreds of milliseconds[60–62]. Other work using activity

manipulations and bulk activity recordings has shown that InsCtx activity is related to

taste, palatability, motivation, valence, and anxiety[19,26,29–31,63,64]. Some of these variables

are likely represented by slower changes across the activity manifold. Nevertheless, the

faster stereotyped changes we discovered appear not to be related to licking, taste

valence, or general reward. Interestingly, a recent study found that activity of anterior

InsCtx projections to the brainstem encodes motivational vigor[30]. Notably, the stereotyped

activity patterns we described here in mid InsCtx likely do not reflect motivational vigor

per se, as they are sensitive to the behavioral task structure, and do not appear to change

with partial satiation, which reduces motivational vigor[30]. Nevertheless, seeing as InsCtx

activity appears to be associated with several different types of information, it will be

important to understand how it can simultaneously represent these different types of

information. Furthermore, it will be important to understand to what extent these different

types of information are eventually integrated before being relayed to downstream brain

regions to affect behavior and bodily physiology. Indeed, different InsCtx projection

populations (e.g., projecting to striatum, thalamus, or brainstem[40,41,65]) may convey

different types of information to downstream regions, and these in turn could differentially

weigh the input from different neurons to simultaneously extract different types of

information.

We previously used linear projections in activity space to suggest that reward

consumption transiently shifts activity towards a future satiety state[33]. Here, we used more

comprehensive analyses of the entire activity manifold and find stereotyped goal-directed reward anticipation activity patterns, which are mostly independent of changes in ongoing activity that reflect physiological state. This further underscores the importance of using both hypothesis-driven, as well as unbiased approaches to explore multi-dimensional neural activity. Thus, while our previous work provided support to a dominant model of interoception[66,67], our analyses here highlight a potentially more general computation InsCtx performs, which could be important for many behaviors InsCtx is involved in, which do not necessarily directly involve interoception.

Activity dynamics on the manifold were remarkably consistent across individuals and motivations, suggesting that these dynamics may reflect an important fundamental aspect of InsCtx function. We speculate that such goal-directed reward anticipation is related to the consistent involvement of InsCtx in cravings during pathological conditions associated with aberrant reward processing, including obesity, binge eating disorder, and drug addiction[65,68–71]. InsCtx has been suggested to be involved in these conditions, as well as in many others, by encoding the anticipated interoceptive value or outcome of actions, which when goes awry can lead to aberrant food or drug cravings[69]. We thus speculate that the stereotyped goal-directed reward anticipation patterns we discovered reflect this fundamental computation in InsCtx. As such, future work using the same analytical framework in mouse models of pathological conditions, such as obesity and drug addiction, could test this idea to form a deeper understanding of the role of InsCtx in pathological conditions that involve aberrant motivations.

Acknowledgements

We thank Mark Andermann, Alon Rubin, Yarden Cohen, Sasha Devore, Stav Shtiglitz, Ayal Lavi, Yael Prilutski, Einav Litvak, Omer Izhaki, Inbar Perets, Daniel Deitch, Tom Talpir, Omer Richmond, Gal Elyasaf, Birgit Jickeli, and Daphna Nachmani for fruitful discussions and helpful comments on the manuscript. YL is supported by research grants from the Israel Science Foundation (#860/21), the European Research Council Starting Grant (ERC StG #101039145), and the Center for New Scientists at the Weizmann Institute of Science.

Methods

Datasets analyzed for this paper were both previously published data[25,33,48], as well as unpublished data. Unpublished datasets included experiments similar to those described in the published datasets (e.g., in water or food restricted mice performing an operant visual discrimination task for water or Ensure rewards), as well as imaging during head-fixed satiation by Ensure consumption and induction of hyperosmotic thirst by intraperitoneal injection of hypertonic saline. We briefly describe the experiments below, see also refs[25,33] for more detailed descriptions.

Behavioral task

To perform the behavioral task, mice were water restricted to ~80% of their pre-restriction body weight or food-restricted to ~85% of their free-feeding body weight. We trained mice to discriminate between square-wave drifting gratings differing in orientation presented on an LCD screen (2 Hz and 0.04 cycles/degree, full-field square wave drifting gratings, 80% contrast; food cue: 0°, aversive cue: 270°, neutral cue: 135°[25,33]. All drifting gratings were presented for 2 sec, after which the mouse had a 2 sec window to respond with a lick. Licking during the visual cue was not punished, but also did not trigger delivery of the water/Ensure/salt-water/quinine. Only the first lick (if any) occurring during the response window triggered delivery of water/Ensure/salt-water/quinine. The lickspout was designed with two adjacent lick tubes (one for each outcome), such that the tongue contacted both tubes on each lick, which served as an effective deterrent for lick responses following aversive cues. Well-trained mice had a high rate of correct water/food cue licking responses (criterion: >80% of trials, usually ~90-95%), and a low rate of licking following aversive cue presentations (criterion: <50%, usually ~20-30%). Each water reward was a ~2-3 µL drop, and Ensure reward was a ~5 µL drop (0.0075 calories). Behavioral training was performed using MonkeyLogic[72] and MonkeyLogic2 (https://monkeylogic.nimh.nih.gov/).

Surgical procedures

Stereotaxic injections AAV8-CaMKII-hM3Dq-mCherry, AAV8-hSyn-DIO-hM3Dq-mCherry, and AAV1-hSyn-GCaMP6f, as well as and implantation of microprisms (2 mm

450   prisms; #MCPH-1.0; Tower Optical; coated with aluminum along their hypotenuse) were
451   performed as previously described[25,33].

Two-photon imaging across different natural and artificial states

Two-photon imaging of GCaMP6f was performed using a resonant-scanning two-photon
microscope with tiltable scanhead (Neurolabware; 31 frames/second; 1154x512 pixels).
All imaging was performed with a 20x 0.45 NA air objective (Olympus) with a 540 x 360
μm2 field of view. All imaged fields of view (FOV) were at a depth of 90-200 μm below
the pial surface, using a Mai Tai DeepSee laser (Newport Corp.) with laser power at 920-
960 nm of 35-80 mW at the front aperture of the objective (power at the sample was likely
substantially less due to partial transmission via the microprism). Imaging depth was
adjusted in between runs (every 30 min) to account for slow drift in the z plane (< 7 μm).

Imaging across thirsty and quenched states

462   We imaged mice during gradual water satiation during consecutive 30-minute runs until
463   the mice voluntarily stopped performing the task. We then performed one more imaging
464   run, the 'quenched' run, in which mice did not respond to the water cue.

Imaging across hungry, satiation, and sated states

We imaged mice in two blocks of trials within a session, one block during food restriction
and a subsequent block immediately following re-feeding. At the start of each imaging
session, food-restricted mice performed the visual cue discrimination task. After ~180
trials (30-min imaging run), we provided the mouse with ad libitum access to Ensure until
voluntary cessation of consumption. Ensure consumption lasted 45-75 minutes. We
triggered delivery of Ensure with every lick, but with a minimum inter-trial interval of 2.5 s
between Ensure deliveries. During this period of time, mice consumed ~3-5 mL of Ensure
and then voluntarily stopped licking for rewards. We then imaged additional ~180 trials
(30 min imaging run) while mice were satiated (operationally defined as the absence of
voluntary licking).

Imaging during chemogenetic activation of SFO^{GLUT} neurons

477   Following imaging during the quenched state (see above), we injected 150 μl 0.9% of
478   either saline or CNO (5 mg/kg), waited 10-15 min and started another imaging run (~180

479   trials, 30 min). For every mouse used for these experiments, we used postmortem
480   histology and immunohistochemistry to verify hM3Dq-mCherry expression in the SFO.

Imaging during following injection of hypertonic saline to induce hyperosmotic thirst

483   Following imaging during the quenched state, we injected ~200 µL of 2M NaCl, waited 5-
484   10min and started another imaging run (~180 trials, 30 min). Mice usually re-engaged in
485   the behavioral task immediately or after 2-3 minutes.

Imaging during chemogenetic activation of AgRP neurons

Following imaging during hunger and satiety, mice were returned to their home-cage with
ad libitum access to regular chow. The next morning, we imaged the same InsCtx field of
view in this satiety state (100-110% of normal body weight) during ~180 trials (30 min.).
We then injected CNO (1-3 mg/kg). Ten minutes later, we initiated an additional imaging
run of ~180 trials. For every mouse used for these experiments, postmortem histology
and immunohistochemistry confirmed hM3Dq-mCherry expression in the hypothalamic
arcuate nucleus.

Pupil videography during two-photon imaging

We acquired data using a GigE Vision camera (Dalsa) with a 60 mm lens (Nikon
MicroNikkor) from a pre-selected region of interest around the eye ipsilateral to the LCD
monitor used to present visual cues (contralateral to the InsCtx microprism). Acquisition
of each frame (frame rate of 15.5 Hz) was triggered on every other frame of two-photon
acquisition (acquired at 31 Hz) using Scanbox software (Neurolabware). The pupil was
backlit with illumination originating from diffusion within the brain of the IR light used for
two-photon excitation during imaging. See below for details of data analysis.

Image registration and time course extraction

First, each acquired image was spatially down sampled by 2X. To correct for motion along
the imaged plane (x-y motion), each frame was registered to an average field-of-view
using efficient subpixel registration methods[73] Within each imaging session, each run (2-
8 runs/session) was registered to the first run of the day. Image stacks were de-noised
using principal components analyses (PCA) of every pixel across time, and by user

identification and removal of noise principal components (low eigenvalues; based on[74]).

Cell masks and calcium activity time courses ('F(t)') were extracted using custom

implementation of common methods[74]. To avoid use of cell masks with overlapping pixels,

we only included the top 75% of pixel weights for a given mask, but users screened each

prospective ROI and could edit the size of the mask, selectively removing the lowest

probability pixels. We then excluded any remaining pixels identified in multiple masks.

We manually verified that all cell masks had typical cell body morphology and size.

Fluorescence time courses were extracted by averaging the pixels within each region-of-

interest ('ROI') mask. Fluorescence time courses for neuropil within a 25 μm annulus

surrounding each ROI (but excluding adjacent ROIs and a protected ring surrounding

each ROI) were also extracted ($F_{neuropil}(t)$: median value from the neuropil ring on each

frame). Fluorescence timecourses were calculated as $F_{neuropil\_corrected}(t) = F_{ROI}(t) -$

$F_{neuropil}(t)$. The change in fluorescence was calculated by subtracting a running estimate

of baseline fluorescence ($F_0(t)$) from $F_{neuropil\_corrected}(t)$, then dividing by $F_0(t)$: $\Delta F/F(t) =$

$(F_{neuropil\_corrected}(t) - F_0(t))/ F_0(t)$. $F_0(t)$ was estimated as the 10th percentile of a 32 sec

sliding window[25,33,48].

Preprocessing of neuronal data

For each collected dataset, we down-sampled $\Delta F/F$ traces by averaging every

consecutive 15 frames (acquired at 31 Hz), resulting in ~0.5 seconds time-bins.

Subsequently, we excluded outlier neurons with abnormally high $\Delta F/F$ values, calibrated

separately for two different microscopes (2000% and 5000%). Following this, we Z-scored

the activity time course for each individual neuron. Each dataset consisted of multiple 30-

minute runs, acquired within the same day. For certain datasets, we applied separate Z-

scoring to each run if substantial extended breaks occurred between runs or if there was

a z-plane shift between the runs.

Intrinsic dimensionality estimation

To estimate the intrinsic dimensionality of the neuronal activity data, we employed the

minimal neighborhood information technique as an estimator[42]. Briefly, the intrinsic

dimension is estimated using the ratio formula:

$$537 \qquad intrinsic\ dimension = -\frac{\log(1 - F(\mu))}{\log(\mu)}$$

Here, μ_i was determined for each data point i, where $\mu_i = \frac{r_{1,i}}{r_{2,i}}$, with $r_{1,i}$ and $r_{2,i}$

representing the Euclidean distances of the nearest and second-nearest data points to

point i, respectively. $F(\mu_i)$ indicates the percentile of μ_i with respect to all other data

points. Subsequently, μ_i was computed for each data point, and the highest 4% of points

were excluded. The intrinsic dimension was defined as the slope calculated from a linear

regression between $\log(1 - F(\mu))$ and $\log(\mu)$ across all data points.

Non-linear dimensionality reduction

To reduce the dimensionality of the neuronal data matrix X, we applied Laplacian

eigenmaps[75] implemented by a pre-established pipeline[76]. Laplacian eigenmaps, a

spectral non-linear dimensionality reduction technique, operates on a data matrix X with

dimensions $N \times T$, by generating a weighted adjacency graph W for the T high-

dimensional data points such as each time point $t \in \mathbb{R}^N$. This adjacency graph is

computed for all pairs of data points, where the weight between a pair of data points

$x_i,\ x_j \in X$ is assigned as $W_{ij} = 1$ if the data points are connected and $W_{ij} = 0$ otherwise.

Data points are considered connected if they fall within the K nearest neighbors of each

other, where K is a user-defined parameter (see below). Following the computation of W,

a diagonal weight matrix D is constructed with $D_{ii} = \sum_j W_{ji}$, which is equivalent to

aggregating the rows of W. Notably, the Laplacian of the original data matrix is $L = D -$

W. The process of obtaining the low-dimensional embedding is achieved by calculating

the eigenvalues and eigenvectors of $Lf = \lambda Df$. After excluding the leading eigenvector[75],

the following m eigenvectors are utilized to obtain a new low-dimensional mapping for

each data point x_i, which is given by[17]

$$560 \qquad x_i \rightarrow (f_1(i), \ldots, f_m(i))$$

with m representing the desired output dimensionality. For every dataset X, we conducted

two rounds[38] of dimensionality reduction using the algorithm described above. In the initial

iteration, a 20-dimensional matrix was generated, with a parameter K set to 7.5% of the

time points in matrix X. The second iteration was performed on the 20-dimensional matrix

obtained from the preceding step. This culminated in the matrix reaching its ultimate
dimension – the mean estimated intrinsic dimension of 6 – with a parameter K set to 2.5%
of the time points in matrix X. The mean estimated intrinsic dimension of 6 was derived
from the process described above.

Topological features analysis and topological similarity

To assess the topological features of the data, we quantified for each dataset its
components (β0) and holes (β1) as a function of the radius threshold, using a pre-
established algorithm[45] (TDAStats: R pipeline for computing persistent homology in
topological data analysis, https://github.com/rrrlw/TDAstats). Due to computational
considerations, we determined the topological characteristics of clustered timepoints
within the reduced data matrix, acquired through K-means clustering with a chosen K
value of 80. Given our utilization of a stochastic variant of K-means, this process was
repeated 20 times for each dataset. For every topological feature extracted from the
clustered data matrix, we quantified its lifespan and derived the lifespan distribution for
all topological features. The topological resemblance between a pair of datasets was
defined as the mean Wasserstein distance[77] between the lifespan distribution of features
from both datasets, covering all 20 iterations of K-means clustering. To ascertain the
significance of differences between dataset pairs, permutation tests were conducted,
following a methodology outlined in prior work[47]. The lifespan distributions of a pair of
datasets were pooled and then systematically rearranged to generate two separate
subsamples, adhering to the original sizes of each distribution. The Wasserstein distance
was computed between these subsamples. This procedure was repeated n=200 times,
and a P-value was established as the percentage of permutations yielding a Wasserstein
distance less than the original distance between the dataset pairs. The ultimate P-value
for every pair of datasets was determined as the average of the P-values derived from
this process across all 20 K-means clustering iterations.

Clustering the activity manifold

To evaluate the activity manifold, our objective was to create a standardized comparison
across datasets through an automated clustering approach. To achieve this, we devised
an unsupervised clustering strategy that recognizes data points that deviate from the
central point cloud, and thus classifies them as distinct neuronal clusters. To do so we

systematically excluded time points that were closest to the centroid of the reduced data
matrix. Starting from 40 times points and progressing to the entire dimensionality of the
data matrix. For each set of excluded timepoints, we conducted 20 iterations of K-means
clustering with K values spanning from 2 to 20. The mean squared error (MSE) was
computed between the resulting clusters and their centroids across all iterations and for
each K value. This procedure produced a two-dimensional grid with its two dimensions
being the fraction of excluded timepoints and the chosen K values for K-means clustering.
The cells of the grid contained MSE values over the 20 clustering repetitions. To identify
the optimal configuration, we scaled both grid dimensions to 1 and determined the grid
cell closest to the origin of a three-dimensional Cartesian system formed by the grid,
where the third dimension was the MSE value within each cell. This process yielded, for
each dataset an optimal configuration consisting of 2 parameters: a K value, and the
number of datapoints to be considered as the central point cloud. Across virtually all
utilized datasets the identified optimal K value was 8, except one dataset in which it was
7. After obtaining the optimal configuration, our aim was to derive deterministic clustering
labels for each dataset. This was accomplished by implementing the optimal
configuration, isolating timepoints deemed part of the central point cloud, and executing
500 iterations of K-means clustering with the optimal K value for the remaining timepoints.
The resulting clustering labels were aggregated into a $500 \times T$ matrix. This matrix was
then subjected to an additional clustering step using the ward agglomerative complete
linkage hierarchical clustering method[78]. This yielded the final labels of the clustered
activity manifold.

Activity manifold parameterization

To evaluate the contribution of behavioral variables to the activity manifold, we devised
labeling for the distinct behavioral variables, considering the total timepoints under
analysis. These variables were categorized into two groups: discrete (binary) and
continuous. Discrete binary variables encompassed the potential classes of trial
outcomes within the behavioral task (e.g., response vs. no response for a given trial type).
For each of the six trial outcome classes, a corresponding label vector was generated,
spanning T timepoints. A value of $Trial\ outcome\ Label = 1$ was assigned for timepoints
ranging from 3 to 6 seconds after the cue onset for each class, while a value of

$Trial\ outcome\ Label = 0$ was allocated to the remaining timepoints. This enabled the

isolation of timepoints exclusively linked to that specific trial outcome, excluding those

related to visual cues. As for continuous variables, namely Pupil Size and Cumulative

Consumed Rewards, their magnitude across the entire experiment was quantified. Pupil

size was quantified using code from previously published work[79]. Two label vectors, each

with a length of T, were established for these variables. To ensure equitable comparison

with discrete variables, both Pupil Size and Cumulative Consumed Rewards were divided

into eight bins.

Mutual information between behavioral variables and the activity manifold

To determine the mutual information between behavioral variables and the activity

manifold, we constructed a joint contingency table that related the two label vectors:

$Cluster$ and $Behavior$. Here, $Cluster$ represents a vector of size T containing different

neuronal cluster labels (including the central point cloud), while Behavior was a T-sized

vector representing behavioral variables (ranging from 0 to 1 for discrete variables and 1

to 8 for continuous variables). Following this, we utilized the contingency table to

formulate a joint probability space, denoted as $P_{Cluster \times Behavior}$. The calculation of mutual

information was carried out according to the formula:

644
$$MI = \sum_{c} \sum_{b} P_{Cluster \times Behavior}(c,b) * \log\left(\frac{P_{Cluster \times Behavior}(c,b)}{P_{Cluster}(c) * P_{Behavior}(b)}\right)$$

Where $P_{Cluster}$ and $P_{Behavioor}$ are the respective marginal probability distributions. To

reduce potential variations caused by different behavioral factors and ensure an equitable

comparison with discrete variables, for the continuous variables we exclusively included

time points unaffected by behavior, following established procedures from our previous

work[33]. These time points corresponded to the final 3 seconds of inter-trial intervals

following the correct rejection of aversive and neutral trials (referred to as non-responded

trials).

Trial structure manifold dynamics

To evaluate manifold dynamics in relation to the trial structure, we compiled the sequential

neuronal cluster label (including the central point cloud) time course for every trial. This

was achieved by capturing a 10-second interval from the onset of the cue for each trial

class. They yielded a sequence of 20 neuronal cluster labels, subsequently combined

into an $M \times 20$ matrix. Here, M denotes the trial count, and each row signifies the

sequence of visited clusters during the respective trial.

Trial similarity

To assess trial dynamics similarity, our focus was on highlighting the sequence of

neuronal clusters. To achieve this, we designated the label -1 to each timepoint

associated with the central point cloud. Meanwhile, other timepoints were allocated labels

ranging from 1 to 7, corresponding to the respective neuronal cluster assignments.

Subsequently, we created a trial structure matrix as previously outlined and computed

pairwise trial similarity using cosine distance.

666
$$Trial\ Similarity(Trial_i, Trial_j) = \frac{Trial_i \cdot Trial_j}{\|Trial_i\| \ \|Trial_j\|}$$

Here, $Trial_i$ represents the timepoint sequence of labels in the ith trial. This procedure

resulted in a $M \times M$ for each dataset. Following that, we concentrated on the lower

triangle of the matrix and pooled all values across all datasets.

Decoding trial outcome across datasets

To enable across-datasets decoding, we generated a trial dynamics matrix using the

same labeling approach employed for assessing trial similarity. Our goal was to enable

decoding between datasets, which necessitated an alignment process. For each dataset,

we initiated alignment by constructing a transition probability matrix between all neuronal

clusters and the central point cloud. This process yielded a 8×8 probability matrix for

each dataset, except for one dataset with 7 clusters that was excluded from this analysis.

Following this, the probabilities underwent clustering through the ward agglomerative

complete linkage hierarchical clustering method[78]. Using the hierarchical clustering

structure, we derived a cluster order based on the similarity of their probabilities. This

order served as a mask across datasets to allow translation of clustering labels among

dataset pairs. Essentially, this step aimed to address the question: Given transition

probabilities, which cluster in dataset A most closely aligns with a cluster in dataset B.

With the labels clusters of dataset B aligned to dataset A, we proceeded to compute the

pairwise trial similarity between each trial pair in both datasets. For each trial a in dataset

A, the decoded class of trial outcome was determined by maximizing the sum of Trial
Similarity values over trials b in dataset B:

687
$$argmax_{class} \sum_{b \in B} Trial\ Similarity(Trial_{b(class)}, Trial_a)$$

This process is analogous to identifying, on average after translation, which class of trials
in dataset B most closely resembles an individual trial in dataset A. To further validate the
decoding accuracy, we established a surrogate shuffle distribution by randomly shuffling
trial classes in both datasets 200 times. We then calculated the decoding accuracy for
each shuffle and computed the P-value by determining the percentile of shuffles that
exhibited a higher decoding accuracy compared to the measured decoding accuracy.
This entire process was repeated for every pairwise combination of datasets.

Decoding trial outcome across datasets based on overall activity levels and thirsty-
quenched linear axis

To decode trial outcomes across dataset based on overall activity levels, we calculated
the overall activity levels within each trial by averaging the collective population activity
across neurons. To decode trial outcomes across dataset based on the thirsty-
quenched linear axis, we defined this axis for each mouse by $\bar{x}_{Thirsty} - \bar{x}_{Quenched}$, where
$\bar{x}$ is the population vector of mean ongoing activity in each state. We projected peri-cue
activity onto this axis by calculating the dot product of this vector with the time-varying
pattern of InsCtx population activity, x(t). We then scaled values along this axis per
mouse, ascribing a value of 1 when x(t) = $\bar{x}_{Thirsty}$, a value of 0 when x(t) = $\bar{x}_{Quenched}$ as
0, and intermediate values for patterns that fall between $\bar{x}_{Thirsty}$ and $\bar{x}_{Quenched}$
projected this data onto a thirsty-quenched linear axis, as outlined in reference[33].
Second, we utilized a similar decoding as explained above. Instead of training it with
aligned trial labels for pairs of trials, we evaluated used the overall activity levels within
each trial or the projection of the trial onto the thirsty-quenched axis as a training
dataset.

Quantification of anticipatory licking onset

The onset of anticipatory licking during a trial was defined as the time point when the first
licking bout took place within the response window after cue presentation. Licking bouts

were considered as instances where mice notably increased their licking rate. To detect
these bouts, for each trial, we calculated the time intervals between consecutive licks
occurring after cue presentation. We then identified the pair of licks with intervals shorter
than a specific threshold:

$$718 \qquad \frac{1}{\mu_{Cue\ licks} + C * \sigma_{Cue\ licks}}$$

719 Where $\mu_{Cue\ licks}$, $\sigma_{Cue\ licks}$ were determined by calculating the average and standard
720 deviation of licks based on the first 15 frames (0.5 sec) of cue presentation across all
721 trials. The constant C varied; it was set as 0.1 for datasets involving water rewards and
722 0.35 for datasets involving food rewards. The time point at which the first pair of licks
723 exceeded this threshold was considered the onset of anticipatory licking.

Quantification of cluster onset

To identify the onset of the cluster sequence, we first determined the two neuronal
clusters with the highest occupancy during the response window for each dataset.
Following this, we identified the initial time point within each trial where a transition
occurred from the central point cloud to each of the identified clusters. The onset was
defined as the first time point in each trial where the transition occurred with the cluster
that exhibited the highest onset correlations with the lick onsets across all trials.

Lick similarity between pairs of trials

732 To define the lick similarity between pairs of trials, for each trial, we smoothed the binary
733 licks vector $Licks_i$. To do so we applied a sliding window average of 100 frames (~3.3
734 sec). The smoothed lick pattern for the ith trial was calculated as follows:

$$735 \qquad SL_{i,a} = \frac{1}{100} \sum_{a}^{a+100} Licks_{i,a}$$

736 Where a is the ath licking frame in the original binary vector and SL_i is the overall
737 smoothed licking pattern. Lick similarity between the ith and jth trials was determined
738 using the formula:

$$739 \qquad Lick\ Similarity(Licks_i, Licks_j) \frac{1}{\left\| SL_i - SL_j \right\|^2}$$

Construction pseudo-trials in freely consuming mice.

To construct pseudo-trials of consumption in freely consuming mice, we identified time points in which mice initiated a licking bout. To do so we identified all the time points within the free consumption epoch of the experiment in which the interval between two consecutive licks was above 1 second, unless this did not yield enough pseudo-trials, in which case we used 0.5 sec as the threshold (2/5 datasets). Every pair of licks that passed these criteria could be used to define the onset of a pseudo-trial. We then took all the time points that occurred two seconds before and four seconds after the second lick in that interval. We used these time points to identify the neuronal cluster dynamics and licking pattern that took place in the experiment and constructed a pseudo-trial matrix. We then removed pseudo-trials that overlapped with each other to avoid inflation of the matrix. To control for behavioral differences between operant task licking and free consumption licking, we repeated the analyses in (Fig. 4j-i) while selecting only pseudo-trials of free consumption in which the average licking rate in the first two seconds of the lick bout were within the mean ± standard deviation of the average licking rate in the first two seconds of operant task licking (Supp. Fig. 7c-e).

References

1. Vyas, S., Golub, M. D., Sussillo, D. & Shenoy, K. V. Computation Through Neural Population
Dynamics. *Annu Rev Neurosci* 249–275 (2020) doi:10.1146/annurev-neuro-092619.

2. Jazayeri, M. & Afraz, A. Navigating the Neural Space in Search of the Neural Code. *Neuron* vol. 93
1003–1014 Preprint at https://doi.org/10.1016/j.neuron.2017.02.019 (2017).

3. Saxena, S. & Cunningham, J. P. Towards the neural population doctrine. *Current Opinion in
Neurobiology* vol. 55 103–111 Preprint at https://doi.org/10.1016/j.conb.2019.02.002 (2019).

4. Schneider, S., Lee, J. H. & Mathis, M. W. Learnable latent embeddings for joint behavioural and
neural analysis. *Nature* **617**, 360–368 (2023).

5. Mitchell-Heggs, R., Prado, S., Gava, G. P., Go, M. A. & Schultz, S. R. Neural manifold analysis of
brain circuit dynamics in health and disease. *Journal of Computational Neuroscience* vol. 51 1–21
Preprint at https://doi.org/10.1007/s10827-022-00839-3 (2023).

6. Langdon, C., Genkin, M. & Engel, T. A. A unifying perspective on neural manifolds and circuits for
cognition. *Nat Rev Neurosci* **24**, 363–377 (2023).

7. Gardner, R. J. *et al.* Toroidal topology of population activity in grid cells. *Nature* **602**, 123–128
(2022).

8. Chaudhuri, R., Gerçek, B., Pandey, B., Peyrache, A. & Fiete, I. The intrinsic attractor manifold and
population dynamics of a canonical cognitive circuit across waking and sleep. *Nat Neurosci* **22**,
1512–1520 (2019).

9. Ajabi, Z., Keinath, A. T., Wei, X.-X. & Brandon, M. P. Population dynamics of head-direction
neurons during drift and reorientation. *Nature* (2023) doi:10.1038/s41586-023-05813-2.

10. Gallego, J. A., Perich, M. G., Chowdhury, R. H., Solla, S. A. & Miller, L. E. Long-term stability of
cortical population dynamics underlying consistent behavior. *Nat Neurosci* **23**, 260–270 (2020).

11. Churchland, M. M. *et al.* Neural population dynamics during reaching. *Nature* **487**, 51–56 (2012).

12. Cechetto, D. F. & Saper, C. B. *Evidence for a Viscerotopic Sensory Representation in the Cortex
and Thalamus in the Rat*. *THE JOURNAL OF COMPARATIVE NEUROLOGY* vol. 262 (1987).

13. Katz, D. B., Simon, S. A. & Nicolelis, M. A. L. *Dynamic and Multimodal Responses of Gustatory
Cortical Neurons in Awake Rats*. (2001).

14. Jones, L. M., Fontanini, A. & Katz, D. B. Gustatory processing: a dynamic systems approach. *Curr
Opin Neurobiol* **16**, 420–428 (2006).

15. Kadohisa, M., Rolls, E. T. & Verhagen, J. V. Neuronal Representations of Stimuli in the Mouth: The
Primate Insular Taste Cortex, Orbitofrontal Cortex and Amygdala. *Chem Senses* **30**, 401–419
(2005).

16. Small, D. M. Flavor is in the brain. *Physiol Behav* **107**, 540–552 (2012).

17. De Araujo, I. E. & Simon, S. A. The gustatory cortex and multisensory integration. *International
Journal of Obesity 2009 33:2* **33**, S34–S43 (2009).

18.   Gardner, M. P. H., Fontanini, A., Gardner, M. P. H. & Fontanini, A. Encoding and Tracking of
795   Outcome-Specific Expectancy in the Gustatory Cortex of Alert Rats. *Journal of Neuroscience* **34**,
796   13000–13017 (2014).

797   19.   Samuelsen, C. L., Gardner, M. P. H. & Fontanini, A. Effects of Cue-Triggered Expectation on
798   Cortical Processing of Taste. *Neuron* **74**, 410–422 (2012).

799   20.   de Araujo, I. E. *et al.* Neural Ensemble Coding of Satiety States. *Neuron* **51**, 483–494 (2006).

800   21.   Rodgers, K. M., Benison, A. M., Klein, A. & Barth, D. S. Auditory, Somatosensory, and
801   Multisensory Insular Cortex in the Rat. *Cerebral Cortex* **18**, 2941–2951 (2008).

802   22.   Stapleton, J. R., Lavine, M. L., Wolpert, R. L., Nicolelis, M. A. L. & Simon, S. A. Rapid Taste
803   Responses in the Gustatory Cortex during Licking. *Journal of Neuroscience* **26**, 4126–4138 (2006).

804   23.   Vincis, R. & Fontanini, A. Associative learning changes cross-modal representations in the
805   gustatory cortex. *Elife* **5**, (2016).

806   24.   Yamamoto, T., Matsuo, R., Kiyomitsu, Y. & Kitamura, R. Sensory inputs from the oral region to the
807   cerebral cortex in behaving rats: an analysis of unit responses in cortical somatosensory and taste
808   areas during ingestive behavior. *https://doi.org/10.1152/jn.1988.60.4.1303* **60**, 1303–1321
809   (1988).

810   25.   Livneh, Y. *et al.* Homeostatic circuits selectively gate food cue responses in insular cortex. *Nature*
811   **546**, 611–616 (2017).

812   26.   Gehrlach, D. A. *et al.* Aversive state processing in the posterior insular cortex. *Nat Neurosci* **22**,
813   1424–1437 (2019).

814   27.   Klein, A. S., Dolensek, N., Weiand, C. & Gogolla, N. Fear balance is maintained by bodily feedback
815   to the insular cortex in mice. *Science (1979)* **374**, 1010–1015 (2021).

816   28.   Dolensek, N., Gehrlach, D. A., Klein, A. S. & Gogolla, N. Facial expressions of emotion states and
817   their neuronal correlates in mice. *Science (1979)* **368**, (2020).

818   29.   Nicolas, C. *et al.* Linking emotional valence and anxiety in a mouse insula-amygdala circuit. *Nat*
819   *Commun* **14**, 5073 (2023).

820   30.   Deng, H. *et al.* A genetically defined insula-brainstem circuit selectively controls motivational
821   vigor. *Cell* **184**, 6344-6360.e18 (2021).

822   31.   Wang, L. *et al.* The coding of valence and identity in the mammalian taste system. *Nature* **558**,
823   127–131 (2018).

824   32.   Egan, G. *et al.* Neural correlates of the emergence of consciousness of thirst. *Proc Natl Acad Sci U*
825   *S A* **100**, 15241–15246 (2003).

826   33.   Livneh, Y. *et al.* Estimation of Current and Future Physiological States in Insular Cortex. *Neuron*
827   **105**, 1094-1111.e10 (2020).

828   34.   Meier, L. *et al.* Thirst-dependent activity of the insular cortex reflects its emotion-related
829   subdivision: a cerebral blood flow study. *Elsevier*.

35. Tataranni, P. A. *et al.* Neuroanatomical correlates of hunger and satiation in humans using
positron emission tomography. *Proc Natl Acad Sci U S A* **96**, 4569–4574 (1999).

36. Jazayeri, M. & Ostojic, S. Interpreting neural computations by examining intrinsic and embedding
dimensionality of neural activity. *Current Opinion in Neurobiology* vol. 70 113–120 Preprint at
https://doi.org/10.1016/j.conb.2021.08.002 (2021).

37. Nieh, E. H. *et al.* Geometry of abstract learned knowledge in the hippocampus. *Nature* **595**, 80–
84 (2021).

38. Rubin, A. *et al.* Revealing neural correlates of behavior without behavioral measurements. *Nat*
Commun **10**, (2019).

39. Humphries, M. D. Strong and weak principles of neural dimension reduction. *Neuron Behav Data*
Anal Theory **5**, 1–28 (2021).

40. Gehrlach, D. A. *et al.* A whole-brain connectivity map of mouse insular cortex. *Elife* **9**, 1–78
(2020).

41. Saper, C. B. The central autonomic nervous system: Conscious visceral perception and autonomic
pattern generation. *Annual Review of Neuroscience* vol. 25 433–469 Preprint at
https://doi.org/10.1146/annurev.neuro.25.032502.111311 (2002).

42. Facco, E., D'Errico, M., Rodriguez, A. & Laio, A. Estimating the intrinsic dimension of datasets by a
minimal neighborhood information. *Sci Rep* **7**, (2017).

43. Deitch, D., Rubin, A. & Ziv, Y. Representational drift in the mouse visual cortex. *Current Biology*
31, 4327-4339.e6 (2021).

44. Rybakken, E., Baas, N. & Dunn, B. Decoding of neural data using cohomological feature
extraction. *Neural Computation* vol. 31 68–93 Preprint at https://doi.org/10.1162/neco_a_01150
(2019).

45. R. Wadhwa, R., F.K. Williamson, D., Dhawan, A. & G. Scott, J. TDAstats: R pipeline for computing
persistent homology in topological data analysis. *J Open Source Softw* **3**, 860 (2018).

46. Rouse, T. C., Ni ID, A. M., Huang, C. & Cohen, M. R. Topological insights into the neural basis of
flexible behavior. *Proceedings of the National Academy of Sciences* **120**, (2023).

47. Robinson, A. & Turner, K. Hypothesis testing for topological data analysis. *J Appl Comput Topol* **1**,
241–261 (2017).

48. Burgess, C. R. *et al.* Hunger-Dependent Enhancement of Food Cue Responses in Mouse Postrhinal
Cortex and Lateral Amygdala. *Neuron* **91**, 1154–1169 (2016).

49. Salkoff, D., Zagha, E., … E. M.-C. & 2020, undefined. Movement and performance explain
widespread cortical activity in a visual detection task. *academic.oup.comDB Salkoff, E Zagha, E*
McCarthy, DA McCormickCerebral Cortex, 2020•academic.oup.com.

50. Musall, S., Kaufman, M., Juavinett, A., Gluf, S. & Churchland, A. Single-trial neural dynamics are
dominated by richly varied movements. *Nat Neurosci* **22**, 1677–1686 (2019).

51. Stringer, C. *et al.* Spontaneous behaviors drive multidimensional, brainwide activity. *Science*
(1979) **364**, (2019).

52. McGinley, M. J. *et al.* Waking State: Rapid Variations Modulate Neural and Behavioral Responses.
Neuron **87**, 1143–1161 (2015).

53. Tenenbaum, J. B., de Silva, V. & Langford, J. C. A Global Geometric Framework for Nonlinear
Dimensionality Reduction. *Science (1979)* **290**, 2319–2322 (2000).

54. Augustine, V., Gokce, S. K. & Oka, Y. Peripheral and Central Nutrient Sensing Underlying Appetite
Regulation. *Trends Neurosci* **41**, 526–539 (2018).

55. Andermann, M. L. & Lowell, B. B. Toward a Wiring Diagram Understanding of Appetite Control.
Neuron **95**, 757–778 (2017).

56. Zimmerman, C. A. & Knight, Z. A. Layers of signals that regulate appetite. *Curr Opin Neurobiol* **64**,
79–88 (2020).

57. Sternson, S. M. & Eiselt, A. K. Three Pillars for the Neural Control of Appetite.
https://doi.org/10.1146/annurev-physiol-021115-104948 **79**, 401–423 (2017).

58. Zimmerman, C. A. *et al.* Thirst neurons anticipate the homeostatic consequences of eating and
drinking. *Nature* **537**, 680–684 (2016).

59. Allen, W. E. *et al.* Thirst regulates motivated behavior through modulation of brainwide neural
population dynamics. *Science (1979)* **364**, (2019).

60. Mazzucato, L., La Camera, G. & Fontanini, A. Expectation-induced modulation of metastable
activity underlies faster coding of sensory stimuli. *Nat Neurosci* **22**, 787–796 (2019).

61. Jones, L., Fontanini, A., Sadacca, B., Miller, P. & Katz, D. Natural stimuli evoke dynamic sequences
of statesin sensory cortical ensembles. *Proc Natl Acad Sci U S A* **104**, 18772–18777 (2007).

62. Lang, L., Camera, G. La & Fontanini, A. Temporal progression along discrete coding states during
decision-making in the mouse gustatory cortex. *PLoS Comput Biol* **19**, (2023).

63. Mukherjee, N., Wachutka, J. & Katz, D. B. Impact of precisely-timed inhibition of gustatory cortex
on taste behavior depends on single-trial ensemble dynamics. *Elife* **8**, (2019).

64. Yiannakas, A. *et al.* Parvalbumin interneuron inhibition onto anterior insula neurons projecting to
the basolateral amygdala drives aversive taste memory retrieval. *Curr Biol* **31**, 2770-2784.e6
(2021).

65. Livneh, Y. & Andermann, M. L. Cellular activity in insular cortex across seconds to hours:
Sensations and predictions of bodily states. *Neuron* vol. 109 3576–3593 Preprint at
https://doi.org/10.1016/j.neuron.2021.08.036 (2021).

66. Barrett, L. F. & Simmons, W. K. Interoceptive predictions in the brain. *Nature Reviews*
Neuroscience vol. 16 419–429 Preprint at https://doi.org/10.1038/nrn3950 (2015).

67. Owens, A. P., Allen, M., Ondobaka, S. & Friston, K. J. Interoceptive inference: From computational
neuroscience to clinic. *Neurosci Biobehav Rev* **90**, 174–183 (2018).

68.   Uddin, L. Q. Salience processing and insular cortical function and dysfunction. *Nature Reviews*
903   *Neuroscience* vol. 16 55–61 Preprint at https://doi.org/10.1038/nrn3857 (2015).

904   69.   Naqvi, N. H., Gaznick, N., Tranel, D. & Bechara, A. The insula: A critical neural substrate for
905   craving and drug seeking under conflict and risk. *Ann N Y Acad Sci* **1316**, 53–70 (2014).

906   70.   Craig, A. D. Interoception: the sense of the physiological condition of the body. *Curr Opin*
907   *Neurobiol* **13**, 500–505 (2003).

908   71.   Gogolla, N. The insular cortex. *Current Biology* 573–591 (2017).

909   72.   Asaad, W. F. & Eskandar, E. N. A flexible software tool for temporally-precise behavioral control
910   in Matlab. *J Neurosci Methods* **174**, 245–258 (2008).

911   73.   Bonin, V., Histed, M. H., Yurgenson, S. & Clay Reid, R. Local diversity and fine-scale organization
912   of receptive fields in mouse visual cortex. *Journal of Neuroscience* **31**, 18506–18521 (2011).

913   74.   Mukamel, E. A., Nimmerjahn, A. & Schnitzer, M. J. Automated Analysis of Cellular Signals from
914   Large-Scale Calcium Imaging Data. *Neuron* **63**, 747–760 (2009).

915   75.   Belkin, M. & Niyogi, P. Laplacian Eigenmaps for Dimensionality Reduction and Data
916   Representation. *Neural Comput* **15**, 1373–1396 (2003).

917   76.   Kraemer, G., Reichstein, M. & Mahecha, M. D. dimRed and coRanking-Unifying Dimensionality
918   Reduction in R. *R J* **10**, 342–358 (2018).

919   77.   Ramdas, A., Garcia, N. & Cuturi, M. On Wasserstein Two Sample Testing and Related Families of
920   Nonparametric Tests. *ArXiv* (2015).

921   78.   Murtagh, F. & Legendre, P. Ward's Hierarchical Agglomerative Clustering Method: Which
922   Algorithms Implement Ward's Criterion? *J Classif* **31**, 274–295 (2014).

923   79.   Winkelmeier, L. *et al.* Striatal hub of dynamic and stabilized prediction coding in forebrain
924   networks for olfactory reinforcement learning. *Nat Commun* **13**, (2022).

Figure 1

Figure 1 – InsCtx activity manifold structure is stereotyped across different mice and motivational states

a. Schematic of the operant visual discrimination task.

b. Two-photon calcium imaging of InsCtx through a microprism, schematic coronal brain section and approach.

c. Illustration of the approach to study population activity in the InsCtx.

d. Three representative planes of dimensionality-reduced neuronal activity over time, from three different thirsty mice. Each point represents the population activity pattern in a 0.5 sec time bin. Note the high qualitative similarity between different mice.

e. Estimation of intrinsic dimensionality for each individual dataset vs. shuffled data. The estimated dimensionality was significantly lower than chance (P < $7 * 10^{-5}$, one-tailed Wilcox signed-rank test).

f. Quantification of topological dissimilarity ($\beta 0$ features) between all datasets (see also Supp. Figure 2). Notably, there was a significant difference observed when comparing either across days or across mice to chance levels (P < 0.0003, one-tailed Wilcox signed-rank test), while comparisons of the topological dissimilarity across days and across mice yielded no significant differences (P > 0.1, two-tailed Wilcox rank sum test).

g. Same as in 'd'., for data obtained from hungry mice. Different mice than those in 'd'.

h. Similar estimated intrinsic dimensionality across hungry and thirsty mice (P > 0.4, two-tailed Wilcox rank-sum test).

i. Similar activity manifold shape (topological dissimilarity, $\beta 0$ features) across hungry and thirsty mice performing a similar behavioral task with different motivations (hunger and thirst) and different rewards (food and water; P > 0.25, two-tailed Wilcox rank-sum test).

j. Comparison of estimated intrinsic dimensionality in InsCtx and cortical visual areas in hungry mice performing the same behavioral task during the same motivational state. Intrinsic dimensionality was significantly higher in visual areas (P < 0.015, one-tailed Wilcoxon rank-sum test).

956   k. Comparison of activity manifold shape (topological dissimilarity, $\beta0$ features) between

957   InsCtx and visual areas in hungry mice performing the same behavioral task. The shape

958   of the activity manifold was significantly more different, i.e., less consistent, in visual areas

959   (P < 0.0035, one-tailed Wilcox rank-sum test).

Figure 2

Figure 2 – InsCtx activity dynamics within the manifold are stereotyped across different mice

a. Example visualization of time-points on the activity manifold that are associated with certain behavioral variables: red points depict time points during denoted trial types. Different colors in the clustered structure reflect the assignment of different timepoints into neuronal clusters based on the automated clustering method (see Supp. Fig 3). Each row depicts a different plane of the activity manifold.

b. Mutual Information between behavioral variables and the activity manifold. Pupil size, cumulative consumed rewards, and rewarded trials had significantly greater mutual information compared to other variables (P < 0.0004, one-tailed Wilcoxon signed-rank test with Bonferroni correction). Cumulative consumed rewards had significantly more mutual information than rewarded trials (P < 0.02, one-tailed Wilcoxon signed-rank test with Bonferroni correction).

c. The most informative variables in 'b' exceeded chance levels, assessed by comparisons to randomly shuffled labels (P < 0.00095, one-tailed Wilcoxon signed-rank test).

d. Example cluster sequence trial dynamics in two different mice. Left: two representative planes of the clustered structure (mouse 1 is the same as in 'a'), with different clusters represented distinctly by different colors. Right: example of the cluster sequence dynamics for different trial types and outcomes (same cluster colors as the structures on the left). Note that for each mouse, cluster indices (colors) are randomly selected during the clustering procedure. Rewarded trials exhibit a consistent sequence of cluster transitions, while dynamics in other trial types appear less consistent.

e. Distribution of trial similarity (cosine similarity) within each different trial type using data pooled from 14 datasets from 5 mice. The distribution of rewarded trials had the highest cosine similarity (0.32 ± 0.37; mean ± SD) and was significantly higher than all other trial types (P < $5 * 10^{-10}$, two-sample Kolmogorov-Smirnov test with Bonferroni correction).

f. Decoding procedure across datasets. The cluster labels of different timepoints are translated across datasets by aligning the order of the transition probability matrix

between the different neuronal clusters. The translated dataset is then used for training

to decode the different trial outcomes of the test dataset.

g. Decoding accuracy for different trial types. We used 14 datasets for decoding across

mice and 12 datasets for decoding across days. Rewarded trials were decoded

significantly better both across days and across mice compared to chance level. Notably,

decoding accuracy across days and across mice did not significantly differ ($P < 0.00075$

for across days and across mice versus shuffle, one-tailed Wilcox signed-rank test with

Bonferroni correction; $P > 0.2$ for comparison between across days and across mice, two-

tailed Wilcox rank-sum test with Bonferroni correction). Other trial types were not decoded

better than chance levels, whether decoding across days or across mice ($P > 0.2$, one-

tailed Wilcox signed-rank test with Bonferroni correction).

a-g. N=14 datasets from 5 mice

1005

Figure 3

1008

Figure 3 – InsCtx activity dynamics within the manifold are stereotyped across different motivational states and rewards

a. Visualization of reward cue trial dynamics in two representative mice from InsCtx and visual areas from hungry mice. Notably, activity in InsCtx displays a consistent and stereotyped pattern of activity in hungry mice during reward cue trials that is not apparent after satiation, similar to that of thirsty mice (Fig. 2). In contrast, activity in visual areas primarily exhibits activity during the presentation of the visual cue and is not affected by whether trials were rewarded or not (black and grey vertical bars, respectively).

b. Decoding accuracy of rewarded trials from hungry mice. The decoder was trained using datasets from InsCtx in thirsty mice or from visual areas in hungry mice. Decoding accuracy was significantly higher when training the decoder on InsCtx of thirsty mice as compared to visual areas of hungry mice ($P < 0.0002$ for decoded hungry data trained on thirsty compared to decoded hungry trained on visual area; $P > 0.35$ for decoded hungry data trained on visual areas compared to shuffled data; $P < 0.0065$ for decoded hungry data trained on thirsty compared to shuffled data, Dunn test with Bonferroni correction). N=9 datasets from 6 mice.

c. Left: Example activity dynamics of reward cue trials. The datasets were obtained before and after artificial induction of hunger using chemogenetic activation of AgRP "hunger neurons". The red arrow indicates the first responded reward cue trial following the reinstatement of behavior due to the artificial induction of the motivational state. Note that the neuronal dynamics of rewarded trials before and after activation are similar.

d. Distribution of pairwise trial similarity (cosine similarity) among rewarded trials across natural and artificially induced hunger. Left: the distributions pooled from all datasets. Right: paired comparisons of the mean pairwise trial similarity for each dataset trials across natural and artificially induced motivational states. The artificially induced motivational states did not significantly alter the dynamics ($P > 0.15$, two-tailed paired t-test).

e. Comparison of the decoding accuracy of rewarded trials across natural and artificially induction of hunger. The artificially induced motivational states did not impact the

decoding accuracy of responded reward trials, which remained significantly higher than chance ($P > 0.3$, for comparisons between natural and artificial data across all states; $P > 0.08$ for comparisons between natural and artificial shuffles across all states; $P < 0.008$ for comparisons between data and shuffle for both natural and artificial; one-tailed Wilcox signed-rank test with Bonferroni correction).

f-h. Same as in 'c'-'e' before and after artificial induction of thirst using chemogenetic activation of SFO "thirst neurons".

i-k. Same as in 'c'-'e' with before and after induction of hypertonic thirst using injection of hypertonic saline.

a-k. Data from N=10 datasets (N=7 mice) for hungry datasets, N=14 datasets (N=5 mice) for thirsty datasets, and N=7 datasets (N=5 mice) for visual areas datasets, N=4 datasets (4 mice) for AgRP activation, N=4 datasets (N=4 mice) for SFO activation, N=3 datasets (N=3) mice for hypertonic saline injection.

 Figure 4

1053

Figure 4 – Stereotyped activity dynamics reflect goal-directed reward anticipation, but not licking, taste, or valence

a. Example of neuronal dynamics during rewarded trials and task-related licking behavior. The onsets of sequential cluster patterns appear to be synchronized with the onset of anticipatory licking.

b. Examples of correlations between anticipatory licking onsets and rewarded trials cluster onsets in 3 datasets from hungry mice (top) and 3 datasets from thirsty mice (bottom). Note relatively high positive correlations in all examples.

c. Left: Distribution of all correlations across all hungry and thirsty datasets (N=10 and N=11, respectively). Right: Scatter plot describing correlation levels as a function of significance levels. All datasets exhibit significant positive correlations between anticipatory licking onset and the rewarded trial sequential cluster pattern onset, as indicated by the dashed vertical line at P=0.05 (P < 0.04, for all datasets, Pearson's product-moment correlation).

d. Example visualization of reward onset, anticipatory licking onset, and activity cluster onset. Left: Cluster dynamics of all rewarded trials. Right: Onsets of anticipatory licking, reward delivery, and cluster dynamics per trial. Note that the onset of the sequential cluster pattern consistently follows the onset of anticipatory licking and appears unrelated to the onset of reward delivery.

e. Distribution of the time difference between rewarded trial cluster onset and anticipatory licking onset. Across all trials, cluster onset occurs approximately 1 second *after* lick onset.

f. Distribution of trial similarity between incorrectly responded aversive and neutral trials ("false alarms") and rewarded trials (purple), compared to the similarity between correctly non-responded aversive and neutral trials ("correct rejections") to rewarded trials (red). Notably, false alarm trials are more similar to rewarded trials compared to correct rejection trials. (Data from N=14 datasets from N=5 thirsty mice).

g. Two examples of cluster sequence trial dynamics during false alarm trials and rewarded trials. False alarm trials were sorted based on their *licking* pattern similarity to the *licking* patterns in rewarded trials. Note that as the licking pattern becomes more similar, false alarm cluster sequence trials become more similar to rewarded trials.

h. Distribution of all correlations between licking similarity and trial similarity in false alarm trials and rewarded trials, as a function of significance levels. Out of 11 datasets, 9 had positive correlations, and 7 of these correlations were statistically significant ($P < 0.05$, assessed using Pearson's product-moment correlation).

i. Example cluster sequence dynamics and licking behavior in two mice during reward consumption while engaged in the operant task (rewarded trials) and during self-paced free consumption behavior. The dynamics were aligned based on the onset of licking. The neuronal cluster dynamics during operant goal-directed behavior appear distinct from those observed when the mice are freely consuming rewards.

j. Left: Distribution of all pairwise comparisons between reward consumption trials during task engagement and free consumption behavior (pooled from N=5 datasets from N=5 mice). Freely consuming mice exhibit distinct cluster dynamic, reflected by the distribution of task versus free behavior trial similarity centered around zero. Right: Comparison of the average trial similarity for each dataset individually when comparing task dynamics to itself and task dynamics versus freely behaving consumption dynamics. Free consumption dynamics are significantly different than task consumption dynamics ($P < 0.009$, one-tailed paired t-test)

Supplemental Figure 1

Supp. Figure S1 – Dimensionality estimation and reduction

a. Dimensionality assessment using a two-NN Estimator[42] on two simulated example datasets. Left: Estimation of a simulated one-dimensional dataset. Right: Estimation of a simulated two-dimensional dataset. In both datasets dimensionality is accurately approximated, with data points aligning parallel to the simulated dimension (i.e., slope).

b. Datasets initially undergo an intermediate dimensionality reduction, followed by an estimation of dimensionality for the intermediate dataset. The final dataset is achieved after an additional iteration of dimensionality reduction, using the mean estimated dimension in the intermediate iteration across all datasets.

c. Estimation of intrinsic dimensionality in shuffled and original data across different intermediate dimensionality reductions and the full embedded data with no intermediate dimensionality reduction. Estimation of the intrinsic dimensionality in the original data converges to ~6 (dashed line, chosen for subsequent dimensionality reduction, see Supp. Figure 2). In contrast, estimation on shuffled data yields an intrinsic dimension close to the intermediate embedding dimension.

d. Left: Estimated dimensionality for 14 different datasets of thirsty mice. The near-constant slope observed across all datasets suggests a consistent intrinsic dimensionality of 6, which suggests the presence of approximately six latent variables within the activity manifold. Right: Same as in `left` for 10 different datasets of hungry mice.

e. The estimated dimension was not correlated with the number of neurons sampled in each dataset for real data and shuffled data. Left and Right are the same as in `d` (P > 0.2, Pearson's product-moment correlation).

f. All dimension pairs in two different mice. Mouse 2 is presented with two different non-linear dimensionality reduction algorithms (right: Isomap; left: Laplacian eigenmaps, used throughout our analyses).

Supplemental Figure 2

1131

Supp. Figure S2 - Assessing similarity between datasets using topological data analysis

1132
1133

a. Illustration of the procedure used to assess the similarity between pairs of datasets. Initially, timepoints are down-sampled through K-means clustering. Subsequently, the distribution of topological features is extracted independently for each dataset, The magnitude of divergence in the feature distributions directly reflects the dissimilarity between the datasets, providing a quantitative measure of their dissimilarity.

b. Description the process for conducting permutation tests to assess the statistical significance of dissimilarity between datasets. Datasets are combined and permuted and redivided 200 times. During each permutation, the divergence of topological features between the randomly permuted and divided datasets is calculated. Subsequently, a p-value is derived by comparing the divergence observed between the original datasets to the distribution of divergences constructed from the permutations. This allows for the assessment of the statistical significance of dataset dissimilarity.

c. P-values calculated based on the shuffle distribution for all pairwise comparisons assessing the topological divergence of $\beta 0$ features.

1146
1147

d. Same as in `c` but for $\beta 1$ features. Shuffle distributions compared to neuronal data had non-significant different results. This observation may be attributed to the sparse nature of the $\beta 1$ features extracted from each dataset.

1148
1149
1150

e. Same as in `c` but also for InsCtx and visual cortex in hungry mice.

1151

c-e. N=100 pairwise comparisons for hungry vs hungry, N=182 pairwise comparisons for thirsty vs thirsty, N=140 pairwise comparisons for thirsty vs hungry mice, N=70 comparisons for Visual vs InsCtx mice, from N=14 datasets of thirsty mice, N=10 datasets of hungry mice, N=7 datasets of visual mice.

1152
1153
1154
1155

1156

Supplemental Figure 3

a

Reward trial (Response) — Reward trial (No response) — Neutral trial (Response) — Neutral trial (No response) — Pupil size (ITI) — Cumulative consumed rwards (ITI)

b

Clusters: 1; Radius: 8%

Clusters: 18; Radius: 17%

Clusters: 4; Radius: 52%

Clusters: 3; Radius: 88%

Clusters: 7; Radius: 61%

c

d log2(MI) = -3.4812

e Cluster × Reward (R) joint probability space

Supp. Figure S3 – Automated clustering of activity manifold; clustered structure relationship with experimental variables

a. Depicting the activity manifold annotated with external/internal variables in an exemple dataset. In each row, a distinct plane of the activity manifold is presented. To illustrate discrete variables (columns 1-4; responded reward trials to non-responded neutral trials) we use red points to indicate the event occurrence. In the case of continuous variables (columns 5-6; pupil size and cumulative consumed rewards) points are color-coded, ranging from blue to yellow according to the variable's values.

b. Illustration of the automated clustering process. For every dataset, we construct a configuration matrix, as depicted. This matrix is generated by performing K-means clustering while iteratively adding points into the central point cloud (rows) and testing with varying cluster numbers (columns). The Mean Squared Error (MSE) is calculated for each K-means configuration during each repetition and stored in the corresponding matrix cell. Several examples of different configurations are outlined with dashed gray lines, whereas the final selected configuration is delineated with a dashed red line (see 'c' for details).

c. Identification of the optimal configuration. The optimal number of points to include in the central point cloud and the optimal cluster count are automatically determined by identifying the deflection point, as detailed in the methods section.

d. Example planes of the clustered structure alongside an external variable of interest

e. Illustration of calculation of Mutual Information. The Mutual Information between the clustered structure and the external variable is obtained by calculating their joint probability space.

Supplemental Figure 4

Supp. Figure S4 – Comparison of linear projection dynamics with activity manifold dynamics

a. Activity manifold from three example mice. Top: Neuronal clusters annotated using automated clustering, with labeled clusters representing ITI water satiety and reward clusters used in 'b'.

b. Left: Physiological state approximated by cumulative consumed water rewards over time (mean across all datasets). Dashed lines indicate the experiment time at which mice reached a certain quenched level (e.g., by the middle of the experiment, mice consumed 70% of the rewards they would consume). Right: neuronal cluster occupancy (percent of time that activity occupied a certain cluster) during reward epochs and quenched phases in relation to gradual changes in physiological state (corresponding to dashed lines in the left cumulative rewards plot). Occupancy of reward clusters remained relatively consistent throughout gradually changing physiological states, and slightly decreased as the number of responded reward trials decreased with water satiation. In contrast, the two most frequently occupied neuronal clusters during the quenched phase were primarily visited when animals ceased water consumption and became quenched, but not before.

c. Top, middle: Timepoints of ongoing activity used to calculate average activity vectors for thirst (blue) and quenched (red) states, indicated by black centroids (following our previously published procedures[33]). Bottom: Trajectory of ten reward trials projected on the activity manifold.

d. The same ten reward trials as in 'c', projected linearly using a neural mode obtained from subtracting the average ongoing activity in thirst and quenched states[33]. During reward trials, population activity briefly shifts from a thirst state towards a quenched state. The trajectory of the same trials can also be observed with respect to the activity manifold described in 'c'.

Supplemental Figure 5

Supp. Figure S5 – Activity dynamics of manifold cluster sequences with two different dimensionality reduction methods

a. All dimension pairs in two different mice annotated by the automated clustering method. Mouse 2 is presented with two different non-linear dimensionality reduction algorithms (Right: Isomap; Left: : Laplacian eigenmaps, used throughout our analyses).

b. Activity dynamics of the same examples as in `a` for reward cue trials. Notice that the stereotyped activity patterns are also observed when using a different non-linear dimensionality reduction algorithm (Isomap).

Supplemental Figure 6

Supp. Figure S6 – Analysis of across dataset decoder performance.

Left: Assessing decoder F1-scores for trial types. A comparison of F1-scores when employing a decoder trained on manifold cluster sequences and another on average activity levels. Notably, only rewarded trials exhibited significantly higher decoding performance above chance. Additionally, decoding using manifold clusters outperformed decoding based on average activity levels. Right: The distribution of P-values for decoding derived from the surrogate shuffle distribution for each pairwise decoding is shown. All trial types exhibited uniform p-values in terms of F1-score significance levels, except for rewarded trials indicating significant decoding.

b. Same as in `a` but now comparing the performance of a decoder trained on manifold cluster sequences versus one trained on the projection of trials onto thirsty-quenched axes (T-Q axis).

c. Same as in `a`, but now comparing between a decoder trained on InsCtx thirsty or visual cortex hungry datasets, and tested on InsCtx hungry datasets accordingly.

d. Comparison of the F1-score for rewarded trials across natural and artificial motivational states.

a-d. N=14 thirsty datasets, N=10 hungry datasets, N=7 visual datasets, N=4 AgRP datasets, N=4 SFO datasets, N=3 hyperosmotic datasets. (Significance levels: * P < 0.05, ** P < 0.01, *** P < 0.001, one-tailed Wilcoxon signed-rank test).

Supplemental Figure 7

1246

Supp. Figure S7 - Comparison of activity dynamics in goal-directed behavioral task and free consumption of rewards

a. Examples of the pseudo-trial construction process to compare reward consumption behavior in two mice. Left: In the visual discrimination task, trials were generated by analyzing the 2 seconds before and the 4 seconds following lick onset in rewarded trials. Right: In free reward consumption, trials were generated by analyzing the 2 seconds before and 4 seconds after self-initiated lick bouts (timepoints in which licking rate sharply increased after a sufficient decrease). Only non-overlapping trials were chosen.

b. Example of activity dynamics during goal-directed behavior (operant visual discrimination task) and free reward consumption (self-initiated) in four different mice. Heatmaps are aligned with the constructed lick trials as described above. Left: Sequence of cluster dynamics observed in constructed lick trials. Middle: Licking in the constructed lick trials. Right: Overall mean activity levels. While goal-directed and self-initiated trials exhibit distinct dynamics, there is an increase in overall activity levels during both goal-directed and self-initiated consumption of rewards.

c. Example cluster sequence dynamics and licking behavior, while controlling for variations in licking behavior between task-based consumption and freely initiated consumption. We addressed these behavioral distinctions by specifically choosing free-consumption trials in which the licking rate within the initial two seconds of the licking onset fell within the mean and standard deviation of the licking rate within the first two seconds of task-based lick bouts.

d. Distribution of all pairwise trial dynamics comparisons between reward consumption trials during task engagement and free consumption behavior, when controlling for behavioral differences in licking (pooled from N=5 datasets from N=5 mice).

e. Comparison of the average trial similarity for each dataset individually when comparing task dynamics to task dynamics, and task dynamics versus freely behaving consumption dynamics, while controlling for behavioral differences in consumption (P < 0.05, one-tailed paired t-test)

1275