Antibacterial activity of human natural killer cells in the absence of accessory cells

against extracellular *Staphylococcus aureus* and hypervirulent *Pseudomonas aeruginosa*

bacteria

Audrey Guilbaud[§], Baptiste Lottin[§], Gwenann Cadiou, Tiffany Beauvais, Sylvia Lambot,

Barbara Mouratou, Nathalie Labarrière, François Davodeau, Frédéric Pecorari*

Nantes Université, Université d'Angers, Inserm, Immunology and New Concepts in

ImmunoTherapy, INCIT, UMR 1302, Nantes, France

[§] Audrey Guilbaud and Baptiste Lottin both contributed equally to this work.

* To whom correspondence should be addressed.

Dr. Frédéric Pecorari, Tel.: +33 2 40 41 28 51; E-mail: frederic.pecorari@univ-nantes.fr

<u>Running title:</u> Antibacterial activity of human natural killer cells against *S. aureus* and

hypervirulent *P. aeruginosa* bacteria

ABSTRACT

Natural killer (NK) cells play a crucial role in the innate immune response to bacterial

infections, including those due to *Pseudomonas aeruginosa* (*P. aeruginosa*) and

Staphylococcus aureus (*S. aureus*). *In vivo*, it has been shown that NK cells are activated by

innate accessory cells that detect the presence of bacteria and activate NK cells via a cytokine

network. *In vitro*, several studies have shown that NK cells can also be activated without the

help of accessory cells by direct contact with some bacteria species such as extracellular *P.*

aeruginosa. Whether this phenomenon of direct activation is restricted to certain bacterial

species, or whether it can be generalized, is still debated, as for example in the case of NK cell

activation by *S. aureus*, which seems to require the intervention of accessory immune cells.

Here, we show with co-incubation experiments between NK cells and two bacterial species,

that, in the absence of accessory cells, NK cells are able to impede bacterial growth. This has

been demonstrated for the *P. aeruginosa* PA14 strain, which is hypervirulent and known for its

deleterious effects on NK cells, as well as for the *S. aureus* Newman strain. The monitoring of

CD107a by flow cytometry suggests that NK cells degranulate after contact with *S. aureus*

bacteria. Our study contributes to the idea that NK cells can be activated in the absence of any

accessory cells by various species of bacteria, even an hypervirulent one, and therefore that

NKs can directly have an antibacterial effect. This important insight may pave the way for new

therapeutic approaches using antibacterial NK-cell engagers.

Keywords: NK cells, bacteria, Pseudomonas aeruginosa, Staphylococcus aureus

INTRODUCTION

NK cells are a type of lymphocyte that plays a crucial role in the innate immune system. They are mostly known for their ability to target and eliminate virus-infected cells and cancer cells. However, the role of NK cells in bacterial infection has been described historically since the 1980s, with their ability to lyse cells infected with *Shigella flexneri*, *Legionella pneumophila* or *Mycobacterium avium* [1–3], and more recently for intracellular *Listeria monocytogenes* [4]. Although the crucial role of NK cells in fighting bacterial infections is quite well documented, some aspects of their activation by bacteria are still partly described. A mechanism of NK activation which has been the most studied, is based on recognition of pathogen-associated molecular patterns (PAMPs) by pathogen-recognition receptors (PRRs) at the surface of accessory cells, such as macrophages, neutrophiles, dendritic cells, and B lymphocytes that detect the presence of bacteria. These activated accessory cells then secrete cytokines, notably IL-15 and IL-18, that play an important role in the development, proliferation, survival and activation of NK cells [5]. NK cell activation is characterized by the expression of CD69 and interferon gamma (IFN-γ) and leads to the polarized release of granules with a cytotoxic lytic effect, thanks to perforin, granulysin and the granzymes they contain. These effectors are able to induce significant physical damage to bacterial walls, leading to their death [6].

About two decades ago, the possibility of NK cells being activated by another mechanism *via* direct contact with bacteria has been questioned [7]. Although increasing evidence show that NK cells can have antibacterial activity independently of accessory cells, this has long remained open to doubt as it contradicts the paradigm that NK cells play their role under the control of a network of signals relayed by accessory cells [8]. However, several studies have reported that NKs can be activated by contact, for instance with mycobateria, *Nocardia farcinica*, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae*, *Escherichia coli*, *Salmonella typhimurium*, *Staphylococcus epidermidis* and *Burkholderia cepacia* [7, 9–13], thereby

inducing an antibacterial effect. Thus, even in the absence of accessory cells, a broad spectrum

of bacteria can activate NKs, including coccus, bacilli, Gram-negative and Gram-positive

bacteria. When studied at the molecular level, bacterial recognition can involve a natural

cytotoxicity receptor (NCR) located on the NK surface [11].

This broad potential of NKs to kill a variety of bacteria could provide the basis for the

development of innovative therapeutic approaches, irrespective of the antibiotic resistance

status of the targeted pathogens. Despite the introduction of more than 30 new antibiotics since

2000, the supply chains that generate new drugs are shrinking for all pathogens [14]. The

emergence of antibiotic-resistant strains is far from being under control, according to the world

health organization (WHO), and cases are multiplying in many countries of diseases for which

there are no longer any non-invalidating treatments. In 2008, *Enterococcus faecium*, *S. aureus*,

Klebsiella pneumoniae, *Acinetobacter baumannii*, *P. aeruginosa* and *Enterobacter* species

were grouped under the ESKAPE acronym as pathogens known to escape the lethal action of

antibiotics [15]. In 2017, the WHO has published the priority pathogen list (PPL) for R&D of

new antibiotics that includes most of those from the ESKAPE group, plus other species such as

Helicobacter pylori and *Campylobacter*[16]. For most of these pathogens, multidrug resistant

clinical isolates are found, and they are the leading cause of nosocomial infections.

Multidrug-resistant (MDR), extensively drug-resistant (XDR) and Pan-resistant (PDR)

strains, including *P. aeruginosa*, have been defined according to their non-susceptibility profile

to various categories of antibiotics, the PDR strains being defined as non-susceptible to all

agents in all antimicrobial categories [17, 18]. *P. aeruginosa* is a Gram-negative bacillus and

an opportunistic pathogen often involved in nosocomial infections, most notably in patients

with cystic fibrosis or those in intensive care units [17]. Upon infection of mice by *P.*

aeruginosa, NKG2D expression is increased and NK cells participate to the clearance of

bacteria [19, 20]. Furthermore, the depletion of NK cells leads to increased mortality rate of

infected mice [21]. *In vitro*, human NK cells have been shown to bind directly to *P. aeruginosa*

via their NKp44 receptor [11]. It was also demonstrated that human NK cells could exert a

microbicide effect via granzymes on extracellular bacteria in the absence of any accessory cells

[13].

S. aureus is also an opportunistic pathogen responsible for a large number of hospital-

acquired infections and community-acquired infections. This Gram-positive coccus can cause

numerous diseases such as skin infections, sepsis, pneumonia, infective endocarditis and

bloodstream infections [22]. During the last decades, the spreading of MDR and XDR strains

have been observed worldwide. Methicillin-resistant *S. aureus* (MRSA) strains are of concern

as they are responsible for infections more difficult to treat, requiring the use of non-beta-lactam

antibiotics, such as vancomycin or daptomycin. However, the increased use of vancomycin as

a treatment of last resort has led to the emergence of vancomycin-resistant strains of *S. aureus*

(VRSA), which poses a very serious threat to human health. The crucial role of NK cells in the

immune response during infection by *S. aureus* was demonstrated with NK-depleted mouse

models and IL-15 knockout mice [23]. The absence of NK cells makes mice susceptible to *S.*

aureus infection, while IL-15 has been shown to have an indispensable effect on NK cells that

cannot be compensated for by other immune cells. *In vitro*, it was reported that NK cells in

presence of *S. aureus* bacteria require a co-stimulation by monocytes to be activated as assessed

by CD69 and IFN-γ levels. A direct contact between *S. aureus* and NK cells was inefficient to

activate NKs and no bacterial viability tests were performed [24]. Surprisingly, a previous study

reported for another *Staphylococcus*, *Staphylococcus epidermidis* (*S. epidermidis*), that NK

cells were sufficient to show a strong antibacterial effect *in vitro* in a contact-dependent manner

according to counting of survival bacteria [12].

Thus, NK cells play a critical role in showing an antibacterial effect, but it seems that

depending on the bacterial species, co-stimulation by accessory cells is either mandatory or not.

The main objective of this work was to determine to which extent NK cells, in the absence

of accessory cells, are able to interfere with pathogenic bacteria growth, whether bacteria are

Gram- bacilli or Gram+ coccus. To this end, in absence of accessory cells, we co-incubated NK

cells with the hypervirulent *P. aeruginosa* PA14 strain and the *S. aureus* Newman strain, both

belonging to the ESKAPE group. The antibacterial activity of NKs was studied at different

NK/bacteria ratios and as a function of incubation time, by determining bacterial survival by

counting colony forming units on petri dishes and by studying NK degranulation using flow

cytometry.

MATERIALS AND METHODS

Bacteria strains

The *S. aureus* Newman strain and the *P. aeruginosa* PA14 strain were kindly provided

by Prof. Timothy Foster (Dublin, Ireland) and by Dr. Alexis Broquet (Nantes, France),

respectively.

NK cell line

The human NK92-CD16 cell line was provided by Dr. Henrie Vié [25] and was cultured

in RPMI 1640 medium with 10% FBS (Eurobio), 2 mM GlutaMAX-I (Gibco), 100 U/mL

penicillin (Gibco) and 0.1 mg/mL streptomycin (Gibco).

Sorting and amplification of NK

PBMC were isolated from heparinized blood from healthy adult volunteers by gradient

centrifugation on Ficoll (Lymphocytes separation medium, Eurobio). All blood donors were

recruited at the Blood Transfusion Center (EFS, France) and informed consent was obtained

from all individuals. NK cells were collected by magnetic cell sorting using the EasySep Human

NK Cell Isolation Kit (Stem Cell, France) according to the manufacturer's instructions.

The purity of each NK population was checked before and after cell sorting and routinely

by flow cytometry with anti-human CD16-BV421 (clone 3G8, BD Pharmagen, France), anti-

human CD56-PeCy7 (clone B159, BD Pharmagen, France), anti-human CD3-FITC (clone

BW264/56, Miltenyi Biotech, France).

After cell sorting, NK cells were expanded on allogeneic irradiated feeder cells

(peripheral blood mononuclear cells and B-EBV B cells) with 150 U/mL human rhIL-2. Briefly,

2000 NK cells/well were distributed in 96-well round bottom plates with 200 µl of RPMI culture

medium containing 8% human serum (Eurobio, France), 2 mM GlutaMAX (Gibco, France),

100 U/mL penicillin (Gibco, France), 0.1 mg/mL streptomycin (Gibco, France), human rIL-2

150 U/ml (Proleukin, Novartis) and irradiated EBV-B feeder cells (1.10^3/well), and allogeneic

PBMC from two donors (1.10^4/well). The culture medium was changed every 2 days. NK cells

were split when their number exceeded $1,5.10^5$/well.

Killing Assay

An isolated colony of *P. aeruginosa* or *S. aureus* was picked from a Luria-Bertani

(LB)/agar petri dish to inoculate 10 mL of LB medium that were incubated 16 h at 37°C with

shaking. Then 1 mL of the culture was used to inoculate 10 mL of LB medium and the culture

was incubated for 2 h at 37°C with shaking. The culture was diluted in RPMI medium to

distribute 5.10^2/well bacteria per well in 96 well round bottom plates. Two days before the

killing assay, NK cells were cultured in RPMI 1640 medium with 10% FBS (Eurobio), 2 mM

GlutaMAX-I (Gibco) without antibiotics. The day of the killing assay, NK cells were washed

3 times in serum and antibiotic free RPMI 1640 media to discard any traces of antibiotics.

Bacteria were co-cultured with or without NK cells (NK92hCD16 or amplified NK - from

$2,5.10^2$/well to 5.10^4/well) in quadruplicate in serum and antibiotic free RPMI 1640 media.

Cultures were incubated at 37°C and 5% CO2 for 2 to 6 h. Co-cultures were diluted in sterile

water and 10 µL were spotted on LB agar plate. Plates were incubated overnight at 37°C and

the colonies were counted and recorded.

Degranulation Assay

NK92hCD16 or amplified NK (2.10^5/well) were stimulated for 4 hours with or without

S. aureus Newman (from 2.10^6 to 2.10^4/well) in the presence of anti-human CD107a-AF647

mAb (clone H4B4, Biolegend, France) added at the beginning of stimulation in serum and

antibiotic free RPMI media in 96 well round bottom plates. Cultures were incubated at 37°C

and 5% CO2 for 4 h. To detect NK, cells were incubated with anti-human CD56 PE

(clone B159, BD Pharmagen, France). Cells were next fixed with PBS 4% paraformaldehyde

for 15 minutes. Stained cells were acquired on a BD FACSCanto II flow cytometer using DIVA

software.

Statistics

Statistical analyses were performed using GraphPad Prism 8 software. One-way ANOVA

with Bonferroni correction was used to evaluate statistical significance, a p-value of less than

0.05 was considered statistically significant (*), $p < 0.01$ (**), $p < 0.001$ (***) and $p < 0.0001$

(****) as highly significant. For non-significant results, (ns) was indicated.

RESULTS

NK cells from healthy donors impede the growth of an hypervirulent strain of *P. aeruginosa*

We first studied the efficacy of NKs against a *P. aeruginosa* strain different from the PAO1 strain studied in the work of Feehan et al [13]. We chose the PA14 strain with the aim to challenge antibacterial efficacity of NK cells, this strain being described as hypervirulent compared to PAO1 [26], and also because PA14 was reported to mediate NK cell apoptosis *in vitro* [27].

NK cells were isolated from healthy donors as described in "Materials and methods" section. Flow cytometry measurements (figure 1) showed that these NK cell populations were highly pure, around 93%.

We then set up co-incubations with a constant number of bacteria per condition (500 CFU) and an increasing number of human NK cells to study the effect of effector:target (E:T) ratio on *P. aeruginosa*. The figure 2A shows that after 6h of co-incubation, PA14 CFU numbers were reduced as the E:T ratio increased compared to the condition without NK cells. The effect on bacterial growth was already optimal at 4 h (figure 2B).

These results confirm that human NK cells impair *P. aeruginosa* growth *in vitro*, even if the bacterial strain is hypervirulent and known to induce NK apoptosis under certain conditions.

NK cells from healthy donors have an activity against *S. aureus* bacteria

We then investigated whether this effect could be observed for a species quite different from *P. aeruginosa* also belonging to the ESKAPE group, i.e. *S. aureus*. Similar co-incubation experiments were performed to study the effect of NK cells on *S. aureus* growth for increasing E:T ratios and co-incubation time (figure 3). Remarkably, an E:T ratio of 0.5 is already

sufficient to have a significant effect on *S. aureus* growth, reflecting a significantly higher

efficacy than against PA14 (figure 2). The time for an optimal effect was observed after 6h of

co-incubation.

NK92 cells also inhibit the growth of *S. aureus* bacteria

Having shown that NK cells from healthy donors impaired the growth of *S. aureus* growth, we

sought to determine if the NK from a cell line (NK92-CD16) could also kill these bacteria.

Indeed, the purity of NK cells from donors, although high (figure 1), is not 100% and the

antibacterial activities we observed could in fact be the result of contaminating accessory cells.

As shown in figure 4, NK92-CD16 cells killed *S. aureus* in a similar way (E:T ratio and time),

demonstrating this effect was solely induced by NK cells, *i.e.* without the help of eventual

contaminant accessory cells.

Assessment of NK cell degranulation

These results led us to study whether this effect on bacteria growth is mediated by degranulation

of NK cells. To address this point, we measured the ability of NK cells to degranulate upon

contact with bacteria, through CD107a expression analyzed by flow cytometry. As shown on

figure 5, for E:T ratio up to 100:1, *P. aeruginosa* bacteria did not induce expression of the

degranulation marker CD107a. In contrast, *S. aureus* bacteria were able to induce the expression

of CD107a even for an E:T ratio as low as 0.1:1 (figure 5). At an E:T = 1:1, up to ~8.3 % of

NK cells were stimulated by *S. aureus* bacteria for the expression of CD107a, reflecting an

important level of degranulation.

DISCUSSION

While NK cells are traditionally associated with viral and tumour responses, emerging evidences highlight their importance in combating bacterial infections. Thus, it was commonly accepted that NKs can inactivate bacteria, but mainly indirectly and with the help of accessory cells. However, several studies demonstrated that NK cells have a direct anti-microbial activity against extracellular *Salmonella typhi, Escherichia coli, Staphylococcus epidermidis, Burkholderia cepacia*, *Mycobacterium kansasii* and *Mycobacterium tuberculosis* bacteria, and [9–12]. In the most recent study, the group of Mody et al. reported this direct effect for two clinical isolates of *P. aeruginosa* and for the lab-strain PAO1 [13]. However, the clinical isolate, CF5, showed a different behaviour, since it was able to multiply better in the presence of NK cells than in their absence. Thus, we wondered how general these observations about a direct effect of NK cells on bacteria could be.

To this end, we first chose to test *P. aeruginosa* strain PA14. PAO1 and PA14 are two strains that were originally isolated from clinical samples and are now widely used in the laboratory to study *P. aeruginosa*. Interestingly, the PAO1 strain displays a moderate level of virulence, while PA14 is described as an hypervirulent strain [26]. This difference is due to several factors including the existence of two pathogenicity islands which encode for effectors such as ExoU via the type III secretion system (T3SS), a potent cytotoxin that induces cell damage and death. This strain also produces a higher level of pyocyanin than PAO1 [28], a toxin that generates reactive oxygen species (ROS) within host cells, leading to oxidative stress and damage to cellular components [29]. It has been reported that pyocyanin attracts neutrophils and induces their apoptosis [30] and that PA14 strain also induces death of NK cells, while a pyocyanin-deficient PA14 strain does not show any cytotoxic effect against NK cells [27]. However, *P. aeruginosa* K (PAK) has a cytotoxic effect on NK cells although it produces pyocyanin at low level, suggesting that other bacterial compounds are involved [27].

Interestingly, our co-incubation experiments show that NK cells from four donors are able to

interfere with PA14 cultures (figure 2). However, large E:T ratios are necessary to decrease

significantly (at least E:T = 200), or to prevent growth (E:T = 400) of bacteria after 6 h of co-

incubation. This effect is notably more visible over time of co-incubation and is maximal in our

setup at 4 hours. These results are very similar to those observed by the group of Mody et al for

PAO1 strain and two other clinical strains (S35004 and CF127) [13]. It should be noted that

after 24 h of co-culture, an effect of NKs against mycobacteria was already observed for an E:T

= 20:1, but a higher ratio (E:T = 100:1) was also necessary to kill them in proportions similar

to those for *P. aeruginosa* [9]. While to our knowledge the levels of pyocyanin productions for

S35004 and CF127 strains are not described in the literature. Given our results for PA14 and

those for PAO1 [13], it seems that the pyocyanin factor is not sufficient to explain why some

strains of *P. aeruginosa* overcome the action of NK cells. The study of Chung et al. [27] aimed

to understand the cytotoxicity mechanisms mediated by *P. aeruginosa*, thus they carried-out

co-incubations experiments with E:T ratios of 0.1:1 and 1:1. The quite high E:T ratios used in

our study and in that of the study of Mody's group necessary to show an antibacterial effect are

probably required to overcome the cytotoxic effect of *P. aeruginosa* strains against NK cells

which is not due to factors specific to PA14 as observed by Chung *et al.* Interestingly, it has

been shown that a contact between NK cells and yeast for at least 90 minutes is necessary to

observe NK cell-induced cytotoxicity [31]. Unlike *S. aureus*, *P. aeruginosa* is mobile which

could impede prolonged contact and explain why higher E:T ratios are required for *P.*

aeruginosa. It will be interesting to further study the mechanisms used by *P. aeruginosa* strains

to overcome the NK cell action in spite of quite high E:T ratios.

P. aeruginosa being a gram-negative bacillus, we then wanted to investigate whether

this effect of NK cells on extracellular bacteria could be observed for a quite different species

that belongs to the ESKAPE group. Thus, we choose *S. aureus* as it is a Gram-positive coccus.

As shown in figure 3, a similar effect to that seen against *P. aeruginosa* could be observed for

S. aureus, regardless of the donor from which the NK cells originate. However, an E:T ratio as

low as 0.5 was already sufficient to get a 50% decrease in viable bacteria, while an E:T = 200:1

was necessary for PA14 strain to gain the same effect. This is also significantly lower than the

minimal efficient E:T ratio of 12.5:1 reported for *Burkholderia cepacia* [10]. Our results

demonstrate the powerful antibacterial activity of NK cells against *S. aureus*. Interestingly,

lower but efficient ratio (E:T < 0.01) have been reported for purified NK from healthy donors

against *Escherichia coli* [12]. Thus, it will be interesting to explore lower E:T ratios than those

we have tested for *S. aureus*. As for *P. aeruginosa*, we found that this effect against *S. aureus*

was maximal after 6 h of co-incubation. A previous work demonstrated that NK cells could be

activated by *S. aureus* as shown by IFN-γ assays, but required a cell contact-dependent co-

stimulation by monocytes [24]. FACS measurements on our NK cells preparations from donors

excluded a contamination with monocytes. Furthermore, our results with NK92 cells exclude

that NK cells need the help of any accessory cells to induce antibacterial activity (figure 4).

Previous studies have shown that the mechanism underlying the antibacterial activity of

NK cells involves their degranulation, and the role of some of their granule components, such

as perforin, granulysin and granzyme, have been demonstrated [9, 10, 13]. Surprisingly, our

results show that NK cells degranulation behaviour differs depending on the bacteria used as

targets. For *P. aeruginosa*, no degranulation was observed (figure 5A), while for *S. aureus*

about up to 8.3 % of NK cells were degranulated in presence of bacteria (figure 5). The work

of Feehan *et al.* demonstrated the role of both granzyme B and H contained in lytic granules for

the killing of *P. aeruginosa*, by using inhibitor compounds and CRISPR Cas9-induced

knockouts [13]. Considering that only quite high E:T ratios are efficient to induce an

antibacterial effect, and that *P. aeruginosa* has been reported to induce a decrease of NKG2D

expression thereby reducing NK cell cytotoxicity [32], it is possible that degranulation occurs

at a such low level that it could be difficult to assess it with CD107a.

Overall, we confirm that NK cells without the help of accessory cells are able to mediate

an antibacterial activity against *P. aeruginosa*, even though it is a hypervirulent strain, and we

found that NK cells are also active against *S. aureus*, but in a much more efficient way. Future

experiments should study the mechanisms by which NK cells are activated by *S. aureus* to exert

their antibacterial activity and which NK effectors are crucial to this effect.

CONCLUSIONS

This work reinforces previous studies showing NK cells can have an antibacterial role against

various species of extracellular bacteria in the absence of accessory cells. How general in term

of bacterial species sensitive to this effect will determine whether NK cells can be considered

as a basis for the development of therapeutic options as an alternative or a complement to

antibiotics. With antibiotic-resistant strains emerging worldwide, we believe that NK-based

therapies could play a role, as they already do in combating tumour cells with the development

of NK-cell engagers.

ACKNOWLEDGEMENT

We thank the core facilitiy of SFR Santé: Cytometry and cell sorting "Cytocell", for expert

technical assistance. We are grateful to Alexis Broquet (Nantes, France) for the gift of the PA14

strain. B.L. received a fellowship from Région Pays-de-la-Loire and from AID (Agence de

l'innovation de défense).

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

1.   Klimpel GR, Niesel DW, Klimpel KD (1986) Natural cytotoxic effector cell activity
against Shigella flexneri-infected HeLa cells. Journal of immunology (Baltimore, Md:
1950) 136:1081–1086

2.   Blanchard DK, Stewart 2nd WE, Klein TW, Friedman H, Djeu JY (1987) Cytolytic
activity of human peripheral blood leukocytes against Legionella pneumophila-infected
monocytes: characterization of the effector cell and augmentation by interleukin 2.
Journal of immunology (Baltimore, Md: 1950) 139:551–556

3.   Katz P, Yeager H, Whalen G, Evans M, Swartz RP, Roecklein J (1990) Natural killer
cell-mediated lysis of Mycobacterium-avium complex-infected monocytes. Journal of
clinical immunology 10:71–77

4.   Walch M, Dotiwala F, Mulik S, Thiery J, Kirchhausen T, Clayberger C, Krensky AM,
Martinvalet D, Lieberman J (2014) Cytotoxic Cells Kill Intracellular Bacteria through
Granulysin-Mediated Delivery of Granzymes. Cell 157:1309–1323

5.   Newman KC, Riley EM (2007) Whatever turns you on: accessory-cell-dependent
activation of NK cells by pathogens. Nat Rev Immunol 7:279–291

6.   Stenger S, Hanson DA, Teitelbaum R, Dewan P, Niazi KR, Froelich CJ, Ganz T, Thoma-
Uszynski S, Melian A, Bogdan C, Porcelli SA, Bloom BR, Krensky AM, Modlin RL
(1998) An antimicrobial activity of cytolytic T cells mediated by granulysin. Science
282:121–5

7.   Chalifour A, Jeannin P, Gauchat JF, Blaecke A, Malissard M, N'Guyen T, Thieblemont
N, Delneste Y (2004) Direct bacterial protein PAMP recognition by human NK cells
involves TLRs and triggers alpha-defensin production. Blood 104:1778–83

8.   Horowitz A, Stegmann K, Riley E (2012) Activation of Natural Killer cells during
microbial infections. Frontiers in Immunology 2:

9.   Lu CC, Wu TS, Hsu YJ, Chang CJ, Lin CS, Chia JH, Wu TL, Huang TT, Martel J, Ojcius
DM, Young JD, Lai HC (2014) NK cells kill mycobacteria directly by releasing perforin
and granulysin. J Leukocyte Biol 96:1119–29

10. Li SS, Saleh M, Xiang RF, Ogbomo H, Stack D, Huston SH, Mody CH (2019) Natural
killer cells kill Burkholderia cepacia complex via a contact-dependent and cytolytic
mechanism. Int Immunol 31:385–396

11. Esin S, Batoni G, Counoupas C, Stringaro A, Brancatisano FL, Colone M, Maisetta G,
Florio W, Arancia G, Campa M (2008) Direct binding of human NK cell natural
cytotoxicity receptor NKp44 to the surfaces of mycobacteria and other bacteria. Infect
Immun 76:1719–27

12. Garcia-Penarrubia P, Koster FT, Kelley RO, McDowell TD, Bankhurst AD (1989)
Antibacterial activity of human natural killer cells. J Exp Med 169:99–113

13. Feehan DD, Jamil K, Polyak MJ, Ogbomo H, Hasell M, Li SS, Xiang RF, Parkins M,
Trapani JA, Harrison JJ, Mody CH (2022) Natural killer cells kill extracellular
Pseudomonas aeruginosa using contact-dependent release of granzymes B and H. PLoS
Path 18:e1010325

14. Butler MS, Blaskovich MA, Cooper MA (2017) Antibiotics in the clinical pipeline at the
end of 2015. J Antibiot (Tokyo) 70:3–24

15. Rice LB (2008) Federal Funding for the Study of Antimicrobial Resistance in Nosocomial
Pathogens: No ESKAPE. The Journal of Infectious Diseases 197:1079–1081

16. WHO publishes list of bacteria for which new antibiotics are urgently needed.
https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-
new-antibiotics-are-urgently-needed. Accessed 18 Oct 2023

17. Magiorakos A-P, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth
S, Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J,
Struelens MJ, Vatopoulos A, Weber JT, Monnet DL (2012) Multidrug-resistant,
extensively drug-resistant and pandrug-resistant bacteria: an international expert proposal
for interim standard definitions for acquired resistance. Clinical Microbiology and
Infection 18:268–281

18. Kunz Coyne AJ, El Ghali A, Holger D, Rebold N, Rybak MJ (2022) Therapeutic
Strategies for Emerging Multidrug-Resistant Pseudomonas aeruginosa. Infect Dis Ther
11:661–682

19. Wesselkamper SC, Eppert BL, Motz GT, Lau GW, Hassett DJ, Borchers MT (2008)
NKG2D Is Critical for NK Cell Activation in Host Defense against Pseudomonas
aeruginosa Respiratory Infection1. The Journal of Immunology 181:5481–5489

20. Borchers MT, Harris NL, Wesselkamper SC, Zhang S, Chen Y, Young L, Lau GW
(2006) The NKG2D-Activating Receptor Mediates Pulmonary Clearance of Pseudomonas
aeruginosa. Infection and Immunity 74:2578–2586

21. Broquet A, Roquilly A, Jacqueline C, Potel G, Caillon J, Asehnoune K (2014) Depletion
of Natural Killer Cells Increases Mice Susceptibility in a Pseudomonas aeruginosa
Pneumonia Model*. Critical Care Medicine 42:e441

22. Lakhundi S, Zhang K (2018) Methicillin-Resistant Staphylococcus aureus: Molecular
Characterization, Evolution, and Epidemiology. Clin Microbiol Rev 31:

23. Small CL, McCormick S, Gill N, Kugathasan K, Santosuosso M, Donaldson N, Heinrichs
DE, Ashkar A, Xing Z (2008) NK cells play a critical protective role in host defense
against acute extracellular Staphylococcus aureus bacterial infection in the lung. J
Immunol 180:5558–68

24. Haller D, Serrant P, Granato D, Schiffrin EJ, Blum S (2002) Activation of Human NK
Cells by Staphylococci and Lactobacilli Requires Cell Contact-Dependent Costimulation
by Autologous Monocytes. Clinical and Vaccine Immunology 9:649–657

25. Clémenceau B, Valsesia-Wittmann S, Jallas A-C, Vivien R, Rousseau R, Marabelle A,
Caux C, Vié H (2015) In Vitro and In Vivo Comparison of Lymphocytes Transduced with
a Human CD16 or with a Chimeric Antigen Receptor Reveals Potential Off-Target
Interactions due to the IgG2 CH2-CH3 CAR-Spacer. J Immunol Res 2015:482089

26. Grace A, Sahu R, Owen DR, Dennis VA (2022) Pseudomonas aeruginosa reference
strains PAO1 and PA14: A genomic, phenotypic, and therapeutic review. Frontiers in
Microbiology 13:

27. Chung JW, Piao Z-H, Yoon SR, Kim MS, Jeong M, Lee SH, Min JK, Kim JW, Cho Y-H,
Kim JC, Ahn JK, Kim KE, Choi I (2009) Pseudomonas aeruginosa Eliminates Natural
Killer Cells via Phagocytosis-Induced Apoptosis. PLOS Pathogens 5:e1000561

28. Sana TG, Lomas R, Gimenez MR, Laubier A, Soscia C, Chauvet C, Conesa A, Voulhoux
R, Ize B, Bleves S (2019) Differential Modulation of Quorum Sensing Signaling through
QslA in Pseudomonas aeruginosa Strains PAO1 and PA14. Journal of Bacteriology
201:10.1128/jb.00362-19

29. Hall S, McDermott C, Anoopkumar-Dukie S, McFarland AJ, Forbes A, Perkins AV,
Davey AK, Chess-Williams R, Kiefel MJ, Arora D, Grant GD (2016) Cellular Effects of
Pyocyanin, a Secreted Virulence Factor of Pseudomonas aeruginosa. Toxins 8:236

30. Allen L, Dockrell DH, Pattery T, Lee DG, Cornelis P, Hellewell PG, Whyte MKB (2005)
Pyocyanin Production by Pseudomonas aeruginosa Induces Neutrophil Apoptosis and
Impairs Neutrophil-Mediated Host Defenses In Vivo1. The Journal of Immunology
174:3643–3649

31. Ogbomo H, Timm-McCann M, Barnes T, Xiang RF, Jamil K, Ganguly A, Stack D,
Huston SM, Li SS, Colarusso P, Mody CH (2018) Granule-Dependent NK Cell Killing of
Cryptococcus Requires Kinesin to Reposition the Cytolytic Machinery for Directed
Cytotoxicity. Cell Rep 24:3017–3032

32. Vourc'h M, David G, Gaborit B, Broquet A, Jacqueline C, Chaumette T, Caillon J,
Roquilly A, Retiere C, Asehnoune K (2020) Pseudomonas aeruginosa Infection Impairs
NKG2D-Dependent NK Cell Cytotoxicity through Regulatory T-Cell Activation.
Infection and Immunity 88:10.1128/iai.00363-20

436
437

FIGURE LEGENDS

Figure 1: Control of NK cell purity from healthy donors. NK cells derived from healthy donor

D3 were stained with FITC-conjugated anti-CD3, PECy7-conjugated anti-CD56 and BV421-

conjugated anti-CD16 to control purity.

Figure 2: NK cells have an effect on PA14 growth. A) Effect of NK cells from healthy donors

on PA14 growth (CFU/mL) after 6 h of co-incubation as a function of E:T ratio; B) For an E:T

= 200:1, study over co-culture time of NK cells on PA14 (control = PA14 alone). CFU was

determined by plate counting. These data were pooled from independent replicates using NK

cells from 4 different donors carried-out (n= 4 and n=6, for the ratio and the kinetic studies,

respectively). ** P≤0.01, **** = P≤0.0001. ns: not significant.

Figure 3: Effect of NK cells from healthy donors on *S. aureus* growth. A) Effect of different

E:T ratios on the *S. aureus* Newman strain growth (CFU/mL) after 6 h of co-incubation; B)

Study over co-culture time of the effect of NK cells on *S. aureus* CFU numbers (control = *S.*

aureus alone); E:T = 5:1). CFU was determined by plate counting. These data were obtained

from independent replicates (n= 7 for the E:T ratio study; n = 6 for the kinetic study) carried-

out on different days using NK cells from 4 different donors. * P≤0.5, **** P≤0.0001. ns: not

significant.

Figure 4: Effect of NK92-CD16 cells on *S. aureus* growth. A) Effect of different E:T ratios

on the *S. aureus* Newman strain growth (CFU/mL) after 6 h of co-incubation; B) Study over

co-culture time of the effect of NK cells on *S. aureus* CFU numbers (control = *S. aureus* alone);

E:T = 25:1). CFU were determined by plate counting. These data were obtained from

independent replicates (n= 5 for the E:T ratio study; n = 6 for the kinetic study) carried-out on

A. Guilbaud and B. Lottin *et al.*   18

different days using NK cells from 4 different donors. * P≤0.5, **** P≤0.0001. ns: not

significant.

Figure 5: Measurement of CD107a expression on NK cells by flow cytometry. A) NK cells

from healthy donors were co-incubated with PA14 *P. aeruginosa* or Newman *S. aureus* bacteria

for 6 h at different E:T ratios and were stained with AF647-conjugated anti-CD107a and with

PE-conjugated anti-CD56. The dot plots for CD107 and CD56-expressing NK cells are

presented. B) The percentages of degranulation of NK cells from four donors for different

NK:*S. aureus* ratios as a function of their potential of degranulation (CD107a+

PMA/Ionomycin) (n = 4). Degranulation controls were performed with NK cells stimulated

with PMA/Ionomycin before analyzing samples by flow cytometry. * P≤0.1, ** P≤0.01, ****

P≤0.0001.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 5