Multifaceted impact of specialized neuropeptide-intensive neurons on the selective

vulnerability in Alzheimer's disease

Manci Li[1,2], Peter A. Larsen[1,2]

Affiliations

[1] Department of Veterinary and Biomedical Sciences, University of Minnesota, St. Paul, MN

55108

[2] Minnesota Center for Prion Research and Outreach, College of Veterinary Medicine,

University of Minnesota, St. Paul MN 55108

Corresponding author:

Manci Li, BSc, Ph.D. candidate

Department of Veterinary and Biomedical Sciences, University of Minnesota

1971 Commonwealth Avenue, St. Paul, MN 55108

email: li000021@umn.edu

phone: 612-626-1694

fax: 612-625-5203

Abstract

INTRODUCTION

Widespread disruption of neuropeptide (NP) networks in Alzheimer's disease (AD) and

disproportionate absence of neurons expressing **h**igh **N**P-**p**roducing, coined as HNP neurons,

have been reported for the entorhinal cortex (EC) of AD brains. Hypothesizing that functional

features of HNP neurons are involved in the early pathogenesis of AD, we aim to understand

the molecular mechanisms underlying these observations.

METHODS

Multiscale and spatiotemporal transcriptomic analysis was used to investigate AD-afflicted and

healthy brains. Our focus encompassed NP expression dynamics in AD, **AD**-associated **NP**s

(ADNPs) trajectories with aging, and the neuroanatomical distribution of HNP neuron.

RESULTS

Findings include that 1) HNP neurons exhibited heightened metabolic needs and an

upregulation of gene expressions linked to protein misfolding; 2) dysfunctions of ADNP

production occurred in aging and mild cognitive decline; 3) HNP neurons co-expressing ADNPs

were preferentially distributed in brain regions susceptible to AD.

DISCUSSION

We identified potential mechanisms that contribute to the selective vulnerability of HNP neurons

to AD. Our results indicate that the functions of HNP neurons predispose them to oxidative

stress and protein misfolding, potentially serving as inception sites for misfolded proteins in AD.

Introduction

Alzheimer's disease (AD) is a neurodegenerative disorder characterized by progressive
cognitive decline and memory loss.[1] The neuropathological features of AD encompass both
"positive"—Aβ plaques and tau tangles, glial responses, and cerebral amyloid angiopathy—and
"negative" lesions, such as the loss of neurons and synapses.[2] The majority of AD cases occur
sporadically, with no clear understanding of their cause or pathogenesis.[1] "Epicenters" are
described as the site exhibiting the peak pathological changes or atrophy within the brain, often
considered coinciding with the initial site of disease onset by network-based
degeneration/spread hypothesis for AD.[3,4] Despite extensive research, gaps in our
understanding remain, particularly concerning the selective vulnerability of specific cell types
and brain regions to protein misfolding.[5,6] Investigating these enigmas is crucial for developing
targeted interventions and advancing our knowledge of AD's pathophysiology.[7]

We recently showed the widespread disruption of neuropeptide (NP) networks in AD and the
disproportionate absence of cells, mostly neurons, expressing higher levels and diversities of
NPs (high NP-producing/HNP cells/neurons) in the entorhinal cortex (EC) of AD brains.[8] As
steps involved in NP production are energy-intensive and HNP neurons dedicated more
transcription power to NP production, we hypothesized that HNP neurons face greater energetic
demands and higher metabolic stress during aging.[8] However, the functional characteristics of
HNP neurons and the relationship between NP co-expression and other established
pathological features of AD are unclear. Although the expression of AD-associated NPs
(ADNPs) decreased in the hippocampus during natural aging,[8] the role of HNP neurons co-
expressing ADNPs in influencing cognitive functions and mediating regional brain vulnerability is
yet to be elucidated.

In this context, we hypothesize that HNP neurons co-expressing ADNPs may represent cellular
substrates implicated in the "epicenters" of cascading protein misfolding events that are the

hallmarks of AD (Figure 1).[4–6] By analyzing single-cell RNA-seq data from the entorhinal cortex

and spatiotemporal RNA-seq data, encompassing 1890 human brain samples, we found that

HNP neurons had increased metabolic demands as well as increased expression of genes

associated with tau misfolding, which was additionally indicated by neurons expressing NPs in

AD showing distinct molecular signatures of protein misfolding. We further report that the

decrease of ADNP expression with aging was specific to early AD-impacted brain regions and

that HNP neurons co-expressing ADNPs were heavily distributed in "epicenters". These results

support the hypothesis that the functional demands of HNP neurons likely predispose them to

oxidative stress during aging, perhaps contributing to their early demise in AD.

Figure 1

Methods

Acquisition and processing of transcriptome data

<u>Single-cell RNA-seq data from the human entorhinal cortex (EC)</u>

Two sets of single-cell EC transcriptome data were included for analysis in this manuscript.[9,10]

Data generated by Grubman et al from six control (CT) and six AD donors (12 total) were

obtained.[9] The detailed documentation of data acquisition, preprocessing steps, and

downstream analyses (dimension reduction and cell identification) can be found in Li and

Larsen (2023).[8] Briefly, cells were filtered based on gene expression and mitochondrial content,

retaining those with 200-2500 expressed genes and less than 5% mitochondrial reads. Data

normalization followed Seurat's (version 4.1.1; R 4.2.3 unless specified) guidelines using a

count per million (CPM) matrix.[11] Using cell identification methods from *BRETIGEA* and

Grubman et al, six primary cell types were identified: astrocytes, microglia, neurons,

oligodendrocytes, oligodendrocyte precursor cells, and endothelial cells. Cells were labeled by

their highest association score, but some were categorized as unidentified or hybrid based on

specific criteria described in detail by Grubman et al as well as Li and Larsen.[8,9] Only cells

identified as neurons were used in the downstream analyses.

In addition, preprocessed and annotated single-cell transcriptome data from human EC

generated by the the MIT ROSMAP Single-Nucleus Multiomics Study

(MIT_ROSMAP_Multiomics) was downloaded from ADKnowledge portal.[12] Three diagnostic

categories from the ROSMAP study were included in the presented study: control (CT or NCI:

no cognitive impairment), mild cognitive impairment (MCI: no other condition contributing to CI),

and AD (Alzheimer's dementia: no other condition contributing to CI (NINCDS/ADRDA Probable

AD)).[12] The clinical study design and detailed diagnostic criteria were described in the original

ROSMAP manuscript and deposited on ADKnowledge portal.[12,13] Individuals with inconsistent

clinical diagnosis, clinical cognitive diagnosis summary, and final consensus cognitive diagnosis

documented in the ROSMAP study were excluded from the analysis. Methods for isolation of

nuclei from frozen post-mortem brain tissue, droplet-based snRNA-seq, and snRNA-seq data

preprocessing are available in detail on ADKnowledge portal.[12] De-identified metadata for

individuals and experiments included in this study were detailed in supplementary materials

(Table S1). To provide equal representation of each condition, we randomly selected eight AD

and CT samples (set.seed = 123) to match the eight available MCI samples. Overall, 41373

neurons (n_neuron = 41373) from eight AD (n_neuron = 16214), MCI (n_neuron = 11732), and

CT (n_neuron = 13427) EC regions were included in the final analysis.

<u>Spatiotemporal bulk RNA-seq data from human brains</u>

RNA-seq transcript matrices (transcript per million (TPM)) of 11 human brain regions from a

cohort of individuals from the general population were obtained from the GTEx portal (Table

S2).[14] Samples from individuals that lacked complete metadata regarding age, sex, or death

classification were excluded, as were those that scored 3 or 4 on the Hardy Scale that indicates

intermediate or slow death. Only samples with an RNA integrity number (RIN) larger than 6

were included in the analysis. The results were visualized with *cerebroViz* (version 1.0; R

3.6.3)[15] and BioRender.

Downstream analyses for single-cell transcriptome data from human EC

<u>Correlational analysis of NP transcripts and ADNPs co-expression</u>

The Spearman rank correlation was utilized to assess the relationship between the levels of

transcripts of NP and the number of co-expressed ADNPs using *cor.test* in R for both human

single-cell transcriptome data. This test is suitable for datasets that do not distribute with

normality. Correlation coefficient (rho) and p-values were reported. The significance cut-off was

set at 0.05.

<u>Differential gene expression (DGE)</u>

DGE analysis was performed for the single-cell dataset generated by Grubman et al. (2019).[9]

The number of co-expressed NPs was used as a proxy for transcript levels of NPs; neurons

were divided into low (0-1), medium (2-5), and high (6+) NP-producing groups (LNP, MNP,

HNP) based on the number of co-expressed NPs in both conditions. DGE analysis was

implemented between LNP and HNP in control neurons as well as MNP groups in control and

AD, using *FindMarkers* function in *Seurat*.[11] Default parameters for DGE in *Seurat* were used

(Wilcoxon Rank Sum test); statistical cut-off was set at 0.05 for a false discovery rate (FDR)

adjusted by Benjamin-Hochberg (BH) method.

<u>Functional enrichment analysis</u>

The output of the DGE analysis from *Seurat* was used as input to the STRING database

(version 11.5).[16] Key enrichment output from STRING analysis was visualized using the

Enrichplot package (version 1.18.4).[17] STRING uses Bonferroni method to correct for multiple

comparisons and provides adjusted p-values.[16] The significance cut-off was set at 0.05 for FDR.

<u>Regression analysis of NP transcripts and ADNPs co-expression</u>

Regression analysis was used to discern the relationship between gene transcript levels and the

presence of NP. Utilizing the *glm* function in R, general linear models were constructed for each

differentially expressed genes (DEG) with transcript levels as the response variable and the

number of co-expressed NPs as the explanatory variable.The BH method was used to adjust for

multiple comparisons for all p-values. 0.05 was used as the significance cut-off for adjusted p-

values. The goal was to identify genes whose expression is notably influenced by the increased

number of co-expressed NPs. Results can be found in Table S3.

<u>Hypergeometric test of cell abundance and ADNP co-expression</u>

The hypergeometric test was employed using the *phyper* function in R to assess the overlap of

genes with decreased expression in AD compared to those that are functionally enriched in

HNP neurons.[18] Specifically, it was defined: 1) the number of overlapped genes as "successes"

in our sample (x = 25); 2) genes with significantly decreased expression in AD MNP neurons as

"successes" in the population (m = 91); 3) the total number of unique genes expressed by AD

MNP and control HNP neurons, minus those with decreased expression in AD MNP neurons, as

"failures" in the population (n = 19430); 4) genes with significantly increased expression in HNP

neurons as the sample size (k = 307). Since phyper is a cumulative distribution function, the

conduction of a one-tailed such analysis (lower.tail = FALSE) would calculate a probability of

observing as extreme and more extreme results in the direction of higher values (p-value).[18] The

significance cut-off was set at 0.05.

<u>Stratified assessment of cognitive condition-neuropeptide abundance and independence</u>

Using the MIT_ROSMAP_Multiomics dataset, Jonckheere-Terpstra test from *DescTools*

package[19] was used to check if there was a statistically significant trend across the continuum

from CT, through MCI, then to AD for both top 50% and top 25% of all ADNP-coexpressing

neurons. 0.05 was used as the significance cut-off. Robust t-test (Yuen Two Sample t-test) with

a trim of 0.1 in R[19] was used as a post-hoc test to compare between conditions; 0.1 was used

as the significance cut-off to ensure all relevant findings were captured. For each condition in

the MIT_ROSMAP_Multiomics dataset (CT, AD, or MCI),[10,12] The number of co-expressed key

NPs (output from ML models as reported previously[8]) was also recorded, ranging from 0 to 3.

To investigate the relationships between the conditions and the number of co-expressed NPs, a

series of Chi-squared tests to assess the independence between each pair of conditions (AD vs

CT, AD vs MCI, and CT vs MCI) were conducted in R. These tests were stratified by the number

of co-expressed NPs to examine how this variable might influence the relationships between the

conditions. A significance level of $\alpha = 0.05$ was used for all tests.

Spatiotemporal correlation analysis between NP gene expression and age

Gene lists of NPs and ADNPs were downloaded from existing publications.[8,20] TPM count matrix

of spatiotemporal RNA-seq data and metadata of human brains were downloaded from GTEx.[14]

All TPM counts were log2 transformed. The correlation between ADNP expression and age was

determined using *PResiduals* package,[21] adjusting for RIN and Hardy Scale. The significance

cut-off was set at 0.05.

Single-cell examination of ADNP-coexpressing HNP (AHNP) neurons across

microdissected brain regions

A comprehensive analysis of AHNP occurrences and cell counts across microdissected brain

regions was performed using single-cell data and an annotated cell cluster (cluster_annotation)

file generated by Siletti et al.[22] File documentation from the study was downloaded from github

link provided by Siletti et al.[22] Firstly, the presence of ADNPs from the file was quantified and

counts for NPs were generated further to calculate non-ADNPs. Based on these counts, we

defined ADNP-HNP (AHNP) neurons as those tagged with > 5 ADNPs and < 3 non-ANDPs,

accounting for the difference in the input NPs.[8,22] Concurrently, we estimated the number of

cells in different brain regions and dissections based on percentage data extracted from the

cluster_annotation file. These estimates were summed across unique regions and dissections to

provide a granular view of cell distribution. Additional analyses were conducted on specific

regions for MEC, where the dataset was grouped by various attributes such as

Neurotransmitter, Subtype, and MTG label,[23], and the number of cells in each group was

summed and visualized.

Results

Alterations of HNP neuronal abundance and functions in AD: overlap of HNP dysfunction and molecular signature of AD

Analyzing the single-cell dataset by Grubman et al,[8,9,24] we showed that the correlation between transcript abundance and the number of co-expressed NPs generally exists for neurons in both control and AD (Figure 2A). Applying the number of co-expressed NPs as a proxy for transcript levels of NPs, we stratified neurons into different groups in AD and controls based on the number of co-expressed NPs, including low (0-1), medium (2-5), and high (6+) NP groups. We observed a similar absence of HNP neurons in AD (Figure 2B; Table S4).[8]

We previously reported a significant reduction of HNP neurons expressing more neuropeptides in the EC of AD brains. As we hypothesized that HNP neurons would experience greater energetic demands and higher metabolic stress, we predicted that HNP neurons would show more metabolic activity. To test this, differentially expressed genes (DEG) were examined between high and low groups of control neurons followed by enrichment analysis.[8,16] We found that all the NPs expressed at significantly higher levels in HNP neurons were AD-associated NPs (ADNPs),[8] and genes required for NP transportation, translation, and metabolic processes were significantly increased in HNP neurons in comparison to LNP neurons (Figure 2C&D; Table S5&6). Although it is widely known that GABA is often co-expressed with NPs,[25] we observed that HNP neurons also participated in other chemical communications, such as histamine and catecholamines, to regulate membrane potential (Figure 2D; Table S5). In addition, ~36% of DEGs regulating membrane potentials were functionally enriched for learning and memory (Figure 2D; Table S5). Surprisingly, the regulation of innate immune response was increased in HNP cells (Figure 2C&D; Table S5).

To elucidate the increase observed across the LNP, MNP, and HNP groups, we sought to identify genes whose expression was significantly influenced by the level of NP co-expression,

which also correlated with the abundance of NP transcripts. We utilized regression models to

examine the relationship between gene expression and NP co-expression then ranked the

increased DEGs in HNP neurons by coefficients and R^2. Excluding NP components, genes

ranked in the top 10 for coefficients or R^2 included several long noncoding RNAs, components

of A-to-I RNA editing, and *NPAS3*–encoding for a transcription factor related to circadian rhythm

regulation[26] (Table S3; Figure S1). Notably, *ERBB4*, the protein products of which respectively

induced tau hyperphosphorylation[27] was among the top genes related to NP co-expression

(Table S3; Figure S1).

Since we discovered that HNP neurons exhibited higher performance in several expected

cellular functions–including transportation, translation, and metabolic processes–and

participated more heavily in other chemical communications, regulation of innate immune

response, and circadian rhythm, we wondered whether dysregulation of processes in these

functions may lead to the loss of neuronal functions and the accumulation of tau pathology,

which are hallmarks of several neurodegenerative disorders including AD.

Speculating that disrupted functions of neurons expressing more NPs are associated with

protein misfolding, we examined the DEGs for neurons stratified by the number of co-expressed

NPs and analyzed the neurons in the medium group (as HNP neurons were virtually absent in

AD brains). Enrichment analysis revealed that these cells displayed molecular characteristics

related to both "positive" and "negative" neuropathology in AD.[2] We note the decreased DEG in

AD cells were significantly enriched for those functionally increased in HNP cells ($p < 0.00001$;

Figure 2E; Table S6&7), indicating that loss of HNP functions participates in the molecular

pathogenesis of AD. Genes with protein products showing significantly decreased expression

included those with functional roles in axons, synapses, and dendrites (Figure 2F; Table S7).

Increased molecular processes included those known to be disturbed in AD, such as negative

regulation of neurogenesis, gliogenesis, and abnormal mitochondrial metabolism (Figure 2F;

Table S8). Notably, genes involved in forming aggresomes and unfolded protein binding were

highlighted,[28] indicating the active occurrence of protein misfolding in AD cells that co-express

NPs (Figure 2F; Table S8).

Figure 2

Loss of ADNP expression participates in early pathogenesis of AD

Aging is the predominant risk factor for developing AD[1]. The decline in cognitive function during natural aging bears resemblance to that seen in the early stages of AD.[29,30] Mild cognitive impairment (MCI) is defined as a stage between normal age-related cognitive changes and pathological cognitive impairments, and it frequently precedes and is considered a potential early sign of Alzheimer's disease.[13] Therefore, examining datasets from both natural aging and MCI would provide a compelling framework to assess whether disruptions in ADNP expression play a role in the early pathogenesis of AD. Our recent report indicated that ADNP expression decreased with age in the hippocampus;[8] however, we do not yet know if the decline of NPs with aging is ADNP- and brain region-specific. To strengthen the evidence supporting the early involvement of ADNPs in AD development, we 1) analyzed bulk RNA-seq data from all brain regions with sufficient sample size generated by GTEx and 2) assessed the abundance of HNP neuron co-expressing ADNPs in the EC of AD, MCI, and control (CT) brains using recent single-cell transcriptome data.[10]

With respect to aging, if our hypothesis were to hold, only brain regions affected by early AD should show age-related changes for ADNP expression and only the accumulative expression of ADNPs, but not other NPs, should decrease with age in the human brain. We first examined the expression of ADNPs during aging among 11 brain regions selected from GTEx (Table 1).[8,14] Supporting our hypothesis, we found that only the hippocampus, frontal cortex, anterior cingulate gyrus, and amygdala—all of which are brain regions affected by early AD[31–36]—showed a significant decrease of ADNP transcription during aging among 11 brain regions selected from GTEx (Table 1; Figure 3A).[8,14,37] We also comprehensively examined the expression of NPs in these brain regions and whether the expression of an individual NP was correlated with age. Using this dataset, we found that ~108 NPs were expressed by each brain region on average (Table S9); the top brain regions that demonstrated a decrease in NP

expression with aging were the hippocampus (25), anterior cingulate gyrus (28), frontal cortex

(16), and amygdala (16) (Table S9). We confirmed that the larger decrease of NP expression in

these brain regions with aging was not attributed to their intrinsic capacity to express NPs;

neither the number nor the average transcript count for NPs differed significantly among

surveyed regions (Table S9). A closer examination of the NPs that demonstrated a decrease in

expression with age in the aforementioned brain regions revealed that these NPs consisted of

expressed ADNPs included in our previous study.[8] To support the specificity of the observed

change in ADNP expression, we analyzed the counts of all non-ADNP NPs in the brain regions

under examination and found no age-related changes in their overall levels (Table S10). In

short, the spatiotemporal transcriptome analysis presented here showed that the decrease of

NPs in the aging human brain was specific to brain regions and NPs implicated in AD,

supporting that age-related cognitive decline shares mechanisms with AD and may be mediated

by loss of ADNP expression during aging.

To further investigate whether loss of HNP neurons co-expressing ADNPs could be associated

with cognitive impairment in early AD, we postulated that brains from individuals with MCI would

maintain more of these neurons than AD but not CT donors. Using single-cell transcriptome

data generated from CT, AD, and MCI EC region (see Methods), we first confirmed that the

correlation between the transcript level of NPs and the number of co-expressed NPs still exist in

this dataset (rho = 0.89, p < 0.0001; Figure S2). We subsequently visualized and analyzed the

distribution of neurons across three conditions, based on the number of co-expressed

neuropeptides (NPs), and marked the median and the 3rd quantile (Q3) values for each

condition (Table S11; Figure 3B). Consistent with previous reports (Figure 1B),[8] distribution of

AD neurons showed right skewness in comparison to CT (Figure 3B). As expected, MCI

neurons also showed right skewness but less so than that in AD (Figure 3B). To quantify the

observation, we analyzed the abundance of neurons above the median ADNP co-expression

levels observed in CT for all three conditions; there was a statistically significant decreasing

trend in the cognitive continuum from CT, MCI, to AD ($p < 0.05$; Figure 3C) and CT ECs had

significantly higher neuronal counts than MCI and AD (Figure 3C). We furthered the comparison

to the third quantile of CT; similarly, the decreasing trend persisted ($p < 0.01$; Figure 3C) and

there were significant differences across these conditions.

Considering subsets of HNP neurons as a representative example, we utilized a combination of

the neuropeptide outputs (key ADNPs: *SST, VGF, VIP*, and *CRH*) derived from three previously

reported machine learning algorithms (Figure 3D; see Methods).[8] For neurons in each condition

among CT, MCI, and AD, we recorded the number of expressed key ADNPs, ranging from 0 to

3. For neurons expressing no key NPs, no significant differences were observed between MCI

and AD but both were more abundant than CT ($p < 0.0001$; Figure 3D; Table S12&13). For the

number of neurons expressing one or three key NPs, significant differences were again only

observed where neuronal counts were higher in CT than the other two conditions ($p < 0.0001$;

Figure 3D; Table S12&13). For neurons expressing two key NPs, neuronal counts were highest

in CT, followed by MCI, and then AD, with significant differences observed between the groups

($p < 0.0001$; Figure 3D; Table S12&13). Overall, our analysis supported the early involvement of

ADNP expression in AD pathogenesis.

Table 1. Significantly decreased expression of ADNPs with aging only occurred in early

AD-impacted regions.

Brain Region	Sample Size	Correlation	p-value	Significance
Hippocampus	103	-0.30	0.00	*
Frontal cortex (BA9)	116	-0.35	0.00	*
Anterior cingulate cortex (BA24)	88	-0.30	0.01	*
Amygdala	73	-0.30	0.02	*
Hypothalamus	109	0.02	0.85	
Caudate basal ganglia	146	-0.08	0.28	
Nucleus accumbens basal ganglia	143	0.00	0.96	
Putamen basal ganglia	126	0.11	0.25	

Substantia nigra	66	-0.23	0.08	
Cerebellar hemisphere	141	-0.03	0.74	
Cerebellum	156	-0.08	0.34	

317   ADNP, Alzheimer's Diseases (AD)-associated neuropeptides; BA, Brodmann area; *,

318   p_value<0.05, α = 0.05.

319   **Figure 3**

Physiological distribution of AHNP neurons may mediate brain region vulnerability to AD

"Epicenters" have been described as brain regions showing the most significant pathological

alterations, hypothesized to be the initial site of disease onset.[3,4] Stressed "nodes" are known as

brain regions with high network traffic, also referred to as "hubs," that experience activity-

induced deterioration that can lead to or exacerbate diseases.[5,38] While both concepts are

instrumental in theories of AD etiology devised by connectome and network-based studies,[3–5,38]

the cell types underlying these "epicenters" and stressed "nodes" are unclear. Based on the

regions that display reduced ADNP expression with aging, we propose that AHNP neurons

serve as one of the cellular components of the "hubs" and "epicenters" leading to the onset

and/or progression of AD. To test our hypothesis, we analyzed the distribution of AHNP neurons

across various regions of the human brain. Two potential observations and implications exist: 1)

AHNP neurons ubiquitously exist in all brain regions but those in early-AD impacted regions are,

regardless of the underlying cause, more susceptible to dysfunction than others, or 2) AHNP

neurons are preferentially distributed in early-AD impacted brain regions to physiologically

perform cognitive functions, but they are more prone to dysfunction than other cell types,

therefore mediating the regional vulnerability for AD with aging-related cell dysfunctions. We

predict that a single-cell transcriptomic survey of brain regions would show that AHNP neurons

are more abundantly distributed in early AD-affected brain regions that engage extensively in

memory and executive functions, such as the entorhinal cortex, hippocampus, and basal

forebrain[39–41]. However, if the first scenario were true, our hypothesis could be negated

altogether.

A recent study published single-cell transcriptome data from ~100 dissections across the

forebrain, midbrain, and hindbrain of human donors, and classified brain cells into 461

clusters.[22] The authors also compared their cell clusters with previous publications that used NP

diversity to classify neurons.[22,23] We first investigated the top brain regions where AHNP

neurons using cluster annotations provided by Siletti et al and calculated the number of neurons

in each top three brain regions and microdissections (see Methods).[22] As anticipated, the

amygdala and hippocampus—where the age-associated decrease of ADNP expressions was

observed[8]—were among the top five brain regions (Figure 4A; Table 2). While existing literature

identified the hypothalamus as considerably relevant in AD,[42,43] it didn't show an age-related

decrease in ADNPs. We also found that the cerebral cortex contained the most AHNP neurons,

but this was likely due to the number of dissections assigned to the cerebral cortex. To

overcome this issue, we set out to identify the distribution of AHNP neurons among cortical

regions separately and ranked the cortical dissections based on the number of AHNP neurons.

Again, we found that medial and lateral EC (MEC and LEC) ranked among the top five cortical

regions (Figure 4A; Table 3). In particular, the MEC harbored the highest number of AHNP

neurons among all micro-dissected regions (Table S14). In contrast, very few AHNP cells were

found in the cerebellum, spinal cord, and medulla, where it is generally thought to be spared by

AD neuropathology (Table S15), further supporting our hypothesis that dysfunction of AHNP

cells mediates the brain region-specific vulnerability to AD.

As the EC—the earliest and most heavily affected region in AD neuropathology—harbored the

most significant population of AHNP neurons, we sought to investigate the specific cell types as

previous investigation has suggested that most ADNP co-expressing neurons were GABA-

ergic.[8] Specifically, we aimed to understand their how they align with established neuronal cell

types. We selected neurons that had MEC in the top three dissections and examined the

proportions of neuron types, including based on the neurotransmitter, subtypes, and transferred

MTG labels (common cell type nomenclatures for the MTG regions of the mammalian brain).[22]

We found that most of the AHNP neurons were indeed GABA-ergic interneurons; however,

about a quarter of the neurons have not been described before (Figure 4B).

Figure 4

Table 2. Top five brain regions ranked by AHNP neuron abundance.

Brain regions	AHNP neuronal count
Cerebral cortex	218145
Amygdala	72256
Basal forebrain	45559
Hippocampus	40870
Hypothalamus	23217

373 **Table 3. Top five cortical regions ranked by AHNP neuron abundance.**

Cortical regions	Full cortical description	AHNP neuronal count
MEC	Anterior parahippocampal gyrus, posterior part (APH) - Medial entorhinal cortex	19206
M1C	Precentral gyrus (PrCG) - Primary motor cortex	11821
MTG	Middle Temporal Gyrus	10654
LEC	Anterior parahippocampal gyrus (AG) - Lateral entorhinal cortex	2625
A40	Supramarginal gyrus (SMG) - A40	1732

Discussion

Why some ubiquitously expressed proteins (e.g., tau and Aβ) exhibit selective accumulation in particular regions of the brain and cells, yet sparing their comparable neighbors, is fundamental to AD research. Examining multiscale and spatiotemporal RNA-seq data from 1890 human brain samples, we aimed to gain a comprehensive understanding of the potential mechanistic roles that high NP-producing (HNP) neurons that co-express AD-associated neuropeptides (ADNPs) play in mediating the selective vulnerability of brains to AD.

To address our previous hypothesis that HNP neurons, given their secretory and peptidal signaling functions, would demand more translation and transportation, leading to increased metabolic vulnerability,[8] we investigated the specific characteristics of HNP neurons to the link between HNP neuron dysfunction and the hallmark molecular indicators of AD. We found that 1) HNP cells were more metabolic active and had gene expression profiles suggesting higher connectivity; 2) MEC AD neurons co-expressing higher levels of NPs showed the molecular signature of AD, including protein misfolding; and 3) the deficiency of AD cells was linked to loss-of-functions of HNP cells. While we anticipated a greater metabolic burden as a source of neuronal vulnerability for HNP neurons, it was surprising to discover that HNP neurons can be predisposed to tau hyperphosphorylation and misfolding.[27,44] In addition to *ERBB4*, recent research has revealed that the impairment of NMD, an elevated process observed in HNP cells and suppressed by cellular stress, mediates tau-induced neurotoxicity.[45] Therefore, we posit that the disruption of cellular processes elevated in HNP neurons could occur more readily/earlier in these cells and contribute to the formation of misfolded proteins within them, subsequently leading to their selective degeneration. This position aligns with the existing knowledge that brain regions with strong anatomical connectivity harbored more misfolded proteins[37,38]; the cell clusters carrying out the functions, such as HNP neurons with greater connectivity and metabolic activity described in this study, are likely more prone to protein

misfolding–the general framework of which was already proposed first as "nodal stress" and

explained in detail by others.[5,6,38] Further, it is generally agreed that brain regions accumulating

most pathology-associated proteins (disease-susceptible brain regions) physiologically contain

the cells that are most susceptible to cytotoxic events and are among the earliest affected in the

disease's progression.[3,4,6] Taken together, at the cellular level, our results indicate that

disproportionate loss of HNP neurons in AD could be a reflection of their earlier involvement in

the disease process as the result of higher chances of cytotoxic events–which are associated

with their functions–occurring, such as propensity to oxidative stress, mitochondrial dysfunction,

and accumulation of misfolded proteins.

To comprehensively demonstrate that the loss of ADNPs expressed by HNP neurons

participates in early AD development and/or progression, we considered perspectives from both

aging and MCI. We analyzed spatiotemporal transcriptomic data of brains from the natural aging

populations generated by the GTEx consortium to reveal that the decrease in ADNP

expressions with aging is specific to early AD-associated brain regions. Supporting our

hypothesis that dysfunction of ADNP expression in HNP (AHNP) neurons appears early in AD,

we observed a significant decline in the abundance of AHNP neurons across the continuum

from CT, progressing through MCI, to AD. These results are consistent with previous studies

that underscored the role of key ADNPs play in processes crucial to the pathogenesis of

neurodegeneration, such as mitochondrial dysfunction, persistent neuroinflammation, and

disrupted circadian rhythm (briefly summarized by Li and Larsen),[8] indicating that alterations in

the abundance and diversity of ADNP-producing neurons likely participate in the onset and/or

progression of cognitive decline and neurodegeneration in AD. Our findings also highlight the

relevance of ADNPs as combinatorial and longitudinal biomarkers to evaluate the risk and

progression of AD development. For instance, an analysis of the CSF proteome identified that

CHGA and VGF exhibited significant differences in abundance among CT, MCI, and AD groups.

Further studies validating these findings and investigating the underlying mechanisms

responsible for the observed decline in AHNP neuron abundance are needed. Evaluating

whether interventions aimed at mimicking and/or preserving the ADNP signaling network could

impede progression of AD is also valuable, especially considering the subtle differences in

GPCR expression observed between CT and AD.[8] Collectively, these observations suggest that

deficiencies of ADNPs with aging contribute to AD development and progression and that these

deficits could be a consequence of losing AHNP neurons during aging, which underlies shared

cognitive decline during aging and in early AD.

Cellular dysfunctions, characterized by distinct molecular pathways and biochemical properties,

play crucial roles in the onset and progression of AD.[6] Extensive research over the past decade

has delved into connectome-based studies of neurodegenerative diseasesand and provided

profound understanding of the brain regions and neural circuitries vulnerable to AD[3,4,7,38];

however, the specific cell types and cellular components that underpin these connectomic

findings in relation to AD remain elusive. Therefore, our presentation of the physiological

distribution of HNP neurons within the brain is fundamental to deciphering their roles in AD; their

preferential localization in AD-susceptible brain regions further signifies their potential role as

cellular elements contributing to the initiation of AD. However, potential confounding variables,

such as transcription factors and epigenetic mechanisms, warrant careful consideration. Recent

research has made tremendous advancements on providing a comprehensive multi-omic brain

atlas for physiological conditions,[22,46,47] our results suggest that such efforts will also be vital to

explore brains of diseased states. By understanding how dysfunction in HNP neurons occurs

with aging and its detrimental effects on interconnected behavioral domains, we can gain

valuable insights into the cell types and dysfunctions at the early stages of AD to develop

effective disease models as well as cell type-specific targeting strategies for prevention and

therapy.

References:

1. 2023 Alzheimer's disease facts and figures. *Alzheimers Dement.* 2023;19(4):1598-1695.

2. Serrano-Pozo A, Frosch MP, Masliah E, Hyman BT. Neuropathological alterations in Alzheimer disease. *Cold Spring Harb Perspect Med.* 2011;1(1):a006189.

3. Zhou J, Gennatas ED, Kramer JH, Miller BL, Seeley WW. Predicting regional neurodegeneration from the healthy brain functional connectome. *Neuron.* 2012;73(6):1216-1227.

4. Vogel JW, Corriveau-Lecavalier N, Franzmeier N, et al. Connectome-based modelling of neurodegenerative diseases: towards precision medicine and mechanistic insight. *Nat Rev Neurosci.* Published online August 24, 2023.

5. Saxena S, Caroni P. Selective neuronal vulnerability in neurodegenerative diseases: from stressor thresholds to degeneration. *Neuron.* 2011;71(1):35-48.

6. Fu H, Hardy J, Duff KE. Selective vulnerability in neurodegenerative diseases. *Nat Neurosci.* 2018;21(10):1350-1358.

7. Yu M, Sporns O, Saykin AJ. The human connectome in Alzheimer disease - relationship to biomarkers and genetics. *Nat Rev Neurol.* 2021;17(9):545-563.

8. Li M, Larsen PA. Single-cell sequencing of entorhinal cortex reveals widespread disruption of neuropeptide networks in Alzheimer's disease. *Alzheimers Dement.* Published online February 24, 2023.

9. Grubman A, Chew G, Ouyang JF, et al. A single-cell atlas of entorhinal cortex from individuals with Alzheimer's disease reveals cell-type-specific gene expression regulation. *Nat Neurosci.* 2019;22(12):2087-2097.

10. Mathys H, Peng Z, Boix CA, et al. Single-cell atlas reveals correlates of high cognitive function, dementia, and resilience to Alzheimer's disease pathology. *Cell.* 2023;186(20):4365-4385.e27.

11. Gribov A, Sill M, Lück S, et al. SEURAT: visual analytics for the integrated analysis of microarray data. *BMC Med Genomics.* 2010;3:21.

12. Bionetworks S. Synapse.

13. Bennett DA, Schneider JA, Arvanitakis Z, Wilson RS. Overview and findings from the religious orders study. *Curr Alzheimer Res.* 2012;9(6):628-645.

14. GTEx Portal.

15. Bahl E, Koomar T, Michaelson JJ. cerebroViz: an R package for anatomical visualization of spatiotemporal brain data. *Bioinformatics.* 2017;33(5):762-763.

16. Szklarczyk D, Gable AL, Lyon D, et al. STRING v11: protein–protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* 2019;47(D1):D607-D613.

17.   Yu G. enrichplot: Visualization of functional enrichment result. R package version
1.10. 2. *Mol Ther Nucleic Acids*.

18.   Johnson NL, Kotz S, Kemp AW. *Univariate Discrete Distributions*. Wiley; 1992.

19.   Signorell A. *DescTools: Tools for Descriptive Statistics and Exploratory Data
Analysis*. Github

20.   Tai KY, Wong K, Aghakhanian F, Parhar IS, Dhaliwal J, Ayub Q. Selected
neuropeptide genes show genetic differentiation between Africans and non-Africans. *BMC
Genet*. 2020;21(1):31.

21.   Liu Q, Shepherd B, Li C. PResiduals: An R Package for Residual Analysis Using
Probability-Scale Residuals. *J Stat Softw*. 2020;94:1-27.

22.   Siletti K, Hodge R, Albiach AM, et al. Transcriptomic diversity of cell types across
the adult human brain. *bioRxiv*. Published online October 14, 2022:2022.10.12.511898.

23.   Jorstad NL, Song JHT, Exposito-Alonso D, et al. Comparative transcriptomics
reveals human-specific cortical features. *bioRxiv*. Published online September 19,
2022:2022.09.19.508480.

24.   Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal
single-cell data. *Cell*. 2021;184(13):3573-3587.e29.

25.   Lizbinski KM, Marsat G, Dacks AM. Systematic Analysis of Transmitter
Coexpression Reveals Organizing Principles of Local Interneuron Heterogeneity. *eNeuro*.
2018;5(5).

26.   Sha L, MacIntyre L, Machell JA, et al. Transcriptional regulation of
neurodevelopmental and metabolic pathways by NPAS3. *Mol Psychiatry*. 2012;17(3):267-
279.

27.   Nie SD, Li X, Tang CE, et al. High glucose forces a positive feedback loop
connecting ErbB4 expression and mTOR/S6K pathway to aggravate the formation of tau
hyperphosphorylation in differentiated SH-SY5Y cells. *Neurobiol Aging*. 2018;67:171-180.

28.   Albaret MA, Textoris J, Dalzon B, et al. HSV-1 cellular model reveals links
between aggresome formation and early step of Alzheimer's disease. *Transl Psychiatry*.
2023;13(1):86.

29.   Small SA, Nava AS, Perera GM, Delapaz R, Stern Y. Evaluating the function of
hippocampal subregions with high-resolution MRI in Alzheimer's disease and aging.
Microsc Res Tech. 2000;51(1):101-108.

30.   Mitchell TW, Mufson EJ, Schneider JA, et al. Parahippocampal tau pathology in
healthy aging, mild cognitive impairment, and early Alzheimer's disease. *Ann Neurol*.
2002;51(2):182-189.

31.   Laakso MP, Soininen H, Partanen K, et al. Volumes of hippocampus, amygdala
and frontal lobes in the MRI-based diagnosis of early Alzheimer's disease: correlation with
memory functions. *J Neural Transm Park Dis Dement Sect*. 1995;9(1):73-86.

524 32. Albert MS, Moss MB. Early Features of Alzheimer's Disease. In: Peters A,
525 Morrison JH, eds. *Cerebral Cortex: Neurodegenerative and Age-Related Changes in*
526 *Structure and Function of Cerebral Cortex*. Springer US; 1999:461-474.

527 33. Jones BF, Barnes J, Uylings HBM, et al. Differential regional atrophy of the
528 cingulate gyrus in Alzheimer disease: a volumetric MRI study. *Cereb Cortex*.
529 2006;16(12):1701-1708.

530 34. Amanzio M, Torta DME, Sacco K, et al. Unawareness of deficits in Alzheimer's
531 disease: role of the cingulate cortex. *Brain*. 2011;134(Pt 4):1061-1076.

532 35. Spalletta G, Piras F, Piras F, et al. Neuroanatomical correlates of awareness of
533 illness in patients with amnestic mild cognitive impairment who will or will not convert to
534 Alzheimer's disease. *Cortex*. 2014;61:183-195.

535 36. Poulin SP, Dautoff R, Morris JC, Barrett LF, Dickerson BC, Alzheimer's Disease
536 Neuroimaging Initiative. Amygdala atrophy is prominent in early Alzheimer's disease and
537 relates to symptom severity. *Psychiatry Res*. 2011;194(1):7-13.

538 37. Braak H, Braak E. Neuropathological stageing of Alzheimer-related changes.
539 *Acta Neuropathol*. 1991;82(4):239-259.

540 38. Buckner RL, Sepulcre J, Talukdar T, et al. Cortical hubs revealed by intrinsic
541 functional connectivity: mapping, assessment of stability, and relation to Alzheimer's
542 disease. *J Neurosci*. 2009;29(6):1860-1873.

543 39. Pennanen C, Kivipelto M, Tuomainen S, et al. Hippocampus and entorhinal
544 cortex in mild cognitive impairment and early AD. *Neurobiol Aging*. 2004;25(3):303-310.

545 40. Fernández-Cabello S, Kronbichler M, Van Dijk KRA, et al. Basal forebrain
546 volume reliably predicts the cortical spread of Alzheimer's degeneration. *Brain*.
547 2020;143(3):993-1009.

548 41. Grothe M, Heinsen H, Teipel SJ. Atrophy of the cholinergic Basal forebrain over
549 the adult age range and in early stages of Alzheimer's disease. *Biol Psychiatry*.
550 2012;71(9):805-813.

551 42. Zhou JN, Hofman MA, Swaab DF. VIP neurons in the human SCN in relation to
552 sex, age, and Alzheimer's disease. *Neurobiol Aging*. 1995;16(4):571-576.

553 43. Cedernaes J, Osorio RS, Varga AW, Kam K, Schiöth HB, Benedict C. Candidate
554 mechanisms underlying the association between sleep-wake disruptions and Alzheimer's
555 disease. *Sleep Med Rev*. 2017;31:102-111.

556 44. Woo RS, Lee JH, Yu HN, Song DY, Baik TK. Expression of ErbB4 in the
557 apoptotic neurons of Alzheimer's disease brain. *Anat Cell Biol*. 2010;43(4):332-339.

558 45. Zuniga G, Levy S, Ramirez P, et al. Tau-induced deficits in nonsense-mediated
559 mRNA decay contribute to neurodegeneration. *Alzheimers Dement*. 2023;19(2):405-420.

560 46. Tian W, Zhou J, Bartlett A, et al. Single-cell DNA methylation and 3D genome
561 architecture in the human brain. *Science*. 2023;382(6667):eadf5357.

47.   Li YE, Preissl S, Miller M, et al. A comparative atlas of single-cell chromatin
accessibility in the human brain. *Science.* 2023;382(6667):eadf7044.

Figure Legends

Figure 1. Hypothetical model illustrating the potential mechanisms underlying the

selective vulnerability of high neuropeptide-producing (HNP) cells and their contribution

to region-specific emergence of Alzheimer's disease (AD). The model proposes that the

unique functions of HNP neurons make them more prone to stress and protein misfolding, and

this susceptibility becomes more evident with advancing age. The regional vulnerability

observed in Alzheimer's disease can be attributed to the distribution and density of these cells,

as well as their excretory functions and interactions with other brain networks. Specifically, it

predicts that (1) temporal, limbic, and prefrontal cortical regions have a higher density of HNP

neurons expressing ADNPs; (2) disruption of cellular processes in HNP neurons leads to a

decrease in ADNPs during aging and localized formation of misfolded proteins in AD, causing

various degrees of cognitive decline and selective degeneration of these neurons; (3) the

dynamic paracrine and secretory activities of HNP cells facilitate the propagation of misfolded

proteins and transneuronal degeneration in adjacent temporal, limbic, and prefrontal cortical

regions, resulting in widespread deposition of misfolded tau in AD.

Figure 2. Changes in high neuropeptide-producing (HNP) neuronal abundance and

function: overlap of dysfunction and AD molecular signatures. (A) The relationship

between transcript abundance and the number of co-expressed neuropeptides (NPs) in neurons

from both control and AD EC. **(B)** The distribution of neurons based on the number of co-

expressed NPs. low: 0-1; medium (mid): 2-5, and high: 6+. Proportion = in-group neuron count

in the condition/total neuron count in the condition. **(C)** Heatplot showing differentially expressed

genes (increased) in HNP neurons (neurons in high NP co-expression group). **(D)** Gene

network plot showing results of functional enrichment for HNP neurons. **(E)** Venn diagram

showing the overlap between gene expressions higher in HNP neurons (neurons in high NP co-

expression group that express 6+ NPs) and significantly decreased in AD neurons co-

expressing 2-5 NPs (MNP neurons). The hypergeometric test was applied to evaluate the

overrepresentation of genes upregulated in control HNP neurons but notably reduced in AD

MNP neurons; α = 0.05. **(F)** Molecular signatures differentially increased and decreased in AD

MNP neurons.

Figure 3. Decreased expression of Alzheimer's disease-associated neuropeptides

(ADNPs) across brain regions during aging as well as cognitive continuum. (A) Brain

regions (medial sagittal view) showing decreased Alzheimer's disease-associated

neuropeptides (ADNPs) with aging. AMY, amygdala; CNG, anterior cingulate cortex; HIP,

hippocampus; BA9, Brodmann area 9. **(B)** Distribution of neurons based on the number of co-

expressed ADNPs from the entorhinal cortex (EC) of control (CT), mild cognitively impaired

(MCI), and AD donor brains. Median and third quantile of the number of co-expressed ADNPs

were marked by dashed lines for each condition. **(C)** The number of neurons above the median

and the third quantiles were plotted for CT, MCI, and AD. The decreasing trend of ADNP-co-

expressing neuronal count in the cognitive continuum was tested by Jonckheere-Terpstra test.

The post-hoc comparisons were tested by Yuen's two-sample t-test. **(D)** Combination of the

neuropeptide outputs (key ADNPs: SST, VGF, VIP, and CRH) derived from three previously

reported machine learning algorithms (color modification) [8] was used to examine the the number

of expressed key ADNPs, ranging from 0 to 3, in the EC of CT, MCI, and AD donor brains.

GLM, General Linear Model; SVM, Support Vector Machine; RF, Random Forest; *SST*,

Somatostatin gene; *VGF*, Nerve Growth Factor Inducible gene; *VIP*, Vasoactive Intestinal

Peptide gene; CRH, Corticotropin-Releasing Hormone gene.

Figure 4. Local density of ADNP-co-expressing high NP-producing (AHNP) neurons may

govern brain vulnerability to AD. (A) Top five brain and cortical regions ranked by AHNP

neuron abundance are visualized. **(B)** Categorization of AHNP neurons predominantly found in

EC by neurotransmitter, subtypes, and transferred MTG labels (common cell type

nomenclatures for the medial temporal gyrus of the mammalian brain).[22]

Consent Statement

Consent from human subjects was not necessary.

Funding

ML is supported by PAL's discretionary funds and the Doctoral Dissertation Fellowship from the Graduate School Fellowship Office at the University of Minnesota.

Author contributions

ML conceived the study. ML performed analyses and data visualization. ML and PAL discussed the results. ML wrote the manuscript. ML and PAL revised the manuscript. All authors approved the final manuscript.

Acknowledgments

Dr. Nicole Flack kindly reviewed the code employed for the analysis presented herein. Dr. Alice Larson provided helpful comments and suggestions that strengthened this manuscript. The data used for the analyses described in this manuscript were obtained from the GTEx Portal in October 2022. The Genotype-Tissue Expression (GTEx) Project was supported by the Common Fund of the Office of the Director of the National Institutes of Health and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. The results published here are in whole or in part based on data obtained from the AD Knowledge Portal. The data available in the AD Knowledge Portal would not be possible without the participation of research volunteers and the contribution of data by collaborating researchers. The results published here are in whole or in part based on data obtained from the AD Knowledge Portal (https://adknowledgeportal.org/). Study data were generated from postmortem brain tissue provided by the Religious Orders Study and Rush Memory and Aging Project (ROSMAP) cohort at Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago. This work was supported in part by the Cure Alzheimer's Fund, NIH grants AG058002, AG062377, NS110453, NS115064, AG062335, AG074003,

NS127187, MH119509, HG008155 (M.K.), RF1AG062377, RF1 AG054321, RO1 AG054012

(L.-H.T.) and the NIH training grant GM087237 (to C.A.B.). ROSMAP is supported by

P30AG10161, P30AG72975, R01AG15819, R01AG17917. U01AG46152, U01AG61356. ML is

supported by the Doctoral Dissertation Fellowship granted through the Graduate School

Fellowship Office at the University of Minnesota. PAL also provided discretionary funds

supporting ML. Figures were organized using BioRender (biorender.com).

Conflict of interest statement

All authors declare no conflict of interest.

Supporting information

Supplementary materials for this manuscript can be found at https://github.com/mancili/HNP.