Branch Length Transforms using Optimal Tree Metric Matching

Shayesteh Arasti,[1][†] Puoya Tabaghi,[2][†] Yasamin Tabatabaee,[3] Alan K. Mayer,[4] and Siavash Mirarab[5*]

[1]*Department of Computer Science and Engineering, UC San Diego, CA 92093, USA*
[2]*Halıcıoğlu Data Science Institute, UC San Diego, CA 92093, USA*
[3]*Department of Computer Science, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA*
[4]*Department of Mathematics, UC San Diego, CA 92093, USA*
[5]*Department of Electrical and Computer Engineering, UC San Diego, CA 92093, USA*
[*]*Corresponding author: smirarab@ucsd.edu*

Abstract

The abundant discordance between evolutionary relationships across the genome has rekindled interest in methods for comparing and averaging trees on a shared leaf set. However, compared to tree topology, where much progress has been made, handling branch lengths has been more challenging. Species tree branch lengths can be measured in various units, often different from gene trees. Moreover, rates of evolution change across the genome, the species tree, and specific branches of gene trees. These factors compound the stochasticity of coalescence times and estimation noise, making branch lengths highly heterogeneous across the genome. For many downstream applications in phylogenomic analyses, branch lengths are as important as the topology, and yet, existing tools to compare and combine weighted trees are limited. In this paper, we address the question of matching one tree to another, accounting for their branch lengths. We define a series of computational problems called Topology-Constrained Metric Matching (TCMM) that seek to transform the branch lengths of a query tree based on a reference tree. We show that TCMM problems can be solved efficiently using a linear algebraic formulation coupled with dynamic programming preprocessing. While many applications can be imagined for this framework, we explore two applications in this paper: embedding leaves of gene trees in Euclidean space to find outliers potentially indicative of errors, and summarizing gene tree branch lengths onto the species tree. In these applications, our method, when paired with existing methods, increases their accuracy at limited computational expense.

Keywords

Phylogenomics, Convex Optimization, Species Tree Branch Length, Outlier Detection, Distance-based phylogenetics

Code and Data availability

The software is available at https://github.com/shayesteh99/TCMM.git. Data is available on Github https://github.com/shayesteh99/TCMM-Data.git.

[†]These authors contributed equally to this work.

Introduction

Discordance among evolutionary histories across the genome (Maddison, 1997) is now understood to be pervasive and has motivated the study of region-specific histories (gene trees) and the speciation history (the species tree). Having alternative trees leads to a need for comparing and combining trees. We need metrics to quantify tree dissimilarity, algorithms for mapping gene trees to the species tree to reconcile them (Doyon et al., 2011), and median tree methods to combine gene trees into a single species tree (Barthélemy and McMorris, 1986). Solutions to these problems have been developed and are widely adopted. Measures of tree difference such as Robinson and Foulds (1981) (RF), quartet distance (Estabrook et al., 1985), and matching split (MS) (Bogdanowicz and Giaro, 2012) are commonly used to quantify gene tree differences. Biologists often use reconciliation methods in comparative genomics to understand the evolution of gene function. Inferring the species tree by finding the tree with the minimum total quartet distance to gene trees (quartet median tree) has been widely adopted due to its statistical consistency under several models of gene tree discordance (Mirarab et al., 2021).

The rich literature on comparing gene trees and species trees has focused on tree topology far more than branch lengths. Discordance, however, is not just limited to topology. One of the main causes of discordance is incomplete lineage sorting (ILS) as modeled by the multi-species coalescent (MSC) model (Degnan and Rosenberg, 2009). Under MSC, internal nodes of a gene tree correspond to coalescence events that predate species divergences. As a result, gene tree branch lengths are guaranteed to differ from the species tree even if there is no topological discordance (a phenomenon Edwards and Beerli (2000) referred to as species versus genic divergences). Beyond true biological differences, challenges in inferring gene trees compound the problem. From sequences alone, we can only measure branch lengths in the unit of the expected number of substitutions per site (SU), which are a function of not just divergence times but also substitution rates, which vary across the tree, across the genome, or both (Kumar, 2005; Lopez et al., 2002; Rasmussen and Kellis, 2007). Furthermore, gene trees are often inferred from short sequences (to avoid recombination within a gene), which makes gene tree estimates noisy, especially for short branches.

Branch lengths are as important as topology to many downstream applications such as dating, diversification studies, comparative genomics, trait evolution, and even microbiome analyses. Nevertheless, the myriad sources of heterogeneity (ILS, rate change, noise) have made it challenging to reconcile gene tree branch lengths with each other and with the species tree. Bayesian co-estimation methods (e.g., Boussau et al., 2013; Heled and Drummond, 2010; Liu, 2008) account for branch lengths by jointly modeling gene trees and species trees and introducing relevant parameters. These methods tend to be accurate (Yang and

⁶⁴ Warnow, 2011) and readily produce a reconciliation of species trees and gene trees. However, they do not

⁶⁵ scale to large datasets (Ogilvie et al., 2017; Zimmermann et al., 2014). Moreover, since complex rate models

⁶⁶ further reduce scalability, these methods often model only limited rate heterogeneity.

⁶⁷ The alternative two-step approach of inferring gene trees and then summarizing them to find the species

⁶⁸ tree often ignores branch lengths. Theoretically justifiable methods that consider branch lengths in inferring

⁶⁹ species trees (e.g., GLASS and STEM; Kubatko et al., 2009; Mossel and Roch, 2010) have been less accurate

⁷⁰ than topology-focused methods given estimated gene trees (DeGiorgio and Degnan, 2014; Yang and Warnow,

⁷¹ 2011), partially due to challenges mentioned earlier (Huang et al., 2010). Others have focused on gene tree

⁷² branch lengths (SU) for rooting them (Mai et al., 2017; Tria et al., 2017), for finding outliers indicative of

⁷³ errors (de Vienne et al., 2012; Mai and Mirarab, 2018), for clustering gene trees (Gori et al., 2016), and

⁷⁴ for summarizing them to infer the species tree branch lengths (Binet et al., 2016; Tabatabaee et al., 2023).

⁷⁵ These attempts have been more successful but lack an underlying mechanism for matching trees that can be

⁷⁶ used across applications. Each deals in some way with the underlying difficulty that two trees that differ in

⁷⁷ both topology and branch length are hard to compare since the same branches are not present in both (see

⁷⁸ Kuhner and Yamato (2015) for tree comparison metrics that include branch length).

⁷⁹ To arrive at a general method for comparing trees while accounting for branch lengths, we propose a

⁸⁰ framework that decouples topological changes and branch length changes to an adjustable extent. We use

⁸¹ a linear algebraic formulation based on patristic distances (pairwise path lengths) to define an optimiza-

⁸² tion problem called Topology-Constrained Metric Matching (TCMM), which is inspired by the least-square

⁸³ distance-based methods (Cavalli-Sforza and Edwards, 1967; Rzhetsky and Nei, 1991, 1992, 1993). The goal

⁸⁴ of TCMM is to assign branch lengths to a query tree such that its patristic distances match a set of one or

⁸⁵ more reference trees as much as possible without changing its topology. Using an additional regularization

⁸⁶ term, we can balance the competing goals of matching the reference tree and keeping the original branch

⁸⁷ length patterns of the query tree. All TCMM problems are convex, require preprocessing doable in quadratic

⁸⁸ time and memory, and are solvable using standard gradient descent techniques. While this ability to align

⁸⁹ the metric structure of one tree with another offers a powerful tool, in this paper, we focus on two particular

⁹⁰ applications: 1) embedding gene tree leaves in a shared space to enhance outlier detection, and 2) transfer-

⁹¹ ring gene tree branch lengths to a species tree. We show that in both applications, using TCMM to match

⁹² gene tree branch lengths, when paired with existing methods, can improve their accuracy.

Material and Methods

Problem formulation

For an unrooted tree $T = (V_T, E_T)$, we let L_T denote the leaf set and $P_T(v, v')$ denote the path between any pairs of vertices $v, v' \in V_T$. For a weighted tree, each edge $e \in E_T$ has length $w_T(e) \in \mathbb{R}^+$. We use $\mathcal{T}(T)$ to denote the set of all weighted trees with the same topology as tree T. A weighted tree T with N vertices indexed by $[N] = \{1, \ldots, N\}$ defines a metric $d_T : V_T \times V_T \to \mathbb{R}^+$ where $d_T(v, v') = \sum_{e \in P_T(v,v')} w(e)$. We define the *weight* vector $w_T = (w_T(e))_{e \in E_T} \in \mathbb{R}^{|E_T|}$ as a vector of all the weights indexed by edges and, with a slight abuse of notation, the *pairwise distance* vector $d_T = (d_T(v, v'))_{v,v' \in L_T, v < v'} \in \mathbb{R}^{\binom{|L_T|}{2}}$ as a vector of all patristic distances indexed by pairs of nodes. We define a binary *path matrix* A_T of size $\binom{|L_T|}{2} \times |E_T|$ where rows correspond to pairs of vertices in L_T and columns represent the edge set. For two distinct vertices $v, v' \in L_T$ and edge $e \in E_T$, an element $((v, v'), e)$ of A_T is one if and only if the edge e is on the path between v and v'. Otherwise, it is equal to zero. The path matrix A_T completely encodes the topology of T.

Consider a query tree T and a reference weighted tree R, both on a shared leafset L. For any tree T' with the same topology as T, let $\text{cost}(T'; R)$ measure the branch length discrepancy of T' to the reference tree R. For any such cost function, we can seek $T^* \in \arg\min_{T' \in \mathcal{T}(T)} \text{cost}(T'; R)$. Since the topology is fixed in this problem, we can reparameterize the cost with respect to the weight vector w_T to get $w^* \in \arg\min_{w'} \text{cost}(w'; R)$, where $w' \in \mathbb{R}^{+|E_T|}$ are constrained to be nonnegative. In this paper, we focus on a particular cost function to quantify the discrepancy of metric information between the two trees and successively refine it. Akin to least squares error distance-based tree estimation, we set

$$\text{cost}(w_T; R) = \sum_{v,v' \in L} |d_R(v, v') - d_T(v, v')|^2 = \|d_R - d_T\|_2^2$$

as the square of the ℓ_2 distance between pairwise distance vectors. As it has long been appreciated (Cavalli-Sforza and Edwards, 1967) (see also Desper and Gascuel, 2003), we can find a separable form for pairwise distances in terms of topology and edge weights. For $v, v' \in L$, we can write $d_T(v, v') = \sum_{e \in P_T(v,v')} w_T(e) = \langle a_{v,v'}, w_T \rangle$ where $a_{v,v'} = \left(1_{e \in P_T(v,v')}\right)_{e \in E_T} \in \{0, 1\}^{|E_T|}$ and 1_x is the indicator function. Note that $a_{v,v'}$ is a row of the path matrix A_T. Thus, for any tree $T = (V_T, E_T)$, we have $d_T = A_T w_T$, and

$$\text{cost}(w_T; R) = \sum_{v,v' \in L} |d_R(v, v') - d_T(v, v')|^2 = \|d_R - A_T w_T\|_2^2 \ . \tag{1}$$

We can now define an ordinary least squares (OLS) problem similar to Cavalli-Sforza and Edwards (1967)

120 (and formulated more clearly by Rzhetsky and Nei (1991)):

121 **Problem 1** (Topology-Constrained Metric Matching— under ℓ_2^2)**.** *Let T, R be weighted trees with edge*
122 *weights w_T and w_R on the common leafset L with at least two elements. The weights of the closest tree to*
123 *R matching T in topology with edge weights lower bounded by $\varepsilon \geq 0$ is:*

$$w^* \in \underset{w \in \mathbb{R}^{|E_T|}, w \geq \varepsilon}{\arg\min} \|d_R - A_T w\|_2^2.$$

The cost function in Problem 1 is agnostic to the original branch lengths of T, w_T, and thus, it can
dramatically alter them.   For many applications, it is desirable to retain the original branch lengths to
some user-adjustable extent. We can accommodate that goal by adding a regularizing term. For an optimal
edge weight vector w_{T*}, we define the *adjustment rate vector* by element-wise dividing w_{T*} and w_T, to get
w_{T*}/w_T. To incorporate w_T in the problem statement, we penalize the deviation of estimated w_{T*} from the
initial weight vector w_T, save for a total scaling factor shared by all branches, as in the following problem.

Problem 2 (Regularized Topology-Constrained Metric Matching — under ℓ_2^2)**.** *Let T, R be weighted trees*
with edge weight functions w_T and w_R on the common leafset L. Let λ be a fixed regularization coefficient
and $\sigma^2(w/w_T)$ denote the variance of the adjustment rates vector w/w_T. We seek the tree $T^* \in \mathcal{T}(T)$ with
edge weights

$$w_{T*} \in \underset{w \in \mathbb{R}^{|E_T|}, w \geq \varepsilon}{\arg\min} \|d_R - A_T w\|_2^2 + \lambda \sigma^2(w/w_T) \, . \tag{2}$$

The regularizing term, which is a quadratic function of the weight vector w, leads to lowered variance
in the adjustment rate vector, w/w_T, and thus, lowered change in *relative* branch lengths compared to T.
Clearly, if the variance of adjustment rate values is 0, the tree T has been simply scaled up or down, without
any real change to its relative branch lengths. Higher λ values encourage less change to the relative branch
lengths of T. As $\lambda \to \infty$, all rate multipliers converge to a solution that simply scales branch lengths of T
with a fixed global rate $r*$:

$$r^* \in \underset{r \cdot \min_e w_T(e) \geq \varepsilon}{\arg\min} \|d_R - r A_T w_T\|_2^2 \, .$$

143 To make λ values more interpretable across all datasets with varying sizes and heights, we first compute
144 the optimal branch lengths w^* according to Problem 1. We then reformulate Problem 2 as:

$$w_{T*} \in \underset{w \in \mathbb{R}^{|E_T|}, w \geq \varepsilon}{\arg\min} \frac{\|d_R - A_T w\|_2^2}{\text{cost}(w^*; R)} + \lambda \sigma^2(w/w_T).$$

where $\text{cost}(w^*; R)$ is defined in Eq. (1). This adjustment ensures that the first term is scale-free and provides
an initial solution that can be used as input for the optimizer. We will use this revised formulation of the

Regularized TCMM Problem in all experiments. We can also extend to multiple inputs.

Problem 3 (Topology-Constrained Consensus Metric Matching)**.** *Let $\{R_k : k \in [K]\}$ be a set of weighted*

reference trees, each with weights w_k. Given a tree T, we seek to find the optimal (adjusted) tree T^* with the

same topology as T and weights w_{T^*} such that:

$$
152 \qquad w_{T^*} \in \underset{\substack{w \in \mathbb{R}^{|E_R|} \\ w \geq \varepsilon}}{\arg\min} \sum_{k \in [K]} \alpha_k \|d_{R_k} - A_T w\|_2^2 + \lambda \sigma^2(w/w_T),
$$

153 *where α_k is a nonnegative weight controlling the contribution of reference tree R_k in the cost function, $\lambda > 0$*

154 *is a regularizing coefficient, $\sigma^2(w/w_T)$ penalizes the weights that result in a large variance for the adjustment*

155 *rate vector w/w_T, and $\varepsilon \geq 0$ is a lower bound for the estimated edge weights.*

156 Remark 1 (supplementary material) shows that problem 3 can be solved by solving Problem 2 after

157 averaging distances across k genes.

A Scalable Convex Optimization Solution

TCMM problems are all convex

160 We can rewrite the cost functions in Problems 1–3 in quadratic form. For simplicity, we use $A = A_T$ and

161 $d = d_R$. Note that Problem 1 minimizes $l_1(w) \stackrel{\text{def.}}{=} \|d - Aw\|_2^2$. We convert l_1 to the quadratic form

$$
162 \qquad l_1(w) = \frac{1}{2} w^\top B w + c^\top w + \text{const}, \tag{3}
$$

163 where $B = 2A^\top A$ is a positive semidefinite matrix, $c = -2A^\top d$, and $\text{const} = \|d\|_2^2$. Since $B \succeq 0$ by

164 construction, l_1 is a convex function of the weight parameter w. Adding the regularization term $\sigma^2(r)$ —

165 where $r = w/w_T$ — does not change the convexity, as we can convert it to the following quadratic form:

$$
166 \qquad l_2(w) \stackrel{\text{def.}}{=} \sigma^2(r) = \frac{1}{|E|}(r - 1\bar{r})^\top (r - 1\bar{r}) = \frac{1}{|E|} r^\top (P_1^\perp)^\top P_1^\perp r = \frac{1}{|E|} r^\top P_1^\perp r
$$

where $1 \in \mathbb{R}^{|E|}$ is the vector of all ones, $\bar{r}$ is a scalar that is the average value of elements in r, $P_1^\perp = I - \frac{1}{|E|} 11^\top$.

It is a known fact that $P_1^\perp$ is an orthogonal projection operator, that is, $(P_1^\perp)^\top P_1^\perp = P_1^\perp$. Furthermore, it is

a positive semidefinite matrix with $0 - 1$ eigenvalues. Therefore, the regularizing term $\sigma^2(w/w_T)$ is indeed a

convex function of the weight parameter w. Thus, Problems 1,2, and 3 all admit globally optimal solutions.

Proposition 1. *The cost functions in Problems 1, 2, and 3 are convex functions of w defined over nonempty,*

closed, and convex constraint sets. Hence, each optimization problem admits a global minimum (though not

necessarily a unique minimizer), and the projected gradient descent method converges to such a solution.

Proofs are provided in the supplementary material, Appendix A.5. By Proposition 1, we can use projected

gradient descent to solve Problems $1-3$ using the gradient terms shown in the proof. We implemented and

optimized the quadratic functions l_1 and l_2 using built-in functions in the Python package `CVXPY`. Our

empirical results suggest a sub-cubic running time ($\approx O(N^{2.74})$) for this implementation (Fig. S1a).

Dynamic programming enables calculating B and c in quadratic time

179 Our formulation does not need to ever compute A_T and d_R — whose dimensions grow cubically with N.

180 Rather, it needs the matrix B, which grows quadratically, and the vector c, which grows linearly with

181 N. Naively computing $A_T^\top A_T$ from its building blocks requires $O(N^4)$ time and $O(N^3)$ memory. Similarly,

182 computing $A_T^\top d_R$ naively requires $O(N^3)$ time and $O(N^3)$ memory. However, we can compute B in quadratic,

183 and c in sub-quadratic time and memory using dynamic programming without ever computing A_T and d_R.

Computing $c := A_T^\top d_R \in \mathbb{R}^{|E_T|}$. Bryant and Waddell (1998) propose a quadratic algorithm to compute $A_T^\top d$ for any arbitrary dissimilarity matrix d. However, in our case, since d_R comes from a tree, we can design a faster algorithm to compute $A_T^\top d_R$. Each edge $e \in E_T$ corresponds to a bipartition of leaves of T. For an arbitrary rooting of T, we define $C_T(e)$ as the set of leaves of T below e. Let e' and e'' be the children of e. As Bryant and Waddell (1998) noted, the element of c that corresponds to e is

$$
\begin{aligned}
c_e &= \sum_{v \in C_T(e)} \sum_{v' \in L_T \setminus C_T(e)} d_R(v, v') \\
&= c_{e'} + c_{e''} - 2 \sum_{v \in C_T(e')} \sum_{v' \in C_T(e'')} d_R(v, v')
\end{aligned}
$$

where the second equation gives a recursive algorithm. For an arbitrary distance matrix, the double-sum

in the recursion requires $O(N^2)$ over the entire tree. While $O(N^2)$ will be dominated by the optimization

step, we note that in Problem 3, that calculation of c will have to be repeated k times, which means without

further improvements, calculation of c could become dominant for large k. Luckily, when d_R is defined on a

given R, we can do better. We have designed an algorithm (see Algorithm S1) to compute that term using

$|C_T(e') + C_T(e'')|$ leaf-to-root traversals on R. Each traversal takes $O(H_R)$, where H_R is the height of R,

and using the recursive equation above, we need exactly $N \cdot H_T$ such traversals, where H_T is the height

of T. Therefore, the total running time of this algorithm is $O(N \cdot H_T \cdot H_R)$. While this algorithm can be

as slow as $O(N^3)$ for unbalanced trees, it would require $O(N \log^2(N))$ for sufficiently balanced trees, often

encountered in practice. Our empirical results (Fig. S1b) show that on average it is sub-quadratic (close

to $O(N^{1.4})$). Note also that for unbalanced input (defined as $H_T \cdot H_R > N$), we switch to the quadratic

algorithm of Bryant and Waddell (1998), thus ensuring the running time is never worse than quadratic.

In the consensus Problem 3, if no distances are missing, we can simply average the terms $\{A_T^\top d_{R_k}\}_{k\in[K]}$ instead of averaging distance vectors $\{d_{R_k}\}_{k\in[K]}$. The consensus cost function allows this:

$$\sum_{k\in[K]} \alpha_k\|d_{R_k} - A_T w\|_2^2 + \lambda\sigma^2(w/w_T) = w^\top A_T^\top A_T w\Big(\sum_{k\in[K]}\alpha_k\Big) - 2\Big(\sum_{k\in[K]}\alpha_k\big(A_T^\top d_{R_k}\big)^\top\Big)w + \lambda\sigma^2(w/w_T) + \text{const}$$

Computing $B = A_T^\top A_T$. Given the definition of the path matrix A_T, it can be easily deduced that for two edges e and e', $B_{e,e'}$ is the number of pairs of leaves $v, v' \in L$ where $e, e' \in P_T(v, v')$. We easily pre-compute $|C_T(e)|$ for each edge $e \in E_T$ in linear time and the least common ancestor (LCA) of all pairs of nodes of T in quadratic time. With these, we can easily compute $B_{e,e'}$ for all $e, e' \in E_T$ in quadratic time using simple calculations given in Algorithm S1 (supplementary material), as confirmed by our empirical analysis (Fig. S1c).

We have assumed so far that query and reference trees have the same species, but this will not always be the case. Gene trees almost always miss some taxa due to gene loss or data incompleteness. Leaves of the reference tree R that are absent from the query tree T do not create a problem as they can be easily pruned from R without affecting the optimization problem. Leaves in T missing from R need care. Appendix A.4 details how missing data are handled in our approach, and Figure S2 provides evidence of its effectiveness.

Application 1: Outlier in gene trees using MCOA

The phylogenomic analysis pipeline introduces various forms of error into gene trees, compounding biological processes such as ILS that create real gene tree discordance. Some errors leave subtle traces, while others distort the inferred trees more dramatically by introducing abnormally long branches or drastically altering the tree topology. These errors can impact downstream analyses (Laurin-Lemay et al., 2012; Philippe et al., 2017; Salichos and Rokas, 2013; Springer and Gatesy, 2018). As a result, several efforts have been made to detect anomalies in gene trees automatically. Mai and Mirarab (2018) proposed the TreeShrink method to identify species with unexpectedly long branches in a tree. Gori et al. (2016) proposed clustering gene trees using various distance metrics. Going further, de Vienne et al. (2012) introduced Phylo-MCOA to use multiple co-inertia analysis (MCOA) to embed all gene trees into a single Euclidean space such that pairwise distances of leaves match those of the input trees as much as possible. Using the principal coordinates of this space enables the detection of outlier genes and species in a two-step process employing Tukey's method (Hubert and Vandervieren, 2008) followed by more complex steps. A recent implementation of Phylo-MCOA is available in an R package called PhylteR (Comte et al., 2023).

Figure 1: (S100-default Dataset) (a) The TCMM pipeline for outlier detection. Each modified gene tree G'_i is the output of running TCMM with the original gene tree G_i as the query and the rest of the gene trees as the reference trees. The modified gene trees are then fed into the PhylteR pipeline, where the multiple co-inertia analysis (MCOA) is performed. Lastly, the principal coordinates (PCs) for each modified gene tree are used for the outlier detection and visualization tasks. (b) An example of the outliers detected by TCMM+PhylteR for one replicate. Left: the true species tree with the two species 37 and 52 and their corresponding subtree highlighted. Right: two outlier genes (16 and 17) for these species. Gene 17 is an outlier due to discordance from the true species tree topology, whereas gene 16 is an outlier due to a long branch (marked in red) separating species 37 and 52 from other species. (c) The PCA plot shows the projections of six species onto the first and second principal coordinates (PCs) with different regularization coefficient (λ) values. The position and the status (outlier or non-outlier) of genes 16 and 17 are marked for species 37 and 52 on each plot. (d) Comparing the Euclidean distance to the median species for true gene trees. (See also Fig. S3).

₂₂₄ We propose to use TCMM to improve PhylteR outlier detection (Fig. 1a). We examine an example

₂₂₅ based on the simulated 100-taxon dataset (see *Datasets*) to motivate the approach (Fig. 1b-d). A species

₂₂₆ tree can have rate variation across branches, including highly diverged clades (Fig. 1b). Additionally, even

₂₂₇ in the absence of error, individual gene trees can have highly discordant topologies (e.g., Gene 17, Fig. 1b) or

₂₂₈ branch lengths (e.g., Gene 16) due to heterotachy (Lopez et al., 2002). These differences can make pairwise

₂₂₉ distances across different gene trees dissimilar and render embeddings noisy (Fig. 1c). In particular, branch

₂₃₀ length variations can lead to outliers even absent any topological discordance (e.g., Gene 16 in Fig. 1c).

₂₃₁ TCMM, by matching gene trees to a shared metric space, reduces the distance of the embeddings to the

₂₃₂ median (consensus) embedding (Fig. 1d). The more concentrated distributions have the potential to better

₂₃₃ reveal *topological* outliers. Moreover, λ defines the degree to which gene trees are matched to the shared

₂₃₄ space, allowing us to identify different types of outliers. With $\lambda = 0$, we match gene trees maximally to

₂₃₅ each other, reducing the noise created by rate heterogeneity both across and within gene trees, resulting in

₂₃₆ more concentrated embeddings and revealing outliers that would otherwise be obscured by noise in the data

₂₃₇ (e.g., Gene 17 becomes an outlier with $\lambda = 0$ in Fig. 1c). As λ increases, the ratios between original branch

₂₃₈ lengths are increasingly maintained (average rate heterogeneity across gene trees is reduced by TCMM). As

₂₃₉ a result, outliers with unexpectedly long branches are detected (e.g., Gene 16 in Fig. 1b,c). A λ value in

₂₄₀ between the two extremes trades off the two aspects.

₂₄₁ We evaluate TCMM's performance on both simulated and biological datasets, where gene trees have

₂₄₂ branch lengths measured in units of expected substitutions per site (SU). We apply the unweighted consen-

₂₄₃ sus version of TCMM to each gene tree, using it as the query tree and treating the remaining gene trees as

₂₄₄ reference trees. The modified gene trees are then fed into the PhylteR pipeline to generate Euclidean embed-

₂₄₅ dings, which are used for visualization and outlier detection. For simplicity, we skip the second step of outlier

₂₄₆ detection used in Phylo-MCOA and report only the outliers identified through Tukey's method. We apply

₂₄₇ this procedure with and without regularization and examine how varying $\lambda \in \{0, 1e-5, 1e-4, \ldots, 10, 100\}$

₂₄₈ influences the outliers detected. The same process is repeated with the original gene trees without running

₂₄₉ TCMM. To measure the effectiveness of outlier detection, we calculate the improvement in the normalized

₂₅₀ RF (nRF) distance from gene trees to the species tree S, compared to changes in nRF by random removal of

₂₅₁ the same number of taxa, leading to a metric called ΔRF (see Appendix A.2). We also use the normalized

₂₅₂ quartet score (nqs), defined as the proportion of shared quartets between two trees, to quantify the similar-

₂₅₃ ity between filtered gene trees and the species tree. We use the true species tree when available (simulated

₂₅₄ datasets) and the ASTRAL-III inferred species tree otherwise. We quantify whether outliers are the same as

₂₅₅ those found by the branch length-based method TreeShrink by reporting the containment Jaccard similarity

₂₅₆ (Appendix A.2) between the outliers detected by TCMM +PhylteR and those detected by TreeShrink. To

evaluate the effect of outlier removal in downstream analyses, we use the filtered gene trees to infer the

species tree using ASTRAL-IV (Zhang et al., 2025). For topology, we infer a species tree from the filtered

gene trees and compare it to the true species tree using normalized RF distance (nRF). We also assess the

accuracy of CU branch lengths, which are a direct function of gene tree discordance (Sayyari and Mirarab,

2016). In this analysis, we fix the species tree topology to the true topology and use the filtered gene trees

to estimate CU branch lengths, expecting underestimation due to errors. We report mean absolute error

(MAE) and bias, defined as the mean difference between estimated and true branch lengths.

Application 2: Species tree branch lengths

Two-step methods of species tree inference produce a species tree topology from a set of gene trees. To be

useful for downstream analyses (e.g., dating), the species tree requires branch lengths. Scalable and accurate

methods such as ASTRAL (Mirarab et al., 2014b) and NJst/ASTRID (Liu and Yu, 2011; Vachaspati and

Warnow, 2015) produce only topology. We can assign coalescent unit (CU) branch lengths for internal

branches of the species tree using gene tree topologies alone, but this approach can lead to systematic

underestimation (Forthman et al., 2022; Sayyari and Mirarab, 2016), and the lack of terminal branch lengths

makes it less useful. Since gene trees have SU branch lengths and many downstream analyses require SU

lengths, we can seek to transfer the SU branch lengths from gene trees onto the species tree. This should be

possible because gene tree branch lengths are a function of species tree SU branch lengths (Fig. 2a). Briefly,

we can assume each branch of the species tree has a substitution rate, and gene tree branches inherit these

rates as well as per-locus rate multipliers (Mallo et al., 2016).

Binet et al. (2016) proposed a method called ERaBLE that uses a least squares optimization applied to

patristic distances of the gene trees to estimate species tree lengths, effectively taking a weighted average.

ERaBLE allows for rate heterogeneity only across loci. It estimates a single scalar rate per gene tree by

matching them simultaneously to the species tree. Tabatabaee et al. (2023) showed that ERaBLE overes-

timates terminal branches because it ignores deep coalescence. They propose a coalescent-based method

called CASTLES to explicitly model the differences between the species and genic divergence times. While

CASTLES and its newer version CASTLES-Pro are more accurate than ERaBLE in MSC simulations, their

accuracy can decrease under high levels of HGT (Tabatabaee et al., 2025). This observation raises the ques-

tion: Can a flexible model-free method assign SU lengths to the species tree while attaining high accuracy

when gene tree heterogeneity is not solely due to ILS?

Here, we use TCMM to summarize gene tree SU branch lengths onto the species tree (Fig. 2b). Unlike

ERaBLE, which estimates only one scalar parameter per gene tree, we assign one parameter for each branch

Figure 2: (a) Assumed data generation model, where species tree branch lengths measured in the unit of generations is multiplied by per-branch mutation rates to obtain SU length. Gene trees inherit these rates, with further per-locus variation. (b) The TCMM pipeline for branch length estimation with the species tree topology as the query and the gene trees with branch lengths in substitution unit (SU) as the reference trees. The final output is the species tree with SU branches.

of each gene tree. Counterintuitively, we use gene trees as the reference and the species tree as the query in one of two ways. One approach, called consensus, uses Problem 3 to jointly match the species tree (as query T) to multiple gene trees (as the reference R). Here, we simply set all the α_k weights to 1. Our second (and default) approach is to run TCMM per gene (i.e., Problem 1), and summarize results at the end. Results provide us with a set of estimated branch lengths $\tilde{w}_{e,k}$ where $e \in E_S$ and $G_k \in \mathcal{G}$. These lengths can then be further summarized into a single value $\hat{w}_e$ for each branch using common averaging methods — arithmetic mean, harmonic mean, and geometric mean. When using arithmetic means, this approach is similar to using the consensus method; Remark 2 (supplementary material) shows that the two are identical if all output lengths are strictly positive. Despite this (near) equivalence, the per-gene approach offers the benefit of enabling outlier removal. We remove both extremely small outliers introduced by TCMM and large outliers, which can arise due to gene tree errors, using the procedures detailed in Appendix A.3. We address missing data using the approaches noted in Appendix A.4.

We can pair TCMM with other branch length estimation tools. In this approach, we take the species tree with branch lengths estimated by another tool (e.g., CASTLES-Pro (Tabatabaee et al., 2025)), and apply the regularized version of TCMM (Problem 2) to this tree. This allows us to refine the input while preserving its branch lengths to an extent controlled by λ. One challenge is selecting the parameter λ. Here, we use an automatic approach to select λ from a predefined set of options. We do this by choosing the λ that minimizes the sum of the estimated species tree branch lengths. This choice is rooted in the minimum-evolution principle (Rzhetsky and Nei, 1993) used by a family of distance-based approaches.

We evaluate TCMM for this application using both simulated and real datasets. On simulated data,

the model species tree branches have true SU lengths (w) obtained by multiplying their lengths (number of generations) with varying mutation rates, creating non-ultrametric trees. We compare the true w values to estimated values $\hat{w}$ to quantify the mean absolute error ($|w - \hat{w}|$) denoted MAE, absolute logarithmic error ($|\log(\hat{w}/w)|$), and bias ($\hat{w} - w$), averaged across all branches of each species tree after excluding the outgroup. We compare TCMM to four leading methods studied by Tabatabaee et al. (2025), all run on the true species tree topology. We include CASTLES-Pro, which accounts for coalescence times predating species divergence times. We examine the commonly used method of running RAxML (Stamatakis, 2006) on concatenated gene alignments to estimate branch lengths of a fixed species tree (referred to as Concat+RAxML in our experiments). We also include two methods that like TCMM use patristic distances. A simple alternative is averaging patristic distance matrices across gene trees and using a distance-based method (FastME, Lefort et al., 2015) to assign branch lengths to S (called Patristic(AVG)+FastME in our experiments). A more advanced version of this idea is ERaBLE (default), mentioned earlier.

Simulated Datasets

For application 1, we studied two simulated datasets, one with introduced errors and one without, and five biological datasets. For application 2, we studied four simulated datasets with different levels of gene tree estimation error, ILS, horizontal gene transfer (HGT), and rate heterogeneity. We also studied three microbial datasets. All biological datasets are detailed in the results section.

S200-perturbed dataset. We started from a 201-taxon Simphy-simulated dataset originally published by Mirarab and Warnow (2015) and perturbed its sequences to introduce errors to study Application 1. We focused on the condition with medium ILS (i.e., 2 million generations and a speciation rate of $1e-7$), which contains 50 replicates, each with 1000 gene trees. We introduced errors as follows. For each replicate, we perform $N + 1$ rounds of perturbation, with $N \sim \text{Poisson}(3)$, leading to at least one round and four rounds on average. In each round $0 \le r \le N$, we select $k_r \sim \text{Poisson}(\lambda = 40)$ genes to be modified uniformly at random, and choose the proportion of taxa to be perturbed by drawing $t_r \sim \text{Beta}(10, 90)$ (mean: 0.1, sd: ≈ 0.03). Instead of selecting these perturbed (i.e., erroneous) species ($\mathcal{E}_r$) uniformly at random, we use a random sampling strategy that encourages $\mathcal{E}_r$ to be closer to each other on the tree than a uniformly random sampling would lead to. To do so, we root the species tree on a randomly chosen taxon and repeatedly traverse from the root to leaves, taking left/right children at random, and add any observed leaf to the set $\mathcal{E}_r$; we repeat until $|\mathcal{E}_r| = \lceil 200 \times t_r \rceil$. To perturb these selected taxa, we first randomly reroot the species tree. Then, for each perturbed gene $1 \le i \le k_r$, we select a "source" taxon $s_{r,i}$ using the same random sampling process used to select $\mathcal{E}_r$, and rerooting the tree at this source between genes. For each gene i,

we then draw $\rho_{r,i} \sim Beta(\alpha = 16, \beta = 13)$ (mean 0.55, sd: 0.09) and replace the first $\rho_{r,i}$ portion of each target sequence in $\mathcal{E}_r$ with the same sites from $s_{r,i}$ in gene i. Thus, each target sequence becomes a chimera between the $s_{r,i}$ and its original sequence. This procedure leads to zero to 81 perturbations per gene (3.2 in expectation: $4 \times 40 \times 0.1 \times 200 \times {}^1/_{1000}$). We then re-infer a gene tree using FastTree-II (Price et al., 2010), given these perturbed sequences. We removed one replicate because the estimated gene trees (prior to introducing errors) contained numerous large polytomies. For each replicate, we used the first 100 gene trees in our analyses to reduce the accuracy of species trees and make the impacts on them more visible.

S100 simulated dataset. We reused the published 101-taxon 1000-gene dataset (S100-default) simulated by Zhang et al. (2018) using Simphy (Mallo et al., 2016), which Tabatabaee et al. (2023) then updated to include species trees with SU branch lengths (see Table S1 for all parameters). Our earlier results, shown as motivation for Application 1, use this dataset, and we use it in Application 2 as well. To recap the simulation procedure, briefly, ultrametric species trees with generation times are simulated using a birth-death model; to get SU length, each branch length is multiplied by a fixed mutation rate across the tree (drawn from a LogNormal distribution with mean 4×10^{-8}) and a rate multiplier drawn from a Gamma distribution (mean: 1; inverse of variance, α, drawn from a LogNormal(1.5,1), leading to variance ranging in our replicates between 0.0153 to 2.6027). The 50 replicates of this dataset are highly heterogeneous in terms of ILS, with the mean nRF distance between the model species tree and the true gene trees at 46%. Gene trees were estimated using FastTree-2 (Price et al., 2010) from 200bp, 400bp, 800bp, and 1600bp alignments, leading to average nRF distance between true and estimated gene trees of 55%, 42%, 31%, and 23%, respectively. In application 1, we only use the first 100 gene trees for each replicate due to computational constraints and to make differences between methods more pronounced.

To tease out the impact of ILS levels and rate heterogeneity modes in Application 2, we used the updated Simphy code to simulate six model conditions of the same S100 dataset (naming them S100-truegt), each condition with 50 replicates (Table S1). In these analyses, intended for validation and sanity checks, rather than biological realism, we used true gene trees; therefore, no sequences were generated. Simphy parameters were kept the same, except for the following changes. For varying ILS, we created three model conditions by changing the population size (low, med, high ILS: N_e = 4e+4, 4e+5, and 4e+6, resp.) and turned off lineage-specific rate variation so that the species tree becomes ultra-metric in SU units. Note that each condition still retains some level of difference in ILS levels across replicates due to differing tree heights. The average nRF distance between true gene trees and the true species tree for low, med, and high ILS model conditions is 8%, 44%, and 89%, respectively. For studying the impact rate heterogeneity, we created three conditions. We refer to the original simulation as `hl+hg+hs`; this condition has rate multipliers per species,

per gene, and per individual branches of individual gene trees. We added `hl+hg` which has a rate multiplier per gene and one multiplier per each branch of the gene tree (`hl+hg`); however, there is no global per species rate, making the species three ultrametric. We also added `hl+hs`, which assigns a rate multiplier per species tree branch, making the tree non-ultrametric, and also assigns a rate per gene.

Results

Application 1: topology focused MCOA

S200-perturbed Simulated Dataset

Using TCMM with PhylteR increases the accuracy of error detection compared to using PhylteR alone for some values of λ (e.g., 0.1 and 1), as reflected in higher true positive rates (TPR) and improved ROC curves (Fig. 3a). The PhylteR threshold k allows us to control sensitivity, with lower k corresponding to increased total number of outliers. Across all k, the Area Under the Curve (AUC) increases from 0.811 with PhylteR alone to 0.884 with PhylteR+TCMM with $\lambda = 0.1$. However, while patterns of performance remain similar across k, choices of k with low false positive rate (FPR) are of particular interest. As default, we use $k = 1.5$, which provides a low FPR and substantial TPR (e.g., FPR = 0.008 and TPR = 0.425 for $\lambda = 0.1$). On this dataset, TreeShrink (with default threshold) has a substantially lower TPR and higher FPR (FPR = 0.026 and TPR = 0.006).

Comparison to PhylteR depends strongly on the choice of λ, with low values (e.g., $\lambda = 0$) showing worse performance than PhylteR alone, and very high values (e.g., $\lambda = 10$) showing only modest improvements over PhylteR. However, for the middle range ($\lambda = 0.1$ or $\lambda = 1$), the improvements are substantial. In this range, the number of removed outliers (roughly 3 per gene tree) is about the same as PhylteR without TCMM (Fig. 3b), showing that the improvements of TCMM are not due to finding more outliers but rather finding more erroneous sequences. With lower λ values, PhylteR+TCMM removes many more topological outliers (e.g., 6.8 with $\lambda = 0$), which leads to increased similarity between the outlier-pruned gene trees and both the true and estimated species trees (Fig. 3c). However, these reductions in discordance come at the cost of removing many more data points (Fig. 3b) and higher FPR (Fig. 3a). To test if the reduction in signal due to pruning by each PhylteR+TCMM variant is justified by their reduction in noise, we used the resulting pruned gene trees to estimate the species tree topology (Fig. 3d) and CU branch lengths (Fig. 3e). For topology, in most cases, removing outliers *increases* the species tree error in terms of nRF distance (Fig. 3d). The only case with (slight) improvement in topological accuracy is $\lambda = 1$, which is among the values that best trades off precision and recall. All the other settings, including PhylteR without TCMM,

Figure 3: (S200-perturbed Dataset) (a) ROC curve for PhylteR+TCMM outlier detection as we change its threshold k (dots). Colors show different inputs to PhylteR. Black shows the original gene trees, whereas other colors are TCMM gene trees with different λ values. $k = 1.5$ is marked with a larger size. Dashed line shows random outlier removal. Area under the curve (AUC) values are shown in the caption. (b) Number of remaining taxa in gene trees after outlier removal. (c) Normalized quartet score of pruned and unfiltered gene trees to the true and estimated species trees. Above the green line shows improvement compared to unfiltered gene trees. (d,e) Species tree accuracy after outlier removal, showing nRF distance of estimated tree topology (d) and mean absolute error (MAE) and bias of estimated coalescent unit (CU) branch lengths (e) to the true species tree. Below the green line shows improvement compared to unfiltered gene trees. Similarity of MAE and bias indicates that the error is due to a systematic under-estimation of lengths, which is reduced after filtering.

result in increased topological error in ASTRAL trees, perhaps because the ASTRAL topology has some level of robustness to errors, and the loss of signal due to mistaken removal of correct sequences (false positives) can outweigh the inclusion of some noise.

Unlike topology, CU branch lengths are highly sensitive to error (Sayyari and Mirarab, 2016) and thus improve consistently as we remove more outliers (Fig. 3e). Any level of filtering, with or without TCMM, reduced error, but the largest reduction was obtained by TCMM+PhylteR with the smallest λ values. The distinct patterns observed between topology and branch length reflect the higher sensitivity of CU branch lengths to gene tree error.

Simulated Dataset (S100-default Dataset)

On the S100-default dataset, which contains no sequence errors (only the typical gene tree estimation error), both PhylteR and TCMM+PhylteR still detect some outliers. However, the number of outliers detected by TCMM+PhylteR on this dataset is substantially lower than on the S200-perturbed dataset (Fig. S3b). For example, with $\lambda = 0.1$, 0.11% of taxa are removed across all conditions of S100-default compared to 1.46%

for S200-perturbed with the same λ. Pairing TCMM with PhylteR results in fewer removals for $\lambda > 0.01$

but not for lower λ (Fig. S3b). While removing these outliers tends to decrease discordance between the

gene trees and the true species tree, these improvements are not substantial (no more than 2%), likely

due to the small number of outliers (Fig. S3c). We observe that decreasing λ makes the outliers detected

by TCMM+PhylteR increasingly dissimilar to those detected by TreeShrink, and the reduced emphasis on

branch lengths leads to the detection of more topologically aberrant outliers (Fig. S4a), a pattern we referred

to earlier with one example (Fig. 1b). Moreover, compared to randomly removing the same number of leaves,

outliers detected by TCMM+PhylteR yield a greater reduction in nRF distance for lower values of λ, with

this pattern observed across all levels of gene tree error and more pronounced in higher-quality gene trees.

These observations are consistent with our expectation regarding the role of λ in shaping the type of outliers

detected by our method (see *Methods*).

Biological Dataset

We next studied six biological datasets (mammals, frogs, plants, and two datasets on Xenacoelomorpha,

Insects) adopted from Mai and Mirarab (2018). These datasets cover a range of number of species, number

of genes, and evolutionary depth (Appendix A.1). Some patterns observed in the simulated datasets persist.

For example, As λ decreases, MCOA embeddings become more concentrated and closer to the median

embedding for each species (Fig. S5). Other patterns differ from the simulated datasets. The impact of λ on

the number of outliers depends on the dataset, with some, like mammals, behaving like simulated data and

others, like insects and plants, showing a reduction in the number of outliers after applying TCMM (Fig. 4a).

The total number of outliers also varies across datasets, ranging from 0.01 on the Xenacoelomorpha dataset

of Rouse et al. (2016) ($\lambda = 1$) to 3.16 for the Frogs dataset ($\lambda = 0.001$).

Removing outliers identified by TCMM+PhylteR generally increases the similarity to the estimated

species tree for most, but not all, datasets (Fig. 4b). The amount of improvement tends to be higher when

the original gene trees are less discordant with the estimated species tree. In particular, for mammals and

frogs, which had low discordance, TCMM+PhylteR improves quartet scores more than using PhylteR alone

with any λ, and more so with lower λ values. For plants and insects, some λ values are better than PhylteR

alone, while others are not, but all cases lead to improved quartet scores compared to no filtering. In the

case of the Xenacoelomorpha datasets of Cannon et al. (2016), pairing TCMM with PhylteR can reduce

improvements compared to PhylteR alone though it still results in improvements. The Xenacoelomorpha

dataset of Rouse et al. (2016), however, was different; removing outliers makes the gene trees more dissimilar

to the species tree with TCMM+PhylteR but not with PhylteR alone or high λ. The gene trees in this

dataset have the highest discordance with the estimated species tree among all the studied biological datasets

Figure 4: (Biological Dataset) (a) Number of remaining taxa in gene trees after outlier removal for each λ. The amount of missing taxa from each dataset is written as a percentage. (b) Normalized quartet score to the ASTRAL estimated species tree. The unfiltered original gene trees are shown in green.

(Fig. 4b), suggesting that gene tree quality and species tree accuracy may impact the effectiveness of outlier detection using TCMM.

Application 2: Species tree branch lengths

Simulated Data

Choosing a version of TCMM

The per-gene approach with arithmetic mean exhibits lower bias compared to the consensus version of TCMM without compromising on MAE (Fig. S6a), making it the preferred choice. Another advantage of the per-gene approach is its ability to perform outlier detection and removal, effectively eliminating extremely small estimated branch lengths that are often present in the consensus approach. In fact, the consensus method can assign branch lengths that are essentially zero, particularly for smaller λ values. However, these underestimations are mitigated in the per-gene method due to outlier removal (Fig. S6bd). Incorporating outlier removal also reduces bias and MAE in terminal branches while slightly increasing MAE for internal branches (Fig. S6a). The consensus approach of TCMM is the most similar to the distance-based method ERaBLE, both in methodology and performance (Fig. S6c). Thus, the increased performance compared to ERaBLE is primarily attributed to the combination of per-gene branch length estimates and outlier detection.

While all three methods (consensus, per-gene, and per-gene with outlier removal) perform similarly when initial branch lengths are available, the per-gene method with outlier removal is adopted as the default version of TCMM due to its performance in the absence of initial branch lengths and the advantages of

⁴⁶⁵ outlier removal in biological datasets. Furthermore, examining varying λ reveals that as λ values increase,

⁴⁶⁶ branch lengths estimated by the default TCMM method become progressively more similar to the initial

⁴⁶⁷ branch lengths (CASTLES-Pro) in terms of their relative values (Fig. S6bd).

Comparison to other methods

⁴⁶⁹ On the S100-truegt dataset, when ILS levels are kept low, all methods run on the true gene trees have

⁴⁷⁰ near-perfect accuracy (Fig. S7). However, when ILS increases, the error goes up for all methods, leading to

⁴⁷¹ TCMM outperforming other distance-based methods but not CASTLES-Pro. Focusing on medium ILS levels

⁴⁷² and changing modes of rate heterogeneity does not change the relative accuracy of methods (Fig. S8). When

⁴⁷³ comparing conditions with ultrametric species trees (`hl+hg`) to non-ultrametric cases (`hl+hs`), we observe

⁴⁷⁴ increased error for all methods, but not to the same degree. TCMM 's normalized MAE increases by 10%,

⁴⁷⁵ while ERaBLE and AVG+FastME show increase by 27% or more (comparisons to `hl+hs+hg` are similar).

⁴⁷⁶ Thus, TCMM is more robust to lineage-specific rate heterogeneity than other distance-based methods, likely

⁴⁷⁷ because methods such as ERaBLE account primarily for genome-wide rate variation.

⁴⁷⁸ On the S100-default dataset (with heterogeneous ILS, simulated sequences, and estimated gene trees),

⁴⁷⁹ TCMM is better than both the commonly used concatenation method (Concat+RAxML) and distance-based

⁴⁸⁰ methods (ERaBLE and AVG+FastME) but not the coalescent-based CASTLES-Pro in terms of overall mean

⁴⁸¹ absolute error and bias (Fig. S9). The improvements are due to much better terminal branch calculation; for

⁴⁸² internal branches, TCMM has a slight disadvantage compared to both distance-based methods (ERaBLE

⁴⁸³ and AVG+FastME), but not concatenation. Combining CASTLES-Pro with TCMM (using automatic λ) is

⁴⁸⁴ substantially better than TCMM alone, yet it does not outperform CASTLES-Pro. This result confirms that

⁴⁸⁵ CASTLES-Pro, which is specifically designed and optimized for ILS, outperforms cause-agnostic distance-

⁴⁸⁶ based methods when gene tree discordance is dominated by ILS.

⁴⁸⁷ The patterns shift in simulations with HGT, where the best method is combining TCMM and CASTLES-

⁴⁸⁸ Pro with automatic λ selection (Fig. 5). The estimation error of all methods increases as the rate of HGT

⁴⁸⁹ increases, especially for the two highest HGT rates. CASTLES-Pro remains better than TCMM alone and

⁴⁹⁰ other methods. TCMM is better than methods other than CASTLES-pro for terminal branches and slightly

⁴⁹¹ worse for internal ones. All methods are biased, and these biases increase as the rate of HGT increases.

⁴⁹² CASTLES-Pro has an underestimation bias, while the other methods have an overestimation bias, which is

⁴⁹³ concentrated on terminal branches. TCMM has the lowest overestimation bias among methods other than

⁴⁹⁴ TCMM+CASTLES-Pro (e.g., on average 50.4% lower bias than concatenation).

⁴⁹⁵ On this dataset, however, TCMM+CASTLES-Pro has the best accuracy for all measures of error and

⁴⁹⁶ bias. On average, the mean absolute error of CASTLES-Pro+TCMM is 8.2% lower than CASTLES-Pro

Figure 5: (HGT Dataset) Comparing TCMM with the alternative methods across various rates of HGT (x-axis). TCMM is run with $\lambda = 0$ (TCMM) and automatic λ selection (CASTLES-Pro+TCMM). The methods are compared using two error metrics (mean absolute error and mean absolute log error) as well as bias.

and 27.3% lower than TCMM alone. The clearest advantage is the reduction in bias, particularly at higher HGT rates: While TCMM reduces bias by 26.7% compared to CASTLES-Pro across all HGT rates, the TCMM+CASTLES-Pro combination achieves reductions of 27.3% and 42.1% over TCMM alone at HGT rates of 2×10^{-7} and 5×10^{-7}, respectively. These results establish TCMM+CASTLES-Pro as a reliable choice for running TCMM across all tested datasets and model conditions (Fig. S10).

No single λ consistently outperforms others across all settings (Fig. S10). This variability makes it useful to select a λ automatically, eliminating the need for manual tuning. Automatic λ selection tends to choose larger λ values (Fig. S11), likely due to TCMM's overestimation bias, which causes smaller λ values to result

Figure 6: (a), (b) The distribution of the TCMM ($\lambda = 0$) estimated length for the branch separating bacteria and archaea per gene tree. Vertical dashed lines show the estimates of TCMM, CASTLES-Pro+TCMM, CASTLES-Pro, and concatenation. The range of the estimated branch lengths for the AB branch by CASTLES-Pro+TCMM for different values of λ is shown in green. (a) Uses the 49 core genes, with $\lambda = 1$ automatically selected for CASTLES-Pro+TCMM; (b) uses the 38 non-ribosomal genes, with $\lambda = 100$. (c), (d) (Zhu et al. (2019) Dataset). (c) The distribution of the outlier genes removed by TCMM ($\lambda = 0$) per branch for different variance improvement thresholds; the default threshold (5) and the AB branch under this threshold are highlighted in red and green, respectively. (d) The distribution of the TCMM estimated AB branch length per gene tree.

in overall higher estimated branch lengths. Note that even with high λ, CASTLES-Pro+TCMM benefits from outlier detection and removal, a feature absent in CASTLES-Pro alone.

Biological data: AB branch length

We analyzed several microbial datasets, focusing on the separation between bacteria and archaea (AB branch). Moody et al. (2022) claimed that lower divergences obtained by some recent studies (Petitjean et al., 2015; Zhu et al., 2019) were the result of highly abundant HGT in microbial datasets (Arnold et al., 2022). They compared results with two alternative datasets of core genes, with less propensity for HGT. Following them, we reanalyzed a 72-taxon tree (Williams et al., 2020) with 49 core genes, including ribosomal proteins and other conserved genes, and a 108-taxon dataset (Petitjean et al., 2015) with 38 genes that only include non-ribosomal proteins. We compared results with a reanalyses of the microbial dataset by Zhu et al. (2019) with 10,575 species sampled evenly from bacteria and archaea and 381 genes (including many non-core genes). The species tree used in this analysis was inferred using ASTRAL and is rooted on the branch separating bacteria from archaea. Gene trees in this dataset lack 28% of taxa on average. On all

datasets, we compare the length of the AB branch inferred using TCMM, CASTLES-Pro, and concatenation, all drawn on species topologies built using ASTRAL-III (Zhang et al., 2018) from the two gene sets. On these datasets, ERaBLE fails to return an output, and AVG+FastME produces an estimated tree with several long negative branches around the root; thus, we exclude these methods.

Using 49 core genes (including ribosomal genes), CASTLES-Pro, TCMM, and CASTLES-Pro+TCMM (automatic λ selected as 1) estimate a length of 1.79, 1.94, and 1.91 SU for the AB branch, respectively, compared to 1.43 SU for concatenation. In contrast, on the 38 non-ribosomal genes (presumably with more HGT), the AB branch is much shorter, as noted by Moody et al. (2022). TCMM and CASTLES-Pro+TCMM (with λ selected as 100) estimate AB branch length to be 0.87 and 0.76, respectively, which are higher than concatenation (0.43) but lower than CASTLES-Pro (1.05). The observation that both gene tree-based methods yield higher branch lengths than concatenation for non-ribosomal genes is consistent with Moody et al. (2022)'s findings that concatenation can underestimate branch length in the presence of HGT. The substantial difference between ribosomal and non-ribosomal genes observed using methods that account for gene tree discordance confirms the fundamentally different patterns of evolution between the two. In per-gene mode, TCMM estimates a separate branch length for each gene tree (Fig. 6 a,b). An examination of the distribution reveals substantial diversity among genes, with some indicating an AB branch length of more than 4 SU, while others suggest a length below 0.5 SU.

On the microbial dataset by Zhu et al. (2019), CASTLES-Pro, TCMM, and CASTLES-Pro+TCMM (λ fixed to 1 due to high running time) estimate branch lengths of 2.68, 1.89, and 1.24 SU, respectively, for the AB branch. As a result, both CASTLES-Pro and TCMM estimate drastically longer branch lengths than concatenation, which estimates 0.15 SU when selecting 100 random sites per gene and 0.12 SU when selecting the 100 most conserved sites (original paper selected 100 sites per locus due to computational limitations of concatenation with > 10000 tips) Once again, the distribution of estimated AB branch lengths across individual gene trees shows significant variation, ranging from as low as 10^{-9} to over 18 SU for some gene trees (Fig. 6d). In computing the final estimate, the outlier removal step of the per-gene method removes 283 out of 381 estimates before summarization (281 low outliers and 2 high outliers) This high level of outlier detection may seem excessive, a topic we will return to in the discussions. On this dataset many branches have large numbers of outliers removed, presumably due to rampant HGT. Compared to all branches in the tree, the AB branch is among those with the highest proportions of outliers (Fig. 6c).

Discussion

We introduce TCMM, a versatile framework for reconciling the branch lengths of a query tree to match one or more reference trees. At its core, TCMM provides a way to match tree metric spaces, thereby decreasing branch length discordance between trees without changing topological discordance. The underlying optimization problem builds on the least-squares formulation long used in distance-based phylogenetics, with added regularization terms. The regularization provides a way to adjust how far the adjustment goes. However, with or without regularization, the output branch lengths do not automatically have a biological meaning; instead, they are intended to serve as input in downstream applications. Our paper focuses on two applications: outlier detection and species tree branch length estimation. In both applications, combining TCMM with existing methods improves accuracy in many but not all conditions.

In the outlier detection application, we use TCMM to match gene trees to each other by harmonizing their metric spaces so that the remaining outliers reflect topological differences. Without such harmonization, embeddings of tree leaves might not align due to divergent rates of evolution in entire genes or some branches of certain genes, rather than errors in data. Our results show that pairing TCMM with PhylteR improves the accuracy of outlier detection compared to both PhylteR and TreeShrink, and that removing these outliers can improve downstream analyses – particularly in settings that are more sensitive to outliers, such as branch length estimation. The regularization term λ helps us control the extent to which cases of substantial rate variation are marked as outliers. For example, with true gene trees, where the only sources of discordance are ILS and rate heterogeneity, we saw that changing λ directly controls whether topological or branch length outliers are detected (Fig. S4a). When simulated sequence data had errors, and in biological data, anomalies may simultaneously affect both topology and branch lengths, blurring the distinction between the two (Fig. S4c). Thus, in practice, it may be helpful to examine a range of λ values. Beyond detecting errors in the data, this type of harmonization may also prove useful in other applications, such as detecting changes in rates of evolution. Note that TCMM outputs a multiplier for every branch of every gene tree, and examining the distributions of these multipliers may prove useful in detecting patterns of rate variation across the tree and across loci.

Our second application matches gene trees to a species tree instead of matching them together to transfer SU branch length from gene trees to the species tree. While TCMM is not as accurate as CASTLES-Pro (which is designed based on the MSC model) under ILS-only conditions, TCMM provides several benefits. Foremost, in its per-gene mode, TCMM provides a length *distribution* for each species tree branch. By outputting a multiplier per branch per gene, it allows for across-genes heterotachy (Lopez et al., 2002), a

feature missing from other methods. ERaBLE uses only one mutation rate per gene, making it far less versatile. CASTLES-Pro models rate heterogeneity across the species tree and overall rate changes across genes; however, it does not estimate a per-gene rate for each species tree branch. Our biological results show evidence (Fig. 6) of heterotachy. Even when outliers are excluded, the remaining loci still provide a range of lengths for the AB branch and other branches across the microbial tree of life, pointing to heterotachy.

The second advantage of TCMM for species tree branch length estimation lies in outlier detection. With input gene trees generated using models assumed by CASTLES-Pro, we observe no advantage in pairing it with TCMM. In contrast, in the presence of HGT, which is not modeled by CASTLES-Pro, we observe a lower bias when CASTLES-Pro is followed by TCMM. HGT creates branch length patterns that can widely vary from those of HGT-free genes, a notion that has been used by some previous methods of HGT detection (e.g., Dessimoz et al., 2008). The outlier detection step can eliminate gene trees with widely incongruent topologies, which, when matched to the species tree, would otherwise lead to extremely short branch lengths. We observe many such cases (283 out of 381) for the AB branch length in the microbial dataset. By omitting these genes, TCMM+CASTLES-Pro is able to estimate a much higher AB length compared to concatenation.

Labeling the majority of genes as outliers, as TCMM does for the AB branch, may seem counterintuitive: How can more data be classified as outliers than not? This point gets at the heart of the debate about the AB branch. By removing enough horizontally transferred genes, we can estimate high values for the AB length, as Moody et al. (2022) argued. Concatenation (or even TCMM without outlier removal) estimates much shorter lengths given a large number of marker genes. Is removing many (or most) genes biologically justifiable? The answer depends on how we interpret the branch lengths and what we use them for. In one view, despite HGT, there is an underlying tree (vertically inherited), providing the backbone of the network representing the full evolutionary history. Then, to find the number of substitutions on a branch of the tree, we should remove horizontally transferred genes, which add not just noise but an underestimation bias. Thus, we do not consider "outliers" merely as data points that deviate from the majority for unknown reasons. Rather, in this case, the high topological discordance due to HGT results in short estimated lengths on the backbone species tree (near zero for TCMM). Thus, the primary role of outlier detection (responsible for removing 281 out of 283 outliers for the AB branch) is to eliminate these near-zero estimates, which are prevalent due to the high levels of HGT around the branch. In a different view, if horizontal evolution is pervasive and concentrated enough to render the backbone tree an inadequate model, one might instead aim to estimate an overall genome-wide level of divergence between two groups of taxa (e.g., archaea and bacteria). In such cases, removing large parts of the genome is less defensible. Regardless, by providing a distribution per branch, TCMM offers a way to better represent the full evolutionary history. A promising direction of future research is to investigate whether these distributions can be incorporated into downstream

applications such as UniFrac calculation (Lozupone and Knight, 2005).

A practical question arises: Should CASTLES-Pro be followed by TCMM for branch length estimation? The answer, based on our findings, is nuanced. If we believe the cause of discordance is mostly incomplete lineage sorting, the answer seems to be no. Additionally, TCMM in its current form is not equipped to handle multi-copy gene trees Chaudhary et al. (2013)— a potential avenue for future development. However, in scenarios involving single-copy gene trees with substantial levels of HGT or other aberrant sources of discordance (e.g., caused by hidden paralogy, alignment errors, or missing data), TCMM can enhance accuracy by detecting and removing outliers. Since homology errors abound in phylogenomic datasets (Springer and Gatesy, 2018; Steenwyk et al., 2023) and outlier loci can leave an outsized impact (Shen et al., 2017), running TCMM with CASTLES-Pro as input to detect the outliers seems prudent in most analyses.

The TCMM framework can have wide-ranging applications beyond those explored in this study. Once the species tree branch lengths are estimated and perhaps calibrated to the unit of time, TCMM can also be used to match gene trees to the species tree (opposite of our approach here) for downstream applications such as visualization (e.g., using Bouckaert (2010)), estimation of average rates, and perhaps even dating individual gene trees, topics that can be explored in the future. The linear algebraic approach employed here effectively compares two trees by matching one to the other. Unlike many commonly used phylogenetic metrics, this matching incorporates both tree topology and branch lengths. Thus, the results of the matching could potentially be translated into new metrics of tree distance and new median tree problems. Future research should investigate these metrics and whether they provide additional insights beyond those offered by existing methods.

Acknowledgment

This work was supported by the National Institutes of Health (1R35GM142725) grant to S.M. This work used Expanse at San Diego Supercomputing Center through allocation ASC150046 from the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program, which is supported by U.S. National Science Foundation grants #2138259, #2138286, #2138307, #2137603, and #2138296.

References

Arnold, B. J., Huang, I.-T., and Hanage, W. P. (2022). Horizontal gene transfer and adaptive evolution in bacteria. *Nature Reviews Microbiology*, 20(4):206–218.

Barthélemy, J.-P. and McMorris, F. R. (1986). The median procedure for n-trees. *Journal of Classification*,
3(2):329–334. Publisher: Springer.

Binet, M., Gascuel, O., Scornavacca, C., P. Douzery, E. J., and Pardi, F. (2016). Fast and accurate branch
lengths estimation for phylogenomic trees. *BMC Bioinformatics*, 17(1):23.

Bogdanowicz, D. and Giaro, K. (2012). Matching split distance for unrooted binary phylogenetic trees.
IEEE/ACM Transactions on Computational Biology and Bioinformatics.

Bouckaert, R. R. (2010). DensiTree: Making sense of sets of phylogenetic trees. *Bioinformatics*, 26(10):1372–
1373. ISBN: 1367-4803.

Boussau, B., Szöllősi, G. J., and Duret, L. (2013). Genome-scale coestimation of species and gene trees.
Genome Research, 23(2):323–330.

Bryant, D. and Waddell, P. (1998). Rapid evaluation of least-squares and minimum-evolution criteria on
phylogenetic trees. *Molecular Biology and Evolution*, 15(10):1346–1346.

Cannon, J. T., Vellutini, B. C., Smith, J. r., Ronquist, F., Jondelius, U., and Hejnol, A. (2016). Xenacoelo-
morpha is the sister group to nephrozoa. *Nature*, 530(7588):89–93.

Cavalli-Sforza, L. L. and Edwards, A. W. (1967). Phylogenetic analysis. Models and estimation procedures.
American journal of human genetics, 19(3 Pt 1):233–57. Publisher: Elsevier.

Chaudhary, R., Burleigh, J. G., and Fernández-Baca, D. (2013). Inferring species trees from incongruent
multi-copy gene trees using the Robinson-Foulds distance. *Algorithms for Molecular Biology*, 8:28. arXiv:
1210.2665.

Comte, A., Tricou, T., Tannier, E., Joseph, J., Siberchicot, A., Penel, S., Allio, R., Delsuc, F., Dray, S., and
de Vienne, D. M. (2023). PhylteR: Efficient Identification of Outlier Sequences in Phylogenomic Datasets.
Molecular Biology and Evolution, 40(11):msad234.

de Vienne, D. M., Ollier, S., and Aguileta, G. (2012). Phylo-mcoa: a fast and efficient method to detect
outlier genes and species in phylogenomics using multiple co-inertia analysis. *Mol Biol Evol*, 29(6):1587–
1598.

DeGiorgio, M. and Degnan, J. H. (2014). Robustness to Divergence Time Underestimation When Inferring
Species Trees from Estimated Gene Trees. *Systematic Biology*, 63(1):66–82.

Degnan, J. H. and Rosenberg, N. A. (2009). Gene tree discordance, phylogenetic inference and
the multispecies coalescent. *Trends in Ecology and Evolution*, 24(6):332–340. ISBN: 0169-5347
Publisher: Department of Human Genetics, University of Michigan, Ann Arbor, MI 48109, USA.
j.degnan@math.canterbury.ac.nz.

Desper, R. and Gascuel, O. (2003). Theoretical Foundation of the Balanced Minimum Evolution Method of

Phylogenetic Inference and Its Relationship to Weighted Least-Squares Tree Fitting. *Molecular Biology and Evolution*, 21(3):587–598. ISBN: 0737-4038 (Print).

Dessimoz, C., Margadant, D., and Gonnet, G. H. (2008). DLIGHT – Lateral Gene Transfer Detection Using Pairwise Evolutionary Distances in a Statistical Framework. In Vingron, M. and Wong, L., editors, *Research in Computational Molecular Biology*, volume 4955, pages 315–330. Springer Berlin Heidelberg, Berlin, Heidelberg. Series Title: Lecture Notes in Computer Science.

Doyon, J.-P. J., Ranwez, V., Daubin, V., and Berry, V. (2011). Models, algorithms and programs for phylogeny reconciliation. *Briefings in Bioinformatics*, 12(5):392–400.

Edwards, S. and Beerli, P. (2000). Perspective: gene divergence, population divergence, and the variance in coalescence time in phylogeographic studies. *Evolution*, 54(6):1839–1854.

Estabrook, G. F., McMorris, F. R., and Meacham, C. A. (1985). Comparison of Undirected Phylogenetic Trees Based on Subtrees of Four Evolutionary Units. *Systematic Biology*, 34(2):193–200.

Feng, Y.-J., Blackburn, D. C., Liang, D., Hillis, D. M., Wake, D. B., Cannatella, D. C., and Zhang, P. (2017). Phylogenomics reveals rapid, simultaneous diversification of three major clades of gondwanan frogs at the cretaceous–paleogene boundary. *Proceedings of the National Academy of Sciences*, 114(29):E5864–E5870.

Forthman, M., Braun, E. L., and Kimball, R. T. (2022). Gene tree quality affects empirical coalescent branch length estimation. *Zoologica Scripta*, 51(1):1–13.

Gatesy, J. and Springer, M. S. (2014). Phylogenetic Analysis at Deep Timescales: Unreliable Gene Trees, Bypassed Hidden Support, and the Coalescence/Concatalescence Conundrum. *Molecular phylogenetics and evolution*, 80:231–266. Publisher: Elsevier Inc.

Gori, K., Suchan, T., Alvarez, N., Goldman, N., and Dessimoz, C. (2016). Clustering Genes of Common Evolutionary History. *Molecular Biology and Evolution*. arXiv: 1510.02356 ISBN: 1537-1719 (Electronic)\r0737-4038 (Linking).

Heled, J. and Drummond, A. J. (2010). Bayesian inference of species trees from multilocus data. *Molecular Biology and Evolution*, 27(3):570–580.

Huang, H., He, Q., Kubatko, L. S., and Knowles, L. L. (2010). Sources of Error Inherent in Species-Tree Estimation: Impact of Mutational and Coalescent Effects on Accuracy and Implications for Choosing among Different Methods. *Systematic Biology*, 59(5):573–583. ISBN: 1076-836X (Electronic)\r1063-5157 (Linking).

Hubert, M. and Vandervieren, E. (2008). An adjusted boxplot for skewed distributions. *Computational Statistics and Data Analysis*, 52(12):5186–5201.

Kubatko, L. S., Carstens, B. C., and Knowles, L. L. (2009). STEM: species tree estimation using maximum

likelihood for gene trees under coalescence. *Bioinformatics*, 25(7):971–973.

Kuhner, M. K. and Yamato, J. (2015). Practical performance of tree comparison metrics. *Systematic Biology*.

ISBN: 1063-5157.

Kumar, S. (2005). Molecular clocks: four decades of evolution. *Nature Reviews Genetics*, 6(8):654–662.

Laurin-Lemay, S., Brinkmann, H., and Philippe, H. (2012). Origin of land plants revisited in the light of

sequence contamination and missing data. *Current Biology*. ISBN: 1879-0445 (Electronic)\r0960-9822

(Linking).

Lefort, V., Desper, R., and Gascuel, O. (2015). FastME 2.0: A comprehensive, accurate, and fast distance-

based phylogeny inference program. *Molecular Biology and Evolution*, 32(10):2798–2800. ISBN: 1537-1719

(Electronic)\r0737-4038 (Linking).

Liu, L. (2008). BEST: Bayesian estimation of species trees under the coalescent model. *Bioinformatics*,

24(21):2542–2543. ISBN: 1367-4803 Publisher: Musuem of Comparative Zoology, Harvard University,

Cambridge, MA, USA. lliu@oeb.harvard.edu.

Liu, L. and Yu, L. (2011). Estimating Species Trees from Unrooted Gene Trees. *Systematic Biology*,

60(5):661–667. ISBN: 1063-5157\r1076-836X.

Lopez, P., Casane, D., and Philippe, H. (2002). Heterotachy, an Important Process of Protein Evolution.

Molecular Biology and Evolution, 19(1):1–7.

Lozupone, C. and Knight, R. (2005). UniFrac : a New Phylogenetic Method for Comparing Microbial

Communities UniFrac : a New Phylogenetic Method for Comparing Microbial Communities. *Applied and*

environmental microbiology, 71(12):8228–8235.

Maddison, W. P. (1997). Gene Trees in Species Trees. *Systematic Biology*, 46(3):523–536.

Mai, U. and Mirarab, S. (2018). TreeShrink: fast and accurate detection of outlier long branches in collections

of phylogenetic trees. *BMC Genomics*, 19(S5):272. ISBN: 9783319679785.

Mai, U., Sayyari, E., and Mirarab, S. (2017). Minimum variance rooting of phylogenetic trees and implications

for species tree reconstruction. *PLOS ONE*, 12(8):e0182238. Publisher: Public Library of Science ISBN:

1111111111.

Mallo, D., De Oliveira Martins, L., and Posada, D. (2016). SimPhy : Phylogenomic Simulation of Gene,

Locus, and Species Trees. *Systematic Biology*, 65(2):334–344.

Mirarab, S., Bayzid, M. S., Boussau, B., and Warnow, T. (2014a). Statistical binning enables an accurate

coalescent-based estimation of the avian tree. *Science*, 346(6215):1250463.

Mirarab, S., Bayzid, M. S., and Warnow, T. (2016). Evaluating Summary Methods for Multilocus Species

Tree Estimation in the Presence of Incomplete Lineage Sorting. *Systematic Biology*, 65(3):366–380. Pub-

lisher: Oxford University Press ISBN: 1063-5157.

Mirarab, S., Nakhleh, L., and Warnow, T. (2021). Multispecies Coalescent: Theory and Applications in

Phylogenetics. *Annual Review of Ecology, Evolution, and Systematics*, 52(1):247–268.

Mirarab, S., Reaz, R., Bayzid, M. S., Zimmermann, T., Swenson, M. S., and Warnow, T. (2014b). ASTRAL:

genome-scale coalescent-based species tree estimation. *Bioinformatics*, 30(17):i541–i548.

Mirarab, S. and Warnow, T. (2015). ASTRAL-II: Coalescent-based species tree estimation with many

hundreds of taxa and thousands of genes. *Bioinformatics*, 31(12):i44–i52.

Misof, B., Liu, S., Meusemann, K., Peters, R. S., Donath, A., Mayer, C., Frandsen, P. B., Ware, J., Flouri,

T., Beutel, R. G., Niehuis, O., Petersen, M., Izquierdo-Carrasco, F., Wappler, T., Rust, J., Aberer, A. J.,

Aspöck, U., Aspöck, H., Bartel, D., Blanke, A., Berger, S., Böhm, A., Buckley, T. R., Calcott, B., Chen,

J., Friedrich, F., Fukui, M., Fujita, M., Greve, C., Grobe, P., Gu, S., Huang, Y., Jermiin, L. S., Kawahara,

A. Y., Krogmann, L., Kubiak, M., Lanfear, R., Letsch, H., Li, Y., Li, Z., Li, J., Lu, H., Machida, R.,

Mashimo, Y., Kapli, P., McKenna, D. D., Meng, G., Nakagaki, Y., Navarrete-Heredia, J., Ott, M., Ou, Y.,

Pass, G., Podsiadlowski, L., Pohl, H., von Reumont, B. M., Schütte, K., Sekiya, K., Shimizu, S., Slipinski,

A., Stamatakis, A., Song, W., Su, X., Szucsich, N. U., Tan, M., Tan, X., Tang, M., Tang, J., Timelthaler,

G., Tomizuka, S., Trautwein, M., Tong, X., Uchifune, T., Walzl, M. G., Wiegmann, B. M., Wilbrandt,

J., Wipfler, B., Wong, T. K. F., Wu, Q., Wu, G., Xie, Y., Yang, S., Yang, Q., Yeates, D. K., Yoshizawa,

K., Zhang, Q., Zhang, R., Zhang, W., Zhang, Y., Zhao, J., Zhou, C., Zhou, L., Ziesmann, T., Zou, S., Li,

Y., Xu, X., Zhang, Y., Yang, H., Wang, J., Wang, J., Kjer, K. M., and Zhou, X. (2014). Phylogenomics

resolves the timing and pattern of insect evolution. *Science*, 346(6210):763–767.

Moody, E. R., Mahendrarajah, T. A., Dombrowski, N., Clark, J. W., Petitjean, C., Offre, P., Szöllősi, G. J.,

Spang, A., and Williams, T. A. (2022). An estimate of the deepest branches of the tree of life from ancient

vertically evolving genes. *eLife*, 11:e66695.

Mossel, E. and Roch, S. (2010). Incomplete lineage sorting: consistent phylogeny estimation from multiple

loci. *IEEE/ACM Transactions on Computational Biology and Bioinformatics (TCBB)*, 7(1):166–171.

Ogilvie, H. A., Bouckaert, R. R., and Drummond, A. J. (2017). StarBEAST2 Brings Faster Species Tree

Inference and Accurate Estimates of Substitution Rates. *Molecular Biology and Evolution*, 34(8):2101–

2114. ISBN: 0737-4038.

Petitjean, C., Deschamps, P., López-García, P., and Moreira, D. (2015). Rooting the domain archaea by

phylogenomic analysis supports the foundation of the new kingdom proteoarchaeota. *Genome biology and*

evolution, 7(1):191–204.

Philippe, H., Brinkmann, H., Copley, R. R., Moroz, L. L., Nakano, H., Poustka, A. J., Wallberg, A., Peterson,

K. J., and Telford, M. J. (2011). Acoelomorph flatworms are deuterostomes related to xenoturbella. *Nature*, 470(7333):255–258.

Philippe, H., Vienne, D. M. d., Ranwez, V., Roure, B., Baurain, D., and Delsuc, F. (2017). Pitfalls in supermatrix phylogenomics. *European Journal of Taxonomy*.

Price, M. N., Dehal, P. S., and Arkin, A. P. (2010). FastTree-2 – Approximately Maximum-Likelihood Trees for Large Alignments. *PLoS ONE*, 5(3):e9490. Publisher: Public Library of Science.

Rasmussen, M. D. and Kellis, M. (2007). Accurate gene-tree reconstruction by learning gene- and species-specific substitution rates across multiple complete genomes. *Genome Research*, 17(12):1932–1942. ISBN: 1088-9051.

Robinson, D. and Foulds, L. (1981). Comparison of phylogenetic trees. *Mathematical Biosciences*, 53(1-2):131–147.

Rouse, G. W., Wilson, N. G., Carvajal, J. I., and Vrijenhoek, R. C. (2016). New deep-sea species of xenoturbella and the position of xenacoelomorpha. *Nature*, 530(7588):94–97.

Rzhetsky, A. and Nei, M. (1991). A simple method for estimating and testing minimum-evolution trees. *Molecular Biology and Evolution*, 9.

Rzhetsky, A. and Nei, M. (1992). Statistical properties of the ordinary least-squares, generalized least-squares, and minimum-evolution methods of phylogenetic inference. *J Mol Evol*, 35(4):367–375.

Rzhetsky, A. and Nei, M. (1993). Theoretical foundation of the minimum-evolution method of phylogenetic inference. *Molecular Biology and Evolution*, 10(5):1073–1095.

Salichos, L. and Rokas, A. (2013). Inferring ancient divergences requires genes with strong phylogenetic signals. *Nature*, 497(7449):327–31. Publisher: Nature Publishing Group ISBN: doi:10.1038/nature12130.

Sayyari, E. and Mirarab, S. (2016). Fast Coalescent-Based Computation of Local Branch Support from Quartet Frequencies. *Molecular Biology and Evolution*, 33(7):1654–1668.

Sayyari, E., Whitfield, J. B., and Mirarab, S. (2017). Fragmentary gene sequences negatively impact gene tree and species tree reconstruction. *Molecular Biology and Evolution*, 34(12):3279–3291.

Shen, X.-x., Hittinger, C. T., and Rokas, A. (2017). Contentious relationships in phylogenomic studies can be driven by a handful of genes. *Nature Ecology & Evolution*, 1(5):0126. Publisher: Macmillan Publishers Limited, part of Springer Nature.

Song, S., Liu, L., Edwards, S. V., and Wu, S. (2012). Resolving conflict in eutherian mammal phylogeny using phylogenomics and the multispecies coalescent model. *Proceedings of the National Academy of Sciences*, 109(37):14942–14947.

Springer, M. S. and Gatesy, J. (2018). On the importance of homology in the age of phylogenomics. *Sys-*

tematics and Biodiversity, 16(3):210–228.

Stamatakis, A. (2006). RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses with thousands

of taxa and mixed models. *Bioinformatics*, 22(21):2688–2690.

Steenwyk, J. L., Li, Y., Zhou, X., Shen, X.-X., and Rokas, A. (2023). Incongruence in the phylogenomics

era. *Nature Reviews Genetics*, 24(12):834–850.

Tabatabaee, Y., Zhang, C., Arasti, S., and Mirarab, S. (2025). Species tree branch length estimation despite

incomplete lineage sorting, duplication, and loss. *bioRxiv*. urlhttps://doi.org/10.1101/2025.02.20.639320.

Tabatabaee, Y., Zhang, C., Warnow, T., and Mirarab, S. (2023). Phylogenomic branch length estimation

using quartets. *Bioinformatics*, 39(Supplement_1):i185–i193.

Tria, F. D. K., Landan, G., and Dagan, T. (2017). Phylogenetic rooting using minimal ancestor deviation.

Nature Ecology & Evolution, 1(7):0193.

Vachaspati, P. and Warnow, T. (2015). ASTRID: Accurate Species TRees from Internode Distances. *BMC*

Genomics, 16(Suppl 10):S3. Publisher: BioMed Central Ltd.

Wang, C. and Xiu, N. (2000). Convergence of the gradient projection method for generalized convex mini-

mization. *Computational optimization and applications*, 16(2):111–120.

Wickett, N. J., Mirarab, S., Nguyen, N., Warnow, T., Carpenter, E., Matasci, N., Ayyampalayam, S.,

Barker, M. S., Burleigh, J. G., Gitzendanner, M. A., Ruhfel, B. R., Wafula, E., Der, J. P., Graham, S. W.,

Mathews, S., Melkonian, M., Soltis, D. E., Soltis, P. S., Miles, N. W., Rothfels, C. J., Pokorny, L., Shaw,

A. J., DeGironimo, L., Stevenson, D. W., Surek, B., Villarreal, J. C., Roure, B., Philippe, H., dePamphilis,

C. W., Chen, T., Deyholos, M. K., Baucom, R. S., Kutchan, T. M., Augustin, M. M., Wang, J., Zhang, Y.,

Tian, Z., Yan, Z., Wu, X., Sun, X., Wong, G. K.-S., and Leebens-Mack, J. (2014). Phylotranscriptomic

analysis of the origin and early diversification of land plants. *Proceedings of the National Academy of*

Sciences, 111(45):E4859–E4868.

Williams, T. A., Cox, C. J., Foster, P. G., Szöllősi, G. J., and Embley, T. M. (2020). Phylogenomics provides

robust support for a two-domains tree of life. *Nature ecology & evolution*, 4(1):138–147.

Yang, J. and Warnow, T. (2011). Fast and accurate methods for phylogenomic analyses. *BMC Bioinformat-*

ics, 12(Suppl 9):S4. Publisher: BioMed Central Ltd.

Zhang,   C.,   Nielsen,   R.,   and   Mirarab,   S.   (2025).   ASTER: A   Package   for   Large-Scale

Phylogenomic   Reconstructions.   *Molecular   Biology   and   Evolution*,   42(8):msaf172.   _eprint:

https://academic.oup.com/mbe/article-pdf/42/8/msaf172/63779203/msaf172.pdf.

Zhang, C., Rabiee, M., Sayyari, E., and Mirarab, S. (2018). ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. *BMC Bioinformatics*, 19(S6):153.

Zhu, Q., Mai, U., Pfeiffer, W., Janssen, S., Asnicar, F., Sanders, J. G., Belda-Ferre, P., Al-Ghalith, G. A.,

Kopylova, E., McDonald, D., Kosciolek, T., Yin, J. B., Huang, S., Salam, N., Jiao, J.-Y., Wu, Z., Xu,

Z. Z., Cantrell, K., Yang, Y., Sayyari, E., Rabiee, M., Morton, J. T., Podell, S., Knights, D., Li, W.-J.,

Huttenhower, C., Segata, N., Smarr, L., Mirarab, S., and Knight, R. (2019). Phylogenomics of 10,575

genomes reveals evolutionary proximity between domains bacteria and archaea. *Nature Communications*,

10(1):5477.

Zimmermann, T., Mirarab, S., and Warnow, T. (2014). BBCA: Improving the scalability of *BEAST using

random binning. *BMC genomics*, 15(Suppl 6):S11.