A neural geometry theory comprehensively explains apparently

conflicting models of visual perceptual learning

Yu-Ang Cheng[1,2], Mehdi Sanayei[3,†], Xing Chen[4], Ke Jia[5,6,7], Sheng Li[8,9,10,11], Fang Fang[8,9,10,11],

Takeo Watanabe[2], Alexander Thiele[3], Ru-Yuan Zhang[1]

[1]Institute of Psychology and Behavioral Science and Shanghai Mental Health Center, Shanghai Jiao Tong University, Shanghai, 200030, China.
[2]Department of Cognitive, Linguistic and Psychological Sciences, Brown University, RI, USA.
[3]Biosciences Institute, Newcastle University, Framlington Place, Newcastle upon Tyne NE2 4HH, UK.
[4]Department of Ophthalmology, University of Pittsburgh, 1622 Locust St, Pittsburgh 15239, USA.
[5]Department of Neurobiology, Affiliated Mental Health Center & Hangzhou Seventh People,
[6]Liangzhu Laboratory, MOE Frontier Science Center for Brain Science and Brain-machine Integration, State Key Laboratory of Brain-Machine Intelligence, Zhejiang University, Hangzhou, China.
[7]NHC and CAMS Key Laboratory of Medical Neurobiology, Zhejiang University, Hangzhou, China.
[8]School of Psychological and Cognitive Sciences, Peking University, Beijing, 100871, China.
[9]Beijing Key Laboratory of Behavior and Mental Health, Peking University, Beijing, 100871, China.
[10]PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, 100871, China.
[11]Key Laboratory of Machine Perception (Ministry of Education), Peking University, Beijing, 100871, China.
[†]Present address: School of Cognitive Sciences, Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

Yu-Ang Cheng (iD) https://orcid.org/0009-0004-0145-1660
Ru-Yuan Zhang (iD) https://orcid.org/0000-0002-0654-715X

Corresponding:
Ru-Yuan Zhang
Zhongyuan 112, Shanghai Jiao Tong University
1954 Huashan Road, Xuhui District
Shanghai, China. 200030
Phone: +86-15674496106
Email: ruyuanzhang@gmail.com

ABSTRACT

Visual perceptual learning (VPL), defined as long-term improvement in a visual task, is considered

a crucial tool for elucidating underlying visual and brain plasticity. However, the identification of

a unified theory of VPL has long been controversial. Multiple existing models have proposed

diverse mechanisms, including improved signal-to-noise ratio, changes in tuning curves, and

reduction of noise correlations, as major contributors to improved neural representations

associated with VPL. However, each model only accounts for specific aspects of the empirical

findings, and there exists no theory that can comprehensively explain all empirical results. Here,

we argue that all neural changes at single units can be conceptualized as geometric transformations

of population response manifolds in a high-dimensional neural space. This approach enables

conflicting major models of VPL to be quantitatively tested and compared within a unified

computational theory. Following this approach, we found that changes in tuning curves and noise

correlations, as emphasized by previous models, make no significant contributions to improved

population representations by visual training. Instead, we identified neural manifold shrinkage due

to reduced trial-by-trial neural response variability, a previously unexplored factor, as the primary

mechanism underlying improved population representations. Furthermore, we showed that

manifold shrinkage successfully accounts for learning effects across various domains, including

artificial neural responses in deep neural networks trained on typical VPL tasks, multivariate

BOLD signals in humans, and multi-unit activities in monkeys. These converging results suggest

that our neural geometry theory offers a quantitative and comprehensive approach to explain a

wide range of empirical results and to reconcile previously conflicting models of VPL.

Keywords: visual perceptual learning, population representations, deep convolutional neural

networks, functional magnetic resonance imaging, multi-unit neural recordings

INTRODUCTION

Adapting to new visual environments is crucial for an organism's survival in its environment. This ability is well exemplified by visual perceptual learning (VPL), which is defined as long-term performance enhancements resulting from visual experience [1,2]. Therefore, elucidating the mechanisms underlying VPL may contribute to an understanding of how organisms adapt to new environments. However, despite years of research in systems neuroscience, psychophysics and machine learning, the mechanisms behind VPL remain mysterious.

It is widely acknowledged that visual training enhances behavioral performance and refines neural representations in neural populations. Previous studies using human neuroimaging and monkey neurophysiology have demonstrated a significant improvement in the fidelity of stimulus encoding within population responses [3-5]. These findings strongly support the theory that enhanced signal-to-noise ratios serve as a potent computational mechanism for refined neural representations associated with VPL (Fig. 1E) [6-8]. However, the precise mechanisms underlying how visual training refines neural representations remain elusive. Several conflicting models have been proposed based on neural changes associated with VPL. One model suggests that VPL is associated with changes in population representations resulting from changes in neuronal tuning curves, as evidenced by sharpened orientation tuning curves in monkey visual cortex [9,10]. Another model assumes that changes in population representations result from a reduction in trial-by-trial co-variation of neuronal firing rate, known as noise correlations, which have been observed in association with VPL in both monkeys and songbirds [11-14].

The primary conceptual gap in reconciling the conflicting models of VPL lies in their focus on mechanisms proposed at the single-unit level (e.g., changes in tuning curves or noise correlations), whereas the effects of these mechanisms in VPL should be evaluated at the population level (i.e., improved population representations). Although VPL is indeed associated with changes in both single-unit responses and improved population representations, it remains controversial whether changes in single-unit responses are the actual *cause* of improved population representations. It is possible that changes in single-unit responses associated with VPL are merely by-products of underlying processes. While these conflicting models of VPL capture certain aspects of the empirical findings, they fail to generate falsifiable predictions about how changes in single-unit responses contribute to improved population representations.

A significant obstacle to comparing the conflicting models of VPL is the complex interactions between different aspects of single-unit responses (e.g., tuning curves and noise correlations) on population representations. Computational neuroscience research has elucidated that the impact of noise correlations on population representations heavily depends on its interaction with tuning curves [15]. It is important to note that reduced noise correlations do not inherently enhance information in a neural population [16-18]. Moreover, the challenge is exacerbated by the fact that their interaction effects are even changing rather than remaining stable throughout a training process, because both noise correlations and tuning curves change. These dynamic changes further complicate the understanding of how training affects their interactions. To overcome this, a comprehensive computational theory is imperative to quantify and disentangle the effects of different changes in single-unit responses, such as sharpened tuning curves and reduced noise correlations, on neural representations at the population level.

To comprehensively explain these conflicting models, we developed a neural geometry theory of VPL. In this theory, trial-by-trial population responses elicited by two stimuli for discrimination form two differentiable manifolds in a high-dimensional neural space. In this space, changes in single-unit responses (e.g., tuning curves, Fano factor, and noise correlations) can be interpreted as changes in several fundamental and measurable geometric properties (e.g., centroids, size, orientations) of neural manifolds. This approach allows quantitative comparisons of conflicting models of VPL and assessments of their contributions to population representations within the same computational framework. Thus, this theory directly bridges single-unit responses and population representations, and offers a normative account of the potential neural mechanisms underlying VPL. Specifically, this theory proposes four possible training-induced geometric changes (signal enhancement, manifold shrinkage, signal rotation, and manifold warping) that can summarize all previous models of VPL. Thus, improved population representations can be achieved by one or a combination of the four interpretable mechanisms.

To assess this neural geometry theory, we trained deep convolutional neural network (DCNN) on two typical VPL tasks—orientation discrimination and motion direction discrimination. First, the DCNNs successfully replicated a wide range of psychophysical and imaging findings in humans, as well as neurophysiological findings in monkeys. Second, analyses of the geometric mechanisms mentioned above suggest that changes in both tuning curves and noise correlations are indeed present in VPL. Third, and most importantly, our analysis further

revealed that neither changes in tuning curves nor changes in noise correlations at the single-unit
level contributed significantly to improved population representations, despite their strong
association with VPL. Surprisingly, we found that neural manifold shrinkage induced by reduced
response variability emerged as the primary mechanism driving VPL. We conducted additional
tests to examine the predictions derived from the geometry theory across different levels of
measurement in different spaces. Remarkably, we found that the geometry theory incorporating
manifold shrinkage aligned closely with BOLD response changes associated with VPL of motion
direction discrimination in humans, as well as the electrophysiological population response
changes associated with VPL of contrast discrimination in monkey V4.

RESULTS

Orientation discrimination learning improves behavioral performance of DCNN

To elucidate the neurocomputational mechanisms of VPL, we trained a DCNN (Fig. 1A) to
perform a classical orientation discrimination task [7]. Similar to the neural network in ref. [19], this
neural network inherits the first five convolutional layers of AlexNet, which was pre-trained on
ImageNet [20]. To emulate the decision stage of orientation discrimination, we added a linear
decoding layer and used the logistic function to classify the activity of the decision unit into a
binary perceptual choice (i.e., clockwise or counterclockwise rotation of the target stimulus
relative to the reference stimulus). Importantly, similar to previous psychophysical studies [7,21],
we systematically manipulated the level of input image noise (Fig. 1B). The network was trained
on stimuli with multiple noise and contrast levels (see Methods and Materials for training details).

To evaluate the performance of the neural network and compare it directly with human
psychophysical results, we assessed orientation discrimination accuracy as a function of stimulus
contrast and noise (i.e., behavioral psychometric functions in Fig. 1C&D), and further derived
contrast thresholds as a function of image noise level (Fig. 1F, Threshold vs. Noise, TvN function).
We found that training significantly improved the network performance in this task in almost all
stimulus contrast and noise conditions. Importantly, training uniformly reduced contrast thresholds
at all noise levels (Fig. 1F). This downshift of TvN functions is consistent with the well-established
psychophysical result (replotted in Fig. 1E) as previously observed in humans [7,8].

Figure 1. A DCNN (**A**) is trained on an orientation discrimination task with Gabor stimuli embedded in different levels of image noise (**B**). Orientation discrimination accuracy is significantly improved from pre- (**C**) to post-test (**D**). Training induces a downshift of the Threshold vs. Noise function (**F**), an effect that is qualitatively similar to existing human psychophysical results (**E**, corresponds to the 70.7% accuracy condition in Fig. 1 of ref. 7). The absolute quantitative differences between **E** and **F** may be due to differences in the overall signal-to-noise ratio or the number of layers and units between the human visual system and the DCNN.

Orientation discrimination learning refines neural population representations

Having established the behavioral signature of VPL in our DCNN, we next sought to understand the effects of visual training on population representations in the network. We simulated many trials elicited by the two orientation stimuli for fine discrimination and performed multivariate decoding analyses in each layer. Training improved decoding accuracy in later layers (layers 3-5, Fig. 2F). The learning effect is more pronounced in later layers possibly because the benefits of visual training accumulate from early to later layers. More formally, we calculated linear Fisher information, a classical metric in computational neuroscience, to quantify stimulus information (i.e., how well the two stimuli can be discriminated based on population responses, see Methods and Materials) in each layer before and after training. The amount of sensory information

176 represented in later layers was indeed enhanced by training (Fig. 2G). Such refined neural
177 representation at the population level is consistent with the decoding results based on both
178 macroscopic cortical activity in humans [3,4,22] and multi-unit spiking activity in monkeys [13,14].

Figure 2. Neural correlates of VPL in humans (***A***), monkeys (***B-E***), and in our DCNN
(***F-J***). Visual training improves stimulus decoding accuracy in related regions in the
human brain (***A***) and decoding scores of Fisher's linear discriminant (FLD) in monkey
V1 (***B***). Visual training sharpens orientation tuning curves of neurons in monkey V4
(***C***), and also reduces Fano factors and interneuron noise correlations (***D&E***). Similar
results are observed in the DCNN: network training also improves decoding accuracy
in layers 3-5 (***F***), and averaged linear Fisher information (aLFI, total information in
each layer divided by the number of units in that layer) (***G***) in layers 3-5. Training
sharpens orientation tuning curves of units in layers 1-4 in the DCNN (results of layer
4 only are shown in ***H***). Similar reduction of Fano factors and noise correlations are
observed in the DCNN (results of layer 1 only are shown in ***I&J***). The data shown in
H-J are the median value across units in a layer. The results of all five layers are shown
in Supplementary Fig. S1. Panels ***A-E*** are reproduced by the data points shown in the
original papers. Error bars and error shadings in panels ***F-J*** represent the S.E.M. across
four reference orientations (error shadings in ***H*** are small and barely visible).

Orientation discrimination learning changes response properties of individual units

In addition to characterizing VPL on the population level, we sought to investigate the neural correlates at the single-unit level.

Importantly, we found that three key neural signatures of VPL as documented in the neurophysiological literature emerge naturally from the neural network training. First, training modestly sharpened the tuning curves of artificial neurons in layers 1 to 4 (Fig. 2H& and Supplementary Fig. S1). The finding of changed tuning curves has been reported previously in refs. 9,10,23 (Fig. 2C, but see also null results in ref. 24). Second, we observed a significant decrease in Fano factor of individual units in all five layers (Fig. 2I and Supplementary Fig. S1), a phenomenon indicating an increased signal-to-noise ratio of individual neuronal responses. This has been reported in both humans 25 and monkeys 11,23 (Fig. 2D). The sharpened tuning curve and reduced Fano factor are also consistent with theoretical modeling 19. Third, training reduced trial-by-trial noise correlations between units in all five layers (Fig. 2J and Supplementary Fig. S1), a finding also consistent with several empirical results in monkeys 11-14. Critically, we also found that the reduction in noise correlation depended on tuning similarity. Learning reduced the noise correlations between units with similar tunings (i.e., positive signal correlations) and increased the noise correlations between units with opposite tunings (i.e., negative signal correlations) (Supplementary Fig. S2). Previous theoretical work has suggested that the former type of noise correlations is detrimental for information coding and the latter type is beneficial 15,16. The pattern of reduced detrimental and increased beneficial noise correlations has been discovered with learning tasks in songbirds 26 and with attention tasks in monkeys 27.

In addition to these classical neurophysiological findings in VPL, our network also captures some important response properties of sensory neurons in the primate early visual system. First, the relationship between the Fano factor and orientation tuning of the artificial neurons bears strong resemblances to the empirical measures of V1 neurons in monkeys 28 (Supplementary Fig. S2). Second, we found a positive relationship between signal correlation and noise correlation among artificial neurons in all layers (Supplementary Fig. S2). This relationship has also recently been documented as a ubiquitous phenomenon in both electrophysiological 29-31 and human imaging 17,18,32 studies.

Taken together, our neural network reproduces a wide range of psychophysical, human imaging, and monkey neurophysiological findings in VPL. Although these similarities are

227 qualitative and not quantitative, these results suggest that our DCNN allows us to explore
228 neurocomputational mechanisms that may be difficult to elucidate in empirical experiments.

229
Figure 3. Four possible mechanisms of VPL in neural populations. *A.* According to classical
signal detection theory, better sensory discriminability can be represented as a larger
separation between two response distributions. Signal enhancement predicts enlarged
distances between the mean of the two distributions while noise reduction predicts reduced
variance between the two response distributions. *B.* Stimulus orientation as a continuous
stimulus variable can evoke population responses with the same shape as individual tuning
curves. *C.* If we continuously sweep the orientation value, the mean of the population

responses forms a manifold in a high-dimensional neural space with the number of
dimensions being equal to the number of units. The mean population responses to the two
stimuli in a discrimination task are two points on the manifold. ***D.*** In realistic population
responses, the trial-by-trial population responses to the two stimuli form two high-
dimensional response distributions (i.e., neural manifolds). The manifolds look elliptical
rather than spherical due to pairwise noise correlations between units. In this high-
dimensional neural space, the signal enhancement mechanism predicts an increased
Euclidean distance (i.e., signal separation, ***E***) between two high-dimensional response
distributions. However, no significant increase in signal separation is observed in any of the
five layers (signal separation decreases in the first two layers, ***F***). The manifold shrinkage
mechanism predicts reduced variance (***G***) of the two neural manifolds. This is observed in
all five layers (***H***). ***I***. The signal rotation mechanism predicts that the positions of the centroid
(i.e., mean) of the two manifolds are changed by training. ***J***. The rotation angle ranges from
$50º$~$70º$ in all five layers, supporting the existence of signal rotation. ***K.*** The manifold
warping mechanism predicts that training changes the shape of the population response
covariance. ***L.*** Indeed, training mostly reduces the variance of the high-variance principal
components of the population responses. The principal components (PC, only show PCs that
account for >99% of the total variance) are ranked from high to low variance. ***M.*** More
importantly, the directions of the principal components rotate significantly from pre- to post-
test.

Neural correlates of four computational mechanisms under the neural geometry theory of VPL

How would improved sensory discrimination manifest in high-dimensional population responses?

In the simplified one-dimensional scenario (Fig. 3A), the classical signal detection theory posits

that better sensory discrimination can be achieved by either increasing the distance between the

means (i.e., signal enhancement) and/or decreasing the variance (i.e., noise reduction) of the two

response distributions, because the former only changes the tuning (i.e., mean) of a response unit

and the latter only reduces the variance around a fixed mean. In multivariate population responses,

the two stimuli to be discriminated instead generate two multivariate response distributions (i.e.,

neural manifold) in a high-dimensional neural space whose dimension corresponds to the

dimension of a population (Fig. 3B&C). In a simplified visualization in 2D space (i.e., assuming

a population of only two units, Fig. 3D), the two distributions are elliptical due to noise correlations

between units. We refer to the vector connecting the mean of the two distributions as the *signal*

vector and its modulus length (i.e., the Euclidean distance between the two manifold centroids) as

the *signal separation*.

In the high-dimensional neural space, our neural geometry theory of VPL proposes that

visual training improves sensory discrimination by shaping some fundamental geometric

properties of the neural manifolds. Here, under this theory, there exist only four possible

mechanisms to separate two neural manifolds (Eq. 4 in Methods and Materials). First, according

to the classical signal detection theory, the *signal enhancement* mechanism predicts an increased

Euclidean distance between the centroids of the two neural manifolds (Fig. 3E). However, we

found that the signal separation between the two manifolds did not significantly increase with

learning in all five layers, and even slightly decreased in the first two layers (Fig. 3F). Second, the

manifold shrinkage mechanism predicts that visual training reduces the trial-by-trial response

variance of units, thereby reducing the size of the manifolds (Fig. 3G). This is what we found in

all five layers (Fig. 3H). These results argue that manifold shrinkage plays a more important role

than signal enhancement in VPL. We further included two novel mechanisms that can only occur

in high-dimensional neural space and lead to increased discriminability of two response

distributions. In the third mechanism, although visual training did not increase signal separation,

it may change the positions of the centroids of the two manifolds and consequently increase

discriminability due to the elliptical shape of the manifolds (Fig. 3I). Interestingly, we found that

the signal vectors in each layer were rotated by ~50°-70° after training (Fig. 3J). We call this

mechanism *signal rotation*. Fourth, visual training can warp the shapes of the high-dimensional

neural manifolds while keeping the size of the manifolds unchanged. As evidenced by the change

of covariance structure, we found that visual training systematically warped the shape (i.e.,

covariance structures) of the high-dimensional neural manifolds (Fig. 3K-M). We refer to this

mechanism as *manifold warping*. Note that manifold warping includes both the changes in

correlation structures and the redistribution of variances across individual units, while holding the

total variance constant. It is manifold shrinkage that attenuates the total variance.

In summary, our neural geometry theory of VPL summarizes possible mechanisms of

improved sensory discrimination into explicit and interpretable changes of geometric properties in

high-dimensional neural space. We present four possible computational mechanisms predicted by

the neural geometry theory of VPL and identify neural evidence for three of them (i.e., except

signal enhancement) in our DCNN training. Note that these four mechanisms (or a combination

thereof) encompass all possible scenarios by which learning can potentially improve sensory

discrimination.

Information-theoretic analyses reveal the key role of manifold shrinkage in orientation

discrimination learning

Given the four possible mechanisms (i.e., signal enhancement, manifold shrinkage, signal rotation, and manifold warping) and their complex interaction effects, how can we delineate their respective contributions to improved population representations? Here, we use linear Fisher information to quantify manifold separability. Besides, we introduce a stepwise approach to further disentangle the respective contributions of the four possible mechanisms. Specially, their respective contributions are assessed by sequentially allowing only one mechanism to occur and quantifying its endowed changes in the linear Fisher information of whole populations (Fig. 4A). For example, as shown in Fig. 4, we first calculate how much information is enhanced by considering only the signal enhancement scenario, then by considering both signal enhancement and manifold shrinkage, and so on until all four mechanisms are included. Here sensory information refers to manifold separability. We emphasize that this stepwise approach is intended only to facilitate the understanding of the computational underpinning of each mechanism. It does not imply that these mechanisms must occur sequentially over the course of visual training.

Interestingly, not all mechanisms have a positive effect on population representations. We found that the effect of signal enhancement is minimal in all five layers. This mechanism even reduces stimulus information in layers 1&2. This is consistent with the reduced Euclidean distance in the first two layers (Fig. 3F). Manifold shrinkage significantly enhances stimulus information in almost all layers. This makes sense because the decreased variance reduces the overlap between the two manifolds. Interestingly, we found that signal rotation appears to enhance stimulus information (Fig. 4B, green bars). This is because rotation of the signal vectors disrupts their relative parallelism to the covariance direction at pre-test, making them more orthogonal. Such changes increase the apparent information. However, the effect becomes minimal when manifold warping is further considered (Fig. 4B, magenta bars) because visual training also warps the covariance direction to realign it with the post-test signal vector, thereby reducing stimulus information.

We want to highlight two important aspects of such stepwise analyses. First, we use signal rotation and manifold warping to reflect changes in the direction of the signal vector and the shape of the covariance, which are theoretically independent (Fig. 3I&K). But what matters for the fidelity of population representations is the relative geometric relationship between covariance and

signal vector rather than their absolute magnitude [15,16] (Fig. 4C-F). Given that learning changes
both signal vector and covariance, we argue that their contributions to improved population
representations should be evaluated together. In Fig. 4B, signal rotation and manifold warping
appear to have strong positive and negative effects on information, respectively. However, their
joint effect is modest (i.e., the magenta bars compared to the blue bars in Fig. 4B). Thus, it can be
argued that manifold shrinkage is the primary contributor to improved stimulus information (also
see Discussion).

Figure 4. **A.** The effects of four mechanisms on population representations are decomposed
into four distinct steps (**A**). **B** shows the effects on information gain by sequentially adding
each of the four mechanisms in each layer. For example, the increase in height from the
brown to the blue bars indicates the positive contribution of manifold shrinkage to encoded
stimulus information. **C-F** illustrate the strong interaction effects between covariance and
signal vector. For distributions with identical covariance (**C** and **E**; **D** and **F**), detrimental (**C**
or **F**) or beneficial (**D** or **E**) effects on discriminability are possible, depending the signal
vector. Similarly, the effects of the signal vector also depend on its relative geometry to the
axis of covariance.

Second, the results are not affected by the order in which the mechanisms are applied changes. As mentioned above, the effects of signal rotation and manifold warping should be evaluated together, but the effects of signal enhancement and manifold shrinkage are independent. Hence, changing the order does not affect our results as long as signal rotation and manifold warping are carried out consecutively. In fact, the apparent positive and negative effects of signal rotation and manifold warping in the stepwise analyses can be reversed when their order is switched, but both orders actually reflect identical underlying changes (see detailed analyses in Supplementary Fig. S6).

Taken together, we propose an interpretable and quantitative neural geometry theory of VPL where visual training refines the geometry of representations in a high-dimensional neural space. Specially, our neural geometry theory predicts four different computational mechanisms of VPL. Our neural network allows us to readily measure the activity of all our simulated units across the visual hierarchy. Using the neural network, we first verified the existence of three of four possible mechanisms, and we performed systematic information-theoretic analyses to disentangle the effects of them. Most importantly, we found that manifold shrinkage in population responses was the key mechanism underlying the improved population representations induced by visual training in the DCNN.

Neural geometric changes of motion direction discrimination learning in DCNN

The above analyses focus only on a classical VPL task—orientation discrimination and a specific neural network structure—a six-layer convolutional neural network. In this section, we switch to motion VPL—another sensory domain that is also widely used in psychophysical [33,34], human imaging [3,4], and neurophysiological studies [35]. Importantly, motion VPL involves the processing of both spatial and temporal signals rather than merely static spatial information in orientation learning. Similarly, we inherited the first six layers of the pre-trained C3D network [36] and trained the neural network to perform a motion direction discrimination task commonly used in psychophysics (see Methods and Materials for stimulus and training details).

In the motion DCNN, we found similar mechanisms as in the orientation discrimination learning task. First, motion direction discrimination training significantly improved the behavioral performance of the network (Fig. 5B). Second, training also enhanced decoding accuracy and

averaged linear Fisher information in later layers (Fig. 5C&D), suggesting that such training refines stimulus representation at the population level. Third, the effects of motion direction discrimination training on individual units in layer 6 are also pronounced (see results for all six layers in Supplementary Fig. S3). We found that training reduced Fano factor (Fig. 5E) and noise correlations (Fig. 5F). Fourth, training did not significantly improve signal separation (Fig. 5G) but markedly reduced response variance (Fig. 5H). In addition, motion direction discrimination training also induced two novel mechanisms—signal rotation (Fig. 5I) and manifold warping (Fig. 5J&K). Most importantly, the four mechanisms induced by the training had similar respective contributions to population representations (Fig. 5L).

Figure 5. DCNN modeling of motion VPL. The DCNN (**A**) of motion VPL uses 3D convolutions to process video stimuli. Here, we simplify the 4D feature maps in each convolutional layer and show them as 3D maps only for illustration purposes. Training significantly improves DCNN direction discrimination performance (**B**), decoding accuracy (**C**), and averaged linear Fisher information (**D**). For single-unit analyses, motion direction discrimination training also reduces the Fano factor (**E**) and noise correlation (**F**) in layer 6. Similar to orientation discrimination training, motion direction discrimination training does not significantly enhance signal separation (**G**), but significantly rotates the position of the two distributions in layer 6 (**I**). Importantly, training clearly reduces the response variance in layer 6 (**H**). Specifically, training reduces the variance of the high variance PCs and rotates the directions of all PCs, indicating a significant effect of manifold warping in layer 6 (**J&K**). The pattern of information gain associated with the four possible mechanisms (**L**) is

406   consistent with that of orientation discrimination training. See results for all six layers in
407   Supplementary Fig. S3. Note that some error bars are very small and barely visible.
408

Neural geometric changes of motion direction discrimination learning in the human brain

Above, our DCNNs help verify, quantify and disentangle the four possible mechanisms proposed

by our neural geometry theory. The converging results in the DCNNs of orientation and motion

direction discrimination, and the remarkable agreement between our DCNNs and existing

empirical neuroscientific findings support the biological plausibility of our DCNNs. However, it

remains unknown whether these predictions are only present in the DCNNs and have no biological

basis in the brain.

To address this question, we analyzed blood oxygenation level dependent (BOLD)

responses in the cortex of human subjects before and after they were trained on a motion direction

discrimination task (Fig. 6A, ref. 37). Twenty-two human subjects participated in the motion VPL

study. Subjects were trained for 10 days on a fine direction discrimination task, and psychophysical

and fMRI tests were performed before and after training.

We identified the early visual areas (V1-V3), the motion-selective regions (V3A and

hMT+), and the decision region (intraparietal sulcus, IPS) using independent functional localizer

experiments (Fig. 6B). We estimated single-trial responses of voxels in these regions and then

performed decoding analyses in these predefined regions, and found that motion training

significantly enhanced decoding accuracy (Fig. 6C) and averaged linear Fisher information (Fig.

6D) in areas V3A and hMT+, a result consistent with several human fMRI studies on motion VPL

[3,4,22].

Figure 6. Trial diagram and training paradigm (*A*), and regions-of-interest in a typical subject (*B*). Motion direction discrimination training in humans significantly improves decoding accuracy (*C*) and averaged linear Fisher information (*D*) in areas V3A and hMT+, a finding consistent with several existing fMRI studies of motion VPL. Note that the four data points in V3A appear as outliers in *C&D*, but the results still hold if these data points are removed. Motion direction discrimination training does not significantly change signal separation in V3A and hMT+ (*E*), but reduces voxel response variance in V3A and hMT+ (*F*) and intervoxel noise correlations in hMT+ (*G*). Similar to the motion DCNNs, Motion direction discrimination training in humans also rotates stimulus distributions (*H*), reduces the variance of high variance PCs (*I*), and warps the covariance directions (*J*). The patterns of information gain associated

with the four mechanisms (K) are consistent with those in the DCNNs of both orientation and motion VPL. The unit "PSC" represents percent signal change of blood oxygenation level dependent signals. Individual data points represent the human subjects. Error bars in all panels represent the S.E.M. across subjects. Significance conventions are: *, $p < 0.05$; **, $p < 0.01$.

We further investigated the coding principles in areas V3A and hMT+, and repeated the above analyses of DCNNs on fMRI data. Note that here we performed the same analyses on voxels instead of artificial neurons in DCNNs. Consistent with the predictions of the DCNNs, motion direction discrimination training in humans did not increase signal separation (Fig. 6E) but markedly reduced voxel response variance (Fig. 6F) in both areas. Motion direction discrimination training also significantly reduced intervoxel correlations in hMT+ (Fig. 6G). The mechanism of signal rotation was also evident, as indicated by the average ~55º rotation of the signal vectors in both areas (Fig. 6H). In addition, training warped the magnitude and direction of the covariance (Fig. 6I&J). Most importantly, the respective contributions of these four mechanisms in both brain regions were similar to the pattern in the DCNNs (Fig. 6K).

Neural geometric changes of contrast discrimination learning in monkey V4

So far, we have verified our neural geometry theory of in DCNNs and in the human brain. However, voxel responses in fMRI studies reflect macroscopic brain activity that aggregates the responses of ~300, 000 – 50, 000 neurons [38]. It remains unclear whether the mechanisms we have discovered so far also exist at the local circuit level of single cells or small clusters of cells. Although in the DCNNs our neural geometry theory generates two novel predictions that (1) visual training induces novel mechanisms (e.g., signal rotation and manifold warping) that can only be revealed by analyzing multivariate responses, and that (2) manifold shrinkage is the primary mechanism of VPL. To our knowledge, these predictions have not been systematically tested using intracranial recording.

Figure 7. We analyzed population responses in area V4 (***A***) of two monkeys while they were trained on a fine contrast discrimination task (***B***). The first four and last four training sessions were grouped as pre- and the post-test conditions, respectively. Contrast discrimination training significantly improved behavioral performance from the early to late stage of training (***C***). All individual data points in ***C*** represent the six target contrast conditions (27, 28, 29, 31, 32, and 33%; the reference contrast is 30%). Each point is averaged over the two monkeys. See plots for individual monkeys in Supplementary Fig. S5. Error bars indicate the S.E.M. across the six conditions. ***D-G*** illustrates the full width at half maximum of the response distributions of four pairs of channels at pre- and post-test (***D&E*** for monkey 1 and ***F&G*** for monkey 2). Solid lines represent 30% reference contrast, and dashed lines represent 29% and 31% target contrast in monkey 1 and monkey 2, respectively. These results show that learning systematically changes the geometries of the multivariate responses.

To further test our hypotheses on neuronal spiking activity, we analyzed the population responses of V4 neurons in two monkeys (Fig. 7A) at the early and the stage of training in which they learned to perform on a fine contrast discrimination task (Fig. 7B, ref. 14). In this task, each monkey was presented sequentially with two identical Gabor patches with different contrast levels. The contrast of the reference (i.e., the first) stimulus was always fixed at 30%, and the contrast of

the target (i.e., the second) stimuli varied systematically near the reference contrast (i.e.,
27/28/29/31/32/33%). This contrast discrimination training significantly improved behavioral
performance (Fig. 7C). Most importantly, responses of multiple channels were continuously
recorded via chronically implanted electrodes in area V4 (21 and 29 channels for monkey 1 and 2,
respectively) throughout training (21 and 23 training sessions for the two monkeys, respectively).
This continuous multi-unit recording is the key to disentangling population-level changes
associated with VPL.

We used the above analyses (previously applied to DCNNs and human fMRI data) and
applied them to the monkey V4 responses, and again found highly consistent results. First, contrast
discrimination training improved stimulus information at the population level (Fig. 8A&B).
Second, at the individual level, contrast discrimination training also significantly reduced Fano
factors (Fig. 8C) and noise correlations (Fig. 8D), consistent with several existing findings.
Interestingly, while the trial-by-trial variance was significantly reduced after training (Fig. 8F), no
apparent change in signal separation was observed (Fig. 8E), suggesting the predominant role of
manifold shrinkage. Importantly, we again observed evidence for signal rotation (Fig. 8G) and
manifold warping (Fig. 8H&I). The stepwise information analyses also qualitatively replicated the
relative contributions of the four mechanisms to the total stimulus information encoded in the
population (Fig. 8J).

Taken together, we proposed a neural geometry theory of VPL that links improved
population representations to four possible neurocomputational mechanisms. We quantitively
verified our theory across species, behavioral tasks, experimental conditions, measurement
modalities, and brain regions. These converging results in humans, monkeys, and DCNNs suggest
that our neural geometry theory can account for VPL in both biological and artificial systems.

Figure 8. Population activity analyses of contrast discrimination learning in monkey
V4. Contrast discrimination training significantly enhanced stimulus information at
the population level (*A&B*). Consistent with VPL in the DCNNs and the human brain,
training monkeys on a contrast discrimination task reduced Fano factors (*C*), noise
correlations (*D*), and response variance (*F*), but had no significant effect on signal
separation (*E*). We also found evidence for signal rotation (*G*) and manifold warping
(*H&I*). The stepwise information analyses also show the similar pattern of the four
mechanisms (*J*). The unit "spk/s" indicates the number of spikes per second (i.e., firing
rate). We calculate aLFI and information gain using stimulus contrast as decimal
values (i.e., 0.29), so they have arbitrary units. The definitions of the individual data
points and error bars are the same as in Fig. 7.

523
524
525

DISCUSSION

Our neural geometry theory bridges response properties at the single-unit level and those at the population level through geometric interpretations in a high-dimensional neural space. This theory allowed us to quantify possible contributions of single-unit responses to population representations. It has been controversial whether single-unit properties such as sharpened tuning curves [9,10] or reduction of noise correlations [11,12] contribute to VPL. Our information-theoretic analysis on neural geometry suggested that, although these properties were indeed observed, they did not contribute significantly to the improved population representations associated with VPL. Rather, we found that the totally overlooked mechanism—the response variance of individual units (i.e., manifold shrinkage)—is the primary contributor to the improved population representations associated with VPL. These results were further tested on deep convolutional neural networks, human fMRI data, and monkey neurophysiological data associated with different VPL tasks and brain regions.

Given the pronounced changes in tuning curves and changes in noise correlations observed after training, why do they not contribute to VPL? Conventional studies treat changes in tuning curves and in noise correlations as two independent factors mediating VPL. However, in our theory, the effects of tuning curve changes partially overlap with those of noise correlations (Eq. 4 in Methods and Materials). According to the neural geometry theory, the effects of tuning curve changes can be decomposed into two parts - signal enhancement independent of noise correlations and signal rotation interacting with noise correlations. We observed minimal evidence for signal enhancement (i.e., comparable magnitude of signal separation before and after training), and it certainly makes minimal contributions to population representations. Although we observed clear evidence for signal rotation and manifold warping, which correspond to the interactions between tuning curve and noise correlation changes, what matters here is the relative geometric relationship between the direction of noise correlations and signal vectors, rather than their absolute changes. Therefore, their respective contributions appeared significant but their overall joint effects were minimal because their effects canceled each other out (Supplementary Figure S6). Our findings challenge previous studies that have only assessed the effects of individual factors without considering the joint effects of multiple factors [11,13]. The joint effects of multiple factors may be quite different from simply summing the effects of individual factors.

The novel finding that manifold shrinkage is the primary contributor to improved

population representations is of unique significance in constraining the model of VPL. In manifold

shrinkage, the total variance of the high-dimensional distributions is scaled down (i.e., λ in Eq. 4

is reduced), but the overall shape of the distributions remains unchanged. Thus, the two stimulus

distributions simply shrink to a "smaller size" (Fig. 4). Note that manifold shrinkage is independent

of any tuning changes and noise correlation changes. We also emphasize that manifold shrinkage

and manifold warping are two different mechanisms. In our theory, manifold warping redistributes

the variance of the high-dimensional distributions in different directions (i.e., $\hat{\lambda}_i$ and ξ_i in Eq. 4

are changed) but, unlike manifold shrinkage, the total amount of variance remains unchanged.

Thus, the shape of the two stimulus distributions is significantly warped.

Our neural geometry theory bridges existing human imaging and monkey

neurophysiological studies by linking changes at both the individual and population levels. Most

existing human imaging studies on VPL have focused only on improved representations at the

population level [3,4,22], whereas most single-unit studies in monkeys can only measure changes in

individual neurons [9,10]. Only two empirical studies have attempted to address the relationship

between the two levels using human fMRI [22] and monkey population recording [13]. In these studies,

researchers analyzed population responses by projecting high-dimensional neural manifolds onto

a one-dimensional optimal decision plane. However, we argue that high-dimensional population

responses, once projected onto a one-dimensional decision plane, no longer contain useful

information about changes in individual units. In their approach, all changes in individual units are

conflated (see our analytical derivations in Supplementary Note 1) and no longer distinguishable.

In contrast, we evaluated all mechanisms within the high-dimensional neural manifolds *per se*

without any dimension reduction operation. Our approach can explicitly disentangle and quantify

the effects of individual factors (see similar analysis of our data in Supplementary Fig. S4). Our

highly consistent findings in both artificial and biological visual systems demonstrate the effective

utility of neural geometry theory in various domains of cognitive neuroscience. Although we only

examined VPL of orientation, motion direction, and contrast discrimination, we argue that our

neural geometry theory *per se* serves as a general framework to reveal whether the same as or

different mechanisms than VPL in this study play a role in VPL for other tasks, possibly involving

distinct brain regions. Neural geometry theory is also highly relevant for understanding the

mechanisms of many other cognitive processes. For example, attention [39] and arousal [40] have

also been shown to be associated with several neural changes (e.g., noise correlations). Similarly,

it remains unclear how these effects contribute to improved population representations.

Elucidating various cognitive processes from the perspective of neural geometry is a promising

direction for future studies.

In summary, the long-standing controversy over whether changed tuning curves or reduced

noise correlation, which are both property changes at the single-unit level, plays a significant role

in VPL has been addressed by our research within the framework of neural geometry theory. Our

converging results from monkey multi-unit recordings, human fMRI recordings, and neural

network applications associated with various VPL tasks indicate that neither changed tuning

curves nor reduced noise correlation affect the population response associated with VPL. Instead,

previously overlooked mechanisms such as manifold shrinkage were identified as significant

drivers of population response changes linked to VPL. This approach sheds light on how properties

at single-unit level are linked to those at the population level in various types of VPL, as well as

in other forms of learning and memory.

METHODS AND MATERIALS

DCNN modeling of orientation VPL

Stimuli. The network was trained to discriminate whether a target stimulus was tilted 1º clockwise or counterclockwise relative to a reference stimulus. All reference stimuli in the orientation discrimination task were Gabor patterns (227 × 227 pixels, spatial frequency=40 pixels/cycle, standard deviation of the Gaussian spatial envelope=50 pixels). The stimuli were varied in contrast (0.1 to 1.0 in 0.1 increments) and image noise level (8 levels: 0.005, 1, 5, 10, 15, 30, 50, 75). Similar to existing psychophysical studies [41], the image noise level is defined as the fraction of pixels randomly selected and replaced by Gaussian noise with a standard deviation of 15 gray level units. To mimic intrinsic sensory noise, we also added Gaussian white noise (standard deviation=10) to each stimulus [19]. To match the spatial frequency of noise and signal, the size of the replaced pixels was set to be 8 × 8. Four reference orientations (35, 55, 125, and 145º) were used and we trained four DCNNs for each of the four reference orientations.

Neural networks and training. A DCNN [20] was used to simulate the orientation VPL. We retained the first five convolutional layers of the pre-trained AlexNet and replaced its three fully connected layers with a single linear fully connected layer for perceptual choice. The network was configured in a Siamese fashion to perform the two-alternative forced-choice (2AFC) task: the same network was fed with both the target and the reference stimuli, producing two scalar outputs, h_t and h_r, respectively. The network then made the final decision with a probability p (classification confidence) calculated by the sigmoid function:

$$p = \frac{e^{h_t - h_r}}{1 + e^{h_t - h_r}} \tag{1}$$

The entire training procedure consisted of two distinct phases: the pre-training phase and the VPL phase. In the pre-training phase, the network was trained on full-contrast noiseless stimulus pairs to understand the task and to establish the pre-test baseline. In the VPL phase, the network was trained on stimulus pairs across all contrasts (10 levels) and noise levels (8 levels). The network was trained for 5000 epochs in the pre-training phase and 500 epochs in the VPL phase using the stochastic gradient descent learning algorithm. The learning rate and the momentum were set to 1e^{-5} and 0.9, respectively. The parameters were updated to minimize the cross-entropy loss between the network outputs and the true stimulus labels. The initial parameters in the fully connected layer were set to zero, as in ref. [19], while those in the convolutional layers

were taken directly from a pre-trained AlexNet available at

http://dl.caffe.berkeleyvision.org/bvlc_AlexNet.caffemodel. We trained one model for each of the

four reference orientations, and the entire procedure was repeated 10 times for each reference

orientation to control for randomness. All model and training procedures were implemented using

PyTorch 1.13.1.

Behavioral and neural changes. The behavioral performance of the network was evaluated

by measuring its classification confidence (Eq. 1) at all combinations of contrast and noise before

and after the training stage. Specifically, in each trial, the firing rate of each artificial neuron was

measured as the output of local response normalization or ReLU layers, averaged over all locations.

All measurements were obtained by simulating 1000 trials for better estimation. To ensure that

units were truly driven by the stimuli, only units with a mean firing rate greater than 0.001 before

and after training were included in the analyses [19].

The behavioral TvN curves of the model were derived for comparison with human

psychophysical results. Specifically, for each noise level, the contrast threshold was obtained by

interpolating the pre- and post-test accuracies of 55% and 70% respectively. To perform

population decoding analyses, we trained a linear classifier on the firing rates of the artificial

neurons to discriminate the target and the reference stimuli. The classifier was trained on half of

the 1000 simulated trials, while the other half served as the test dataset.

To characterize the response properties of individual units, we measured orientation-

selective tuning curves by sweeping the orientation of high-contrast stimuli from 0 to 180º. The

tuning curves were derived by averaging 100 simulated trials for each orientation. The resulting

tuning curves were then smoothed with a 10º Gaussian kernel. To control the heterogenous

response range across units, we then normalized the tuning curves of each unit by its maximum

response and averaged the tuning curves across units to obtain the group-level tuning curves. The

group-level tuning curves were then fitted with a Gaussian function and rescaled to 0~1 for better

comparison.

To calculate the Fano factor of each unit, we simulated 1000 trials for each reference

orientation. The Fano factor of each artificial neuron is defined as the ratio of the variance of the

firing rate to its mean. Similarly, noise correlations between artificial neurons were calculated as

the correlations between unit firing rates over the 1000 simulated trials for each reference

orientation. We took the median of the Fano factor across units in each layer to generate the data

plot (Fig. 2I). We took the median of the lower triangle of the noise correlation matrix in each

layer to generate the data plot (Fig. 2J). The error bars in Fig. 2I&J represent the standard errors

across four reference orientations.

Linear Fisher information analyses. To understand how neural activation contributes to

behavioral improvements, we applied linear Fisher information analysis to population responses.

We considered the firing rates of the same groups of units under the reference and the target

stimulus conditions as two distributions in a high-dimensional neural space. We refer to the *signal*

vector as the vector connecting the mean of the two distributions. A signal vector is calculated as

the difference between the mean firing rates of units to two stimuli. The signal separation is

referred to as the modulus length of the signal vector, and the angle of the signal vectors before

and after training is referred to as the signal rotation angle.

To measure how much information was contained in a layer per unit, we calculated the

average linear Fisher information (aLFI) as follows:

677
$$aLFI = \frac{1}{n} * \frac{df^T * \overline{\Sigma}^{-1} * df}{\Delta\theta^2} \tag{2}$$

678
$$\overline{\Sigma} = \frac{\Sigma_1 + \Sigma_2}{2}$$
$$= \frac{V_1^T * C_1 * V_1 + V_2^T * C_2 * V_2}{2} \tag{3}$$

where n is the number of units in a layer, $\Delta\theta$ is the separation between the target stimulus and the

reference stimulus (i.e., 1º). df is the signal vector. $\overline{\Sigma}$ is the mean of the covariance matrices (i.e.,

Σ_1 and Σ_2) of units responding to the two stimuli. V is a diagonal matrix with the variance of the

units as the diagonal terms. C is the correlation matrix of the population with all diagonal elements

equal to 1.

To further elaborate on the potential mechanisms of the improved LFI, we performed an

eigen decomposition on the covariance matrix $\overline{\Sigma}$, where we obtained λ_i, the eigenvalue of $\overline{\Sigma}$ and

ξ_i, its corresponding normalized eigenvector. The aLFI can be rewritten as follows:

687
$$aLFI = \frac{1}{n\Delta\theta^2} * \sum_{i=1}^{N} \frac{(df * \xi_i)^2}{\lambda_i}$$
$$= \frac{1}{n\Delta\theta^2} * \frac{|df|^2}{\overline{\lambda}} * \sum_{i=1}^{N} \frac{(df * \xi_i)^2}{\lambda_i} \tag{4}$$

where $\underline{\lambda}$ is the mean variance, and $\lambda_i = \underline{\lambda} * \hat{\lambda}_i$. $\frac{df}{|df|}$ is the unit vector with length of 1 and direction

as the same as the signal vector df. According to Eq. 4, we disentangled the potential mechanisms

of improved LFI into four subparts: signal enhancement, reflected by the modulus length $|df|$;

manifold shrinkage, reflected by the mean variance of $\bar{\lambda}$; signal rotation, reflected by the direction

of the signal vector df, and manifold warping, reflected by the relative angle of both ξ_i and $\hat{\lambda}_i$.

We applied a stepwise approach to assess their respective contributions by sequentially allowing

only one mechanism to occur and calculating the resulting changes in $aLFI$. Specifically, we first

calculated $aLFI$ at pre-test as:

$$aLFI_{pre} = \frac{1}{n\Delta\theta^2} * \frac{|df_{pre}|^2}{\underline{\lambda}_{pre}} * \sum_{i=1}^{N} \frac{\left(\frac{df_{pre}}{|df_{pre}|} * \xi_i^{pre}\right)^2}{\hat{\lambda}_i^{pre}} \tag{5}$$

698 Only considering the effect of signal enhancement, we can calculate its effect as $aLFI_{se}$:

$$aLFI_{se} = \frac{1}{n\Delta\theta^2} * \frac{|df_{post}|^2}{\underline{\lambda}_{pre}} * \sum_{i=1}^{N} \frac{\left(\frac{df_{pre}}{|df_{pre}|} * \xi_i^{pre}\right)^2}{\hat{\lambda}_i^{pre}} \tag{6}$$

700 Note that the only difference here is that the $|df_{pre}|^2$ in Eq. 5 is replaced by the $|df_{post}|^2$ in Eq. 6.

701 The difference between $aLFI_{se}$ and $aLFI_{pre}$ is considered as the information gain introduced by

702 the signal enhancement mechanism (i.e., the brown bars in in Fig. 4B). Following this idea, we

703 can calculate the stepwise $aLFI$ by one-by-one considering the effects of manifold shrinkage

704 $aLFI_{ms}$, signal rotation $aLFI_{sr}$, and manifold warping ($aLFI_{mw}$ or $aLFI_{post}$) as:

$$aLFI_{ms} = \frac{1}{n\Delta\theta^2} * \frac{|df_{post}|^2}{\underline{\lambda}_{post}} * \sum_{i=1}^{N} \frac{\left(\frac{df_{pre}}{|df_{pre}|} * \xi_i^{pre}\right)^2}{\hat{\lambda}_i^{pre}} \sum_i \frac{\left(\frac{df_{pre}}{|df_{pre}|} * \xi_i^{pre}\right)^2}{\hat{\lambda}_i^{pre}} \tag{7}$$

$$aLFI_{sr} = \frac{1}{n\Delta\theta^2} * \frac{|df_{post}|^2}{\underline{\lambda}_{post}} * \sum_{i=1}^{N} \frac{\left(\frac{df_{post}}{|df_{post}|} * \xi_i^{pre}\right)^2}{\hat{\lambda}_i^{pre}} \sum_i \frac{\left(\frac{df_{post}}{|df_{post}|} * \xi_i^{pre}\right)^2}{\hat{\lambda}_i^{pre}} \tag{8}$$

$$aLFI_{mw} = aLFI_{post} = \frac{1}{n\Delta\theta^2} * \frac{|df_{post}|^2}{\underline{\lambda}_{post}} * \sum_{i=1}^{N} \frac{\left(\frac{df_{post}}{|df_{post}|} * \xi_i^{post}\right)^2}{\hat{\lambda}_i^{post}} \tag{9}$$

708 The information gain in Fig. 4B indicates the difference between $aLFI_{se}$, $aLFI_{ms}$, $aLFI_{sr}$, and

709 $aLFI_{mw}$ (i.e., $aLFI_{post}$) as compared to the pre-test baseline $aLFI_{pre}$. They are shown as brown,

710 blue, green, and magenta bars in Fig. 4B, respectively.

DCNN modeling of motion VPL

Stimuli. The experiment used random dot motion (RDM) stimuli, which consist of a cloud of independent moving dots with some degree of coherence in a given moving direction [42]. The network was trained to discriminate whether the moving direction of a target RDM stimulus was 4º clockwise or counterclockwise relative to its corresponding reference RDM stimulus. To meet the network's specifications, the motion stimuli were 16-frame videos (112×112 pixels per frame). Within each frame, ~100 dots were displayed, with each dot represented by a cross of 3 pixels in both height and width. We set 8 coherence levels (8.84, 12.5, 17.7, 25, 35.3, 50, 70.7, and 100%) and four reference directions (45, 135, 225, and 315º). The motion speed was 7.5 pixels/frame. All non-coherently moving dots appeared randomly in the image. The display of each frame was limited to a centered circle with a diameter of 112 pixels, with the surrounding areas displayed in black.

Neural network architecture and training. Our DCNN is a 3D convolutional neural network inherited from the C3D network for action recognition [36]. The original C3D consists of 10 convolutional layers and 3 fully connected layers. The main difference between C3D and AlexNet is that C3D uses 3D convolutional kernels to process spatiotemporal information. We kept the first 6 convolutional layers from the pre-trained C3D and replaced the 3 fully connected layers with a fully connected layer that outputs a single scalar. The number of layers was chosen to (1) keep roughly similar number of parameters to the orientation DCNN, and (2) to roughly match the number of ROIs in the human neuroimaging experiment. Similar to the orientation DCNN, the motion DCNN was also configured in a Siamese fashion to perform the 2AFC task based on the sigmoid function.

Similar to the orientation DCNN, the entire training procedure consisted of two phases: the pre-training phase and the VPL phase. During the pre-training phase, the network was trained on full-coherence noiseless RDM pairs, while during the VPL phase, the network was trained on stimulus pairs across all coherence levels (8 levels). The network was trained for 1000 epochs in the pre-training phase and 2000 epochs in the training phase using stochastic gradient descent (SGD) with a learning rate of $1e^{-7}$, the momentum of 0.9, and a weight decay of 0.0005. The parameters were updated to minimize the cross-entropy loss between the network outputs and the true stimulus labels. The initial parameters in the fully connected layer were normally randomized while those in the convolutional layers were taken directly from a pre-trained C3D available at

. The entire procedure

was repeated 10 times for each reference direction to control for randomness.

Behavioral and neural analyses. The behavioral performance of the network was also

evaluated by its classification confidence (Eq. 1) at all coherence levels before and after the visual

training phase. In addition, the firing rates of artificial neurons were measured on each trial as the

output of the ReLU layers, averaged over all locations and time points. All measurements were

taken over 1000 simulated trials. To ensure that units were truly driven by the stimuli, only units

with a mean firing rate greater than 0.001 before and after training were included in subsequent

analyses.

To perform decoding analyses, we trained a linear classifier on the firing rates of the

artificial neurons to discriminate between the target and the reference stimuli. To assess the

performance of the classifier, we split all trials half-half as training and test datasets, and used the

average performance of the test-set. For comparison with the electrophysiological data, we

calculated the Fano factor of each unit as the ratio of the variance of the firing rate to its mean, and

the noise correlations as the correlation between the firing rates of units when viewing the same

RDM stimulus. In addition, to measure how much information was contained in a layer per unit,

we calculated the aLFI (see above).

We further validated the computational mechanisms in the motion direction discrimination

task. To this end, the firing rates of the same group of units under the reference and the target

stimuli were also considered as two distributions in a high-dimensional neural space. In the high-

dimensional neural space, we defined signal vector, signal separation, variance, correlation, signal

rotation angle, PC strength, and PC rotation angle as above.

Again, we computed linear Fisher information using a stepwise approach. For all models,

we sequentially added signal enhancement, manifold shrinkage, signal rotation, and manifold

warping to the calculation of linear Fisher information and see how the information within units

varies with all four mechanisms. Fig. 5L shows the results of the stepwise analysis in layer 6.

Supplementary Fig. S3 shows the results in all six layers of the motion DCNN.

Human fMRI experiment

The human fMRI experiment data have been published in ref. [37] for different research questions.

The core analyses in this study beyond preprocessing and ROI definitions are novel. We provide

relevant methods as follows and more detailed methods in Supplementary Note 2 to avoid cross-

referencing.

Subjects and Experimental procedures. A total of 22 human subjects (10 males and 12

females, ages 17-25) participated in the experiment. All participants had normal or correct-to-

normal vision. None of the participants were aware of the study's objectives. All participants

provided written informed consent, and the study obtained approval from the local ethics

committee at Peking University.

All subjects were trained on a direction discrimination task (Fig. 6A, see Supplementary

Note 2 for apparatus and stimulus details). The whole experiment consisted of three phases: pre-

test (2 days), training (10 days) and post-test (2 days). On day 1 at pre-test and day 2 at post-test,

subjects were tested on direction discrimination around 45º and 135º (angular difference 4º, 120

trials for each direction) to assess their behavioral performance before and after training. Subjects

were trained on the fine direction discrimination task for 10 days. Half of the subjects were trained

at 45º and the other half at 135º (see training details in Supplementary Note 2). Training-induced

behavioral improvements have been reported in our previous work [37].

To assess the neural changes induced by visual training, two identical fMRI sessions were

performed on day 1 at pre-test and day 2 at post-test, respectively. In each fMRI session, subjects

completed four runs of the motion direction discrimination task. Each run contained 30 trials for

45º and 135º (i.e., a total of 120 trials for each direction). Each run also contained 15 fixation trials

and the trial order was randomized.

MRI data acquisition. All MRI data were acquired using a 12-channel phase array coil on

a Siemens Trio 3T scanner at Peking University. The T1-weighted anatomical data with a

resolution of $1 \times 1 \times 1$ mm³ were collected for each subject. Echo-planar imaging (EPI) functional

data were collected for the motion direction discrimination task, retinotopic mapping, and motion

localizer experiments. EPI data were acquired using gradient echo-pulse sequences from 33 axial

slices, covering the whole brain. The standard EPI sequence used for data acquisition was as

follows: a repetition time of 2000 ms, an echo time of 30 ms, a flip angle of 90º, and a resolution

of $3 \times 3 \times 3$ mm³. The slice order was interleaved ascending.

In addition to the four runs of the motion direction discrimination task, we also collected
one or two retinotopic mapping runs [52,55] and a motion localizer run [43] to define regions-of-
interest.

MRI data analyses. In Brain Voyager QX, the anatomical data were transformed into the
Talairach coordinate space. For all functional data, the first four volumes of each functional run
were discarded to allow the longitudinal magnetization to reach a steady state. The functional data
underwent several standard preprocessing procedures, including slice timing correction, head
motion correction, spatial smoothing, temporal high-pass filtering (GLM with Fourier basis set at
2 cycles), and linear trend removal. Brain Voyager QX was also used to preprocess the data of the
retinotopic mapping experiment and the motion localizer experiment. We used the standard phase-
encoding method to define the retinotopic visual areas V1, V2, V3, and V3A [44,45]. A generalized
linear model (GLM) was then applied to the motion localizer data to define the motion-selective
voxels (hMT+ and motion-selective voxels in IPS).

The functional data of the motion direction discrimination task were preprocessed using
SPM12 (www.fil.ion.ucl.ac.uk/spm). The data were aligned to the first volume of the first run of
the first session, corrected for acquisition delay, and then normalized to the MNI coordinate space
using an EPI template. We used the GLMdenoise package developed in ref. [46] without evoking
multirun denoise procedures to estimate the single trial activity of voxels.

Voxel population response analyses. We adapted the analysis previously used for artificial
neurons in neural networks to the single-trial fMRI response estimates. To improve signal-to-noise
ratio, we selected the 60 most responsive voxels in each ROI at pre-test. We first investigated
which ROI was involved in motion VPL by measuring the discriminability between two different
motion conditions (trained direction, e.g., 45º vs. untrained direction, e.g., 135º) before and after
training. We trained a linear classifier on the fMRI data to discriminate between the two motion
conditions. To assess the performance of the classifier, we performed a leave-one-trial-out cross-
validation, and the average performance on the leave-out test trial was used as the discriminability
measure. We also computed the average linear Fisher information (see equations above) between
the 45º vs. 135º conditions to quantify stimulus discriminability. We found that motion direction
discrimination training significantly improved stimulus discriminability in V3A and hMT+.
Therefore, we only included V3A and hMT+ voxels in the subsequently analyses.

Similar to the analyses in the DCNNs, we defined the signal vector, the signal separation,

the variance, the intervoxel correlations, the signal rotation angle, the PC strength, and the PC

rotation angle in the multivoxel high-dimensional space using the same method defined above (Fig.

6). In addition, we applied the same stepwise analysis approach of calculating aLFI to the fMRI

data (Fig. 6K).

Monkey multi-unit recording experiment

Part of the monkey psychophysical and neurophysiological data have been published in refs. 14·47.

These previous studies showed qualitatively similar results of the learning-induced reduction in

Fisher information, Fano factor, and noise correlations via different analysis methods. Other results

and analyses on the characteristics of population responses in this study (i.e., Figs. 7&8), especially

the validation of signal rotation and manifold warping mechanisms, as well as the stepwise

information analyses, are novel. We provide relevant methods as follows and more detailed

methods in Supplementary Note 3 to avoid cross-referencing.

Ethics statement and Data collection. The Newcastle University Animal Welfare Ethical

Review Board approved all procedures in this study. All experimental procedures were carried out

in accordance with the European Communities Council Directive RL 2010/63/EC, the US National

Institutes of Health Guidelines for the Care and Use of Animals for Experimental Procedures and

the UK Animals Scientific Procedures Act. This study included two male monkey monkeys (5 and

14 years of age).

Experimental preparation. The surgical procedure is described in ref. 48 and

Supplementary Note 3. The headpost and electrode implementations are also described in

Supplementary Note 3. Briefly, in monkey 1, two 4×5 grids of microelectrodes were implanted

in area V4; in monkey 2, one 5×5 grid was implanted in V4. These chronically implanted

electrodes allowed us to record population activity in area V4 over the course of visual training.

Importantly, we were able to record stably from a few small multi-unit clusters. The stability of

the recording is shown in ref. 14. Stable recording of multi-channel neuronal activity allows

analyses of changes in population responses induced by training.

Behavioral task and monkey training. The monkeys were trained in a contrast

discrimination task in which subjects were asked to decide whether the contrast of a test stimulus

was higher or lower as compared to that of a reference stimulus by making a saccade to one of two

distinct locations (Fig. 7B). On each trial, the subject first kept fixation on the center-of-screen for

512ms. After 539 ms of fixation, a vertically oriented reference Gabor stimulus with 30% contrast

was presented, centered at the V4 Receptive Field (RF) coordinates. The outer diameter of the

Gabor stimulus was truncated at 16° for monkey 1 and 14° for monkey 2. Following the Gabor

stimulus, monkey 2 experienced an interstimulus interval of 512 ms. In contrast, monkey 1

experienced a randomly chosen interstimulus interval, ranging from 512 to 1024 ms. During the

interstimulus interval, only the fixation dot was presented. A test stimulus was then presented for

512 ms. This test stimulus was identical in size and orientation to the reference stimulus but

differed in contrast, with the contrast level chosen pseudorandomly. The test stimulus was

followed by another blank period of 512 ms during which only the fixation dot was visible. After

the fixation dot, two target squares, one black and one white with a size of 0.5° in size, appeared

to the left and right of the location where the reference and test stimuli were previously presented.

The monkeys were cued to make a decision once the fixation dot disappeared. The monkeys were

required to make a saccade to the white square within a 2° × 2° window if the test stimulus had a

higher contrast than the reference stimulus. Conversely, they were expected to make a saccade to

the black square if the test stimulus had a lower contrast than the reference stimuli. A correct

saccade was rewarded with a fluid reward, while an incorrect saccade led to no reward and a 0.2 s

timeout period.

The two monkeys were first trained on an easy version (target contrast 5% or 90%) of the

contrast discrimination task. After they were fully familiar with the easy task, the target contrast

increased from 2 to 8, 12, and 14 levels. The data correspond to the 14 levels of target contrast (10,

15, 20, 25, 27, 28, 29, 31, 32, 33, 35, 40, 50 or 60%; Supplementary Note 3). We only focus on

target contrast levels (27, 28, 29, 31, 32, and 33%) near the reference contrast (i.e., 30%) according

to the definition of linear Fisher information.

Dataset and preprocessing. We used chronically implanted Utah arrays to record spiking

activity. We refer to small multi-unit neuronal clusters recorded from a given electrode as

'channels'. Twenty-nine and Twenty channels were recorded in monkey 1 and monkey 2,

respectively. These channels exhibited good responses (signal-to-noise ratio [SNR] >1) on over

80% of the recording sessions (see SNR computation in Supplementary Note 3). Baseline activity

matching was performed between sessions for multi-unit activity (MUA) data in order to obtain

comparable activity levels across sessions.

Behavioral and neural analyses. We noticed that the relationship between neural activity
and discriminability can change drastically during the stimulus presentation period, and through
training, the improvement in discriminability can also vary over the course of the training period.
We chose the first 4 and the last 4 training sessions as the early and the late phase of training. This
choice ensures an overall sufficient and comparable number of trials at both pre- and post-test for
further analyses.

To determine the time window, we systematically varied the time window and trained a
linear classifier to discriminate between the reference and target stimuli, and obtained its
performance through 10-fold cross-validation. We chose the time window with the largest change
in decoding accuracy between the reference stimulus (30% contrast) and the target stimuli (29%
or 31% contrast). For monkey 1, the chosen time window was 30-130 ms after stimulus onset. For
monkey 2, the time window was 130-230 ms after stimulus onset. Note that this choice aims to
maximize training effects on population representations (similar to the decoding analyses for first
identifying V3A and hMT+ as the ROIs where learning effects are most pronounced in the human
fMRI study), but does not guarantee the underlying mechanisms such as signal separation
enhancement and manifold shrinkage. Also, varying the time window did not qualitatively change
our results. We used a simple multivariate Poisson log-normal (MPLN) model (Supplementary
Note 3, see also refs. 49-52) to estimate the trial-by-trial variability of population firing rates. We
further use the estimated firing rates and covariance to compute all neural metrics mentioned above.
We report all results in Figs. 7&8 for visual comparison with the DCNN and fMRI results above.

915
916
CONFLICT OF INTERESTS
XC is a cofounder and shareholder of a neurotechnology start-up, Phosphoenix (Netherlands).
Other authors declare no competing financial interests.

ACKNOWLEDGEMENT
We thank Shuguang Kuai, Duje Tadin, Oh-Sang Kwon for valuable comments on the manuscripts.
This work was supported by the National Natural Science Foundation of China (32100901 to R.-

Y.Z, 3230085 to K.J.) and Natural Science Foundation of Shanghai (21ZR1434700 to R.-Y.Z),

The monkey work was supported by the Medical Research Council, UK, G0700976.

AUTHOR CONTRIBUTIONS

Y-A. C. and R-Y.Z. conceived and designed the study. Y-A. C. implemented the neural networks.

J.K., S.L., and F.F., prepared and provided the preprocessed fMRI data. A.T., X.C., and M.S.

recorded, organized, and preprocessed the monkey physiological data. Y-A. C. and R-Y.Z.

performed in-depth analyses on neural networks, human fMRI data, and monkey

electrophysiological data. Y-A. C. and R-Y.Z. wrote the first draft of the manuscript. All authors

revised the manuscript and provided valuable feedback to the final manuscript.

RFERENCES

1   Lu, Z. L., Yu, C., Watanabe, T., Sagi, D. & Levi, D. Perceptual learning: functions,
mechanisms, and applications. *Vision Res*. **50**, 365-367 (2010).
2   Watanabe, T. & Sasaki, Y. Perceptual learning: toward a comprehensive theory. *Annu*
Rev Psychol. **66**, 197-221 (2015).
3   Chen, N. *et al.* Sharpened cortical tuning and enhanced cortico-cortical communication
contribute to the long-term neural mechanisms of visual motion perceptual learning.
Neuroimage. **115**, 17-29 (2015).
4   Shibata, K. *et al.* Decoding reveals plasticity in V3A as a result of motion perceptual
learning. *PLoS One*. **7**, e44003 (2012).
5   Jia, K. *et al.* Recurrent Processing Drives Perceptual Plasticity. *Curr Biol*. **30**, 4177-4187
e4174 (2020).
6   Lu, Z. L. & Dosher, B. A. Characterizing observers using external noise and observer
models: assessing internal representations with external noise. *Psychol Rev*. **115**, 44-82
(2008).
7   Dosher, B. A. & Lu, Z. L. Perceptual learning reflects external noise filtering and internal
noise reduction through channel reweighting. *Proc Natl Acad Sci U S A*. **95**, 13988-13993
(1998).
8   Dosher, B. & Lu, Z. L. Visual Perceptual Learning and Models. *Annu Rev Vis Sci*. **3**, 343-
363 (2017).
9   Schoups, A., Vogels, R., Qian, N. & Orban, G. Practising orientation identification
improves orientation coding in V1 neurons. *Nature*. **412**, 549-553 (2001).
10   Yang, T. & Maunsell, J. H. The effect of perceptual learning on neuronal responses in
monkey visual area V4. *J Neurosci*. **24**, 1617-1626 (2004).
11   Ni, A. M., Ruff, D. A., Alberts, J. J., Symmonds, J. & Cohen, M. R. Learning and
attention reveal a general relationship between population activity and behavior. *Science*.
359, 463-465 (2018).
12   Gu, Y. *et al.* Perceptual learning reduces interneuronal correlations in macaque visual
cortex. *Neuron*. **71**, 750-761 (2011).
13   Yan, Y. *et al.* Perceptual training continuously refines neuronal population codes in
primary visual cortex. *Nat Neurosci*. **17**, 1380-1387 (2014).
14   Sanayei, M. *et al.* Perceptual learning of fine contrast discrimination changes neuronal
tuning and population coding in macaque V4. *Nat Commun*. **9**, 4238 (2018).
15   Kohn, A., Coen-Cagli, R., Kanitscheider, I. & Pouget, A. Correlations and Neuronal
Population Information. *Annu Rev Neurosci*. **39**, 237-256 (2016).
16   Moreno-Bote, R. *et al.* Information-limiting correlations. *Nat Neurosci*. **17**, 1410-1417
(2014).
17   Zhang, R. Y., Wei, X. X. & Kay, K. Understanding multivariate brain activity:
Evaluating the effect of voxelwise noise correlations on population codes in functional
magnetic resonance imaging. *PLOS Computational Biology*. **16**, e1008153 (2020).
18   Cheng, Z. J., Yang, L., Zhang, W. H. & Zhang, R. Y. Representational Geometries
Reveal Differential Effects of Response Correlations on Population Codes in
Neurophysiology and Functional Magnetic Resonance Imaging. *J Neurosci*. **43**, 4498-
4512 (2023).
19   Wenliang, L. K. & Seitz, A. R. Deep Neural Networks for Modeling Visual Perceptual
Learning. *J Neurosci*. **38**, 6028-6044 (2018).

20 Krizhevsky, A., Sutskever, I. & Hinton, G. E. Imagenet classification with deep
convolutional neural networks. *Advances in neural information processing systems*. **25**,
1097-1105 (2012).
21 Dosher, B. A. & Lu, Z. L. Mechanisms of perceptual learning. *Vision Res*. **39**, 3197-3221
(1999).
22 Chen, N., Cai, P., Zhou, T., Thompson, B. & Fang, F. Perceptual learning modifies the
functional specializations of visual cortical areas. *Proc Natl Acad Sci U S A*. **113**, 5724-
5729 (2016).
23 Raiguel, S., Vogels, R., Mysore, S. G. & Orban, G. A. Learning to see the difference
specifically alters the most informative V4 neurons. *J Neurosci*. **26**, 6589-6602 (2006).
24 Ghose, G. M., Yang, T. & Maunsell, J. H. Physiological correlates of perceptual learning
in monkey V1 and V2. *J Neurophysiol*. **87**, 1867-1888 (2002).
25 Jehee, J. F., Ling, S., Swisher, J. D., van Bergen, R. S. & Tong, F. Perceptual learning
selectively refines orientation representations in early visual cortex. *J Neurosci*. **32**,
16747-16753a (2012).
26 Jeanne, J. M., Sharpee, T. O. & Gentner, T. Q. Associative learning enhances population
coding by inverting interneuronal correlation patterns. *Neuron*. **78**, 352-363 (2013).
27 Ruff, D. A. & Cohen, M. R. Attention can either increase or decrease spike count
correlations in visual cortex. *Nat Neurosci*. **17**, 1591-1597 (2014).
28 Hennequin, G., Ahmadian, Y., Rubin, D. B., Lengyel, M. & Miller, K. D. The Dynamical
Regime of Sensory Cortex: Stable Dynamics around a Single Stimulus-Tuned Attractor
Account for Patterns of Noise Variability. *Neuron*. **98**, 846-860 e845 (2018).
29 Ecker, A. S. *et al.* Decorrelated neuronal firing in cortical microcircuits. *Science*. **327**,
584-587 (2010).
30 Gutnisky, D. A. & Dragoi, V. Adaptive coding of visual information in neural
populations. *Nature*. **452**, 220-224 (2008).
31 Huang, X. & Lisberger, S. G. Noise correlations in cortical area MT and their potential
impact on trial-by-trial variation in the direction and speed of smooth-pursuit eye
movements. *J Neurophysiol*. **101**, 3012-3030 (2009).
32 van Bergen, R. S. & Jehee, J. F. M. Modeling correlated noise is necessary to decode
uncertainty. *Neuroimage*. **180**, 78-87 (2018).
33 Liu, Z. Perceptual learning in motion discrimination that generalizes across motion
directions. *Proc Natl Acad Sci U S A*. **96**, 14085-14087 (1999).
34 Ball, K. & Sekuler, R. Direction-specific improvement in motion discrimination. *Vision
Res*. **27**, 953-965 (1987).
35 Law, C. T. & Gold, J. I. Neural correlates of perceptual learning in a sensory-motor, but
not a sensory, cortical area. *Nat Neurosci*. **11**, 505-513 (2008).
36 Tran, D., Bourdev, L., Fergus, R., Torresani, L. & Paluri, M. in *Proceedings of the IEEE
international conference on computer vision*. 4489-4497.
37 Jia, K. *et al.* Visual perceptual learning modulates decision network in the human brain:
The evidence from psychophysics, modeling, and functional magnetic resonance
imaging. *J Vis*. **18**, 9 (2018).
38 Thomson, H. L. *et al.* Haemodynamics and left ventricular mass regression: a comparison
of the stentless, stented and mechanical aortic valve replacement. *Eur J Cardiothorac
Surg*. **13**, 572-575 (1998).

39 Cohen, M. R. & Maunsell, J. H. Attention improves performance primarily by reducing
interneuronal correlations. *Nat Neurosci*. **12**, 1594-1600 (2009).
40 Reimer, J. *et al.* Pupil fluctuations track fast switching of cortical states during quiet
wakefulness. *Neuron*. **84**, 355-362 (2014).
41 Lu, Z. L., Chu, W. & Dosher, B. A. Perceptual learning of motion direction
discrimination in fovea: separable mechanisms. *Vision Res*. **46**, 2315-2327 (2006).
42 Britten, K. H., Shadlen, M. N., Newsome, W. T. & Movshon, J. A. The analysis of visual
motion: a comparison of neuronal and psychophysical performance. *J Neurosci*. **12**,
4745-4765 (1992).
43 Huk, A. C., Dougherty, R. F. & Heeger, D. J. Retinotopy and functional subdivision of
human areas MT and MST. *J Neurosci*. **22**, 7195-7205 (2002).
44 Engel, S. A., Glover, G. H. & Wandell, B. A. Retinotopic organization in human visual
cortex and the spatial precision of functional MRI. *Cereb Cortex*. **7**, 181-192 (1997).
45 Sereno, M. I. *et al.* Borders of multiple visual areas in humans revealed by functional
magnetic resonance imaging. *Science*. **268**, 889-893 (1995).
46 Kay, K. N., Rokem, A., Winawer, J., Dougherty, R. F. & Wandell, B. A. GLMdenoise: a
fast, automated technique for denoising task-based fMRI data. *Front Neurosci*. **7**, 247
(2013).
47 Chen, X., Sanayei, M. & Thiele, A. Perceptual learning of contrast discrimination in
macaca mulatta. *J Vis*. **13**, 22 (2013).
48 Thiele, A., Delicato, L. S., Roberts, M. J. & Gieselmann, M. A. A novel electrode-pipette
design for simultaneous recording of extracellular spikes and iontophoretic drug
application in awake behaving monkeys. *J Neurosci Methods*. **158**, 207-211 (2006).
49 Goris, R. L., Movshon, J. A. & Simoncelli, E. P. Partitioning neuronal variability. *Nat
Neurosci*. **17**, 858-865 (2014).
50 Henaff, O. J. *et al.* Primary visual cortex straightens natural video trajectories. *Nat
Commun*. **12**, 5982 (2021).
51 Rabinowitz, N. C., Goris, R. L., Cohen, M. & Simoncelli, E. P. Attention stabilizes the
shared gain of V4 populations. *Elife*. **4**, e08998 (2015).
52 Zhu, R. J. B. & Wei, X. X. Unsupervised approach to decomposing neural tuning
variability. *Nat Commun*. **14**, 2298 (2023).
1057