Can a flower color ancestral polymorphism transcend speciation?

Mercedes Sánchez-Cabrera[1], Eduardo Narbona[2], Montserrat Arista[1], Pedro L. Ortiz[1],

Francisco J. Jiménez-López[3], Amelia Fuller[4], Benjamin Carter[5], and Justen B. Whittall[6*]

[1]Department of Plant Biology and Ecology, Universidad de Sevilla, Seville, Spain

41012

[2]Department of Molecular Biology and Biochemical Engineering, Pablo de Olavide

University, Seville, Spain 41013

[3]Departamento de Biología, Geología, Física y Química Inorgánica, Universidad Rey

Juan Carlos, Móstoles, Spain

[4]Department of Chemistry and Biochemistry, Santa Clara University, Santa Clara, CA

95053

[5]Department of Biological Sciences, San Jose State University, San Jose, CA 95182

[6]Department of Biology, Santa Clara University, Santa Clara, CA 95053

[*]**Corresponding Author:** jwhittall@scu.edu

Abstract

Polymorphisms are common in nature, but they are rarely shared among closely related

species. They could originate through convergence, ancestral polymorphism, or

introgression. Although shared neutral genomic variation is commonplace, very few

examples of shared functional traits exist. The blue-orange petal color polymorphisms

in two closely related species, *Lysimachia monelli* and *L. arvensis* were investigated

with UV-vis reflectance spectra, flavonoid biochemistry, and transcriptome

comparisons followed by climate niche analysis. The similarities in reflectance spectra,

biochemistry, and transcriptomes suggest a single shift from blue-to-orange shared by

both lineages is possible. Transcriptome comparisons reveal two orange-specific genes

are directly involved in both blue-orange color polymorphisms: *DFR-2* specificity

redirects flux from the malvidin to the pelargonidin while *BZ1-2* stabilizes the

pelargonidin with glucose, producing the orange pelargonidin 3-glucoside. The climate

niches for each color morph are the same between the two species for three temperature

characteristics but differ for four precipitation variables. We suggest that this persistent

flower color polymorphism may represent an ancestrally polymorphic trait that has

transcended speciation with some unique ecological effects.

Keywords: ancestral polymorphism, anthocyanin, flavonoid, *Lysimachia arvensis*,

Lysimachia monelli, petal transcriptome

Introduction

Since the Modern Synthesis, we have assumed that most phenotypic variation emerges within populations and is subsequently acted upon by evolutionary forces within species (Dobzhansky, 1937; Grant, 1981). Herein, we describe an ancestral flower color polymorphism that may have transcended the species boundary providing a concrete example of unsorted ancient phenotypic variation in two species with contrasting ecological effects. This study addresses how shared traits evolve across lineages, a central topic in understanding the evolution of biological diversity.

Polymorphisms, or variation among individuals of the same species, provide rare windows into the process of adaptation and speciation (Allison, 1961; Igić et al., 2008; Martin et al., 2013; Comeault et al., 2015). Polymorphisms among individuals in a population or among populations of a single species, are omnipresent across the tree of life (McKinnon & Pierotti, 2010; McLean & Stuart-Fox, 2014) and can be maintained by a diversity of forces including negative frequency-dependent selection (Madsen et al., 2022), heterozygote advantage (Sirugo et al., 2014), genetic drift (Wright, 1943, but see Schemske & Bierzychudek, 2007), gene flow (Martin et al., 2013) and spatially or temporally variable selection (Svensson, 2017). Some polymorphisms are phylogenetically dispersed (e.g. bird plumage color or flower color; Hugall & Stuart-Fox, 2012), while in others reoccur in very closely related species (e.g. heterostyly in *Primula*, shell chirality in *Amphidromus*, cryptic body color in *Timema*, and wing patterning in *Heliconius*; reviewed in Jamie & Meier, 2020). The molecular underpinnings, persistence mechanics and evolutionary forces acting on these rare cases of shared variation are largely unexplored.

Some studies have proposed that polymorphisms promote the utilization of a wide range of environmental resources and can act as a precursor to speciation which then become fixed after divergence (Corl et al., 2010; Hugall & Stuart-Fox, 2012; McLean & Stuart-Fox, 2014). However, polymorphisms that persist across species (Jamie & Meier, 2020) represent evolutionary enigmas. How can polymorphisms transcend species boundaries? Convergent evolution (Gould, 1989; Conway, 2005) including developmental convergence (i.e. independent changes in gene expression), introgression (Mallet, 2005; Nolte et al., 2009), and the maintenance of an ancestral polymorphism (Igić & Kohn,

2001; Igić et al., 2008) are three plausible evolutionary avenues leading to shared

polymorphisms among closely related species.

Color polymorphisms are widespread in nature but only a few cases of trans-specific

color polymorphisms have been reported, and mostly restricted to animals (reviewed in

McKinnon & Pierotti, 2010; Martin et al., 2013; McLean & Stuart-Fox, 2014; McRobie

et al., 2019). Few examples have been investigated in plants. The closest situation for

flower color is the repeated transitions from blue to red flowers (Wessinger & Rausher,

2013) associated with shifts from bee- to hummingbird-pollination (Thompson &

Wilson, 2008; Muchhala et al., 2014; Wessinger et al., 2019) across a diversity of

angiosperm lineages. However, in this case intraspecfic blue-red flower color

polymorphisms (variation within populations) are very rare since either (1) multiple

mutations are required to accomplish this shift and/or (2) the color is associated with a

pollinator shift leading to reproductive isolation between the color types thereby fixing

the color differences in newly formed species (but see Stankowski and Streisfeld 2015).

Unlike the blue to red flower color transitions leading to speciation, the blue-orange

flower color polymorphism in the closely related species *Lysimachia arvensis* and *L.*

monelli (*La* and *Lm* herein, Figure 1A) does not confer a pollinator shift (Jiménez-

López et al., 2019; Sánchez-Cabrera, 2023). Instead, in *La* the color polymorphism is

driven by abiotic non-pollinator agents of selection such as drought and sunlight

intensity (Arista et al., 2013, Ortiz et al., 2015) and temperature (Jiménez-López et al.,

2023; B. Carter et al., unpublished data) where blue flowered individuals are fitter than

orange ones in drier, sunnier and hotter environments. *La*, is an annual native to the

Mediterranean Basin and central and northern Europe. A recent autopolyploid origin has

been suggested for *La* for three reasons: (1) due to the lack of ITS polymorphisms

(Jiménez-López et al., 2022), (2) the presence of four identical copies of one set of

chromosomes (Monein et al., 2003), (3) and the phylogenetic configuration of the group

(Jiménez-López et al., 2022). Color morph frequencies range from 0% to 100% with a

steep cline in central-southern Mediterranean region where mixed populations are

common (e.g., Portugal, Spain, Italy and France; Arista et al., 2013). The cline and

resulting association with abiotic forces strongly argue against pollinator mediated

selection and instead point to abiotic factors which have been substantiated in a

greenhouse study (Arista et al., 2013). In this species, the shift from blue to orange

(Sánchez-Cabrera et al., 2021) correlates with a biochemical transition from malvidin 3-

rhamnoside to pelargonidin 3-glucoside (Harborne, 1968; Sánchez-Cabrera et al., 2021).

Orange flowers have increased expression of *DFR-2*, a duplicate gene only found in

orange-flowered individuals with non-synonymous SNPs suggesting substrate

specificity for dihydrokaempferol drawing flux down the pelargonidin branch of the

anthocyanin biosynthetic pathway (ABP) (Sánchez-Cabrera et al., 2021).

Herein, we investigate the shared blue-orange flower color polymorphism in the closely

related *Lm*, a diploid perennial, that co-occurs with *La* in some populations (Figure 1).

Blue flowered populations of *Lm* are more common in drier habitats of central and

southwestern Iberian Peninsula and eastern North Africa; while orange-flowered

populations are more often found in wetter habitats of northeastern Iberian Peninsula,

western North Africa, and Sardinia (Gibbs & Talavera, 2001; Sánchez-Cabrera, 2022)

(Figure 1A). In *Lm*, no populations have both color morphs (Sánchez-Cabrera, 2023).

To determine the origin and maintenance of the shared blue-orange flower color

variation in this species of *Lysimachia* and how it relates to the strikingly similar blue-

orange flower color variation in *La*, we investigate the similarities in petal reflectance

spectra, flavonoid biochemistry and compare petal transcriptomes allowing us to weigh

four possible hypotheses explaining the origin and persistence of this flower color

polymorphism: convergence, introgression, ancestral polymorphism or the non-

monophyly of species (Figure 2).

Results

Petal UV-Vis Spectra

Reflectance spectra of petals for both species are very similar across the UV and visible

wavelengths when comparing the same color morphs, yet quite different between color

morphs (Figure 1B; Appendix 1—figure 1). Spectra distances between color morphs are

~7x larger than spectra distances within morphs between species (Permutational

Manova, $R^2 = 0.75$ vs. 0.15; Appendix 1—figure 1). Orange petals have double UV

reflectance peaks that are absent in blue petals and differ in their primary inflection

points (380, 570nm for orange and 480, 660nm for blue).

Petal Biochemical Comparisons

Biochemical analyses of petal extracts in acidified methanol using HPLC-MS-DAD

revealed the underlying biochemical differences between blue and orange morphs are

nearly identical between the two species. Blue petals of both species are composed of

malvidin 3-rhamnoside, however small amounts of the aglycones malvidin and

delphinidin, and a flavonol derivative were also detected in both species (Figure 1C;

Appendix 1—table 1; Sánchez-Cabrera et al., 2021). In contrast, orange petals of both

species accumulate principally pelargonidin 3-glucoside and to a lesser degree, other

pelargonidin derivatives. There were no significant differences in the relative content of

malvidin 3-rhamnoside and pelargonidin 3-glucoside derivatives when comparing the

blue and orange samples of these two species, respectively (Figure 1C).

Transcriptome comparisons

Differential expression

Transcriptome results for *Lm* petals identified 37,552 distinct genes (see Sánchez-

Cabrera et al., 2021 for *La* results including the absence of any evidence of divergent

homologous copies of these genes). Of those, 336 were differentially expressed between

blue and orange flowers (1.19x more genes with O > B expression; Chi-Square =

14307; p = < 2.2e-16). A total of 128 flavonoid biosynthetic pathway (FBP) genes were

detected, although only five had significant differential expression, four of which had O

> B expression in *Lm* (Appendix 1—table 2), three of which are in the core ABP (Figure

3C)

In comparing FBP gene expression in *Lm* to *La,* three genes (*DFR-2*, *BZ1-2* and

Caffeoyl CoA-1) were significantly differentially expressed in both species (FDR<10^{-5}

and log$_2$FC>1; Figure 3C; Appendix 1—table 2; see Sánchez-Cabrera et al., 2021). Two

key ABP genes had high expression in orange petals and were nearly undetectable in

blue petals (Figure 3A, C). First, *DFR-2* with substrate specificity for pelargonidin (see

references in Sánchez-Cabrera et al., 2021) had high expression in orange petals

(*LmDFR-2* orange mean = 157.18; *LaDFR-2* orange mean = 99.80; Figure 3A). The

second is *BZ1-2*, a gene responsible for glycosylation that stabilizes the orange pigment

(*LmBZ1-2* orange mean = 2651.84; *LaBZ1-2* orange mean = 294.55; Figure 3A).

However, the reciprocal decrease in expression for the primarily blue paralogue (*DFR-1*

and *BZ1-1*) in orange flowers was only significant in *La*, but trends in the same

direction at similar magnitude for *LmDFR-1* (blue is ~2x higher than orange) (Figure

3A). The third gene, *Caffeoyl CoA-1*, is an early gene in the ABP that functions before

the first dedicated step in the ABP (not previously reported to be involved in this type of

color change; Appendix 1—figure 2B). It had higher expression in blue petals of both

species and was nearly undetectable in orange petals of both species - the opposite

pattern as that described above for *DFR-2* and *BZ1-2*. In addition to not having known

function in the biochemistry of blue and orange, this gene is also unlikely responsible

for the color change because this copy of Caffeoyl CoA has the lowest expression of the

three copies expressed in petals (Appendix 1—figure 2B).

Some additional ABP genes likely involved in this color shift showed significant

differential expression between colors of *La* and consistent, yet non-significant trends,

in *Lm*. For example, *DFR-1* and *F3'5'H* had higher expression in blue petals than in

orange petals in both species, but was only significant in *La* (2.10x and 2.45x more in

*La*B and 1.80x and 4.71x more in *Lm*B, respectively; Figure 3A). Another example of

an ABP gene with significant DEG in *La* and a similar trend, only significant with a

Mann-Whitney U test, in *Lm* is *MYB4-2* which has higher expression in orange petals

compared to blue petals (25.75x more in *La*O and 3.83x more in *Lm*O; Figure 3A).

Alternatively, *CHS* showed significant differential expression between colors of *Lm*, but

not significant nor consistent in *La* (1.95x more in *La*B, but 2.36x more in *Lm*O; Figure

3A) and is unlikely involved in the transition from blue to orange. Finally, a third *DFR*

(*DFR-3*; Figure 4) was detected in both species, but exhibited very low expression

(maximum TMM = 1.07) indicating it is not the primary gene copy at this step in the

ABP in petals (*DFR-1* and *DFR-2* have >100x higher expression).

Phylogenetic analysis of ABP sequences

We compared coding sequences of a total of 22 ABP genes that were either

differentially expressed in one of the species or potential candidate genes to determine

whether species were monophyletic (convergence) or if color was monophyletic and

transcended speciation (ancestral polymorphism). Individual phylogenetic analyses for

all 22 ABP genes (10 structural and 12 regulatory) consistently recovered a

monophyletic clade of *Lysimachia* samples suggesting these were broadly orthologous

comparisons (>70% bootstrap support; Appendix 1—table 3 and figure 2). For two key

ABP genes in the transition from blue to orange (*DFR-2* and *BZ1-2*), all orange samples

of both species are strongly supported as monophyletic (99% and 100% bootstrap,

respectively; Figure 4; Appendix 1—table 3 and figure 2A). Furthermore, the regulatory

locus, *bHLH12* is reciprocally monophyletic for species and within species color

morphs (Appendix 1—figure 2J). In a combined phylogenetic analysis of all ABP loci,

we recovered monophyletic species and monophyletic colors within each species except

for paraphyletic *Lm*O (Appendix 1—figure 3).

SNP analysis

To determine if there is an association between genotype and phenotype, we examined

SNPs across the FBP loci for all samples. Genes responsible for color differences in one

or both species could have non-synonymous (NS) SNPs in functional regions that

correlate with color within species if causal changes are in coding regions. Further, if

these are shared between species this would provide further evidence supporting a

single origin (ancestral polymorphism). We detected the exact same 13 NS SNPs in

DFR-2 of orange *Lm* that differentiate it from *DFR-1* which overlap with known

functional regions providing substrate specificity for pelargonidin (Figure 3B) as we

detected previously in *La* (see references in Sánchez-Cabrera et al., 2021).

Broadening our analysis to all FBP/ABP genes examined, only two loci have

completely fixed differences between color morphs in both species (one NS SNP in

DFR-1 and two NS SNPs in *MYB4-2*; Appendix 1—table 4), but none are located in

known functional sites.

By comparing NS SNPs that differentiate the color morphs with the NS SNPs that

differentiate the two species, we can begin to hone in on the loci responsible for the

color change to eventually address if it was a single event (ancestral polymorphism or

introgression) or more than one event (convergence). We find 16 NS SNPs that

differentiate the two species in eight genes that we don't predict are causing the color

change (Appendix 1—table 4). Noticeably absent from this list are the loci we have

evidence are involved in the color shift (*DFR-1*, *DFR-2*, *BZ1-2*, *MYB 4-2*), and only one

NS SNP in the extreme 5' region of *F3'5'H* (Appendix 1—table 4). Only *F3'H* showed

color-specific NS SNPs, higher in *Lm* than in *La* (five vs. one), however there are even

more NS SNPs between the species (seven) (Appendix 1—table 4).

Structural and regulatory gene interaction

To help discern if the shared expression differences in orange petals of the two species for *DFR-2* and *BZ1-2* are controlled by the same or different regulatory genes, we looked for patterns of correlated expression with the MYB-bHLH-WD40 regulatory complex. The rationale for investigating expression correlations is that if color has evolved via independent changes in expression (i.e. developmental convergence), then we expect gene expression correlations in the gene involved to be species specific. Using a hypothesis testing approach (alpha = 0.05) in both species, *DFR-2* and *BZ1-2* were significantly positively correlated with one another (R = 0.73, p = 8.8e-05 in *La*; R = 0.62 and p = 2.9e-04 in *Lm*) and with *TTG1*, a WD40 known to activate ABP structural genes (across both genes and species p < 0.0051; Appendix 1—figure 4). Furthermore, *DFR-1* expression is negatively correlated with *MYB4-2* expression, a known repressor of ABP structural genes (R = -0.68 and p = 2.5e-04 in *La*; R = -0.62 and p = 1.4e-04 in *Lm*; Appendix 1—figure 4). *MYB4-2* is positively correlated with *DFR-2* (R = 0.39 and p = 0.37 in *La*; R = 0.47 and p = 0.0062 in *Lm*), but non-significant after Bonferroni correction.

Reproductive isolation between *La* and *Lm*

None of the crosses performed between *La* and *Lm* produced fruits irrespective of crossing direction and flower color (n=53 pollinations, see Appendix 1—table 5). In contrast, morphs within species were interfertile. Between morph crosses in *Lm* produced a mean fruit-set of 47.86% ± 0.39% (n=575) and a mean number of seeds per fruit of 16.25 ± 9.16 (n=287) (Sánchez-Cabrera, 2023). Whereas in *La*, 100% of between morph crosses produced fruits containing 16.95 ± 7.19 seeds per fruit (n= 97) (Jiménez-López et al., 2023).

Climate niche modeling

The four morphs were statistically different from one another across most of the 19 BIOCLIM variables examined (Appendix 1—table 6). The orange and blue *La* morphs differed significantly from each other for 17 of the 19 climate variables, and the two *Lm* morphs differed for nine variables. The two orange morphs differed from one another for 16 variables and the two blue morphs differed across 11 variables.

Nine variables were significantly different between the two blue morphs and between the two orange morphs of both species. Of these, two temperature variables differed in

parallel (e.g., for "isothermality" orange morphs showed lower values than blue morphs

for both species) (Appendix 1—figure 5A), while three temperature and four

precipitation variables differed in opposing directions (e.g., for "annual precipitation"

orange *La* were wetter than blue *La*, but orange *Lm* were drier than blue *Lm*) (Appendix

1—figure 5C).

The logistic regression results showed that the two best models using AIC did not

employ any of the same variables. The best model for *La* included (in decreasing order

of importance based on standardized coefficients) precipitation seasonality,

isothermality and temperature seasonality, while the model for *Lm* consisted of

precipitation of the wettest month and mean temperature of the wettest quarter

(Appendix 1—table 7).

Since the ancestral state in *Lysimachia* is blue flowers, we hypothesized that blue would

be less likely to diverge in their climate niche between the two species compared to

orange morphs of the two species. We tested this with a Monte Carlo procedure that

randomized colors within species and then compared observed climatic differences with

differences from the null distributions. Across the 19 BIOCLIM variables, blues were

more divergent between the two species than expected by chance for only one variable

while orange samples were more divergent between the two species for 15 variables

(~79% of variables studied), and neither color morph was significantly different for

three variables (Appendix 1—table 8).

Discussion

There are many similarities (and some differences) between color morphs of *L. arvensis*

and *L. monelli* that can be used to assess the alternative hypotheses (Table 1), however

definitive evidence excluding all but one of these hypotheses remains elusive. If orange

evolved independently in each species (i.e. convergence; Figure 2A), we would expect

distinct molecular causes (Streisfeld & Rausher, 2009; Larter et al. 2018; Ng & Smith,

2016), which we don't have any strong evidence for in these two species of *Lysimachia*,

but the exact mutation(s) responsible for the color shift in each species awaits further

genetic dissection. If the remarkably similar upregulation of *DFR-2* was activated by

different MYBs in orange petals of these two species, we would expect distinct

correlations in expression. Instead, *DFR-2* and *MYB4-2* are positively correlated in both

species (but not significant after Bonferroni correction; Figure S5). Alternatively, the likelihood of a single shift from blue to orange followed by introgression into the other species (Figure 2B) as clearly documented in the *Diplacus* (*Mimulus*) *aurantiacus* complex (Stankowski and Streisfeld 2015, Short and Streisfeld 2023) is unlikely in *Lysimachia* for two reasons. First, although the two species co-occur, hybrids with intermediate flower size and plant size have never been observed (Pujadas, 1997) unlike in the *Diplacus aurantiacus* complex where hybrid zones are well documented. Second, numerous attempted crosses of the two species reveal complete reproductive isolation (Table S7), presumable because of the ploidy differences – whereas the members of the *Diplacus aurantiacus* complex are at least partially interfertile (Stankowski and Streisfeld 2015). But, we can only test for the likelihood of genetic exchanges between the two species in the present – there is always the possibility that there was some gene exchange soon after the (auto)polyploid speciation event. Even in that scenario, the single origin followed by introgression would require the long-term persistence of this polymorphism even though it may not have transcended the speciation event.

Given the evidence presented herein, the most parsimonious explanation starts with a single origin of orange from blue in a common ancestor of *La* and *Lm* - creating an ancestral polymorphism that then transcends the speciation event separating these two closely related species (Figure 2C). In this case, both blue and orange individuals would have persisted through the speciation process including polyploidy, loss of self-incompatibility, significant reduction in flower size, and a shift from perennial to annual. Although the persistence of polymorphisms across species has been documented in a few exceptional animals (see Jamie & Meier, 2020 for a review), this is the first example in plants with regards to a flower color polymorphism, however an analogous pattern of trans-specific, shared polymorphism is well documented for self-incompatibility alleles (Igić et al. 2008). Furthermore, this flower color example appears to be driven by non-pollinator agents of selection – specifically the distinct climatic niches in each species, especially those involving precipitation variables (Arista et al., 2013; Ortiz et al., 2015; Sánchez-Cabrera, 2023), yet does not appear to be maintained by negative frequency dependent selection, the most common mechanism maintaining polymorphisms over long periods of time in other species of plants and animals.

A final alternative to ancestral polymorphism could involve a more complex

evolutionary history where these two species are not reciprocally monophyletic

(Jiménez-López et al., 2022; Figure 2D). This is unlikely given the combined

phylogenetic analysis of 22 ABP loci which strongly supports the monophyly of both

species (Appendix 1—figure 3). Moreover, in this scenario, a single origin of orange

would have to be followed by repeated origins of polyploidy, independent losses of self-

incompatibility, recurrent reduction in flower size and separate shifts to the annual habit

which is substantially less parsimonious than the persistence of an ancestral

polymorphism (Figure 2C vs. D).

The color morphs of these two *Lysimachia* species have nearly identical biochemical

and molecular underpinnings as measured thus far. The evolution of orange

pelargonidin 3-glucoside from blue malvidin 3-rhamnoside requires (1) redirecting ABP

flux down the pelargonidin branch and (2) stabilizing the newly formed orange

anthocyanidin with glucose instead of rhamnose. Both steps are accomplished in the

same way in both species - via the recruitment of orange-specific paralogues that are

undetectable in transcriptomes of blue petals. In fact, the copy of *DFR* unique to orange

petals (*DFR-2*) has nearly the same overall gene expression levels as the shared copy

(*DFR-1*), but its expression is undetected in blue petals of both species. Furthermore,

DFR-2 of both species contain the same 13 NS SNPs in the substrate specificity region

that differentiate it from *DFR-1*, which likely allows it to outcompete *F3'5'H* (and

F3'H) for dihydrokaempferol thereby shunting flux down the pelargonidin branch

leading to orange petals (Johnson et al., 2001; Petit et al., 2007; Katsu et al., 2017). The

substantial genetic distance between *DFR-1* and *DFR-2* suggests that this duplication

predated the genus and maybe even the Primulaceae family – a molecular toolkit

deployed when pelargonidin provides a selective advantage over malvidin. Although the

exact regulatory locus (loci) responsible for these shifts in expression will require

further genetic analysis, we hypothesize it involves *TTG1* (pelargonidin-specific

activator) and *MYB4-2* (malvidin-specific negative regulator and potential positive

activator of *DFR-2*). We find no evidence of independent NS SNPs in either *TTG1*, nor

MYB4-2, suggesting that the regulatory control of this flower color polymorphism has

only evolved once. Instead, we see remarkably similar expression patterns and even

some shared non-synonymous SNPs in both regulatory loci that are the most likely

candidates for the molecular basis for this color shift.

The coordinated changes in ABP gene expression is like other blue to red flower color

shifts that involve the recruitment of substrate specific *DFR* copies with correlated

downregulation of the alternative side-branch gene expression (e.g., *F3'H* and *F3'5'H*)

(Zufall & Rausher, 2004; Des Marais & Rausher, 2010). We also find decreased

expression in *F3'5'H* (but not *F3'H*) in orange petals of both species, however, this is

only significant in *La* due to high expression variation among the blue *Lm* samples

(Figure 3). Regardless, the shared expression and nearly identical coding sequences of

orange *DFR-2* in *La* and *Lm* are strong evidence of a single origin that predates this

speciation event and has persisted to the present in these distinct lineages.

Similarly, orange petals employ an orange-specific glycosyltransferase (*BZ1-2*),

undetected in blue petals, glycosylating the orange pigment in the same way in both

species. The shift from blue to orange is correlated with a dramatic increase in

expression of the glycosyltransferase *BZ1-2* (similar to the *DFR*s mentioned above),

which is likely involved in the stability, solubility, storage and biological activity of this

particular anthocyanin (Yonekura-Sakakibara et al., 2008). There are no NS SNPs

between orange samples of *La* and *Lm* (same evolutionary history), however two *BZ1*

paralogues are highly divergent (>60% nucleotide divergence) and unalignable

indicating a relatively ancient duplication event). The orange-specific *BZ1-2* of both

species contains NS SNPs correlating with glucose-specificity as characterized in *Vitis*

vinifera and *Medicago truncata* (He et al., 2006; Wang, 2009). In particular, there are

20 NS SNPs distinguishing these two paralogues in the PSPG-box, a conserved 44

amino acid region found in all plant UFGTs (Caputi et al., 2012) that likely confers

glucose-specificity of *BZ1-2*. We infer that the two paralogues likely have different

functions regarding which sugar they add and their efficiency in doing so (Sun et al.,

2016), but within a color morph for both species, their sequence similarity suggest they

perform the same function. In contrast to *BZ1-2*, *BZ1-1* has variable expression in both

color morphs - *La* has significantly higher expression in blue than orange as expected,

but mean expression is similar in blue and orange *Lm* (and not significant). Regardless

of whether there is compensatory downregulation in orange petals of *BZ1-1*, we know

that *BZ1-2* has 10-50x higher expression in orange and we predict that it has higher

efficiency in adding glucose to pelargonidin of orange petals than *BZ1-1* (Sun et al., 2016). Although there are other examples of closely related species exhibiting the same flower color because they accumulated similar major categories of anthocyanidins (i.e. aglycones) (e.g. Ng & Smith, 2016; Larter et al., 2018; Ogutcen et al., 2020), the specific anthocyanins they accumulate are generally distinct due to the enormous diversity of biochemical decorations in the flavonoids (Passeri et al., 2016; Berardi et al., 2021). In these two *Lysimachia* species, orange morphs use the same *3GT* correlated with the predominant glycosylation of pelargonidin clearly indicating a single ancestral evolutionary transition shared by both species that has transcended the speciation event.

If *DFR-2* and *BZ1-2* are similarly upregulated in orange *La* and *Lm*, we predict they will be controlled by the same regulatory gene(s) in both species if there was a single ancestral polymorphism. In fact, expression of *DFR-2* and *BZ1-2* in orange petals of both species is positively correlated with the same regulatory gene *TTG1* (WD40) known to control these two ABP structural genes (Tiang & Wang, 2020) (Appendix 1—figure 4). Looking at the sequence of this regulatory gene, there is only one color differentiating NS SNP in *La* and none in *Lm* strongly suggesting another trans-acting regulatory gene is likely responsible for the differential expression of *TTG1* in blue and orange morphs of these two *Lysimachia* species. However, *MYB4-2* has two NS SNPs shared by both species that positively correlates with *DFR-2* expression (Appendix 1—figure 4).

For both *Lysimachia* species, *DFR-1* expression in orange petals is negatively correlated with *MYB4-2* expression (Appendix 1—figure 4). Moreover, in *La* this increase of *MYB4-2* is correlated with a decrease in *F3'5'H* in orange petals, facilitating the redirection of ABP flux to the pelargonidin branch of the pathway. Although *MYB4-2* has low expression in both morphs of *Lm* and is not significantly correlated with *DFR-1* nor *F3'5'H*, it exhibited significant differential expression between morphs (Mann-Whitney U test, U = 19, p = 0.021). In *Arabidopsis thaliana* and *Vitis vinifera*, *MYB4* represses anthocyanin biosynthesis either through direct binding to the promoter regions of ABP genes (*ANS*, *DFR* and *UFGT*) or by displacing the activator MYB in the MBW complex (Pérez-Díaz et al., 2016; Chen et al., 2019). However, how could *MYB4-2* repress *DFR-1* and not *DFR-2*? Like in *Lotus* (Yoshida et al., 2010) and *Fagopyrum* (Katsu et al., 2017) where different *DFR* paralogues are independently regulated by

their unique promoter sequences, we predict that *MYB4-2* has differential regulatory effects on the two primary *DFR* paralogues. Overall, we argue that *MYB4-2* contributes to redirecting ABP flux down the pelargonidin branch of the pathway by negatively regulating the non-orange-specific paralogues at the critical branchpoint (*DFR-1* and *F3'5'H*) in both species similarly.

Previous experimental work and our climate niche modelling indicate that there may be pleiotropic effects of petal color. Orange morphs of both species are found in locations with colder winters than blue morphs (Appendix 1—figure 5A). However, the two morphs appear to respond in opposing directions to primarily moisture-related climate niche variables (Appendix 1—figure 5C). Orange *Lm* is found in habitats with a wide range of precipitation while orange *La* is found in wet habitats. Although the climate niche of the blue morphs of the two species are often distinct, the orange samples are clearly driving these orthogonal responses to precipitation. Why a biochemically and genetically similar polymorphism shared between different species would have contrasting ecological side-effects remains unclear. Previous experimental work in *La* shows that blue-flowered individuals (containing rhamnose stabilized malvidin) inhabit environments with lower precipitation and higher solar radiation than orange-flowered plants with glucose-bound pelargonidin, potentially linking abiotic stress with differential glycosylation (Arista et al., 2013; Sánchez-Cabrera, 2022). If *BZ1-2* is only found in orange petals and is glucose-specific, then does the type of sugar confer physiological or ecologically-relevant adaptations in *Lysimachia*? The coupling of rhamnose to malvidin found in *Lysimachia* also explains the blue color of *Petunia hybrida* (Kroon et al., 1994), *Lobelia erinus* (Hsu et al., 2017) and *Parochetus communis* (Tatsuzawa, 2020) and is known to provide stress tolerance (e.g. UV response) when rhamnose is bound to flavonoids (Li et al., 2017; Jiang et al., 2021). However, in *Lm*, the opposite is true for several climate niche parameters, especially with regard to precipitation (Appendix 1—table 7 and figure 5C). It appears this shared color polymorphism that may have transcended speciation confers unique climate niches.

Further evidence of gene function, enzyme kinetics and genetic association must be conducted before any final conclusions can be made regarding the molecular causes and ecological consequences of this flower color variation. However, the data presented

herein lean toward a single biochemical and molecular footprint shared by both species.

If the shift from blue to orange happened twice (i.e. convergence), it would be

remarkable since it would most likely be in the same trans-acting regulator of *TTG1*.

Additional experimental approaches (genetic dissection, transformation, knock-out,

DFR specificity enzyme assays, etc.) promise to identify the type, number and order of

mutations responsible for these flower color changes, yet given the phenotypic,

biochemical and transcriptome similarities described here, we have yet to find any

strong evidence rejecting the ancestral polymorphism hypothesis.

Methods

Population sampling

We collected stem cuttings from blue-flowered populations from southern Spain (seven

of *Lm* and one of *La*) and from orange-flowered populations from northern Spain,

Morocco and Italy (five of *Lm* and one of *La*) (Appendix 1—table 9). Cuttings were

rooted in an aeroponic cloner under glasshouse conditions. We also sowed seeds from

blue-flowered plants from Portugal in glasshouses at the University of Seville. Voucher

specimens were deposited in the University of Seville Herbarium (SEV, Appendix 1—

table 9).

We sampled all five petals per flower from a mean of 19 flowers per plant (10 - 24

flowers) from ten blue- and ten orange-flowered plants of *Lm*, and a mean of 30 flowers

per plant from eight flowered plants of each color in *La* (see Sánchez-Cabrera et al.

(2021) for more details on *La* sampling). In all cases the bullseye at the petal base was

avoided. All samples were taken from first-day anthesis flowers and immediately, petals

were flash frozen and stored at -80ºC. For flavonoid profiling, we collected 10 samples

per flower color for *Lm* (flowers from different plants of the same population per

sample), including those present in the RNA-Seq analysis (Appendix 1—table 9). *La*

samples were obtained from 14 blue- and 11 orange-flowered individuals (see Sánchez-

Cabrera et al. 2021). For both species, each sample consisted of two to five flowers

without the bullseye.

UV-Vis petal spectra

Reflectance spectra of the adaxial surface of 44 blue and 67 orange petals of *La* and 75

blue and 50 orange petals of *Lm* were captured using a JAZ A1465 double-beam

spectrophotometer (Ocean Optics, Florida, USA). The deuterium-tungsten light source

provided reflectance between 300 and 700 nm.

We tested the UV-vis spectra for the relative amounts of variation in two tests (1)

between color morphs within a species versus (2) between species within a color morph

using permutational multivariate analysis (PERMANOVA test) (Maia & White, 2018).

For each test, we computed the Euclidean distances based on the difference in

reflectance at each wavelength between all pairs of individual spectra and conducted

permutational MANOVA tests (Anderson, 2001) to assess statistical differences (9999

permutations) making quantitative comparative statements between the R^2 values of the

two tests (adonis2 in vegan in R). We visualized the distances between the spectra using

metaMDS in vegan in R with 500 tries.

Identification of flavonoid compounds

Flavonoids were extracted by placing petal samples (see Population sampling section;

Appendix 1—table 9) in 1.5 mL microfuge tubes containing 500µl of MeOH with 1%

HCl and stored at -80°C. Homogenization was performed with a BeadBeater using 3 x

3.2 mm steel balls for 30 seconds, then centrifuged for 10 min to pellet cellular debris.

Samples of the supernatant were loaded on a Dionex UltiMate3000 ultra-high pressure

liquid chromatography system equipped with a diode array detector and connected to a

Thermo Fleet LCQ mass spectrometer (Thermo-Fisher, Waltham, MA USA). Data were

analyzed using Xcalibur software (Thermo Fisher Scientific). Flavonoid identification

was based on the compound's retention time, UV-Vis spectra and whenever possible,

chromatographic comparisons with authentic standards: quercetin, luteolin, kaempferol,

isorhamnetin, malvidin, pelargonidin, cyanidin and delphinidin (Extrasynthese, Genay,

France), previously reported for other *Lysimachia* species (see Sánchez-Cabrera et al.,

2021). Relative contents (%) of the primary anthocyanins of each color morph sample,

based on the total area under the curve, were calculated and used in statistical

comparisons between the two species applying a Mann-Whitney U test in R.

RNA extraction and reads filtering

Collected petals (see Population sampling section) were homogenized using a mortar
and pestle. Total RNA was then extracted following the Qiagen RNeasy Plant Mini Kit
protocol (Qiagen, Germany) with the addition of PEG 20,000 mol. wt. (550µl, 2%;
Gehrig et al. 2000) before the first filtering step. The addition of PEG was essential to
achieve reasonable RNA concentrations for library preparation and sequencing. RNA
samples were stored at -80ºC until further analysis. RNA concentration and purity was
initially assessed with a Nanodrop Nd-1000 (ThermoFisher) and agarose gel, and then
confirmed with a Bioanalyzer (Agilent, Santa Clara, CA, USA) before sequencing.
Individual libraries were barcoded, multiplexed and sequenced as 150 bp paired-end
reads using two lanes on an Illumina Hi-Seq 2000 (Illumina, San Diego, CA, USA)
through Novogene (Beijing, China). Raw paired-end Illumina reads were assessed for
quality using FastQC (Andrews, 2010) and were processed using Rcorrector v1.0.4
(Song and Florea, 2015) to correct random sequencing errors. Then, reads were trimmed
with Trimmomatic v0.39 (Bolger et al., 2014) to remove any read containing bases with
Phred scores lower than 20, low quality reads less than 50 bp long, and any adapter or
other Illumina-specific sequences that were still present (Appendix 1—table 10).

Identification of differentially expressed genes (DEGs)

To determine DEGs for *Lm*, we used the *La* transcriptome assembly with known ABP
genes identified (Sánchez-Cabrera et al., 2021) as reference for mapping *Lm* RNA-Seq
samples with Bowtie2 software (Langmead & Salzberg, 2012). We calculated Trimmed
mean of M-values (TMM, mean of log-expression ratios; Robinson & Oshlack, 2010)
for each gene using RSEM software (Li & Dewey, 2011). Then, with EdgeR package
(Robinson et al., 2009) in R v4.0.0 (R Core Team, 2020) we determined statistically
significant differentially expressed genes (DEGs) between blue and orange petal
samples, applying the conservative thresholds for DEG identification: the false
discovery rate (FDR) less than 10^{-5} and the expression difference threshold was greater
than one $\log_2$ fold-change ($\log_2$FC), following Sánchez-Cabrera et al. (2021). To keep
biologically interesting genes for differential expression analysis, we considered those
genes with more than one count per million (CPM) in a minimum of four samples.
Some regulatory genes have very low expressions and magnitudes were unreliable. For
those loci, we performed the non-parametric Mann-Whitney U test to test for
differential expression.

Genes and isotigs selection for analysis and correlated expression analysis

We selected 22 ABP color differentiating structural and regulatory genes with

differential expression between colors in either both species, just in one species or not

differentially expressed but was a copy of a DEG (see Results section). We established

a criterion to select one isotig per gene: (1) samples with less than 20% ambiguous

nucleotides in the CDS, (2) samples with more than 100 reads mapped to the sequence,

(3) samples with the longest CDS, (4) samples with higher expression. Finally, we

selected the isotig which was present in a higher number of samples after applying the

criteria. We tested for correlated expression among the 22 structural and regulatory

genes in both species using Kendall correlations with Bonferroni correction for multiple

testing in R.

Phylogenetic analyses and non-synonymous SNP identification

We conducted phylogenetic analyses on the selected isotigs for all ABP structural and

regulatory genes. We followed the methods described in Sánchez-Cabrera et al. (2021)

to map to reference the reads using Geneious v9.0.4 (Kearse et al., 2012). For the

outgroups, we used top BLASTn hits (filtering for *Cyclamen* sp., *Camellia* sp.,

Vaccinium sp., *Actinidia* sp. and *Rhododendron* sp. when available) and, when possible,

top tBLASTx from *Camellia sinensis*, *Embelia ribes* and *Primula veris* genomes. We

reconstructed the relationships using the RAxML 7.2.8 plug-in from within Geneious

v9.0.4, searching for the maximum likelihood tree with 1000 bootstrap replicates. We

added normalized expression values (TMM) to the phylogenetic trees with "ggtree" (Yu

et al., 2017) and "phytools" (Revell, 2012) packages in R v4.0.0 (R Core Team, 2020).

We concatenated alignments of all ABP structural and regulatory loci and performed a

combined phylogenetic analysis using the same methodology as above but partitioning

the data by locus and determining the best fit model of nucleotide substitution.

We calculated the NS SNP rate between blue and orange petals of each species, with a

cutoff of 75% difference between colors. We used the isotigs selected for NS SNP

search. We focused on genes with orange monophyly to either one or both species,

because orange is derived from blue and therefore, we expect genes responsible for the

blue to orange shift to show a single common ancestor for the orange samples.

Differentiating the four hypotheses

To differentiate the four hypotheses (Figure 2) we used the following logic. First, the

convergence hypothesis predicts separate origins of the blue to orange color shift which

would produce separate clades of orange samples for both species. Under this

hypothesis, and with enough data, *La* and *Lm* should be reciprocally-monophyletic. The

introgression hypothesis involves a single origin of the color change in both species

(after the speciation event) which would appear as very close relationships for the color

differentiating locus between orange *Lm* and *La* since they are the same gene, just

shared across species boundaries (for the color causing gene(s) the species should not be

monophyletic), yet for non-color causing loci, species may be monophyletic. However,

the species must be interfertile. Third, the ancestral polymorphism hypothesis produces

a single origin of the color change (before the speciation event) creating separate

monophyletic blue and orange clades. Finally, if taxonomic confusion has blurred the

identity of the species (non-monophyletic species hypothesis, Jiménez-López et al.,

2022), then there could be a monophyletic color clade (across species boundaries) – a

topology expected for all loci, not just those contributing to the color shift.

Reproductive isolation between La and Lm

581 To determine reproductive isolation between *La* and *Lm*, we performed controlled

582 crosses in plants from different populations grown in glasshouse. Flowers were

583 emasculated before anthesis, and hand-pollinations were made during the first day of

584 anthesis within and between species and color morphs. Pollinated flowers were left to

585 set fruits and the fruit production and the number of seeds per fruit were counted.

Climate niche modeling

To test whether the color morphs of each species differed ecologically across their

ranges, we constructed an occurrence dataset using a combination of personal

observations and iNaturalist (inaturalist.org) records (manually verified for species

identification and color; Appendix 1—figure 6). The dataset comprised 4287 records.

All 19 BIOCLIM variables were then extracted for each record based on GPS

coordinates at the highest available spatial resolution (30 second) (Fick and Hijmans,

2017). Occurrences with no climate data and duplicated occurrences (multiple

occurrences of the same species with the same flower color within a pixel) were deleted.

There were 3309 occurrences in the final dataset (*La* blue = 641, *La* orange = 1941, *Lm*

blue = 662, *Lm* orange = 65).

To test for univariate climatic differences among the four morphs, a separate Kruskall-

Wallis test with a post-hoc Dunn test (non-parametric analogs of ANOVA with post-hoc

Tukey's test) was applied to each variable. Significance tests used Bonferroni

adjustment for 76 comparisons (19 climate variables across four species/flower color

combinations).

To determine whether similar climate variables were most important for separating

color morphs in the two species, logistic regression models were built for each species

on a subset on non-correlated climate variables. First, strongly correlated climate

variables were removed and a dataset of five variables that were both relatively

uncorrelated and putatively biologically meaningful based on field observations and

experimental manipulations (Arista et al. 2013; J. Jiménez-Lopez et al., unpublished

data) were retained: precipitation of wettest month, precipitation seasonality,

isothermality, temperature seasonality and mean temperature of wettest quarter. Data

were normalized using the 'normalize' procedure in the 'decostand' function from the

vegan R package (Oksanen et al. 2019) to facilitate later comparison of regression

coefficients. Best fit models were selected using AIC scores derived from the 'dredge'

function from the MuMIn R package (Bartón, 2020).

To test whether the two orange morphs were more divergent than the two blue morphs

across the 19 BIOCLIM variables, a randomization procedure was used. The specific

null hypothesis was no difference between the distance between the two orange morph

means and the distance between the two blue morph means. To build a null model for

each of the 19 variables, species identities were retained but color morphs were

randomized 999 times. For each distribution, one tail indicated that differences between

orange populations was greater while the other tail indicated greater differences

between blue populations. A two-tailed test indicated whether observed orange

differences were greater than expected, whether blue differences were greater than

expected or whether neither color was more divergent.

Data Availability

Acknowledgements

We would like to thank the support staff in the Department of Biology and the

Department of Chemistry and Biochemistry at Santa Clara University. Boris Igić

provided thoughtful insights on the evolutionary history of color in these species. We

would like to thank M. Abdelaziz for his help in organizing the field collecting trip and

during the sampling of Moroccan *L. monelli* plants. Grants PID2020-116222GB-100 to

M.A. and E.N., CGL2015-63827 to M.A. and P.L.O. and BES-C-2016-0023 to M.S.C

funded by MCIN/AEI/ 10.13039/501100011033 and by "European Union

NextGenerationEU/PRTR".

Conflict of interest

None

Author contributions

MS-C, EN, MA, PO, and JW designed and performed research; AF conducted the

UHPLC technique; BC conducted the climate niche analyses; FJ-L performed plant

crosses and supplied *L. arvensis* reflectance data. MS-C, EN, and JW analyzed data; and

MS-C and JW wrote and revised the paper with assistance from coauthors.

References

Allison AC. 1961. Genetic factors in resistance to malaria. *Ann. N. Y. Acad. Sci.* **91**:
710–729.

Anderson MJ. 2001. A new method for non-parametric multivariate analysis of
variance. *Austral Ecol.* **26**: 32–46.

Arista M, Talavera M, Berjano R, Ortiz PL. 2013. Abiotic factors may explain the
geographical distribution of flower colour morphs and the maintenance of colour
polymorphism in the scarlet pimpernel. *J. Ecol.* **101**: 1613–1622.

Bartón K. 2020. *MuMIn: Multi-Model Inference.* R package version 1.43.17.

Berardi AE, Esfeld K, Jäggi L, Mandel T, Cannarozzi GM, Kuhlemeier C. 2021.
Complex evolution of novel red floral color in *Petunia*. *Plant Cell* 33: 2273–2295.

Caputi L, Malnoy M, Goremykin V, Nikiforova S, Martens S. 2012. A genome-wide
phylogenetic reconstruction of family 1 UDP-glycosyltransferases revealed the
expansion of the family during the adaptation of plants to life on land. *Plant J.* **69**:
1030–1042.

Chen C, Zhang K, Khurshid M, Li J, He M, et al. 2019. MYB transcription
repressors regulate plant secondary metabolism. *Crit. Rev. Plant Sci.* **38**: 159–170.

Comeault AA, Flaxman SM, Riesch R, et al. 2015. Selection on a genetic
polymorphism counteracts ecological speciation in a stick insect. *Curr. Biol.* **25**: 1975–
1981.

Conway MS. 2005. *Life's solution: inevitable humans in a lonely universe.* Cambridge,
UK: Cambridge University Press.

Corl A, Davies AR, Kuchta SR, Sinervo B. 2010. Selective loss of polymorphic
mating types is associated with rapid phenotypic evolution during morphic speciation.
Proc. Natl. Acad. Sci. U.S.A. **107**: 4254–4259.

Des Marais DL, Rausher MD. 2010. Parallel evolution at multiple levels in the origin
of hummingbird pollinated flowers in *Ipomoea*. *Evolution* **64**: 2044–2054.

Dobzhansky T. 1937. *Genetics and the origin of species.* New York, NY: Columbia
University Press.

Fick SE, Hijmans RJ. 2017. WorldClim 2: new 1km spatial resolution climate surfaces
for global land areas. *Inter. J. Climat.* **37**: 4302–4315.

Gibbs PE, Talavera S. 2001. Breeding system studies with three species of *Anagallis*
(Primulaceae): self-incompatibility and reduced female fertility in *A. monelli* L. *Ann.*
Bot. **88**: 139–144.

Gould SJ. 1989. *Wonderful life: the Burgess shale and the nature of history.* New York,
NY: W.W. Norton.

Grant V. 1981. *Plant speciation*, 2nd ed. New York, NY: Columbia University Press.

Harborne JB. 1968. Comparative biochemistry of the flavonoids. VII. Correlations
between flavonoid pigmentation and systematics in the family Primulaceae.
Phytochemistry **7**: 1215–1230.

He XZ, Wang X, Dixon RA. 2006. Mutational analysis of the *Medicago*
glycosyltransferase UGT71G1 reveals residues that control regioselectivity for
(iso)flavonoid glycosylation. *J. Biol. Chem.* **281**: 34441–34447.

Hsu YH, Tagami T, Matsunaga K, Okuyama M, Suzuki T, et al. 2017. Functional
characterization of UDP-rhamnose-dependent rhamnosyltransferase involved in
anthocyanin modification, a key enzyme determining blue coloration in *Lobelia erinus*.
Plant J. **89**: 325–337.

Hugall AF, Stuart-Fox D. 2012. Accelerated speciation in colour-polymorphic birds.
Nature **485**: 631–634

Igić B, Kohn JR. 2001. Evolutionary relationships among self-incompatibility RNases.
Proc. Natl. Acad. Sci. U.S.A. **98**: 13167–13171.

Igić B, Lande R, Kohn JR. 2008. Loss of self-incompatibility and its evolutionary
consequences. *Int. J. Plant Sci.* **169**: 93–104.

Jamie GA, Meier JI. 2020. The persistence of polymorphisms across species
radiations. *Trends Ecol. Evol.* **35**: 795–808.

Jiang N, Dillon FM, Silva A, Gomez-Cano L, Grotewold E. 2021. Rhamnose in
plants-from biosynthesis to diverse functions. *Plant Sci.* **302**: 110687.

Jiménez-López FJ, Matas L, Arista M, Ortiz PL. 2019. Flower colour segregation
and flower discrimination under the bee vision model in the polymorphic *Lysimachia*
arvensis. *Plant Biosyst.* **154**: 535–543.

Jiménez-López FJ, Viruel J, Arista M, Ortiz PL, Talavera M. 2022. Molecular
approaches reveal speciation between red- and blue-flowered plants in the
Mediterranean *Lysimachia arvensis* and *L. monelli* (Primulaceae). *Bot. J. Linn. Soc.*
199: 557–577.

Jiménez-López FJ, Arista M, Talavera M, Morellato LP, Pannell JR, Viruel J,
Ortiz PL. 2023. Multiple pre- and post-zygotic componets of reproductive isolation
between two co-occurring *Lysimachia* species. *New Phytol.* **238**: 874–887.

Johnson ET, Ryu S, Yi HK, Shin B, Cheong H, Choi G. 2001. Alteration of a single
amino acid changes the substrate specificity of dihydroflavonol 4-reductase. *Plant J.* **25**:
325–333.

Katsu K, Suzuki R, Tsuchiya W, Inagaki N, Yamazaki T, et al. 2017. A new
buckwheat dihydroflavonol 4-reductase (DFR), with a unique substrate binding
structure, has altered substrate specificity. *BMC Plant Biol.* **17**: 1–14.

Kroon J, Souer E, De Graaff A, Xue Y, Mol J, Koes R. 1994. Cloning and structural
analysis of the anthocyanin pigmentation locus Rt of *Petunia hybrida*: characterization
of insertion sequences in two mutant alleles. *Plant J.* **5**: 69–80.

Larter M, Dunbar-Wallis A, Berardi AE, Smith SD. 2018. Convergent evolution at
the pathway level: predictable regulatory changes during flower color transitions. *Mol.*
Biol. Evol. **35**: 2159–2169.

Li P, Li YJ, Zhang FJ, Zhang GZ, Jiang XY, et al. 2017. The *Arabidopsis* UDP-
glycosyltransferases UGT79B2 and UGT79B3, contribute to cold, salt and drought
stress tolerance via modulating anthocyanin accumulation. *Plant J.* **89**: 85–103.

Madsen T, Stille B, Ujvari B, Bauwens D, Endler JA, 2022. Negative frequency-
dependent selection on polymorphic color morphs in adders. *Curr. Biol*. **32**: 3385–3388.

Maia R, White TE. 2018. Comparing colors using visual models. *Behav. Ecol.* **29**:
649–659

Mallet J. 2005. Hybridization as an invasion of the genome. *Trends Ecol. Evol.* **20**:
229–237.

Martin SH, Dasmahapatra KK, Nadeau NJ, Salazar C, Walters JR, et al. 2013.
Genome-wide evidence for speciation with gene flow in *Heliconius* butterflies. *Genome*
Res. **23**: 1817–1828.

McKinnon JS, Pierotti ME. 2010. Colour polymorphism and correlated characters:
genetic mechanisms and evolution. *Mol. Ecol.* **19**: 5101–5125.

McLean CA, Stuart-Fox D. 2014. Geographic variation in animal colour
polymorphisms and its role in speciation. *Biol. Rev*. **89**: 860–873.

McRobie HR, Moncrief ND, Mundy, NI. 2019. Multiple origins of melanism in two
species of North American tree squirrel (*Sciurus*). *BMC Evol. Biol.* **19**: 1–14.

Monein A, Atta II, Shehata AA. 2003. On the delimitation of *Anagallis arvensis* L.
(Primulaceae) 1. Evidence based on micromorphological characters, palynological
features and karyological studies. *Pakistan J. Biol. Sci.* **6**: 29–35.

Muchhala N, Johnsen S, Smith, SD. 2014. Competition for hummingbird pollination
shapes flower color variation in Andean Solanaceae. *Evolution* **68**: 2275–2286.

Ng J, Smith SD. 2016. How to make a red flower: the combinatorial effect of pigments.
AoB Plants 8: plw013.

Nolte AW, Gompert Z, Buerkle CA. 2009. Variable patterns of introgression in two
sculpin hybrid zones suggest that genomic isolation differs among populations, *Mol.*
Ecol. **18**: 2615–2627.

Ogutcen E, Durand K, Wolowski M, Clavijo L, Graham C, Glauser G, Perret M.
2020. Chemical basis of floral color signals in Gesneriaceae: the effect of alternative
anthocyanin pathways. *Front. Plant Sci*. **11**: 604389.

Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, et al. 2019. *vegan:*
community ecology package. R package version 2.5-6. https://CRAN.R-
project.org/package=vegan

Ortiz PL, Berjano R, Talavera M, Rodríguez-Zayas L, Arista M. 2015. Flower
colour polymorphism in *Lysimachia arvensis*: how is the red morph maintained in
Mediterranean environments? *Perspect. Plant Ecol. Evol. Syst.* **17**: 142–150.

Passeri V, Koes R, Quattrocchio FM. 2016. New challenges for the design of high
value plant products: stabilization of anthocyanins in plant vacuoles. *Front. Plant Sci*. **7**:
153.

Petit P, Granier T, d' Estaintot BL, Manigand C, Bathany K, Schmitter JM, et al.
2007. Crystal structure of grape dihydroflavonol 4-reductase, a key enzyme in flavonoid
biosynthesis. *J. Mol. Biol.* **368**: 1345–1357.

Pérez-Díaz JR, Pérez-Díaz J, Madrid-Espinoza J, González-Villanueva E, Moreno
Y, Ruiz-Lara S. 2016. New member of the R2R3-MYB transcription factors family in
grapevine suppresses the anthocyanin accumulation in the flowers of transgenic
tobacco. *Plant Mol. Biol.* **90**: 63–76.

Pujadas A. 1997. *Anagallis* L. In: Castroviejo S, Aedo C, Laínz M, Morales R, Muñoz
Garmendia F, et al., eds. *Flora Iberica, V. Ebenaceae-Saxifragaceae*. Madrid, Spain:
Real Jardín Botánico, CSIC, 57–62.

Revell LJ. 2012. phytools: An R package for phylogenetic comparative biology (and
other things). *Methods Ecol. Evol.* **3**: 217–223.

Sánchez-Cabrera M. 2023. *Flower color polymorphism and speciation: Lysimachia*
monelli and Lysimachia arvensis as study systems. Ph.D. thesis, University of Seville,
Seville, Spain.

Sánchez-Cabrera M, Jiménez-López FJ, Narbona E, Arista M, Ortiz PL, Romero-
Campero FJ, et al. 2021. Changes at a critical branchpoint in the anthocyanin
biosynthetic pathway underlie the blue to orange flower color transition in *Lysimachia*
arvensis. *Front. Plant Sci.* **12**: 247.

Schemske DW, Bierzychudek P. 2007. Spatial differentiation for flower color in the
desert annual *Linanthus parryae*: was Wright right? *Evolution* **61**: 2528–2543.

Short AW, Streisfeld MA. 2023. Ancient hybridization leads to the repeated evolution
of red flowers across a monkeyflower radiation. Evol. Lett. **7**: 293–304.

Sirugo G, Predazzi IM, Bartlett J, Tacconelli A, Walther M, Williams SM. 2014.
G6PD A-deficiency and severe malaria in The Gambia: heterozygote advantage and
possible homozygote disadvantage. *Am. J. Trop. Med. Hyg.* **90**: 856.

Stankowski S, Streisfeld MA. 2015. Introgressive hybridization facilitates adaptive
divergence in a recent radiation of monkeyflowers. *Proceedings of the Royal Society B:*
Biological Sciences. **282**: 20151666.

Streisfeld MA, Rausher MD. 2009. Genetic changes contributing to the parallel
evolution of red floral pigmentation among *Ipomoea* species. *New Phytol.* **183**: 751–
763.

Sun W, Liang L, Meng X, Li Y, Gao F, Liu X, et al. 2016. Biochemical and
molecular characterization of a flavonoid 3-O-glycosyltransferase responsible for
anthocyanins and flavonols biosynthesis in *Freesia hybrida*. *Front. Plant Sci.* **7**: 410.

Svensson EI. 2017. Back to basics: using colour polymorphisms to study evolutionary
processes. *Mol. Evol.* **26**: 2204–2211.

Tatsuzawa F. 2020. Flavonoids in the blue flowers of *Parochetus communis* Buch.-
Ham. ex D. Don (Leguminosae). *Biochem. Syst. Ecol.* **92**: 104–108.

Thomson JD, Wilson P. 2008. Explaining evolutionary shifts between bee and
hummingbird pollination: convergence, divergence, and directionality. *Int. J. Plant Sci.*
169: 23–38.

Tian H, Wang S. 2020. TRANSPARENT TESTA GLABRA1, a key regulator in plants
with multiple roles and multiple function mechanisms. *Int. J. Mol. Sci.* **21**: 4881.

Wang X. 2009. Structure, mechanism and engineering of plant natural product
glycosyltransferases. *FEBS Lett.* **583**: 3303–3309.

Wessinger CA, Rausher MD, Hileman LC. 2019 Adaptation to hummingbird
pollination is associated with reduced diversification in *Penstemon*. *Evol. Lett.* **3**: 521–
533.

Wessinger CA, Rausher MD. 2013. Predictability and irreversibility of genetic
changes associated with flower color evolution in *Penstemon barbatus*. *Evolution* **68**:
1058–1070.

Wright S. 1943. An analysis of local variability of flower color in *Linanthus parryae.*
Genetics **28**: 139–156.

Yonekura-Sakakibara K, Nakayama T, Yamazaki M, Saito K. 2008. Modification
and stabilization of anthocyanins. In: Winefield C, Davies K, Gould K, eds.
Anthocyanins. New York, NY: Springer.

Yoshida K, Iwasaka R, Shimada N, Ayabe S, Aoki T, Sakuta M. 2010.
Transcriptional control of the dihydroflavonol 4-reductase multigene family in *Lotus*
japonicus. *J. Plant Res.* **123**: 801–805.

Yu G, Smith D, Zhu H, Guan Y, Lam TT. 2017. ggtree: an R package for
visualization and annotation of phylogenetic trees with their covariates and other
associated data. *Methods Ecol. Evol.* **8**: 28–36.

Zufall RA, Rausher MD. 2004. Genetic changes associated with floral adaptation
restrict future evolutionary potential. *Nature* **428**: 847–850.

Figure 1. Flower color characterization of blue and orange *Lysimachia arvensis* and *L. monelli*. (A) Scaled images and geographic distributions of the two flower color morphs of both species. Coexistence of the *L. arvensis* color morphs is indicated with diagonal striping. Populations sampled for transcriptome comparisons are indicated with black circles (blue) and black stars (orange) (numbers indicate >1 geographically adjacent populations were sampled for *L. monelli*). (B) UV-vis reflectance spectra of blue and orange petals of both species showing mean curves with 95% confidence intervals in colored shading above and below the mean curve. (C) Molecular structures and relative content from the biochemical analysis of the primary anthocyanins in both species. Non-significant differences (ns) in relative content are indicated using a Mann-Whitney U test. In *L. arvensis*, orange samples include pelargonidin 3-glucoside derivatives. Error bars represent standard error (SEM).

845
846

Figure 2. Four evolutionary hypotheses of petal color evolution in *L. arvensis* and *L.*

monelli. Assuming the ancestral state is blue and diploid, the most parsimonious color

shifts are indicated by a triangle and ploidy changes are indicated with a circle (placed

at the node since they likely drive divergence). In (A), convergent evolution of orange

petals is coupled with a single origin of the tetraploid lineage predicts independent and

distinct molecular causes for the color change in each lineage. In (B), a single origin of

orange petals in *L. monelli* followed by introgression to *L. arvensis* requires only one

shift to orange and one polyploid event but requires the lineages to be able to hybridize.

In (C), a single origin of orange in the common ancestor creates an ancestral

polymorphism that transcends the speciation event and persists to the present in both

species, requiring one polyploidy event. In (D), non-monophyletic species allow for a

single origin of orange petals but requires independent origins of the tetraploid lineages.

860
Figure 3. Gene expression comparisons of blue and orange morphs of two species of

Lysimachia. (A) Suggested Anthocyanin Biosynthetic Pathway (ABP) producing blue

(malvidin 3-rhamnoside) and orange (pelargonidin 3-glucoside) flower colors in these

two species. Mean expression (TMM +/- standard error) in both species (La and Lm)

are shown for each ABP gene (green font). Significantly differentially expressed genes

(DEGs) are indicated with a black box around the bar plots. (B) Diagrams of the *DFR*

coding sequence showing color-differentiating non-synonymous SNPs in pink (>70%

frequency difference) between blue and orange *DFR-1* of *L. arvensis* and *L. monelli*,

and differences between *DFR-1* (blue and orange) and *DFR-2* (orange) of both species.

Gray areas indicate substrate specificity and active sites of *DFR* genes. The magnified

region comparing *DFR-1* and *DFR-2* shows many NS SNPs (pink) in the largest

substrate specificity region. (C) Volcano plot comparing the entire transcriptomes of *L.*

arvensis and *L. monelli* highlighting ABP genes with significant differential expression.

875
876

Figure 4. Maximum likelihood cladogram of the coding sequences of three *DFR*
paralogues expressed in two *Lysimachia* species. Outgroups were selected from top
BLASTn hits and, when available, from the genomes of closely related species
(*Camelia sinensis*, *Embelia ribes* and *Primula veris*). Outgroups are indicated by
scientific name followed by Genbank Accession numbers. Bootstrap values are
provided to the right of the nodes when greater than 70%. The bar plot shows the
expression level (TMM) for blue and orange flowers of *L. arvensis* (solid) and *L.*

884 *monelli* (black stripes). The monophyletic clade of orange petal samples for *DFR-2* is

885 indicated with an asterisk (*). A phylogram (inset) is provided with the samples in the

886 same order as the cladogram to compare relative branch lengths. Scale bar is in

887 substitutions per site.

Table 1. Evidence for a single origin versus convergent evolution of the blue to orange flower color shift in *Lm* and *La*.

Evidence	Evidence for trans-specific polymorphism	Evidence for convergent evolution
UV-vis spectra	7x more variation between color morphs than between species (Figure 1B; Appendix 1—figure 1)	In the UV, two peaks for *Lm*-O vs. one peak + a shoulder for *La*-O and in the visible wavelengths there are some small differences in the inflection points (Figure 1B)
Biochemistry	Identical anthocyanins detected in both species and no significant differences in the relative concentrations for each color morph between species (Figure 1C)	None
Expression – structural genes	Significant upregulation of *DFR-2* and *BZ1-2* in orange morphs of both species (Figure 3A)	Compensatory downregulation of the *F3'5'H*, *DFR-1* and *BZ1-1* in orange morphs is only significant in *La* (but *F3'5'H* and *DFR-1* trend toward downregulation in orange *Lm*) (Figure 3A); *CHS, OMT, 3GT* and *UGT72E* are DEGs in *La*, but not *Lm* (Figure 3A)
Expression – regulatory genes	*MYB4-2* (negative regulator of ABP structural genes) has higher expression in orange morphs of both species (but is only significant in *La*; Figure 3A)	No single known ABP regulatory locus with significant expression differences between blue and orange morphs for both species.

Expression correlations	*DFR-2* and *BZ1-2* are strongly correlated with each other in both species and *BZ1-2* is significantly correlated with the positive regulator *TTG1* in both species; *DFR-1* is negatively correlated with *MYB4-2* in orange morphs of both species (Appendix 1—figure 4)	Correlation between *DFR-2* and *TTG1* is only significantly in *La* after Bonferroni correction (but they have a strong positive association in *Lm*)
SNPs – structural genes	*DFR-2* of orange morphs of both species have the same 13 NS SNPs which overlap with a known pelargonidin-specific domain (Figure 3B)	*bHLH12* exhibits reciprocal monophyly of species (Appendix 1—figure 2J), however no NS SNPs differentiate color morphs in either species
SNPs – regulatory genes	*MYB4-2* has two NS SNPs in both species	*MYB4-2*'s NS SNPs do not overlap with known functional domains; *F3'H* has five NS SNPs in *Lm*-orange and only one in *La*-orange
Climate niche modeling	Three temperature-related variables have parallel effects on color morphs for both species (orange = cooler than blue; Appendix 1—table 4A)	Six variables (mostly precipitation-related) have opposing effects on color morphs in the two species (orange *Lm* = dry vs. orange *La* = wet; Appendix 1—table 4A).

Appendix 1

Appendix 1—figure 1. NMDS plot of UV-Vis reflectance spectra for blue and orange

petalled individuals of *L. arvensis* (LYAR) and *L. monelli*. Statistical analysis was

assessed with permutational manova (Anderson, 2001) with the following results (1)

restricting permutations within color (comparing across species), $R^2 = 0.15113$, F =

41.662, p < 1e-04 vs. (2) restricting permutations within species (comparing across

colors), $R^2 = 0.74999$, F = 701.98, p < 1e-04. Very similar results were found after

applying a brightness correction – subtracting the minimum reflectance value of each

sample from all wavelengths (not shown).

BZ1-2 and -3

Caffeoyl CoA-1,-2 and *-3.*

 CHS

F3'5'H

F3'H

AN11a

bHLH1

bHLH2-2

bHLH2-3

bHLH12

COP1

MYB4-1

929   *MYB4-2*

931   *MYB16-1*

933   *MYB16-2*

935   *SPA1*

937   *TTG1*

Appendix 1—figure 2. Maximum likelihood phylogenetic analysis of the coding

sequence for eight structural (A-E) and 12 regulatory (F-Q) loci from blue and orange

petals of *L. arvensis* and *L. monelli*. (A) *BZ1-2* and *-3*. (B) *Caffeoyl CoA-1,-2* and *-3*.

(C) *CHS*. (D) *F3'5'H*. (E) *F3'H*. (F) *AN11a*. (G) *bHLH1*. (H) *bHLH2-2*. (I) *bHLH2-3*.

(J) *bHLH12*. (K) *COP1*. (L) *MYB4-1*. (M) *MYB4-2*. (N) *MYB16-1*. (O) *MYB16-2*. (P)

SPA1. (Q) *TTG1*. Outgroups were included when available from the genomes of closely

945   related species (*Camelia sinensis, Embelia ribes* and *Primula veris*). Bootstrap values

946   above 70% are provided to the right of the nodes. Inset phylograms are provided for

947   branch length comparisons with samples in same order as the larger cladogram and

948   contain a scale bar in substitutions per site. The bar plots show the gene expression level

949   (TMM values) of blue and orange flowers.

Appendix 1—figure 3. Combined maximum likelihood phylogenetic analysis of 24
ABP structural and regulatory loci. Phylogram on the left with branchlengths and
cladogram on the right with bootstrap values indicated at the nodes. Cladogram
annotated with petal color (blue and orange brackets). Lysimachia arvensis (LYAR and
La samples) and L. monelli (LYMO and Lm samples) are strongly supported as
monophyletic as are the color types within each species except for the paraphyly of
LYMO orange.

Appendix 1—figure 4. Kendall correlations of the expression level (TMM values) between the most relevant differential expressed structural genes (*DFR-1*, *DFR-2*, *F3'5'H*, *BZ1-2*, *CaffCoA-1*, *CHS*) and the regulatory genes (*MYB4-1*, *MYB4-2*, *MYB16-1*, *MYB16-2*, *bHLH1*, *bHLH12*, *bHLH2-1*, *bHLH2-2*, *bHLH2-3*, *TTG1*, *TTG2*, *SPA1*,

964   *AN11a*, *COP1*) present in blue and orange *L. arvensis* (A) and *L. monelli* (B).

965   Correlation coefficients (R) and P values are listed below the diagonal and the raw data

966   above the diagonal. Black boxes indicate those correlations that are significant

967   following a Bonferroni correction.

A. Parallel, both significant.

B. Parallel, non-significant (one species without significant within-species differences)

C. Intersecting, significant

D. Intersecting, non-significant (one species without significant within-species differences)

Appendix 1—figure 5. Climate niche modeling results for blue and orange morphs of

L. arvensis and *L. monelli.* Grouped the 19 variables into 4 categories. (A) Parallel

significant: slopes are similar (both + or both -), with orange/blue *L. monelli*

significantly different AND orange/blue *L. arvensis* significantly different. (B) Parallel

non-significant: slopes are similar, one or both within-species comparison is non-

significant. (C) Intersecting significant: slopes are different (one +, one -), within-

species differences are significant. (D) Intersecting non-significant: Slopes are different,

one within-species comparisons is non-significant.

Appendix 1—figure 6. Occurrence data used in climate niche model from individually

reviewed iNaturalist records and other sources (herbarium records, personal

observations). For *L. arvensis*, there are 1941 orange occurrences and 641 blue (A). For

L. monelli, there are 662 blues and 65 oranges.

Appendix 1—table 1. Putative flavonoid identifications of blue and orange petal extracts of *L. monelli* from the UHPLC–MS biochemical

analysis (MS analysis was acquired in positive mode). Flavonoids <1% (trace) are not shown.

Petal color	Flavonoid class	Putative flavonoid identification	N samples**	t_R (min)	λ_{max} (nm)	Mass (m/z)	Relative content (%)***
Blue	Anthocyanin	Malvidin 3-rhamnoside	10/10	5.7	228, 278, 526	476.87	94.4
(**n** = 10 plants)	Anthocyanin	Delphinidin	10/10	2.1	229, 300, 521	variable	1.1
	Flavonol	Kaempferol 3-glucoside / Quercetin 3-rhamnoside / Isorhamnetin 3-pentoside / Luteolin 7-glucoside	5/10	7.7	226, 258, 291, 350	448.26	3.7
Orange	Anthocyanin	Pelargonidin 3-diglucoside	7/9	2.4	230, 277, 329, 426, 501	594.98	4.1
(**n** = 9 plants)	Anthocyanin	Pelargonidin 3-glucoside	9/9	3.1	231, 279, 429, 499	433.03	95.6

Identification was based on retention time (t_R), maximum absorbance wavelengths (λ_{max}) observed in the UV-Vis spectra of the compound that

eluted at the t_R indicated, and molecular mass comparisons with standards and values of previously reported flavonoids for *Lysimachia* spp.

(Sánchez-Cabrera et al., 2021 and the database http://metabolomics.jp).

*The fraction of samples containing each flavonoid.

**Relative content of each putative compound in each flower color, calculated from the area under the UV-Vis peaks.

Appendix 1—**table 2**. Flavonoid biosynthetic pathway genes with significant differential expression between blue- and orange-flowered *L. monelli*.

Gene abbreviation	Gene name	Blue expression (TMM)	Orange expression (TMM)	Log fold-change	Pathway
BZ1-2	*Anthocyanin 3-O-glucosyltransferase*	0.31	2651.84	-13.05	ABP
DFR-2	*Dihydroflavonol 4 reductase*	0.01	157.18	-12.53	ABP
CHS	*Chalcone synthase*	331.90	784.13	-1.24	ABP
Beta-glucosidase	*Beta-glucosidase*	0.09	4.13	-5.37	Non-ABP
Caffeoyl CoA-1	*Caffeoyl CoA-O-methyltransferase*	12.73	0.32	5.29	Non-ABP

Appendix 1—table 3. Summary of individual phylogenetic analyses of ABP structural
and regulatory genes with high bootstrap support (>70%) indicating monophyletic
clades in specific groups (genus, species, color). Genes showing orange monophyly
(either for *L. arvensis*, *L. monelli* or both species) are highlighted in bold.

Genes	*Lysimachia*	La	La-O	La-B	Lm	Lm-O	Lm-B	O	B
Structural genes									
*BZ1-2***	100	98	-	-	-	-	-	100	-
BZ1-3	100	-	-	-	-	-	-	-	-
*Caff. CoA-1**	100	79	-	-	98	-	-	-	100
Caff. CoA-2	100	-	-	-	-	-	-	-	-
Caff. CoA-3	100	100	99	98	85	-	-	-	-
CHS	100	-	-	-	70	-	-	-	-
DFR-1	100	-	91	100	-	92	-	-	-
*DFR-2***	99	-	-	-	-	-	-	99	-
F3'5'H	100	-	98	86	93	-	-	-	-
F3'H	100	100	93	99	-	-	83	-	-
Regulatory genes									
AN11a	100	-	-	-	-	-	-	-	-
bHLH1	100	78	-	91	-	-	-	-	-
bHLH2-3	100	-	-	99	-	-	-	-	-
bHLH2-2	100	-	99	-	-	-	-	-	-
bHLH12	100	70	-	100	94	-	-	-	-
COP1	100	-	-	99	-	-	-	-	-
MYB4-1	99	-	-	-	-	-	-	-	-
MYB4-2	100	-	-	-	-	-	-	-	-

MYB16-1	100		-	87	-	-	-	-	-
MYB16-2	100	-	-	98	-	-	-	-	-	-
SPA1	100	-	-	-	-	-	-	-	-
TTG1	100	-	-	100	-	-	-	-	-

999   La (*L. arvensis*), Lm (*L. monelli*), O (orange), B (blue)

1000   * Genes only present in blue petals

1001   ** Genes only present in orange petals

Appendix 1—table 4. Fixed non-synonymous SNPs in ABP loci that differentiate color morphs and those that differentiate species.

Gene	Position (aa)	Blue AA	Orange AA	BLOSUM62
Fixed Between Color Morphs				
DFR-1	279/287[a]	E	Q	+2
MYB4-2	361	M	I	+1
MYB4-2	376	P	S	-1
Fixed Between Species				
BZ1-3	435	K/A[b]	T	-1/0
CaffCoA-3	36	V	A	0
CaffCoA-3	200	F	Y	+3
CaffCoA-1 (only in blue)[c]	2	T	A	0
CaffCoA-1 (only in blue)	46	E	A	-1
CaffCoA-1 (only in blue)	215	T	A	0
CHS	363	R	L	-2
F3'H-1	19	S	Y	-2
F3'H-1	262	I	M	+1
F3'H-1	324	Q	H	0
F3'H-1	326	K	E	+1
F3'H-1	338	A	V	0
F3'H-1	341	R	L/P	-2/-2
F3'5'H	2	E	D	+2
MYB4-1	71	S	T	-1
MYB16-1	18	T	I	+1

[a] Two positions separated by a slash indicates the alignment position in *L. arvensis*/*L. monelli* (because of indels).

[b] Two amino acids separated by a slash indicates variable in one species (BLOSUM62 values are reported with a slash in the same order).

[c] "Only in blue" is reported when the locus was not detected in orange samples (still fixed between species, but only for blue).

1011 **Appendix 1**—**table 5**. Results of hand-crosses between L. arvensis and L. monelli

1012 coming from different populations.

Mother species	Father species	Mother plant*	Father plant*	Mother color	Father color	Fruit production
L. monelli	*L. arvensis*	630/14_1	HIN_1.15	Blue	Blue	0
L. monelli	*L. arvensis*	630/14_2	HIN_1.15	Blue	Blue	0
L. monelli	*L. arvensis*	630/14_3	HIN_1.15	Blue	Blue	0
L. monelli	*L. arvensis*	630/14_4	HIN_2.21	Blue	Blue	0
L. monelli	*L. arvensis*	630/14_5	HIN_2.21	Blue	Blue	0
L. monelli	*L. arvensis*	630/14_1	COR_1.6	Blue	Orange	0
L. monelli	*L. arvensis*	630/14_2	COR_1.6	Blue	Orange	0
L. monelli	*L. arvensis*	630/14_3	COR_1.6	Blue	Orange	0
L. monelli	*L. arvensis*	630/14_4	COR_2.5	Blue	Orange	0
L. monelli	*L. arvensis*	630/14_5	COR_2.5	Blue	Orange	0
L. monelli	*L. arvensis*	151/14_1	HIN_1.15	Blue	Blue	0
L. monelli	*L. arvensis*	151/14_3	HIN_1.15	Blue	Blue	0
L. monelli	*L. arvensis*	151/14_4	HIN_2.21	Blue	Blue	0
L. monelli	*L. arvensis*	151/14_6	HIN_2.21	Blue	Blue	0
L. monelli	*L. arvensis*	151/14_7	HIN_8.7	Blue	Blue	0
L. monelli	*L. arvensis*	151/14_8	HIN_8.7	Blue	Blue	0
L. monelli	*L. arvensis*	151/14_1	COR_1.6	Blue	Orange	0
L. monelli	*L. arvensis*	151/14_3	COR_1.6	Blue	Orange	0
L. monelli	*L. arvensis*	151/14_4	COR_1.6	Blue	Orange	0
L. monelli	*L. arvensis*	151/14_6	COR_1.52	Blue	Orange	0
L. monelli	*L. arvensis*	151/14_7	COR_1.52	Blue	Orange	0
L. monelli	*L. arvensis*	151/14_8	COR_1.60	Blue	Orange	0
L. arvensis	*L. monelli*	HIN_1.15	630/14_1	Blue	Blue	0
L. arvensis	*L. monelli*	HIN_2.21	630/14_4	Blue	Blue	0
L. arvensis	*L. monelli*	HIN_2.21	630/14_5	Blue	Blue	0
L. arvensis	*L. monelli*	COR_1.6	630/14_1	Orange	Blue	0
L. arvensis	*L. monelli*	COR_2.5	630/14_5	Orange	Blue	0
L. arvensis	*L. monelli*	HIN_1.15	151/14_1	Blue	Blue	0
L. arvensis	*L. monelli*	HIN_1.15	151/14_3	Blue	Blue	0

L. arvensis	*L. monelli*	HIN_2.21	151/14_4	Blue	Blue	0
L. arvensis	*L. monelli*	HIN_8.7	151/14_8	Blue	Blue	0
L. arvensis	*L. monelli*	COR_1.6	151/14_1	Orange	Blue	0
L. arvensis	*L. monelli*	COR_1.6	151/14_3	Orange	Blue	0
L. arvensis	*L. monelli*	COR_1.52	151/14_6	Orange	Blue	0
L. arvensis	*L. monelli*	COR_1.60	151/14_8	Orange	Blue	0
L. monelli	*L. arvensis*	485/14_1	HIN_1.15	Orange	Blue	0
L. monelli	*L. arvensis*	485/14_1	HIN_1.15	Orange	Blue	0
L. monelli	*L. arvensis*	485/14_2	HIN_2.21	Orange	Blue	0
L. monelli	*L. arvensis*	485/14_2	HIN_2.21	Orange	Blue	0
L. monelli	*L. arvensis*	485/14_3	HIN_8.7	Orange	Blue	0
L. monelli	*L. arvensis*	485/14_3	HIN_8.7	Orange	Blue	0
L. monelli	*L. arvensis*	485/14_1	COR_1.6	Orange	Orange	0
L. monelli	*L. arvensis*	485/14_1	COR_1.6	Orange	Orange	0
L. monelli	*L. arvensis*	485/14_2	COR_1.52	Orange	Orange	0
L. monelli	*L. arvensis*	485/14_2	COR_1.52	Orange	Orange	0
L. monelli	*L. arvensis*	485/14_3	COR_1.60	Orange	Orange	0
L. monelli	*L. arvensis*	485/14_3	COR_1.60	Orange	Orange	0
L. arvensis	*L. monelli*	HIN_1.15	485/14_1	Blue	Orange	0
L. arvensis	*L. monelli*	HIN_2.21	485/14_2	Blue	Orange	0
L. arvensis	*L. monelli*	HIN_8.7	485/14_3	Blue	Orange	0
L. arvensis	*L. monelli*	COR_1.6	485/14_1	Orange	Orange	0
L. arvensis	*L. monelli*	COR_1.52	485/14_2	Orange	Orange	0
L. arvensis	*L. monelli*	COR_1.60	485/14_3	Orange	Orange	0

*Population information:

- HIN: L. arvensis. Spain, Huelva, Hinojos (37º17'42.6"N, 6º58'05"W)

- COR: L. arvensis. France, Corsica, Solenazara (41°51'19.6"N, 9°21'43.7"E)

- 630/14: L. monelli. Spain. Cádiz. Barbate. Breña de Barbate (36°10'39.3"N,

5°58'56.1"W)

- 485/14: L. monelli. Spain. Tarragona. Mont-Roig (41º5'18"N, 0º56'5"E)

- 151/14: L. monelli. Portugal. Natural Reserve das Lagoa de St André

(38º0'55.2"N, 8º49'11.1"W)

1021

1022   **Appendix 1—table 6**. Climatic differences among morphs. Comparisons indicate significance of differences among morphs (e.g. between the

1023   two orange morphs or between the two color morphs within *L. arvensis*) and were Bonferroni adjusted for 76 comparisons (nineteen variables x

1024   four morph comparisons).

	Variable means (SE)				Morph Comparisons (Kruskal Wallis p values)			
	arvensis orange	arvensis blue	monelli orange	monelli blue	arvensis difference	orange difference	blue difference	monelli difference
AnMnTmp	11.839 (0.067)	15.912 (0.093)	14.508 (0.35)	15.21 (0.076)	< 0.001	< 0.001	< 0.001	1.0
MnTmpWrmQtr	18.71 (0.07)	22.96 (0.083)	22.027 (0.32)	21.659 (0.077)	< 0.001	< 0.001	< 0.001	1.0
MnTmpCldQtr	5.534 (0.084)	9.604 (0.129)	7.8 (0.432)	9.464 (0.11)	< 0.001	< 0.001	1.0	0.04
AnnPrcp	768.135 (5.65)	541.901 (8.085)	534.675 (25.145)	663.231 (7.37)	< 0.001	< 0.001	< 0.001	0.002
PrcpWetMo	96.007 (0.794)	87.062 (1.458)	77.805 (3.642)	101.085 (1.169)	< 0.001	0.003	< 0.001	< 0.001
PrcpDryMo	34.857 (0.427)	8.609 (0.441)	8.078 (1.151)	7.946 (0.243)	< 0.001	< 0.001	1.0	1.0
PrcpSeas	32.321 (0.425)	59.064 (0.891)	52.067 (1.748)	56.902 (0.445)	< 0.001	< 0.001	1.0	1.0
PrcpWetQt	266.99 (2.216)	234.572 (4.038)	211.078 (10.287)	281.789 (3.318)	< 0.001	< 0.001	< 0.001	< 0.001
PrcpDryQt	121.847 (1.295)	41.244 (1.606)	47.494 (4.703)	42.091 (0.853)	< 0.001	< 0.001	1.0	1.0
PrcpWmrQt	152.319 (1.603)	51.462 (1.897)	58.494 (5.645)	47.816 (0.897)	< 0.001	< 0.001	1.0	1.0
PrcpCldQt	221.595 (2.408)	206.694 (4.206)	172.61 (10.381)	265.361 (3.44)	0.582	0.022	< 0.001	< 0.001
DiurnRng	8.312 (0.038)	10.038 (0.077)	10.438 (0.217)	9.737 (0.08)	< 0.001	< 0.001	1.0	0.698
Isoth	36.593 (0.107)	40.213 (0.167)	39.281 (0.491)	41.305 (0.108)	< 0.001	< 0.001	0.002	0.05
TmpSeas	541.797 (2.946)	549.092 (4.9)	584.242 (13.062)	498.631 (5.108)	1.0	1.0	< 0.001	< 0.001
MxTmpWmMo	24.626 (0.087)	29.884 (0.11)	29.34 (0.366)	28.619 (0.13)	< 0.001	< 0.001	< 0.001	1.0
MnTmpCoMo	1.703 (0.081)	4.712 (0.142)	2.594 (0.454)	4.862 (0.126)	< 0.001	1.0	1.0	< 0.001
TmpAnRng	22.923 (0.104)	25.172 (0.195)	26.747 (0.56)	23.757 (0.219)	< 0.001	< 0.001	< 0.001	< 0.001

MnTmpWetQtr	10.711 (0.098)	12.542 (0.128)	11.798 (0.465)	10.655 (0.097)	< 0.001	1.0	< 0.001	1.0
mnTmpDryQtr	13.224 (0.178)	21.795 (0.179)	21.802 (0.297)	21.399 (0.08)	< 0.001	< 0.001	1.0	1.0

1025

1026

Appendix 1—table 7. Logistic regression models for each species on a subset on non-correlated climate variables. The best models using AIC for *L. arvensis* (Arv) and *L. monelli* (Mon) are presented. "Good models" have lower, but close AIC values and reinforce the importance of the variables included in the best model.

L. arvensis **Best Model**

Coefficients	Estimate	Std. Error	z value	p
Intercept	14.5510859	0.8911108	16.33	< 0.0001
PrcpSeas	-0.0453151	0.0028234	-16.05	< 0.0001
Isoth	-0.1980946	0.0168407	-11.76	< 0.0001
TmpSeas	-0.0070202	0.0005796	-12.11	< 0.0001

L. arvensis **Good Models**

Intercept	Isoth	MeanTemp WarmQtr	PrcpSeas	PrcWarm Mon	TmpSeas	df	Delta AIC
47.92	20.240		-37.950	2.044	-46.670	5	1.20
58.550	18.270	1.649	-39.18		-57.010	5	1.96
45.220	19.610	5.531	-37.94	2.695	-44.090	6	2.80

L. monelli **Best Model**

Coefficients	Estimate	Std. Error	z value	p
Intercept	-1.568856	0.635342	-2.469	0.0135
PrcpWetMo	-0.035533	0.005195	-6.840	< 0.0001
MnTmpWetQtr	0.229809	0.049652	4.628	< 0.0001

L. monelli **Good Models**

Intercept	Isoth	MeanTemp WarmQtr	PrcpSeas	PrcWarm Mon	TmpSeas	df	Delta AIC
22.280	-81.070	215.70	20.42	-24.93	-20.80	6	1.68
46.480	-89.320	210.80		-24.22	-44.92	5	5.67
-0.02067	-70.320	210.30		-17.75		4	5.82

1040 **Appendix 1—table 8**. Significance values from randomization tests indicating whether

1041 orange or blue populations are more different from one another climatically than

1042 expected by chance.

1043

	Orange divergent	Blue divergent	Conclusion
AnMnTmp	< 0.001	0.999	Orange more divergent
MnTmpWrmQtr	< 0.001	0.999	Orange more divergent
MnTmpCldQtr	< 0.001	0.999	Orange more divergent
AnnPrcp	< 0.001	0.999	Orange more divergent
PrcpWetMo	0.205	0.794	Not Significant
PrcpDryMo	< 0.001	0.999	Orange more divergent
PrcpSeas	< 0.001	0.999	Orange more divergent
PrcpWetQt	0.039	0.96	Orange more divergent
PrcpDryQt	< 0.001	0.999	Orange more divergent
PrcpWmrQt	< 0.001	0.999	Orange more divergent
PrcpCldQt	0.796	0.203	Not significant
DiurnRng	< 0.001	0.999	Orange more divergent
Isoth	< 0.001	0.999	Orange more divergent
TmpSeas	0.702	0.297	Not significant
MxTmpWmMo	< 0.001	0.999	Orange more divergent
MnTmpCoMo	0.410	0.958	Not significant
TmpAnRng	< 0.001	0.999	Orange more divergent
MnTmpWetQtr	0.999	< 0.001	Blue more divergent
MnTmpDryQtr	< 0.001	0.999	Orange more divergent

1044

1045

Appendix 1—**table 9.** Plant population locations and herbarium accession numbers for *Lysimachia monelli* samples used for RNA-seq and

UHPLC-MS (flavonoid profiling) (marked with a X). Voucher specimens were deposited in the University of Seville Herbarium (SEV).

Population	Color	Country	Location	Lat.	Long.	RNA-Seq	UHPLC-MS	**Accession #**
1/17-MS	Blue	Spain	Huelva. Almonte	37.055472	-6.541722	X	X	SEV285033
2/17-MS	Blue	Spain	Huelva. Pinares de Hinojos	37.217917	-6.417889	X	X	SEV287947
3/17-MS	Blue	Spain	Sevilla. Bormujos	37.371861	-6.082139	X	X	SEV287606
4/17-MS	Blue	Spain	Sevilla. Bellavista	37.330694	-5.964528		X	SEV287605
6/17-MS	Blue	Spain	Sevilla. Dos Hermanas	37.275042	-5.932889	X	X	SEV286835
7/17-MS	Blue	Spain	Cádiz. Carretera entre Borneo y Espera	36.836028	-5.752167		X	SEV286849
10/17-MS	Blue	Spain	Sevilla. Entre los Molares y Utrera. Km 5	37.162364	-5.760236	X	X	NA
LM_8	Blue	Spain	Cádiz. Barbate. Breña de barbate	36.177583	-5.982250	X	X	SEV286494
LM_24	Blue	Portugal	Carrapateira	37.193333	-8.905278	X	X	SEV286463
LM_28	Blue	Portugal	Sainto André	38.015333	-8.819750		X	SEV284791
ES_19	Orange	Spain	Tarragona. Urb. Sant Jordi D'Alfama	40.924883	0.813133	X		SEV285130
SAR_2	Orange	Italy	Sardinia. Isola de Sant Pietro, Cala Fico	39.154967	8.228783	X	X	SEV287100

SAR_3	Orange	Italy	Sardinia. Isola de Sant Pietro, Cabo Sandalo, Faro	39.144100	8.224050	X	X	SEV287102
SAR_6	Orange	Italy	Sardinia. Sa Duchesa	39.372417	8.597500	X	X	SEV287085
MAR_2	Orange	Morocco	Carretera de El Hajeb a Azrou	33.527778	-5.306944		X	SEV289309
MAR_4	Orange	Morocco	Rif. Chefchaouene	35.183767	-5.224550		X	SEV289310
MAR_8	Orange	Morocco	Medio Atlas. Kenifra. Entre Boufekran y Kenifra	33.618217	-5.430767	X	X	SEV289311
MAR_10	Orange	Morocco	Rif. Ouezzane	34.880000	-5.530000		X	SEV289312
MAR_12	Orange	Morocco	Medio Atlas. Kenifra	33.027933	-5.324650		X	SEV289314
MAR_30	Orange	Morocco	Zerhoun	34.069083	-5.455567		X	SEV289315

1048

1049

1050

 Appendix 1—table 10. Summary of sequencing and assembly results for the 20 *L. monelli* petal samples.

Sample	Population	Flower color	Number of reads	Number of filtered reads	Filtered reads aligned to the assembly (%)	Number of transcripts	Number of trinity genes*
Lymo_1	ES_19	orange	19,356,011	18,421,349	92.48	47,377	26,986
Lymo_2	SAR_3	orange	17,662,750	16,866,036	93.30	45,323	27,137
Lymo_3	SAR_2	orange	18,038,810	17,237,553	93.16	48,194	29,126
Lymo_7	SAR_6	orange	17,084,149	16,312,231	92.11	49,194	29,265
Lymo_18	SAR_6	orange	17,166,042	16,255,988	90.86	48,459	28,927
Lymo_23	MAR_8	orange	16,537,229	15,503,176	92.47	48,701	28,213
Lymo_24	MAR_8	orange	16,709,072	15,886,953	92.23	49,265	28,074
Lymo_34	ES_19	orange	17,230,125	16,466,487	92.21	49,477	28,830
Lymo_35	ES_19	orange	23,467,043	22,697,727	92.73	46,959	27,097
Lymo_36	SAR_2	orange	16,981,334	15,773,601	92.57	52,169	31,542
Lymo_40	2/17-MS	blue	18,359,772	17,602,178	93.03	46,380	26,228

Lymo_45	LM_8	blue	17,828,507	16,984,718	92.32	48,951	28,143
Lymo_8	LM_24	blue	16,850,071	16,124,259	91.26	45,973	26,624
Lymo_9	1/17-MS	blue	18,050,225	17,338,160	92.84	45,891	26,354
Lymo_11	6/17-MS	blue	19,073,958	18,182,529	92.42	47,224	26,815
Lymo_12	LM_8	blue	18,350,299	17,810,293	93.54	44,695	25,856
Lymo_13	10/17-MS	blue	17,255,892	16,498,600	92.63	46,720	26,591
Lymo_17	3/17-MS	blue	17,405,546	16,803,173	93.02	47,313	27,096
Lymo_25	3/17-MS	blue	16,982,762	15,953,718	92.29	48,271	27,650
Lymo_29	6/17-MS	blue	16,331,233	15,578,920	93.01	45,869	25,935

1052   * Trinity genes were limited to those having TMM > 1.

1053