Novel gut probiotic engages microbiota for recovery and pathobiont clearance while preventing

inflammation

Authors: Vitor Cabral[1,*,†], Rita A. Oliveira[1,3,*,†], Margarida B. Correia[1], Miguel F. Pedro[1,4], Carles Ubeda[2], Karina

B. Xavier[1,†]

Affiliations:

[1]Instituto Gulbenkian Ciência, 2780-156 Oeiras, Portugal

[2]FISABIO, València, Spain

[3]Current address: Duchossois Family Institute, The University of Chicago, Chicago, IL, USA

[4]Current address: LEAF, TERRA, Instituto Superior de Agronomia, University of Lisbon, Portugal

* These authors contributed equally and share co-first authorship of this publication

[†]To whom correspondence should be addressed:

vcabral@igc.gulbenkian.pt; kxavier@igc.gulbenkian.pt; aaoliveira@uchicago.edu

Keywords: gut microbiota; recovery; colonization resistance; probiotic; *Klebsiella*; inflammatory bowels

disease; butyrate; inflammation; intestinal pathogens; *Escherichia coli*; AIEC; *Enterobacteriaceae*

Abstract

Intestinal inflammation is at the root of several acute and chronical diseases, many times with unclear etiology where the gut microbiota, genetic predisposition, and/or environmental conditions all play a role. Its recurrence locks patients in long-lasting disease cycles that greatly diminish quality of life and might lead to life-threatening conditions.

Current treatments focus on attenuation of inflammatory episodes, through antibiotics or anti-inflammatory medication. However, while mitigating inflammation, such therapies also promote major changes in the intestinal microbiota with consequent loss of its beneficial functions, such as protection against pathogenic bacteria, leading to infections that can reactivate intestinal inflammation, resulting in relapse. Alternative and complementary treatments that can break the inflammation-treatment-infection-inflammation cycle are urgently needed as the next therapeutic breakthrough for intestinal inflammatory conditions.

Here, we use an Inflammatory Bowel Disease (IBD) model that is susceptible to inflammatory episodes to demonstrate how the use of a protective native gut commensal bacterium can break the disease cycle.

Our probiotic therapy, centered of the gut commensal bacterium *Klebsiella* sp. ARO112, can transiently colonize an imbalanced and infected microbiota, preventing the relapse of intestinal inflammation, by promoting a faster recovery of the gut microbial community. This ARO112-induced recovery is driven by the resurgence of native bacteria and butyrate production, halting the relapse of inflammation, and displacing the pathobiont Adherent and Invasive Escherichia coli (AIEC), thereby halting the disease cycle by avoiding the need for further treatments.

This promising biotherapy could be used to break the inflammation-treatment-infection-inflammation cycle that traps most not only IBD patients, but also in many other inflammatory based conditions, allowing for their gut microbiota to recover from debilitating diseases and increasing their quality of life.

Introduction

The gut microbiota plays a crucial role in human health and disease. An abundant and diverse intestinal community of microorganisms performs essential functions, including providing colonization resistance to potential pathogenic external bacteria. This protective capacity can be significantly compromised by environmental perturbations that alter the population composition, such as changes in diet or medical treatments, like antibiotics [1]–[3]. These shifts in populational dynamics, induced by dietary changes and antibiotic treatments, can be challenging to rectify, leaving patients more susceptible to infections and other medical complications [4].

The pursuit of interventions to restore dysbiotic gut microbiotas has gained momentum. Currently, both basic research and clinical studies encompass a range of approaches, from introducing single protective bacterial species to well-defined bacterial communities or even complete microbiota, as in fecal microbiota transplants (FMT) [5]. Although the United States Food and Drug Administration has recently approved FMT for the successful treatment of recurrent *Clostridioides difficile* infections in humans [6], the undefined nature of these procedures poses safety risks [7], [8]. These risks underscore the need for the identification of defined strategies, which could involve either single- or multi-strain bacterial cocktails. Next-generation probiotic bacteria, defined as bacterial species native to the host, which upon ingestion and/or colonization, provide benefits to the host, offer promise for restoring microbiota functions. This requires an extensive understanding of the benefits, potential drawbacks, and the organism's safety profile, all of which are important to its therapeutic importance.

Patients with high susceptibility to infections often undergo extensive antibiotic treatments, rendering them more susceptible to recurrent infections. These patients require innovative approaches to complement current treatments and break the cycle of disease recurrence. Inflammatory Bowel Diseases (IBD) are a group of multi-factorial pathologies with a complex etiology, where genetic predisposition, gut microbiota, and environmental conditions play a role [9]. Three outcomes emerge as key aspects of disease cycles that afflict IBD patients: gut microbiota imbalances [10], increased and persistent intestinal inflammation [11], and susceptibility to intestinal infections [12]. To manage inflammation flares, IBD patients are often prescribed antibiotics or anti-inflammatory medication [13]. However, these treatment options can inadvertently harm the microbiota, often exacerbating the proinflammatory intestinal environment [14], thus perpetuating the cycle. The consequent loss of natural protection provided by a balanced microbiota often leads to infections by *Enterobacteriaceae* bacteria, like the adherent-invasive *Escherichia coli* (AIEC), which can also trigger inflammatory episodes [15].

Probiotics are often considered as potential alternative or complementary treatments for intestinal diseases, but their benefits for complex pathologies like IBD have been limited [16]. *E. coli* Nissle 1917 has

been proposed as a potential therapy for IBD [17]. However, despite various proposals and even clinical trials [18], the European Society for Clinical Nutrition and Metabolism, recommends against probiotics for treatment nor prevention of Crohn's Disease [19], owing to lack of promising results obtained thus far. Therefore, therapeutic probiotics warrant further research, as their efficacy in patients has been inconsistent [20]–[23].

In previous work, we have identified a commensal intestinal bacterium, *Klebsiella* sp. ARO112 from the *Enterobacteriaceae* family. ARO112 provides partial colonization resistance against an invading *Escherichia coli* K-12 MG1655 strain and *Salmonella enterica* Typhimurium in a streptomycin-induced intestinal dysbiosis murine model [24]. ARO112 can also displace the invader *E. coli*, when administered as a treatment, i.e., after colonization by the invading *E. coli*, in the presence but also in the absence of other microbiota members (in ex-germ-free mice). The ability to displace an already established bacterial invader is not a standard feature, even among probiotic strains. Research has shown that commercially available probiotics can partially displace certain pathogenic bacteria but with varying efficacy [25]. Our previous findings have prompted us to test the potential of ARO112 strain as a next-generation probiotic, capable of delaying or eliminating colonization by other *Enterobacteriaceae* invading the gut community during dysbiotic events.

In this study, we assessed the safety of ARO112 strain by evaluating its pathogenic potential for an assembly of pathogenic traits commonly associated with *Enterobacteriaceae* clinical isolates. Our results show that ARO112, compared to other well-characterized *Enterobacteriaceae,* has low pathogenic potential, limited efficiency in acquiring or maintaining plasmids carrying antibiotic resistance, and does not induce host inflammation, unlike *E. coli* Nissle 1917. Simultaneously, ARO112 displayed a competitive advantage for colonization against different species from the Pseudomonadota phylum (formerly known as Proteobacteria). Its competitive properties and lack of inflammatory potential had a significant impact in an IBD mouse model, where ARO112 exhibited strong potential as a next-generation probiotic, reducing infectious bacterial loads, preventing gut inflammation, and promoting butyrate-producing microbiota recovery from antibiotic-induced dysbiosis.

Results

ARO112 encodes fewer predicted pathogenic traits when compared with other *Enterobacteriaceae*

To address the potential of ARO112 as a probiotic, we performed a genome comparison analysis with

fifteen *Enterobacteriaceae* strains to assess their pathogenic potential, including five *E. coli* strains, five

Klebsiella pneumoniae strains, and five non-*pneumoniae Klebsiella* strains, including ARO112 (Supplementary

table 1). These strains were selected in order to incorporate human (clinical) and murine bacterial isolates,

including pathogenic, commensal, and one established probiotic strain, as well as type or laboratory strains

for these groups. With this broad strain selection, we expected to involve a wide range of encoded virulence

traits and bacteria with different pathogenic potential.

A whole genome-based phylogenetic tree segregates the fifteen strains into three major clades,

differentiating among *E. coli*, *K. pneumoniae*, and non-*pneumoniae Klebsiella* strains (Fig. 1a). The non-

pneumoniae Klebsiella clade shows *Klebsiella* sp. ARO112 clustering with *Klebsiella grimontii* (Kgr) type strain

and close to *K. michiganensis* (Kmi) type strain, and *Klebsiella oxytoca* MBC022 mouse commensal clustering

with the *K. oxytoca* DSM5175 (Kox) type strain. This placement of *Klebsiella* sp. ARO112 closer to Kgr than

Kmi had already been shown by others [26], [27], where ARO112 was compared to other non-*pneumoniae*

Klebsiella strains isolated from humans. Osbelt and colleagues have shown that ARO112 strain is close, but

not part of, the cluster of 3 *K. grimontii* tested [26], raising the hypothesis that ARO112 might not be a *K.*

grimontii species. In the *E. coli* clade, the probiotic *E. coli* Nissle 1917 (EcN) and the pathobiont AIEC (LF82)

strains were phylogenetically closer to each other than to both non-pathogenic *E. coli* K-12 (MG1655) and *E.*

coli B (EcB) laboratory strains and to the multi-resistant clinical isolate *E. coli* (Ec1898) (Fig. 1a). In the *K.*

pneumoniae clade, the three clinical isolates cluster together (MH258, Kp1012, and Kp834), while the other

two Kpn strains (NCTC9633 and ATCC43816) form a distinct cluster.

To predict the genome-encoded pathogenic potential of these strains we resorted to 9 databases

reported in Bacterial and Viral Bioinformatics Resource Center [28] for virulence factors, drug targets and

transporters, and antibiotic resistance genes. The presence/absence of the pathogenic properties segregated

the three clades represented in the phylogenetic tree (Fig. 1b,c). We analyzed the pathogenic properties that

are ubiquitous (present in all 5 strains) or non-ubiquitous (not present in all 5 strains) of each group

(Supplementary Table 2). In the non-*pneumoniae Klebsiella* strains, there are 64 absent properties when

compared to *K. pneumoniae* strains, of which 48% encode for transporters and membrane proteins,

indicating an overall lower pathogenic potential predicted for the non-*pneumoniae Klebsiella* group. The

difference in absent predicted virulence properties of the non-*pneumoniae Klebsiella* strains was even higher

when compared with *E. coli* strains (403), which mainly included genes encoding for transporters, membrane

proteins, and genes potentially conferring resistance to antibiotics (Fig. 1b, Supplementary Table 2). There

are other properties absent in the non-*pneumoniae Klebsiella* strains, but present in the *E. coli* strains that

are notable, such as genes related to lipopolysaccharide (LPS) biosynthesis and assembly, flagellum, type IV

pilus, toxins, ethanolamine utilization, and hemolysin; all of which having been associated with virulence and

pathogenesis [29]–[32].

We investigated the gene products and predicted genome-encoded pathogenic traits in common

within each clade, i.e., shared by all 5 strains tested per clade. A comparison between gene products, which

consider the whole genome (Supplementary Table 3), and the predicted genome-encoded pathogenic

properties (Supplementary Table 2), shows that the three clades share a higher percentage of predicted

pathogenic traits (47.36%) than gene products (14.23%; Extended Data Fig. 1a). Also of note is the high

percentage of gene products that are not shared by the tested strains of the three clades (>75%), when

compared to the lower percentage of predicted pathogenic traits that are not shared by all members of the

tested clades (<20%).

To assess the presence of gene products that might have clinical relevance, but not be included in

the available pathogenic properties databases, we analyzed the genomes of each strain for the number of

gene products related to the following categories: toxin-antitoxin, multidrug resistance, siderophores, LPS,

natural competence, conjugation, transposase, and phages/integrases. With this analysis, ARO112 shows

consistently lower number of gene products within these categories, and no gene product related to

conjugation or conjugal proteins, which are important for bacteria to acquire antibiotic resistance genes

(Extended Data Fig. 1b). Moreover, even though the genome analysis displays Kgr as the closest tested

relative to ARO112 (Fig. 1a) and the predicted pathogenic traits analysis hints at Kmi as closely related to

ARO112 (Fig. 1b), both Kgr and Kmi clearly differ from ARO112 when focusing on gene products for such

clinically-relevant categories (Extended Data Fig. 1b), indicating that ARO112 might be a different species

altogether.

Upon adding data for the abundance of each pathogenic property (Supplementary Table 4), instead

of the binary analysis of presence/absence (Fig. 1c), the strains were no longer segregated by taxonomy, and

instead segregated in two clusters along PCo 2 (Fig. 1d). One cluster included a known multi-drug resistant

(MDR) *K. pneumoniae* strain (MH258) and three strains isolated from hospitalized patients. The top 10

pathogenic traits driving the separation of the two clusters include several antimicrobial resistance-

associated properties (6/10; Supplementary Table 5), and accordingly these properties were enriched in the

cluster including the MDR *K. pneumoniae* strain. ARO112 was in the second cluster together with MBC022,

both isolated from non-antibiotic-treated gut microbiomes [24], and together with known non-MDR strains

like MG1655 (Fig. 1d). These results show that the predicted pathogenic properties encoded by the different

bacteria correlate with the taxonomy. However, the abundance of these properties segregates by the drug

resistance-encoded genes of individual isolates, regardless of taxonomy. ARO112 separated from known

MDR strain MH258 and clinical isolates Kp1012, Kp834, and Ec1898, and showed overall less predicted
pathogenic properties than the ensemble of other tested strains.

ARO112 shows fewer traits associated with nosocomial infections

Given that genome predictions do not always directly translate into phenotypes, we decided to
perform several in vitro tests for traits that are generally associated with *Enterobacteriaceae* pathogenicity
in nosocomial infections and gut colonization (Fig. 2a). For that, we selected a subset of the aforementioned
strains, including four *E. coli* strains (non-pathogenic K-12 MG1655 strain, pathobiont AIEC, probiotic EcN,
and clinical isolate Ec1898), three *K. pneumoniae* strains (clinical isolates MH258, Kp1012, and Kp834), and
two non-*pneumoniae Klebsiella* strains from our strain collection (MBC022, ARO112). We performed these
tests under two different laboratory conditions (nutrient rich – LB – and nutrient poor media – M9 salts
minimal medium with glucose) to evaluate if these pathogenic traits vary in different environments.

We started by testing antibiotic resistance, as this is the main driver of clinical problems associated
with *Enterobacteriaceae* [33], due to their efficient ability to acquire and share antibiotic resistance genes
from other bacteria through mechanisms like horizontal gene transfer (HGT) [34]. Our results showed that
ARO112, similar to MG1655, EcN, and MBC022, is not MDR since it displays resistance to less than three
classes of antibiotics (Fig. 2b). The other five strains tested showed resistance to three or less (AIEC),
depending on the medium, or more than three (Ec1898, Kp1012, Kp834, MH258) classes of antibiotics,
including, as expected, the known MDR strain MH258 in addition to the clinical isolates from hospitalized
patients Ec1898, Kp1012, and Kp834 (Fig. 2b), confirming their MDR status (Fig. 1d). Most of all, these results
confirm that ARO112 is not MDR and highlight that bacterial phenotypes are context dependent, with
minimal medium, frequently, potentiating resistance phenotypes.

Biofilms are microbial communities enclosed in extracellular matrix and adherent to a substrate and
can contribute to increased recalcitrance to antibiotic treatments [35]. We found that Kp1012 was the strain
that produced the most robust biofilm biomass, in minimal medium. In rich medium, most strains resulted in
low (EcN, Kp834, MBC022, ARO112) or very low (Ec1898, MH258) biofilm biomass (Fig. 2c), with only
MG1655, AIEC, and Kp1012 showing slightly higher biofilm biomass (Fig. 2c). Under minimal medium,
ARO112, Kp834, and MBC022 presented some capacity for biofilm formation, but much lower than Kp1012.
All *E. coli* strains and MH258 showed lower biofilm biomass (Fig. 2c).

Urease production is a common trait in *K. pneumonia* and *K. oxytoca* and can be problematic in
urinary tract infections, but might also be a good competitive trait for gut colonization [36]. In cultures grown
in rich medium there was no detectable urease activity, except for *K. oxytoca* MBC022 and, to a lesser extent,
ARO112, both displaying low levels of urease (Fig. 2d). In cultures grown in minimal medium all *Klebsiella*

strains, but not *E. coli* strains, showed significantly higher urease activity, with Kp1012 and MBC022 presenting the highest and Kp834, MH258, and ARO112 the lowest activities, with the latter with similar levels to rich medium (Fig. 2d). The absence of urease activity in the *E. coli* strains is expected, since this species is a known non-urease producer, albeit some clinical isolates of the pathogenic Enterohemorrhagic *E. coli* have been shown to have urease activity [37]. Strains proficient in both biofilm formation and urease production, as is the case for Kp1012, can become problematic and often associated with recurrent urinary tract infections [38].

In rich medium, which is not limited in iron, all strains showed a reduced production of iron chelators, siderophores (Fig. 2e). Interestingly, in iron-limited conditions, siderophore production was not increased in ARO112, MG1655, nor MBC022, as opposed to the rest of the strains, where a significantly higher siderophore production was observed (Fig. 2e).

When testing resistance to serum-mediated killing of bacteria grown in rich medium, only the *K. pneumoniae* strains and MBC022 showed significant resistance to and replication ability in the presence of human serum (Fig. 2f), indicating their potential capacity to survive and thrive in the bloodstream of patients. On the contrary, most *E. coli* strains (MG1655, AIEC, EcN) are very susceptible to human serum, while Ec1898 and ARO112 are not killed by nor do they replicate in human serum (Fig. 2f). Interestingly, testing these strains grown in minimal medium resulted in a significantly higher replication for MH258, increasing from 40- (in rich medium) to 140-fold. More surprising, however, was the change observed for EcN; according to what has been reported, EcN shows strong susceptibility to human serum [39] when grown in rich medium, but a strong resistance and replication capacity when grown in minimal medium (70-fold increase; Fig. 2f). Reports on this phenotype are usually performed using rich media to grow bacteria [40], but here we found that some bacteria, like EcN and MH258, show increased proliferation when grown in minimal medium. Finally, ARO112 shows no evidence of being killed by the human serum, nor does it replicate to the levels seen for most strains (Fig. 2f). Since this virulence trait directly affects the host, as it potentially allows bacteria to proliferate in the bloodstream, with life-threatening risks, ARO112 inability to proliferate is a desirable trait for a potential probiotic.

The evaluation of the pathogenic potential based on phenotypic assays gives us the resolution to the strain level regarding this selection of virulence traits. While the genome-based approach showed us that the virulence traits predicted to be present or absent in the genome correlate to the taxonomy, the selective testing of clinically-important phenotypes allows us to distinguish distinctly active virulence traits within a taxon, such as the heterogeneity within both the *E. coli* and the *K. pneumoniae* clades, but also to pinpoint strains with overall less pathogenic traits associated with nosocomial infections. Of note, ARO112 stands out with low pathogenic traits commonly associated with nosocomial infections.

ARO112 is resistant to acquisition and efficient in curing plasmids bearing antibiotic resistance

The lack of resistance found in the commensal non-clinical strains does not exclude the potential for antibiotic resistance acquisition by HGT, which is common among *Enterobacteriaceae* within complex communities, like the gut microbiota [41]. Indeed, the acquisition of antibiotic resistance, in particular by HGT events, has been shown to be problematic, with *Enterobacteriaceae* at the forefront of this increasingly concerning medical issue [34]. Therefore, we assessed the capacity of ARO112 to acquire resistance from donor bacteria through HGT, in comparison with the probiotic EcN and the MDR *K. pneumoniae* strain that had typical phenotypes of causative agents of nosocomial infections, Kp1012. We tested the transfer of a commonly used conjugative plasmid (RP4) from a donor strain (MG1655 or Kgr) into the aforementioned recipient strains. Interestingly, the already MDR strain Kp1012 had the highest acquisition rate, with a median of 83% and 25% of cells acquiring the plasmid from the MG1655 and Kgr donors, respectively (Fig. 2g). EcN presented an intermediate transfer rate, with a median of 25% and 4% acquiring the plasmid and consequent antibiotic resistance from MG1655 and Kgr, respectively (Fig. 2g). In contrast, ARO112 showed a strong resistance to plasmid acquisition, with a median of less than 4% of the population acquiring the RP4 plasmid from MG1655 donor strain, and less than 0.5% from Kgr donor strain (Fig. 2g). Next, we tested the capacity to retain the acquired plasmids, after five days of daily passages in rich or minimal media in the absence of antibiotic selection. Remarkably, ARO112 showed the highest plasmid loss, with 87% and 65% of the population losing the plasmid in rich and minimal media, respectively (Fig. 2h). In contrast, the MDR strain Kp1012 lost the plasmid in 27% and 2% of the population, in rich and minimal media, respectively, and EcN lost the plasmid in 61% and 34% of the population in rich and minimal media, respectively (Fig. 2h).

Even plasmids that are non-self-transmissible, can be transferred among bacteria, through conjugation events [42] in which non-conjugative plasmids take advantage of conjugative plasmids' machinery from the donor strain. We have also tested the capacity of ARO112, EcN, and Kp1012 to retain a non-conjugative plasmid (pMP7605) carrying an antibiotic resistance received by electroporation. Kp1012 barely lost the plasmid, both in rich (7%) and minimal media (11%; Fig. 2i), and 42% of EcN colonies lost the non-conjugative plasmid in rich medium (Fig. 2i), while no loss was observed in minimal medium (Fig. 2i). Remarkably, ARO112 showed the highest plasmid loss, with 83% and 51% of the population losing the non-conjugative plasmid when grown in rich or minimal media, respectively (Fig. 2i).

Natural plasmids are known to be more stably maintained as they often confer some fitness advantages that drive their persistence [43]. We also tested the retention capacity of a Kanamycin resistance-bearing natural plasmid(s) transferred from one of the MDR clinical isolates, Kp834, into ARO112 and EcN (Kp1012 was not tested due to its MDR complicating counterselection). EcN did not lose the plasmid, while

ARO112 lost the plasmid in over 15% (rich medium) and 30% (minimal medium) of the population (Extended Data Fig. 2b).

In summary, even though ARO112 shares with other *Enterobacteriaceae* strains the capacity to acquire plasmids, when compared to the probiotic strain EcN and the *K. pneumoniae* clinical isolate Kp1012, ARO112 displays a significantly lower efficiency in plasmid acquisition and retention, a main source of antimicrobial resistance in *Enterobacteriaceae* and, therefore, presents a reduced risk of becoming a clinical problem.

Bacterial pathogenic phenotypes can be altered upon gut colonization

Having observed that the tested phenotypes varied depending on the culture medium used, we questioned which of the tested laboratory media, rich or minimal, was resembling the phenotypical profiles of bacteria in the mammalian gut more closely. Therefore, we tested the phenotypes whose assays do not require bacterial growth, in bacteria collected directly from fecal samples from mono-colonized mice (Fig. 3a). We compared the phenotypes in ARO112 with EcN and Kp1012 collected from fecal samples of ex-germ-free mice colonized for five days with each of these bacteria, without selection or growth. Regarding urease activity, Kp1012 presented the same high urease activity detected in minimal medium-grown cultures and EcN showed the expected low urease activity also observed in all laboratory-grown cultures (Fig. 3b and 2d), ARO112 presents a variable phenotype, with half the samples showing low urease activity, similar to the results obtained with cultures grown in the laboratory, while the other half showed increased urease activity (Fig. 3b and 2d). This result, while surprising due to the low levels of urease activity in culture-grown ARO112, indicates that in this bacterium this phenotype is regulated differently in the gut environment. Interestingly, fecal siderophore measurements showed low values for all three strains, but with a similar tendency observed in minimal medium of lower values for ARO112, when compared with both EcN and Kp1012 (Extended Data Fig. 3a and Fig. 2e).

When testing for resistance to human serum after gut colonization, similar results were obtained with bacteria extracted directly from feces of mono-colonized mice (Fig. 3c) to the ones of laboratory-grown cultures (Fig. 2f). When compared to both Kp1012 and EcN, ARO112 strain is more sensitive human serum, even when grown in mice (Fig. 3c), therefore being a safer strain for clinical settings than the probiotic EcN.

We also checked the non-conjugative plasmid retention in the murine gut by having each strain colonizing germ-free mice carrying the non-conjugative plasmid pMP7605, and quantifying how much of the population retained the plasmid after five days of colonization. EcN lost the plasmid in only 11% of the population, while Kp1012 lost in 51% (Fig. 3d). Remarkably, ARO112 lost the plasmid in more than 99% of cells (Fig. 3d), showing that the recalcitrance of ARO112 to maintaining acquired antibiotic resistance is potentiated in the murine gut.

Next, we tested the host reaction to the colonization by each of the three aforementioned strains, by assessing the fecal levels of Lipocalin-2 (Lcn2), as a marker for gut inflammation [44], [45]. After 5 days of mono-colonization, fecal levels of Lcn2 in mice colonized with ARO112 were similar to germ-free mice, providing no evidence for ARO112 inducing intestinal inflammation in mono-colonized animals (Fig. 3e). This contrasted with colonization with EcN and, to an even greater extent, with Kp1012, which significantly induced Lcn2 when compared to the germ-free mice levels (Fig. 3e). Since Lcn2 decreases bacterial growth through siderophore sequestration, we then decided to test how these three strains were able to maintain growth with increasing levels of Lcn2, which is commonly increased in the inflamed intestinal environment. ARO112, EcN, or Kp1012 were grown in rich or minimal media in the absence or in the presence of increasing concentrations of Lcn2. ARO112 showed an overall higher resistance than EcN to increased levels of Lcn2 (Extended Data Fig. 3b), suggesting a better ability to survive in an inflamed gut environment.

Together, these results indicate that ARO112 could be an optimum next-generation probiotic since it has reduced resistance to human serum and low plasmid retention and low pathogenic potential. Moreover, it has low plasmid retention, as opposed to other *Enterobacteriaceae* strains. Additionally, ARO112's ability to endure high levels of Lcn2 with no inflammatory induction, as opposed to EcN, further highlights how this commensal strain is potentially safer and better suited to compete with invading bacteria in intestinal inflammatory environments.

ARO112 promotes a faster recovery from AIEC infection in an IBD mouse model

Having shown that ARO112 has less potentially harmful features than an MDR *K. pneumoniae* and the probiotic EcN and that in previous work we also see that ARO112 is able to partially displace *E. coli* MG1655 from the gut of in mono-colonized and antibiotic-treated mice [24], here, we decided to test the potential protective role of ARO112 against AIEC. The pathobiont AIEC is particularly problematic under inflammatory conditions, such as those in IBD patients. Therefore, we reasoned that the low inflammatory potential of ARO112 and its capacity to endure Lcn2 could be beneficial in the IBD context.

It was previously shown that a genetic-based murine model for IBD, Nod2$^{-/-}$ mutant mice, is more susceptible to AIEC colonization after antibiotic treatment, similar to IBD patients, as clearance of this pathobiont is slower in the Nod2$^{-/-}$ animals than in WT animals [46]. Additionally, like IBD patients, Nod2$^{-/-}$ animals are more prone to gut inflammation increases [47], [48]. Therefore, we started by comparing susceptibility of WT and Nod2$^{-/-}$ mice to colonization by AIEC in our animal facility. All mice were treated by oral gavage with a combination of two non-absorbed antibiotics used in clinical settings on IBD patients [49], vancomycin and gentamicin, once daily for three consecutive days [46] (Fig. 4 and Extended Data Fig. 4a). After antibiotic treatment, mice were gavaged with AIEC, and AIEC colonization was followed through

selective plating. As observed by others [46], Nod2$^{-/-}$ mice were more susceptible to AIEC colonization in comparison with WT mice, with ~50% of the Nod2$^{-/-}$ mice still being colonized with AIEC after 20 days of infection (Fig. 4b, control mice), while all WT mice had cleared the infection in that timeframe (Extended Data Fig. 4a, control mice). To test the therapeutic potential of ARO112, one day following AIEC infection, Nod2$^{-/-}$ control mice (PBS) were compared with Nod2-/- mice gavaged with either ARO112 strain or the probiotic EcN strain (Fig. 4a). AIEC decolonization was faster in mice treated with ARO112, with 50% of the animals being AIEC-free at 8 days of infection (7 days of probiotic treatment), compared to mice that received no probiotic treatment, in which 14 days were needed for 50% of the mice to resolve the infection, and 7 out of 15 mice still remained infected at the end of treatment (Fig. 4b,c). Surprisingly, treatment with the probiotic EcN did not help decolonization, but actually resulted in a worse outcome, with all but one tested mouse remaining infected with high levels of AIEC for the 20 days of the experiment (Fig. 4b,c). We highlight that the one mouse of this group that displaced AIEC infection, was never colonized with EcN and therefore, the displacement observed in this animal is not attributed to the EcN treatment (Extended Data Fig. 4b). When assessing the number of mice that resolve the infection in each condition, it is evident that treatment with EcN promotes a continued infection with AIEC, while the absence of probiotic treatment led to the infection being resolved in half of the population (Fig. 4b,c), thus the outcome obtained with the treatment with ARO112, in which 10 out of 12 mice (>83%) became AIEC-free by the end of the experiment, was more efficient than the control (Fig. 4b). Additionally, we observed that ARO112 is itself cleared from the intestinal microbiota throughout the experiment after AIEC, with a median of 10 days post-AIEC inoculation for the ARO112 loads to drop below detection level in 50% of mice (Extended Data Fig. 4b), having disappeared in 9 out of the 12 mice tested (75%) by the end of the experiment (Extended Data Fig. 4c). Interestingly, the protective effect of ARO112 observed in Nod2$^{-/-}$ mice was not seen in WT mice treated with the same antibiotic regimen, where AIEC decolonization was observed in all the animals independently of the treatment (Extended Data Fig. 4a).

We have previously shown [24] that ARO112 could partially displace MG1655 in WT mice with a more severe antibiotic treatment of two weeks of streptomycin in the drinking water, which drastically affects the microbiota composition and facilitates *Enterobacteriaceae* colonization that compete for the available carbon sources. We tested the effect of ARO112 and EcN on AIEC in streptomycin-treated WT mice where, in contrast to what we had observed with MG1655, neither ARO112 nor EcN were able to displace AIEC (Extended Data Fig. 4d). Moreover, in the absence of a microbiome in mice mono-colonized with AIEC, a limited displacement of AIEC was obtained upon co-colonization with ARO112 (~4 fold; Extended Data Fig. 4e), which was similar to what was observed when co-colonized with other *Enterobacteriaceae* like the MDR clinical isolates Ec1898 and Kp1012 (>4 and ~6, respectively; Extended Data Fig. 4f,h), but lower than the >10 fold seen previously against MG1655 [24] or against other Pseudomonadota species like *Vibrio cholerae* (>38;

Extended Data Fig. 4g). Interestingly, when co-cultured in vitro in minimal medium to test for competition for carbon sources, ARO112 had minimal effect on AIEC growth as opposed to EcN that seemed to highly reduce AIEC loads (Extended Data Fig. 4i). Together, these results show that ARO112 does not strongly compete directly with AIEC in these environments and indicate that the mechanisms explaining the displacement of AIEC in the Nod2$^{-/-}$ animals do not seem to rely on a direct ARO112 effect on AIEC colonization and rather indicate the potential involvement of the gut microbiota. This contrasts from what we showed before for MG1655, where we could demonstrate that the reduction in MG1655 colonization mediated by ARO112, was largely mediated by the direct interaction between these 2 species through nutrient competition as the ARO112-dependent inhibition of MG1655 could be observed with in vitro cultures and in germ-free mono-colonized mice [24]. This highlights that ARO112 can have a broad range effect on different Pseudomonadota pathogens, and most likely with different mechanisms to outcompete other bacteria.

ARO112 treatment promotes microbiota recovery, namely potential butyrate-producing bacteria depleted by antibiotic treatment

Whether ARO112 effect on AIEC in the IBD model is a result of the direct interplay with the host genetic background, the gut microbiota, the inflammatory status, or all of the above, is not clear. However, our results indicate that this outcome does not seem to rely on a direct ARO112 effect on AIEC growth. Nonetheless, the fact that we observed that in highly damaged gut microbiotas due to severe antibiotic regimens (14 days of streptomycin) or in the absence of a gut microbiota (ex-germ-free animals), ARO112's effect on AIEC was greatly diminished, together with the fact that ARO112 itself was removed from the gut after the AIEC displacement in the Nod2$^{-/-}$ mice, led us to evaluate a possible role of the gut microbiota in this process. We analyzed the microbiota composition of Nod2$^{-/-}$ mice by 16s rRNA gene sequencing before (pre-AB) and after (post-AB) antibiotic treatment, as well as post-AIEC colonization and after treatments with PBS, EcN, or ARO112. We observed that mice treated with ARO112 showed a significant recovery in richness and a tendency in recovery of diversity, unlike the long-lasting decrease of both metrics in the mice that received no probiotic treatment or EcN, which persisted as low as with antibiotic treatment (Fig. 5a,b). This microbiota recovery promoted by the treatment with ARO112 can help explain both the better displacement of AIEC in this group, but also the displacement of ARO112 itself (Extended Data Fig. 4b,c), as we expect the recovering microbiota to develop colonization resistance.

We then analyzed which of the most abundant taxa (37 taxa representing an average of 98% abundance of all taxa) were affected by the antibiotic treatment. Out of these 37 taxa, 22 (59%) were depleted by the antibiotics, while 7 (19%) were enriched and 8 (22%) remained unchanged during the

antibiotic treatment (Fig. 5c). The mice from the control group or treated with EcN were able to recover 14% (3 of 22) taxa depleted by the antibiotic treatment (Fig. 5c). When treated with ARO112 strain, mice were able to recover 41% (9 of 22) taxa depleted by the antibiotic treatment (Fig. 5c).

A deeper analysis of the groups of bacteria affected by antibiotic treatment showed that of the 8 taxa of potential butyrate producers depleted by the antibiotics, 4 recovered upon almost 20 days post-treatment with ARO112 (*Dorea*, *Lachnospiraceae*, *Oscillibacter* and *Ruminococcaceae*), while only 1 was recovered in the control (*Lachnospiraceae*) and another with EcN treatment (*Alistipes*) (Fig. 5c). Overall, we can see a better and faster recovery of these 4 taxa upon treatment with ARO112, with 3 of them showing significantly increased levels as early as 4 days post-treatment (day 5 post-infection with AIEC), one of which, *Lachnospiraceae*, already recovered to pre-AB levels (Fig.5d). Animals that were not treated with ARO112 either show a delayed recovery or no recovery at all of these potential butyrate producers (Fig. 5d).

PCoA analysis shows that before antibiotic treatment (pre-AB) samples cluster tightly together, whereas after (post-AB) samples were more dispersed (Extended Data Fig. 5a), revealing a stochastic effect of the antibiotics. Treatment with PBS (Ctrl) increased the dispersion of the samples, with 29% (4/14) of samples positioning close to pre-AB samples, 50% (7/14) remaining close to the post-AB samples, and 21% (3/14) in neither proximity. Treatment with EcN resulted in 90% (9/10) of samples clustering tightly and away from both pre- and post-AB samples, while 1 sample was positioned with the pre-AB samples (this was the aforementioned mouse that was resisted colonization by EcN). Interestingly, treatment with ARO112 probiotic showed 40% (4/10) of samples placed close to the pre-AB samples, while 50% (5/10) remained in close to post-AB samples, and only 1 sample occupying a region apart from both pre- and post-AB samples (Extended Data Fig. 5a). Plotting only the 4 taxa of putative butyrate producers that are depleted by antibiotics and recover after treatment (*Dorea*, *Oscillibacter*, *Lachnospiraceae*, and *Ruminococcaceae*), we noticed that, unlike the treatment with EcN, the microbiota composition treated with PBS or ARO112 probiotic cluster closely with pre-AB samples (Extended Data Fig. 5b).

IBD patients experience blooms of Pseudomonadota, in particular after antibiotic treatment, a reason of concern due to the several members of this phylum that present pathogenic potential. Accordingly, we also observed 3 taxa of Pseudomonadota, *Escherichia/Shigella*, *Enterobacteriaceae*, and *Proteus,* that are usually considered pathobionts, that bloomed after antibiotics. This bloom was transient in the control group and the group treated with ARO112, but not with EcN treatment (Fig. 5c; Extended Data Fig. 5d).

In summary, here we show how in a mouse model susceptible to infection and prone to inflammation, ARO112 targeted several of the microbiota-based IBD hallmarks, by displacing the invading pathobiont AIEC, while promoting recovery of the dysbiotic microbiota, by reducing Pseudomonadota and increasing potential butyrate producers.

Probiotic therapy with ARO112 enhances microbiota production of butyrate and prevents intestinal inflammation

As described in the above section, treatment with ARO112 leads to a better recovery of microbiota richness and diversity, including potential butyrate-producing bacteria. Studies have tagged short-chain fatty acids (SCFAs) like butyrate, acetate, and propionate as important metabolites in IBD, with depletion of butyrate-producing gut bacteria by antibiotic treatments being considered a clinical outcome in these patients [14], [50]. Moreover, oral administration of butyrate to improve IBD outcomes in patients has been proposed [51]. Therefore, our observation that ARO112 treatment promotes potential butyrate producers in the microbiota, possibly leading to an increased and sustained production of butyrate within the microbiota and gut epithelium, instead of the need for continuous exogenous administration, could be very relevant. For that reason, we have assessed SCFA concentrations in fecal samples of mice treated with probiotics or untreated (Fig. 6a). In agreement with our previous results showing the increase of potential butyrate-producing bacteria, treatment with ARO112 resulted in a clear increase of fecal levels of butyrate when compared to mice treated with PBS or with EcN (Fig. 6a). An increase in isobutyrate (an isomer of butyrate) and acetate was also observed with the ARO112 treatment in comparison with the control group and EcN-treated mice, although not statistically significant. Propionate levels were significantly higher with ARO112 than with PBS, but similar to the levels in EcN group. On the other hand, succinate showed higher, but not significantly, values in mice treated with EcN, when compared to mice treated with PBS or ARO112 probiotic (Fig. 6a).

Butyrate has been linked to healthy gut states, with anti-inflammatory properties [52], while succinate has been touted as an inflammation marker [53]. Therefore, we used Lcn2 to evaluate the intestinal inflammatory response of the Nod2$^{-/-}$ mice infected with AIEC and treated with PBS, ARO112, or EcN. Interestingly, treatment with EcN not only led to persistent infection by AIEC but also resulted persistently high Lcn2 levels (Fig. 6b), unlike PBS treatment, in which Lcn2 levels varied between animals, having a significant but transient increase at day 5 of infection (Fig. 6b). More importantly, mice treated with ARO112 showed persistently low levels of Lcn2, never significantly varying from the levels prior to antibiotic treatment (Fig. 6b).

When assessing metrics for infection and recovery, such as microbiota diversity and richness, infection duration, intestinal inflammation, and intestinal butyrate levels, we have noticed that in the control group there is a significant negative correlation between infection duration and both diversity and richness, supporting a link between microbiota recovery and clearance of infection (Supplementary Table 7). On the other hand, mice treated with ARO112 do not show that significant correlation, demonstrating that the

infection clearance is not solely linked to microbiota recovery, with other therapeutic effects in play
(Supplementary Table 7).

The reduced Lcn2 levels observed upon treatment with ARO112 is befitting the previous data
showing increased butyrate in the gut, since a link between this SCFA and low inflammatory states has been
made before [52]. Similarly, the higher levels of succinate present in mice treated with EcN are also in
agreement with their higher Lcn2 levels. However, because the preventive effect on inflammation promoted
by ARO112 treatment was seen as early as 5 days post-infection, we can speculate that the recovery of
Lachnospiraceae seen with ARO112 by day 5 might be responsible for the anti-inflammatory effect of
butyrate at this timepoint. These results indicate that ARO112 treatment leads to a prevention of
inflammation, which in turn can also promote recovery of the microbiota, including butyrate producers,
whose increased production of butyrate might further contribute to the anti-inflammatory intestinal
environment and further recovery of the microbiota from pathobiont expansion and dysbiosis. Moreover,
the possible pro-inflammatory potential of EcN also highlights ARO112 as a safer probiotic in both non-
inflammatory or inflammatory host states.

The ensemble of these results, namely the displacement of AIEC, the prevention of inflammation,
and the recovery of the dysbiotic microbiota and butyrate, all obtained upon treatment with ARO112, show
a three-pronged therapy targeting the three hallmarks of IBD.

Discussion

The main characteristic of a healthy intestinal microbiome is its richness and diversity of microbes
and metabolites that sustain its protective role. Its robustness, however, is greatly affected by extrinsic
factors, such as medical treatments and changing diets. Our understanding of the functional aspects driving
this protection by a balanced microbiota is still limited. The importance of identifying protecting bacteria to
be used as therapeutics is unprecedented and research has been driven by the discovery and characterization
of the beneficial effects those bacteria can provide.

Here, we showed that our previously identified microbiota member of the non-*pneumoniae*
Klebsiella complex, *Klebsiella* sp. ARO112 strain [24], is able to displace the pathobiont AIEC from an IBD
mouse model, while preventing inflammatory flares and mediating the recovery of an imbalanced
microbiota. The therapeutic effect of ARO112 is multifactorial, affecting all three main hallmarks of intestinal
inflammatory disorders: gut inflammation, dysbiosis, and susceptibility to infection. Our results show that in
the presence of ARO112 there is a faster recovery of microbiota members producers of butyrate which is
reflected in an increase in intestinal butyrate levels, a SCFA known to be important for gut health, including
in mediating clearance of MDR *Enterobacteriaceae* by direct inhibition [54]. Other studies have shown that

some members of the microbiota (belonging to *Lactobacillus* genus) can contribute to promote clearance of MDR *K. pneumoniae* by driving the recovery of *Clostridiales* [55]. Similarly, a study focusing on human samples showing colonization resistance against intestinal pathogens with high risk of bacteremia found an enrichment of non-*pneumoniae Klebsiella* species, whose protective effect was then validated in murine models [27]. A previous study had already pinpointed, and validated, *K. oxytoca* strains present in children's microbiota with protection against invading *K. pneumoniae* [26], in the aftermath of our previous publication showing ARO112 as protective against *E. coli* and *Salmonella* Typhimurium [24]. These three studies highlighting the protective role of non-*pneumoniae Klebsiella* species against *Enterobacteriaceae* invasion, display the important role of nutrient competition in this interaction. Nevertheless, all these studies are focused on colonization resistance and pathogen displacement. Here, we show that ARO112 therapeutic effect goes beyond nutrition competition, even being able to clear a clinically-important pathobiont, AIEC, in a disease model for intestinal inflammation, not relying on this direct competition effect, with an astounding >83% success rate. Our results, however, are not solely focused on the important and decisive pathogen clearance, but also showing that ARO112 probiotic treatment engages the native butyrate producers depleted by antibiotic treatment into microbiota recovery and prevent inflammatory episodes in the process.

Even though some protective capabilities seem to be shared by different non-*pneumoniae Klebsiella* species, when considering bacteria as potential probiotics, their pathogenic potential and safety also need to be considered. Our pipeline for strains' characterization shows that their pathogenic potential can be context-dependent, emphasizing how important it is to test different conditions before asserting the virulence status of a potential probiotic. We show how ARO112 presents a reduced pathogenic potential and is safer than other potential protective bacteria, such as *K. oxytoca* or *E. coli* Nissle 1917, both by displaying less pathogenic phenotypes, but also by being more recalcitrant to antibiotic resistance acquisition and maintenance compared to *E. coli* Nissle 1917, therefore not contributing to the currently increasing antibiotic resistance pandemic.

Having a safe protective microbiota member that can be used to promote a faster recovery of an imbalanced microbiota and displace pathogens or pathobionts, while preventing inflammatory flares in disease context is of great therapeutic importance, and a major leap in current probiotic research knowledge and capabilities. Further research is needed to deepen the understanding of the mechanisms driving ARO112 protection, which will open up new avenues for the identification and characterization of more microbiota members and consortia with safe therapeutic potential for clinical use, which can then be engineered to tackle a myriad of clinically relevant disorders.

Acknowledgements

We thank the members of the Bacterial Signalling Laboratory at Instituto Gulbenkian Ciência and Tanja Dapa, and João Xavier for critically reading the manuscript. In particular, we thank Joana Amaro for the help processing samples for genomic sequencing and metabolomics. We also thank Instituto Gulbenkian Ciência's Animal Facility, in particular Joana Bom, and the Genomics Facility, namely João Sobral, for the help provided during this project's execution. The authors would like to thank the Microbiology Service from the Hospital La Fé in Valência (Spain) for isolating and providing the strains Ec1898, Kp834, and Kp1012. I would like to thank Gabriel Nuñez for providing the AIEC strain (LF82), and Roberto Balbontín-Soria for providing the plasmid pMP7605. We would also like to acknowledge Nassos Typas for the suggestion to test conjugation from a clinical isolate.

This work was funded by Fundação para a Ciência e Tecnologia (FCT) projects [PTDC/BIA-MIC/6990/2020] and [SFRH/BPD/116806/2016], as well as by Municipality of Oeiras Proof-of-Concept InnOValley 2022 [IOVPoC-2022-21] and a European Commission grant [MSCA-IF-2018-843183]. Additionally, this project was also funded by InfectERA-ERANET- through FCT grant (InfectERA/0004/2015) and *Acciones complementarias* grant [PCIN-2015-094] from the Spanish Ministerio de Economía y Competitividad and the 7th Research framework program from EU, and a grant from the Spanish MICINN [PID2020-120292RB-I00], granted to KBX and CU, respectively.

Author contributions

V.C. and R.A.O conceptualized and designed the research studies. M.B.C. performed in vivo experiments with *V. cholerae* and *Ec1898*, under the supervision of R.A.O. and K.B.X.. M.F.P. performed the NMR under the supervision of K.B.X.. R.A.O. executed in vitro experiments with support from V.C.. V.C. executed in vivo experiments with AIEC and Kp1012, in SPF wildtype, SPF Nod2$^{-/-}$, and GF wildtype models, with support from R.A.O.. R.A.O. analyzed sequencing data. V.C., and R.A.O. analyzed and interpreted all the data, under the supervision of K.B.X.. V.C. and R.A.O. wrote the first draft of the manuscript with valuable input from K.B.X. All authors critically read and approved the manuscript.

Competing interests

A provisional patent application has been filed by the Gulbenkian Foundation based on this work.

References

[1] P. J. Turnbaugh, M. Hamady, T. Yatsunenko, B. L. Cantarel, A. Duncan, R. E. Ley, M. L. Sogin, W. J. Jones, B. A. Roe, J. P. Affourtit, M. Egholm, B. Henrissat, A. C. Heath, R. Knight, and J. I. Gordon, "A core gut microbiome in obese and lean twins.," *Nature*, vol. 457, no. 7228, pp. 480–484, Jan. 2009, doi: 10.1038/nature07540.

[2] E. B. Hollister, C. Gao, and J. Versalovic, "Compositional and functional features of the gastrointestinal microbiome and their effects on human health.," *Gastroenterology*, vol. 146, no. 6, pp. 1449–1458, May 2014, doi: 10.1053/j.gastro.2014.01.052.

[3] R. A. Oliveira, V. Cabral, and K. B. Xavier, "Microbiome–diet interactions drive antibiotic efficacy," *Nat. Microbiol.*, vol. 6, no. 7, pp. 824–825, 2021, doi: 10.1038/s41564-021-00926-8.

[4] A. Andremont, J. Cervesi, P.-A. Bandinelli, F. Vitry, and J. de Gunzburg, "Spare and repair the gut microbiota from antibiotic-induced dysbiosis: state-of-the-art," *Drug Discov. Today*, vol. 26, no. 9, pp. 2159–2163, 2021, doi: https://doi.org/10.1016/j.drudis.2021.02.022.

[5] R. A. Oliveira and E. G. Pamer, "Assembling symbiotic bacterial species into live therapeutic consortia that reconstitute microbiome functions.," *Cell Host Microbe*, vol. 31, no. 4, pp. 472–484, Apr. 2023, doi: 10.1016/j.chom.2023.03.002.

[6] N. Jain, T. P. Umar, A.-F. Fahner, and V. Gibietis, "Advancing therapeutics for recurrent clostridioides difficile infections: an overview of vowst's FDA approval and implications.," *Gut Microbes*, vol. 15, no. 1, p. 2232137, 2023, doi: 10.1080/19490976.2023.2232137.

[7] Z. DeFilipp, P. P. Bloom, M. Torres Soto, M. K. Mansour, M. R. A. Sater, M. H. Huntley, S. Turbett, R. T. Chung, Y.-B. Chen, and E. L. Hohmann, "Drug-Resistant E. coli Bacteremia Transmitted by Fecal Microbiota Transplant.," *N. Engl. J. Med.*, vol. 381, no. 21, pp. 2043–2050, Nov. 2019, doi: 10.1056/NEJMoa1910437.

[8] S. Gupta, B. H. Mullish, and J. R. Allegretti, "Fecal Microbiota Transplantation: The Evolving Risk Landscape.," *Am. J. Gastroenterol.*, vol. 116, no. 4, pp. 647–656, Apr. 2021, doi: 10.14309/ajg.0000000000001075.

[9] A. Goethel, K. Croitoru, and D. J. Philpott, "The interplay between microbes and the immune response in inflammatory bowel disease.," *J. Physiol.*, vol. 596, no. 17, pp. 3869–3882, Sep. 2018, doi: 10.1113/JP275396.

[10] P. Qiu, T. Ishimoto, L. Fu, J. Zhang, Z. Zhang, and Y. Liu, "The Gut Microbiota in Inflammatory Bowel Disease.," *Front. Cell. Infect. Microbiol.*, vol. 12, p. 733992, 2022, doi: 10.3389/fcimb.2022.733992.

[11] M. Fakhoury, R. Negrulj, A. Mooranian, and H. Al-Salami, "Inflammatory bowel disease: clinical
aspects and treatments.," *J. Inflamm. Res.*, vol. 7, pp. 113–120, 2014, doi: 10.2147/JIR.S65979.

[12] P. M. Irving, S. de Lusignan, D. Tang, M. Nijher, and K. Barrett, "Risk of common infections in people
with inflammatory bowel disease in primary care: a population-based cohort study.," *BMJ open*
Gastroenterol., vol. 8, no. 1, Feb. 2021, doi: 10.1136/bmjgast-2020-000573.

[13] A. B. Pithadia and S. Jain, "Treatment of inflammatory bowel disease (IBD).," *Pharmacol. Rep.*, vol.
63, no. 3, pp. 629–642, 2011, doi: 10.1016/s1734-1140(11)70575-8.

[14] P. T. Santana, S. L. B. Rosas, B. E. Ribeiro, Y. Marinho, and H. S. P. de Souza, "Dysbiosis in
Inflammatory Bowel Disease: Pathogenic Role and Potential Therapeutic Targets.," *Int. J. Mol. Sci.*,
vol. 23, no. 7, Mar. 2022, doi: 10.3390/ijms23073464.

[15] C.-L. N. Small, S. A. Reid-Yu, J. B. McPhee, and B. K. Coombes, "Persistent infection with Crohn's
disease-associated adherent-invasive Escherichia coli leads to chronic inflammation and intestinal
fibrosis.," *Nat. Commun.*, vol. 4, p. 1957, 2013, doi: 10.1038/ncomms2957.

[16] B. P. Abraham and E. M. M. Quigley, "Probiotics in Inflammatory Bowel Disease.," *Gastroenterol.*
Clin. North Am., vol. 46, no. 4, pp. 769–782, Dec. 2017, doi: 10.1016/j.gtc.2017.08.003.

[17] M. Schultz, "Clinical use of E. coli Nissle 1917 in inflammatory bowel disease.," *Inflamm. Bowel Dis.*,
vol. 14, no. 7, pp. 1012–1018, Jul. 2008, doi: 10.1002/ibd.20377.

[18] H. Matthes, T. Krummenerl, M. Giensch, C. Wolff, and J. Schulze, "Clinical trial: probiotic treatment
of acute distal ulcerative colitis with rectally administered Escherichia coli Nissle 1917 (EcN).," *BMC*
Complement. Altern. Med., vol. 10, p. 13, Apr. 2010, doi: 10.1186/1472-6882-10-13.

[19] S. C. Bischoff, P. Bager, J. Escher, A. Forbes, X. Hébuterne, C. L. Hvas, F. Joly, S. Klek, Z. Krznaric, J.
Ockenga, S. Schneider, R. Shamir, K. Stardelova, D. V. Bender, N. Wierdsma, and A. Weimann,
"ESPEN guideline on Clinical Nutrition in inflammatory bowel disease," *Clin. Nutr.*, vol. 42, no. 3, pp.
352–379, 2023, doi: https://doi.org/10.1016/j.clnu.2022.12.004.

[20] J. Suez, N. Zmora, G. Zilberman-Schapira, U. Mor, M. Dori-Bachash, S. Bashiardes, M. Zur, D. Regev-
Lehavi, R. Ben-Zeev Brik, S. Federici, M. Horn, Y. Cohen, A. E. Moor, D. Zeevi, T. Korem, E. Kotler, A.
Harmelin, S. Itzkovitz, N. Maharshak, *et al.*, "Post-Antibiotic Gut Mucosal Microbiome Reconstitution
Is Impaired by Probiotics and Improved by Autologous FMT.," *Cell*, vol. 174, no. 6, pp. 1406-
1423.e16, Sep. 2018, doi: 10.1016/j.cell.2018.08.047.

[21] E. Montassier, R. Valdés-Mas, E. Batard, N. Zmora, M. Dori-Bachash, J. Suez, and E. Elinav,
"Probiotics impact the antibiotic resistance gene reservoir along the human GI tract in a person-
specific and antibiotic-dependent manner.," *Nat. Microbiol.*, vol. 6, no. 8, pp. 1043–1054, Aug. 2021,

doi: 10.1038/s41564-021-00920-0.

[22]   J.-P. Nougayrède, C. V Chagneau, J.-P. Motta, N. Bossuet-Greif, M. Belloy, F. Taieb, J.-J. Gratadoux, M. Thomas, P. Langella, and E. Oswald, "A Toxic Friend: Genotoxic and Mutagenic Activity of the Probiotic Strain Escherichia coli Nissle 1917.," *mSphere*, vol. 6, no. 4, p. e0062421, Aug. 2021, doi: 10.1128/mSphere.00624-21.

[23]   M. Olier, I. Marcq, C. Salvador-Cartier, T. Secher, U. Dobrindt, M. Boury, V. Bacquié, M. Pénary, E. Gaultier, J.-P. Nougayrède, J. Fioramonti, and E. Oswald, "Genotoxicity of Escherichia coli Nissle 1917 strain cannot be dissociated from its probiotic activity.," *Gut Microbes*, vol. 3, no. 6, pp. 501–509, 2012, doi: 10.4161/gmic.21737.

[24]   R. A. Oliveira, K. M. Ng, M. B. Correia, V. Cabral, H. Shi, J. L. Sonnenburg, K. C. Huang, and K. B. Xavier, "Klebsiella michiganensis transmission enhances resistance to Enterobacteriaceae gut invasion by nutrition competition.," *Nat. Microbiol.*, Jan. 2020, doi: 10.1038/s41564-019-0658-4.

[25]   M. C. Collado, J. Meriluoto, and S. Salminen, "Role of commercial probiotic strains against human pathogen adhesion to intestinal mucus.," *Lett. Appl. Microbiol.*, vol. 45, no. 4, pp. 454–460, Oct. 2007, doi: 10.1111/j.1472-765X.2007.02212.x.

[26]   L. Osbelt, M. Wende, É. Almási, E. Derksen, U. Muthukumarasamy, T. R. Lesker, E. J. C. Galvez, M. C. Pils, E. Schalk, P. Chhatwal, J. Färber, M. Neumann-Schaal, T. Fischer, D. Schlüter, and T. Strowig, "Klebsiella oxytoca causes colonization resistance against multidrug-resistant K. pneumoniae in the gut via cooperative carbohydrate competition.," *Cell Host Microbe*, vol. 29, no. 11, pp. 1663-1679.e7, Nov. 2021, doi: 10.1016/j.chom.2021.09.003.

[27]   J. Schluter, A. Djukovic, B. P. Taylor, J. Yan, C. Duan, G. A. Hussey, C. Liao, S. Sharma, E. Fontana, L. A. Amoretti, R. J. Wright, A. Dai, J. U. Peled, Y. Taur, M.-A. Perales, B. A. Siranosian, A. S. Bhatt, M. R. M. van den Brink, E. G. Pamer, *et al.*, "The TaxUMAP atlas: Efficient display of large clinical microbiome data reveals ecological competition in protection against bacteremia," *Cell Host Microbe*, vol. 31, no. 7, pp. 1126-1139.e6, 2023, doi: https://doi.org/10.1016/j.chom.2023.05.027.

[28]   R. D. Olson, R. Assaf, T. Brettin, N. Conrad, C. Cucinell, J. J. Davis, D. M. Dempsey, A. Dickerman, E. M. Dietrich, R. W. Kenyon, M. Kuscuoglu, E. J. Lefkowitz, J. Lu, D. Machi, C. Macken, C. Mao, A. Niewiadomska, M. Nguyen, G. J. Olsen, *et al.*, "Introducing the Bacterial and Viral Bioinformatics Resource Center (BV-BRC): a resource combining PATRIC, IRD and ViPR.," *Nucleic Acids Res.*, vol. 51, no. D1, pp. D678–D689, Jan. 2023, doi: 10.1093/nar/gkac1003.

[29]   L. Craig, M. E. Pique, and J. A. Tainer, "Type IV pilus structure and bacterial pathogenicity.," *Nat. Rev. Microbiol.*, vol. 2, no. 5, pp. 363–378, May 2004, doi: 10.1038/nrmicro885.

[30]   P. Thiennimitr, S. E. Winter, M. G. Winter, M. N. Xavier, V. Tolstikov, D. L. Huseby, T. Sterzenbach, R.
M. Tsolis, J. R. Roth, and A. J. Bäumler, "Intestinal inflammation allows Salmonella to use
ethanolamine to compete with the microbiota.," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 108, no. 42, pp.
17480–17485, Oct. 2011, doi: 10.1073/pnas.1107857108.

[31]   C. Palmela, C. Chevarin, Z. Xu, J. Torres, G. Sevrin, R. Hirten, N. Barnich, S. C. Ng, and J.-F. Colombel,
"Adherent-invasive Escherichia coli in inflammatory bowel disease.," *Gut*, vol. 67, no. 3, pp. 574–
587, Mar. 2018, doi: 10.1136/gutjnl-2017-314903.

[32]   A. G. Buret, J.-P. Motta, T. Allain, J. Ferraz, and J. L. Wallace, "Pathobiont release from dysbiotic gut
microbiota biofilms in intestinal inflammatory diseases: a role for iron?," *J. Biomed. Sci.*, vol. 26, no.
1, p. 1, Jan. 2019, doi: 10.1186/s12929-018-0495-4.

[33]   J. P. 3rd Lynch, N. M. Clark, and G. G. Zhanel, "Escalating antimicrobial resistance among
Enterobacteriaceae: focus on carbapenemases.," *Expert Opin. Pharmacother.*, vol. 22, no. 11, pp.
1455–1473, Aug. 2021, doi: 10.1080/14656566.2021.1904891.

[34]   N. A. Lerminiaux and A. D. S. Cameron, "Horizontal transfer of antibiotic resistance genes in clinical
environments.," *Can. J. Microbiol.*, vol. 65, no. 1, pp. 34–44, Jan. 2019, doi: 10.1139/cjm-2018-0275.

[35]   G. Wang, G. Zhao, X. Chao, L. Xie, and H. Wang, "The Characteristic of Virulence, Biofilm and
Antibiotic Resistance of Klebsiella pneumoniae.," *Int. J. Environ. Res. Public Health*, vol. 17, no. 17,
Aug. 2020, doi: 10.3390/ijerph17176278.

[36]   H. L. Mobley and R. P. Hausinger, "Microbial ureases: significance, regulation, and molecular
characterization.," *Microbiol. Rev.*, vol. 53, no. 1, pp. 85–108, Mar. 1989, doi: 10.1128/mr.53.1.85-
108.1989.

[37]   M. Nakano, T. Iida, and T. Honda, "Urease activity of enterohaemorrhagic Escherichia coli depends
on a specific one-base substitution in ureD.," *Microbiology*, vol. 150, no. Pt 10, pp. 3483–3489, Oct.
2004, doi: 10.1099/mic.0.27280-0.

[38]   S. Clegg and C. N. Murphy, "Epidemiology and Virulence of Klebsiella pneumoniae," *Microbiol.
Spectr.*, vol. 4, no. 1, Feb. 2016, doi: 10.1128/microbiolspec.UTI-0005-2012.

[39]   U. Sonnenborn and J. Schulze, "The non-pathogenic Escherichia coli strain Nissle 1917 – features of
a versatile probiotic," *Microb. Ecol. Health Dis.*, vol. 21, no. 3–4, pp. 122–158, Jan. 2009, doi:
10.3109/08910600903444267.

[40]   R. Podschun, D. Sievers, A. Fischer, and U. Ullmann, "Serotypes, hemagglutinins, siderophore
synthesis, and serum resistance of Klebsiella isolates causing human urinary tract infections.," *J.
Infect. Dis.*, vol. 168, no. 6, pp. 1415–1421, Dec. 1993, doi: 10.1093/infdis/168.6.1415.

[41]   C. Ronda, S. P. Chen, V. Cabral, S. J. Yaung, and H. H. Wang, "Metagenomic engineering of the
mammalian gut microbiome in situ," *Nat. Methods*, vol. 16, no. 2, 2019, doi: 10.1038/s41592-018-
0301-y.

[42]   A. S. Millan, R. Peña-Miller, M. Toll-Riera, Z. V Halbert, A. R. McLean, B. S. Cooper, and R. C.
MacLean, "Positive selection and compensatory adaptation interact to stabilize non-transmissible
plasmids," *Nat. Commun.*, vol. 5, no. 1, p. 5208, 2014, doi: 10.1038/ncomms6208.

[43]   A. Alonso-del Valle, R. León-Sampedro, J. Rodríguez-Beltrán, J. DelaFuente, M. Hernández-García, P.
Ruiz-Garbajosa, R. Cantón, R. Peña-Miller, and A. San Millán, "Variability of plasmid fitness effects
contributes to plasmid persistence in bacterial communities," *Nat. Commun.*, vol. 12, no. 1, p. 2653,
2021, doi: 10.1038/s41467-021-22849-y.

[44]   B. Chassaing, G. Srinivasan, M. A. Delgado, A. N. Young, A. T. Gewirtz, and M. Vijay-Kumar, "Fecal
lipocalin 2, a sensitive and broadly dynamic non-invasive biomarker for intestinal inflammation.,"
PLoS One, vol. 7, no. 9, p. e44328, 2012, doi: 10.1371/journal.pone.0044328.

[45]   A. R. Moschen, T. E. Adolph, R. R. Gerner, V. Wieser, and H. Tilg, "Lipocalin-2: A Master Mediator of
Intestinal and Metabolic Inflammation.," *Trends Endocrinol. Metab.*, vol. 28, no. 5, pp. 388–397, May
2017, doi: 10.1016/j.tem.2017.01.003.

[46]   M. Drouet, C. Vignal, E. Singer, M. Djouina, L. Dubreuil, A. Cortot, P. Desreumaux, and C. Neut, "AIEC
colonization and pathogenicity: influence of previous antibiotic treatment and preexisting
inflammation.," *Inflamm. Bowel Dis.*, vol. 18, no. 10, pp. 1923–1931, Oct. 2012, doi:
10.1002/ibd.22908.

[47]   A. Couturier-Maillard, T. Secher, A. Rehman, S. Normand, A. De Arcangelis, R. Haesler, L. Huot, T.
Grandjean, A. Bressenot, A. Delanoye-Crespin, O. Gaillot, S. Schreiber, Y. Lemoine, B. Ryffel, D. Hot,
G. Nùñez, G. Chen, P. Rosenstiel, and M. Chamaillard, "NOD2-mediated dysbiosis predisposes mice
to transmissible colitis and colorectal cancer.," *J. Clin. Invest.*, vol. 123, no. 2, pp. 700–711, Feb.
2013, doi: 10.1172/JCI62236.

[48]   P. R. de Souza, F. R. Guimarães, H. Sales-Campos, G. Bonfá, V. Nardini, J. E. L. Chica, W. M. Turato, J.
S. Silva, D. S. Zamboni, and C. R. de B. Cardoso, "Absence of NOD2 receptor predisposes to intestinal
inflammation by a deregulation in the immune response in hosts that are unable to control gut
dysbiosis.," *Immunobiology*, vol. 223, no. 10, pp. 577–585, Oct. 2018, doi:
10.1016/j.imbio.2018.07.003.

[49]   R. Lev-Tzion, O. Ledder, E. Shteyer, M. L. N. Tan, H. H. Uhlig, and D. Turner, "Oral Vancomycin and
Gentamicin for Treatment of Very Early Onset Inflammatory Bowel Disease.," *Digestion*, vol. 95, no.

734   4, pp. 310–313, 2017, doi: 10.1159/000475660.

735   [50]   K. Aden, A. Rehman, S. Waschina, W.-H. Pan, A. Walker, M. Lucio, A. M. Nunez, R. Bharti, J.
736   Zimmerman, J. Bethge, B. Schulte, D. Schulte, A. Franke, S. Nikolaus, J. O. Schroeder, D. Vandeputte,
737   J. Raes, S. Szymczak, G. H. Waetzig, *et al.*, "Metabolic Functions of Gut Microbes Associate With
738   Efficacy of Tumor Necrosis Factor Antagonists in Patients With Inflammatory Bowel Diseases.,"
739   *Gastroenterology*, vol. 157, no. 5, pp. 1279-1292.e11, Nov. 2019, doi: 10.1053/j.gastro.2019.07.025.

740   [51]   A. Di Sabatino, P. Cazzola, R. Ciccocioppo, R. Morera, P. Biancheri, L. Rovedatti, L. Cantoro, A. Vanoli,
741   F. P. Tinozzi, S. Tinozzi, and G. R. Corazza, "Efficacy of butyrate in the treatment of mild to moderate
742   Crohn's disease," *Dig. Liver Dis. Suppl.*, vol. 1, no. 1, pp. 31–35, 2007, doi:
743   https://doi.org/10.1016/S1594-5804(08)60009-1.

744   [52]   J. Chen and L. Vitetta, "The Role of Butyrate in Attenuating Pathobiont-Induced
745   Hyperinflammation.," *Immune Netw.*, vol. 20, no. 2, p. e15, Apr. 2020, doi: 10.4110/in.2020.20.e15.

746   [53]   J. Connors, N. Dawe, and J. Van Limbergen, "The Role of Succinate in the Regulation of Intestinal
747   Inflammation.," *Nutrients*, vol. 11, no. 1, Dec. 2018, doi: 10.3390/nu11010025.

748   [54]   M. T. Sorbara, K. Dubin, E. R. Littmann, T. U. Moody, E. Fontana, R. Seok, I. M. Leiner, Y. Taur, J. U.
749   Peled, M. R. M. van den Brink, Y. Litvak, A. J. Baumler, J.-L. Chaubard, A. J. Pickard, J. R. Cross, and E.
750   G. Pamer, "Inhibiting antibiotic-resistant Enterobacteriaceae by microbiota-mediated intracellular
751   acidification.," *J. Exp. Med.*, vol. 216, no. 1, pp. 84–98, Jan. 2019, doi: 10.1084/jem.20181639.

752   [55]   A. Djukovic, M. J. Garzón, C. Canlet, V. Cabral, R. Lalaoui, M. García-Garcerá, J. Rechenberger, M.
753   Tremblay-Franco, I. Peñaranda, L. Puchades-Carrasco, A. Pineda-Lucena, E. M. González-Barberá, M.
754   Salavert, J. L. López-Hontangas, M. Á. Sanz, J. Sanz, B. Kuster, J.-M. Rolain, L. Debrauwer, *et al.*,
755   "Lactobacillus supports Clostridiales to restrict gut colonization by multidrug-resistant
756   Enterobacteriaceae.," *Nat. Commun.*, vol. 13, no. 1, p. 5617, Sep. 2022, doi: 10.1038/s41467-022-
757   33313-w.

758   [56]   K. Wilson, "Preparation of genomic DNA from bacteria.," *Curr. Protoc. Mol. Biol.*, vol. Chapter 2, p.
759   Unit 2.4, Nov. 2001, doi: 10.1002/0471142727.mb0204s56.

760   [57]   A. Reisner, K. A. Krogfelt, B. M. Klein, E. L. Zechner, and S. Molin, "In vitro biofilm formation of
761   commensal and pathogenic Escherichia coli strains: impact of environmental and genetic factors.,"
762   *J. Bacteriol.*, vol. 188, no. 10, pp. 3572–3581, May 2006, doi: 10.1128/JB.188.10.3572-3581.2006.

763   [58]   G. A. O'Toole, "Microtiter dish biofilm formation assay.," *J. Vis. Exp.*, no. 47, Jan. 2011, doi:
764   10.3791/2437.

765   [59]   D. B. Alexander and D. A. Zuberer, "Use of chrome azurol S reagents to evaluate siderophore

production by rhizosphere bacteria," *Biol. Fertil. Soils*, vol. 12, no. 1, pp. 39–45, 1991, doi: 10.1007/BF00369386.

[60] S. Das and M. Barooah, "Characterization of siderophore producing arsenic-resistant Staphylococcus sp. strain TA6 isolated from contaminated groundwater of Jorhat, Assam and its possible role in arsenic geocycle," *BMC Microbiol.*, vol. 18, no. 1, p. 104, 2018, doi: 10.1186/s12866-018-1240-6.

[61] J. A. Thompson, R. A. Oliveira, A. Djukovic, C. Ubeda, and K. B. Xavier, "Manipulation of the Quorum Sensing Signal AI-2 Affects the Antibiotic-Treated Gut Microbiota," *Cell Rep.*, vol. 10, no. 11, pp. 1861–1871, Mar. 2015, doi: 10.1016/j.celrep.2015.02.049.

[62] J. G. Caporaso, C. L. Lauber, W. A. Walters, D. Berg-Lyons, C. A. Lozupone, P. J. Turnbaugh, N. Fierer, and R. Knight, "Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample," *Proc. Natl. Acad. Sci.*, vol. 108, no. supplement_1, pp. 4516–4522, Mar. 2011, doi: 10.1073/pnas.1000080107.

[63] J. G. Caporaso, C. L. Lauber, W. A. Walters, D. Berg-Lyons, J. Huntley, N. Fierer, S. M. Owens, J. Betley, L. Fraser, M. Bauer, N. Gormley, J. A. Gilbert, G. Smith, and R. Knight, "Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms," *ISME J.*, vol. 6, no. 8, pp. 1621–1624, 2012, doi: 10.1038/ismej.2012.8.

[64] E. Pruesse, C. Quast, K. Knittel, B. M. Fuchs, W. Ludwig, J. Peplies, and F. O. Glöckner, "SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB.," *Nucleic Acids Res.*, vol. 35, no. 21, pp. 7188–7196, 2007, doi: 10.1093/nar/gkm864.

[65] B. J. Haas, D. Gevers, A. M. Earl, M. Feldgarden, D. V Ward, G. Giannoukos, D. Ciulla, D. Tabbaa, S. K. Highlander, E. Sodergren, B. Methé, T. Z. DeSantis, J. F. Petrosino, R. Knight, and B. W. Birren, "Chimeric 16S rRNA sequence formation and detection in Sanger and 454-pyrosequenced PCR amplicons.," *Genome Res.*, vol. 21, no. 3, pp. 494–504, Mar. 2011, doi: 10.1101/gr.112730.110.

[66] Q. Wang, G. M. Garrity, J. M. Tiedje, and J. R. Cole, "Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy.," *Appl. Environ. Microbiol.*, vol. 73, no. 16, pp. 5261–5267, Aug. 2007, doi: 10.1128/AEM.00062-07.

[67] J. Barroso-batista, M. F. Pedro, J. Sales-dias, and C. J. G. Pinto, "Specific eco-evolutionary contexts in the mouse gut reveal Escherichia coli metabolic versatility."

      Figure Legends

Fig. 1 | ARO112 genomic comparison with other *Enterobacteriaceae* strains from the *Klebsiella* and
***Escherichia* genera for their pathogenic potential. a,** Phylogenetic tree based on whole-genome analysis of
15 strains (5 *E. coli* strains, 5 *K. pneumoniae* strains, and 5 non-*pneumoniae Klebsiella* strains). Strains in bold
were tested experimentally; detailed description of the different the strains in Supplementary Table 1. **b,**
Heatmap, **c,** Principal Coordinate Analysis representing presence/absence of pathogenic properties and **d,**
Principal Coordinate Analysis of the different strains using the abundance of pathogenic properties. **c-d,**
ellipses represent 90% confidence.

Fig. 2 | Phenotypical evaluation of ARO112 and other *Enterobacteriaceae* strains for nosocomial-
related pathogenic factors important for host invasion and colonization. a, Schematic representation of the
pipeline used to test bacterial strains for pathogenic traits. A set of 9 strains (4 *E. coli*, 3 *K. pneumoniae*, 2
non-*pneumoniae Klebsiella*) were tested, both in rich and minimal media, for: **b,** antibiotic resistance, MDR
Enterobacteriaceae strains are represented in bold (resistant to >3 classes of antibiotics); **c,** biofilm
formation; **d,** urease activity; **e,** siderophore production; and **f,** capacity to resist killing or inhibition by human
serum. Selected strains (ARO112, EcN, and Kp1012) were tested for their capacity to **g**, acquire the
conjugative plasmid RP4 from MG1655 donor strain or Kgri donor strain, or **h**, retain the plasmid in rich and
minimal media. **i**, The capacity to retain the non-conjugative plasmid pMP7605 in rich and minimal media
was also tested. In **c-i**, bars represent median values. **b,** includes data from a total of 6 replicates per group,
from 2 independent experiments; **c**, from a total of 15 replicates per group, from 3 independent experiments;
d, from a total of 6 replicates per group, from 2 independent experiments; **e,** more than 8 replicates per
group, from 3 independent experiments; **f,** from a total of 6 replicates per group, from 2 independent
experiments; **g,** from a total of 15 (from MG1655 RP4) or 10 (From Kgri RP4) replicates per group, from 2-4
independent experiments; **h,** from a total of 6 replicates per group, from 2 independent experiments; **i,** a
total of 7 replicates per group, from 2 independent experiments. In **c-g,** statistical analysis was done using
Kruskal-Wallis' test with Dunn's correction for multiple comparisons. In **e-f,** different letters denote
significant differences among strains within the same medium (p<0.05) and # denotes significant differences
between different media for the same strain (p<0.05). In **h** and **i**, data were analyzed using Two-way ANOVA
with Sidak's correction for multiple comparisons. * p<0.05; ** p<0.01; **** p<0.0001.

Fig. 3 | Phenotypic testing of ARO112, EcN, and Kp1012 isolated from fecal samples of mono-
colonized mice. a, Schematic representation of the pipeline used to test bacterial strains for pathogenic traits
after gut colonization. Germ-free (GF) mice were mono-colonized with each of selected strains (ARO112, EcN,
Kp1012) for five days, after which bacteria were isolated from the fecal samples and tested for **b,** urease
activity and **c,** resistance to human serum. **d,** Bacteria were also evaluated for plasmid retention upon
intestinal colonization. **e,** Fecal supernatants were evaluated for Lipocalin-2/NGAL levels, as a marker for
intestinal inflammation. In **b**-**e**, data were analyzed using Kruskal-Wallis' test with Dunn's correction for
multiple comparisons (* $p<0.05$; ** $p<0.01$; **** $p<0.0001$). Urease activity and human serum resistance
from gut isolated bacteria were tested in a total of 10 (ARO112), 8 (EcN), or 4-7 (Kp1012) replicates, from at
least 2-3 independent experiments. Lcn2 was tested in germ-free animals before colonization (GF, N= 24),
and after colonization with ARO112 (N=10), EcN (N=9), or Kp1012 (N=15), from at least 3 independent
experiments. Non-conjugative plasmid retention in gut-isolated bacteria was tested in a total of 10 (ARO112),
9 (EcN), or 7 (Kp1012) replicates, from at least 3 independent experiments. In **b**-**e**, bars and lines represent
median values.

Fig. 4 | Evaluation of therapeutical potential capacities of ARO112 and EcN against AIEC
colonization in an IBD mouse model previously treated with antibiotics. a, Experimental setup of IBD mouse
model Nod2$^{-/-}$ mice treated with antibiotics, followed by AIEC infection and treated with ARO112, EcN or PBS
(Ctrl). **b,** Percentage of mice colonized with AIEC in the different groups throughout the experiment. **c,** AIEC
loads during the experiment. Thin lines represent AIEC loads in individual mice over time, while thick lines
represent median values for each treatment over time. In **b** and **c**, data were analyzed using Two-way ANOVA
with Tukey's correction for multiple comparisons (* $p<0.05$; ** $p<0.01$; **** $p<0.0001$). AIEC fecal loads and
colonization were tested in a total of 15 (Ctrl), 12 (+ARO112), or 10 (+EcN) animals, from 2-3 independent
experiments.

Fig. 5 | Intestinal microbiota composition resultant from treatment with probiotic strains (ARO112,
EcN) or lack thereof (Ctrl). a, Richness and **b,** diversity of gut microbiota in fecal samples collected before
(pre-AB, day -4), or after (post-AB, day 0) antibiotic treatment and 5 and 20 days after AIEC infections of mice
untreated (Ctrl) or treated with ARO112 or EcN. In **a** and **b**, lines represent median values for each treatment
over time. **c,** Changes in the relative abundances of most prevalent taxa at day 20 after AIEC infection in
comparison with the levels before antibiotics (day -4). **d,** Changes in relative abundances of potential butyrate
producers at day 5 after AIEC infection that were depleted by antibiotics, but recovered during experiment.
In **c** and **d**, different colors display severely depleted (<-5), depleted (<-1), unchanged or recovered (-1 to 1),

enriched (>1) and severely enriched (>5) $\log_2$ fold changes to pre-AB levels of each mouse. In **a-d**, data were analyzed using Two-way ANOVA with Dunnett's correction for multiple comparisons (* $p<0.1$; ** $p<0.01$; *** $p<0.001$; **** $p<0.0001$). Fecal microbiota composition analyses were performed in a total of 14 (Ctrl), 10 (+ARO112), or 4-10 (+EcN) samples, from 2-4 independent experiments.

Fig. 6 | SCFA levels and intestinal inflammation profiles of mice upon different probiotic treatments (ARO112, EcN) or lack thereof (Ctrl). a, Metabolomic profile of butyrate, isobutyrate, acetate, propionate, and succinate in fecal samples of IBD mouse model Nod2$^{-/-}$ mice not treated (Ctrl) or treated with probiotics ARO112 (+ARO112) or EcN (+EcN), at the end of the experiment. **b,** Intestinal inflammation was measured via Lcn2 proxy in fecal supernatants, throughout the experiment. Thin lines represent Lcn2 levels in individual mice overtime, while thick lines represent median values for each treatment overtime. In **a**, bars represent median values and data were analyzed using Kruskal-Wallis test with Dunn's correction for multiple comparisons. In **b**, data were analyzed using Two-way ANOVA with Tukey's correction for multiple comparisons (* $p<0.05$; ** $p<0.01$; *** $p<0.001$; **** $p<0.0001$). Metabolomics and Lcn2 were tested in 12 (Ctrl), 8 (+ARO112) and 10 (+EcN) samples.

Extended Data Figures Legends

Extended Data Fig. 1 | Quantification and comparison of genome-encoded gene products and predicted virulence factors' properties. Genome analyses were performed in PATRIC BV-BRC software, from where the number of hits for the different virulence factors databases were obtained. **a,** Venn diagram of total gene products, shared by every member of each clade and Venn diagram of virulence properties' hits shared by every member of each clade. **b,** Number of gene products encoded on the genome of each strain related to each of 8 clinically-relevant categories, per clade. In **c**, data were analyzed using Two-way ANOVA with Tukey's correction for multiple comparisons (* $p<0.05$; ** $p<0.01$; **c**).

Extended Data Fig. 2 | Comparison and correlations of phenotypically tested virulence traits. a, Bacterial growth in rich and minimal media. **b,** ARO112 and EcN were tested for their capacity to retain a natural antibiotic-resistant plasmid acquired by conjugation from the clinical isolate Kp834. In **a** and **b**, bars represent median values. In **b**, data were analyzed using Two-way ANOVA test with Sidak's correction for multiple comparisons (**$p<0.01$; **** $p<0.0001$). Growth was tested in 6-11 replicates, from 2 independent

experiments. Natural conjugative plasmid retention was tested in a total of 5 replicates per group, from 2
independent experiments.

Extended Data Fig. 3 | Siderophore production in mono-colonized mice and resistance to Lcn2. a,
Siderophores were measured in fecal supernatants collected from mice mono-colonized with each of
selected strains (ARO112, EcN, Kp1012) for 5 days. Bars represent median values. **b,** Selected strains
(ARO112, EcN, Kp1012) were tested for their capacity to grow in the presence of increased concentrations of
Lcn2 in rich or minimal media. Growth capacity was categorized as resistant (>80%), intermediate (>40%), or
sensitive (<40%) when compared to the absence of Lcn2. In **a**, siderophore production in fecal samples was
tested in a total of 10 (ARO112), 9 (EcN), or 7 (Kp1012) samples, from at least 3 independent experiments. In
b, Lcn2 resistance was tested in a total of 6 replicates, from 2 independent experiments.

Extended Data Fig. 4 | Probiotic colonization and evaluation of protective capacities of ARO112
and EcN against different invading bacteria in different mouse models. a, Loads of AIEC in fecal samples of
WT mice treated with vancomycin and gentamicin and treated with PBS (Ctrl) or ARO112 probiotic
(+ARO112). **b-c,** data from experiment shown in Fig. 4, where **b,** shows the loads of ARO112 or EcN in fecal
samples and **c,** the percentage of mice colonized by ARO112 or EcN during (13 timepoints tested) the
experiment in Nod2$^{-/-}$ mice. **d,** Loads of AIEC in fecal samples of WT mice treated with streptomycin and
treated with ARO112 (+ARO112) or EcN (+EcN) probiotics. **e,** Loads of AIEC in germ-free WT mice treated
with PBS (Ctrl) or ARO112 probiotic (+ARO112). Loads of **f,** Ec1898, **g,** *V. cholerae***,** and **h,** Kp1012 after 4 days
of mono-colonization (pre-treat) or after 8 days of treatment with ARO112 (+ARO112), in germ-free WT mice.
i, Loads of AIEC after 24h of growth without or with ARO112 or EcN, in rich or minimal media. AIEC loads in
WT SPF models were tested in 3 replicates per group (**a**) or 7 replicates per group (**d**), from 1 or 2 independent
experiments, respectively. Probiotic loads were tested in 12 (+ARO112) or 10 (+EcN) replicates from 2-3
independent experiments (**b**). AIEC loads in WT GF model were tested in 3 (Ctrl) and 9 (+ARO112) replicates
in 1-2 independent experiments (**e**). Ec1898 loads were assessed in 5 replicates from 2 independent
experiments (**f**). *V. cholerae* loads were assessed from 6 replicates from 2 independent experiments (**g**).
Kp1012 loads were assessed in 8 replicates from 2 independent experiments (**h**). In vitro growths were tested
in 10 (Ctrl), 7 (+ARO112), or 3 (+EcN) replicates in minimal medium, and 9 (Ctrl), 19 (+ARO112), or 19 (+EcN)
replicates in rich medium, from 1-4 independent experiments (**i**), and two-way ANOVA was tested with
Tukey's multiple comparison correction (* $p<0.05$; **** $p<0.0001$). In f-h, data were analyzed using Mann-
Whitney tests (* $p<0.05$; *** $p<0.001$). In **b**, thin lines represent ARO112 and EcN loads in individual mice
over time, while in **a**, **b**, **d**, **e**, thick lines and bars represent median values.

Extended Data Fig. 5 | Comparison between treatments of different microbiota-related parameters in experiment in Nod2$^{-/-}$ mice. a, Principal Coordinate Analysis displaying clustering of samples before antibiotic treatment (pre-AB; 33 samples), after antibiotic treatment (post-AB; 34 samples), and after treatment with PBS (Ctrl; 14 samples) or probiotics (+ARO112 or +EcN; 10 samples per group), using microbiota composition or **b,** selective four potential butyrate producers (*Dorea*, *Oscillibacter*, unc_*Runimococcacea*, unc_*Lachnospiraceae*). Ellipses in **a** and **b** represent 90% confidence intervals. **c,** Relative abundance of Pseudomonadota taxon enriched after antibiotic treatment. Values were tested from 14 (Ctrl), 11 (+ARO112) or 4-10 (+EcN) mice. In **c**, lines represent median values and data were analyzed using Two-way ANOVA with Tukey's correction for multiple comparisons.

Material and Methods

Bacterial strains, plasmids and culture conditions

See Supplementary Table 1 for all species, strains, and plasmids. Unless otherwise specified bacteria
were cultured in Lysogenic Broth (LB, here refereed as rich medium) or M9 minimal medium (47.74mM
NaHPO$_4$-7H$_2$O, 22.04mM KH$_2$PO$_4$, 8.554mM NaCl, 18.694mM NH$_4$Cl, 2mM MgSO$_4$, 0.1mM CaCl$_2$, 1mM
thiamine) with 0.5% glucose media at 37ºC with aeration (shaking at 240 rpm).

Whole genome sequencing

One *Klebsiella oxytoca* MBC022 clone was isolated from the feces of a WT specific pathogen-free
(SPF) C57BL/6J mice bred in the animal house facility at Instituto Gulbenkian Ciência and grown in LB at 37
°C with agitation for subsequent DNA isolation using a previously described protocol [56]. A DNA library was
constructed and sequenced at the IGC Genomics facility. Paired-end sequencing of each sample was
performed using an Illumina MiSeq Benchtop Sequencer (Illumina), which produced datasets of 250 bp read
pairs. *K. oxytoca* genome has been deposited in GenBank (see Data availability).

Ec1898, Kp1012, and Kp834 were isolated from leukemia patients from Hospital La Fe in Valencia
(Spain). Genome sequence was obtained as described above in the FISABIO Sequencing facility. Sequences
have been deposited in GenBank.

Method for phylogenetic tree

Phylogenetic tree was performed using the whole genomes of strains uploaded to the PATRIC BV-
BRC browser software (v3.29.20). Parameters used included 100 single-copy genes, using Multiple Alignment
using Fast Fourier (MAFFT) Transform alignment program and RAxML Fast Bootstrapping branch support
method (v8.2.11).

Genome-encoded predicted pathogenic properties

Genomes of all strains were uploaded to the PATRIC BV-BRC [28] browser software (v3.29.20) and
analyzed. For each genome, a table containing all predicted pathogenic properties from 9 available
databases: 3 for Virulence Factors (Victors, PATRIC_VF, VFDB), 3 for Drug Targets and Transporters
(DrugBank, TCDB, TTD), and 3 for antibiotic resistance (PATRIC, CARD, NDARO). The ensemble of these data
was downloaded and two merged tables were compiled with the information for all the strains: one
displaying the presence or absence of each pathogenic property in each strain, and another including the
abundance of each pathogenic property in each strain. Principal Coordinates analyses were performed using
either presence/absence of pathogenic properties (Bray-Curtis dissimilarity index for binary/categorical data;
one-way PERMANOVA with Bonferroni correction) or abundance of said pathogenic properties (Euclidean
distances for continuous data; one-way PERMANOVA with Bonferroni correction).

 Antibiotic profile

Cultures were grown in either rich or minimal medium for 24h at 37ºC with shaking. Cultures were
washed once in sterile PBS and adjusted to a final OD_{600} of 0.05 in 150µl/well in the appropriate medium with
different concentrations of antibiotics (as specified). These cultures were then grown for 24h in a 96-well
plate in a plate shaker, after which OD_{600} measurements were performed in a plate reader Multiskan Sky.
Strains were considered resistant if in the presence of antibiotic could reach at least 80% of the growth
obtained in the absence of antibiotic. If growth in the presence of antibiotic was more than 40% and less
than 80% of the control growth, the strain was considered intermediate for resistance, and sensitive if grown
less than 40% of control growth. Resistance or intermediate resistance to one antibiotic of any class is
considered resistance or intermediate resistance to that class of antibiotics. Sensitivity to a class of antibiotics
means that bacteria were sensitive to all antibiotics tested of that class. A strain was considered multidrug
resistant (MDR) if it was non-susceptible (sensitive or intermediate) to at least 1 antimicrobial agent in 3 or
more antimicrobial classes, in both tested media, as defined before [55].

Class	Antibiotic	Concentration (µg/ml)
β-lactam	ampicillin	100
	carbenicillin	50
Aminoglycosides	streptomycin	100
	kanamycin	50
	gentamicin	15
	neomycin	200
	apramycin	50
	spectinomycin	50
Chloramphenicol	chloramphenicol	25
Macrolides	erythromycin	125
Rifamycins	rifampicin	20
Tetracyclines	tetracycline	10
Fluoroquinolones	ciprofloxacin	0.5
	trimethoprim	4

Naphthyridone	nalidixic acid	32
Cephalosporins	cefotoxidine	8

<u>Biofilm formation</u>

Biofilm formation assays were performed by adapting a previously published method [57], [58]. In
summary, bacterial cultures were grown for 24h in either rich or minimal media at 37ºC with shacking. The
following day, each culture was washed once in sterile PBS, and final OD_{600} adjusted to 0.05 in the appropriate
medium and 200 µl of culture were dispensed per well in 48-well plates. Plates were centrifuged for 5 min at
4000 rpm at room temperature, after which plates were incubated at 37ºC for 90 min for adhesion. After
adhesion, supernatants were carefully removed and discarded, wells were washed once with 200 µl of sterile
PBS, and 500 µl of sterile fresh appropriate medium were carefully dispensed per well. Plates were incubated
at 37ºC for 24h, after which attached biofilms (biomass) was quantified with Crystal Violet (CV). For that
supernatants were removed and discarded and well bottoms were carefully washed with 500 µl of sterile
PBS. Each well was carefully stained with 0.1% CV solution and plates were incubated at room temperature
for 20 min protected from light. After incubation, each well was washed with PBS and plates were incubated
open and inverted in a paper towel for 30 min protected from light, to dry. Each well was then de-stained
with 200µl of a 33% glacial acetic acid solution and incubated for 15 min at room temperature protected
from light. Supernatants were removed from each well into a new plate and read for OD_{580} in a Multiskan Sky
plate reader (peak for CV staining), to quantify biofilm biomass.

<u>Urease activity</u>

Cultures were grown in either rich or minimal medium for 24h at 37ºC with shaking. Cultures were
washed once in sterile PBS and adjusted to an OD_{600} of 1, after which each culture was diluted 20x in urea
medium (3g/L L-tryptophan, 5 g/L NaCl, 1g/L H_2KO_4P, 1g/L $H_2K_2O_4P$, 20g/L urea, 0,012g/L phenol red) and
incubated for 24h at 37ºC with shaking. Urea degradation leads to the medium color to change from
yellow/orange to pink/red. 100 µl of each culture were then transferred into a 96-wellplate and measured
at OD_{560} in a Multiskan Sky plate reader to quantify urease activity. For urease activity measurement of
bacteria extracted from fecal samples of monocolonized mice, cell suspensions of 2.5×10^7 cells/ml were used,
instead of the grown culture.

<u>Siderophore production</u>

Cultures were grown in either rich or minimal medium for 24h at 37ºC with shaking. Cultures or fecal
samples diluted in PBS were centrifuged to recover supernatants. Culture and fecal supernatants were
filtered in 0.22 µm filters. Cell-free supernatants (100 µl) were mixed 1:1 (v:v) with Cas-PIPES solution (100

μl) [59], [60] in a 96-well plate and incubated at room temperature for 20 min, after which OD_{630} was

measured in Multiskan Sky plate reader to calculate percentage of siderophore production, using sterile

media or PBS as reference.

1017
$$\% \ siderophores = \frac{(ref - sample) * 100}{ref}$$

<u>Resistance to pooled human serum</u>

Bacterial susceptibility to human serum was tested as described [40] with modifications. Bacterial

strains were grown in either rich or minimal medium for 24h at 37ºC with shaking, or extracted directly from

fecal pellets. Cultures were washed once in sterile PBS and adjusted to 2×10^7 cells/ml, after which 25 μl of

cell suspension in PBS were added to 75 μl of pooled human serum solution (P30-2401, PAN-Biotech).

Cultures were plated in LB agar plates for CFUs at 0h and after 3h of static incubation at 37ºC, to assess killing,

resistance, or proliferation of bacterial cells in the presence of human serum.

<u>Plasmid conjugation</u>

Cultures were grown in rich medium for 24h at 37ºC with shaking. Cultures were washed once in

sterile PBS. Donor (MG1655 RP4/RK2, Kgri RP4/RK2) and recipient (ARO112, EcN, Kp1012) strains were

thoroughly mixed at a 1:1 ratio (total of 10^8 cells), centrifuged for 30 sec at 14,000 rpm and resuspended in

20 μl of PBS. Two drops of 10 μl each were placed on an agar plate, air-dried, and incubated at 37ºC for 24h.

After incubation, both drops were removed with a sterile loop and resuspended in 1 ml of sterile PBS. Serial

dilutions were performed and plated to count CFUs for recipient strain with (selective plating with plasmid-

borne antibiotic resistance) and without (non-selective or selective plating for recipient's antibiotic resistance

plating) tested plasmid. Conjugative plasmid (RP4) harbors resistance to tetracycline (10 μg/ml) and

kanamycin (50 μg/ml).

<u>Plasmid retention</u>

Strains harboring tested plasmids (RP4 acquired through conjugation from MG1655 RP4 donor strain;

pMP7605 acquired through electroporation; Kanamycin resistance-bearing natural plasmid acquired through

conjugation from Kp824) were grown in either rich or minimal medium with antibiotics (for plasmid-borne

antibiotic resistance; kanamycin (50 μg/ml) and tetracycline (10 μg/ml) for RP4; gentamicin (30 μg/ml) for

pMP7605; kanamycin (50 μg/ml) for natural plasmid) for 24h at 37ºC with shaking. Cultures were washed

once in sterile PBS and adjusted to a final OD_{600} of 0.05 in 150 μl in the appropriate medium without

antibiotics, and serial dilutions were plated in selective and non-selective LB agar plates, to assess CFUs with

(plasmid retention) and without (plasmid loss) antibiotic resistance. Cultures were incubated at 37ºC with

shaking for 24h. Every 24h, cultures were diluted 1:100 in fresh appropriate medium and re-incubated. After

5 days of passages, serial dilutions of cultures were plated as mentioned before.

Animal experimentation

All of the mouse experiments performed at Instituto Gulbenkian Ciência (IGC) were approved by the

Institutional Ethics Committee and the Portuguese National Entity (Direção Geral de Alimentação e

Veterinária; 015190), which complies with European Directive 86/609/EEC of the European Council.

C57BL/6J male mice, either WT or Nod2$^{-/-}$, were used at 6–8 weeks of age and were randomly

assigned to experimental and control groups. C57BL/6J mice were used for all experiments. None of the

animal experiments were performed blinded. Sample size was chosen according to institutional directives

and in accordance with the guiding principles underpinning humane use of animals in research. No statistical

analyses were performed to predetermine the sample sizes. All of the experiments were performed at least

twice, except when stated otherwise.

Fecal bacterial extraction

WT C57BL/6J germ-free mice were orally gavaged with 100 μL containing ~10^8 CFUs of ARO112, EcN,

or Kp1012 strains bearing the non-conjugative plasmid pMP7605. Five days later, fecal pellets were collected

for live gut bacteria extraction, as previously described [41]. Briefly, pellets were weighed and 500 μL of

sterile PBS were added before mechanical disruption with a motorized pellet pestle, after which another 500

μL of sterile PBS were added. Samples were subjected to four iterations of vortex mixing for 15 sec,

centrifugation at 1,000 rpm for 30 sec at room temperature and recovery of 750 μL of the debris-free cell-

containing supernatants into a new tube, replacing that volume with sterile PBS before the next iteration.

After recovery of 3 ml per sample, isolated cells were centrifuged at 4,000 g for 5 min at room temperature,

discarding the supernatant and resuspending the cells fractions in 1 ml of sterile PBS supplemented with

glycerol and cysteine at final concentrations of 20% and 0.1%, respectively. Samples were frozen and stored

at -80ºC, to measure the phenotypes listed below.

Nod2$^{-/-}$ and WT SPF mice experiments

C57BL/6J Nod2$^{-/-}$ and WT mice bred under specific pathogen-free conditions in the animal house

facility at the Instituto Gulbenkian Ciência were kept individually in sterile ISOcages (Tecniplast) at our specific

opportunistic pathogen-free facility. Mice were orally gavaged with gentamicin (3 mg/kg/d) and vancomycin

(40 mg/kg/d) once daily for three consecutive days [46]. Two days after the last gavage with antibiotics, mice

were orally gavaged with 100μL containing ~10^8 CFUs of AIEC strain (LF82-strepR), and the following day were

orally gavaged either with 100 μL of PBS (control), or 100 μL containing ~10^8 CFUs of ARO112 or EcN strains.

Fecal samples were collected at the timepoints indicated in the corresponding graph and plated in selective

medium to assess colonization levels of AIEC. Additionally, fecal samples were also used to measure levels of Lcn2 (Lipocalin-2/NGAL ELISA) and to assert the intestinal microbiota composition.

C57BL/6J WT mice bred under SPF conditions in the animal house facility at the Instituto Gulbenkian Ciência were kept individually in ventilated cages with high-efficiency particulate air filters in our animal facility. Streptomycin (5 g/l) was maintained ad libitum in the drinking water for 15 days, being replaced every 3 days, after which mice were kept in non-supplemented water. Four days after halting streptomycin treatment, AIEC strain (LF82-strepR) was orally gavaged (~10^8 CFUs/mouse in 100 µL). Three days after that, either ARO112 strain or EcN strain were orally gavaged (~10^8 CFUs/mouse in 100 µL). Fecal samples were collected at the timepoints indicated in the corresponding graph and plated in selective medium to assess colonization levels of AIEC.

WT gnotobiotic mice experiments

C57BL/6J germ-free mice were bred and raised in IGC's Gnotobiology Unit in axenic isolators (La Calhene/ORM) and were later transferred into sterile ISOcages (Tecniplast). Animals were individually housed and orally gavaged with 100 µL containing ~10^8 CFUs of AIEC (LF82-strepR), Kp1012, *V. cholerae*, or Ec1898 strains. The following day, mice were gavaged with 100 µL containing ~10^8 CFUs of either ARO112 or EcN strains, or 100µl of PBS. Fecal samples were collected at the timepoints indicated in the corresponding graphs and plated in selective media to assess colonization levels of AIEC, Kp1012, *V. cholerae*, or EC1898 strains.

16S rRNA sequencing

All fecal samples from SPF mice experiments stored at −80 °C were processed for DNA extraction as previously described [61]. In brief, DNA was extracted using a combination of the QIAamp Fast DNA Stool Mini Kit (Qiagen) according to the manufacturer's instructions, mechanical disruption using a motorized pellet pestle and elution to a final volume of 100 µL in ATE buffer. For each sample, the 16S rRNA gene was amplified using the 515F/806R (V4 regions) primer pairs recommended by the Earth Microbiome Project under the following PCR cycling conditions: 94 °C for 3 min; 35 cycles of 94°C for 60s, 50°C for 60s and 72°C for 105s; and extension at 72°C for 10min [62], [63]. After library preparation, 2 × 250 bp sequencing was performed at the IGC Genomics Unit using an Illumina MiSeq Benchtop Sequencer.

Fecal microbiota analysis

Mothur v.1.32.1 was used to process sequences as previously described [61], with some modifications. Sequences were converted to FASTA format. Sequences that were shorter than 220 bp containing homopolymers of longer than 8 bp or undetermined bases with no exact match with the forward and reverse primers, and barcodes that did not complement each other or that did not align with the

appropriate 16S rRNA variable region, were not included in the analysis. A quality score above 30 (range, 0 to 40, where 0 represents an ambiguous base) was used to process sequences, which were trimmed using a sliding-window technique over a 50 bp window. Sequences were trimmed from the 3' end until the quality score criterion was met, and were merged after that. Between 20,000 and 50,000 sequences were obtained per sample. 16S rRNA gene sequences were aligned using SILVA template reference alignment [64]. ChimeraSlayer [65] was used to remove potential chimeric sequences. Sequences with distance-based similarity of greater than 97% were joined into the same OTU using the average-neighbour algorithm. All of the samples were rarefied to the same number of sequences (10,000) for diversity analyses. Samples below 10,000 reads were removed from the analysis. A Bayesian classifier algorithm with a 60% bootstrap cut-off was used for each sequence [66]; sequences were assigned to the genus level where possible or otherwise to the closest genus-level classification. All OTUs were used, and taxa plotted are the result of the merge of OTUs with the same classification, being collapsed to 37 different taxa, with lower represented OTUs/taxa being summed and plotted as "other". Principal Coordinate Analyses were performed using the 37 taxa or a subset of 4 taxa of potential butyrate producers (*Dorea*, *Lachnospiraceae*, *Oscillibacter*, *Ruminococcaceae*), representing the Bray-Curtis dissimilarity index (one-way PERMANOVA with Bonferroni correction).

SCFA quantification

We performed 1H-NMR (proton nuclear magnetic resonance) analysis on fecal contents to determine the abundance of SCFA in the mouse gut of Nod2$^{-/-}$ treated with ARO112 and EcN, or untreated (control), as previously described [67]. Fecal samples (1 to 3 fecal pellets) were diluted in 1mL of PBS. To remove large debris, the samples were pelleted by centrifugation at 18000 g for 15 min at 4°C. Supernatant was collected and filtered through a 0.22 mm filter (Milipore). Next, samples were filtered with 3 KDa filters (Vivaspin 500) by centrifugation at 15000 g and 4°C, for 3 h (or until 150 μL of filtrate was obtained). Filtered samples were stored at −80°C until spectrum acquisition. For spectrum acquisition, samples were thawed at room temperature and 150 μL mixed with 60 μL of 350 mM phosphate buffer (pH 7.09 with 2% NaN3, 10 mL of a 0.05% (w/v) 3-(Trimethylsilyl)propionic-2,2,3,3-d4 (TSP-d4, Sigma-Aldrich) solution, and 380 μL of deuterated water - D$_2$O) to a total volume of 600 μL. The mixture was transferred to a 5 mm glass NMR tube. All solutions were prepared with D$_2$O. Samples were homogenized by inversion and the spectra were acquired after pH measurement. Acquisitions were performed on a Bruker NEO 500MHz instrument equipped with QXI H-C/N/P 5 mm probe-head with z-gradients. 1H-NMR spectra were acquired using 1D NOESY pulse sequence with pre-saturation (noesypr1d) under the following conditions: 90 pulse for excitation mixing time 100 ms, acquisition time 4 s, and relaxation delay 1 s. All spectra were acquired with 200 scans at 25C, with 48018 data points and 6002 Hz (12 ppm) spectral width (Chenomx acquisition parameters). The recorded 1H-NMR spectra were phase corrected using Bruker TopSpin 4.0.7 and spectra were then analyzed using Chenomx

NMR Suite 8.1. Compounds were identified by manually fitting reference peaks to spectra in database

Chenomx 500 MHz Version 10. Quantification was based on internal standard peak integration (TSP-d4).

Graphical representation and Statistical analyses

Phylogenetic tree (Fig. 1a) was done using iTOL (itol.embl.de). Cluster heatmap (Fig. 1b) was done

using SRPlot (https://www.bioinformatics.com.cn), using bidirectional clustering, complete cluster method

and Euclidean distances. All statistical graphs and statistical analyses presented were done using Graphpad

Prism 8. Correlation tables (Extended Data Fig. 2 and 5) were drawn in Microsoft Excel for Mac, using results

obtained with Graphpad Prism 8. Principal Coordinates analyses (Fig. 1, Extended Data Fig. 5) were obtained

with Past4.02 software (https://folk.universitetetioslo.no/ohammer/past) as explained above.

1150

1151

1152

Fig. 1

Fig. 2

a
1 Culture bacterial isolates
2 Separate cells from media
3 Phenotypic assays

Rich medium   Minimal medium

Supernatant → Siderophores

Cells →
Biofilm
Urease − +
Human serum resistance →

Antibiotic resistance
Plasmid retention
Conjugation

b
Rich medium   Minimal medium

β-lactam
aminoglycosides
chloramphenicol
macrolides
rifamycins
tetracyclines
cephalosporins
fluoroquinolones

MG1655 AIEC EcN **Ec1898 Kp1012 Kp834 MH258** MBC022 ARO112

Resistant **Intermediate** **Sensitive**

c Biofilm formation
Crystal Violet; ΔOD$_{580}$

d Urease activity
OD$_{560}$

e Siderophores production
% Siderophores units

f Human serum resistance
log$_{10}$ CFUs/ml fold change to untreated
Detection limit

▼ MG1655 ⬛ AIEC ● EcN ▲ Ec1898 *E. coli* strains
⬤ Kp1012 ⬛ Kp834 △ MH258 *K. pneumoniae* strains
⬛ MBC022 ● ARO112 Non-*pneumoniae Klebsiella* strains

g Plasmid acquisition
Donor strain: MG1655 RP4
Donor strain: Kgr RP4
Population frequency (log$_{10}$)
ARO112 EcN Kp1012
**** ** * *

h Conjugative plasmid retention
Population frequency (log$_{10}$)
Rich medium Minimal medium
● ARO112 ● EcN ● Kp1012
** **** *

i Non-conjugative plasmid retention
Population frequency (log$_{10}$)
Rich medium Minimal medium
● ARO112 ● EcN ● Kp1012
**** * ** **** ** **

Fig. 3

Fig. 4

Fig. 5

a

Richness

Chao Index

····○···· Ctrl
——●—— +ARO112
— —○— — +EcN

****

****

pre-AB post-AB 5 10 15 20
Time (days post-AIEC gavage)

b

Diversity

Shannon Index

····○···· Ctrl
——●—— +ARO112
— —○— — +EcN

****

pre-AB post-AB 5 10 15 20
Time (days post-AIEC gavage)

c

Verrucomicrobia → *Akkermansia*

Pseudomonadota		post-AB	+ARO112	Ctrl	+EcN
	Escherichia/Shigella		****	****	
	Proteus		****	****	***
	Unc. *Enterobacteriaceae*		*	**	
	Parasutterella		*	**	
	Enterobacter/Klebsiella		***		
	Unc. *Alcaligenaceae*				****
	Alphaproteobacteria incertae sedis				
	Desulfovibrio			**	

Bacteroidota					
	Parabacteroides		*		
	Unc. *Prevotellaceae*		*		*
	Alistipes				****
	Unc. Bacteroidota				
	Unc. Bacteroidales				
	Bacteroides		***	*	****
	Unc. *Porphyromonadaceae*				
	Muribaculaceae				***
	Prevotella				

Bacillota					
	Unc. Clostridiales		**		****
	Lactobacillus			*	
	Unc. *Erysipelotrichaceae*		***		**
	Blautia		**	***	****
	Unc. *Clostridia*				
	Papillibacter				
	Holdemania				
	Allobaculum				
	Syntrophococcus				
	Ruminococcus				
	Butyricicoccus				
	Anaerovorax				
	Dorea		*	*	
	Unc. Bacillota				
	Acetanaerobacterium			*	
	Turicibacter				*
	Oscillibacter		***		
	Unc. *Lachnospiraceae*		****	****	
	Unc. *Ruminococcaceae*		****	****	
	Other				

0.054   **
**

d

	post-AB	+ARO112	Ctrl	+EcN
Alistipes				
Dorea				
Oscillibacter		*		
Unc. *Lachnospiraceae*		****	****	
Unc. *Ruminococcaceae*		***	**	

Legend:

Potential butyrate producers

■ Severely enriched
■ Enriched
□ Unchanged / Recovered
■ Depleted
■ Severely depleted

Fig. 6

a

b

Extended Data Fig. 1

a

Extended Data Fig. 2

a

Bacterial growth

b

Donor strain:
Kp834 natural plasmid
Conjugative plasmid retention

Extended Data Fig. 3

a

Siderophore production

b   Lipocalin-2 resistance

Extended Data Fig. 5

a

Microbiota (37 taxa)

b

Potential butyrate producers subset (4 taxa)

c

Enriched Pseudomonadota