Dietary restriction and life-history trade-offs: insights into mTOR pathway regulation and reproductive investment in birds

Gebrehaweria Kidane Reda[1,2,3] *, Sawadi Fransisco Ndunguru[1,2,3], Brigitta Csernus[3], Gabriella Gulyás[1], Renáta Knop[1], Csaba Szabó[4], Levente Czeglédi[1], Ádám Z. Lendvai[3*]

[1]*Department of Animal Science, Institute of Animal Science, Biotechnology and Nature Conservation, Faculty of Agricultural and Food Sciences and Environmental Management, University of Debrecen, 4032 Debrecen, Hungary*
[2]*Doctoral School of Animal Science, University of Debrecen, 4032 Debrecen, Hungary*
[3]*Department of Evolutionary Zoology and Human Biology, University of Debrecen, 4032 Debrecen, Hungary*
[4]*Department of Animal Nutrition and Physiology, Faculty of Agriculture and Food Sciences and Environmental Management, University of Debrecen, 4032 Debrecen, Hungary*

* Corresponding authors: email: gebrek2000@gmail.com, AZ.Lendvai@gmail.com

Abstract

Resources are needed for growth, reproduction and survival, and organisms must trade-off limited

resources among competing processes. Nutritional availability in organisms is sensed and

monitored by nutrient-sensing pathways that can trigger physiological changes or alter gene

expression. Previous studies have proposed that one such signalling pathway, the mechanistic

target of rapamycin (mTOR), underpins a form of adaptive plasticity when individuals encounter

constraints in their energy budget. Despite the fundamental importance of this process in

evolutionary biology, how nutritional limitation is regulated through the expression of genes

governing this pathway and its consequential effects on fitness remains understudied, particularly

in birds. We used dietary restriction to simulate resource depletion and examined its effects on

body mass, reproduction and gene expression in Japanese quails (*Coturnix japonica*). Quails were

subjected to *ad libitum* (ADL) feeding or 20%, 30%, and 40% restriction levels for two weeks. All

restricted groups exhibited reduced body mass, whereas reductions in the number and mass of eggs

were observed only under more severe restrictions. dietary restriction led to decreased expression

of *mTOR* and insulin-like growth factor 1 (*IGF1*), whereas the ribosomal protein S6 kinase 1

(*RPS6K1*) and autophagy-related genes (*ATG9A* and *ATG5*) were upregulated. The pattern in

which mTOR respond to restriction was similar to what has been seen in body mass. Regardless

of the treatment, proportionally higher reproductive investment was associated with individual

variation in *mTOR* expression. These findings reveal the connection between dietary intake and

the expression of *mTOR* and related genes in this pathway.

Keywords: dietary restriction, gene expression, mTOR pathway, reproduction, trade-off,

Coturnix japonica

1. INTRODUCTION

Resource availability is a key driver of resource allocation decisions which define life-history trade-offs (Ng'oma et al., 2017; Zera and Harshman, 2001). When food is abundant, animals allocate resources towards current reproduction against somatic maintenance and further reproduction. The bias for reproductive investment may be accompanied by physiological costs, including oxidative stress and a reduced immune potential (Chang van Oordt et al., 2022; Metcalfe and Monaghan, 2013), which in turn affects future reproduction performance and the health span of the organism (Hassan et al., 2003; Mahrose et al., 2022). This phenomenon changes under limited resources when organisms must divert energy from reproduction to somatic maintenance (Carlsson et al., 2021; Flatt et al., 2013).

Dietary restriction (DR) is an intervention that mimics the depletion of resources, which regulates life-history traits almost uniformly in model organisms ranging from yeast to humans (Colman et al., 2014; Inness and Metcalfe, 2008; Simons et al., 2013). Reducing calorie intake affects the investment into self-maintenance, growth and reproduction antagonistically (McCracken et al., 2020; Regan et al., 2020). The classical resource allocation theory predicts that DR should lead to a linear decrease in reproduction in favour of self-maintenance (Shanley and Kirkwood, 2000). However, modest DR can maintain or even improve reproductive performance by activating cell recycling mechanisms such as apoptosis and autophagy (Adler and Bonduriansky, 2014; Mahrose et al., 2022). When the level of DR becomes more severe, the organism must shift energy away to meet basal energetic requirements; thus, the reproduction rate will decline (Moatt et al., 2016; Ottinger et al., 2005; Shanley and Kirkwood, 2000).

Resource availability at the organismal level is monitored by the neuroendocrine system that dynamically responds to internal signals via changes in physiology or gene expression (Maruska et al., 2018). The key mediators of this process are the nutrient-sensing pathways governed by insulin-like growth factor-1 (IGF-1) and mechanistic target of rapamycin (mTOR) (Johnson, 2018; Kapahi et al., 2017). The IGF-1/mTOR signalling pathway is activated by high nutrient availability and triggers growth and reproduction while downregulating cellular processes that maintain organismal and cellular homeostasis (e.g. apoptosis and autophagy) (Montoya et al., 2022; Papadopoli et al., 2019). In response to growth hormone and energy availability, IGF-1 is released into the bloodstream and binds to its membrane receptor (IGF-1R), which activates further

molecular components (PI3K and Akt) that will, in turn, trigger mTOR activation (Feng and
Levine, 2010).

mTOR serves as the central regulator of the nutrient-sensing pathway, integrating
intracellular nutrient availability and extracellular signals to govern essential cellular processes,
including metabolism, growth, proliferation, and survival, thereby influencing tissue and organ
growth (Papadopoli et al., 2019; Rabanal-Ruiz and Korolchuk, 2018). The mTOR exists in two
distinct complexes known as mTORC1 and mTORC2, each with different functions and regulatory
mechanisms. mTORC1 is primarily involved in regulating cell growth and protein synthesis in
response to various environmental cues such as nutrient availability, energy status, and growth
factors. On the other hand, mTORC2 has diverse roles in cell survival, cytoskeletal organization,
and metabolism (Szwed et al., 2021).

DR has been shown to downregulate the mTORC1 pathway through various mechanisms,
primarily involving its upstream effectors, such as IGF-1 and intracellular amino acid deprivation
(Sancak et al., 2010; Speakman and Mitchell, 2011). Furthermore, DR has a complex impact on
mTORC2 activity. On the one hand, it has been observed that DR downregulates mTORC2 by
inhibiting insulin/Akt signalling (Saxton and Sabatini, 2017). On the other hand, there is evidence
to suggest that DR can upregulate mTORC2 through the activation of adenosine monophosphate-
activated protein kinase (AMPK) (Fu and Hall, 2020).

The activity of mTOR and its downstream effectors is further regulated through
transcriptional regulation/mRNA expression and post-translational modifications (Deng et al.,
2014; Mierziak et al., 2021; Rollins et al., 2019). While post-translational modifications and
specific amino acid availability induce mTOR activation, general resource availability is
responsible for adaptive changes in gene expression (Efeyan et al., 2015; Mierziak et al., 2021;
Sandri et al., 2013). Although the final activity of mTOR is influenced by multiple factors, higher
expression of the mTOR gene can potentially increase the pool of available mTOR protein for
activation, while lower gene expression leads to reduced protein production and availability
(Buccitelli and Selbach, 2020). Studies on model organisms have predominantly focused on the
post-translational activation of mTOR and its downstream effectors (Laplante and Sabatini, 2012;
Papadopoli et al., 2019). However, the impact of DR on differential expression of nutrient
singsing genes and how they mediat fitness remain to be fully elucidated.

The mTORC1 performs its effect on fitness traits through a number of downstream
effectors, including ribosomal protein S6 kinase 1 (RPS6K1) (Guo and Yu, 2019; Nojima et al.,
2003) and autophagy-related genes (Ma et al., 2018). Under excessive food availability, the
mTORC1/RPS6K1 pathway facilitates protein synthesis and subsequent cell growth and
proliferation (Fenton and Gout, 2011). Under DR, the mTORC1/autophagy pathway recycles cell
contents for energy substitution and reduces oxidative stress (Chung and Chung, 2019). Hence,
RPS6K1, *ATG9A* and *ATG5* are the best representative candidate genes of the mTOR downstream
pathways. Studying the expression of these genes and their relationship to fitness traits under
different DR levels is critically important to understand their role beyond post-translational
regulation.

Previous studies on DR have primarily focused on model organisms other than birds, and
except for some production-intentioned experiments on chicken (Deng et al., 2014; Hao et al.,
2021; She et al., 2019), understanding the mechanism of mTOR signalling in avian biology
remains largely unclear. Studies conducted in mammals have indicated that a higher nutritional
intake and metabolic rate upregulate the mTOR pathway, leading to a subsequent downregulation
of autophagy. This, in turn, exposes organisms to oxidative damage and cellular senescence.
However, DR without malnutrition has been found to mitigate these effects (Carroll and
Korolchuk, 2018). On the contrary, birds have higher metabolic rates, circulating glucose levels
and body temperature than mammals, while they live twice as long as size-matched mammals
(Barja, 1998). Despite the fact that a higher metabolic rate contributes to oxidative and
glycoxidative damages, birds suffer less compared to mammals at a given body size (Jimenez et
al., 2019). This metabolic paradox may be due to a different diet-fitness relationship in birds, which
can be elucidated using dietary manipulation experiments.

At this stage, we aimed to observe the effects of DR gradient on liver *mTOR* mRNA
expression, as well as its main upstream and downstream effectors, using adult female Japanese
quails (*Coturnix japonica*) as an experimental avian model system. The liver serves as a primary
site for the intricate nutrient metabolic pathway, which significantly influences the proper
functioning of the entire body. Being a major organ responsible for regulating homeostasis, the
liver plays a crucial role in nutrient regulation, protein synthesis, and detoxification processes.
Nearly all genes involved in the nutrient-sensing pathway exhibit differential expression in the

liver and display a robust correlation with the overall functioning of the body, ultimately determining fitness traits (Baloni et al., 2019; Gokarn et al., 2018).

We hypothesize that under DR, the expression of target genes involved in the mTOR pathway plays a crucial role in mediating resource availability and influencing fitness traits. We predict reduced *mTOR*, *IGF1* and *RPS6K1* expression and upregulation of autophagy-related genes. Furthermore, we anticipate that variations in *mTOR* expression will be associated with proportional differences in reproductive investment. These findings will provide insights into the relationship between gene expression, resource allocation, and fitness traits.

2. Materials and methods

2.1. Experimental animals and housing

The experiment was approved by the Ethical Committee for animal use of the University of Debrecen, Hungary (Protocol No. 5/2021/DEMAB) and followed all institutional and national regulations.

We purchased four weeks old Japanese quail (*Coturnix japonica*) chicks from a commercial quail breeder (Budai Fürjészet, Hungary) and housed them in the animal house of the Institute of Agricultural Research and Educational Farm of the University of Debrecen. The birds were kept in cages in groups of 10 for an additional four weeks (until they reached maturity) before being subjected to the experimental treatment. At the age of eight weeks, 32 female birds with similar body mass were selected and housed in individual cages (18.5 cm long × 21 cm wide × 18.5 cm high) for seven days of acclimation period on *ad libitum* (ADL) feed and water. The experimental room was maintained at a temperature of 24±3 °C and 60 - 75% relative humidity. Photoperiod was fixed at 12:12 L:D daily cycle and regulated using a LED Lighting Dimming System. The basal feed for experimental quails was formulated as a breeder quail ration (20% CP; 12.13 MJ/kg ME, (NRC (National Research Council), 1994)) based on corn, soybean, and wheat (Table S1).

2.2. Experimental design

During the acclimation period, the daily feed intake of each individual was measured for seven consecutive days. Approximately 50 g feed was weighed on a digital scale (± 0.1 g) and provided in a 200 g capacity plastic feeder each morning between 8:00-9:00 am. The following day, 24h

later, the weight of the remaining feed was weighed again, and the feeders were replenished with

fresh feed. Daily feed intake was measured as the difference between the weight of the offered and

the remaining food. Due to the design of the feeders, food spillage was negligible. The average

daily feed intake was calculated as the mean of the seven measurements. The average feed intake

during the acclimation period was 29.67 ± 3.73 g. We also measured the live body mass of each

bird at the beginning and at the end of the acclimation period to analyse mass change. At the start

of the experimental treatment, there were no significant temporal pattern of either body mass or

feed intake. Birds were regularly laying eggs during the acclimation period.

After acclimation, 32 female birds were randomly assigned to four dietary treatments. The

birds were fed with 80% (DR20), 70% (DR30), and 60% (DR40) of their average individual feed

intake, while the control group was fed ad libitum (ADL). The experiment lasted for 14 days. The

amount of feed left daily in the ADL group was measured and analysed to detect any significant

changes in temporal intake. However, no significant changes were observed.

178   2.3. Measurements and sampling

Immediately after lights-on in the morning, we removed all the feeders to maintain similar feeding

conditions (empty gut) between the ADL and restricted birds during the measurement and

sampling points. We measured body mass at the beginning of the experiment (day 0) and on days

7 and 14 of the restriction period. We collected eggs daily, measured their mass and recorded the

identity of the respective hen. Body mass and egg mass were measured using a digital scale (± 0.1

g). On day 14, all birds were euthanised, and immediately a liver sample was collected and rapidly

frozen on dry ice, and immediately stored at -80 °C until further assays.

187   2.4. RNA Extraction and cDNA Synthesis

Total RNA from liver tissue was isolated using the TRIzol reagent (Direct-zol™ RNA MiniPrep,

Zymo Research Corporation, USA) according to the manufacturer's protocol, including DNA

digestion step. The RNA concentration and purity were measured using HTX Synergy Multi-Mode

Microplate Reader spectrophotometer (Agilent BioTek, BioTek Instruments Inc, USA). RNA

integrity was checked by 1% agarose gel electrophoresis stained with ethidium bromide (see

supplementary material for more detailed protocol). Reverse transcription was performed using

the qScript cDNA synthesis kit, following the manufacturer's protocol (Quantabio Reagent

Technologies, QIAGEN Beverly Inc., USA) in PCRmax Alpha Thermal Cycler (Cole-Parmer

Ltd., Vernon Hills, IL, USA) (see supplementary material for more detailed protocol).

2.5. Real-time PCR (qPCR)

The real-time quantitative PCR was performed using EvaGreen qPCR Mix (*Solis BioDyne*,

Teaduspargi, Estonia) according to the manufacturer's protocol. Intron-spanning gene-specific

primer pairs for quails were designed using Oligo7 software and obtained from Integrated DNA

Technologies (BVBA-Leuven, Belgium) (Table S2). We checked for target identity using Primer-

Blast software of the National Centre for Biotechnology Information (NCBI)

(http://www.ncbi.nlm.nih.gov) (see Supplementary Material for more detailed protocol).

Among the most frequently used reference genes in birds, beta-actin (*ACTB*),

glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) and 18S ribosomal RNA (RN18S), we

selected the best reference gene, *ACTB*, using NormFinder, BestKeeper and deltaCt algorithms.

The $2^{-\Delta\Delta Ct}$ method was employed to analyse the relative changes in mRNA expression of target

genes (*mTOR*, *RPS6K1*, *IGF1*, *IGF1R, ATG9A* and *ATG5*) (Livak and Schmittgen, 2001).

2.6. Statistical analysis

All analyses were performed using R v. 4.1.2 'Bird Hippie' (R Core Team, 2021). We fitted four

models to analyse our data depending on the data source and relationship of variables. We used a

linear model to analyse single-time data such as relative mRNA expression, hereafter gene

expression, and the regression of one-to-one parameters. To analyse body mass across restriction

time and restriction levels, we used linear mixed-effects models using 'lme4' (Bates et al., 2015)

and 'lmerTest' packages v. 3.1-3 (Kuznetsova et al., 2017) considering individuals and

experimental blocks as a random intercept (Table S3). To analyse the egg mass across days and

restriction level, we used generalised additive mixed-effects models using the 'mgcv' package v.

1.8-40 (Wood, 2017) to incorporate nonlinear forms of the predictor restriction days. We used

generalised linear mixed-effects models of the family logit using package 'aod' v. 1.3.2 (Lesnoff

and Lancelot, 2012) to analyse binary response variable (daily egg laying). In mixed models, the

individual bird ID was included as a random intercept to control for the repeated measures.

Akaike's information criterion (AICc) was used to choose the best-supported models (Burnham

and Anderson, 2010). The log of fold change was used to analyse the relative gene expression. The

Tukey test was used as *post hoc* test with p < 0.05 significance level and bars set as means ± SEM.

To see the multivariate regression of genes' mRNA expression against the fitness traits and

resource allocation strategy, we used principal component analysis (PCA) using 'prcomp' function

from 'stats' package to avoid multicollinearity between the predictor variables (R Core Team,

2021). We used the 'ggbiplot' package to visualise the clustered data against treatments (Vu,

2011). We used Kaiser's rule to retain PCs for further analysis (Kaiser, 1960). We analysed

individual genes' impact on each fitness traits using linear regression. We analysed resource

allocation strategy using the proportional change in egg mass in function of change in body mass

compared to the pre-treatment status. We used trigonometric calculation to determine the direction

of the resulting vector in radians ($-\pi/2$ - $\pi/2$) that reflects resource allocation strategy. A value of

zero corresponds to no re-allocation, positive and negative values indicate re-allocation towards

reproduction and self-maintenance, respectively. This method allowed us to analyse the

relationship between individual gene expression and resource allocation strategy across treatment

groups.

3. Results

3.1. Dietary restriction affects body mass

Experimental groups did not differ in their initial body mass, but DR reduced it significantly

(treatment: $F_{3,28} = 13.832$, $p < 0.0001$; time: $F_{2,56} = 61.826$, $p < 0.0001$; treatment × time

interaction: $F_{6,56} = 12.262$, $p < 0.0001$). Birds grouped under all restriction levels showed a

significant reduction in body mass compared to the *ad libitum*-fed birds (ADL) at both week 1 and

week 2. The DR40 also resulted in significantly lower body mass than the DR20 at both time

points, while the other restricted groups did not differ significantly (Figure 1, Table S4). All

restricted groups showed significantly lower body mass at both weeks compared to initial body

mass. The DR30 and DR40 groups showed a significant and marginally significant body mass

reduction from week 1 to week 2, respectively, while the DR20 group showed no further

significant variation from week 1 to week 2 restriction time points (Table S5).

FIGURE 1 The effect of different dietary restriction levels on body mass of female quails at different time points. Data are represented by the mean ± SE from 8 birds per group. ns: none significant at p < 0.05; *** significantly different at p < 0.001; ADL: *ad libitum*; DR20: 20% restriction; DR30: 30% restriction; DR40: 40% restriction.

3.2. Severe dietary restriction reduces reproductive traits

The treatment groups differed significantly in the total number of eggs laid during the 14 days restriction period ($F_{3,24} = 5.448$, $p < 0.005$). The DR40 group laid significantly fewer eggs than the ADL group ($t = 3.992$, $p = 0.002$), while the DR20 and DR30 did not show significant reduction (Figure 2a). The level of DR and restriction period significantly explained daily egg-laying probability. Restricted birds decreased daily egg-laying probability in comparison to the ADL group. The restriction period also significantly reduced the overall probability of egg laying (Table 1, Figure S1).

The time-dependent trend indicated that egg mass was significantly reduced in DR30 and DR40 groups starting from day 5 ($p < 0.003$). Like the ADL group, egg mass from the DR20 group showed no change throughout the restriction period. On the last days of the experiment, egg mass from the DR30 and DR40 groups showed improvement to the level that they no longer differed from the ADL and DR20 groups (Figure 2b, Table S6).

274
FIGURE 2. Effect of dietary restriction on egg production and egg mass. DR40 significantly
reduced total number of eggs (a), egg mass reduced at DR30 and DR40 five sixth day. Data are
represented by the mean ± SE from 8 birds per group. Means followed by a common letter are not
significantly different at p < 0.05. ns: none significant; * significantly different at p < 0.05; **
significantly different at p < 0.01. Abbreviations follow Figure 1.

TABLE 1 Generalised linear mixed model of the family logit predicting the probability of daily
egg laying as a function of dietary restriction levels restriction days with restriction treatment and
restriction days as fixed effect and individual bird as random effect.

Predictors	*Log-Odds*	*SE*	*z value*	*p-value*
Intercept	2.82	0.53	5.29	0.0001
DR20	-1.57	0.65	-2.39	0.0154
DR30	-1.69	0.65	-2.57	0.0089
DR40	-1.58	0.64	-2.44	0.0132
Day	-0.06	0.02	-2.41	0.0151
Random effects				
birdID Variance	1.127			
Observations	480			

285 birdID: individual bird, SE: standard error
286

3.3. Dietary restriction affects gene expression

The *mTOR* gene expression showed a significant and gradual decrease across the DR levels ($F_{3,28}$

$= 15.424$, $p < 0.0001$; Figure 3a, Table S7). However, the expression of *RPS6K1* was increased in

response to the treatments ($F_{3,28} = 7.522$, $p < 0.001$), showing an increasing trend with the severity

of the restrictions (Figure 3b, Table S7).

The DR treatment also significantly decreased *IGF1* gene expression ($F_{3,28} = 8.998$, $p <$

0.0002). All the restricted groups had significantly lower *IGF1* gene expression than the ADL-fed

controls. Contrary to *mTOR*, the downregulation of *IGF1* did not intensify with the severity of the

treatment (Figure 3e, Table S7). Despite a similar trend, *IGF1R* gene expression remained

statistically undistinguishable among the four groups ($F_{3,28} = 1.108$, $p = 0.362$, Figure 3f).

Furthermore, the restriction treatment significantly increased *ATG9A* and *ATG5* gene expression

(ATG9A: $F_{3,28} = 5.726$, $p < 0.01$, ATG5: $F_{3,28} = 4.117$, $p < 0.05$, Figure 3c,d).

299
FIGURE 3 The 14 days dietary restriction showed a significant effect on relative mRNA
expression (log of fold change) of (a) mechanistic target of rapamycin (*mTOR*), (b) ribosomal
protein S6 kinase 1 (*RPS6K1*), (c) autophagy-related 9A (*ATG9A*), (d) autophagy-related 5
(*ATG5*), (e) insulin-like growth factor 1 (*IGF1*), whereas had no significant effect on (f) insulin-
like growth factor 1 receptor (*IGF1R*). Data are represented by the mean ± SE from 8 birds per
group. Means followed by a common letter are not significantly different at p < 0.05. Dots on top
of letters indicates marginaly significant difference (p<0.1). Abbreviations follow Figure 1.
307

3.4. Gene expression is related to fitness

The PC analysis indicates that cumulatively 63.7% of the variation is explained by PC1 and PC2

with an eigenvalue greater than 1 (Table S9). PC1 reflects the expression of *ATG9A*, *RPS6K1*,

ATG5, *mTOR*, and *IGF1*, whereas PC2 reflects mainly the *IGF1R*, *mTOR, ATG9A* and *IGF1*

expression (Figure 4). *mTOR* and *IGF1* expressions contributed positively, while *RPS6K1, ATG9A*

and *ATG5* expression negatively to variation in PC1 (Table S8). While *mTOR* and *IGF1* are

positively correlated, both are negatively correlated with *RPS6K1* and *ATG9A* (Figure S4).

FIGURE 4 Dimensional indication of PCs of genes expression as independent variables in line
with dietary restriction levels clustering. Abbreviations follow Figure 1.
318

Variation in both PC1 and PC2 significantly explained body mass. However, reproductive

parameters (egg number and egg mass) were related to only PC1 (Table 2). Therefore, a significant

increase in the positive contributor variables of PC1, the *mTOR* and *IGF1*, and a decrease in the

negative contributor variables, the *RPS6K1, ATG9A* and *ATG5*, lead to an increase in body mass,

egg number and egg mass. A significant decrease in the major contributors to the PC2, *IGF1R* and

mTOR significantly decrease the body mass.

Individually, the expression of mTOR and IGF1 significantly explains all fitness variables

positively. The RPS6K1 expression showed a negative relationship with body mass and egg mass,

while the ATG9A expression was negatively related to body mass and egg number (Figure S3).

As mTOR plays a crucial role in gene transcription, we conducted tests to examine its relationship

with the expression of the other genes. Our findings indicate that it significantly accounts for the

expression of these genes (Figure S5).

TABLE 2 Output of the multiple linear regression of PCs from gene expression predicting body
mass, egg number and egg mass.

Response variable	Predictor	Coefficient	SE	t	P-value	R^2	RSE
	Intercept	246.83	4.13	59.70			
Body mass	PC1	12.22	2.68	4.56	<0.0001	0.48	23.39
	PC2	-9.12	3.59	-2.54	0.02		
	Intercept	10.72	0.41	26.34			
Egg number	PC1	0.70	0.26	2.75	0.011	0.22	2.23
	PC2	-0.082	0.35	-0.24	0.82		
	Intercept	11.64	0.18	64.89			
Egg mass	PC1	0.36	0.12	3.13	0.004	0.32	0.96
	PC2	-0.23	0.15	-1.53	0.14		

3.5. mTOR expression is associated with resource re-allocation

On week 1, only the DR20 group increased relative reproductive investment (p = 0.02), while the

ADL controls and the two other treatment groups did not deviate significantly from zero re-

allocation, and all treatment groups had similar strategy by the end of the second week (p > 0.4,

Figure 5). At the end of the experiment, individual variation in allocation strategy was only related

to mTOR expression (t = –3.118, p = 0.004). Irrespective of the treatment, individuals with lower

mTOR values were more likely to invest proportionally more in reproduction than individuals with

higher mTOR activity (Figure 5).

FIGURE 5 Effect of dietary restriction on resource allocation decision. (a) A conceptual figure
illustrating resource allocation decision. The x- and y-axis show the proportional change in body
mass and egg-mass, respectively, during the experimental period (compared to the pre-treatment
body mass and egg mass, respectively, thus, all vectors start from the origo). The angle of the
vectors (radians) illustrates the allocation strategy. The solid black line shows where y = x, i.e.
when there is no re-allocation: a unit change in body mass is accompanied by the same amount of
correction in egg mass. The space above the identity line indicates reproductive re-allocation: in
response to a change of available resources, the individual allocates more to reproduction at the
cost of self-maintenance. In contrast, the space under the identity line indicates re-allocation
towards self-maintenance because the same change in body mass is associated with a
proportionally larger reduction in reproductive investment. (b) Individual allocation strategy is
related to mTOR expression. (c-d) The thin lines show individual data points, the thick vectors
show the median response of the respective treatment group. In the first week (c), DR20 and DR30
groups invest more in reproduction under a loss of body mass, while DR40 group tends to re-
allocate resources to self-maintenance. (d) Over two weeks, all restricted groups show
reproductive re-allocation, albeit to different degrees. The longer vectors in the DR40 also
illustrate that this treatment imposed a higher cost than in the other two groups (where the lengths
of the vectors are similar). The ADL control group remains unchanged over time. Abbreviations
follow Figure 1.

4. Discussion

366 Understanding the evolution of resource allocation and its underlying mechanisms remains a major

367 challenge in biology (Moatt et al., 2020; Ng'oma et al., 2017). In this study, we decreased resource

availability along a gradient of varying severity and investigated the body mass change, reproductive performance and expression of key genes in the nutrient-sensing IGF-1/mTOR pathway. Our study provided four key results.

First, we found that the severity of the feed restriction affected the body mass differently (Figure 1), indicating that our manipulation of resource availability was successful. Even a mild (20%) feed restriction decreased body mass in the first week, but birds could stabilise their body mass in the second week. The 30% and 40% restrictions resulted in stronger initial body mass losses and birds continued to lose mass in the second week. Birds with higher metabolic rate could be more sensitive to variation in food availability (Brzęk et al., 2012; Zhang et al., 2018).

Second, despite the effective treatments, reproductive performance dropped considerably only when food restriction was more severe. Under 20% food restriction, only the probability of daily egg laying was reduced, indicating that birds were more likely to skip some days in laying, whereas controls reliably laid an egg every day. However, the total number of eggs and the egg mass throughout the study remained similar in DR20 to the ADL-fed birds. In DR30 birds, while the number of eggs remained similar to ADL, the probability of egg laying and the mass of the eggs were reduced, especially in the second half of the experiment. In the most severely restricted hens (DR40), all three reproductive parameters were affected: the probability of egg laying decreased during the study, resulting in fewer and smaller eggs than the controls. However, even under the highest restriction level, the birds continued reproduction, indicating that despite an overall reduction in their resource pool, they managed to keep their reproductive performance, in some cases (DR20) even at the level of the *ADL* controls. However, depending on the magnitude of reduction in the available energy, birds had to face different trade-offs. At a low restriction level (DR20), individuals had to allocate more resources from a limited budget to reproduction, but they could do it without compromising egg size (Figure S2). When resources became more limiting (DR30), birds had to trade-off quality for quantity of reproduction. Under even more challenging conditions (DR40), egg number and mass and also body mass are compromised. Our analysis of resource allocation strategy supports the idea that birds invest in reproduction at moderate restriction (DR20), whereas favour self-maintenance at more severe restriction (DR40) (Figure 5). These results corroborate previous findings (Li et al., 2011; Mahrose et al., 2022), indicating that moderate restriction improves egg production at the expense of egg mass and body mass. Moderate DR has also shown a positive effect on preserving reproductive capacity in mammals (Sun et al.,

2021). Mild DR has for years been used in the poultry sector to avoid rapid growth and maintain
reproductive life- and health span (Holmes and Ottinger, 2003). A study on rainbow trout indicated
that a 20% restriction led to the production of bigger eggs than the ADL fed (Cardona et al., 2019)
and suggested that organisms fed ADL seem to invest much of their energy for growth, while the
moderately restricted organisms favour investing in their reproductive success.

Third, in response to our treatment, we found characteristic signatures in gene expression
patterns. DR downregulated both *mTOR* and *IGF1*, albeit in different ways. The pattern of change
in *mTOR* gene expression across treatments groups mirrored the variation in body mass loss and
showed a dose-dependent reaction, where the downregulation of *mTOR* gene expression was
proportional to body mass loss (Figure 1, 3a, Table S4, S8). In contrast, *IGF1* expression was
affected equally under all restricted groups. Nutritional stress has been found to downregulate the
expression of the *IGF1* gene, resulting in a decrease in circulating levels of IGF-1. The deficiency
of IGF-1 has been observed to have pleiotropic effects (Lodjak and Verhulst, 2020). The result
also indicates that the mTOR gene expression is more sensitive to a gradient of nutritional
deficiency. Although the specific mechanisms underlying the effect of DR on mTOR gene
expression have not been thoroughly investigated, our findings shed light on the similarity between
the effect of DR on gene expression and the previously studied effect on the abundance of activated
mTORC1 (Velingkaar et al., 2020).

The expression of the mTOR gene is crucial for the cellular production of the mTOR
protein, which is then assembled into mTORC1 and mTORC2 complexes along with other
component proteins (Szwed et al., 2021). The reduced *mTOR* gene expression could contribute to
a lower mTORC1 abundance for activation. Concurrently, DR can downregulate the expression of
potential mTORC1 upstream activators. On the other hand, in normal nutritional conditions,
activated mTORC1 suppresses mTORC2 activation by phosphorylating RPSK1 (Oh and Jacinto,
2011). Consequently, reduced mTOR gene expression during DR may have a positive impact on
the activation of mTORC2 and subsequent cell survival under dietary stress. mTORC2 is also
activated by the energy stress sensor AMPK under restriction conditions (Szwed et al., 2021).

Contrary to our assumption, we also found that *RPS6K1* expression increased with the
severity of restriction (Figure 3b). In response to phosphorylation by mTOR, RPS6K1 initiates
ribosomal translation, consequently promoting cell growth and differentiation (Saxton and
Sabatini, 2017). Previous studies have reported reduced *RPS6K1* expression in the liver of overfed

geese (Han et al., 2015) and in the brain of restricted mice (Ma et al., 2015). Therefore, we expected

that downregulation of *mTOR* would reduce the expression of *RPS6K1* and subsequently reduces

body mass (Bae et al., 2012). Despite the predicted decrease in *mTOR* expression, body mass and

RPS6K1 expression were negatively correlated (Figure S3). While a consistent response is

expected (Buccitelli and Selbach, 2020), expression of *RPS6K1* gene and the phosphorylated

RPS6K1 protein might respond differently to upstream factors. Protein expression of RPS6K1

kinase alone is not sufficient to initiate ribosomal protein translation, rather, it needs to be

phosphorylated by the activated mTORC1 kinase (Holz et al., 2005). Hence, the mechanism of

action of *mTOR* and *RPS6K1* gene expression, their total protein expression, their phosphorylated

expression, their impact on shaping fitness traits are of future research interests. Although there is

growing evidence for the correlation between gene expression and protein abundance

(Koussounadis et al., 2015; Nie et al., 2006), post-transcriptional modification may alter the

biological function of these genes. Currently, the paucity of research reporting the effect of DR on

RPS6K1 gene expression hinders the generalisation of the observed patterns. Nevertheless, our

results suggest that *RPS6K1* may be critical in resource allocation decisions. The other genes of

interest are the autophagy related genes, *ATG9A* and *ATG5*, the genes involved in autophagosome

formation, elongation and closure. Both genes showed a tendency to be upregulated under all

restriction groups (Figure 3c).

mTOR expression was positively related to *IGF1* expression and its signalling receptor

IGF1R (Figure S5). The mTOR pathway not only affects translation but is also a key regulator of

gene transcription by regulating the activity of specific transcription factors, epigenetic

mechanisms or by affecting RNA stability (Laplante and Sabatini, 2013). The modification of

transcription factors is important for activation, translocation, interaction, stability and binding

(Filtz et al., 2014; Sukumaran et al., 2020). The mTORC1 phosphorylates transcription factors in

response to resource availability, which in turn regulate several essential genes. Evidence shows

that mTORC1 itself can function as a transcription factor when it is localised in the nucleus (Jiang,

2010; Tsang et al., 2010). Therefore, activated mTORC1 can upregulate the transcription of the

IGF1, *IGF1R* and the *mTOR* gene itself. In the case of DR, the inhibition of mTOR can lead to a

decrease in the expression of genes involved in growth and reproduction, including *IGF1*.

Contrarily, under scarce resources (DR), the downregulation of mTORC1 allows the nuclear

localisation and activity of Transcription Factor EB (TFEB) and upregulates autophagosome

formation through coordinating the expression of genes involved in autophagy such as *ATG9A* and

ATG5 (Martina et al., 2012; Napolitano and Ballabio, 2016). These transcriptional factors are

mainly related to maintenance of cellular homeostasis by regulating autophagy and lysosomal

genes at the transcriptional level during nutritional deficiency (Inoki et al., 2012; Martina et al.,

2012). The correlation analysis in the present study revealed that the expression of *ATG9A* and

ATG5 is negatively related to *mTOR* expression (Figure S4), indicating that the downregulation of

mTOR mediates the upregulatory effect of DR on autophagy genes.

Finally, we found that variation in the gene expression pattern was coordinated and related

to fitness parameters and resource allocation. At a severe DR, the reduced egg number and egg

mass are aligned with low *IGF1* and *mTOR* expression, suggesting that these genes are associated

with the effect of DR on reproduction (Figure S3). However, individual variation in resource

allocation strategy was only related to *mTOR* expression. Stronger restrictions induced an

increasing reduction of *mTOR* expression, but irrespective of the treatment, individuals with

relatively lower *mTOR* expression had a proportionally larger reproductive investment. This may

seem surprising because mTORC1 is required for and thought to promote reproduction (Guo et

al., 2018; McLaughlin et al., 2011). The resource re-allocation hypothesis suggests that organisms

shift resources between reproduction and somatic maintenance when faced with limited resources

(Moatt et al., 2020; Regan et al., 2020), a process mediated by the mTOR pathway. When nutrition

is limited, mTORC1 activity is downregulated, triggering alternative pathways (Johnson et al.,

2013; Li et al., 2015). In our study, the higher resource re-allocation to reproduction at a lower

individual *mTOR* expression may have triggered upregulation of the autophagy pathway and

recycling of damaged cell contents as an energy substitution for the nutrient deficit (Adler and

Bonduriansky, 2014; Chung and Chung, 2019). The upregulated cellular maintenance helps to

preserve the follicle pool and maintain reproductive potential (English and Bonsall, 2019), while

activation of autophagy-related genes promotes oocyte maturation (Zhou et al., 2019). Rapamycin

treatment, which downregulates mTORC1 was also found to stimulate oocyte maturation by

increasing the expression of autophagy-related genes (Lee et al., 2015). In our study, un-inhibition

(i.e. upregulation) of recycling mechanisms may have channelled resources towards reproduction.

In conclusion, this study revealed that resource limitation-induced allocation trade-offs are

associated with a differential expression of nutrient-sensing genes. A limited energy budget

induces a lower expression of *mTOR* and *IGF1* and a higher expression of *RPS6K1, ATG9A* and

ATG5 differently at different restriction levels, and leads to overall lower fitness values. However,

individuals showing relatively lower *mTOR* expression invest proportionally more in reproduction,

which contradicts the established premise that mTOR mediates resource allocation towards

reproduction. This apparent paradox may be resolved by a deeper understanding of mTOR's

stimulatory, suppressive and permissive functions.

Data accessibility statements

499 All data will be archived in Dryad upon acceptance. R software code for analysis will be archived
500 on relevant database upon acceptance.

Funding information

503 The study was supported by the National Development, Research and Innovation Office, Hungary
504 (OTKA K139021).

Acknowledgements

507 We would like to thank James Kachungwa Lugata, Angyal Eszter and Fadella Nur Almira for their
508 help during the experiment. Our thanks also go to Zoltán Németh for valuable discussion and
509 inputs.

Authors' contributions

GKR.: conceptualisation, data curation, formal analysis, investigation, visualisation, methodology,
writing—original draft, writing—review and editing; SFN.: conceptualisation, methodology,
writing—review and editing; BC: data curation, methodology and writing—review and editing;
RK: methodology and writing—review and editing; CSz.: methodology, resource and writing—
review and editing; GG: methodology and writing—review and editing; LC: conceptualisation,
investigation, supervision, funding acquisition, project administration, resources, validation and
writing—review and editing; AZL: conceptualisation, data curation, formal analysis, visualisation,
methodology, supervision, funding acquisition, project administration, resources, validation and
writing—review and editing.
All authors gave final approval for publication and agreed to be held accountable for the
work performed therein.

Conflict of interest declaration

525 The authors declare no competing interests.
526

References

528 Adler, M. I.and Bonduriansky, R. (2014): Why do the well-fed appear to die young? *Bioessays*. 36(5),
529   439-450. https://doi.org/10.1002/bies.201300165
530 Bae, E. J., Xu, J., Oh, D. Y., Bandyopadhyay, G., Lagakos, W. S., Keshwani, M.and Olefsky, J. M.
531   (2012): Liver-specific p70 S6 kinase depletion protects against hepatic steatosis and systemic insulin

resistance. *Journal of Biological Chemistry*. 287(22), 18769-18780.
https://doi.org/10.1074/jbc.M112.365544

Baloni, P., Sangar, V., Yurkovich, J. T., Robinson, M., Taylor, S., Karbowski, C. M., Hamadeh, H. K., He, Y. D.and Price, N. D. (2019): Genome-scale metabolic model of the rat liver predicts effects of diet restriction. *Sci Rep*. 9(1), 9807. https://doi.org/10.1038/s41598-019-46245-1

Barja, G. (1998): Mitochondrial free radical production and aging in mammals and birds. *Annals of the New York Academy of Sciences*. 854(1), 224-238. https://doi.org/10.1111/j.1749-6632.1998.tb09905.x

Bates, D., Mächler, M., Bolker, B.and Walker, S. (2015): Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*. 67(1), 1-48. https://doi.org/10.18637/jss.v067.i01

Brzęk, P., Książek, A., Dobrzyń, A.and Konarzewski, M. (2012): Effect of dietary restriction on metabolic, anatomic and molecular traits in mice depends on the initial level of basal metabolic rate (BMR). *The Journal of Experimental Biology*. 215(18), 3191–3199.
https://doi.org/10.1242/jeb.065318

Buccitelli, C.and Selbach, M. (2020): mRNAs, proteins and the emerging principles of gene expression control. *Nature Reviews Genetics*. 21(10), 630-644. https://doi.org/10.1038/s41576-020-0258-4

Burnham, K. P.and Anderson, D. R. (2010): "Model selection and multimodel inference: a practical information-theoretic approach," Springer, New York, NY.

Cardona, E., Bugeon, J., Guivarc'h, F., Goardon, L., Panserat, S., Labbé, L., Corraze, G., Skiba-Cassy, S.and Bobe, J. (2019): Positive impact of moderate food restriction on reproductive success of the rainbow trout Oncorhynchus mykiss. *Aquaculture*. 502, 280-288.
https://doi.org/10.1016/j.aquaculture.2018.12.057

Carlsson, H., Ivimey-Cook, E., Duxbury, E. M., Edden, N., Sales, K.and Maklakov, A. A. (2021): Ageing as "early-life inertia": Disentangling life-history trade-offs along a lifetime of an individual. *Evolution Letters*. 5(5), 551-564. https://doi.org/10.1002/evl3.254

Carroll, B.and Korolchuk, V. I. (2018): Nutrient sensing, growth and senescence. *The FEBS journal*. 285(11), 1948-1958. https://doi.org/ 10.1111/febs.14400

Chang van Oordt, D. A., Taff, C. C., Ryan, T. A.and Vitousek, M. N. (2022): Timing of breeding reveals a trade-off between immune investment and life history in tree swallows. *Integrative and Comparative Biology*. 62(6), 1629-1639. https://doi.org/10.1093/icb/icac033

Chung, K. W.and Chung, H. Y. (2019): The Effects of Calorie Restriction on Autophagy: Role on Aging Intervention. *Nutrients*. 11(12), 2923. https://doi.org/10.3390/nu11122923

Colman, R. J., Beasley, T. M., Kemnitz, J. W., Johnson, S. C., Weindruch, R.and Anderson, R. M. (2014): Caloric restriction reduces age-related and all-cause mortality in rhesus monkeys. *Nature communications*. 5(1), 3557. https://doi.org/10.1038/ncomms4557

Deng, H., Zheng, A., Liu, G., Chang, W., Zhang, S.and Cai, H. (2014): Activation of mammalian target of rapamycin signaling in skeletal muscle of neonatal chicks: effects of dietary leucine and age. *Poultry Science*. 93(1), 114-121. https://doi.org/10.3382/ps.2013-03287

Efeyan, A., Comb, W. C.and Sabatini, D. M. (2015): Nutrient-sensing mechanisms and pathways. *Nature*. 517(7534), 302-310. https://doi.org/10.1038/nature14190

English, S.and Bonsall, M. B. (2019): Physiological dynamics, reproduction-maintenance allocations, and life history evolution. *Ecology and Evolution*. 9(16), 9312-9323. https://doi.org/10.1002/ece3.5477

Feng, Z.and Levine, A. J. (2010): The regulation of energy metabolism and the IGF-1/mTOR pathways by the p53 protein. *Trends in Cell Biology*. 20(7), 427-434. https://doi.org/10.1016/j.tcb.2010.03.004

Fenton, T. R.and Gout, I. T. (2011): Functions and regulation of the 70 kDa ribosomal S6 kinases. *The International Journal of Biochemistry & Cell Biology*. 43(1), 47-59.
https://doi.org/10.1016/j.biocel.2010.09.018

Filtz, T. M., Vogel, W. K.and Leid, M. (2014): Regulation of transcription factor activity by interconnected post-translational modifications. *Trends in Pharmacological Sciences*. 35(2), 76-85.
https://doi.org/10.1016/j.tips.2013.11.005

Flatt, T., Amdam, G. V., Kirkwood, T. B. L.and Omholt, S. W. (2013): Life-history evolution and the polyphenic regulation of somatic maintenance and survival. *The Quarterly Review of Biology*. 88(3), 185-218. https://doi.org/10.1086/671484

Fu, W.and Hall, M. N. (2020): Regulation of mTORC2 signaling. *Genes*. 11(9), 1045. https://doi.org/10.3390/genes11091045

Gokarn, R., Solon-Biet, S. M., Cogger, V. C., Cooney, G. J., Wahl, D., McMahon, A. C., Mitchell, J. R., Mitchell, S. J., Hine, C., de Cabo, R., et al. (2018): Long-term dietary macronutrients and hepatic gene expression in aging mice. *J Gerontol A Biol Sci Med Sci*. 73(12), 1618-1625. https://doi.org/10.1093/gerona/gly065

Guo, J., Zhang, T., Guo, Y., Sun, T., Li, H., Zhang, X., Yin, H., Cao, G., Yin, Y., Wang, H., et al. (2018): Oocyte stage-specific effects of MTOR determine granulosa cell fate and oocyte quality in mice. *PNAS*. 115(23), E5326-E5333. https://doi.org/10.1073/pnas.1800352115

Guo, Z.and Yu, Q. (2019): Role of mTOR signaling in female reproduction. *Frontiers in endocrinology*. 10, 692. https://doi.org/10.3389/fendo.2019.00692

Han, C., Ye, F., Shen, X., Liu, D., He, F., Wei, S., Xu, H., Li, L.and Liu, H. (2015): Change of the mTOR pathway in tissues of overfed geese. *Brazilian Journal of Poultry Science*. 17(3), 293-300. https://doi.org/10.1590/1516-635x1703293-300

Hao, E.-Y., Wang, D.-H., Chen, Y.-F., Zhou, R.-Y., Chen, H.and Huang, R.-L. (2021): The relationship between the mTOR signaling pathway and ovarian aging in peak-phase and late-phase laying hens. *Poultry Science*. 100(1), 334-347. https://doi.org/10.1016/j.psj.2020.10.005

Hassan, S. M., Mady, M. E., Cartwright, A. L., Sabri, H. M.and Mobarak, M. S. (2003): Effect of early feed restriction on reproductive performance in Japanese quail (Coturnix coturnix japonica). *Poultry Science*. 82(7), 1163-1169. https://doi.org/10.1093/ps/82.7.1163

Holmes, D. J.and Ottinger, M. A. (2003): Birds as long-lived animal models for the study of aging. *Experimental Gerontology*. 38(11-12), 1365-1375. https://doi.org/10.1016/j.exger.2003.10.018

Holz, M. K., Ballif, B. A., Gygi, S. P.and Blenis, J. (2005): mTOR and S6K1 mediate assembly of the translation preinitiation complex through dynamic protein interchange and ordered phosphorylation events. *Cell*. 123(4), 569-580. https://doi.org/10.1016/j.cell.2005.10.024

Inness, C. L. W.and Metcalfe, N. B. (2008): The impact of dietary restriction, intermittent feeding and compensatory growth on reproductive investment and lifespan in a short-lived fish. *Proceedings of the Royal Society B: Biological Sciences*. 275(1644), 1703-1708. https://doi.org/10.1098/rspb.2008.0357

Inoki, K., Kim, J.and Guan, K.-L. (2012): AMPK and mTOR in cellular energy homeostasis and drug targets. *Annual Review of Pharmacology and Toxicology*. 52(1), 381-400. https://doi.org/10.1146/annurev-pharmtox-010611-134537

Jiang, Y. (2010): mTOR goes to the nucleus. *Cell Cycle*. 9(5), 868. https://doi.org/10.4161/cc.9.5.11070

Jimenez, A. G., O'Connor, E. S., Tobin, K. J., Anderson, K. N., Winward, J. D., Fleming, A., Winner, C., Chinchilli, E., Maya, A.and Carlson, K. (2019): Does cellular metabolism from primary fibroblasts and oxidative stress in blood differ between mammals and birds? The (lack-thereof) scaling of oxidative stress. *Integrative and Comparative Biology*. 59(4), 953-969. https://doi.org/10.1093/icb/icz017

Johnson, S. C. (2018): Nutrient sensing, signaling and ageing: The role of IGF-1 and mTOR in ageing and age-related disease. *In* "Biochemistry and Cell Biology of Ageing: Part I Biomedical Science", Vol. 90, pp. 49-97. Springer Singapore.

Johnson, S. C., Rabinovitch, P. S.and Kaeberlein, M. (2013): mTOR is a key modulator of ageing and age-related disease. *Nature*. 493(7432), 338-345. https://doi.org/10.1038/nature11861

Kaiser, H. F. (1960): The application of electronic computers to factor analysis. *Educational and Psychological Measurement*. 20(1), 141-151. https://doi.org/10.1177/00131644600200

Kapahi, P., Kaeberlein, M.and Hansen, M. (2017): Dietary restriction and lifespan: Lessons from invertebrate models. *Ageing Research Reviews*. 39, 3-14. https://doi.org/10.1016/j.arr.2016.12.005

Koussounadis, A., Langdon, S. P., Um, I. H., Harrison, D. J.and Smith, V. A. (2015): Relationship
between differentially expressed mRNA and mRNA-protein correlations in a xenograft model
system. *Scientific Reports*. 5(1), 10775. https://doi.org/10.1038/srep10775
Kuznetsova, A., Brockhoff, P. B.and Christensen, R. H. B. (2017): LmerTest package: Tests in linear
mixed effects models. *Journal of Statistical Software*. 82(13), 1–26.
https://doi.org/10.18637/jss.v082.i13
Laplante, M.and Sabatini, D. M. (2012): mTOR signaling in growth control and disease. *Cell*. 149(2),
274-293. https://doi.org/10.1016/j.cell.2012.03.017
Laplante, M.and Sabatini, D. M. (2013): Regulation of mTORC1 and its impact on gene expression at a
glance. *Journal of Cell Science*. 126(Pt 8), 1713-1719. https://doi.org/10.1242/jcs.125773
Lee, J., Park, J.-I., Im Yun, J., Lee, Y., Yong, H., Lee, S. T., Park, C.-K., Hyun, S.-H., Lee, G.-S.and Lee,
E. (2015): Rapamycin treatment during in vitro maturation of oocytes improves embryonic
development after parthenogenesis and somatic cell nuclear transfer in pigs. *Journal of Veterinary*
Science. 16(3), 373-380. https://doi.org/10.4142/jvs.2015.16.3.373
Lesnoff, M.and Lancelot, R. (2012): aod: Analysis of Overdispersed Data. R package version 1.3.2.
https://cran.r-project.org/package=aod
Li, F., Xu, L. M., Shan, A. S., Hu, J. W., Zhang, Y. Y.and Li, Y. H. (2011): Effect of daily feed intake in
laying period on laying performance, egg quality and egg composition of genetically fat and lean
lines of chickens. *British poultry science*. 52(2), 163-168.
https://doi.org/10.1080/00071668.2011.559455
Li, L., Fu, Y.-C., Xu, J.-J., Lin, X.-H., Chen, X.-C., Zhang, X.-M.and Luo, L.-L. (2015): Caloric
restriction promotes the reserve of follicle pool in adult female rats by inhibiting the activation of
mammalian target of rapamycin signaling. *Reproductive Sciences*. 22(1), 60-67.
https://doi.org/10.1177/1933719114542016
Livak, K. J.and Schmittgen, T. D. (2001): Analysis of relative gene expression data using real-time
quantitative PCR and the 2−ΔΔCT method. *Methods*. 25(4), 402-408.
https://doi.org/10.1006/meth.2001.1262
Lodjak, J.and Verhulst, S. (2020): Insulin-like growth factor 1 of wild vertebrates in a life-history context.
Molecular and Cellular Endocrinology. 518, 110978. https://doi.org/10.1016/j.mce.2020.110978
Ma, L., Dong, W., Wang, R., Li, Y., Xu, B., Zhang, J., Zhao, Z.and Wang, Y. (2015): Effect of caloric
restriction on the SIRT1/mTOR signaling pathways in senile mice. *Brain Research Bulletin*. 116, 67-
72. https://doi.org/10.1016/j.brainresbull.2015.06.004
Ma, Y., Vassetzky, Y.and Dokudovskaya, S. (2018): mTORC1 pathway in DNA damage response.
Biochimica et Biophysica Acta - Molecular Cell Research. 1865(9), 1293-1311.
https://doi.org/10.1016/j.bbamcr.2018.06.011
Mahrose, K. M., Abol-Ela, S., Amin, R. M.and Abou-Kassem, D. E. (2022): Restricted feeding could
enhance feed conversion ratio and egg quality of laying Japanese quail kept under different stocking
densities. *Animal Biotechnology*. 33(1), 141-149. https://doi.org/10.1080/10495398.2020.1810059
Martina, J. A., Chen, Y., Gucek, M.and Puertollano, R. (2012): mTORC1 functions as a transcriptional
regulator of autophagy by preventing nuclear transport of TFEB. *Autophagy*. 8(6), 903-914.
https://doi.org/10.4161/auto.19653
Maruska, K. P., Butler, J. M.and Field, K. E. (2018): Alternative Reproductive Phenotypes Within
Species. *In* "Encyclopedia of Reproduction", pp. 41-49. Elsevier.
McCracken, A. W., Adams, G., Hartshorne, L., Tatar, M.and Simons, M. J. P. (2020): The hidden costs of
dietary restriction: Implications for its evolutionary and mechanistic origins. *Science Advances*. 6(8),
eaay3047. https://doi.org/10.1126/sciadv.aay3047
McLaughlin, M., Patrizio, P., Kayisli, U., Luk, J., Thomson, T. C., Anderson, R. A., Telfer, E. E.and
Johnson, J. (2011): mTOR kinase inhibition results in oocyte loss characterized by empty follicles in
human ovarian cortical strips cultured in vitro. *Fertility and Sterility*. 96(5), 1154-9.e1.
https://doi.org/10.1016/j.fertnstert.2011.08.040

Metcalfe, N. B.and Monaghan, P. (2013): Does reproduction cause oxidative stress? An open question. *Trends in Ecology & Evolution*. 28(6), 347-350. https://doi.org/10.1016/j.tree.2013.01.015

Mierziak, J., Kostyn, K., Boba, A., Czemplik, M., Kulma, A.and Wojtasik, W. (2021): Influence of the bioactive diet components on the gene expression regulation. *Nutrients*. 13(11), 3673. https://doi.org/10.3390/nu13113673

Moatt, J. P., Nakagawa, S., Lagisz, M.and Walling, C. A. (2016): The effect of dietary restriction on reproduction: a meta-analytic perspective. *BMC Evolutionary Biology*. 16(1), 199. https://doi.org/10.1186/s12862-016-0768-z

Moatt, J. P., Savola, E., Regan, J. C., Nussey, D. H.and Walling, C. A. (2020): Lifespan extension via dietary restriction: time to reconsider the evolutionary mechanisms? *Bioessays*. 42(8), 1900241. https://doi.org/10.1002/bies.201900241

Montoya, B., Tóth, Z., Lendvai, Á. Z., Stier, A., Criscuolo, F., Zahn, S.and Bize, P. (2022): Does IGF-1 shape life-history trade-offs? Opposite associations of IGF-1 with telomere length and body size in a free-living bird. *Frontiers in Ecology and Evolution*. 10, 853674. https://doi.org/10.3389/fevo.2022.853674

Napolitano, G.and Ballabio, A. (2016): TFEB at a glance. *Journal of Cell Science*. https://doi.org/10.1242/jcs.146365

Ng'oma, E., Perinchery, A. M.and King, E. G. (2017): How to get the most bang for your buck: the evolution and physiology of nutrition-dependent resource allocation strategies. *Proceedings of the Royal Society B: Biological Sciences*. 284(1857), 20170445. https://doi.org/10.1098/rspb.2017.0445

Nie, L., Wu, G.and Zhang, W. (2006): Correlation of mRNA expression and protein abundance affected by multiple sequence features related to translational efficiency in Desulfovibrio vulgaris: A quantitative analysis. *Genetics*. 174(4), 2229-2243. https://doi.org/10.1534/genetics.106.065862

Nojima, H., Tokunaga, C., Eguchi, S., Oshiro, N., Hidayat, S., Yoshino, K.-I., Hara, K., Tanaka, N., Avruch, J.and Yonezawa, K. (2003): The mammalian target of rapamycin (mTOR) partner, raptor, binds the mTOR substrates p70 S6 kinase and 4E-BP1 through their TOR signaling (TOS) motif. *Journal of Biological Chemistry*. 278(18), 15461-15464. https://doi.org/10.1074/jbc.c200665200

NRC (National Research Council). (1994): "Nutrient Requirements of Poultry," 9th Edition/Ed. National Academy Press, Washington, D. C.

Oh, W. J.and Jacinto, E. (2011): mTOR complex 2 signaling and functions. *Cell cycle*. 10(14), 2305-2316. https://doi.org/10.4161/cc.10.14.16586

Ottinger, M. A., Mobarak, M., Abdelnabi, M., Roth, G., Proudman, J.and Ingram, D. K. (2005): Effects of calorie restriction on reproductive and adrenal systems in Japanese quail: are responses similar to mammals, particularly primates? *Mechanisms of Ageing and Development*. 126(9), 967-975. https://doi.org/10.1016/j.mad.2005.03.017

Papadopoli, D., Boulay, K., Kazak, L., Pollak, M., Mallette, F., Topisirovic, I.and Hulea, L. (2019): mTOR as a central regulator of lifespan and aging. *F1000Research*. 8, 998. https://doi.org/10.12688/f1000research.17196.1

R Core Team. (2021): R: A language and environment for statistical computing. Computer software version 4.1.2. Vienna, Austria. https://R-project.org/.

Rabanal-Ruiz, Y.and Korolchuk, V. (2018): MTORC1 and nutrient homeostasis: The central role of the lysosome. *International Journal of Molecular Sciences*. 19(3), 818. https://doi.org/10.3390/ijms19030818

Regan, J. C., Froy, H., Walling, C. A., Moatt, J. P.and Nussey, D. H. (2020): Dietary restriction and insulin-like signalling pathways as adaptive plasticity: A synthesis and re-evaluation. *Functional Ecology*. 34(1), 107-128. https://doi.org/10.1111/1365-2435.13418

Rollins, J. A., Shaffer, D., Snow, S. S., Kapahi, P.and Rogers, A. N. (2019): Dietary restriction induces posttranscriptional regulation of longevity genes. *Life Science Alliance*. 2(4), e201800281. https://doi.org/10.26508/lsa.201800281

Sancak, Y., Bar-Peled, L., Zoncu, R., Markhard, A. L., Nada, S.and Sabatini, D. M. (2010): Ragulator-
Rag complex targets mTORC1 to the lysosomal surface and is necessary for its activation by amino
acids. *Cell*. 141(2), 290-303. https://doi.org/10.1016/j.cell.2010.02.024
Sandri, M., Barberi, L., Bijlsma, A., Blaauw, B., Dyar, K., Milan, G., Mammucari, C., Meskers, C.,
Pallafacchina, G.and Paoli, A. (2013): Signalling pathways regulating muscle mass in ageing skeletal
muscle. The role of the IGF1-Akt-mTOR-FoxO pathway. *Biogerontology*. 14(3), 303-323.
https://doi.org/10.1007/s10522-013-9432-9
Saxton, R. A.and Sabatini, D. M. (2017): mTOR signaling in growth, metabolism, and disease. *Cell*.
169(2), 361-371. https://doi.org/10.1016/j.cell.2017.03.035
Shanley, D. P.and Kirkwood, T. B. L. (2000): Calorie restriction and aging: A life-history analysis.
Evolution. 54(3), 740-750. https://doi.org/10.1111/j.0014-3820.2000.tb00076.x
She, Y., Deng, H., Cai, H.and Liu, G. (2019): Regulation of the expression of key signalling molecules in
mTOR pathway of skeletal muscle satellite cells in neonatal chicks: Effects of leucine and glycine-
leucine peptide. *Journal of Animal Physiology and Animal Nutrition*. 103(3), 786-790.
https://doi.org/10.1111/jpn.13090
Simons, M. J. P., Koch, W.and Verhulst, S. (2013): Dietary restriction of rodents decreases aging rate
without affecting initial mortality rate – a meta-analysis. *Aging cell*. 12(3), 410-414.
https://doi.org/10.1111/acel.12061
Speakman, J. R.and Mitchell, S. E. (2011): Caloric restriction. *Molecular Aspects of Medicine*. 32(3),
159-221. https://doi.org/10.1016/j.mam.2011.07.001
Sukumaran, A., Choi, K.and Dasgupta, B. (2020): Insight on transcriptional regulation of the energy
sensing AMPK and biosynthetic mTOR pathway genes. *Frontiers in Cell and Developmental
Biology*. 8, 671. https://doi.org/10.3389/fcell.2020.00671
Sun, J., Shen, X., Liu, H., Lu, S., Peng, J.and Kuang, H. (2021): Caloric restriction in female
reproduction: is it beneficial or detrimental? *Reproductive Biology and Endocrinology*. 19(1), 1.
https://doi.org/10.1186/s12958-020-00681-1
Szwed, A., Kim, E.and Jacinto, E. (2021): Regulation and metabolic functions of mTORC1 and
mTORC2. *Physiological Reviews*. 101(3), 1371-1426. https://doi.org/10.1152/physrev.00026.2020
Tsang, C. K., Liu, H.and Zheng, X. F. S. (2010): mTOR binds to the promoters of RNA polymerase I-
and III-transcribed genes. *Cell Cycle*. 9(5), 953-957. https://doi.org/10.4161/cc.9.5.10876
Velingkaar, N., Mezhnina, V., Poe, A., Makwana, K., Tulsian, R.and Kondratov, R. V. (2020): Reduced
caloric intake and periodic fasting independently contribute to metabolic effects of caloric restriction.
Aging cell. 19(4), e13138. https://doi.org/10.1111/acel.13138
Vu, V. Q. (2011): ggbiplot: A ggplot2 based biplot. R package version 0.55.
http://github.com/vqv/ggbiplot
Wood, S. N. (2017): "Generalized additive models," Chapman and Hall/CRC, New York.
Zera, A. J.and Harshman, L. G. (2001): The Physiology of Life History Trade-Offs in Animals. *Annual
Review of Ecology, Evolution, and Systematics*. 32(1), 95-126.
https://doi.org/10.1146/annurev.ecolsys.32.081501.114006
Zhang, Y., Yang, K., Yang, P., Su, Y., Zheng, W.and Liu, J. (2018): Food restriction decreases BMR,
body and organ mass, and cellular energetics, in the Chinese Bulbul (Pycnonotus sinensis). *Avian
Research*. 9(1), 39. https://doi.org/10.1186/s40657-018-0131-8
Zhou, J., Peng, X.and Mei, S. (2019): Autophagy in ovarian follicular development and atresia.
International Journal of Biological Sciences. 15(4), 726. https://doi.org/10.7150/ijbs.30369
775
776