1 **Acentric chromosome congression and alignment on the metaphase plate occur via**
2 **kinetochore-independent forces**
3
Abstract

Chromosome congression and alignment on the metaphase plate involves lateral and

microtubule plus-end interactions with the kinetochore. Here we take advantage of our ability to

efficiently generate a GFP-marked acentric X chromosome fragment in *Drosophila* neuroblasts

to identify forces acting on chromosome arms that drive congression and alignment. We find

acentrics efficiently align on the metaphase plate, often more rapidly than kinetochore-bearing

chromosomes. Unlike intact chromosomes, the paired sister acentrics oscillate as they move to

and reside on the metaphase plate in a plane distinct and significantly further from the main

mass of intact chromosomes. Consequently, at anaphase onset acentrics are oriented either

parallel or perpendicular to the spindle. Parallel-oriented sisters separate by sliding while those

oriented perpendicularly separate via unzipping. This oscillation, together with the fact that in

monopolar spindles acentrics are rapidly shunted away from the poles, indicates that distributed

plus-end directed forces are primarily responsible for acentric migration. This conclusion is

supported by the observation that reduction of EB1 preferentially disrupts acentric alignment. In

addition, reduction of Klp3a activity, a gene required for the establishment of pole-to-pole

microtubules, preferentially disrupts acentric alignment. Taken together these studies suggest

that plus-end forces mediated by the outer pole-to-pole microtubules are primarily responsible

for acentric metaphase alignment. Surprisingly, we find that a small fraction of sister acentrics

are anti-parallel aligned indicating that the kinetochore is required to ensure parallel alignment

of sister chromatids. Finally, we find induction of acentric chromosome fragments results in a

global reorganization of the congressed chromosomes into a torus configuration.

Introduction

Chromosome fragments lacking a centromere and a telomere are known as acentrics. Due to their lack of a kinetochore, acentrics are unable to make canonical attachments to microtubules. Kinetochore-microtubule interactions play key roles in mediating chromosome congression to the spindle equator, alignment on the metaphase plate, sister chromosome separation during the metaphase-to-anaphase transition and segregation during anaphase [Navarro & Cheeseman, 2021]. Thus, acentrics might be expected to fail to align properly on the metaphase plate, display separation and segregation defects in anaphase resulting in exclusion from daughter nuclei and the formation of micronuclei [Kanda & Wahl, 2000; LaFountain et al., 2001; Fenech et al., 2011]. However, these expectations are contradicted by the finding that acentric chromosome fragments often display proper sister chromosome separation, poleward migration, and inclusion into daughter nuclei [Royou et al., 2010; Bajer, 1958; Kanda et al., 1998; Vicars et al., 2021; Karg et al., 2017; Bretscher & Fox, 2016; Karg et al., 2015; Warecki & Sullivan, 2018; Khodjakov et al., 1996; Liang et al., 1993; Warecki et al., 2020]. Proposed mechanisms include neo-centromere formation and direct association of acentrics with microtubules or a kinetochore-bearing chromosome [Ishii et al., 2008; Platero et al., 1999, Kanda et al., 2001; Ohno et al., 2016; Karg et al., 2017; Warecki & Sullivan, 2020]. Studies have shown that separation of sister acentric chromosomes during early anaphase relies on Topoisomerase II activity as well as microtubule plus-end pushing forces [Vicars et al., 2021]. Additional work has shown that the anaphase poleward segregation of acentric chromosomes involves Klp3a mediated microtubule-based movement [Karg et al., 2017].

Here we explore the mechanisms that drive congression and alignment of acentric chromosome fragments at the metaphase plate as this also provides insight into the non-kinetochore forces driving congression of intact chromosomes. As with sister chromosome separation and segregation, the kinetochore plays a key role in chromosome congression and alignment on the metaphase plate [Barisic et al., 2014; Ye et al., 2015; Chmatal et al., 2015].

Following nuclear-envelope breakdown, chromosomes positioned within the microtubule arc

emanating from opposing centrosomes quickly establish lateral interactions with the plus-end

kinetochore-associated motor protein CENP-E (kinesin-7) and are transported to the metaphase

plate [Yen, 1992; McEwen et al., 2001; Kapoor et al., 2006]. This is referred to as direct

congression. Polar ejection forces mediated by plus-end microtubule dynamics and

chromokinesins acting on the chromosomes also drive chromosomes away from the poles

toward the equator [Maiato et al., 2017; Tanaka, 2008]. Once at the equator, lateral interactions

are converted to plus-end microtubule interactions and the sister kinetochores establish

biorientation with microtubules attached to opposing poles [Krenn & Musacchio, 2015].

Opposing forces at the sister kinetochores and chromosome arms result in oscillations but

maintain the chromosomes at the metaphase plate. Congression of chromosomes located at

the spindle periphery and outside of the arc described by the opposing centrosomes and the

metaphase plate require an additional step, known as peripheral congression. Chromosomes

must first rely on kinetochore-associated Dynein for microtubule minus-end directed transport to

the spindle pole. Once at the pole, the chromosomes engage kinetochore kinesin CENP-E for

transport to the metaphase plate. [McEwen et al., 2001; Kapoor et al., 2006].

Despite the key role of the kinetochore in driving chromosome movement, acentric

chromosome fragments are capable of moving toward and aligning on the metaphase plate

[Carlson, 1938a; Royou et al., 2010; Kaye et al., 2004; Bajer, 1958; Vicars et al., 2021; Karg et

al., 2017; Rieder et al., 1986]. This was dramatically demonstrated through live analysis of

acentric fragments generated via laser ablation. These fragments rapidly moved away from the

poles at a rate similar to intact chromosomes [Rieder et al., 1986]. Subsequent live analysis of

X-chromosome acentric fragments generated through endonuclease induction in *Drosophila*

neuroblasts revealed they also experience robust anti-poleward forces and are capable of

aligning on the metaphase plate [Royou et al., 2010; Vicars et al., 2021; Karg et al., 2017;

Warecki et al., 2020].

Left unresolved are the mechanisms driving congression and metaphase plate alignment

of chromosome fragments lacking a kinetochore. Here we explore this issue by taking

advantage of our ability to efficiently generate GFP-labeled acentric fragments in the genetically

tractable *Drosophila* neuroblasts. The GFP tag enables us to distinguish the acentric from the

kinetochore bearing chromosomes and facilitates tracking of the acentric as it moves to and

aligns on the metaphase plate. We find acentrics efficiently align on the metaphase plate more

rapidly than kinetochore-bearing chromosomes. Multiple lines of evidence indicate that proper

acentric alignment on the metaphase plate relies on pole-to-pole microtubules and plus-end

directed microtubule-based forces. Surprisingly, we find that a small but significant fraction of

sister acentrics exhibit an anti-parallel alignment. This suggests that the kinetochore is required

to ensure parallel alignment of sister chromatids. In addition, we discovered that induction of

acentric chromosome fragments results in a global reorganization of the congressed

chromosomes into a torus configuration.

Results

Acentric sister chromatids undergo rapid movement at the metaphase plate

As previously described, acentric chromosome fragments have a remarkable ability to

congress to the metaphase plate, separate and segregate from one another in anaphase, and

incorporate into the reforming daughter nuclei, all while lacking canonical kinetochore-

microtubule attachments [Vicars et al., 2021; Warecki et al., 2020]. However, it has been difficult

to track an unmarked acentric chromosome fragment during congression and metaphase

alignment when the acentric is embedded within the entire chromosome complement. Here we

address this issue by taking advantage of a GFP-marker that specifically tags the *Drosophila* X

chromosome. In *Drosophila*, the male-specific lethal (MSL3) complex plays a major role in

dosage compensation by upregulating genes on the male X chromosome [Sural et al., 2008].

MSL3-GFP preferentially binds the male X euchromatin and thus is suitable for distinguishing

the X chromosome from the other chromosomes in live studies. Acentrics are generated using

heat-shock-induced expression of the I-CreI endonuclease, which targets rDNA repeats

embedded in the X-chromosome centric heterochromatin [Rong et al., 2002; Maggert & Golic,

2005; Paredes & Maggert, 2009; Golic & Golic, 2011]. This produces a kinetochore-bearing

heterochromatic chromosome fragment and a euchromatic acentric chromosome fragment. As

MSL3-GFP specifically localizes to the euchromatin, it preferentially labels the acentric

chromosome fragment and is readily tracked over the course of a mitotic division (Figure 1). The

MSL3-GFP marker does not disrupt acentric chromosome alignment, segregation, or

micronuclei formation when compared to control cells with acentrics alone. In control cells

expressing MSL3-GFP but not I-CreI, all of the intact X-chromosomes align normally (N=10).

Upon I-CreI induction, 95% (N=18) of the GFP-marked acentrics align on the metaphase plate

like intact chromosomes.

Live analysis 80 seconds prior to anaphase onset was performed to determine the rate

on which intact and acentric chromosomes move to the metaphase plate (Table 1, see

accompanying schematic). This analysis reveals that on average acentric X chromosomes

move at a significantly faster rate while actively aligning at the metaphase plate (11.7 ± 3.5

nm/s, N=14) when compared to intact X chromosomes (8.4 ± 3.3 nm/s, N=12, P=0.01, Mann-

Whitney U Test) (Table 1). Intact autosomes (8.5 ± 2.4 nm/s, N=28) align at the metaphase

plate at a rate in accordance with the intact X chromosomes (8.4 ± 3.3 nm/s, N=12, P=0.6,

Mann-Whitney U Test) (Table 1).

Acentric induction causes a global reorganization of congressed chromosomes such

that acentrics localize to the periphery of the main chromosome mass

Three-dimensional rendering of metaphase images of labeled intact X chromosomes

reveal that they are aligned with the autosomes in a planar columnar configuration (Figure 2A,

N=5). In addition, the intact X chromosome is consistently positioned at the outer edge of the

column of congressed chromosomes (Figure 2A, N=5). Unexpectedly, we found that I-CreI

induction of the acentric X chromosome results in a global reorganization of the congressed

chromosomes. Rather than organized in a columnar configuration, the chromosomes are

organized into a distinct torus with an absence of chromatin in the center (Figure 2B, N=5). We

quantified this global organization by measuring the circularity of the entire congressed mass of

chromosomes at metaphase. Using the circularity function in ImageJ, we define perfect circles

as having a circularity of 1. The circularity measurements demonstrate that the columnar

formations in cells in which I-CreI is not expressed are significantly distinct from the torus

shapes seen in neuroblasts expressing I-CreI (Figure 2C). The average circularity of metaphase

congressed chromosomes with and without acentrics is 0.95 (SD=0.013, N=6) and 0.44

(SD=0.183, N=6), respectively (Figure 2C).

As with the intact X chromosome, the acentric is peripherally localized. However, the

extent to which the acentric X is positioned away from the main chromosome mass is much

greater than that of the intact X chromosome. We quantified peripheral positioning by

calculating the distance of the intact X chromosome and acentric from the center of

chromosome mass (Figure 2C). On average, acentric X chromosomes are positioned

significantly farther (3.60 ± 1.34 µm, N=14) from the chromosomal mass center when compared

to intact X chromosomes (1.89 ± 0.74 µm, N=16, P=0.01, Mann-Whitney U Test). The center of

the mass of chromosomes is determined by defining the upper, lower, left, and right boundaries

of the chromosome mass and calculating the center point. Thus, while the acentric X

chromosome fragments efficiently congress to the metaphase plate, they reside in a position

distinct from the main chromosome complement (Figure 2).

Acentrics oscillate while aligning on the metaphase plate

In prometaphase, kinetochore-bearing replicated sister chromosomes remain paired

while congressing to the metaphase plate. This is due to the presence of cohesin protein

complexes and DNA catenations between sisters [Guacci et al., 1997; Michaelis et al., 1997;

Porter & Farr, 2004]. Similarly, sister acentric X chromosome fragments travel to the metaphase

plate as a pair, held together via cohesin and DNA catenations [Vicars et al., 2021].

Previous work demonstrated that the orientation of sister acentrics at the metaphase-to-

anaphase transition correlates with their mode of separation during late anaphase [Vicars et al.,

2021]. Congressed intact chromosomes always align on the metaphase plate perpendicular to

the spindle. In contrast, sister acentrics align either perpendicular or parallel to the spindle.

Sister acentrics aligned perpendicular to the spindle, separate via unzipping from one another,

while those aligned parallel to the spindle slide past one another to opposing poles. Here we

provide insight into the trajectories and mechanisms that result in acentrics reaching the plate in

multiple orientations. We discovered that acentrics oscillate while aligning on the metaphase

plate. To quantify this, we measure the orientation angles of acentric and intact X chromosomes

with respect to the cell equator at 5 second intervals during the 80-second interval prior to

anaphase onset (Figure 3).

Kinetochore-bearing sister chromatids align at the metaphase plate undergoing no

oscillations, most often remaining perpendicular to the mitotic spindle (Figure 3A, 3C). However,

acentric sisters oscillate as they line up at the metaphase plate, alternating between orienting

parallel and perpendicular to the spindle and division axis (Figure 3B, 3C). These results

suggest that the parallel and perpendicular orientations of the acentric at the metaphase plate

are due to the fact that they oscillate while aligning on the plate prior to anaphase onset. During

anaphase, acentric sisters that are oriented parallel separate by sliding, while those oriented

perpendicularly separate via unzipping [Vicars et al., 2021].

Anti-parallel alignment of sister acentrics

To more precisely determine acentric orientation during metaphase alignment, neuroblasts expressing inducible I-CreI, the histone marker H2Av-RFP, and the telomere marker HOAP-GFP [Cenci et al., 2003] were imaged live. Marked telomeres enabled us to determine the orientation of paired sister acentrics with respect to one another as they aligned on the metaphase plate. A surprising outcome of this analysis is that a low but notable frequency of sister acentrics exhibited anti-parallel alignment. As expected, 100% (25/25) of intact chromosomes have telomeres paired and aligned (Figure 4). However, in 12% (4/34) of paired sister acentrics, the telomeres are oriented in opposing directions, with the remaining 88% (30/34) exhibiting normal telomere-to-telomere alignment. This implies that stable pairing of sister chromatids is possible despite the absence of a gene-by-gene alignment. These findings indicate the kinetochore plays a key role in ensuring the fidelity of sister chromosome alignment in *Drosophila*. Additionally, we find acentrics are oriented with their telomeres facing the periphery of the mass of congressed chromosomes (N=10, Figure 4C & 4D).

Acentric alignment requires microtubule plus-ends

Previous studies in *Drosophila* reported the microtubule-stabilizing protein Map205 and the microtubule plus-end associated protein EB1 are required for acentric sister separation during anaphase initiation [Vicars et al., 2021]. To determine if acentric alignment on the metaphase plate relies on microtubule plus-ends, we re-analyze published data of cells expressing RNAi against the microtubule plus-end tracking protein EB1 [Vicars et al., 2021]. Our analysis reveals 27% (8/30) of acentric sisters unable to align with kinetochore-bearing chromosomes (Figure 5). In control wild-type cells expressing only I-CreI, only 5% of acentric sisters fail to congress to the metaphase plate (Figure 5).

A separate study demonstrated that peripheral interpolar microtubules play a critical role

in the poleward segregation of acentric chromosome fragments [Karg et al., 2017]. Disruption of

interpolar microtubule organization via knockdown of Klp3a (kinesin-4), disrupts the poleward

segregation of the acentric, but not intact chromosomes. Additional laser-ablation experiments

demonstrated that segregating acentrics are physically connected to the interpolar microtubules

[Karg et al., 2017]. To determine if interpolar microtubules are involved in acentric movement to

and alignment on the metaphase plate, we re-analyzed previously published acentric movies in

a *klp3a* knockdown. *Drosophila* neuroblasts homozygous for hypomorphic alleles of *klp3a*,

exhibited defects in congression of the acentrics, but not the intact chromosomes (Figure 6B &

6C). In I-CreI expressing wild-type and *klp3a* mutant neuroblasts, 5% (1/18) and 30% (6/20) of

acentrics fail to align with kinetochore-bearing chromosomes at the metaphase plate

respectively (Figure 6C).

Examination of chromosome dynamics on monopolar spindles provides a means to

readily determine the extent to which microtubule plus-end forces act on chromosomes [Muscat

et al., 2015]. Unfortunately, we discovered the small molecule inhibitors commonly used to

induce monopolar spindles do not work in *Drosophila*. However, we find that RNAi knockdown

of the chromokinesin Klp31E (kinesin 4) in *Drosophila* neuroblasts resulted in monopolar spindle

formation. In these *Drosophila* neuroblasts with monopolar spindles, the acentric chromosomes,

as well as the intact chromosomes, are pushed to the cell periphery (Figure 7). This finding

indicates acentric chromosomes can be driven to microtubule plus ends prior to the metaphase-

to-anaphase transition without requiring kinetochores. It remains to be determined if the

acentrics are directly interacting with microtubule plus ends or utilizing motor proteins to travel to

the microtubule plus ends.

Surprisingly, despite the presence of monopolar spindles when Klp31E is partially

knocked down, neuroblasts form an acentrosomal bipolar spindle allowing cells to successfully

divide. 92% (14/15) of acentrics are able to align on the metaphase plate, with 80% (12/15) of

acentric sisters successfully separating and segregating from one another and moving to

opposite cell poles (Figure 7C & 7D).

Discussion
242
Taking advantage of our ability to fluorescently tag the *Drosophila* X-chromosome

acentric, we define the forces acting on the chromosome arms and the role of the kinetochore in

chromosome congression. These studies revealed that during the late stages of congression,

acentrics move faster than their neighboring kinetochore-bearing chromosomes. This indicates

that kinetochore microtubule interactions act as a brake during congression slowing the action

of plus-end directed forces acting on chromosome arms. In a similar fashion, the minus-end

directed kinesin, Ncd (kinesin-14), acts as a brake on the action of the plus-end directed

KLP61F (kinesin-5) in sliding anti-parallel microtubules during spindle pole separation [Tao et

al., 2006]. Thus, the kinetochore serves to modulate and reduce the motion of the

chromosomes during congression and alignment on the metaphase plate.

Our analysis also demonstrates that in contrast to intact chromosomes, the paired sister

acentrics oscillate during the late stages of congression and upon arrival at the metaphase

plate. The most likely explanation is that in the absence of a kinetochore, polar ejection forces

unevenly distributed along the length of the chromosome generate oscillating movements. Thus,

the kinetochore serves to stabilize chromosome orientation during alignment on the metaphase

plate. As a result of these oscillations, sister acentrics orient either parallel or perpendicular to

the spindle at anaphase onset. The former orientation results in acentric sister separation by

sliding past one another while the later results in acentrics separating via "unzipping" with

separation starting at one end of the chromosome [Vicars et al., 2021]

We also find that acentric sisters are positioned at the periphery of the metaphase plate

and on a plane distinct from the rest of the chromosome complement. Previous studies

demonstrated that the peripheral interpolar microtubules play a key role in acentric segregation

during anaphase [Karg et al., 2017]. The positioning of the acentrics at the edge of the

metaphase plate, and its association with interpolar microtubules, indicates that congression

and alignment of the acentric chromosomes relies on interpolar microtubules. This conclusion is

supported by the finding that knockdown of Klp3a, a kinesin required for establishing the

population of interpolar microtubules, preferentially disrupts congression of acentric

chromosomes [Karg et al., 2017]. *Drosophila* neuroblasts homozygous for hypomorphic alleles

of *klp3a*, exhibit a dramatic increase in acentric congression defects (Figure 6). Since Klp3a is a

chromokinesin, it is also possible that it interacts directly with acentrics propelling them along

interpolar microtubules to the metaphase plate. Our finding that acentrics oscillate argues

against this as direct lateral interactions between the acentric and microtubules would inhibit

oscillation. In addition, the fact that the acentrics move faster than intact chromosomes suggests

that they are not restricted by direct interactions with motor proteins. Acentrics may

preferentially rely on interpolar microtubules due to the fact that kinetochore microtubules

maintain intact chromosomes in a limited central region of the spindle. Lacking a kinetochore,

acentrics are shunted to the interpolar microtubules.

280   Our studies also demonstrate that polar ejection forces play significant roles in acentric

congression. This is most dramatically illustrated by the fact that reduction of EB1 activity

preferentially disrupts alignment of the acentric on the metaphase plate but not the intact

chromosomes. EB1 has been found to be essential for generating anti-polar forces on

chromosomes and stabilizing microtubules [Tirnauer & Bierer, 2000; Vitre et al., 2008]. This

finding is also supported by the finding that in monopolar spindles acentrics are rapidly pushed

away from the poles. As described above, the oscillation of acentrics as they align at metaphase

is readily explained by distributed plus-end directed forces acting along the chromosome arms.

288   One of the most surprising findings of this study emerges from live imaging paired sister

acentrics with GFP-labeled telomeres. The labeled telomeres enable us to determine the

relative orientation of each sister chromatid in the acentric pair as they align to the metaphase

plate. As expected, intact sister chromatids exhibit parallel alignment across the entire length of

the chromosome. This gene-for-gene alignment of paired sisters likely facilities their stable

cohesion. In addition, it is essential for successful mitotic recombination [Lisby & Rothstein,

2015; Cohen-Fix, 2001]. Surprisingly, we find that a small but significant fraction of sister

acentrics align in an anti-parallel orientation, indicating that the kinetochore is required to

prevent antiparallel alignment of sister chromatids. The mechanism by which the kinetochore

facilitates parallel alignment remains unknown as it is likely established during interphase

immediately following completion of S-phase [Skibbens, 2008].

Another unexpected finding from this analysis is that the presence of the acentric

chromosome fragment induces a global reorganization of congressed chromosomes. In wild-

type *Drosophila* neuroblasts, fully congressed chromosomes are roughly organized as a column

at metaphase. However, in the presence of the acentric chromosome fragment, the

chromosomes arrange themselves in a ring-like configuration. With a lack of chromatin in the

center of the ring the congressed chromosomes form a torus. Whether it is the presence of the

acentric or damaged DNA that causes the global reorganization of the congressed chromosome

remains unclear. Interestingly, this torus configuration of chromosomes at metaphase is well-

documented in human cells [Paulson et al., 2021]. 3D-reconstructions of human metaphase

chromosomes reveal their arrangement in a circular shape with kinetochores organized around

the center and chromosome arms pushed towards the cell periphery [Paulson et al., 2021]. It

should be noted, that human cancer cell lines often contain damaged DNA [Lewis & Golsteyn,

2016; Valencia-González et al., 2019]. As with acentric induction in *Drosophila*, the presence of

DNA damage may induce the toroidal shape of the chromosome mass at metaphase.

Studies in other species reveal distinct mechanisms can result in congressed

chromosomes forming a torus. For example, a similar torus configuration of congressed

chromosomes has been observed in diatoms. In these species, interpolar microtubules gather in

the center of the cell at metaphase, connecting spindle poles, and push chromosomes to either

side of the interpolar microtubule bundle [Pickett-Heaps et al., 1975]. Additionally, in

prometaphase newt pneumocytes, bundles of keratin filaments push chromosomes to the cell

periphery [Hayden et al., 1990]. However, in *Drosophila* neuroblasts expressing I-CreI, the

mitotic spindle is arranged with microtubules connecting to congressed chromosomes in the

torus formation and an absence of microtubules in the center of the torus (Supplemental Figure

3). Whether the torus configuration has functional consequences or is simply an outcome of the

mechanisms by which the spindle is formed is unknown. Taken together, these data support a

model in which acentrics congress to and align on the metaphase plate via plus-end microtubule

interactions stabilized by outer pole-to-pole microtubules and microtubule-associated proteins.

326
Materials and methods

Fly stocks

All stocks were raised on standard *Drosophila* media at room temperature (20–22˚C) as

previously described [Royou et al., 2008]. For generating acentrics, a transgenic fly line bearing

the I-CreI endonuclease under heat-shock 70 promoter were kindly provided by Kent Golic at

The University of Utah.

Live analysis of acentric behavior in *Drosophila* third instar neuroblasts

As previously described, acentric chromosome fragments were induced by I-CreI expression

(under heat shock 70 promoter) in 3rd instar larvae by a 1-hour 37˚C heat shock followed by a

1-hour recovery period at room temperature [Royou et al., 2010]. The larval brains from third

instar larvae were then dissected in PBS and placed on a slide with 20 µl of PBS. A coverslip

was placed on the brain in PBS and the excess PBS was wicked out from the edges of the

coverslip to induce squashing of brain between the slide and coverslip. Then, the edge of

coverslip was sealed with halocarbon and was imaged for live analysis as described below.

Neuroblast divisions in images for Figures 1-4, 6 and 7, Table 1, and Supplemental Figure 1

were from male 3rd instar larvae. Neuroblast divisions in images for Figure 5 and Supplemental

Figure 2 were from female 3rd instar larvae.

Microscopy and image acquisition

Wide-field microscopy. Time-lapse imaging for Figures 5B and 5C was performed using a

Leica DM16000B wide-field inverted microscope equipped with a Hamamatsu electron-

multiplying charge coupled device camera (ORCA 9100–02) with a binning of 1 and a

100x Plan-Apochromat objective with NA 1.4. Successive time points were filmed at 20 second

intervals. RFP (585 nm) and GFP (508 nm) fluorophores were imaged. Samples were imaged in

PBS and at room temperature (20–22˚C). Widefield images were acquired with Leica

Application Suite Advanced Fluorescence Software and 3D deconvolved using AutoQuant

X2.2.0 software.

Spinning-disk microscopy. Images in Figures 1-4, 5A, 6, and 7, and Supplemental Figures 1

and 2 were acquired with an inverted Nikon Eclipse TE2000-E spinning disk (CSLI-X1) confocal

microscope equipped with a Hamamatsu electron-multiplying charge coupled device camera

(ImageEM X2) with a 100X 1.4 NA oil-immersion objective. Samples were imaged in PBS and

at room temperature (20–22˚C). Images were acquired with MicroManager 1.4 software. Time-

lapse fluorescent images of neuroblasts divisions were done with 120 and 100 ms exposures for

GFP and RFP respectively with 0.5 µm Z-steps. Time-lapse videos with both GFP and RFP

were done every 5 seconds and time-lapse movies with RFP alone were done every 5 seconds.

Figures were assembled in Adobe Illustrator. Selected stills (both experimental and control)

were processed with ImageJ (http://rsb.info.nih.gov/ij/)).

Measurements

In Figure 2, circularity measurements of chromosomes (H2Av-RFP) were done using the

circularity function in ImageJ of the region outlined around acentrics and the main mass of

chromosomes. For statistical analyses, two-sided Mann-Whitney U tests were used. Two-sided

Mann-Whitney U tests were performed in R (R Core Team) and Prism Version 8 (GraphPad

Software). 3D renderings in Figures 2 and 4 and Supplemental Figure 2 were created using

Imaris software.

References
375
376 1. Bajer A. Cine-micrographic studies on chromosome movements in β-irradiated cell.
377   Chromosoma.1958; 9:319–331. PMID: 13608834.
378
379 2. Barisic, M, Aguiar, P, Geley, S, Maiato, H. Kinetochore motors drive congression of
380   peripheral polar chromosomes by overcoming random arm-ejection forces. Nat Cell Biol.
381   2014; 16:1249–1256. https://doi.org/10.1038/ncb3060
382
383 3. Bretscher HS, Fox DT. Proliferation of Double Strand Break Resistant Polyploid Cells
384   Requires Drosophila FANCD2. Dev Cell. 2016; 37(5):444–57.
385   https://doi.org/10.1016/j.devcel.2016.05.004 PMID: 27270041; PubMed Central PMCID:
386   PMC4901310.
387
388 4. Carlson JG Mitotic behavior of induced chromosomal fragments lacking spindle
389   attachments in the neuroblasts of the grasshopper. Proc Natl Acad Sci USA. 1938a;
390   24:500–507. https://doi.org/10.1073/pnas.24.11.500
391
392 5. Cenci G, Siriaco G, Raffa GD, Kellum R, Gatti M. The Drosophila HOAP protein is
393   required for telomere capping. Nat Cell Biol. 2003; 5(1):82–4.
394   https://doi.org/10.1038/ncb902 PMID: 12510197.
395
396 6. Chmatal L., Yang K., Schultz R.M., Lampson M.A. Spatial Regulation of Kinetochore
397   Microtubule Attachments by Destabilization at Spindle Poles in Meiosis I. Curr.
398   Biol. 2015; 25:1835–1841. doi: 10.1016/j.cub.2015.05.013.
399
400 7. Cohen-Fix O. The making and breaking of sister chromatid cohesion. Cell. 2001 Jul
401   27;106(2):137-40. doi: 10.1016/s0092-8674(01)00439-1. PMID: 11511341.
402
403 8. Fenech M, Kirsch-Volders M, Natarajan AT, Surralles J, Crott JW, Parry J, et al.
404   Molecular mechanisms of micronucleus, nucleoplasmic bridge and nuclear bud
405   formation in mammalian and human cells. Mutagenesis. 2011; 26(1):125–32.
406   https://doi.org/10.1093/mutage/geq052 PMID: 21164193
407
408 9. Golic MM, Golic KG. A simple and rapid method for constructing ring-X chromosomes in
409   Drosophila melanogaster. Chromosoma. 2011; 120(2):159–64.
410   https://doi.org/10.1007/s00412-010-0297-2 PMID:21085980; PubMed Central PMCID:
411   PMC4454366.
412
413 10. Guacci V, Koshland D, Strunnikov A. A Direct Link between Sister Chromatid Cohesion
414   and Chromosome Condensation Revealed through the Analysis of MCD1 in S.

cerevisiae. Cell. 1997; 91(1):47–57. https://doi.org/10.1016/s0092-8674(01)80008-8 PMID: 9335334; PubMed Central PMCID: PMC2670185.

11. Hayden JH, Bowser SS, Rieder CL. Kinetochores capture astral microtubules during chromosome attachment to the mitotic spindle: direct visualization in live newt lung cells. J Cell Biol. 1990 Sep;111(3):1039-45. doi: 10.1083/jcb.111.3.1039. PMID: 2391359; PMCID: PMC2116290.

12. Ishii K, Ogiyama Y, Chikashige Y, Soejima S, Masuda F, Kakuma T, et al. Heterochromatin Integrity Affects Chromosome Reorganization After Centromere Dysfunction. Science. 2008; 321(5892):1088–91. https://doi.org/10.1126/science.1158699 PMID: 18719285.

13. Kanda T, Otter M, Wahl GM. Mitotic segregation of viral and cellular acentric extrachromosomal molecules by chromosome tethering. J Cell Sci. 2001; 114:49–58. PMID: 11112689.

14. Kanda T, Sullivan KF, Wahl GM Histone-GFP fusion protein enables sensitive analysis of chromosome dynamics in living mammalian cells. Curr Biol. 1998; 8:377–385. https://doi.org/10.1016/s0960-9882(98)70156-3

15. Kanda T, Wahl GM. The dynamics of acentric chromosomes in cancer cells revealed by GFP- based chromosome labeling strategies. J Cell Biochem Suppl. 2000; Suppl 35:107–14. https://doi.org/10.1002/1097-4644(2000)79:35+<107::aid-jcb1133>3.0.co;2-y PMID: 11389539.

16. Kapoor T.M., Lampson M.A., Hergert P., Cameron L., Cimini D., Salmon E.D., McEwen B.F., Khodjakov A. Chromosomes can congress to the metaphase plate before biorientation. Science. 2006;311:388–391. doi: 10.1126/science.1122142.

17. Karg T, Elting MW, Vicars H, Dumont S, Sullivan W. The chromokinesin Klp3a and microtubules facilitate acentric chromosome segregation. J Cell Biol. 2017; 216(6):1597–608. https://doi.org/10.1083/jcb.201604079 PMID: 28500183; PubMed Central PMCID: PMC5461011.

18. Karg T, Warecki B, Sullivan W. Aurora B–mediated localized delays in nuclear envelope formation facilitate inclusion of late-segregating chromosome fragments. Mol Biol Cell. 2015; 26(12):2227–41. https://doi.org/10.1091/mbc.E15-01-0026 PMID: 25877868; PubMed Central PMCID: PMC4462941.

19. Kaye JA, Melo JA, Cheung SK, Vaze MB, Haber JE, Toczyski DP. DNA breaks promote genomic instability by impeding proper chromosome segregation. Curr Biol. 2004; 14:2096–2106. https://doi.org/10.1016/j.cub.2004.10.051

20. Khodjakov A, Rieder CL. Kinetochores moving away from their associated pole do not exert a significant pushing force on the chromosome. J Cell Biol. 1996; 135:315–327. https://doi.org/10.1083/jcb.135.2.315

21. Krenn V, Musacchio A. The Aurora B Kinase in Chromosome Bi-Orientation and Spindle Checkpoint Signaling. Front Oncol. 2015 Oct 16;5:225. doi: 10.3389/fonc.2015.00225. PMID: 26528436; PMCID: PMC4607871.

467   22. LaFountain JR, Oldenbourg R, Cole RW, Rieder CL. Microtubule Flux Mediates
468   Poleward Motion of Acentric Chromosome Fragments during Meiosis in Insect
469   Spermatocytes. Mol Biol Cell. 2001; 12 (12):4054–65.
470   https://doi.org/10.1091/mbc.12.12.4054 PMID: 11739800; PubMed Central PMCID:
471   PMC60775.
472
473   23. Lewis CW, Golsteyn RM. Cancer cells that survive checkpoint adaptation contain
474   micronuclei that harbor damaged DNA. Cell Cycle. 2016 Nov 16;15(22):3131-3145. doi:
475   10.1080/15384101.2016.1231287. Epub 2016 Sep 16. PMID: 27636097; PMCID:
476   PMC5134707.
477
478   24. Liang H, Wright WH, Cheng S, He W, Berns MW. Micromanipulation of chromosomes in
479   PTK2 cells using laser microsurgery (optical scalpel) in combination with laser-induced
480   optical force (optical tweezers). Experimental cell research. 1993; 204(1):110–20.
481   https://doi.org/10.1006/excr.1993.1015 PMID: 8416789.
482
483   25. Lisby M, Rothstein R. Cell biology of mitotic recombination. Cold Spring Harb Perspect
484   Biol. 2015 Mar 2;7(3):a016535. doi: 10.1101/cshperspect.a016535. PMID: 25731763;
485   PMCID: PMC4355273.
486
487   26. Maggert KA, Golic KG. Highly Efficient Sex Chromosome Interchanges Produced By I-
488   CreI Expression in Drosophila. Genetics. 2005; 171(3):1103–14.
489   https://doi.org/10.1534/genetics.104.040071 PMID:16020774; PubMed Central PMCID:
490   PMC1456814.
491
492   27. Maiato H, Gomes AM, Sousa F, Barisic M. Mechanisms of Chromosome Congression
493   during Mitosis. Biology (Basel). 2017 Feb 17;6(1):13. doi: 10.3390/biology6010013.
494   PMID: 28218637; PMCID: PMC5372006.
495
496   28. McEwen B.F., Chan G.K., Zubrowski B., Savoian M.S., Sauer M.T., Yen T.J. CENP-E is
497   essential for reliable bioriented spindle attachment, but chromosome alignment can be
498   achieved via redundant mechanisms in mammalian cells. Mol. Biol. Cell. 2001; 12:2776–
499   2789. doi: 10.1091/mbc.12.9.2776.
500
501   29. Michaelis C, Ciosk R, Nasmyth K. Cohesins: Chromosomal Proteins that Prevent
502   Premature Separation of Sister Chromatids. Cell. 1997; 91(1):35–45.
503   https://doi.org/10.1016/s0092-8674(01)80007-6 PMID: 9335333.
504
505   30. Muscat CC, Torre-Santiago KM, Tran MV, Powers JA, Wignall SM. Kinetochore-
506   independent chromosome segregation driven by lateral microtubule bundles. Elife. 2015
507   May 30;4:e06462. doi: 10.7554/eLife.06462. PMID: 26026148; PMCID: PMC4481507.
508
509   31. Navarro AP, Cheeseman IM. Kinetochore assembly throughout the cell cycle. Semin
510   Cell Dev Biol. 2021 Sep; 117:62-74. doi: 10.1016/j.semcdb.2021.03.008. Epub 2021 Mar
511   19. PMID: 33753005; PMCID: PMC8384650.
512
513   32. Ohno Y, Ogiyama Y, Kubota Y, Kubo T, Ishii K. Acentric chromosome ends are prone to
514   fusion with functional chromosome ends through a homology-directed rearrangement.
515   Nucleic Acids Res. 2016; 44(1):232–44. https://doi.org/10.1093/nar/gkv997 PMID:
516   26433224; PubMed Central PMCID: PMC4705696.
517

33. Paredes S, Maggert KA. Ribosomal DNA contributes to global chromatin regulation. Proc Natl Acad Sci USA. 2009; 106(42):17829–34. https://doi.org/10.1073/pnas.0906811106 PMID: 19822756; PubMed Central PMCID: PMC2764911.

34. Paulson JR, Hudson DF, Cisneros-Soberanis F, Earnshaw WC. Mitotic chromosomes. Semin Cell Dev Biol. 2021 Sep;117:7-29. doi: 10.1016/j.semcdb.2021.03.014. Epub 2021 Apr 6. PMID: 33836947; PMCID: PMC8406421.

35. Pickett-Heaps JD, McDonald KL, Tippit DH. Cell division in the pennate diatom Diatom vulgare. Protoplasma. 1975; 86, 205-242.

36. Platero JS, Ahmad K, Henikoff S. A Distal Heterochromatic Block Displays Centromeric Activity When Detached from a Natural Centromere. Molecular Cell. 1999; 4(6):995–1004. https://doi.org/10.1016/s1097-2765(00)80228-2 PMID: 10635324.

37. Porter ACG, Farr CJ. Topoisomerase II: untangling its contribution at the centromere. Chromosome Res. 2004; 12(6):569–83. https://doi.org/10.1023/B:CHRO.0000036608.91085.d1 PMID: 15289664.

38. Rieder CL, Davison EA, Jensen LC, Cassimeris L, Salmon ED. Oscillatory movements of monooriented chromosomes and their position relative to the spindle pole result from the ejection properties of the aster and half-spindle. J Cell Biol. 1986 Aug;103(2):581-91. doi: 10.1083/jcb.103.2.581. PMID: 3733881; PMCID: PMC2113830.

39. Rong YS, Titen SW, Xie HB, Golic MM, Bastiani M, Bandyopadhyay P, et al. Targeted mutagenesis by homologous recombination in D. melanogaster. Genes Dev. 2002; 16(12):1568–81. https://doi.org/10.1101/gad.986602 PMID: 12080094; PubMed Central PMCID: PMC186348.

40. Royou A, Gagou ME, Karess R, Sullivan W. BubR1- and Polo-Coated DNA Tethers Facilitate Poleward Segregation of Acentric Chromatids. Cell. 2010; 140(2):235–45. https://doi.org/10.1016/j.cell.2009.12.043 PMID: 20141837; PubMed Central PMCID: PMC2969851.

41. Royou A, McCusker D, Kellogg DR, Sullivan W. Grapes (Chk1) prevents nuclear CDK1 activation by delaying cyclin B nuclear accumulation. J Cell Biol. 2008; 183(1):63–75. https://doi.org/10.1083/jcb.200801153 PMID: 18824564; PubMed Central PMCID: PMC2557043.

42. Skibbens RV. Mechanisms of sister chromatid pairing. Int Rev Cell Mol Biol. 2008;269:283-339. doi: 10.1016/S1937-6448(08)01005-8. PMID: 18779060.

43. Sural TH, Peng S, Li B, Workman JL, Park PJ, Kuroda MI. The MSL3 chromodomain directs a key targeting step for dosage compensation of the Drosophila melanogaster X chromosome. Nat Struct Mol Biol. 2008 Dec;15(12):1318-25. doi: 10.1038/nsmb.1520. Epub 2008 Nov 23. PMID: 19029895; PMCID: PMC2636508.

44. Tanaka TU. Bi-orienting chromosomes: acrobatics on the mitotic spindle. Chromosoma. 2008 Dec;117(6):521-33. doi: 10.1007/s00412-008-0173-5. Epub 2008 Aug 2. PMID:

18677502.

45. Tao L, Mogilner A, Civelekoglu-Scholey G, Wollman R, Evans J, Stahlberg H, Scholey
JM. A homotetrameric kinesin-5, KLP61F, bundles microtubules and antagonizes Ncd in
motility assays. Curr Biol. 2006 Dec 5;16(23):2293-302. doi: 10.1016/j.cub.2006.09.064.
PMID: 17141610.

46. Tirnauer JS, Bierer BE. EB1 proteins regulate microtubule dynamics, cell polarity, and
chromosome stability. J Cell Biol. 2000; 149(4):761–6.
https://doi.org/10.1083/jcb.149.4.761 PMID: 10811817; PubMed Central PMCID:
PMC2174556.

47. Valencia-González HA, Ruíz G, Ortiz-Sánchez E, García-Carrancá A. Cancer Stem
Cells from Tumor Cell Lines Activate the DNA Damage Response Pathway after Ionizing
Radiation More Efficiently Than Noncancer Stem Cells. Stem Cells Int. 2019 Apr
3;2019:7038953. doi: 10.1155/2019/7038953. PMID: 31073313; PMCID: PMC6470433.

48. Vicars H, Karg T, Warecki B, Bast I, Sullivan W. Kinetochore-independent mechanisms
of sister chromosome separation. PLOS Genetics. 2021; 17(1): e1009304.
doi:10.1371/journal.pgen.1009304 PMID: 33513180

49. Vitre B, Coquelle FM, Heichette C, Garnier C, Chre´tien D, Arnal I. EB1 regulates
microtubule dynamics and tubulin sheet closure in vitro. Nat Cell Biol. 2008; 10(4):415–
21. https://doi.org/10.1038/ncb1703 PMID: 18364701.

50. Warecki B, Sullivan W. Micronuclei formation is prevented by Aurora B-mediated
exclusion of HP1a from late-segregating chromatin in Drosophila. Genetics. 2018;
210(1):171–187. https://doi.org/10.1534/genetics.118.301031 PMID: 29986897; PubMed
Central PMCID: PMC6116970.

51. Warecki B, Sullivan W. Mechanisms driving acentric chromosome transmission.
Chromosome Res. 2020; 28(3-4):229-246. https://doi.org/10.1007/s10577-020-09636-z
PMID: 32712740.

52. Ye A.A., Deretic J., Hoel C.M., Hinman A.W., Cimini D., Welburn J.P., Maresca T.J.
Aurora A Kinase Contributes to a Pole-Based Error Correction Pathway. Curr.
Biol. 2015; 25:1842–1851. doi: 10.1016/j.cub.2015.06.021.

53. Yen T.J., Li G., Schaar B.T., Szilak I., Cleveland D.W. CENP-E is a putative kinetochore
motor that accumulates just before mitosis. Nature. 1992;359:536–539.
doi: 10.1038/359536a0.

Figure Legends

Table 1: Average velocities of chromosomes at the metaphase plate. (A) Average velocities of individual chromosomes as they align at the metaphase plate. Velocity is measured in nanometers per second. Asterisks indicate statistical significance (*P=0.005, **P=0.01) as determined by a two-sided Mann-Whitney U test. (B) Schematic showing how chromosome velocity is determined. Velocity is defined as beginning 80 seconds prior to anaphase onset and ends one frame (5 s) prior to anaphase onset.

Figure 1: Acentrics align on the metaphase plate. (A) Still frames of a time-lapse movie of a mitotic neuroblast labeled with H2Av-RFP (magenta) and MSL3-GFP (green) not expressing I-CreI. (B) Still frames of a time-lapse movie of a mitotic neuroblast with I-CreI induced acentrics. Sister acentrics (white arrowheads) undergo movement at the metaphase plate while aligning (yellow brackets), lag on the spindle equator but eventually separate, segregate, and are incorporated into daughter nuclei. Bars, 2 µm. Time in seconds.

Figure 2: Acentric induction induces a global reorganization of congressed chromosomes such that the acentric localizes to the periphery of the main chromosome mass (A) Still images of a 3D rendering of a neuroblast at metaphase labeled with H2Av-RFP (magenta) and an intact MSL3-GFP (green) tagged X chromosome (I-CreI not induced). (B) Still frames of a 3D rendering of a neuroblast at metaphase with I-CreI induced acentrics. This results in a dramatic global reorganization of the congressed chromosomes into a ring with an absence of chromatin in the center. The acentric localizes to the outer most edge of the chromosome ring. Bars, 2 µm. Images are rotated 180°. (C) Box plot showing the distances of X chromosomes from the chromosome mass center. Distances are measured in µm. Acentric X chromosomes are positioned significantly farther from the chromosome mass center when compared to the intact X chromosome (*P=0.01, Mann-Whitney Test).

Figure 3: Acentrics undergo varied orientations during metaphase (A) Still frames of a time-lapse movie of a mitotic neuroblast labeled with H2Av-RFP (magenta) and MSL3-GFP (green), not expressing I-CreI, during metaphase. Zoomed in images show only the X chromosomes marked by MSL3-GFP (gray). (B) Still frames of a time-lapse movie of a mitotic neuroblast with I-CreI induced acentrics during metaphase. Zoomed in images show only the X chromosomes marked by MSL3-GFP (gray). Bars, 2 µm. Time in seconds. (C) Line graph showing the change in orientation of acentric (green) and intact (black) X chromosomes over time, prior to anaphase onset. Data points every 5 seconds. Shaded regions represent one- and two-times the standard deviation.

Figure 4: Acentrics align at the metaphase plate with telomeres either paired or unpaired. (A) Still frames of a time-lapse movie of a mitotic neuroblast labeled with H2AvRFP (magenta) and HOAP-GFP (green) expressing I-CreI with acentric sister telomeres aligned and paired (arrowheads and yellow outline) at metaphase. (B) Still frames of a time-lapse movie of a mitotic neuroblast expressing I-CreI with acentric sister telomeres opposed (arrowheads and yellow outline) at metaphase. Most acentrics are oriented with their telomeres paired during metaphase. Bars, 2 µm. Time in seconds. (C) Still images of a 3D rendering of a mitotic neuroblast with H2Av-RFP (magenta) and HOAP-GFP (green) (N=5). (D) Still images of a 3D rendering of a mitotic neuroblast with I-CreI induced chromosome fragments. All chromosomes are arranged in a semi-circular formation with distally oriented telomeres immediately prior to anaphase onset (N=10). Bars, 2 µm. Images are rotated 180°. (E) Schematic showing frequency of sister acentrics with telomeres paired vs. telomeres opposing.

Figure 5: EB1 required for acentric metaphase plate alignment. (A) Still frames of a time-lapse movie of a mitotic neuroblast labeled with H2Av-RFP (gray) with I-CreI induced acentrics (white arrowheads). (B) Still frames of a time-lapse movie of a mitotic neuroblast expressing I-CreI and EB1 RNAi. Sister acentrics fail to align with the chromosome mass at metaphase and segregate to the same cell pole. Bars, 2 µm. Time in seconds. (C) Bar graph showing congression failure rates. Successful chromosome congression is in gray. Failure of chromosomes to congress is in magenta.

Figure 6: The chromokinesin Klp3a is required for aligning acentrics to the metaphase plate. (A) Still images from a time-lapse movie of a neuroblast expressing I-CreI alone (control). (B) Still images from a time-lapse movie of a neuroblast expressing I-CreI in *klp3a* mutant background. Acentric sisters (white arrows) are positioned near the spindle pole at metaphase. Bars, 2 μm. Time in seconds. (C) Bar graph showing the rate of chromosome alignment at the metaphase plate. Successful chromosome alignment at the metaphase plate is in gray. Failure of chromosomes to align is in magenta.

Figure 7: Acentrics are pushed by microtubule plus-ends (A) Still frames of a time-lapse movie of a mitotic neuroblast expressing H2AvRFP (magenta) and Jupiter-GFP-labeled microtubules (green) with I-CreI induced acentrics. Sister acentrics (white arrowheads) align on the metaphase plate and lag behind at the spindle equator but eventually separate, segregate, and are incorporated into daughter nuclei. (B) Still frames of a time-lapse movie of a mitotic neuroblast with I-CreI induced acentrics (white arrowheads) and expressing Klp31E RNAi. All chromosomes are pushed to the cell periphery by the mono-polar mitotic spindle. Bars, 2 μm. Time in seconds. (C) Percentages of acentric sisters that fail to congress to the metaphase plate. (D) Percentages of acentric sisters that fail to separate from one another during late anaphase.

Figure 8: Schematic showing proposed mechanism of acentric chromosome congression. Chromosomes in magenta, microtubules in green, dynein in purple, chromokinesins in orange, CENP-E in blue, and EB1 in yellow. (A) Normal chromosomes congress to the metaphase plate either through direct congression or peripheral congression. Direct congression relies on chromokinesins localized at chromosome arms and end-on kinetochore-microtubule attachments. Peripheral congression involves chromosomes initially moving to the spindle pole by the minus-end motor protein dynein. Once at the pole, the chromosomes engage chromokinesins and the kinetochore-associated kinesin CENP-E for transport to the metaphase plate. (B) Acentric chromosome fragments congress by using EB1-driven microtubule plus-end pushing forces and lateral attachments with microtubules (potentially via chromokinesins).

Table 1

	Avg velocity ± SD (nm/s)	N (individual chromosomes)
Intact X chromosomes	8.4 ± 3.3	12
Non-X chromosomes	8.5 ± 2.4	28
Acentric X chromosomes	11.7 ± 3.5	14

B

80 seconds   5 seconds

Time prior to anaphase onset

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A I-CreI — 0 s, 30, 40, 50, 100 — H2Av-RFP

B EB1 RNAi, I-CreI — 0 s, 660, 720, 780, 920 — H2Av-RFP

C

Failure to congress
Successful congression

% of acentric sisters

Control I-CreI Alone: 5%, 95%, N=18
I-CreI, EB1 RNAi: 27%, 73%, N=30

Figure 6

A I-CreI — 0 sec, 50, 95, 140, 250 — H2Av-RFP

B I-CreI, *klp3a²* — 0 sec, 120, 155, 275, 320

C

Failure to align
Successful alignment

% of acentric sisters

Control I-CreI alone: 5%, 95%, N=18
I-CreI, *klp3a²*: 30%, 70%, N=21

Figure 7

Figure 8

A   Normal chromosome congression

B   Acentric chromosome congression

Dynein

Chromokinesins

CENP-E

EB1