Prior Fc Receptor activation primes macrophage for increased sensitivity to IgG via long term and short term mechanisms

Annalise Bond[1], Sareen Fiaz[1], Kirstin R Rollins[1], Jazz Elaiza Q Nario[1], Samuel J Rosen[1], Alyssa Granados[1], Maxwell Z Wilson[1], Meghan A Morrissey[1,2]*

[1] Molecular Cellular and Developmental Biology Department, University of California, Santa Barbara, Santa Barbara CA

[2] Lead contact
*morrissey@ucsb.edu

Keywords: macrophage, phagocytosis, Fc Receptor, IgG

Summary

Macrophages measure the 'eat-me' signal IgG to identify targets for phagocytosis. We wondered if prior encounters with IgG influence macrophage appetite. IgG is recognized by the Fc Receptor. To temporally control Fc Receptor activation, we engineered an Fc Receptor that is activated by light-induced oligomerization of Cry2, triggering phagocytosis. Using this tool, we demonstrate that Fc Receptor activation primes macrophages to be more sensitive to IgG in future encounters. Macrophages that have previously experienced Fc Receptor activation eat more IgG-bound cancer cells. Increased phagocytosis occurs by two discrete mechanisms – a short- and long-term priming. Long term priming requires new protein synthesis and Erk activity. Short term priming does not require new protein synthesis and correlates with an increase in Fc Receptor mobility. Our work demonstrates that IgG primes macrophages for increased phagocytosis, suggesting that therapeutic antibodies may become more effective after initial priming doses.

Introduction

Macrophages eat pathogens and infected, cancerous or dying cells via phagocytosis. To select targets for phagocytosis, macrophages measure 'eat-me' signals, like IgG antibodies. IgG is recognized by the Fc Receptor (FcR), which is phosphorylated and recruits the kinase Syk, triggering downstream signaling[1,2]. Therapeutic IgGs like Rituximab or Trastuzumab trigger Antibody-dependent Cellular Phagocytosis (ADCP) or Antibody-dependent Cellular Cytotoxicity (ADCC) to reduce cancer growth[2–7]. Even many antibodies originally designed to block the function of their target actually activate the FcR for full efficacy[8,9]. Given the therapeutic importance, there is substantial interest in understanding how to boost macrophage phagocytosis.

What affects macrophage appetite? One important parameter is how sensitive a macrophage is to 'eat-me' signals. Antibody-dependent phagocytosis requires the coordinated activation of a sufficient number of FcRs[10]. Targets with a subthreshold amount of IgG are not phagocytosed, despite triggering the initial steps in the phagocytosis signaling pathway[10]. In other macrophage signaling pathways, low levels of activating signal do not elicit any response on their own, but prime macrophages for rapid and intense response to future stimuli[11]. Whether subthreshold FcR signaling has any effect on macrophage appetite is not clear.

During an immune response or treatment with a therapeutic antibody, macrophages encounter multiple potential targets for phagocytosis sequentially, leading to bursts of FcR activation. Some encounters with antibody-opsonized cells result in phagocytosis of the entire cell[3,5], but many cells do not have sufficient antibodies to trigger phagocytosis. Instead macrophages may trogocytose, or nibble, a target cell or simply ignore it[12–15]. In some circumstances, prior phagocytosis increases macrophage appetite[16,17]. In contrast, other studies demonstrated that phagocytosing several whole cancer cells reduces macrophage appetite[18]. There is no clear, unifying model explaining these differences, which could be dependent on the specific 'eat-me' signal presented, the time since phagocytosis, the intensity of the 'eat-me' signal, digestion of the internalized particle, or any number of other factors[19].

To unravel how prior IgG exposure affects macrophage appetite, we need to precisely control the timing and intensity of activating specific phagocytic receptors. Delivering a temporally controlled, homogenous antibody stimuli to a population of cells is very difficult with the current tools. Because soluble IgG does not activate the FcR, IgG must be presented on antibody-bound targets. Controlling precisely the number of targets a macrophage encounters, or the timing of these encounters is very difficult and low-throughput.

To quantitatively control the duration and strength of FcR activation, we developed an optogenetic FcR (optoFcR). We found that prior FcR activation primes macrophages for greater responses to subsequent stimuli. Counterintuitively, low levels of optoFcR

activation induced stronger priming than high levels of optoFcR activation. Macrophage
priming is controlled by two independent mechanisms, one short-term (<1 hour) and
one long-term response (starting at 4 hours, and lasting up to 3 days). The short term
response is associated with an increase in FcR mobility, accelerated initiation of
phagocytosis and increased phagocytic cup formation. The long term response requires
activation of Erk to drive a transcriptional response. These data suggest that
macrophages can integrate signaling from previous encounters with IgG to modify the
response to the next target. This study provides insight into how macrophage appetite is
regulated, and may suggest strategies to enhance antibody-dependent cellular
phagocytosis.
81
82 **Results**
83
84 **Optogenetic FcR recapitulates native FcR signaling for precise temporal control**
85 **over signaling**
86
To precisely control the temporal pattern of FcR activation across an entire field of
cells, we sought to design an optogenetic FcR that could be turned on and off with light.
Prior work has shown that the FcR clusters upon IgG binding. Although the FcR has no
inherent kinase activity, clustering promotes phosphorylation of the intracellular
Immunoreceptor Tyrosine-based Activation Motifs (ITAMs) and phagocytosis[20–22]. We
hypothesized that clustering may be sufficient to induce FcR activation. To test this, we
designed an optoFcR construct that consists of a myristoylation sequence for
membrane localization, the intracellular domain for the native FcR gamma-chain and a
light activatable peptide CRY2 (Figure 1a). We selected the variant CRY2olig because it
forms multimeric, rapidly reversible clusters[23]. Upon blue light (450nm) stimulation the
optoFcR oligomerizes (Figure 1b; Video S1). Clusters dissociate when the cells are
returned to the dark (Figure 1b). Clustering also results in the recruitment of the
downstream effector protein Syk, suggesting that clustering is sufficient to induce FcR
phosphorylation (Figure 1c,d; Video S2).
We next sought to determine if clustering of the optoFcR is sufficient to trigger
phagocytosis. We incubated ICAM-1 conjugated beads with macrophages expressing
either the optoFcR or membrane tethered mCh (mCh-CAAX) and exposed them to 15
min of light stimulation (Figure 1e). ICAM-1 mediates bead binding to the macrophage,
but does not trigger phagocytosis of otherwise unopsonized beads[24,25]. Macrophages
expressing the optoFcR engulfed three times as many beads as control macrophages
when stimulated with the highest intensity light and twice as many beads when
stimulated with medium intensity light (Figure 1f,g; Figure S1; Video S3). Low intensity
light did not activate phagocytosis. This dose response is similar to the dose response
seen in IgG mediated phagocytosis (Figure S1). Together, these data demonstrate that

clustering of the FcR ITAM domain is sufficient to initiate phagocytosis in macrophages
without a specific 'eat-me' signal.
113
114 **Prior FcR activation enhances phagocytosis of IgG coated beads.**
115
With the ability to temporally control FcR activation in bone marrow-derived
macrophages, we next sought to determine if prior FcR activation influences phagocytic
ability. We stimulated macrophages with low intensity blue light to activate the optoFcR
for 15 minutes which is around the same timescale that a macrophage interacts with a
phagocytic target. We then waited 1 or 12 hours before adding IgG opsonized beads,
which is sufficient time for the optoFcR to completely decluster. Then we measured the
number of beads engulfed per cell using microscopy (Figure 2a). We found that prior
optoFcR activation increased the amount of eating roughly 2-fold compared to cells that
either did not receive prior light stimulation or did not express the optoFcR (Figure 2b).
This demonstrates that prior FcR activation primes macrophages to respond to future
IgG.
Next, we systematically varied the light intensity and stimulation time used to activate
the optoFcR as well as the delay between activating the optoFcR and measuring
phagocytosis of IgG coated beads. In all cases, we found that low doses of light, and
thus less FcR activity, led to the highest macrophage priming (Figure S2). The amount
of light that best primed macrophages was not sufficient to activate phagocytosis on its
own (Figure 1h). This suggests that a sub-threshold level of FcR activation, not
sufficient to activate phagocytosis, primes macrophages for future encounters with IgG.
We then wanted to know if we could induce macrophage priming using IgG and the
endogenous FcR. Based on our data from the optoFcR, we predicted that a low level of
IgG exposure would enhance phagocytosis of a second dose of IgG-coated beads. To
test this, we preincubated macrophages with beads containing an IgG density that did
not activate phagocytosis (Figure S1) or unopsonized beads. We then extensively
washed the macrophages to remove these beads and added a second color of IgG-
bound beads (Figure 2c). We found that macrophages preincubated with IgG-coated
beads phagocytosed more than macrophages pre-incubated with unopsonized beads
when the second dose was above the threshold for inducing phagocytosis (Figure 2d).
This suggests that the native IgG and FcR system primes macrophages like the
optoFcR system.
145
146 **Prior FcR activation specifically enhances phagocytosis of IgG coated beads, not**
147 **phosphatidylserine or unopsonized beads.**
148
We next wanted to know if prior FcR activation specifically primed macrophages to
phagocytose IgG-coated targets or if it broadly enhanced phagocytosis. We first

determined if prior optoFcR activation increased non-specific phagocytosis of
unopsonized targets. Primed macrophages did not phagocytose more unopsonized
beads than control macrophages (Figure 2e). We then determined if prior optoFcR
activation increased efferocytosis, the engulfment of apoptotic cells. The molecular
regulators of efferocytosis partially overlap with antibody-dependent phagocytosis, but
the processes require some unique signaling pathways including different 'eat-me'
signal receptors[26]. By integrating phosphatidylserine (PS), an efferocytic 'eat-me' signal,
into the lipid mixture on our silica bead targets, we can recapitulate apoptotic corpse
engulfment *in vitro*[27]. We incubated PS-coated beads with macrophages at 1 and 12
hours post light stimulation and measured the amount of eating. There was no change
in the amount of eating at either timepoint (Figure 2f). These data suggest that prior FcR
activation primes macrophages to specifically react to IgG. This also suggests that the
molecular regulators of priming are unique to FcR signaling, rather than in one of the
pathways shared by efferocytosis and antibody-dependent phagocytosis.
We then investigated if prior FcR stimulation changed macrophage sensitivity for IgG,
lowering the threshold of IgG required for initiating phagocytosis. Alternatively, priming
could increase macrophage capacity, the maximum number of targets each
macrophage can engulf. To do this, we added beads with various concentrations of IgG
to BMDMs and calculated the phagocytic index (Figure S2). Primed macrophages show
enhanced eating of beads with low concentration of IgG. The total capacity for
phagocytosis is not significantly changed in primed macrophages. This indicates that
priming primarily alters macrophage sensitivity to low levels of IgG, rather than overall
capacity for phagocytosis.
174
175 **Primed macrophages phagocytose more antibody opsonized cancer cells.**
176
We next sought to determine if primed macrophages can increase whole cell eating of
opsonized cancer cell targets. In addition to phagocytosis, macrophages often
trogocytose or nibble target cells, stripping the cancer cells of target antigen without
killing them[15]. Prior clinical studies have shown that the anti-CD20 therapeutic antibody
(rituximab) is most effective when administered more frequently at a low dose, which
minimizes antigen shaving[28]. Given our results, we hypothesized that prior FcR
activation might also enhance phagocytosis of cancer cells. We first measured
phagocytosis of Raji B cells incubated with increasing concentrations of anti-CD20
antibody to find an antibody concentration where we could detect a change in
macrophage sensitivity (Figure S3). To measure both the amount of whole cell
phagocytosis and trogocytosis, we incubated IgG opsonized Raji cell targets with
primed or unprimed optoFcR-expressing BMDMs and analyzed the cells with timelapse
microscopy (Figure 3a-b; Video S4-6). We found that the number of Raji cells
phagocytosed, and the percent of phagocytic macrophages increased in primed

macrophages (Figure 3c-d). The percentage of primed macrophages that trogocytosed
was not significantly increased compared to unprimed macrophages (Figure 3e). This
suggests that primed macrophages are better at phagocytosing antibody-opsonized
cancer cells.
195
196   **Macrophage priming occurs through a short-term and long-term mechanism.**
197
We next sought to determine the molecular mechanism for enhanced phagocytosis
after FcR activation. We observed enhanced phagocytosis at both 1 hour and 12 hours
after optoFcR stimulation (Figure 2b). While 12 hours post-stimulation is likely enough
time for changes in transcription or translation to affect macrophage phenotypes, 1 hour
is likely too short for this mechanism. We decided to carefully assay when macrophage
priming occurred. To do this we activated the optoFcR and varied the time before
presenting a second stimulus of IgG coated beads and measuring phagocytosis. We
saw robust priming occurring in two discrete waves: a short-term response that peaks
around 1 hour after FcR activation, and a long-term response that begins at 4 hours
after FcR activation and persists for at least 24 hours (Figure 4a).
As the long-term priming following optoFcR activation persists for at least 72 hours,
we speculated that this response requires *de novo* protein production rather than a
more transient post-translational modification mechanism. We evaluated priming in
macrophages treated with cycloheximide (CHX) and actinomycin D (AD) to inhibit
translation and transcription respectively. Treatment with either CHX or AD significantly
reduced phagocytosis in primed macrophages compared to DMSO treated control
macrophages at 4 and 6 hours post light stimulation (Figure 4b). This suggests that *de*
novo mRNA and protein synthesis is required for a long-term memory response.
Blocking new protein synthesis did not significantly reduce priming at 1 hour post
stimulation, suggesting that short-term memory is not reliant on new protein production
(Figure 4b). Overall this suggests that there are two mechanisms for macrophage
priming – one that operates on a short timescale and does not require synthesis of new
proteins, and one that operates on a long time scale and requires changes in gene
expression.
222
223   **Erk activation is required for long-term priming.**
224
Because long term priming requires new protein production, we sought to dissect
which transcriptional programs were being executed by the macrophages. Erk, a
nuclear kinase, functions downstream of the FcR and regulates the macrophage dose
dependent response to LPS as well as many other immune signaling pathways[11]. To
determine if Erk contributes to macrophage priming, we used PD0325901 to block Erk
activity. We then stimulated the optoFcR for 15 minutes, waited 1 or 12 hours, and
measured phagocytosis of IgG-coated beads. Inhibiting Erk signaling blocked long-term

memory with no effect on short-term memory (Figure 4c). These results indicate that
long-term priming requires a transcriptional response mediated by Erk activation.
234
235 **Initiation of engulfment proceeds faster and has a higher success rate in primed**
236 **macrophages.**
237
Having demonstrated that short-term priming does not require new protein synthesis
like long-term priming, we sought to determine the mechanism for short term priming.
To do this, we first wanted to isolate which step in the phagocytic process was
enhanced by prior sub-threshold activation of the FcR. We quantified the kinetics of
engulfment using live cell imaging, breaking the process of phagocytosis into three
steps: target binding, initiation of phagocytosis, and completion[21] (Figure 5a,b; Video
S7). We then quantified the time between each step in primed and unprimed
macrophages, and the percent of bead contacts that successfully progressed from one
step to the next without membrane retraction and target release. Primed macrophages
were more likely to initiate phagocytosis of bound beads, and the time between bead
binding and initiation was less (Figure 5c,d). In contrast, after initiation the chance of
successfully completing phagocytosis and the speed of phagocytosis were the same in
primed and unprimed macrophages (Figure 5e,f). Overall, the percent of bead contacts
that result in successful phagocytic events is significantly increased in primed
macrophages (Figure 5g). Stimulated control macrophages that do not express the
optoFcR did not show a difference in any measure of phagocytosis compared to
unstimulated macrophages (Figure S4). These data indicate that prior sub-threshold
FcR activation primes macrophages for faster target recognition and more frequent
signal initiation, implicating early phagocytic machinery.
257
258 **Priming is associated with enhanced FcR mobility.**
259
As initiation of phagocytosis is faster in primed macrophages, and our prior data
indicated that the molecular regulator of priming was specific to the FcR pathway, we
decided to look at FcR mobility. Clustering of IgG, and subsequently FcR, increases the
frequency and speed of initiating phagocytosis but not the speed of cup closure, similar
to the phenotype we observed in our phagocytosis kinetics analysis[21]. FcR cluster
formation and subsequent activation is dependent on the lateral mobility of FcRs, which
is constrained by a heterogeneous F-actin 'fence' and other mechanisms[29–32]. Increased
FcR mobility correlates with increased binding of IgG-coated targets and
phagocytosis[29,31]. We hypothesized that primed macrophages may have higher FcR
mobility, which could explain the increased speed and frequency of initiating
phagocytosis. To determine if optoFcR priming increases receptor mobility, we tracked
single FcR molecules on optoFcR primed and unprimed cells (Figure 6a). On average,
the FcR molecules on primed macrophages had a higher mean jump distance (MJD;

average distance traveled between frames) than on unprimed macrophages (Figure
6b). Graphing the MJD of individual tracks, primed macrophages had a multi-modal
distribution of track MJDs, suggesting there may be a more mobile population in the
primed macrophages (Figure 6c).
Previous studies have described FcR motion as free or confined[29,31]. Confined FcRs
have limited mobility and may not be available to form signaling microclusters required
for phagocytosis. To see if primed macrophages had more free FcRs, we categorized
FcR tracks by motion type using a moment scaling spectrum analysis. Consistent with
previous studies, we found a population of confined Fc Receptors that were restricted to
small microdomains or 'corals' within the membrane (Figure 6d,e). This population was
decreased in primed macrophages, suggesting that more FcRs may be available to join
signaling microclusters. In addition, the average diffusion coefficient, indicative of how
far each receptor travels, was significantly higher in primed macrophages (Figure 6f).
Overall, this data suggests that FcRs are more mobile on primed macrophages, which
may increase formation of FcR signaling clusters and the probability of initiating
phagocytosis.
289
290   **Discussion**
291
Our work suggests that the macrophage response to IgG integrates information from
sequential encounters with IgG coated targets. Prior FcR activation makes
macrophages more likely to phagocytose an IgG-coated target. This enhanced
phagocytosis correlates with an increase in FcR mobility in the short term. Long term
priming requires Erk signaling and transcription of new proteins.
Clustering of the FcR has previously been linked to its activation and enhanced
phagocytosis[21,22,33,34]. We recapitulate this clustering using an optogenetic method and
find that clustering of the optogenetic FcR is sufficient to drive Syk recruitment, which is
indicative of phosphorylation of the intracellular Immunoreceptor Tyrosine-based
Activation Motifs (ITAMs). The FcR has no inherent kinase activity, so why does
clustering promote receptor activation? ITAM phosphorylation is controlled by the
opposing actions of Src family kinases, which favor activation, and transmembrane
phosphatases like CD45 that deactivate the receptor[1]. Prior work has shown that
particle binding drives CD45 exclusion from the phagocytic synapse because the bulky
extracellular domain is sterically excluded from the tight membrane-membrane
interface[35-38]. This allows Src family kinases to dominate, tipping the kinase-
phosphatase balance in favor of FcR activation[39]. Our prior work has shown that FcR
clustering enhances ITAM phosphorylation independent of avidity effects that would
increase receptor binding to the target[21]. This may be mediated by the formation of lipid
ordered domains that segregate kinases and phosphatases, favoring immune receptor
activation[40,41]. The optoFcR is tethered to the membrane via a myristoylation motif,

favoring inclusion in lipid ordered domains, which may promote activation by Src family
kinases and segregation from CD45 without the need for particle binding. Alternatively,
the geometry of the clustered intracellular FcR domains may promote activation through
another mechanism.
In the hour following optoFcR activation, the mobility of native FcRs on the
macrophage surface increases. FcR mobility is constrained by the actin cytoskeleton
and actin-associated transmembrane pickets[29,31,32,42]. Syk kinase activation rearranges
the actin cytoskeleton allowing for less constrained FcR diffusion and greater mobility[31].
Since the optoFcR also recruits Syk, this may underlie the change in FcR mobility in our
system. Our working model is that increased FcR mobility increases the formation of
receptor clusters that promote initiating and continuing phagocytosis. This could
enhance macrophage sensitivity by enabling formation of signaling clusters with a lower
overall IgG density.
Prior work has shown a critical threshold of FcR activation is required for
macrophages to commit to phagocytosis[10]. FcR signaling that is below this threshold
activates the initial steps in the phagocytic signaling pathway, including recruitment of
the downstream effector kinase Syk[10]. Whether this low level of activation has any effect
on macrophages was not clear. Our data suggests that this low level of activation may
have a purpose in the macrophage, allowing the cell to prepare for future encounters
with IgG-bound targets.
Our data show that lower levels of optoFcR stimulation prime phagocytosis better than
high levels. Prior work has shown that low levels of TLR activation prime macrophages
for a rapid and strong response to future stimuli, without activating an inflammatory
response alone[11]. Our study suggests this may be true for the FcR pathway as well. The
effect of high levels of FcR activation may be different because it is associated with
receptor internalization, which can lead to decreased phagocytic capacity[18]. Previous
studies have shown that phagocytosing many antibody-coated cancer cells causes
macrophage 'hypophagia' or reduced phagocytosis. This suggests that the signaling
consequences of successful phagocytosis and sub-threshold FcR activation may be
quite different.
While the adaptive immune system is traditionally thought of as the source of
immunological memory, a growing body of evidence shows that the innate immune
system also remembers prior infections and threats. This is often called "trained
immunity" and occurs via epigenetic reprogramming of myeloid cells to increase or
decrease their transcriptional response to reinfection[16,43,44]. Trained immunity can
persist for years if myeloid progenitor cells are affected. Most of the work on trained
immunity has focused on a memory of pathogenic molecules or inflammatory cytokines.
Our work builds on this, suggesting that FcR activation also elicits a long term molecular
memory. In contrast, the short term priming we describe is distinct from prior
descriptions of trained immunity since it does not involve changes in gene expression.

The FcR is required for the full efficacy of many cancer immunotherapies, including popular immunotherapies like PD-1 and CTLA-4 blockades[9,45]. Some therapies, like the anti-CD20 antibody Rituximab, heavily rely on antibody-dependent cellular phagocytosis as a mechanism for eliminating cancer cells[4]. Interestingly, more frequent low dose treatments of Rituximab are more effective at treating Chronic Lymphocytic Leukemia (CLL) patients than higher dose treatments[28]. A key reason for this dosing schedule is to mitigate antigen shaving, or trogocytosis of target antigen. Enhancing phagocytosis without increasing antigen shaving is important but difficult. Our data shows that primed macrophages are better at phagocytosing whole cancer cells, but equally likely to trogocytose. This suggests that the current dosing regimen of frequent, low doses may already benefit from the effects of macrophage priming. Additionally, monocytes expressing Chimeric Antigen Receptors that signal through the FcR intracellular signaling domain are an exciting new avenue of cancer research[46–49]. How can we engineer hungrier macrophages to attack cancer cells? Our studies reveal that macrophage priming could enhance phagocytosis or other anti-cancer signaling pathways in these macrophages.

Acknowledgements
We thank members of the Morrissey lab for critical feedback on this manuscript. This work was supported by the UCSB Academic Senate, the National Institute of General Medical Sciences of the National Institutes of Health (R35 GM146935) and the UC Cancer Research Coordinating Committee (C23CR5592) to M.A.M. and the Eunice Kennedy Shriver National Institute of Child Health and Human Development of the National Institutes of Health (R01 HD108803-01) to M.Z.W. A.B. was supported by the Karl Storz Imaging fellowship. J.E.Q.N. was supported by a supplement to C23CR5592, the EUREKA scholars program and the MARC scholars program. A.G. was supported by the MARC scholars program. Schematics were created with BioRender.com. We thank the NRI-MCDB Microscopy Facility at UCSB, especially the director Ben Lopez for providing advice.

Author contributions
Conceptualization, A.B., M.Z.W. and M.A.M.; Methodology, A.B., M.Z.W. and M.A.M.; Software, K.R. and S.J.R; Validation, A.B.; Formal Analysis, A.B., S.F., K.R. and S.J.R.; Investigation, A.B, S.F., E.N. and A.G.; Resources, A.B. and M.A.M.; Writing - Original Draft, A.B. and M.A.M.; Writing - Review and Editing, A.B. and M.A.M.; Visualization, A.B.; Supervision, M.A.M.; Funding Acquisition, M.A.M.

Competing interests
The authors A.B., M.Z.W. and M.A.M. have filed a patent relating to this material. The authors have no other competing interests.

Figure legends

Figure 1. Optogenetic Fc Receptor (optoFcR) controls phagocytosis.
A) Schematic of optoFcR design, containing a myristoylation signal for membrane localization, the ITAM-containing intracellular domain from the FcR gamma-chain for signaling, iRFP fluorophore, and the homo-oligomerizing peptide - cryptochrome 2 (CRY2olig). B) Representative images of optoFcR in bone marrow derived macrophages (BMDMs). The optoFcR starts dispersed on the membrane (T0m; n=0/22 cells with visible clusters), and clusters after 15 minutes of constant high intensity light (T15m; n=21/22 cells with visible clusters). The optoFcR is declustered after 30 minutes in the dark (T45m; n=0/22 cells with visible clusters). See also supplemental movie 1. C) Representative images of downstream effector protein, SYK, colocalized with optoFcR in Raw macrophages stably expressing both the optoFcR and SYK-mNeonGreen. On the right, a line scan shows the Syk and optoFcR intensity at the position indicated by the yellow arrow in the inset. See also supplemental movie 2. D) Graph shows the Pearson's correlation coefficient for Syk and optoFcR at the cell cortex. Each data point represents a cell with an ROI drawn around a membrane region. The same ROI was used for both T0 (OFF) and T2 (ON). E) Schematic of experimental design for f-g. F) Representative images show optoFcR (green) expressing macrophages and ICAM conjugated beads visualized by atto390 in the supported lipid bilayer (magenta) before (OFF) and after (ON) 15 min of optoFcR activation with high intensity light. Internalized beads are labeled with a yellow star. See also supplemental movie 3. G) Quantification of phagocytosis in BMDMs after 15 minutes of optoFcR stimulation at low (5 uW/cm^2), medium (90 uW/cm^2), and high (1390 uW/cm^2) intensity light compared to control cells that do not express the optoFcR but receive the high intensity light stimulus. Both medium and high intensity light stimulate phagocytosis. Each data point represents the mean of an independent experiment. Data collected in the same replicate are denoted by symbol shape. In all graphs, bars represent the mean and SEM. * indicates p<0.05, *** indicates p<0.0005 using a paired t-test (d), one way ANOVA with dunnett correction (g). Scale bars are 10 um. See also supplemental figure 1.

Figure 2. Prior FcR activation specifically enhances macrophage sensitivity to IgG.
A) Schematic of experimental design for light-induced priming. Control (mCherry-CAAX) and optoFcR expressing BMDMs were stimulated with light for 15 min. The cells were returned to the dark for either 1 or 12 hrs, then targets opsonized with 1 nM IgG were introduced. After 15 minutes of phagocytosis, beads were washed out. Cells were imaged and the number of targets phagocytosed per cell was counted. B) Quantification of experiment described in (a). OptoFcR expressing cells that received light phagocytosed significantly more than cells that did not receive light, or cells that received light but did not express the optoFcR. C) Schematic of experimental design for

priming macrophages with IgG beads. Wildtype BMDMs were given either beads with
0.1 nM or 0 nM IgG for 15 min as a priming stimulus. Those beads were then washed
out and the cells were allowed to recover for 1 hr. A second set of different color beads
ligated to IgG at the indicated concentration were added and the amount of
phagocytosis was determined after 15 min. D) Quantification of phagocytosis for
experiment described in (c). Macrophages that were preincubated with 0.1 nM IgG-
conjugated beads phagocytosed significantly more than macrophages preincubated
with unopsonized beads. E) Phagocytosis of bead targets without an 'eat-me' signal.
Experiment was performed as described in (a) except the beads did not contain IgG.
Phagocytosis is normalized to unstimulated control cells. F) Phagocytosis of bead
targets with the efferocytic 'eat-me' signal, phosphatidylserine. Phagocytosis is
normalized to unstimulated control cells. Experiment was performed as described in (a)
except the beads were covered in a 10% phosphatidylserine bilayer to mimic an
apoptotic corpse. In all graphs, each point represents the mean of an independent
experiment. Data collected in the same replicate are denoted by symbol shape. Bars
represent the mean and SEM. * indicates $p<0.05$, **indicates $p<0.005$, **** indicates
$p<0.0001$ by two way anova with Sidak corrections (b), multiple T-tests with Holm-Sidak
corrections (d), and unpaired t-test (e,f). See also supplemental figure 2.
453
Figure 3. Macrophages are primed for increased cancer cell engulfment.
A) Schematic of experimental design for cancer cell targets. optoFcR cells were
exposed to light for 15 minutes, then returned to the dark. Raji B cells opsonized with 5
ng/ml IgG were added after 12 hours, and the cancer cell-macrophage interactions were
captured by timelapse microscopy. B) Representative images of optoFcR expressing
macrophages completing phagocytosis and trogocytosis. Raji B cells were visualized
using cell trace far red (magenta) and macrophages were visualized using optoFcR-
mSc (green). Yellow arrows point to phagocytosis of a whole cancer cell and white
arrows point to trogocytosis of cancer cell fragments. See also supplemental movies 4-
6. C) Percent of BMDMs that engulfed a whole target cell increased with prior FcR
stimulation. D) The number of whole raji cells engulfed was greater in primed
macrophages. E) Percent of BMDMs that trogocytosed a raji target was not significantly
different with priming. Each data point represents the mean of an independent
experiment, denoted by symbol shape, and bars represent the mean and SEM.
**indicates $p<0.005$ using an unpaired t-test. See also supplemental figure 3.
469
Figure 4. FcR mediated priming occurs via a short- and long-term mechanism.
A) Phagocytosis of 1nM IgG beads added at the indicated time post 15 min low intensity
light stimulation. Blue line denotes macrophages expressing the optoFcR and red
denotes macrophages expressing a control mCh-CAAX. Enhanced phagocytosis occurs
in two discrete peaks and lasts for at least 72 hrs. Points denote the mean of 4

independent replicates. Phagocytosis is normalized to unstimulated control cells. B)
Macrophages were treated with Actonomycin D (AD, 10nM) or Cycloheximide (CHX,
10ug/ml) to block transcription or translation respectively starting 7 hours before
phagocytosis. Macrophages were stimulated with light to activate the optoFcR at 1, 4 or
6 hours before phagocytosis. AD and CHX did not eliminate priming at 1 hour post-
stimulation. AD and CHX eliminated the enhanced phagocytosis phenotype at 4 and 6
hours post stimulation. C) Macrophages were treated with Erk inhibitor (PD0325901,
0.5uM) or DMSO control for 16 hrs before measuring phagocytosis. optoFcR
macrophages stimulated with light 1 hour before bead addition still showed enhanced
phagocytosis. Macrophages stimulated with light 12 hours before bead addition
phagocytosed the same number of beads as controls. Phagocytosis is normalized to
unstimulated control cells. Each data point represents the mean of an independent
experiment. Data collected in the same replicate are denoted by symbol shape. Bars
represent the mean and SEM. *indicates $p < 0.05$, **indicates $p < 0.005$, ***indicates
$p < 0.0005$, **** indicates $p < 0.0001$ using a two way anova with Sidak corrections (a-c).
490
Figure 5. Initiation of phagocytosis is faster and more likely in primed
macrophages.
A) optoFcR macrophages received a 15 min light stimulus 1 hour before bead addition
(primed) and were compared to optoFcR macrophages that did not receive light
stimulation (unprimed). Timelapse imaging was used to quantify the kinetics of
phagocytosis. Schematic shows each step in phagocytosis: binding (target contact with
cell), initiation (formation of the phagocytic cup), and completion (cup closure and bead
internalization). Below, the schematic shows percent of macrophage-bead contacts that
progress from one stage to the next for primed and unprimed macrophages. B)
Representative images of each stage. Scale bar is 10 um. See also supplemental movie
7. C) Percent of bead contacts that initiate phagocytosis is higher in primed
macrophages. D) Time from binding to initiation in optoFcR primed macrophages is
decreased compared with unprimed macrophages. E) Probability of completing
phagocytosis after initiation is comparable in primed and unprimed macrophages. F)
The speed of cup closure is comparable in primed and unprimed macrophages. G) The
overall success rate of phagocytosis is higher in primed macrophages compared to
unprimed macrophages. Each filled data point represents the mean of an independent
experiment, denoted by symbol shape, and corresponding outlined data points
represent individual bead times. Bars represent the mean and SEM. *indicates $p < 0.05$,
**indicates $p < 0.005$, ***indicates $p < 0.0005$ using an unpaired t-test on the means of
each replicate. See also supplemental figure 4.
512

Figure 6. Priming increases FcR mobility.
A) optoFcR macrophages were primed with a 15 minute light stimulus or were not stimulated. One hour later, single FcR molecules were labeled with QDots and tracked for 60 seconds. Tracks from a representative primed and unprimed image are shown. Scale bar is 1 um. B) Quantification of the per image mean jump distance (MJD) of all tracks. FcRs from primed macrophages had increased MJD compared to unprimed macrophages. C) Histogram of MJD from all acquired tracks from primed and unprimed macrophages. Primed macrophages show a population of FcRs that have a greater MJD than unprimed macrophages. Histogram was smoothed in prism using a second order polynomial with 6 neighbors on each size. D) Examples of 'free' and 'confined' tracks. E) Quantification of motion type analysis shows the proportion of tracks categorized as 'free' (circles) or 'confined' (triangles) for individual images. The percent of tracks defined as 'free' increases in optoFcR expressing macrophages exposed to light is compared to optoFcR macrophages with no light exposure. F) Quantification of the per image mean diffusion coefficients. Tracks from primed macrophages have a higher diffusion coefficient compared to unprimed macrophages. Each filled data point represents the mean from a single image. Images were acquired in three separate experiments, denoted by symbol shape (b and e). Bars represent the mean and SEM. *indicates $p < 0.05$, **indicates $p < 0.005$ using an unpaired t-test (b and f) or a one-way anova (e).

Methods

Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Meghan Morrissey (morrissey@ucsb.edu).

Bone-marrow derived macrophage cell culture.

Six- to ten-week-old C57BL/6 mice were sacrificed by CO2 inhalation. Hips and femurs were dissected and bone marrow was harvested as described in Weischenfeldt and Porse[50]. Macrophage progenitors were differentiated for seven days in RPMI-1640, 10% FBS, 1% PSG supplemented with 20% L929- conditioned media. Macrophage differentiation was confirmed by flow cytometry identifying CD11b and F4/80 double positive cells. Differentiated BMDMs were used for experiments from days 7 to 11.

Lentivirus production and infection.

All constructs were expressed in BMDMs using lentiviral infection. Lentivirus was produced in HEK293T cells transfected with pMD2.G (Gift from Didier Trono, Addgene plasmid # 12259 containing the VSV-G envelope protein), pCMV-dR8.2[51] (Gift from Bob Weinberg, Addgene plasmid #8455), and a lentiviral backbone vector containing the construct of interest using lipofectamine LTX (Invitrogen, Catalog # 15338–100). The media was harvested 72 h post-transfection, filtered through a 0.45 μm filter (Millapore, Catalog #SLHVM33RS) and concentrated using LentiX (Takara Biosciences, Catalog #631232). Concentrated lentivirus was added to cells on day 2 of differentiation. Cells were analyzed between days 7-11.

Optogenetic stimulation.

Cells receiving low intensity light (5 uW/cm^2) and medium intensity light (190 uW/cm^2) were stimulated using a LITOS LED illumination plate[52] for 15 min. Cells receiving high intensity light were stimulated using the 488 laser at 75% laser power on a spinning disc confocal microscope for 1 s at 20 s intervals for a total of 15 m (1,389 uW/cm^2). Intensity was determined using a slide power meter set to measure 450 nm light. Light used to prime cells was low intensity (5 uW/cm^2) unless otherwise indicated.

Clustering and colocalization analysis.

optoFcR clustering

50,000 BMDMs expressing the optoFcR were plated in one well of a 96-well glass bottom MatriPlate (Brooks, Catalog # MGB096-1-2-LG-L) between 12 and 24 h prior to the experiment. Cells were then continuously imaged with high intensity light stimulation for 30 min and then imaged for another 60 min without light stimulation. Clustering was determined by the presence of visible puncta in the cells.

SYK and optoFcR colocalization

50,000 RAW264.7 macrophages virally infected with both the optoFcR and SYK-NeonGreen (Addgene, Plasmid # 176610) were plated in one well of a 96-well glass bottom MatriPlate (Brooks, Catalog # MGB096-1-2-LG-L) between 12 and 24 h prior to the experiment. Cells were then continuously imaged with high intensity light stimulation for 30 min. Colocalization was determined by a pearsons correlation coefficient for the same membrane region at the first and last timepoints using the JaCoP plugin in ImageJ[53].

ICAM-1 protein purification.

ICAM-tagBFP-His$_{10}$[54] was expressed in SF9 or HiFive cells using the Bac-to-Bac baculovirus system as described previously[55]. Insect cell media containing secreted proteins was harvested 72 h after infection with baculovirus. His10 proteins were purified by using Ni-NTA agarose (QIAGEN, Catalog # 30230), followed by size exclusion chromatography using a Superdex 200 10/300 GL column (GE Healthcare, Catalog # 17517501). The purification buffer was 150 mM NaCl, 50 mM HEPES pH 7.4, 5% glycerol, 2 mM TCEP.

Supported lipid bilayer coated beads.

SUV preparation

For IgG conjugated beads the following chloroform-suspended lipids were mixed and desiccated overnight to remove chloroform: 98.8% POPC (Avanti, Catalog # 850457), 1% biotinyl cap PE (Avanti, Catalog # 870273), 0.1% PEG5000-PE (Avanti, Catalog # 880230, and 0.1% atto390-DOPE (ATTO-TEC GmbH, Catalog # AD 390–161) or 0.1% atto647-DOPE (ATTO-TEC GmbH, Catalog # AD 647–161). The lipid sheets were resuspended in PBS, pH7.2 (GIBCO, Catalog # 20012050) at 10 mM concentration and stored under inert nitrogen gas.

For ICAM-1 conjugated beads, the following chloroform-suspended lipids were mixed and desiccated overnight to remove chloroform: 97.8% POPC (Avanti, Catalog # 850457), 2% DGS-NTA (Avanti, Catalog # 790404), 0.1% PEG5000-PE (Avanti, Catalog # 880230, and 0.1% atto390-DOPE (ATTO-TEC GmbH, Catalog # AD 390–161). The lipid sheets were resuspended in PBS, pH7.2 (GIBCO, Catalog # 20012050) and stored under inert gas.

For PS beads the following chloroform-suspended lipids were mixed and desiccated overnight to remove chloroform: 89.8% POPC (Avanti, Catalog # 850457), 10% DOPS (Avanti, Catalog # 840035), 0.1% PEG5000-PE (Avanti, Catalog # 880230, and 0.1% atto390-DOPE (ATTO-TEC GmbH, Catalog # AD 390–161). The lipid sheets were resuspended in PBS, pH7.2 (GIBCO, Catalog # 20012050) and stored under inert gas.

613   For all SUVs, the lipids were broken into small unilamellar vesicles via several rounds of
614   freeze-thaws. The lipids were then stored at -80°C under argon for up to six months. To
615   remove aggregated lipids, the solution was diluted to 2 mM and filtered through a 0.22
616   uM filter (Millapore, Catalog # SLLG013SL) immediately prior to use.
617
Bead preparation
Silica beads with a 4.89 μm diameter (10% solids, Bangs Labs, Catalog # SS05003, Lot
13427) were washed with PBS, mixed with 1mM SUVs in PBS and incubated at room
temperature for 30 min with end-over-end mixing to allow for bilayer formation. Beads
were then washed with PBS to remove excess SUVs and incubated in 0.2% casein
(Sigma, Catalog # C5890) in PBS for 15 min before protein coupling (IgG and ICAM-1
beads). For IgG conjugated beads, anti-biotin AlexaFluor647-IgG (Jackson
ImmunoResearch Laboratories Catalog # 200-602-211, Lot # 156182) was added at 1
nM to a 10x dilution of beads (1% solids), unless otherwise indicated. For ICAM-1
conjugated beads, ICAM-1 was added at 10nM. Proteins were coupled to the bilayer for
30 min at room temperature with end-over-end mixing.
629
630   Bead engulfment assay
631   50,000 BMDMs were plated in one well of a 96-well glass bottom MatriPlate (Brooks,
632   Catalog # MGB096-1-2-LG-L) between 12 and 24 h prior to the experiment. ~8 × 10^5
633   beads were added to wells and engulfment was allowed to proceed for 15 min.
634
Inhibitors
For transcription and translation inhibited priming, 10 nM actinomycin D (Cell signaling,
Catalog # 15021s) or 10 ug/ml cycloheximide (Cell signaling, Catalog # 2112s) were
added to cells 7 hours prior to the start of the experiment. For ERK inhibited priming, 0.5
uM PD0325901 (Sigma, Catalog # PZ0162) was added to cells 16 hours prior to the
start of the experiment.
641
Microscopy and analysis
Images were acquired on a spinning disc confocal microscope (Nikon Ti2-E inverted
microscope with a Yokogawa CSU-W1 spinning disk unit and an Orca Fusion BT scMos
camera) equipped with a 40 × 0.95 NA Plan Apo air and a 100 × 1.49 NA oil immersion
objective. The microscope was controlled using Nikon Elements. Internalized particles
were counted in ImageJ by a blinded analyzer using Blind-Analysis-Tools-1.0 ImageJ
plug in.
649
Bead priming
50,000 BMDMs were plated in 1 well of a 96-well glass bottom plate 12-24 hrs prior to
the start of the experiment. A priming dose of ~8 × 10^5 atto390 beads conjugated to

either 1 nM or 0 nM IgG were added to the wells for 15 min. Any unengulfed beads
were washed out 5x with media, then checked with a dissecting microscope to confirm
the majority of engulfed beads had been removed. Then the cells were allowed to
recover for 1 hour. ~8 × 10^5 atto647 beads prepared with the indicated IgG
concentrations were then added to the wells and engulfment was allowed to proceed for
15 min. Cells were dyed with CellTrace CSFE (Thermo, Catalog # C34570) imaged and
the number of atto647 beads engulfed per cell were counted.
660
Raji eating assay.
40,000 BMDMs were plated in 1 well of a 96-well glass bottom plate 24 hrs prior to the
experiment and stimulated with low intensity LITOS illumination 12 hrs prior to the
experiment. Raji cells were dyed with CellTrace Far Red (Thermo, C34572), incubated
with a human-mouse hybrid aCD20 (InvivoGen hcd20-mab10, 5 ng/ml), added to wells
at 40,000 cells per well, and imaged immediately. 25 positions per well were
automatically selected and imaged every 3 min for 10 hrs. Unless otherwise noted, 100
macrophages were randomly selected and scored by a blind analyzer. Phagocytic
macrophages were characterized as BMDMs that engulfed 1 or more whole Raji cell
targets. Trogocytic macrophages were characterized as BMDMs that engulfed portions
of Raji targets. The number of Raji cells engulfed per 100 macrophages was also
counted.
673
Kinetics of engulfment.
BMDMs were plated as described in the bead engulfment assay 12-24 hrs prior to the
experiment and stimulated with low intensity LITOS illumination 1 hr prior to the
experiment. Using ND acquisition in Elements, 2-3 positions per well were manually
selected. Approximately 4 x 10^5 beads were added and phagocytosis was imaged at 20
s intervals through 7 z planes for 15 min. Only beads that bound within the first 12 min
were counted.
681
Receptor labeling and single particle tracking.
<u>Fab generation.</u>
Fabs from rat anti-mouse FcR (Cell signaling, 101307) and rabbit anti-rat biotin
conjugated (Invitrogen, 13-4813-85) were generated using the Pierce Fab Preparation
Kit (Thermo, 44985) according to the manufacture's protocol. In brief, antibodies were
run through a Zeba desalting column before cleavage with immobilized papain for 3
hours with end over end mixing and digestion was confirmed via SDS-PAGE. Fab
fragments were then purified using a NAb Protein A column.
690
<u>Receptor labeling and imaging.</u>

Single FcRs were labeled as previously described[31]. In brief, cells were blocked for 5
min in RPMI supplemented with 5% goat serum. Then, cells were incubated with
primary fab fragments for 10 min in blocking medium. Next, cells were incubated with
biotinylated secondary fab fragments for 10 min. Finally, cells were washed with
blocking media and incubated with streptavidin-congugated Qdot 655 (Thermo,
Q10123MP) for 4 min and immediately imaged. Images were acquired at 10 fps for 1
min using ND acquisition in Elements on a spinning disc confocal microscope.
699
Tracking.
Tracks and particle mean jump distance were generated using the trackmate plugin on
ImageJ[56,57]. Tracks that were less than 50 frames or that contained gaps greater than 3
frames were discarded from analysis. Motion types and diffusion coefficients were
determined using moment scaling spectrum analysis using the formula described in
Ewers et al, 2005[31,58–60].
706
707 **Quantification and statistical analysis.**
708 Statistical analysis was performed in Prism 8 (GraphPad). The statistical test used is
709 indicated in the relevant figure legend. Sample sizes were predetermined and indicated
710 in the relevant figure legend.
711

References

1. Freeman, S.A., and Grinstein, S. (2014). Phagocytosis: Receptors, signal integration, and the cytoskeleton. Immunological Reviews *262*, 193–215. 10.1111/imr.12212.

2. Nimmerjahn, F., and Ravetch, J.V. (2008). Fcγ receptors as regulators of immune responses. Nature Reviews Immunology *8*, 34–47. 10.1038/nri2206.

3. Gül, N., Babes, L., Siegmund, K., Korthouwer, R., Bögels, M., Braster, R., Vidarsson, G., ten Hagen, T.L.M., Kubes, P., and van Egmond, M. (2014). Macrophages eliminate circulating tumor cells after monoclonal antibody therapy. J Clin Invest *124*, 812–823. 10.1172/JCI66776.

4. Weiskopf, K., and Weissman, I.L. (2015). Macrophages are critical effectors of antibody therapies for cancer. mAbs *7*, 303–310. 10.1080/19420862.2015.1011450.

5. Montalvao, F., Garcia, Z., Celli, S., Breart, B., Deguine, J., Rooijen, N.V., and Bousso, P. (2013). The mechanism of anti-CD20–mediated B cell depletion revealed by intravital imaging. J Clin Invest *123*, 5098–5103. 10.1172/JCI70972.

6. Uchida, J., Hamaguchi, Y., Oliver, J.A., Ravetch, J.V., Poe, J.C., Haas, K.M., and Tedder, T.F. (2004). The innate mononuclear phagocyte network depletes B lymphocytes through Fc receptor-dependent mechanisms during anti-CD20 antibody immunotherapy. Journal of Experimental Medicine *199*, 1659–1669. 10.1084/jem.20040119.

7. Watanabe, M., Wallace, P.K., Keler, T., Deo, Y.M., Akewanlop, C., and Hayes, D.F. (1999). Antibody dependent cellular phagocytosis (ADCP) and antibody dependent cellular cytotoxicity (ADCC) of breast cancer cells mediated by bispecific antibody, MDX-210. Breast Cancer Research and Treatment *53*, 199–207. 10.1023/A:1006145507567.

8. Chen, X., Song, X., Li, K., and Zhang, T. (2019). FcγR-Binding Is an Important Functional Attribute for Immune Checkpoint Antibodies in Cancer Immunotherapy. Frontiers in Immunology *10*, 292. 10.3389/fimmu.2019.00292.

9. Dahan, R., Sega, E., Engelhardt, J., Selby, M., Korman, A.J., and Ravetch, J.V. (2015). FcγRs Modulate the Anti-tumor Activity of Antibodies Targeting the PD-1/PD-L1 Axis. Cancer Cell *28*.

10. Zhang, Y., Hoppe, A.D., and Swanson, J.A. (2010). Coordination of Fc receptor signaling regulates cellular commitment to phagocytosis. Proceedings of the National Academy of Sciences *107*, 19332–19337. 10.1073/pnas.1008248107.

11. Gottschalk, R.A., Martins, A.J., Angermann, B.R., Dutta, B., Ng, C.E., Uderhardt, S., Tsang, J.S., Fraser, I.D.C., Meier-Schellersheim, M., and Germain, R.N. (2016). Distinct NF-κB and MAPK Activation Thresholds Uncouple Steady-State Microbe Sensing from Anti-pathogen Inflammatory Responses. Cell Systems *2*, 378–390. 10.1016/j.cels.2016.04.016.

12. Bettadapur, A., Miller, H.W., and Ralston, K.S. (2020). Biting off what can be chewed: Trogocytosis in health, infection, and disease. Infection and Immunity *88*. 10.1128/IAI.00930-19/ASSET/50880292-BFB8-40CC-A318-88FA45997A88/ASSETS/GRAPHIC/IAI.00930-19-F0006.JPEG.

13. Velmurugan, R., Challa, D.K., Ram, S., Ober, R.J., and Ward, E.S. (2016). Macrophage-Mediated Trogocytosis Leads to Death of Antibody-Opsonized Tumor Cells. Molecular Cancer Therapeutics *15*, 1879–1889. 10.1158/1535-7163.MCT-15-0335.

14. Beum, P.V., Peek, E.M., Lindorfer, M.A., Beurskens, F.J., Engelberts, P.J., Parren, P.W.H.I., van de Winkel, J.G.J., and Taylor, R.P. (2011). Loss of CD20 and bound CD20 antibody from opsonized B cells occurs more rapidly because of trogocytosis mediated by Fc receptor-expressing effector cells than direct internalization by the B cells. Journal of immunology (Baltimore, Md.□: 1950) *187*, 3438–3447. 10.4049/JIMMUNOL.1101189.

15. Pham, T., Mero, P., and Booth, J.W. (2011). Dynamics of Macrophage Trogocytosis of Rituximab-Coated B Cells. PLoS One *6*, e14498. 10.1371/journal.pone.0014498.

16.   Weavers, H., Evans, I.R., Martin, P., and Wood, W. (2016). Corpse Engulfment
Generates a Molecular Memory that Primes the Macrophage Inflammatory Response. Cell
165, 1658–1671. 10.1016/j.cell.2016.04.049.
17.   Sharma, P., Vijaykumar, A., Raghavan, J.V., Rananaware, S.R., Alakesh, A., Bodele, J.,
Rehman, J.U., Shukla, S., Wagde, V., Nadig, S., et al. (2022). Particle uptake driven
phagocytosis in macrophages and neutrophils enhances bacterial clearance. Journal of
Controlled Release *343*, 131–141. 10.1016/j.jconrel.2022.01.030.
18.   Pinney, J.J., Rivera-Escalera, F., Chu, C.C., Whitehead, H.E., Vandermeid, K.R.,
Nelson, A.M., Barbeau, M.C., Zent, C.S., and Elliott, M.R. (2020). Macrophage hypophagia
as a mechanism of innate immune exhaustion in mAb-induced cell clearance. Blood *136*,
2065–2079. 10.1182/BLOOD.2020005571.
19.   Zent, C.S., and Elliott, M.R. (2017). Maxed out macs: physiologic cell clearance as a
function of macrophage phagocytic capacity. The FEBS Journal *284*, 1021–1039.
10.1111/febs.13961.
20.   Goodridge, H.S., Underhill, D.M., and Touret, N. (2012). Mechanisms of Fc Receptor
and Dectin-1 Activation for Phagocytosis. Traffic *13*, 1062–1071. 10.1111/j.1600-
0854.2012.01382.x.
21.   Kern, N., Dong, R., Douglas, S.M., Vale, R.D., and Morrissey, M.A. (2021). Tight
nanoscale clustering of Fcγ receptors using DNA origami promotes phagocytosis. eLife *10*.
10.7554/eLife.68311.
22.   Sobota, A., Strzelecka-Kiliszek, A., Gładkowska, E., Yoshida, K., Mrozińska, K., and
Kwiatkowska, K. (2005). Binding of IgG-Opsonized Particles to FcγR Is an Active Stage of
Phagocytosis That Involves Receptor Clustering and Phosphorylation. The Journal of
Immunology *175*, 4450–4457. 10.4049/jimmunol.175.7.4450.
23.   Taslimi, A., Vrana, J.D., Chen, D., Borinskaya, S., Mayer, B.J., Kennedy, M.J., and
Tucker, C.L. (2014). An optimized optogenetic clustering tool for probing protein interaction
and function. Nat Commun *5*, 4925. 10.1038/ncomms5925.
24.   Springer, T.A., and Dustin, M.L. (2012). Integrin inside-out signaling and the
immunological synapse. Current Opinion in Cell Biology *24*, 107–115.
10.1016/j.ceb.2011.10.004.
25.   Morrissey, M.A., Kern, N., and Vale, R.D. (2020). CD47 Ligation Repositions the
Inhibitory Receptor SIRPA to Suppress Integrin Activation and Phagocytosis. Immunity *53*,
290-302.e6. 10.1016/j.immuni.2020.07.008.
26.   Vorselen, D., Kamber, R.A., Labitigan, R.L.D., Loon, A.P. van, Peterman, E., Delgado,
M.K., Lin, S., Rasmussen, J.P., Bassik, M.C., and Theriot, J.A. (2022). Cell surface receptors
TREM2, CD14 and integrin αMβ2 drive sinking engulfment in phosphatidylserine-mediated
phagocytosis. Preprint at bioRxiv, 10.1101/2022.07.30.502145 10.1101/2022.07.30.502145.
27.   Williamson, A.P., and Vale, R.D. (2018). Spatial control of Draper receptor signaling
initiates apoptotic cell engulfment. J Cell Biol *217*, 3977–3992. 10.1083/jcb.201711175.
28.   Williams, M.E., Densmore, J.J., Pawluczkowycz, A.W., Beum, P.V., Kennedy, A.D.,
Lindorfer, M.A., Hamil, S.H., Eggleton, J.C., and Taylor, R.P. (2006). Thrice-Weekly Low-
Dose Rituximab Decreases CD20 Loss via Shaving and Promotes Enhanced Targeting in
Chronic Lymphocytic Leukemia1. The Journal of Immunology *177*, 7435–7443.
10.4049/jimmunol.177.10.7435.
29.   Freeman, S.A., Vega, A., Riedl, M., Collins, R.F., Ostrowski, P.P., Woods, E.C.,
Bertozzi, C.R., Tammi, M.I., Lidke, D.S., Johnson, P., et al. (2018). Transmembrane Pickets
Connect Cyto- and Pericellular Skeletons Forming Barriers to Receptor Engagement. Cell
172, 305-317.e10. 10.1016/j.cell.2017.12.023.
30.   Kusumi, A., Fujiwara, T.K., Chadda, R., Xie, M., Tsunoyama, T.A., Kalay, Z., Kasai,
R.S., and Suzuki, K.G.N. (2012). Dynamic Organizing Principles of the Plasma Membrane
that Regulate Signal Transduction: Commemorating the Fortieth Anniversary of Singer and

Nicolson's Fluid-Mosaic Model. Annual Review of Cell and Developmental Biology *28*, 215–
250. 10.1146/annurev-cellbio-100809-151736.
31.   Jaumouillé, V., Farkash, Y., Jaqaman, K., Das, R., Lowell, C.A., and Grinstein, S.
(2014). Actin Cytoskeleton Reorganization by Syk Regulates Fcγ Receptor Responsiveness
by Increasing Its Lateral Mobility and Clustering. Developmental Cell *29*, 534–546.
10.1016/j.devcel.2014.04.031.
32.   Andrews, N.L., Lidke, K.A., Pfeiffer, J.R., Burns, A.R., Wilson, B.S., Oliver, J.M., and
Lidke, D.S. (2008). Actin restricts FcepsilonRI diffusion and facilitates antigen-induced
receptor immobilization. Nat Cell Biol *10*, 955–963. 10.1038/ncb1755.
33.   Duchemin, A.M., Ernst, L.K., and Anderson, C.L. (1994). Clustering of the high affinity Fc
receptor for immunoglobulin G (Fc gamma RI) results in phosphorylation of its associated
gamma-chain. J Biol Chem *269*, 12111–12117.
34.   Lin, J., Kurilova, S., Scott, B.L., Bosworth, E., Iverson, B.E., Bailey, E.M., and Hoppe,
A.D. (2016). TIRF imaging of Fc gamma receptor microclusters dynamics and signaling on
macrophages during frustrated phagocytosis. BMC Immunology *17*, 5. 10.1186/s12865-016-
0143-2.
35.   Goodridge, H.S., Reyes, C.N., Becker, C.A., Katsumoto, T.R., Ma, J., Wolf, A.J., Bose,
N., Chan, A.S.H., Magee, A.S., Danielson, M.E., et al. (2011). Activation of the innate
immune receptor Dectin-1 upon formation of a 'phagocytic synapse.' Nature *472*, 471–475.
10.1038/nature10071.
36.   James, J.R., and Vale, R.D. (2012). Biophysical mechanism of T-cell receptor triggering
in a reconstituted system. Nature *487*, 64–69. 10.1038/nature11220.
37.   Bakalar, M.H., Joffe, A.M., Schmid, E.M., Son, S., Podolski, M., and Correspondence,
D.A.F. (2018). Size-Dependent Segregation Controls Macrophage Phagocytosis of Antibody-
Opsonized Targets. Cell *174*, 131–142. 10.1016/j.cell.2018.05.059.
38.   Freeman, S.A., Goyette, J., Furuya, W., Woods, E.C., Bertozzi, C.R., Bergmeier, W.,
Hinz, B., Van Der Merwe, P.A., Das, R., and Grinstein, S. (2016). Integrins Form an
Expanding Diffusional Barrier that Coordinates Phagocytosis. Cell *164*, 128–140.
10.1016/j.cell.2015.11.048.
39.   Davis, S.J., and van der Merwe, P.A. (2006). The kinetic-segregation model: TCR
triggering and beyond. Nature Immunology *7*, 803–809. 10.1038/ni1369.
40.   Shelby, S.A., and Veatch, S.L. (2023). The Membrane Phase Transition Gives Rise to
Responsive Plasma Membrane Structure and Function. Cold Spring Harb Perspect Biol,
a041395. 10.1101/cshperspect.a041395.
41.   Stone, M.B., Shelby, S.A., Nññez, M.F., Wisser, K., and Veatch, S.L. (2017). Protein
sorting by lipid phase-like domains supports emergent signaling function in b lymphocyte
plasma membranes. eLife *6*, 1–33. 10.7554/eLife.19891.
42.   Jaumouillé, V., Grinstein, S., Jaumouill?, V., and Grinstein, S. (2011). Receptor mobility,
the cytoskeleton, and particle binding during phagocytosis. Current Opinion in Cell Biology
23, 22–29. 10.1016/j.ceb.2010.10.006.
43.   Netea, M.G., Domínguez-Andrés, J., Barreiro, L.B., Chavakis, T., Divangahi, M., Fuchs,
E., Joosten, L.A.B., van der Meer, J.W.M., Mhlanga, M.M., Mulder, W.J.M., et al. (2020).
Defining trained immunity and its role in health and disease. Nat Rev Immunol *20*, 375–388.
10.1038/s41577-020-0285-6.
44.   Tehrani, S.S.H., Kogan, A., Mikulski, P., and Jansen, L.E.T. (2023). Remembering foods
and foes: emerging principles of transcriptional memory. Cell Death Differ, 1–11.
10.1038/s41418-023-01200-6.
45.   Simpson, T.R., Li, F., Montalvo-Ortiz, W., Sepulveda, M.A., Bergerhoff, K., Arce, F.,
Roddie, C., Henry, J.Y., Yagita, H., Wolchok, J.D., et al. (2013). Fc-dependent depletion of
tumor-infiltrating regulatory t cells co-defines the efficacy of anti-CTLA-4 therapy against
melanoma. Journal of Experimental Medicine *210*, 1695–1710. 10.1084/jem.20130579.

46.   Morrissey, M.A., Williamson, A.P., Steinbach, A.M., Roberts, E.W., Kern, N., Headley,
M.B., and Vale, R.D. (2018). Chimeric antigen receptors that trigger phagocytosis. eLife *7*,
e36688. 10.7554/eLife.36688.
47.   Klichinsky, M., Ruella, M., Shestova, O., Lu, X.M., Best, A., Zeeman, M., Schmierer, M.,
Gabrusiewicz, K., Anderson, N.R., Petty, N.E., et al. (2020). Human chimeric antigen
receptor macrophages for cancer immunotherapy. Nature Biotechnology *38*, 947–953.
10.1038/s41587-020-0462-y.
48.   Sloas, C., Gill, S., and Klichinsky, M. (2021). Engineered CAR-Macrophages as
Adoptive Immunotherapies for Solid Tumors. Front Immunol *12*, 783305.
10.3389/fimmu.2021.783305.
49.   Wang, S., Yang, Y., Ma, P., Zha, Y., Zhang, J., Lei, A., and Li, N. (2022). CAR-
macrophage: An extensive immune enhancer to fight cancer. eBioMedicine *76*.
10.1016/j.ebiom.2022.103873.
50.   Weischenfeldt, J., and Porse, B. (2008). Bone Marrow-Derived Macrophages (BMM):
Isolation and Applications. Cold Spring Harb Protoc *2008*, pdb.prot5080.
10.1101/pdb.prot5080.
51.   Stewart, S.A., Dykxhoorn, D.M., Palliser, D., Mizuno, H., Yu, E.Y., An, D.S., Sabatini,
D.M., Chen, I.S.Y., Hahn, W.C., Sharp, P.A., et al. (2003). Lentivirus-delivered stable gene
silencing by RNAi in primary cells. RNA *9*, 493–501. 10.1261/rna.2192803.
52.   Höhener, T.C., Landolt, A.E., Dessauges, C., Hinderling, L., Gagliardi, P.A., and Pertz,
O. (2022). LITOS: a versatile LED illumination tool for optogenetic stimulation. Sci Rep *12*,
13139. 10.1038/s41598-022-17312-x.
53.   Bolte, S., and Cordelières, F.P. (2006). A guided tour into subcellular colocalization
analysis in light microscopy. Journal of Microscopy *224*, 213–232. 10.1111/j.1365-
2818.2006.01706.x.
54.   O'Donoghue, G.P., Pielak, R.M., Smoligovets, A.A., Lin, J.J., and Groves, J.T. (2013).
Direct single molecule measurement of TCR triggering by agonist pMHC in living primary T
cells. eLife *2*, e00778. 10.7554/eLife.00778.
55.   Hui, E., and Vale, R.D. (2014). In vitro membrane reconstitution of the T-cell receptor
proximal signaling network. Nat Struct Mol Biol *21*, 133–142. 10.1038/nsmb.2762.
56.   Ershov, D., Phan, M.-S., Pylvänäinen, J.W., Rigaud, S.U., Le Blanc, L., Charles-Orszag,
A., Conway, J.R.W., Laine, R.F., Roy, N.H., Bonazzi, D., et al. (2022). TrackMate 7:
integrating state-of-the-art segmentation algorithms into tracking pipelines. Nat Methods *19*,
829–832. 10.1038/s41592-022-01507-1.
57.   Tinevez, J.-Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E.,
Bednarek, S.Y., Shorte, S.L., and Eliceiri, K.W. (2017). TrackMate: An open and extensible
platform for single-particle tracking. Methods *115*, 80–90. 10.1016/j.ymeth.2016.09.016.
58.   Ferrari, R., Manfroi, A.J., and Young, W.R. (2001). Strongly and weakly self-similar
diffusion. Physica D: Nonlinear Phenomena *154*, 111–137. 10.1016/S0167-2789(01)00234-2.
59.   Ewers, H., Smith, A.E., Sbalzarini, I.F., Lilie, H., Koumoutsakos, P., and Helenius, A.
(2005). Single-particle tracking of murine polyoma virus-like particles on live cells and
artificial membranes. Proceedings of the National Academy of Sciences *102*, 15110–15115.
10.1073/pnas.0504407102.
60.   Jaqaman, K., Kuwata, H., Touret, N., Collins, R., Trimble, W.S., Danuser, G., and
Grinstein, S. (2011). Cytoskeletal control of CD36 diffusion promotes its receptor and
signaling function. Cell *146*, 593–606. 10.1016/j.cell.2011.06.049.

Figure 1: optogenetic FcR controls phagocytosis

A. Myristoyl lipid, ITAM, Fluorophore, CRY2. 450 nm light. Dispersed: inactive / Clustered: active. Phosphorylation.

B. OFF - T0m. ON - T15m. OFF - T45m.

C. optoFcR, SYK, MERGE, INSET. OFF - T0m / ON - T2m. OFF / ON linescan plots: Normalized intensity vs Linescan Position (um). optoFcR (green), SYK (magenta).

D. Co-localization of optoFcR and SYK. Pearsons Correlation Coefficient. OFF / ON. **

E. Stimulate macrophages with light in presence of nonopsonized bead → Image and calculate phagocytic index.

F. optoFcR / ICAM Beads. OFF / ON. ★ internalized bead.

G. Phagocytosis of ICAM coated beads. Fold Change Phagocytosis. Light Intensity: Low, Medium, High, Control. * / ***

Figure 2: Prior FcR activation specifically enhances macrophage sensitivity to IgG

Figure 3: Primed macrophages phagocytose more antibody opsonized cancer cells

Figure 4: FcR mediated priming occurs via a short and long term mechanism

Figure 5: Initiation of phagocytosis is faster and the probability of completing phagocytosis is higher in primed macrophages

Figure 6: priming increases FcR mobility