TIPE drives a cancer stem-like phenotype by promoting glycolysis via PKM2/HIF-1α axis in melanoma

Maojin Tian[1#], Le Yang[2#], Ziqian Zhao[3#], Jigang Li[1], Lianqing Wang[1], Qingqing Yin[2], Wei Hu[1], Yunwei Lou[4], Jianxin Du[1*], Peiqing Zhao[1*]

[1] *Center of Translational Medicine, Zibo Central Hospital Affiliated to Binzhou Medical University, Zibo, 255036, PR China*
[2] *Shandong First Medical University, Jinan,250117, PR China*
[3] *The Second Medical College, Xinjiang Medical University, Urumqi, 830092, PR China*
[4] *School of Laboratory Medicine, Xinxiang Medical University, Xinxiang,453003, PR China*
[#] Contributed equally
* Correspondence to: Peiqing Zhao, Center of Translational Medicine, Zibo Central Hospital Affiliated to Binzhou Medical University, Zibo, China, 54 Gongqingtuan Xi Road, Zibo 255036, PR China. Jianxin Du, Center of Translational Medicine, Zibo Central Hospital Affiliated to Binzhou Medical University, Zibo, China, 54 Gongqingtuan Xi Road, Zibo 255036.
Tel: +86 533 2365217 Fax: +86 533 2360819
E-mail: bzjzzpq@sdu.edu.cn or bzjzzpq@163.com (Peiqing Zhao); jianxindu2005@126.com (Jianxin Du)

Abstract

TIPE (TNFAIP8) has been identified as an oncogene and participates in tumor biology. However, how its role in the metabolism of tumor cells during melanoma development remains unclear. Here, we demonstrated that TIPE promoted glycolysis by interacting with pyruvate kinase M2 (PKM2) in melanoma. We found that TIPE induced PKM2 dimerization, thereby facilitating its translocation from the cytoplasm to the nucleus. TIPE-mediated PKM2 dimerization consequently promoted HIF-1α activation and glycolysis, which contributed to melanoma progression and increased its stemness features. Notably, TIPE specifically phosphorylated PKM2 at Ser 37 in an ERK-dependent manner. Consistently, the expression of TIPE was positively correlated with the levels of PKM2 Ser37 phosphorylation and cancer stem cell markers in melanoma tissues from clinical samples and tumor bearing mice. In summary, our findings indicate that the TIPE/PKM2/HIF-1α signaling pathway plays a pivotal role in promoting cancer stem cell properties by facilitating the glycolysis, which would provide a promising therapeutic target for melanoma intervention.

Keywords: TIPE, Glycolysis, PKM2, HIF-1α, Melanoma

Introduction

Melanoma is derived from the aberrant growth of melanocytes and is classified into several subgroups such as cutaneous melanoma (CM), uveal melanoma (UM), and mucosal melanoma (MM), based on its tissue locations(Han et al., 2021). Cutaneous melanoma (CM) is the major subgroup of melanoma. Accumulating evidence has revealed that metabolic reprogramming contributes to the tumorigenesis and progression of melanoma. Owing to its low incidence rate, targeting metabolism might be a promising therapeutic approach for the treatment of melanoma and offers a deeper understanding of its pathobiology(Brunner and Finley, 2022; Faubert et al., 2020). However, melanoma-altered metabolic reprogramming remains poorly understood.

The Warburg effect, also known as aerobic glycolysis, is a part of metabolic reprogramming. It increases lactate levels regardless of oxygen availability in tumor cells, providing cancer cell tumorigenicity, progression, and chemoresistance(Guo et al., 2022). Pyruvate kinase isoform M2 (PKM2) is the key enzyme that irreversibly converts phosphoenolpyruvate (PEP) to pyruvate in the last step of glycolysis (Li et al., 2016; Liu et al., 2021). It is shifted between catalytically inactive

dimeric and active tetrameric forms, and the PKM2 dimer predominantly

serves as a key glycolytic enzyme that provides advantages for tumor

progression due to the Warburg effect(Liu et al., 2017). Post-translational

modifications of PKM2, such as FGFR1-mediated Tyr105

phosphorylation and ERK1/2-dependent Ser37 phosphorylation, can

promote PKM2 from a tetrameric to dimeric form, accelerating its

translocation into the nucleus (Christofk et al., 2008; Li et al., 2016). In

the nucleus, PKM2 acts as a protein kinase that phosphorylates

STAT3(Chen et al., 2022a; Dhanesha et al., 2022) and Histone H3(Niu et

al., 2020). Interestingly, nuclear PKM2 also functions as a coactivator for

several transcription factors, such as HIF-1α(Luo et al., 2011; Wang et al.,

2014) or Oct4(Yang et al., 2018), by which regulates metabolic gene

transcription and promotes tumorigenesis or sustains cancer stem cell

populations. Furthermore, HIF-1α is constitutively activated in melanoma

under both normal and hypoxic conditions and is accompanied by

increased glycolysis(D'Aguanno et al., 2021; Malekan et al., 2021).

Therefore, the precise mechanisms underlying PKM2 activity regulation

and its function as an HIF-1α coactivator might provide a molecular basis

for the melanoma targeted therapy.

The tumor necrosis factor alpha-induced protein 8 (TNFAIP8, TIPE)

family is composed of four members: TIPE, TIPE1, TIPE2, and

TIPE3(Yang et al., 2022). Accumulating evidence suggests that TIPE is

strongly associated with the development of various cancers by affecting cell proliferation, apoptosis, invasion, and metastasis(Goldsmith et al., 2017; Niture et al., 2018). Therefore, TIPE is broadly considered pro-cancerous(Padmavathi et al., 2018). However, we previous revealed that TIPE functions as a tumor suppressor in colitis and colitis-associated colon cancer(Lou et al., 2022), indicating that TIPE might serve a multifarious function in malignancies. Hitherto, the role of TIPE in melanoma remains unclear. In this study, we revealed that TIPE-mediated PKM2 Ser37 phosphorylation accelerates PKM2 translocation from the cytoplasm to the nucleus, thereby elevating its binding ability to HIF-1α and hypoxia response elements (HREs), thus promoting metabolic reprogramming. TIPE-mediated metabolic reprogramming can confer tumorigenesis and stem-like phenomena to melanoma cells. This process represents an important regulator of the TIPE family switching between oxidative phosphorylation and aerobic glycolysis, paving the way for cancer-specific metabolism in response to low-oxygen challenge.

Results

TIPE increases aerobic glycolysis in melanoma cells

To characterize how TIPE performs biological functions in melanoma, we examined its expression levels in melanoma cell lines. The results showed that A375 and A875 cells had relatively higher TIPE expression

than G361 and MM96L cells. Therefore, we downregulated TIPE expression in A375 cells and overexpressed TIPE in G361 cells to perform further experiments (**Fig.S1**). Furthermore, we demonstrated that TIPE promotes melanoma tumorigenesis both *in vitro* and *in vivo*(**Fig.S2**).

Consequently, we performed RNA sequencing analysis by overexpressing TIPE in G361 cells (**Fig. 1A**) to examine the underlying mechanisms involved in the progression of melanoma. The results revealed that the overexpression of TIPE dramatically promoted glycolysis metabolites and the HIF-1α pathway (**Fig. 1B, C**). Since metabolic reprogramming, especially aerobic glycolysis, is tightly dependent on the HIF-1α pathway(Baik et al., 2019), we assumed that TIPE promotes melanoma progression might via the regulation of metabolic reprogramming. Therefore, we performed a metabolomic analysis after interfering with TIPE in A375 cells. Interestingly, the results indicated that the aerobic glycolysis pathway, including pyruvate and lactate, was significantly inhibited in melanoma cells after TIPE silencing (**Fig. 1D**), confirming that TIPE facilitates glycolytic metabolism in melanoma. Consequently, ATPase activity, lactate secretion, and ATP content assays were performed to further confirm TIPE-induced glycolytic changes. The results demonstrated that TIPE decreased ATPase activity and ATP content while promoting lactate

production (**Fig. 1E-G**). Additionally, we measured the extracellular acidification rate (ECAR) after the overexpression or downregulation of TIPE using a Seahorse Bioscience Flux Analyzer. The results also revealed that TIPE promoted glycolysis and glycolytic capacity in melanoma cells (**Fig. 1H-K**).

Furthermore, we investigated whether TIPE affected HIF-1α transcriptional activity or its expression levels. The results demonstrated that TIPE increased the expression of HIF-1α at both mRNA and protein levels(**Fig. S3**). Interestingly, TIPE can dramatically activate the hypoxia response element(HRE) reporter activity, with which HIF-1α is more prone to recognize the specific DNA motif to regulate metabolic gene transcription(Chen et al., 2022b)(**Fig. 1L**).

Collectively, above results demonstrate that TIPE plays a crucial role in promoting melanoma glycolysis. However, the impact of TIPE on the Warburg effect by regulating HIF-1α activity requires further investigation.

TIPE interacts with PKM2, promotes its dimerization and nuclear import

To further study the functions of TIPE in melanoma, we used the Co-IP/MS method to bait the potential binding partners of TIPE (the candidate interacting proteins of TIPE are listed in **Table S1**). Interestingly, we found that PKM2, the key enzyme in aerobic glycolysis,

might be a new partner of TIPE in melanoma (**Fig. 2A**) (**Fig. S4**). Next, we performed Co-IP assays to confirm the interactions between TIPE and PKM2. The results revealed that exogenously overexpressed Flag-tagged PKM2 interacted with exogenously overexpressed HA-tagged TIPE (**Fig. 2B, C**). Furthermore, endogenous interactions between PKM2 and TIPE were confirmed in the A375 cells (**Fig. 2D, E**). GST-pull down assays demonstrated that TIPE directly bound to PKM2 *in vitro* (**Fig. 2F**). Moreover, we performed Doulink assay to further demonstrate their endogenous interactions. The results also showed that endogenous interactions between TIPE and PKM2 were confirmed in G361 cells (**Fig. 2G**). Based on these interactions, we identified the binding regions between TIPE and PKM2. We generated four fragments of PKM2, including amino acids 1-390(NA1BA2), 45-390(A1BA2), 1-219(NA1B) and 220-390(A2), based on their structural domains (**Fig. 2H**). The results showed that except for the A2 domain, the NA1BA2, A1BA2, and NA1B domains of PKM2 were necessary for its interaction with TIPE (**Fig. 2I**). Finally, the TIPE protein was divided into two fragments, including amino acids 1-100 (M1) and 101-198 (M2). The data showed that the amino acids 1-100(M1) and 101-198(M2) of TIPE interacted with PKM2 (**Fig. 2J**). Taken together, these data revealed that TIPE interacts with PKM2.

Since TIPE interacts with PKM2 in melanoma, we determined whether

TIPE increases aerobic glycolysis by affecting PKM2 expression or its activity. The results showed that TIPE did not influence the expression of PKM2 at either the mRNA or the protein level (**Fig. S5**). Intriguingly, we demonstrated that dimeric PKM2 was reduced in TIPE-silenced A375 cells (**Fig. 2K**) and was increased following TIPE overexpression in G361 cells (**Fig. 2L**). Normally, a dimeric form of PKM2 with low affinity for its substrate, phosphoenolpyruvate (PEP), and a tetrameric form have a high PEP affinity. Moreover, the dimeric form of PKM2 is more prone to translocation into the nucleus and thus exerts protein kinase activity, which confers the Warburg effect and tumorigenesis(Zhou et al., 2022). Western blot analysis of nuclear and cytosolic fractions from A375 cells revealed that PKM2 primarily resided in the cytoplasm in which TIPE was downregulated. Likewise , the nuclear level of PKM2 was increased after overexpression of TIPE in G361 cells (**Fig. 2M**). Furthermore, we examined the distribution of PKM2 after TIPE overexpression by using a Thermo Fisher EVOS microscopic analysis. As shown in **Fig.2N**, PKM2 was predominantly present in the nucleus after TIPE overexpression, and this phenomenon was diminished by the administration of a PKM2 dimer formation suppressor (TEPP-46, 100μM). It has been shown that dimeric PKM2 promotes tumor progression by regulating the Warburg effect(Zhou et al., 2022). Thus, we speculated that TIPE restores the nuclear levels of PKM2 to promote

melanoma tumorigenicity via a dimeric PKM2-dependent Warburg effect.

TIPE requires PKM2 for transcriptional activation of HIF-1α

Accumulating evidence has revealed that tetrameric PKM2 is

enzymatically active towards pyruvate and acts as a transcriptional

coactivator, while it dissociates into a dimer(Wei et al., 2020).

Consequently, the dimeric form of PKM2 becomes a coactivator of HIF-

1α to transcriptionally activate glycolytic genes in favor of the Warburg

effect (**Fig.3A**)(Ouyang et al., 2018). As PKM2 and HIF-1α both play

important roles in the metabolic reprogramming process, it is reasonable

to believe that the interactions between TIPE and these two molecules

could affect the glycolysis in melanoma. Thus, we investigated whether

TIPE promoted the transcriptional activation of HIF-1α in a PKM2-

dependent manner.

To confirm the effect of TIPE activation on HIF-1α transcriptional

activity, we tested the ability of TIPE to activate the HRE reporter

activity in HRE-luciferase plasmid transfected HEK-293T cells. Not

surprisingly, significant augmentation of HRE promoter activation by

TIPE was observed, especially when combined with PKM2 (**Fig.3B**).

Consequently, TIPE activated HRE reporter activity in a dose-dependent

manner, and this phenomenon was reversed after PKM2 knockdown,

demonstrating that TIPE has critical effects on the activation of HIF-1α

transcription in a PKM2-dependent manner (**Fig.3C**). HIF-1α targeted

genes such as *LDHA* and *GLUT1* were also upregulated by TIPE in a dose-dependent manner and were reversed by interfering with PKM2 (**Fig.3D, E**). Due to PKM2 cooperated with HIF-1α to activate glycolytic genes during the Warburg effect, we used the Duolink method to confirm whether TIPE affects the endogenous interaction between HIF-1α and PKM2. The results showed that the interaction between HIF-1α and PKM2 was decreased by TIPE interference, whereas the interaction was increased when TIPE was overexpressed (**Fig.3F, G**). More interestingly, TIPE enhanced the PKM2/HIF-1α interaction in the nucleus, further indicating that TIPE induced PKM2 nuclear translocation and thus promoted its interaction with HIF-1α (**Fig.3H**). In addition, the Co-IP assay revealed that TIPE elevated this interaction in a dose-dependent manner (**Fig.3I**).

Moreover, we analyzed the correlation between TIPE and hypoxia scores using TCGA dataset. Interestingly, we found that the hypoxia score was significantly related to the expression of TIPE (**Fig.3J**) (**Table S2**), and the TIPE higher expression group had a higher hypoxia score than the lower expression group in melanoma (**Fig.3K**).

Collectively, these results demonstrated that TIPE has critical effects on the transcriptional activation of HIF-1α in a PKM2-dependent manner.

TIPE promotes HIF-1α transcriptional activation via PKM2 Ser37 phosphorylation

The presence of PKM2 in the nucleus enhances glucose metabolism by

regulating HIF-1α(Luo et al., 2011). Our results preliminarily

demonstrate that TIPE promotes PKM2 dimerization and nuclear import,

thereby increasing HIF-1α transcriptional activation and promoting

aerobic glycolysis. Normally, Erk1/2-dependent phosphorylation of

PKM2 at serine 37 (Ser37) and FGFR1-dependent phosphorylation of

PKM2 at tyrosine 105 (Tyr105)(Li et al., 2016; Zhou et al., 2018) are

indicative of its dimerization and translocation, further promoting the

Warburg effect and serving as a transcription factor

coactivator(Novoyatleva et al., 2019; Wei et al., 2020). As shown in

Fig.4A, interference with TIPE in A375 cells blocked PKM2 Ser37

phosphorylation, and overexpression of TIPE in G361 cells caused Ser37

phosphorylation compared to controls. However, the phosphorylation of

PKM2 Tyr105 did not show any obvious changes. Consequently, we

hypothesized that this transformation caused by Ser37 phosphorylation

affects the interaction between TIPE and PKM2. As expected, the

mutation of PKM2 at S37A blocked its interaction with TIPE, but not

with the Y105D mutation (**Fig.4B**). Interestingly, this mutation also

interfered with the interaction between PKM2 and HIF-1α, which was

elevated by TIPE overexpression (**Fig.4C**). A previous study has revealed

that TIPE promotes chemoresistance in acute myeloid leukemia (AML)

by activating the ERK signaling pathway(Pang et al., 2020). In addition,

phosphorylation of PKM2 at serine 37 is ERK1/2-dependent

manner(Yang et al., 2012). TIPE does not serve as a phosphorylase to

perform such post-translational modifications directly. Taken together,

these results indicate that TIPE promotes PKM2 dimerization depending

on ERK signaling. As expected, the ERK inhibitor, U0126, inhibited

PKM2 Ser37 phosphorylation induced by TIPE overexpression (**Fig.4D**).

To investigate whether TIPE affects the binding affinity of PKM2 to the

Hif-1α specific binding site of the *LDHA* and *GLUT1* promoter using

chromatin immunoprecipitation *q*PCR (ChIP-*q*PCR) analysis, we found

that TIPE increased the binding of PKM2 to the *LDHA* and *GLUT1*

promoters. This phenomenon, caused by the overexpression of TIPE, was

diminished after the administration of TEPP-46, indicating that TIPE

promoted the binding of PKM2 to the *LDHA* and *GLUT1* promoters in a

dimeric PKM2-dependent manner (**Fig.4E, F**). Finally, we demonstrated

that the augmentation of TIPE on HRE promoter activation or HIF-1α

targeted genes such as *LDHA* and *GLUT1* was blocked by the S37A

mutation (**Fig.4G-I**).

In brief, we revealed that TIPE promoted HIF-1α transcriptional

activation via ERK-dependent PKM2 Ser37 phosphorylation.

TIPE facilitates melanoma tumorigenesis and aerobic glycolysis in a

dimeric PKM2-dependent manner

Several studies have demonstrated that dimeric PKM2 increases tumor

progression by regulating the Warburg effect. Therefore, we speculated that TIPE promotes melanoma growth through a dimeric PKM2-dependent Warburg effect. To demonstrate this, we knocked down TIPE in A375 cells and found that cell proliferation was decreased. Interestingly, we found that Pyridoxine, a PKM2 dimer formation inducer (Wei et al., 2020), was sufficient to abrogate the TIPE silencing-mediated cell proliferation (**Fig.5A**). Consistently, TEPP-46, a PKM2 dimer formation suppressor, was sufficient to reverse the increased proliferative capacity induced by TIPE-overexpression in G361 cells (**Fig.5B**). Clone formation ability (**Fig.5C, D**) and *in vivo* experiments (**Fig.5E-G**) also demonstrated this phenomenon. Interestingly, we found that decreased ECAR, glycolysis, and glycolytic capacity induced by TIPE interference could be fully antagonized by Pyridoxine in A375 cells (**Fig.5H-J**). In addition, TEPP-46 reversed the ECAR, glycolysis, and glycolytic capacity induced by TIPE overexpression in G361 cells (**Fig.5K-M**).

Lastly, to further demonstrate how TIPE affects the PKM2/HIF-1α pathway during the Warburg effect, we revealed that inhibition of PKM2 dimerization can rescue TIPE-induced cell proliferation. However, this phenomenon was diminished by the activation of HIF-1α, demonstrating that TIPE promotes melanoma cell proliferation via dimeric PKM2 dependent HIF-1α activation (**Fig.5N, O**). More interestingly, we evidenced that suppression of PKM2 dimerization inhibited the TIPE-

mediated Warburg effect. In addition, this phenomenon was diminished by the activation of HIF-1α (**Fig.5P, Q**).

Taken together, these data suggest that TIPE regulates cell growth and the Warburg effect by inducing the formation of dimeric PKM2, and thus activating the HIF-1α pathway.

TIPE accelerates melanoma cells stem-like phenomenon via promoting PKM2 dimerization under normoxic conditions

Increasing evidence has shown that the HIF-1α pathway is associated with acquisition of cancer stem-like properties(Weinstein et al., 2022; Zhu et al., 2022). To confirm the effects of TIPE on the stemness of melanoma cells under normoxic conditions, we examined the stemness-associated features *in vitro*. First, we measured cancer stem-like phenotype markers, including *NANOG*, *NOTCH*, *OCT3/4*, *SOX2* and *BMI-1*. The results showed that the markers were increased after overexpression of TIPE in G361 cells (**Fig.6A**) and were reduced after TIPE interference in A375 cells by using *q*PCR method (**Fig.6B**). We further revealed that TIPE increased the migration ability and increased sphere formation (**Fig.6C, D**). In addition, TIPE significantly increased the chemosensitivity of melanoma cell to sorafenib (**Fig.6E, F**). Next, we examined the tumorigenicity of melanoma cells with TIPE knockdown *in vivo* by injecting Ctrl-sh or TIPE-sh A375 cells subcutaneously into nude mice at three dilutions (1×10^6, 1×10^5 and 1×10^4) and allowing them to

grow for four weeks. The Confidence intervals (CIs) for 1/ (stem cell frequency) using extreme limiting dilution were calculated as previous reported(Hu and Smyth, 2009; Niu et al., 2021). The estimated CI for the frequency of CSCs in the TIPE knockdown group was 202749, compared with 18441 in the control group (*P*<0.001) (**Fig.6G**) (**Table S3, 4**). Moreover, CD44$^+$ subpopulations were significantly elevated, as detected using a fluorescence-activated cell sorting (FACS) assay (**Fig.6H, I**).

Because TIPE promotes HIF-1α transcriptional activation in a dimeric PKM2-dependent manner, we performed a sphere formation assay by overexpressing TIPE in G361 cells. The results showed that TEPP-46 partially reversed mammosphere formation induced by TIPE overexpression. Moreover, overexpression of HIF-1α diminished this phenomenon, indicating that TIPE-induced sphere formation depended on dimeric PKM2-dependent HIF-1α activation (**Fig.6J**). In addition, the increased CD44$^+$ subpopulations of G361 cells induced by TIPE were decreased by the administration of TEPP-46, and could be rescued by overexpression of HIF-1α(**Fig.6K**).

These findings strongly suggest that TIPE maintains melanoma cell stemness via dimeric PKM2-dependent HIF-1α activation.

TIPE is positively correlated with the levels of PKM2 Ser37 phosphorylation and cancer stem cell markers

We analyzed the expression of TIPE in melanoma and benign skin

tumors, and the results showed that the expression of TIPE was

dramatically increased compared to controls using a tissue chip (**Fig.7A,**

B). To determine if there was a correlation between TIPE and PKM2

Ser37 levels, TIPE and PKM2 Ser37 expression was analyzed by IHC

staining. The results showed that TIPE expression was positively

correlated with PKM2 Ser37 expression ($P < 0.001$) (**Fig.7C, D**).

Consistent with the above findings, immunohistochemistry (IHC)

results showed that TIPE, PKM2 Ser37, LDH, and CSCs marker CD44

were inhibited in A375 TIPE-interfering cell-derived xenografts (**Fig.7E**).

Lastly, TCGA dataset revealed that TIPE has a positive relationship with

CSCs markers including *BMI1*, *NANOG*, *NOTCH1*, and

OCT3/4(*POU5F1*) in melanoma (**Fig.7F-I**).

These results further demonstrate that TIPE maintains cancer stem cell

phenotypes in a PKM2 dimerization manner.

375 **Discussion**

376 Accumulating evidence has revealed that TIPE serves as an oncogene and

377 a negative regulator of apoptosis. Its overexpression is positively

378 associated with the development of various cancers, including liver, lung,

379 and breast cancers(Lou et al., 2022; Niture et al., 2018). However, Chen

380 et al. reported that TIPE initially inhibits cancer cell proliferation but

promotes tumor migration and metastasis(Li et al., 2021). In addition, a previous report has demonstrated that TIPE functions as a tumor suppressor in certain inflammatory carcinogenesis(Lou et al., 2022). These results suggest that TIPE plays a variety of roles in cancer progression. Interestingly, Niture et al. reported that TIPE promotes prostate cancer progression by regulating oxidative phosphorylation and glycolysis(Niture et al., 2021). However, the mechanism through which TIPE regulates metabolic reprogramming remains unclear. Here, we demonstrate that TIPE facilitates the Warberg effect by promoting dimeric PKM2-mediated HIF-1α transactivation in melanoma. Therefore, targeting TIPE may offer a novel therapeutic approach for melanoma treatment.

Mammalian pyruvate kinase consists of four isoforms, M1, M2, L, and R, which are encoded by two distinct genes (PKM and PKLR) (Valentini et al., 2002). Although these isoforms share similar features, only PKM2 exhibits the lowest basal enzymatic activity and is capable of configuration transformation between the tetramer and dimer, meeting differential metabolic demands(Morgan et al., 2013; Puckett et al., 2021). Cancer cells shift towards the preferential expression of the specific isoform of PKM2, especially its dimeric form, and serve as a pivotal metabolic enzyme that promotes the activation of oncogenic pathways in tumors(Icard et al., 2022). Normally, PKM2 enzymatic activity is

accurately modulated in a variety of ways, such as post-translational

modifications or metabolite binding, to maintain balanced levels in tumor

cells(Puckett et al., 2021). Here, we showed that TIPE promoted

melanoma tumorigenesis by promoting glycolysis. The results showed

that TIPE interacts with PKM2 and may participate in glycolysis.

However, TIPE did not influence the expression of PKM2 at either the

mRNA or the protein level. Because of the modulation of PKM2 kinetic

properties primarily via its dimer and tetramer configurations, we

demonstrated that TIPE could promote its dimeric form transformation,

increasing PKM2 residing in the nucleus. However, the downstream

signaling pathways that are mediated by TIPE in melanoma remain

largely unknown.

The nuclear PKM2 routinely serves as a coactivator to transcriptionally

activate certain metabolic genes(Cluntun et al., 2017; Zhang et al., 2019).

Because TIPE binds PKM2 and transports it into the nucleus, the dimeric

form of PKM2 more easily enters the nucleus to become a coactivator of

HIF-1α, resulting in the upregulation of HIF-1α targeted genes such as

LDHA and, *GLUT1*. Thus, we elucidate these relationships in the next

step. To verify whether TIPE activates HIF-1α transcriptional activity, we

found that TIPE significantly increased the activation of the HRE

promoter, especially when combined with PKM2. However, this

activation was diminished by interfering with PKM2, indicating that

TIPE has critical effects on the activation of HIF-1α transcription in a

PKM2-dependent manner. Because PKM2 interacts with HIF-1α and

potentiates PKM2-mediated HIF-1α transcription, we found that TIPE

increased the interaction between HIF-1α and PKM2. In TCGA dataset,

the hypoxia score was positively correlated with the mRNA expression of

TIPE in melanoma, and the TIPE higher expression group had a higher

hypoxia score than the lower expression group.

The mechanisms associated with PKM2 translocation are diverse, the

most important of which is Erk1/2-dependent phosphorylation of PKM2

at serine 37 (Ser37) and FGFR1-dependent phosphorylation of PKM2 at

tyrosine 105 (Tyr105)(Li et al., 2016; Zhou et al., 2018). Consequently,

we demonstrated that TIPE increased the levels of PKM2 Ser37 but not

PKM2 Tyr105. Furthermore, we conjectured that this transformation

caused by Ser37 phosphorylation affects the interaction between TIPE

and PKM2. The results showed that mutation of PKM2 at S37A blocked

its interaction with TIPE but not with Y105D. Because phosphorylation

of PKM2 at serine 37 is dependent on ERK1/2(Yang et al., 2012) and

TIPE promotes AML chemoresistance by activating the ERK signaling

pathway(Pang et al., 2020), we used an ERK inhibitor to further

demonstrate the activation of HIF-1αinduced by TIPE. Not surprisingly,

we found that TIPE facilitates the Warberg effect predominantly by

driving PKM2 Ser37 phosphorylation. Furthermore, we demonstrated

that the augmentation of TIPE on HRE promoter activation or HIF-1α

targeted genes such as *LDHA* and *GLUT1* was blocked by the S37A

mutation, as previously reported. These results demonstrated that TIPE

promotes HIF-1α transcriptional activation via ERK-dependent PKM2

Ser37 phosphorylation.

Increasing evidence has shown that HIF-1α promotes tumor

progression, metastasis, and therapeutic drug resistance by maintaining

cancer stemness(Cui et al., 2017; Niu et al., 2021). Here, we observed

that TIPE significantly enhanced tumorigenesis, glycolysis, and CSC

properties under normoxia in melanoma, whereas suppression of PKM2

dimerization rescued these TIPE-induced effects. Interestingly, we

revealed that decreasing of PKM2 dimerization can rescue TIPE induced

cell proliferation, glycolysis, and CSC properties. However, this

phenomenon was diminished by activation of HIF-1α, demonstrating that

TIPE promotes melanoma cells proliferation, glycolysis, and CSC

properties via dimeric PKM2 dependent HIF-1α activation. Furthermore,

using a tissue ship, we showed that the expression of TIPE in melanoma

tissues was dramatically increased compared to that in benign skin tumor

tissues. To determine if there was a correlation between TIPE and PKM2

Ser37 levels, the results also demonstrated that TIPE expression was

positively correlated with PKM2 Ser37 levels. Consistently, the

expression of TIPE has a positive relationship with PKM2 Ser37, LDH,

and the CSCs marker CD44 in TIPE-overexpressing cell-derived

xenografts. TCGA dataset revealed that TIPE has a positive relationship

with CSCs markers, including BMI1, NANOG, NOTCH1, and

OCT3/4(POU5F1) in melanoma. These results further demonstrate that

TIPE maintains melanoma cell stem phenotypes in a PKM2 dimerization

manner.

In summary, we identified that TIPE serves as an oncogenic candidate

for melanoma and determined that TIPE promotes the Warburg effect and

maintains the stemness of melanoma. TIPE binds to PKM2, inducing its

dimeric transformation by enhancing ERK-dependent PKM2 Ser37

phosphorylation, and thus activating the transcription of HIF-1α,

providing tumorigenicity and cancer stem-like phenomena for melanoma

progression. The newly identified TIPE/PKM2/HIF-1α axis is an ideal

therapeutic target for melanoma and opens an avenue for the development

of a novel strategy for precision therapy.

Materials and methods

Cell culture and plasmids construction

Human A375, A875, G361, MM96L and HEK-293T cell lines were

purchased from the Shanghai Institute of Cell Biology (Chinese Academy

of Sciences, Shanghai, China) and were maintained in Dulbecco's

modified Eagle's medium (DMEM, Gibco) supplemented with 10% fetal

bovine serum (FBS, Gibco) at 37°C under 5% CO_2. The TIPE

overexpression lentiviral vector (TIPE), TIPE expression knockdown

vector (TIPE-sh) and their control vectors were constructed by

GeneChem company (Shanghai, China). HIF-1α, PKM2-Flag, TIPE-HA

and their truncated vectors were constructed in our laboratory. The mutant

vectors such as PKM2 S37A were synthesized by Tsingke Biology Co.,

Ltd.(Beijing, China). All plasmids were transfected into the cells

according to the manufacturer's protocol.

Cell proliferation, cell cycle and colony formation assays

Cell proliferation was measured using the Cell Counting Kit-8 method

(Dojindo, Shanghai, China). For the cell cycle assay, the cells were

stained with propidium iodide (PI; Sigma, USA) and were measured with

a flow cytometer (FACS Aria 2, BD, USA). For the colony formation

assay, cells were seeded, fixed and stained as previously described(Yang

et al., 2022).

Transwell assay

Tranwell assay was performed as we previous described(Yang et al.,

2022). Photographs of at least three randomly selected fields were

captured and were counted.

Bioinformatics, transcriptomics and metabolomics analyses

The correlations between TIPE and HIF1A, Hypoxia score, cancer stem

cells (CSCs) markers including *BMI1*, *NANOG*, *NOTCH1* and

OCT3/4(POU5F1) were estimated by using the ASSISTANT for Clinical Bioinformation website (https://www.aclbi.com/static/index.html#/). Transcriptomics and metabolomics were performed and analyzed by GeneChem company (Shanghai, China). Overexpression of TIPE into G361 cells, interfering TIPE in A375 cells and their control group were randomly divided into three groups to perform transcriptomic analyses, respectively. Interfering TIPE in A375 cells and its control group were randomly divided into six groups to perform metabolomic analyses.

ATP content, ATPase activity, pyruvate kinase activity and lactic acid content measurement

The concentration of ATP was determined 72h after overexpression or interfering of TIPE by using an ATP assay kit (Beyotime, S0026) according to the manufacturer's instructions. ATPase activity was measured by using an ATPase Assay Kit (Sigma-Aldrich, MAK113). Pyruvate kinase activity and lactate production after overexpression or interfering of TIPE were measured using a Pyruvate Kinase Activity Assay Kit (Nanjing Jiancheng, A076-1-1) and Lactic Acid Assay Kit (Nanjing Jiancheng, A019-2-1) according to the manufacturer's instruction, respectively.

ECAR

The Extra Cellular Acidification Rate (ECAR) of melanoma cells were determined using the Seahorse XF extracellular flux analyzer (Agilent,

535 Santa Clara, CA, USA). The cells were plated, washed before analysis.

536 The cells were then treated at specific time points: glucose (10 mM),

537 followed by oligomycin (1 μM) and 2-DG (50 mM). Finally, the ECAR

538 were measured by the Seahorse XF software.

539 Chromatin Immunoprecipitation (ChIP) Quantitative PCR (ChIP-

540 qPCR)

Melanoma cells were cross-linked with 1% formaldehyde, lysed with

sodium dodecyl sulfate buffer and sonicated. Sheared DNA was

immunoprecipitated with PKM2 antibody and negative control anti-IgG,

and quantified using SYBR Green Realtime PCR analysis (ABI, USA).

Primer sequences are listed in **Table S5**. Fold enrichment was calculated

based on Ct as $2^{-\Delta(\Delta Ct)}$, where $\Delta Ct = Ct_{IP} - Ct_{Input}$ and $\Delta(\Delta Ct) = \Delta Ct_{antibody} -$

ΔCt_{IgG}.

548 Sphere formation assay

Melanoma cells transfected with TIPE or TIPE-sh were plated onto 24-

well low attachment surface polystyrene plates (Corning, USA). Cells

were grown in DMEM/F12 medium (Invitrogen, USA) for approximately

10 days supplemented with 20ng/mL EGF (Bioworld, BK0026), 2% B27

(Invitrogen, 17504044), 20 ng/mL basic FGF (PeproThech, #100-18B),

and 1% N2(Invitrogen, 17502048).

555 Luciferase reporter assay

556 Human HEK-293T cells were seeded onto 48-well plates and transiently

transfected with pHRE-Firefly luciferase reporter, TIPE-HA and PKM2-Flag or sh-RNA vector targeting PKM2 plasmids together with internal control reporter pTK (Thymidine kinase)-Renilla luciferase by Lipofectamine™ 2000 reagent (Thermo, USA). FLuc and RLuc activities were determined using the Dual-Luciferase Assay System (Promega, USA).

qPCR

mRNA was isolated and cDNAs were synthesized by using a Prime Script RT Reagent Kit (TaKaRa, China) according to manufacturer's instruction. *q*PCR was performed using Real SYBR Mixture (CoWin Bioscience, China) on an ABI PRISM 7500 instrument (ThermoFisher, USA). The sequences of the primers used in this study are listed in **Table S5**.

Duolink proximity ligation assay (PLA)

Duolink® In Situ PLA® kit (Sigma-Aldrich, DUO92101) was performed to demonstrate the endogenous interactions between TIPE and PKM2, and to interpret how TIPE impact the interactions between HIF-1α and PKM2 according to we described previously(Yang et al., 2022; Zhao et al., 2019). Quantification analysis was performed using the ImageJ software.

Nuclear and cytosolic fractionation

Nuclear and cytosolic fractions were prepared using NE-PER Extraction

Reagents (Thermo, 78833) according to the manufacturer's

recommendation. Finally, both cytosolic and nuclear subfractions were

stored in -80°C prior to western blot.

Western blot

The prepared lysates were separated by 10% sodium dodecyl sulfate

polyacrylamide gel electrophoresis and transferred onto PVDF

membranes (Bio-Rad, USA). The membranes were then blocked and

incubated overnight at 4°C with the primary antibodies (as listed in **Table**

S6). After incubated with secondary antibodies, the protein bands were

visualized using a Tanon 5200 Imager (Shanghai, China).

Immunohistochemical and immunofluorescence

Melanoma tissue chip was purchased from Zhongke Guanghua (Xi'an,

China) Intelligent Biotechnology Co., Ltd.(#K063Me01), and the

clinicopathological characteristics of melanoma specimens are

summarized in **Table S7**. Immunohistochemistry (IHC) staining was

performed as previously reported. The samples were finally analyzed by

using semiquantitative scoring criteria. The staining index values (0-12)

were obtained and calculated(Yang et al., 2022). For

immunofluorescence, cells were seeded onto 24-well plates at a density

of 5,000 cells per well. Then, the cells were fixed using 4% formaldehyde

solution and permeabilized with 0.5% Triton X-100 solution. Finally,

cells were incubated overnight with antibodies prior to blocked with 1%

bovine serum albumin (BSA) in phosphate-buffered saline containing

0.2% Tween-20. The fluorescence images of the cells were acquired with

a fluorescence microscope equipped with appropriate filter combinations.

Co-immunoprecipitation (Co-IP) and Co-immunoprecipitation mass

spectrometry (Co-IP/MS)

For Co-IP, the cells were seeded, harvested, and were then washed twice

with PBS and digested and lysed in lysis buffer. The lysed supernatant

was incubated with Tag-conjugated beads to bait the potential interacted

proteins according to the manufacturer's instruction. For Co-IP/MS, the

beads consequently were washed, eluted, resolved, and then stained with

Coomassie brilliant blue. The bands were excised and digested with

chymotrypsin and subjected to LC-MS/MS sequencing and data analysis

by OBiO Technology Corp.,Ltd. (Shanghai, China).

Flow Cytometry Analysis

Cell apoptosis was measured by using a Cell Apoptosis Analysis Kit

(Engreen Biosystem Co, Ltd., Beijing, China). The cells were then

incubated in serum-free medium for starvation prior to analysis and fixed

using 70% ethanol for 12 h. Consequently, cells were washed, incubated

with propidium iodide, and analyzed by a FACS scanner (Aria II, BD,

USA). For CD44 analyzing, the cells were seeded, harvested, and

incubated with anti-CD44 fluorescent antibody. Isotype-matched human

immunoglobulins served as controls. Finally, the cells were counted with

the FACS scanner.

624 Mouse xenograft tumor study

Xenograft tumor studies were conducted utilizing the 6-week-old male

BALB/c nude mice purchased from the Shanghai SLAC Laboratory

Animal Co., Ltd.. The total number of mice was randomly divided into

respective groups and maintained in a standard pathogen free

environment. All the animal experiments were approved by the Animal

Ethics Committee of Zibo Central Hospital, Binzhou Medical University.

Tumor growth was monitored regularly and the length, width, and height

measurements taken every 7 days. All mice were killed by 10 weeks post

injection. For the rescue test, one week after injection, mice bearing

TIPE-vector were treated with TEPP-46. TEPP-46 (50mg/kg) or vehicle

control (0.9% NaCl) was administrated via intraperitoneal injection. At

the end of experiment, mice were euthanized and tumors were collected.

Tumors were fixed with 4% paraformaldehyde for further

immunohistochemistry (IHC) analysis.

639 Statistical analysis

GraphPad Prism (GraphPad Software, San Diego, CA) was used for

statistical analyses. Two-tailed Student's t-tests were used to compare two

groups. Linear regression analysis was explored to analyze the correlation

of TIPE with other indicators. All the data are shown as the mean with

standard deviations (SD) from at least three independent experiments. A

P value of <0.05 was considered statistically significant for all tests.

Data availability Statement

648 All the data needed to evaluate the conclusions of the study are presented

649 in this paper. Additional data related to this study may be requested from

650 the corresponding author.

Acknowledgements

This work was supported by the National Natural Science Foundation of

China (Nos. 81972002, 12304241, 81971491), Natural Science

Foundation of Shandong Province (ZR2023QC68, ZR2023MC136,

ZR2021MC083, ZR2021MC165), Medicine and Health Science

Technology Development Projects of Shandong Province (202102050803

and 2018WS534), and Qingdao Postdoctoral Application Innovation

Project (QDBSH20230102068).

Ethics approval

660 This study has been approved by the Medical Ethics Committee, the

661 Animal Care and Use Committee of Zibo Central Hospital, Binzhou

662 Medical University.

Author contributions

Peiqing Zhao: Conceptualization, Supervision. Jianxin Du: Methodology,

Writing-Original draft preparation. Maojin Tian: Data curation,

Investigation. Le Yang: Investigation. Ziqian Zhao: Investigation. Jigang

Li: Validation. Qingqing Yin: Software. Lianqing Wang: Data curation.

Wei Hu: Investigation. Yunwei Lou: Visualization.

Declaration of competing interest

The authors declare that they have no conflicts of interest concerning this

article.

References

Baik, S.H., S. Kang, W. Lee, H. Choi, S. Chung, J.I. Kim, and I. Mook-Jung. 2019. A

Breakdown in Metabolic Reprogramming Causes Microglia Dysfunction in

Alzheimer's Disease. *Cell Metab*. 30:493-507 e496.

Brunner, J.S., and L.W.S. Finley. 2022. Metabolic determinants of tumour initiation. *Nat Rev*

Endocrinol.

Chen, C., W. Zhang, T. Zhou, Q. Liu, C. Han, Z. Huang, S. Chen, Q. Mei, C. Zhang, K. Zhang,

H. Ma, R. Zhou, W. Jiang, W. Pan, and S. Zhu. 2022a. Vitamin B5 rewires Th17 cell

metabolism via impeding PKM2 nuclear translocation. *Cell Rep*. 41:111741.

Chen, Y., C. Cattoglio, G.M. Dailey, Q. Zhu, R. Tjian, and X. Darzacq. 2022b. Mechanisms

governing target search and binding dynamics of hypoxia-inducible factors. *Elife*. 11.

Christofk, H.R., M.G. Vander Heiden, N. Wu, J.M. Asara, and L.C. Cantley. 2008. Pyruvate

kinase M2 is a phosphotyrosine-binding protein. *Nature*. 452:181-186.

Cluntun, A.A., M.J. Lukey, R.A. Cerione, and J.W. Locasale. 2017. Glutamine Metabolism in

Cancer: Understanding the Heterogeneity. *Trends Cancer*. 3:169-180.

Cui, C.P., C.C. Wong, A.K. Kai, D.W. Ho, E.Y. Lau, Y.M. Tsui, L.K. Chan, T.T. Cheung, K.S.

Chok, A.C.Y. Chan, R.C. Lo, J.M. Lee, T.K. Lee, and I.O.L. Ng. 2017. SENP1

promotes hypoxia-induced cancer stemness by HIF-1alpha deSUMOylation and

SENP1/HIF-1alpha positive feedback loop. *Gut*. 66:2149-2159.

D'Aguanno, S., F. Mallone, M. Marenco, D. Del Bufalo, and A. Moramarco. 2021. Hypoxia-

dependent drivers of melanoma progression. *J Exp Clin Cancer Res*. 40:159.

Dhanesha, N., R.B. Patel, P. Doddapattar, M. Ghatge, G.D. Flora, M. Jain, D. Thedens, H.

Olalde, M. Kumskova, E.C. Leira, and A.K. Chauhan. 2022. PKM2 promotes

neutrophil activation and cerebral thromboinflammation: therapeutic implications for

ischemic stroke. *Blood*. 139:1234-1245.

Faubert, B., A. Solmonson, and R.J. DeBerardinis. 2020. Metabolic reprogramming and

cancer progression. *Science*. 368.

Goldsmith, J.R., S. Fayngerts, and Y.H. Chen. 2017. Regulation of inflammation and

tumorigenesis by the TIPE family of phospholipid transfer proteins. *Cell Mol Immunol*.

14:482-487.

Guo, D., Y. Tong, X. Jiang, Y. Meng, H. Jiang, L. Du, Q. Wu, S. Li, S. Luo, M. Li, L. Xiao, H.

He, X. He, Q. Yu, J. Fang, and Z. Lu. 2022. Aerobic glycolysis promotes tumor

immune evasion by hexokinase2-mediated phosphorylation of IkappaBalpha. *Cell*

Metab. 34:1312-1324 e1316.

Han, A., Z.T. Schug, and A.E. Aplin. 2021. Metabolic Alterations and Therapeutic

Opportunities in Rare Forms of Melanoma. *Trends Cancer*. 7:671-681.

Hu, Y., and G.K. Smyth. 2009. ELDA: extreme limiting dilution analysis for comparing

depleted and enriched populations in stem cell and other assays. *J Immunol Methods*.

347:70-78.

Icard, P., L. Simula, L. Fournel, K. Leroy, A. Lupo, D. Damotte, M.C. Charpentier, C. Durdux,

M. Loi, O. Schussler, G. Chassagnon, A. Coquerel, H. Lincet, V. De Pauw, and M.

Alifano. 2022. The strategic roles of four enzymes in the interconnection between

metabolism and oncogene activation in non-small cell lung cancer: Therapeutic

implications. *Drug Resist Updat*. 63:100852.

Li, M., X. Li, J.R. Goldsmith, S. Shi, L. Zhang, A. Zamani, L. Wan, H. Sun, T. Li, J. Yu, Z.

Etwebi, M.J. Bou-Dargham, and Y.H. Chen. 2021. Decoupling tumor cell metastasis

from growth by cellular pilot protein TNFAIP8. *Oncogene*. 40:6456-6468.

Li, N., L. Feng, H. Liu, J. Wang, M. Kasembeli, M.K. Tran, D.J. Tweardy, S.H. Lin, and J.

Chen. 2016. PARP Inhibition Suppresses Growth of EGFR-Mutant Cancers by

Targeting Nuclear PKM2. *Cell Rep*. 15:843-856.

Liu, F., F. Ma, Y. Wang, L. Hao, H. Zeng, C. Jia, Y. Wang, P. Liu, I.M. Ong, B. Li, G. Chen, J.

Jiang, S. Gong, L. Li, and W. Xu. 2017. PKM2 methylation by CARM1 activates

aerobic glycolysis to promote tumorigenesis. *Nat Cell Biol*. 19:1358-1370.

Liu, F., M. Ma, A. Gao, F. Ma, G. Ma, P. Liu, C. Jia, Y. Wang, K. Donahue, S. Zhang, I.M.

Ong, S. Keles, L. Li, and W. Xu. 2021. PKM2-TMEM33 axis regulates lipid

homeostasis in cancer cells by controlling SCAP stability. *EMBO J*. 40:e108065.

Lou, Y., X. Tian, C. Sun, M. Song, M. Han, Y. Zhao, Y. Song, X. Song, W. Zhang, Y.H. Chen,

and H. Wang. 2022. TNFAIP8 protein functions as a tumor suppressor in

inflammation-associated colorectal tumorigenesis. *Cell Death Dis*. 13:311.

Luo, W., H. Hu, R. Chang, J. Zhong, M. Knabel, R. O'Meally, R.N. Cole, A. Pandey, and G.L.

Semenza. 2011. Pyruvate kinase M2 is a PHD3-stimulated coactivator for hypoxia-

inducible factor 1. *Cell*. 145:732-744.

Malekan, M., M.A. Ebrahimzadeh, and F. Sheida. 2021. The role of Hypoxia-Inducible Factor-

1alpha and its signaling in melanoma. *Biomed Pharmacother*. 141:111873.

Morgan, H.P., F.J. O'Reilly, M.A. Wear, J.R. O'Neill, L.A. Fothergill-Gilmore, T. Hupp, and

M.D. Walkinshaw. 2013. M2 pyruvate kinase provides a mechanism for nutrient

sensing and regulation of cell proliferation. *Proc Natl Acad Sci U S A*. 110:5881-5886.

Niture, S., X. Dong, E. Arthur, U. Chimeh, S.S. Niture, W. Zheng, and D. Kumar. 2018.

Oncogenic Role of Tumor Necrosis Factor alpha-Induced Protein 8 (TNFAIP8). *Cells*.

8.

Niture, S., M. Lin, J.O. Odera, J. Moore, H. Zhe, X. Chen, S. Suy, S.P. Collins, and D. Kumar.

2021. TNFAIP8 drives metabolic reprogramming to promote prostate cancer cell

proliferation. *Int J Biochem Cell Biol*. 130:105885.

Niu, J., W. Li, C. Liang, X. Wang, X. Yao, R.H. Yang, Z.S. Zhang, H.F. Liu, F.Y. Liu, S.H. Pei,

W.Q. Li, H. Sun, D. Fang, and S.Q. Xie. 2020. EGF promotes DKK1 transcription in

hepatocellular carcinoma by enhancing the phosphorylation and acetylation of

histone H3. *Sci Signal*. 13.

Niu, Y., Z. Lin, A. Wan, L. Sun, S. Yan, H. Liang, S. Zhan, D. Chen, X. Bu, P. Liu, C. Chen, W.

He, X. Lu, and G. Wan. 2021. Loss-of-Function Genetic Screening Identifies Aldolase

A as an Essential Driver for Liver Cancer Cell Growth Under Hypoxia. *Hepatology*.

74:1461-1479.

Novoyatleva, T., N. Rai, N. Weissmann, F. Grimminger, H.A. Ghofrani, W. Seeger, H. Gall,

and R.T. Schermuly. 2019. Is PKM2 Phosphorylation a Prerequisite for Oligomer

Disassembly in Pulmonary Arterial Hypertension? *Am J Respir Crit Care Med*.

200:1550-1554.

Ouyang, X., S.N. Han, J.Y. Zhang, E. Dioletis, B.T. Nemeth, P. Pacher, D. Feng, R. Bataller,

J. Cabezas, P. Starkel, J. Caballeria, R.L. Pongratz, S.Y. Cai, B. Schnabl, R. Hoque,

Y. Chen, W.H. Yang, I. Garcia-Martinez, F.S. Wang, B. Gao, N.J. Torok, R.G. Kibbey,

and W.Z. Mehal. 2018. Digoxin Suppresses Pyruvate Kinase M2-Promoted HIF-

1alpha Transactivation in Steatohepatitis. *Cell Metab*. 27:339-350 e333.

Padmavathi, G., K. Banik, J. Monisha, D. Bordoloi, B. Shabnam, F. Arfuso, G. Sethi, L. Fan,

and A.B. Kunnumakkara. 2018. Novel tumor necrosis factor-alpha induced protein

eight (TNFAIP8/TIPE) family: Functions and downstream targets involved in cancer

progression. *Cancer Lett*. 432:260-271.

Pang, Y., Y. Zhao, Y. Wang, X. Wang, R. Wang, N. Liu, P. Li, M. Ji, J. Ye, T. Sun, J. Li, D. Ma,

F. Lu, and C. Ji. 2020. TNFAIP8 promotes AML chemoresistance by activating ERK

signaling pathway through interaction with Rac1. *J Exp Clin Cancer Res*. 39:158.

Puckett, D.L., M. Alquraishi, W. Chowanadisai, and A. Bettaieb. 2021. The Role of PKM2 in

Metabolic Reprogramming: Insights into the Regulatory Roles of Non-Coding RNAs.

Int J Mol Sci. 22.

Valentini, G., L.R. Chiarelli, R. Fortin, M. Dolzan, A. Galizzi, D.J. Abraham, C. Wang, P.

Bianchi, A. Zanella, and A. Mattevi. 2002. Structure and function of human

erythrocyte pyruvate kinase. Molecular basis of nonspherocytic hemolytic anemia. *J*

Biol Chem. 277:23807-23814.

Wang, H.J., Y.J. Hsieh, W.C. Cheng, C.P. Lin, Y.S. Lin, S.F. Yang, C.C. Chen, Y. Izumiya,

J.S. Yu, H.J. Kung, and W.C. Wang. 2014. JMJD5 regulates PKM2 nuclear

translocation and reprograms HIF-1alpha-mediated glucose metabolism. *Proc Natl*

Acad Sci U S A. 111:279-284.

Wei, Y., M. Lu, M. Mei, H. Wang, Z. Han, M. Chen, H. Yao, N. Song, X. Ding, J. Ding, M. Xiao,

and G. Hu. 2020. Pyridoxine induces glutathione synthesis via PKM2-mediated Nrf2

transactivation and confers neuroprotection. *Nat Commun*. 11:941.

Weinstein, A.G., I. Godet, and D.M. Gilkes. 2022. The rise of viperin: the emerging role of

viperin in cancer progression. *J Clin Invest*. 132.

Yang, M., Y. Zhang, G. Liu, Z. Zhao, J. Li, L. Yang, K. Liu, W. Hu, Y. Lou, J. Jiang, Q. Liu,

and P. Zhao. 2022. TIPE1 inhibits osteosarcoma tumorigenesis and progression by

regulating PRMT1 mediated STAT3 arginine methylation. *Cell Death Dis*. 13:815.

Yang, W., Y. Zheng, Y. Xia, H. Ji, X. Chen, F. Guo, C.A. Lyssiotis, K. Aldape, L.C. Cantley,

and Z. Lu. 2012. ERK1/2-dependent phosphorylation and nuclear translocation of

PKM2 promotes the Warburg effect. *Nat Cell Biol*. 14:1295-1304.

Yang, Y.C., M.H. Chien, H.Y. Liu, Y.C. Chang, C.K. Chen, W.J. Lee, T.C. Kuo, M. Hsiao, K.T.

Hua, and T.Y. Cheng. 2018. Nuclear translocation of PKM2/AMPK complex sustains

cancer stem cell populations under glucose restriction stress. *Cancer Lett*. 421:28-40.

Zhang, Z., X. Deng, Y. Liu, Y. Liu, L. Sun, and F. Chen. 2019. PKM2, function and expression

and regulation. *Cell Biosci*. 9:52.

Zhao, P., X. Pang, J. Jiang, L. Wang, X. Zhu, Y. Yin, Q. Zhai, X. Xiang, F. Feng, and W. Xu.

2019. TIPE1 promotes cervical cancer progression by repression of p53 acetylation

and is associated with poor cervical cancer outcome. *Carcinogenesis*. 40:592-599.

Zhou, Q., Y. Yin, M. Yu, D. Gao, J. Sun, Z. Yang, J. Weng, W. Chen, M. Atyah, Y. Shen, Q.

Ye, C.W. Li, M.C. Hung, Q. Dong, C. Zhou, and N. Ren. 2022. GTPBP4 promotes

hepatocellular carcinoma progression and metastasis via the PKM2 dependent

glucose metabolism. *Redox Biol*. 56:102458.

Zhou, Z., M. Li, L. Zhang, H. Zhao, O. Sahin, J. Chen, J.J. Zhao, Z. Songyang, and D. Yu.

2018. Oncogenic Kinase-Induced PKM2 Tyrosine 105 Phosphorylation Converts

Nononcogenic PKM2 to a Tumor Promoter and Induces Cancer Stem-like Cells.

Cancer Res. 78:2248-2261.

Zhu, B., W. Zhong, X. Cao, G. Pan, M. Xu, J. Zheng, H. Chen, X. Feng, C. Luo, C. Lu, J. Xiao,

W. Lin, C. Lai, M. Li, X. Du, Q. Yi, and D. Yan. 2022. Loss of miR-31-5p drives

hematopoietic stem cell malignant transformation and restoration eliminates leukemia

stem cells in mice. *Sci Transl Med*. 14:eabh2548.

Figure Legends

Figure 1. Effect of TIPE on melanoma cell glycolysis. A, B. Transcriptomics analysis by unsupervised hierarchical clustering and Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis showed that TIPE increased glycolysis and HIF-1α pathways. **C.** GSEA analysis of glycolysis showed that TIPE enhanced glycolysis compared to the control. **D.** Untargeted metabolomics analysis indicated that interfering with TIPE decreased the glycolysis pathway. **E-G.** TIPE decreases ATPase activity and ATP content and increases lactate levels. **H, I.** Overexpression of TIPE promotes glycolysis and glycolytic capacity according to ECAR analysis. **J, K.** Interfering with TIPE decreased glycolysis and glycolytic capacity using ECAR analysis. **L.** TIPE significantly activated HRE activity. *$P<0.05$; **$P<0.01$; ***$P<0.001$.

Figure 2. TIPE interacts with PKM2 to govern its nuclear import in a dimeric form dependent manner. A. Co-IP/MS analysis demonstrated that PKM2 interacted with TIPE in A375 cells. **B, C.** Immunoprecipitation (IP) and western blot analysis of the exogenous TIPE/PKM2 proteins interaction in the HEK-293T cells co-transfected

with Flag-tagged PKM2 and HA-tagged TIPE. **D, E.** IP and western blot analysis of the endogenous TIPE/PKM2 proteins interaction in the A375 cells. **F.** GST-pull down assay analysis of TIPE/PKM2 proteins interaction using purified GST-tagged PKM2 and TIPE-HA. **G.** TIPE (rabbit source, $TIPE_R$) endogenously interacted with PKM2(mouse source, $PKM2_M$) in G361 cells by using the Doulink assay. The results suggested that TIPE interacted with PKM2 to form a red color complex. The red color signal is generated only when the two proteins are too close to interaction. **H, I.** IP and western blot analysis of HA-tagged TIPE and Flag-tagged PKM2 fragment protein interaction in HEK-293T cells. **J.** IP and western blot analysis of the Flag-tagged PKM2 and HA-tagged TIPE fragment protein interaction in HEK-293T cells. **K.** Interfering TIPE decreased PKM2 dimeric formation but increased tetramer formation, as analyzed by BN-PAGE with β-actin as a loading control. **L.** Overexpression of TIPE increased PKM2 dimeric formation and decreased tetramer formation. **M.** Western blot showed that TIPE promoted PKM2 translocation into the nucleus. **N.** TIPE enhanced PKM2 translocation into the nucleus, and this phenomenon was diminished by the administration of TEPP-46 (100μM). *$P<0.05$; **$P<0.01$.

Figure 3. TIPE promotes HIF-1α transcription in a PKM2-dependent manner. A. Proposed molecular mechanism by which dimeric PKM2

regulates cell proliferation and glycolysis by modulating HIF-1α activity.

B. TIPE, especially when combined with PKM2, boosts relative HRE

luciferase activity, as examined by luciferase reporter assay. **C.** TIPE

promoted HRE activity in a dose- and PKM2-dependent manner. **D, E.**

TIPE increases HIF-1α targeted genes, including *LDHA* and *GLUT1*, in a

dose- and PKM2-dependent manner. **F, G.** TIPE promoted endogenous

interaction between PKM2 and HIF-1α in melanoma cells using a

Doulink assay. Interference of TIPE in A375 cells promoted the

interaction between PKM2 and HIF-1α(upper) compared to that

overexpression of TIPE in G361 cells decreased their interaction (lower).

The density of the red color signaling means the interactive strength

between PKM2 and HIF-1α affected by TIPE. **H.** TIPE enhanced the

PKM2/HIF1a interaction in the nucleus. **I.** TIPE increased the exogenous

interaction between PKM2 and HIF-1α in a dose- dependent manner in

HEK-293T cells. **J.** TCGA dataset revealed that TIPE has a positive

relationship with hypoxia score in melanoma. **K.** Higher expression of

TIPE is associated with a relatively higher hypoxia score in melanoma.

*$P<0.05$; **$P<0.01$; ***$P<0.001$.

Figure 4. TIPE increases HIF-1α activity depending on PKM2 Ser37

phosphorylation. A. TIPE enhanced PKM2 Ser37 phosphorylation, but

not that of Tyr105. **B.** PKM2 Ser37, but not Tyr105, increased its

interaction with TIPE. **C.** PKM2 Ser37 mutation(S37A) hampered its interaction with HIF-1α promoted by TIPE. **D.** TIPE elevated PKM2 Ser37 phosphorylation in an ERK-dependent manner. **E, F.** TIPE enhanced PKM2 binding to the HRE for *LDHA* and *GLUT1* in a dimeric form dependent manner. **G.** PKM2 Ser37 mutation(S37A) inhibited the HRE activity induced by TIPE. **H, I.** PKM2 Ser37 mutation(S37A) decreased the expression of *LDHA* and *GLUT1* that promoted by TIPE. **$P<0.01$; ***$P<0.001$.

Figure 5. TIPE promotes melanoma cells proliferation and glycolysis depending on PKM2 dimerization. A. Pyridoxine (which facilitates PKM2 dimerization) treatment reversed the inhibition of cell proliferation that induced via interfering TIPE. **B.** Administration of TEPP-46 rescued TIPE overexpression induced cell proliferation. **C, D.** TIPE promoted melanoma clone formation in a dimeric PKM2-dependent manner. **E-G.** *In vivo* experiments showed that TIPE promoted melanoma proliferation via the dimerization of PKM2. **H-J.** Activation of PKM2 dimerization reversed the inhibition of glycolysis and glycolytic capacity after TIPE interfering. In contrast, inhibition of PKM2 dimerization altered this phenomenon (**K-M**). **N.** Pyridoxine increased the cell proliferation that was inhibited by TIPE-sh. It was rescued by inhibition of HIF-1α. **O.** TEPP-46 inhibited the TIPE-induced cell proliferation, and was rescued

by overexpression of HIF-1α. **P.** Pyridoxine enhanced the Warberg effect that was suppressed by TIPE-sh, and it was diminished via administration of PX-478(a HIF-1α inhibitor). **Q.** TEPP-46 can suppression of the Warberg effect that was increased by overexpression of TIPE, this phenomenon was diminished by overexpression of HIF-1α. *P<0.05; ***P<0.001.

Figure 6. TIPE fostered PKM2 dimerization increases melanoma stem-like phenomenon. A, B. TIPE increased the cancer stem-like phenotype markers, including *NANOG*, *NOTCH*, *OCT3/4*, *SOX2* and *BMI-1*, measured by *q*PCR. **C-F.** The effects of TIPE on the stemness of melanoma cells are shown by *in vitro* cell migration (**C**), sphere formation (**D**), and chemoresistance (**E, F**). **G.** Limiting dilution xenograft formation of A375 cells with TIPE interference. **H, I.** TIPE increased the expression of CD44^{+} cells in melanoma. **J, K.** The effects of TIPE on the stemness of melanoma cells following the administration of TEPP-46, and this phenomenon was reversed by overexpression of HIF-1α, as evidenced by sphere formation (**J**) and CD44^{+} cell population (**K**). *P<0.05; **P<0.01; ***P<0.001.

Figure 7. TIPE positively corelated with cancer stem cell markers and the levels of p-PKM2(Ser37). A, B. Higher expression of TIPE was

observed in melanoma tumor tissues than in the control, as evidenced by

immunohistochemistry. **C, D.** The expression of TIPE correlated well

with p-PKM2(Ser37) in melanoma tumor tissues. **E.** Similarly, a good

correlation was observed between TIPE, p-PKM2(Ser37), LDH, and

CD44 in mouse xenografts. **F-I.** In addition, the expression of TIPE was

positively correlated with CSCs markers, including *BMI1*, *NANOG*,

NOTCH1, and *OCT3/4*(*POU5F1*) in TCGA dataset. **J.** A brief model

depicting the functional impact of TIPE on metabolic reprogramming in

melanoma. ***P<0.001.