1 **A putative rhamnogalacturonan-II CMP-β-Kdo transferase identified using**
2 **CRISPR/Cas9 gene edited callus to circumvent embryo lethality.**

4 Yuan Zhang[1,2], Deepak Sharma[3,6], Yan Liang[1,2], Nick Downs[1,2], Fleur Dolman[4], Kristen
5 Thorne[3,6], Ian M. Black[3], Jose Henrique Pereira[1,2], Paul Adams[1,2], Henrik V. Scheller[1,2,5],
6 Malcolm O'Neill[3], Breeanna Urbanowicz[3,6*], Jenny C. Mortimer[1,2,4*]
7

[1]Joint BioEnergy Institute California, USA,
[2]Lawrence Berkeley National Laboratory, USA,
[3]Complex Carbohydrate Research Center, The University of Georgia, USA,
[4]University of Adelaide, Australia,
[5]University of California, Berkeley, USA
[6]Department of Biochemistry and Molecular Biology, University of Georgia
14

15 *Authors for correspondence:
16 **A/Prof Jenny C. Mortimer,** jenny.mortimer@adelaide.edu.au;  **A/Prof B. Urbanowicz**,
17 breeanna@uga.edu

## Abstract

Rhamnogalacturonan II (RG-II) is a structurally complex and conserved domain of the pectin present in the primary cell walls of vascular plants. Borate crosslinking of RG-II is required for plants to grow and develop normally. Mutations that alter RG-II structure also affect crosslinking and are lethal or severely impair growth. Thus, few genes involved in RG-II synthesis have been identified. Here we developed a method using CRISPR/Cas9-mediated gene to generate callus carrying loss-of-function mutations in the *MPG2* gene that encodes a putative family GT29 glycosyltransferase. Plants homozygous for this mutation do not survive. We show that in the callus mutant cell walls, RG-II does not crosslink normally because it lacks 3-deoxy-D-manno-octulosonic acid (Kdo) and thus cannot form the $\alpha$-L-Rha*p*-(1→5)-$\alpha$-D-kdo*p*-(1→ sidechain. We suggest that MGP2 encodes an inverting CMP-β-<u>K</u>do transferase (RCKT1). Our discovery provides further insight into the role of sidechains in RG-II dimerization.

The primary wall surrounding growing plant cells is a dynamic structure comprised primarily of
cellulose, hemicellulose and pectin [1]. The synthesis of these polysaccharides involves a large
number of genes, which together may account for up to 10% of a plant's genome. The Arabidopsis
genome encodes at least 567 predicted glycosyltransferases (GTs) across 44 Carbohydrate
Active Enzyme (CAZy) families. Many of these GTs are involved in cell wall glycan synthesis.
However, only a small number of them have been functionally characterized [2].

Rhamnogalacturonan II (RG-II) is a structurally complex and conserved domain of the
pectin present in the primary cell walls of all vascular plants. Borate crosslinking of this RG-II is
required for these plants to grow and develop normally [3–7]. Mutations that result in alterations to
RG-II structure also affect crosslinking and may be lethal or severely impair growth. Thus, few
genes involved in RG-II synthesis have been identified [8–11]. RG-II has a backbone of
homogalacturonan (HG) that is decorated with four structurally distinct side chains (A-D) and two
arabinofuranosyl (Ara*f*) substituents (**Fig. 1a**). The side chains are composed of 12 different
monosaccharides linked by at least 20 different glycosidic bonds. Thus, RG-II is the most
structurally complex polysaccharide yet identified in Nature [4,12–14].

Plants carrying mutations in genes that affect RG-II synthesis and structure typically
exhibit severe growth defects. For example, the Arabidopsis *mur1* mutant, which is impaired in its
ability to synthesize GDP-L-fucose (Fuc), is unable to form a normal side-chain A and has a dwarf
phenotype [15,16]. Similar phenotypes were obtained by silencing the major Golgi-localized GDP-L-
Fuc transporter in the *gfp1* mutant [17], and by mutations affecting the synthesis and transport of
other monosaccharides required to form RG-II, including GDP-L-galactose (Gal) [18,19]; UDP-D-
glucuronic acid (GlcA), a precursor for UDP-D-apiose (Api) [20]; UDP-D-Api/UDP-D-xylose (Xyl) [21];
and CMP-β-3-deoxy-D-manno-octulosonic acid (Kdo) [22].

Identifying the GTs involved in the formation of RG-II is required to understand the role of
this polysaccharide in cell wall integrity and plant growth. Reverse genetic approaches have had
limited success in identifying RG-II-specific GTs. This is primarily due to the deleterious effects
that even minor modifications to RG-II structure have on plant growth. To date only two GTs have
been implicated in RG-II synthesis. Rhamnogalacturonan Xylosyltransferase (RGXT) 1-4, a family
of GTs is believed to add xylose to side chain A [23–26]. Galacturonosyltransferase 4 (GAUT4) may
be responsible for forming the homogalacturonan backbone [27]. Loss-of-function mutations in
GAUT4 and RGXT4 are lethal. Thus, alternate strategies were used to gain insight into their
functions: RNA interference (RNAi) knockdown of *GAUT4* in switchgrass and poplar caused a
reduction in the amounts and crosslinking of RG-II [27]; A pollen-rescue strategy enabled the
generation of a homozygous RGXT4 (also known as MGP4) loss-of-function mutant, which

showed decreased xylose content and reduced RG-II dimerization [26]. Nevertheless, both

strategies have their limitations. Knockdown of target gene expression may be insufficient and

obscure the consequences of gene loss-of-function. The pollen rescue strategy is effective only

if lethality is associated with the defective male gametophyte. Both methods require extensive

crossbreeding and screening yet offer no assurance of achieving stable lines. Hence, there is a

need to establish a facile system to generate viable plant lines lacking expression of functional

GTs.

Glycosylinositol phosphorylceramides (GIPCs) are an important class of plasma

membrane lipids in plants (**Fig. 1b**) [28]. Previously, we observed that plants carrying mutations that

affect the glycosylation of GIPCs had severe growth defects, yet their null mutant callus grew well

[29–31]. GIPCs have been proposed to have a role in cell-cell adhesion [31,32] as has the formation of

a pectin network by borate crosslinking of RG-II. For example, reduced RG-II crosslinking in the

*mur1* mutant results in defective cell adhesion, specifically at the interface between stele and

cortex cells [33]. The "*rotten ear*" maize mutant also exhibits decreased cell adhesion as well as

disrupted cell wall organization in the ear that has been attributed to incomplete borate cross-

linking of RG-II [34]. This led us to hypothesize that engineered callus tissue may provide an

opportunity to identify genes responsible for glycan synthesis that are harmful to normal plant

growth and development. Thus, we developed CRISPR-mediated tissue transformation to

generate callus carrying loss-of-function GT mutations. We validated this approach with the well

characterized GIPC mannosyl-transferase1 (AtGMT1) (**Fig. 1b)** [30]. We then generated a loss-of-

function mutant of a CAZy family GT29 protein (Male gametophyte defective 2, MGP2) currently

annotated as a sialyltransferase-like protein in TAIR (arabidopsis.org). The callus mutant together

with NMR spectroscopic and mass spectrometric analysis of its RG-II provided evidence that

MGP2 is the CMP-β-Kdo transferase (RCKT1) required to form the $\alpha$-L-Rha$p$-(1→5)-$\alpha$-D-Kdo$p$-

(1→ side chain of RG-II.

# Results

**Arabidopsis GT29 family members are highly co-expressed with primary cell wall- and
GIPC-related genes**

To identify GTs implicated in cell-to-cell adhesion processes, we used bioinformatics and publicly

available arrays of 113 developmental RNA seq datasets, representing 11 distinct tissue and

organ types to identify genes that are co-expressed with Arabidopsis *GMT1* [35,36]. Disrupting this

gene causes severe plant growth defects and reduces cell-cell adhesion [30,32]. We then used SUBA

(Subcellular Location Database for Arabidopsis Proteins) to refine our list to 75 candidates with

likely Golgi localization (**Supplementary Data 1**). These candidates were regarded as putative

GTs relevant to cell-cell adhesion. Somewhat unexpectedly, all three Arabidopsis family GT29

members - At1g08280 (GALT29A), At1g08660 (RCKT1, also known as MGP2), and At3g48820

(SIA2) - were included. They are annotated as sialyltransferase-like (ST-like) proteins since they

possess motifs characteristic of mammalian family GT29 STs. However, the plant enzymes have

no discernible ST activity [37]. This is not entirely unexpected since plants produce little if any sialic

acid [38].

We then considered the possibility that the plant GT29 enzymes use a nucleotide sugar

donor substrate structurally related to CMP-Sialic acid, the donor for STs. Indeed, *RCKT1* and

*SIA2* have been proposed to encode enzymes that catalyze the transfer of 2-keto-3-deoxy-D-

manno-octulosonic acid (Kdo) or 2-keto-3-deoxy-D-lyxo-heptulosaric acid (Dha) [39,40]. The

activated form of Dha remains to be identified. CMP-$\beta$-Kdo is the activated form of KDO and the

family GT29 proteins utilize an inverting mechanism for glycosyl transfer, indicating they would

be able to form the $\alpha$-2,3-bond that links Kdo to the RG-II backbone and initiates formation of

sidechain C (**Figure 1**).

**Creating biallelic mutant calli carrying GT knockouts**

No homozygous mutants of *GALT29A*, *RCKT1*, (*MGP2*), and *SIA2* have been isolated,

suggesting they are lethal or severely harm the mutant plants. This also suggests that these GTs

have an essential role in plant development. This led us to develop a CRISPR/Cas9-mediated

tissue culture transformation strategy to generate loss-of-function callus to study the function of

GTs. CRISPR/Cas9 technology is a powerful tool for targeted genome editing in plants and has

considerable potential for generating knockout mutants. Here, we optimized a tissue culture-

based transformation method [41], and used it to introduce the CRISPR/Cas9 system with the aim

of circumventing the need to produce whole plants and viable seeds (**Supplementary Fig. 1**).

The isolated clonal knockout lines can be indefinitely propagated as callus tissue, which provides

material suitable for biochemical and functional characterization of plants with primary cell wall

chemotypes.

We used a modular cloning system [42], which includes a T-DNA destination vector, a Cas9

entry vector, and a guide RNA (gRNA) entry vector (**Supplementary Fig. 2**). To increase the rate

of editing, each construct contained three sgRNAs for the target gene (**Fig. 2a, Supplementary**

**Table 1**). Since CRISPR/Cas9 editing typically generates small insertions and deletions (indels)

through non-homologous end joining (NHEJ), the disruption of gene function primarily stems from

frameshift mutations. To maximize the impact of such disruptions, the sgRNAs were designed to
target the protein-coding region proximal to the 5'-untranslated region (5'-UTR).

**Validating the system with a known GT as the target for knockout**

We first targeted *AtGMT1* to test the effectiveness of our CRISPR/Cas9 toolbox and tissue culture
transformation method in creating GT knockout mutant calli. We targeted three unique sites in
*AtGMT1*: two in Exon1 and one in Exon2 (**Fig. 2b**). The primary roots of ten-day-old Arabidopsis
seedlings were used for *Agrobacterium*-mediated transformation, yielding 16 viable calli out of
300 infected root explants (5.3%). The stable integration of the CRISPR/Cas9 cassette in these
transformants was indicated by their resistance to kanamycin and the detection of DsRed
fluorescence, and further validated via PCR.

To characterize the editing events in each transformant, the genomic region spanning the
three target sites was amplified by PCR, Sanger sequenced, and then analyzed using DECODR
v3.0 (https://decodr.org/) [43]. Cas9-induced mutations were introduced into 13 lines (81%) at the
sgRNA3 target site in Exon2. This gave us 7 homozygous mutants (with two identical mutant
alleles), 1 biallelic mutant (with two distinct mutant alleles) and 5 chimeric mutants (carrying two
or more types of alleles, with or without mutation) (**Fig. 2c**). The 7 editing events detected in the
homozygous lines consisted of 5 frameshifting indel mutations, expected to result in truncated
proteins, and 2 in-frame deletion mutations that would lead to the loss of one amino acid (**Fig.
2c**). The single biallelic mutant had one allele with a frameshift insertion, and another with a frame-
preserving deletion (**Fig. 2c**). All the transgenic lines were unedited at the sgRNA1 and sgRNA2
target sites of Exon1, while 3 lines completely escaped targeted editing of *AtGMT1* by the
CRISPR/Cas9 complex. Thus, our strategy enables the efficient creation of clonal knockout lines
in the form of callus, where every individual cell carries the same frameshift mutation within the
target gene.

**Validating the function of *gmt1* by GIPC analysis**

We analyzed the GIPCs of the callus lines to assess the suitability of homozygous frameshift
mutant calli for decreasing or eliminating GT activity. A transgenic callus line without CRISPR
editing was included as a non-edited (NE) control. We also examined callus derived from Col-0
seedling roots and a *gmt1-3* T-DNA insertional mutant.

GIPCs were extracted from callus tissue grown in liquid media and the degree of
glycosylation was determined using thin layer chromatography (TLC) [44]. The GIPC profiles of the
*gmt1-4* and *gmt1-3* mutants were clearly different from those of Col-0 and the NE control

(**Supplementary Fig. 3).** Notably, the GIPC bands exhibited a larger shift in both *gmt1* mutants,
indicating a reduction in molecular weight due to the absence of mannosylation [30]. Furthermore,
the bands displayed a compact and tightly clustered appearance in contrast to the dispersed
bands observed in the negative control lines. This distinctive feature marked a significant
decrease in the diversity of GIPC species when normal mannosylation of the sugar head group
is absent.

We next determined the mannose content of the extracted GIPCs. The monosaccharides
were released by hydrolysis with trifluoroacetic acid (TFA) and quantified by high-performance
anion exchange chromatography coupled with pulsed amperometric detection (HPAEC-PAD) [30].
The mannose content was substantially reduced (3-7 fold) in both *gmt1* mutants compared to that
in the wild-type and NE control (**Supplementary Table 2**). Our qualitative TLC assay and glycosyl
residue composition analysis demonstrate that the *gmt1* CRISPR callus mutant recapitulates the
GPIC chemotype of the *gmt1-3* T-DNA insertional mutant.

**Creating homozygous *RCKT1* knockout mutant callus**

We next targeted the *RCKT1* gene identified in our bioinformatic screen. The tri-gRNA module in
the transformation vector was designed to target 3 distinct sites within the *RCKT1* coding region
(Exon 2, 3, and 5) (**Fig. 2d**). We obtained 16 viable calli out of 410 infected root explants (3.9%
efficiency). Cas9-mediated editing was restricted to the sgRNA2 target site of *RCKT1*. Genotyping
of the transgenic lines identified the presence of 1 homozygous mutant, 2 heterozygous/biallelic
mutants, 11 chimeric mutants, and 1 NE control line. The homozygous mutant *rckt1-21* carried a
"-49bp/-49bp" biallelic mutation, disrupting the open reading frame and introducing an early stop
codon (**Fig. 2e**). Consequently, the predicted product of the *RCKT1* mutant allele in *rckt1-21* is a
truncated protein consisting of 104 amino acid residues at the N-terminus, with 73 residues
preserved and 31 altered. This mutant RCKT1 variant is expected to lack the catalytic domain,
rendering *rckt1-21* an ideal knockout line for cell wall analysis. In contrast, the two heterozygous
mutants, *rckt1-1* and *rckt1-9*, harbored a "-1/-3" and "+1/-6" biallelic mutation, respectively (**Fig.
2e**). These mutations resulted in one allele carrying a frameshift indel mutation (-1 or +1), while
the other allele containing a frame-preserving deletion (-3 or -6), which led to the loss of one or
two amino acid(s). It's unknown whether such biallelic mutations would cause the complete loss
of *RCKT1* function in these heterozygous mutants. The chimeric mutants possessed wild-type
alleles at varying percentage (6-52%), which remained recalcitrant to CIRSPR/Cas9 editing.

The homozygous *rckt1-21*, the heterozygous/biallelic *rckt1-1* and *rckt1-9* are gene edited

lines that can be used to study AtRCKT1 function. The non-edited *rckt1-6*, with a T-DNA insertion

but lacking any *AtRCKT1* editing is a control.

**The absence of RG-II Kdo from the *rckt1* mutant**

To explore the effect of *RCKT1* mutation on cell wall glycosyl residue composition, we grew the

*rckt1* callus lines and controls in liquid media, and then isolated cell walls for all lines as alcohol-

insoluble residue (AIR). The monosaccharide compositions of the walls from the *rckt1* mutants

and the wild-type and the non-edited control lines were similar (**Fig. 3a-b, Supplementary Data**

**2**). However, we were unable to detect Kdo in intact AIR, thus we isolated RG-II from the callus

walls. To this end we treated the AIR with an endopolygalacturonase (EPG) and separated the

solubilized products by size-exclusion chromatography (SEC) [45] to obtain an RG-II rich fraction

(**Fig. 3c**). This procedure also shows the relative abundance of the RG-II monomer and dimer

present in the walls at the time of extraction. The dimer accounts for ~80% of the RG-II in Col-0

and the NE control (**Fig. 3c**, **Supplementary Table 3),** a result consistent with previous reports

[9,19,46]. However, in both the homozygous *rckt1-21* mutant and the heterozygous/biallelic *rckt1-9*

mutant the monomer accounted for at least 74% of the RG-II (**Fig. 3c, Supplementary Table 3**),

showing that the ability to form the dimer is diminished in the *rckt1* mutants.

The absence of Kdo in the RG-II from the two *rckt1* mutants was first established by

solution state nuclear magnetic resonance spectroscopy (NMR). To facilitate the detection of

signals corresponding to Kdo and Dha, the RG-II (dimer) was treated with dilute HCl to generate

the monomer. Resonances diagnostic for Kdo (H3, δ 1.88 ppm) and Dha (H3, δ 1.78 ppm) were

clearly discernible in the wild-type RG-II (**Fig. 4a-b**). The Kdo signal was absent in the *rckt1*

mutants (**Fig. 4c-d**). A 2D TOCSY NMR experiment showed that only the wild-type RG-II

contained resonances characteristic of H3 and H3' (δ 1.88 and 2.13 ppm) correlated with H4 (δ

4.13) of Kdo, whereas the Dha H3 and H3' resonance (δ 1.78 and 2.05 ppm) correlated with H4

(δ 4.08) were present in both the controls and the *rckt1* mutants (**Fig. 5**).

We next prepared the trimethylsilyl methyl-ester methyl glycoside (TMS) derivatives of the

RG-IIs for analysis by gas-chromatography with electron impact mass spectrometry (GC-EI-MS)

[45] . Dha is present in the controls and the *rckt1* mutants, whereas Kdo is detected in the controls,

but absent in the *rckt1-21* homozygous mutant and the *rckt1-9* heterozygous/biallelic mutant

(**Supplementary Fig. 4**). In summary, our NMR spectroscopic and glycosyl residue composition

analyses    conclusively    demonstrate    that    the    homozygous    mutant    and    the    *rckt1-9*

heterozygous/biallelic mutant produce RG-II that lacks discernible amounts of Kdo. This evidence
supports our hypothesis that RCKT1 is the Kdo transferase required to form RG-II side-chain C.

**Predicted structure of RCKT1 in the presence of CMP-β-Kdo**

Determining the biochemical properties of RCKT1 is challenging since its acceptor is unknown
and the CMP-β-Kdo donor has an estimated half-life of ~34 minutes [47,48]. Thus, we used
AlphaFold [49] to predict the conformation RCKT1 and gain some molecular insight into its function.
The model shows a N-terminal transmembrane domain from residue 1 to 36 with the
transmembrane α-helix between residues 12 and 36. RCKT1 is a mix of αβ folds, comprising 1
central β-sheet consisting of 7 parallel β-strands surrounded by 14 α-helices and 2 small β-sheets
(**Fig. 6a-b**). The mammalian α-2,3-sialyltransferase ST3Gal I (PDB ID 2WNB) from *Sus Scrofa*
(wild pig) [50] shares 32% amino acid sequence identity with RCKT1 and represents the closest
structural homolog in the Protein Data Bank. Both enzymes are members of the GT29 family,
whose structurally characterized members have been shown to adopt a modified GT-A fold, but
lack the characteristic DxD motif [51]. The structure of the GT42 sialyltransferase CstII (PDB ID
1RO7) from *Campylobacter jejuni* [52], which has similar folding to RCKT1, was solved in complex
with CMP-3FNeuAc. A superposition of CStII and RCKT1, allowed us to identify the likely position
of the CMP binding site and consequently model CMP-β-Kdo into the substrate donor active site
of RCKT1. This putative binding site suggests that the main and side chain atoms of Phe306,
Thr307 and Thr316 directly contact the cytidine nucleotide. Moreover, Trp315 makes aromatic
stacking interactions. The ribose portion of the substrate interacts with the main chain N atoms of
Asn186 and Gly287, and the Kdo sugar region showed 2 hydrogen bonds with Thr236 (**Fig. 6c**).
It will be important to test the importance of these predicted residues in future experiments.
253

# Discussion

**RG-II structure integrity is important for its function**

It is widely recognized that the structure of RG-II must be maintained to ensure it can form a dimer
normally and that this is a key factor in its functions related to plant growth, development, and
viability. This view has primarily been supported by research on plant mutants deficient in RG-II
specific GTs, enzymes and transporters responsible for RG-II specific nucleotide sugars, and
boron uptake/transport mutants. Knockdown of the RGXTs involved in side-chain formation [26],
and disruption of the synthesis and transport of RG-II nucleotide sugars, such as GDP-L-Fuc [15–

[17], GDP-L-Gal [18,19], UDP-D-Api [20], all lead to severe growth and developmental defects in plants.

The truncation of side-chain A, due to the absence of D-Xyl, L-Fuc, L-Gal or D-Api is believed to

be responsible for the defective RG-II dimerization. These data are consistent with the notion that

the borate cross-link formed between two RG-II monomers involves the chain A by D-Api residue

[3,4].

Our study of RCKT1 introduces a new perspective by showing that preventing the

formation of side-chain C, a disaccharide seemingly distant from the boron-binding D-Api residue,

also disrupts dimer formation. This broadens the prevailing hypothesis that any alteration to the

RG-II structure may have a substantial impact on its dimerization and compromise the formation

of a normal pectic network *in muro*. Our CRISPR-mediated strategy provides a basis to test this

hypothesis. Generating loss-of-function mutants of RG-II GTs, would allow us to create a diverse

array of RG-II glycoforms *in situ*, allowing for an exploration of the relationship between RG-II

structure and functionality. Beyond RG-II dimerization, we can also delve into whether structural

modifications of RG-II influence potential interaction with other pectic domains and various cell

wall components. This would open new avenues for us to probe how the pectic network

contributes to the mechanical strength and physical properties of the plant primary wall, which

play a substantial role in shaping the growth and development of plant cells, tissues, and organs

[53–55]. The formation of a pectic network involves multiple levels of crosslinking, including but not

limited to the backbone glycosidic linkages of the three pectic domains, the borate ester

crosslinking of RG-II, and the calcium crosslinking of HG [56]. Although the specific role of RGII-

boron cross-linking in maintaining the physical property of plant cell wall is far from clear, it has

been suggested to regulate wall pore size [57], mechanical strength [58] and extensibility [1].

**Challenges to confirm the enzymatic activity of RCKT1 *in vitro***

Our data have shed light on the role of RCKT1 in the addition of Kdo to RG-II. An *in vitro* assay

for enzymatic activity will be required to increase our understanding of RCKT1's function as an

RGII-specific Kdo transferase. However, this is challenging for several reasons including the

difficulties of heterologously expressing a functional RCKT1 enzyme, the availability of the donor

substrate CMP-β-Kdo, and the availability of appropriate RG-II glycoforms as acceptor substrates.

Plant cell wall GTs have been challenging to express as stable and active recombinant

proteins in bacteria and yeast [59,60]. Our attempts to express RCKT1 using the *Escherichia coli*

Origami B strain was not successful. The Human Embryonic Kidney (HEK293) cell system [61],

provides an alternative technology as it has been used to produce several plant GTs [62–68].

However, an equally large challenge arises from the instability of CMP-β-Kdo, the donor substrate

for the RCKT1 enzyme. CMP-β-Kdo is rapidly hydrolyzed in aqueous buffers (half-life of ~34 mins)

resulting in the formation of CMP and free Kdo [47,48]. To address this, several groups have

generated CMP-β-Kdo *in situ* using CTP, Kdo and a CMP-β-Kdo synthetase (KdsB) purified from

*E. coli*. This strategy has proven successful in studies focused on β-Kdo-transferases involved in

bacterial glycolipid biosynthesis [69–71]. Finally, an appropriate RG-II glycoform, capable of

accepting Kdo from the nucleotide sugar donor CMP-β-Kdo, is required. It is possible that an RG-

II lacking side-chain C could serve as an ideal acceptor for Kdo. The *RCKT1* callus mutant

generated in this study could provide this glycoform. Alternatively, sidechain C could be removed

using the bacterial glycanase described by Ndeh et al [14]. However, the possibility cannot be

discounted that a glycoform lacking other side-chains in addition to side-chain C is the acceptor.

Tools are now available to modify RG-II structure *in vitro* and generate different glycoforms in

amounts suitable for GT assays [14]. Nevertheless, identifying the appropriate acceptor molecule

remains a considerable challenge.

It will also be important to obtain structural information to understand the molecular

mechanism of its catalytic activity. This will allow comparison of the sialytransferase motifs from

the closely related animal STs with the plant RCKT1 to identify potential residues responsible for

substrate specificity. Future work will include generating a series of RCKT1 variants for kinetic

analysis. It will also be interesting to compare this protein to the previously identified AtKDTA

KDO transferase candidate, a mitochondria-localized protein which is thought to have a role in

glycosylating an as-yet unidentified lipid signaling molecule [72].

**Advantages of, and potential extensions to, a callus-based gene editing method**

We have demonstrated the utility of our callus-based gene editing method for generating KO

mutations in two GTs that are required for normal plant growth and development. We believe that

this approach could serve as a tool for both reverse and forward genetic investigations. One

application would be the creation of callus carrying mutations in multiple different GTs. Gene

redundancy significantly hampers the characterization of plant GTs [59]. For example, the GAUT

family of Arabidopsis has 15 members. Only a few of these have yielded mutants with observable

phenotypes that assist in revealing gene function [27,73,74]. The multiplexed CRISPR/Cas9 toolbox

employed in our study allows the assembly of up to eight sgRNAs into a single T-DNA vector, a

crucial feature that can enable the simultaneous targeted editing of multiple genes [42].

Our method can be adapted for use in forward screens to identify candidate GTs involved

in specific biological processes, such as cell-cell adhesion. Known cell adhesion mutants exhibit

a decreased regeneration phenotype [75]. It should be possible to create a library of CRISPR callus

mutants for the 75 candidate genes in our target clade, and assess organ regeneration capability.
The CRISPR/Cas9 system has been used with rice to generate a large-scale mutant collection
[76].

Improvements to gene editing efficiency would enable this method to be more widely
implemented. Only one of the three bioinformatically predicted top-performing sgRNAs created a
CRISPR mutation at the target sites of *GMT1* and *RCKT1*. Liang and colleagues have developed
a prescreening method to assess sgRNA efficiency via the tobacco (*Nicotiana benthamiana*)
transient expression system [77], that could be applied to select highly efficient sgRNAs for the
target genes. This is a rapidly developing area of research, with recent improvements including
the use of stronger promoters to drive higher expression of the sgRNAs and Cas9, the utilization
of intronized Cas9 [78], or alternative Cas enzymes, can be used to further optimize its editing
efficiency in plant species.
340
341

## Materials and Methods

**Plant material and incubation conditions**

*Arabidopsis thaliana* ecotype Col-0 and *gmt1-3* seeds were surface sterilized for 10 minutes using a 10% (v/v) commercial bleach (Clorox, 6.5%, v/v) in aq. Ethanol (100% v/v), and then washed twice with 100% ethanol, and allowed to air dry overnight. Sterilized seeds were placed on B5 agar media (**Supplementary Table 3**) and stratified for three days at 4°C in the absence of light. The plates were then transferred to a growth chamber (8,000 lux, 16:8 L:D, 22°C, 75% humidity). After 10 days, the primary roots were isolated from the seedlings for root transformation as described below. *gmt1-3* allele is as described in [30]. The callus culture of *gmt1-3* was derived from roots as described in [79].

Calli were maintained by sub-culturing every 14 days onto freshly prepared callus induction media (CIM) (**Supplementary Table 3**). The callus selected for subculture should be creamy and white, indicating healthy status. Calli plates were kept in the dark in a growth chamber (25°C, 50% humidity). To grow cell suspension culture in liquid media, callus tissue (approx. 100 mg) was inoculated into 1 L flask containing 250 mL liquid CIM media and shaken at 100 rpm in the dark at 23°C, a method modified from [80]. Callus from liquid cultures was harvested after 14 days of growth. The tissue was collected into falcon tubes and lyophilized at -50°C for 5 days. Fully lyophilized callus tissue was stored at -80°C for downstream biochemical analysis.

**Screening GTs important for plant cell-cell adhesion by a bioinformatic pipeline**

To identify putative GTs involved in the plant cell-cell adhesion process, we set up a bioinformatics pipeline based on the following filtering process.

Filter I: Arabidopsis GT genes were obtained from the CAZy database (http://www.cazy.org/). Genes corresponding to family GT1 proteins were removed since they are largely involved in the glycosylation of small molecules and are unlikely to be involved in cell wall biosynthesis. 441 genes from 49 GT families were identified.

Filter II: At*GMT1*, which encodes a GIPC mannosyltransferase was selected as the query gene for co-expression analysis. Co-expression analysis used 113 developmental RNA seq datasets across 11 tissue and organ types of Arabidopsis, and ThaleMine (https://bar.utoronto.ca/thalemine/begin.do) [81]. Genes with similar expression patterns across the datasets were grouped based on hierarchical clustering using Cluster 3.0 (http://bonsai.hgc.jp/~mdehoon/software/cluster/). The gene clusters were visualized using TreeView (https://jtreeview.sourceforge.net/) and the clade containing the query gene *GMT1* was extracted.

Filter III: Genes encoding GTs localized in Golgi. The subcellular localization of the extracted GTs was obtained by SUBA (https://suba.live/). This step resulted in 75 Golgi-localized candidates as putative GTs most likely to be relevant to plant cell-cell adhesion.

**Guide RNA design and transformation vector construction**

For the three sgRNA designed to target each gene, DNA oligonucleotides were synthesized by Integrated DNA technologies (**Supplementary Table 4**) and then annealed and phosphorylated. The phosphorylated sgDNA were individually cloned into the BsmBI site of the Golden Gate entry vectors pYPQ13XA or pYPQ13XB to create pYPQ131A-AtU6-gRNA1, pYPQ132B-AtU3-gRNA2,

and pYPQ133A-AtU6-gRNA3. The three gRNA expression cassettes were then cloned into the recipient vector pYPQ143Amp by Golden Gate assembly to create a vector that contains the Tri-sgRNA module. The Tri-sgRNA module and Cas9 were assembled into the pTkan destination vector by Gateway cloning.

This modular cloning toolkit for CRISPR/Cas9 vector construction was originally developed by Lowder et al. [42], and modified in this study. The antibiotic selection markers of the Golden Gate recipient vector pYPQ143 and the Cas9 entry vector pYPQ150 from the original toolkit were replaced by the ampicillin resistance marker (AmpR) to be compatible with the pTKan destination vector we had in house.

Whole plasmid sequencing was performed to validate the sequences of the different components in the transformation vectors targeting AtGM1 and AtRCKT1. The final constructs were transformed into *Agrobacterium fabrum* (previously known as *Agrobacterium tumefaciens*) GV3101 for Arabidopsis root transformation.

**Arabidopsis root transformation**
*Arabidopsis thaliana* ecotype Col-0 roots were transformed with *A. fabrum* GV3101 carrying the CRISPR/Cas9 vectors. The method was modified from Vergunst et al. [41]. Sterilized Col-0 seeds were sown on B5 agar media (**Supplementary Table 3**) and grown in a growth chamber (8,000 lux, 16:8 L:D, 22C°, 75% humidity). Primary roots were isolated from 10d old seedlings using a sterile razor blade and transferred onto CIM media (**Supplementary Table 3**) and grown for three days. *A. fabrum* GV3101 carrying the transformation vector was grown for two days at 30°C on LB solid media with antibiotics (rifampin, gentamicin, spectinomycin). A single *A. fabrum* colony was inoculated into liquid LB media with antibiotics (rifampin, gentamicin, spectinomycin) and cultured overnight at 30°C with shaking at 200 rpm. *A. fabrum* cells were pelleted and then suspended in liquid B5 media (final $OD_{600}$ = 0.1). The isolated primary roots were soaked for five minutes in the *A. fabrum* suspension, then sliced into 0.5-1 cm explants using a sterile razor blade. The root explants were blot dried on sterile filter paper and plated onto CIM media containing D-glucose (1.8 g/L) and acetosynringone (0.1mM). Root explants were co-cultivated with *A. fabrum* for three days at 25°C under dim light (2,000 lux, 16:8 L:D). The root explants were then washed four times with Timentin-supplemented (100mg/L) B5 liquid media and plated onto CIM media supplemented with antibiotics (50 mg/L Kanamycin and 100mg/L Timentin). After 14 days, the explants were transferred to fresh CIM media containing Kanamycin and Timentin. Calli resistant to Kanamycin were collected and sub-cultured every 14 days.

**CRISPR editing analysis of the transgenic callus mutants**
Callus tissue was sampled from each independent transformation event, flash frozen in liquid nitrogen, and then milled with metallic beads using a TissueLyzer (Qiagen) for subsequent genomic DNA (gDNA) extraction. Genomic DNA was extracted from callus using the QIAGEN DNeasy Plant Mini Kit®. Primers that amplified the genomic region of the guide RNA (gRNA) target sites were used to genotype the material (**Supplementary Table 5**). PCR amplicons were column-purified using the Zymo Research DNA Clean & Concentrator® kit, and Sanger sequenced by Azenta (https://www.azenta.com/). Decodr v3.0 (https://decodr.org/) was used to deconvolute the sequenced amplicon for mutation analysis.

**GIPC purification and analysis by TLC**

GIPC analysis by TLC was performed as described by Jing et al. [44]. Lyophilized callus tissue (~0.5 g dry weight) harvested from three distinct liquid cultures were combined and ground to a powder with metallic beads using a TissueLyzer (Qiagen). Sphingolipids were extracted from the ground tissue by using the lower layer of the isopropanol/hexane/water (55:20:25 v/v/v) extract followed by deesterification with 4 mL of 33% methylamine solution in ethanol/water (7:3 v/v) as previously described [82]. The water used in this experiment is HPLC-grade. To enrich the GIPCs, the dried extracts were suspended in 1 mL of enrichment solution (chloroform/ethanol/ammonia/water [10:60:6:24 v/v/v/v]) and kept overnight at room temperature with agitation (2 rpm). The GIPCs were then isolated using a weak anion exchange SPE cartridge (Strata X-AW; Phenomenex) as described [29]. GIPCs bound to the cartridge were eluted with 3 mL of enrichment solution and dried under a flow of nitrogen gas or a stream of clean dry air. The dried GIPCs were suspended in solution A {chloroform/methanol/[4M ammonium hydroxide in 1.8M ammonium acetate] (9:7:2 v/v/v)} and separated by thin layer chromatography (TLC). The separations were performed on high-performance TLC Silica gels on glass plates (Merck) with solvent A as the mobile phase. GIPCs were stained with primuline (10 mg in 100 mL acetone/water [8:2 v/v]), and visualized under a fluorescent blue light (460 nm) [83].

**GIPC glycan profiling by HPAEC-PAD**

Purified GIPCs were treated for 1 hour at 120°C with 2M trifluoroacetic acid (TFA, 1 mL), cooled on ice, and dried overnight using a vacuum concentrator. The residues were dissolved (2-5 fold) in HPLC grade water and the liberated monosaccharides analyzed by high-performance anion-exchange chromatography with a pulsed amperometric detector (HPAEC-PAD) as described [30].

**Preparation of plant cell walls as alcohol insoluble residue (AIR)**

AIR was prepared as previously described [29]. Briefly, lyophilized callus was milled into a fine powder and then suspended in 96% (v/v) ethanol. The suspension was kept for 30 min at 70°C, and the suspension then cooled and centrifuged (15 mins, 18,500 rcf). The pellet was then suspended in ethanol (96% v/v), vortexed, and then centrifuged again (15 mins, 18,500 rcf). The pellet was then suspended in methanol:chloroform (2:3 v/v) and kept overnight at 4 °C with agitation (300 rpm). Following centrifugation (15 min, 18,500 rcf), the pellet was treated for 1h at 4°C methanol:chloroform (2v:3v) (1h, 300 rpm,). Following centrifugation (15 min, 18,500 rcf), the pellet was washed with ethanol: 100% (v/v), 65% (v/v), 70% (v/v), 80% (v/v), 100% (v/v). Each wash involved a 5-minute centrifugation (18,500 rcf) and removal of the supernatant in between, with a thorough vortex to suspend the pellet. The resulting alcohol insoluble residue (AIR) was dried at 50 °C overnight in a vacuum concentrator or oven.

Starch was removed by treating the AIR (~10 mg) for 10 mins at 75°C with agitation at 300 rpm) with $\alpha$-amylase (3 U/mL) in 50 mM MOPS, pH 7.3). Pullulanase (1 U/mL) and amyloglucosidase (3 U/mL) X mM Na acetate, pH 4.5, were then added, and the suspensions kept for a further two hours at 40°C (with agitation at 300 rpm). Following centrifugation, the pellet was washed with the following ethanol series: 100% (v/v), 70% (v/v), 70% (v/v). The destarched AIR was dried overnight at 30°C in a vacuum concentrator.

**Cell wall monosaccharide composition analysis**

Non-cellulosic sugars were liberated from de-starched AIR by TFA hydrolysis for one hour at 120°C. The supernatant containing the liberated monosaccharides was collected after centrifugation. Residual monosaccharides in the remaining pellet were further extracted using water and combined with the supernatant. The combined supernatant was dried overnight at 30°C in a vacuum concentrator. The residue was suspended in HPLC-grade water and filtered through a nitrocellulose filter (0.45 µM). The non-cellulosic sugars in the filtrate were diluted (4-5 fold) with water and filtered through a nitrocellulose filter (0.45 µM) prior to HPAEC-PAD analysis [30].

The TFA-insoluble AIR pellet was then subjected to Saeman hydrolysis to hydrolyze cellulose. The pellet was suspended in 72% (v/v) sulfuric acid and kept for one hour at room temperature. Water was added to dilute the sulfuric acid to 1M and the suspensions then heated for 3h at 100°C. The suspensions were then neutralized by adding barium carbonate and centrifuged. The supernatant containing was dried overnight at 30°C in a sample concentrator. The residue was suspended in water and filtered through a nitrocellulose filter (0.45 µM). The sugars in the filtrate were diluted with water and filtered prior to HPAEC-PAD analysis. The water used in this experiment is HPLC-grade only.

**Isolation of RG-II from AIR of controls and the *rckt1* mutants**

The callus AIR (500-700mg) extracted from three distinct liquid cultures were combined and suspended in NaOAc (50 mM, pH 5.0) at a concentration of 25mg/ml, and then treated for 24 hrs at 37 °C with 5 U/g endopolygalacturonase (EPG) and 10 U/g of pectin methyl esterase (PME) with shaking at 150 rpm. The EPG solublized material and the insoluble residue were separated by filtration through a nylon mesh (100 µm). The insoluble material retained on the mesh were treated again with EPG and then filtered as before. The EPG-soluble materials were obtained by combining the filtrates after the two treatments. The x mL solution in total were then fractionated by preparative size-exclusion chromatography (SEC) on a column Superdex Increase 75, (10/300 GL, Cytiva, USA, column L × I.D. 30 cm × 10 mm, 9 µm) with refractive index (RI) detection. The column was eluted with ammonium formate (50 mM, pH 5.0) at 0.5 mL/min. Fractions enriched in RG-II were collected manually, dialyzed (3500 Dalton MWCO) against deionized water, and freeze-dried.

**RG-II glycosyl residue composition analyses**

To specifically detect Kdo and Dha, the glycosyl residue of RG-II were analyzed as their trimethylsilyl (TMS) methyl glycoside derivatives by gas-chromatography with electron impact mass spectrometry (GC-EI-MS) as described in [45]. In general, RG-II (100 µg) was suspended in 300 µL methanolic HCl (1 M) in screw-top glass tubes secured with Teflon-lined caps and heated at 80 °C for 18 hrs. After cooling to room temperature, the solutions were concentrated to dryness under the nitrogen stream. The released methyl glycosides and methyl glycoside methyl-esters were then reacted for 30 min at 80 °C with Tri-Sil® (ThermoFisher, USA). GC-EI-MS analysis of the TMS methyl glycosides was performed on an Agilent 7890A GC interfaced to an Agilent 5975C mass selective detector, with a Supelco Equity-1 fused silica capillary column (30 m × 0.25 mm ID).

**Preparation of RG-II monomer for NMR spectroscopic analysis**

The RG-II dimer was converted to the monomer as previously described [45]. Briefly, the RG-II enriched material obtained by SEC was treated for 1 h at room temperature with HCl (0.1 M). The solution was then dialyzed (3500 Dalton MWCO) against deionized water and freeze-dried.

**1D $^1$H-NMR and 2D TOCSY spectroscopy of RG-II monomer**

RG-II (1-2 mg) was dissolved in $D_2O$ (0.2 mL, 99.9%; Cambridge Isotope Laboratories, Tewksbury, MA, USA) and placed in a 3 mm NMR tube [45]. Spectra were obtained using a Bruker NMR spectrometer (Agilent Technologies) operating at 900 MHz using a 5 mm cold probe and a sample temperature of 25⁰C.  The 1D $^1$H and 2D TOCSY spectra recorded using standard Bruker pulse programs. Chemical shifts of the RG-II glycosyl residue were measured relative to internal $D_2O$ (δH 4.707). Data were processed using MestReNova software (Mestrelab Research S.L., Santiago de Compostela, Spain).

**Statistical analysis**

One-way ANOVA were used for statistical analysis.

**Accession Numbers**

Sequence data from this article can be found at The Arabidopsis Information Resource (TAIR) under accession number At3g55830 (*AtGMT1*), At1g08660 (*AtRCKT1*), AT3G48820 (*AtSIA2*), and AT1G08280 (*AtGALT29A*).

## Figures and Tables

**Fig. 1**| Structure of rhamnogalacturonan-II (RG-II) and Glycosylinositol Phosphorylceramide (GIPC). **a**, Schematic structure of RG-II (modified from Sechet *et al.*, 2018). Monosaccharides are shown using the Symbol Nomenclature for Glycans [84,85]. **b**, Example structure of a GIPC showing the ceramide (black), inositol phosphate (yellow), glucuronic acid (GlcA, blue), and mannose (Man, green) (modified from Mortimer and Scheller, 2022). Man-GlcA-IPC is the dominant headgroup structure in wild-type Arabidopsis callus tissue. Inset: The synthesis of Man-GlcA-IPC catalyzed by Arabidopsis GMT1 (AtGMT1). Man and GlcA are shown using the same Symbol Nomenclature for Glycans as in **a**.

**Fig. 2| Creation of CRISPR/Cas9-edited Arabidopsis callus lines with different mutations on *GMT1* and *RCKT1*. a**, Illustration of the T-DNA transformation vector used for CRISPR editing of the target gene. Arabidopsis U6 (AtU6) promoter is used to express gRNA1 and gRNA3, while Arabidopsis U3 (AtU3) promoter is used to express gRNA2. **b**, Schematic diagram of the *GMT1* gene structure with gRNA targeting sites labeled. The black boxes represent exons; the grey boxes represent 5'-UTR and 3'-UTR; the black lines represent introns. CRISPR/Cas9 editing was only detected at the gRNA3 (red) target site. **c**, Gene-editing events at the gRNA3 target site of the *gmt1* callus mutants. Seven homozygous lines were identified: five are *gmt1* knockout mutants with frameshifted and truncated GMT1 protein, and two are mutants with the loss of a single amino acid from the protein sequence. One heterozygous/biallelic line was identified with two mutant alleles, one with a 1 bp insertion causing premature termination of the GMT1 protein, and the other with a 3 bp deletion resulting in the loss of one amino acid from the protein sequence. **d**, Schematic diagram of the *RCKT1* gene structure with gRNA targeting sites labeled. CRISPR/Cas9 editing was only detected at the gRNA2 (red) target site. **e**, Gene-editing events at the gRNA2 target site of the *rckt1* callus mutants. The homozygous knockout mutant *rckt1-21* has a 49 bp deletion within the *RCKT1* gene, creating a frameshifted and truncated RCKT1 protein. The *rckt1-1* heterozygous/biallelic line has two mutant alleles, one with a 3 bp deletion causing the loss of one amino acid from the RCKT1 protein, and the other with a 1 bp deletion resulting in frameshift and premature stop codon in the protein sequence. The *rckt1-9* heterozygous line has two mutant alleles, one with a 6 bp deletion causing the loss of two amino acids from the RCKT1 protein, and the other with a 1 bp insertion resulting in frameshift and premature stop codon in the protein sequence.

559
**Fig. 3| Cell wall polysaccharides and pectin analysis of *rckt1* callus mutants. a**,
Monosaccharide composition of non-cellulosic cell wall polysaccharides **b**, Cellulose-derived
glucose. Data in **a** and **b** are means of three biological replicates $\pm$ SD. No statistically significant
difference (ANOVA test) was observed between Col-0 wild-type, Non-edited (NE) Control, and
the homozygous and heterozygous/biallelic *rckt1* mutants. **c**, Separation of pectic domains using
size-exclusion chromatography (SEC). The representative SEC profile of endopolygalacturonase
(EPG) and pectin methyl esterase (PME) treated alcohol-insoluble residues (AIR) are shown for
Col-0 wild-type (black), NE Control (green), *rckt1-9* (light blue), and *rckt1-21* (dark blue). The
position of rhamnogalacturonan I (RG-I), RG-II dimer and monomer, and the oligogalacturonides
(OGAs) are shown.

**Fig 4| [1]H-NMR analysis of RG-II isolated from a**, Col-0 wildtype, **b**, NE control, **c**, *rckt1-9* and
**d**, *rckt1-21*. The isolated RG-II (dimers) was treated with dilute HCl to generate monomers for the
analysis. The diagnostic resonance of Kdo is shown in **a** and **b**, but absent from **c** and **d**. The Dha
signal is detected in all the samples.

576
577
578 **Fig 5| 2D TOCSY NMR analysis of RG-II isolated from a**, Col-0 wildtype, **b**, NE control, **c**,
579 *rckt1-9* and **d**, *rckt1-21*. The isolated RG-II (dimers) was treated with dilute HCl to generate
580 monomers for the analysis. The diagnostic resonance of Kdo is shown in **a** and **b**, but absent
581 from **c** and **d**. The Dha signal is detected in all the samples.

582
583
**Fig. 6| Structure of Arabidopsis RCKT1 predicted by AlphaFold modeling. a,** Cartoon
representation of general architecture of RCKT1 model. The predicted N-terminal transmembrane
domain correspondent from residue 1 to 36 is not shown. The nucleotide donor substrate, CMP-
β-Kdo, is positioned atop a central β-sheet comprising seven parallel β-strands surrounded by 14
α-helices and 2 small β-sheets. **b,** Surface representation of RCKT1 structure in complex with
CMP-β-Kdo. It reveals the potential CMP-β-Kdo binding site and the surrounding binding cleft,
which is a likely location for the binding of the acceptor substrate. **c**, Zoom in on the CMP-β-Kdo
binding site showing all the residues interacting via hydrogen bond and aromatic interactions
between RCKT1 and CMP-β-Kdo ligand.
593
594

## Data availability

The data supporting the findings of this study are available within the paper and its supplementary information files.

**A putative rhamnogalacturonan-II CMP-β-Kdo transferase identified using CRISPR/Cas9 gene edited callus to circumvent embryo lethality.**

## Reference

1.  Cosgrove, D. J. Building an extensible cell wall. *Plant Physiol.* **189**, 1246–1277 (2022).
2.  Ulvskov, P. & Scheller, H. V. Cell walls have a new family. *Nat. Plants* **4**, 635–636 (2018).
3.  Kobayashi, M., Matoh, T. & Azuma, Ji. Two Chains of Rhamnogalacturonan II Are Cross-Linked by Borate-Diol Ester Bonds in Higher Plant Cell Walls. *Plant Physiol.* **110**, 1017–1020 (1996).
4.  O'Neill, M. A., Ishii, T., Albersheim, P. & Darvill, A. G. Rhamnogalacturonan II: structure and function of a borate cross-linked cell wall pectic polysaccharide. *Annu. Rev. Plant Biol.* **55**, 109–139 (2004).
5.  Yang, Y. & Anderson, C. T. Biosynthesis, localisation, and function of pectins in plants. in *Pectin: technological and physiological properties* (ed. Kontogiorgos, V.) 1–15 (Springer International Publishing, 2020). doi:10.1007/978-3-030-53421-9_1.
6.  Bar-Peled, M., Urbanowicz, B. R. & O'Neill, M. A. The Synthesis and Origin of the Pectic Polysaccharide Rhamnogalacturonan II - Insights from Nucleotide Sugar Formation and Diversity. *Front. Plant Sci.* **3**, 92 (2012).
7.  González-Fontes, A. Why boron is an essential element for vascular plants. *New Phytol.* **226**, 1228–1230 (2020).
8.  Ishii, T. & Matsunaga, T. Isolation and characterization of a boron-rhamnogalacturonan-II complex from cell walls of sugar beet pulp. *Carbohydr. Res.* **284**, 1–9 (1996).
9.  Ishii, T., Matsunaga, T. & Hayashi, N. Formation of rhamnogalacturonan II-borate dimer in pectin determines cell wall thickness of pumpkin tissue. *Plant Physiol.* **126**, 1698–1705 (2001).
10. Matsunaga, T. *et al.* Occurrence of the primary cell wall polysaccharide rhamnogalacturonan II in pteridophytes, lycophytes, and bryophytes. Implications for the evolution of vascular plants. *Plant Physiol.* **134**, 339–351 (2004).
11. Funakawa, H. & Miwa, K. Synthesis of borate cross-linked rhamnogalacturonan II. *Front. Plant Sci.* **6**, 223 (2015).
12. Mohnen, D. Pectin structure and biosynthesis. *Curr. Opin. Plant Biol.* **11**, 266–277 (2008).
13. Scheller, H. V., Jensen, J. K., Sørensen, S. O., Harholt, J. & Geshi, N. Biosynthesis of pectin. *Physiol. Plant.* **129**, 283–295 (2006).
14. Ndeh, D. *et al.* Complex pectin metabolism by gut bacteria reveals novel catalytic functions. *Nature* **544**, 65–70 (2017).
15. O'Neill, M. A., Eberhard, S., Albersheim, P. & Darvill, A. G. Requirement of borate cross-linking of cell wall rhamnogalacturonan II for Arabidopsis growth. *Science* **294**, 846–849 (2001).
16. Pabst, M. *et al.* Rhamnogalacturonan II structure shows variation in the side chains monosaccharide composition and methylation status within and across different plant species. *Plant J.* **76**, 61–72 (2013).
17. Rautengarten, C. *et al.* The Arabidopsis Golgi-localized GDP-L-fucose transporter is required for plant development. *Nat. Commun.* **7**, 12119 (2016).
18. Qi, T. *et al.* GDP-D-mannose epimerase regulates male gametophyte development, plant growth and leaf senescence in Arabidopsis. *Sci. Rep.* **7**, 10309 (2017).

19.  Sechet, J. *et al.* Suppression of Arabidopsis GGLT1 affects growth by reducing the L-
galactose content and borate cross-linking of rhamnogalacturonan-II. *Plant J.* **96**, 1036–
1050 (2018).
20.  Reboul, R. *et al.* Down-regulation of UDP-glucuronic acid biosynthesis leads to swollen
plant cell walls and severe developmental defects associated with changes in pectic
polysaccharides. *J. Biol. Chem.* **286**, 39982–39992 (2011).
21.  Ahn, J.-W. *et al.* Depletion of UDP-D-apiose/UDP-D-xylose synthases results in
rhamnogalacturonan-II deficiency, cell wall thickening, and cell death in higher plants. *J.*
*Biol. Chem.* **281**, 13708–13716 (2006).
22.  Delmas, F. *et al.* The synthesis of the rhamnogalacturonan II component 3-deoxy-D-
manno-2-octulosonic acid (Kdo) is required for pollen tube growth and elongation. *J. Exp.*
*Bot.* **59**, 2639–2647 (2008).
23.  Egelund, J. *et al.* Arabidopsis thaliana RGXT1 and RGXT2 encode Golgi-localized (1,3)-
alpha-D-xylosyltransferases involved in the synthesis of pectic rhamnogalacturonan-II.
*Plant Cell* **18**, 2593–2607 (2006).
24.  Egelund, J. *et al.* Functional characterisation of a putative rhamnogalacturonan II specific
xylosyltransferase. *FEBS Lett.* **582**, 3217–3222 (2008).
25.  Petersen, B. L. *et al.* Assay and heterologous expression in Pichia pastoris of plant cell
wall type-II membrane anchored glycosyltransferases. *Glycoconj. J.* **26**, 1235–1246
(2009).
26.  Liu, X.-L. *et al.* Male gametophyte defective 4 encodes a rhamnogalacturonan II
xylosyltransferase and is important for growth of pollen tubes and roots in Arabidopsis.
*Plant J.* **65**, 647–660 (2011).
27.  Biswal, A. K. *et al.* Sugar release and growth of biofuel crops are improved by
downregulation of pectin biosynthesis. *Nat. Biotechnol.* **36**, 249–257 (2018).
28.  Mortimer, J. C. & Scheller, H. V. Synthesis and function of complex sphingolipid
glycosylation. *Trends Plant Sci.* **25**, 522–524 (2020).
29.  Mortimer, J. C. *et al.* Abnormal glycosphingolipid mannosylation triggers salicylic acid-
mediated responses in Arabidopsis. *Plant Cell* **25**, 1881–1894 (2013).
30.  Fang, L. *et al.* Loss of inositol phosphorylceramide sphingolipid mannosylation induces
plant immune responses and reduces cellulose content in arabidopsis. *Plant Cell* **28**,
2991–3004 (2016).
31.  Ishikawa, T. *et al.* GLUCOSAMINE INOSITOLPHOSPHORYLCERAMIDE
TRANSFERASE1 (GINT1) Is a GlcNAc-Containing Glycosylinositol Phosphorylceramide
Glycosyltransferase. *Plant Physiol.* **177**, 938–952 (2018).
32.  Singh, S. K., Eland, C., Harholt, J., Scheller, H. V. & Marchant, A. Cell adhesion in
Arabidopsis thaliana is mediated by ECTOPICALLY PARTING CELLS 1--a
glycosyltransferase (GT64) related to the animal exostosins. *Plant J.* **43**, 384–397 (2005).
33.  Voxeur, A. *et al.* Altered lignification in mur1-1 a mutant deficient in GDP-L-fucose
synthesis with reduced RG-II cross linking. *PLoS ONE* **12**, e0184820 (2017).
34.  Chatterjee, M. *et al.* The boron efflux transporter ROTTEN EAR is required for maize
inflorescence development and fertility. *Plant Cell* **26**, 2962–2977 (2014).
35.  Jarvis, M. C., Briggs, S. P. H. & Knox, J. P. Intercellular adhesion and cell separation in
plants. *Plant Cell Environ.* **26**, 977–989 (2003).
36.  *Plant cell separation and adhesion.* (Blackwell Publishing Ltd, 2007).
doi:10.1002/9780470988824.
37.  Daskalova, S. M., Pah, A. R., Baluch, D. P. & Lopez, L. C. The Arabidopsis thaliana
putative sialyltransferase resides in the Golgi apparatus but lacks the ability to transfer
sialic acid. *Plant Biol (Stuttg)* **11**, 284–299 (2009).
38.  Zeleny, R., Kolarich, D., Strasser, R. & Altmann, F. Sialic acid concentrations in plants
are in the range of inadvertent contamination. *Planta* **224**, 222–227 (2006).

39.   Angata, T. & Varki, A. Chemical Diversity in the Sialic Acids and Related α-Keto
Acids: An Evolutionary Perspective. *Chem. Rev.* **102**, 439–470 (2002).
40.   Paccalet, T. *et al.* Engineering of a sialic acid synthesis pathway in transgenic plants by
expression of bacterial Neu5Ac-synthesizing enzymes. *Plant Biotechnol. J.* **5**, 16–25
(2007).
41.   Vergunst, A. C., de Waal, E. C. & Hooykaas, P. J. Root transformation by Agrobacterium
tumefaciens. *Methods Mol. Biol.* **82**, 227–244 (1998).
42.   Lowder, L. G. *et al.* A crispr/cas9 toolbox for multiplexed plant genome editing and
transcriptional regulation. *Plant Physiol.* **169**, 971–985 (2015).
43.   Bloh, K. *et al.* Deconvolution of Complex DNA Repair (DECODR): Establishing a Novel
Deconvolution Algorithm for Comprehensive Analysis of CRISPR-Edited Sanger
Sequencing Data. *The CRISPR Journal* **4**, 120–131 (2021).
44.   Jing, B. *et al.* The Arabidopsis thaliana nucleotide sugar transporter GONST2 is a
functional homolog of GONST1. *Plant Direct* **5**, e00309 (2021).
45.   Barnes, W. J. *et al.* Protocols for isolating and characterizing polysaccharides from plant
cell walls: a case study using rhamnogalacturonan-II. *Biotechnol. Biofuels* **14**, 142 (2021).
46.   Reuhs, B. L. *et al.* l-Galactose replaces l-fucose in the pectic polysaccharide
rhamnogalacturonan II synthesized by the l-fucose-deficient mur1 Arabidopsis mutant.
*Planta* **219**, 147–157 (2004).
47.   Wen, L., Zheng, Y., Li, T. & Wang, P. G. Enzymatic synthesis of 3-deoxy-d-manno-
octulosonic acid (KDO) and its application for LPS assembly. *Bioorg. Med. Chem. Lett.*
**26**, 2825–2828 (2016).
48.   Sugai, T., Lin, C. H., Shen, G. J. & Wong, C. H. CMP-KDO synthetase: overproduction
and application to the synthesis of CMP-KDO and analogs. *Bioorg. Med. Chem.* **3**, 313–
320 (1995).
49.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,
583–589 (2021).
50.   Rao, F. V. *et al.* Structural insight into mammalian sialyltransferases. *Nat. Struct. Mol.*
*Biol.* **16**, 1186–1188 (2009).
51.   Meng, L. *et al.* Enzymatic basis for N-glycan sialylation: structure of rat α2,6-
sialyltransferase (ST6GAL1) reveals conserved and unique features for glycan sialylation.
*J. Biol. Chem.* **288**, 34680–34698 (2013).
52.   Chiu, C. P. C. *et al.* Structural analysis of the sialyltransferase CstII from Campylobacter
jejuni in complex with a substrate analog. *Nat. Struct. Mol. Biol.* **11**, 163–170 (2004).
53.   Braybrook, S. A. & Peaucelle, A. Mechano-chemical aspects of organ formation in
Arabidopsis thaliana: the relationship between auxin and pectin. *PLoS ONE* **8**, e57813
(2013).
54.   Levesque-Tremblay, G., Pelloux, J., Braybrook, S. A. & Müller, K. Tuning of pectin
methylesterification: consequences for cell wall biomechanics and development. *Planta*
**242**, 791–811 (2015).
55.   *Pectin: technological and physiological properties.* (Springer International Publishing,
2020). doi:10.1007/978-3-030-53421-9.
56.   Caffall, K. H. & Mohnen, D. The structure, function, and biosynthesis of plant cell wall
pectic polysaccharides. *Carbohydr. Res.* **344**, 1879–1900 (2009).
57.   Fleischer, A., O'Neill, M. A. & Ehwald, R. The Pore Size of Non-Graminaceous Plant Cell
Walls Is Rapidly Decreased by Borate Ester Cross-Linking of the Pectic Polysaccharide
Rhamnogalacturonan II. *Plant Physiol.* **121**, 829–838 (1999).
58.   Zhou, T. *et al.* Low-Boron Tolerance Strategies Involving Pectin-Mediated Cell Wall
Mechanical Properties in Brassica napus. *Plant Cell Physiol.* **58**, 1991–2005 (2017).

59.    Amos, R. A. & Mohnen, D. Critical review of plant cell wall matrix polysaccharide
glycosyltransferase activities verified by heterologous protein expression. *Front. Plant*
*Sci.* **10**, 915 (2019).
60.    Welner, D. H. *et al.* Plant cell wall glycosyltransferases: High-throughput recombinant
expression screening and general requirements for these challenging enzymes. *PLoS*
*ONE* **12**, e0177591 (2017).
61.    Moremen, K. W. *et al.* Expression system for structural and functional studies of human
glycosylation enzymes. *Nat. Chem. Biol.* **14**, 156–162 (2018).
62.    Urbanowicz, B. R., Peña, M. J., Moniz, H. A., Moremen, K. W. & York, W. S. Two
Arabidopsis proteins synthesize acetylated xylan in vitro. *Plant J.* **80**, 197–206 (2014).
63.    Urbanowicz, B. R. *et al.* Structural, mutagenic and in silico studies of xyloglucan
fucosylation in Arabidopsis thaliana suggest a water-mediated mechanism. *Plant J.* **91**,
931–949 (2017).
64.    Voiniciuc, C. *et al.* Identification of key enzymes for pectin synthesis in seed mucilage.
*Plant Physiol.* **178**, 1045–1064 (2018).
65.    Amos, R. A. *et al.* A two-phase model for the non-processive biosynthesis of
homogalacturonan polysaccharides by the GAUT1:GAUT7 complex. *J. Biol. Chem.* **293**,
19047–19063 (2018).
66.    Soto, M. J. *et al.* AtFUT4 and AtFUT6 Are Arabinofuranose-Specific
Fucosyltransferases. *Front. Plant Sci.* **12**, 589518 (2021).
67.    Engle, K. A. *et al.* Multiple Arabidopsis galacturonosyltransferases synthesize polymeric
homogalacturonan by oligosaccharide acceptor-dependent or de novo synthesis. *Plant J.*
**109**, 1441–1456 (2022).
68.    Amos, R. A. *et al.* Polymerization of the backbone of the pectic polysaccharide
rhamnogalacturonan I. *Nat. Plants* **8**, 1289–1303 (2022).
69.    Gronow, S., Brabetz, W. & Brade, H. Comparative functional characterization in vitro of
heptosyltransferase I (WaaC) and II (WaaF) from Escherichia coli. *Eur. J. Biochem.* **267**,
6602–6611 (2000).
70.    Ovchinnikova, O. G. *et al.* Bacterial β-Kdo glycosyltransferases represent a new
glycosyltransferase family (GT99). *Proc Natl Acad Sci USA* **113**, E3120-9 (2016).
71.    Lanz, N. D. *et al.* Characterization of the β-KDO Transferase KpsS, the Initiating Enzyme
in the Biosynthesis of the Lipid Acceptor for Escherichia coli Polysialic Acid. *Biochemistry*
(2021) doi:10.1021/acs.biochem.1c00088.
72.    Séveno, M. *et al.* Characterization of a putative 3-deoxy-D-manno-2-octulosonic acid
(Kdo) transferase gene from Arabidopsis thaliana. *Glycobiology* **20**, 617–628 (2010).
73.    Caffall, K. H., Pattathil, S., Phillips, S. E., Hahn, M. G. & Mohnen, D. Arabidopsis
thaliana T-DNA mutants implicate GAUT genes in the biosynthesis of pectin and xylan in
cell walls and seed testa. *Mol. Plant* **2**, 1000–1014 (2009).
74.    Sterling, J. D. *et al.* Functional identification of an Arabidopsis pectin biosynthetic
homogalacturonan galacturonosyltransferase. *Proc Natl Acad Sci USA* **103**, 5236–5241
(2006).
75.    Johnson, K. L., Kibble, N. A. J., Bacic, A. & Schultz, C. J. A fasciclin-like
arabinogalactan-protein (FLA) mutant of Arabidopsis thaliana, fla1, shows defects in
shoot regeneration. *PLoS ONE* **6**, e25154 (2011).
76.    Meng, X. *et al.* Construction of a Genome-Wide Mutant Library in Rice Using
CRISPR/Cas9. *Mol. Plant* **10**, 1238–1241 (2017).
77.    Liang, Y. *et al.* A screening method to identify efficient sgRNAs in Arabidopsis, used in
conjunction with cell-specific lignin reduction. *Biotechnol. Biofuels* **12**, 130 (2019).
78.    Grützner, R. *et al.* High-efficiency genome editing in plants mediated by a Cas9 gene
containing multiple introns. *Plant Commun.* **2**, 100135 (2021).

79.    Czako, M. & Marton, L. In.* Arabidopsis: The Compleat Guide*(D. Flanders, and C.
799            Dean eds). AFRC PMB Programme. (1992).
800    80.    Prime, T. A., Sherrier, D. J., Mahon, P., Packman, L. C. & Dupree, P. A proteomic
801            analysis of organelles from Arabidopsis thaliana. *ELECTROPHORESIS* (2000).
802    81.    Krishnakumar, V. *et al.* Thalemine: A warehouse for arabidopsis data integration and
803            discovery. *Plant Cell Physiol.* **58**, e4 (2017).
804    82.    Markham, J. E. & Jaworski, J. G. Rapid measurement of sphingolipids from Arabidopsis
805            thaliana by reversed-phase high-performance liquid chromatography coupled to
806            electrospray ionization tandem mass spectrometry. *Rapid Commun. Mass Spectrom.* **21**,
807            1304–1314 (2007).
808    83.    Skipski, V. P. [47] Thin-layer chromatography of neutral glycosphingolipids. in *Lipids Part
809            B* vol. 35 396–425 (Elsevier, 1975).
810    84.    Varki, A. *et al.* Symbol nomenclature for graphical representations of glycans.
811            *Glycobiology* **25**, 1323–1324 (2015).
812    85.    Neelamegham, S. *et al.* Updates to the Symbol Nomenclature for Glycans guidelines.
813            *Glycobiology* **29**, 620–624 (2019).
814
815

## Acknowledgements

This work was part of the DOE Joint BioEnergy Institute (http://www.jbei.org) supported by the U. S. Department of Energy, Office of Science, Office of Biological and Environmental Research, through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U. S. Department of Energy. This work was also supported in part by an establishment grant to JCM from the University of Adelaide. Work conducted at the Complex Carbohydrate Research Center, of the University of Georgia, was supported by the Division of Chemical Sciences, Geosciences, and Biosciences, Office of Basic Energy Sciences of the United States Department of Energy through Grant DESC0008472 for funding structural studies of RG-II and DE-SC0015662 to the Complex Carbohydrate Research Center. We thank Maartje van Kregten and Annette Vergunst for the insightful discussion on the Agrobacterium-mediated root transformation of Arabidopsis. We thank Yu Gao and Yi-Chun Chen for their support with HPAEC-PAD experiments. We thank Jutta Dalton and Mi Yeon Lee for their support with growth chamber maintenance. We thank Ramana Pidatala for his advice on plant tissue culture. We thank Ariel Orellana for the insightful discussion on the putative CMP-Kdo transporter, homologous to CMP-sialic acid transporter identified in mammals. We thank Parastoo Azadi and the analytical service laboratory at the Complex Carbohydrate Research Center for technical support.

## Contributions

Y.Z. and J.C.M. conceived and designed the study. Y.Z. conducted the gene editing, transformation and growth experiments, assisted by N.D., Y.L., and F.D. YZ conducted GIPC and cell wall analysis. D.S., K.T., M.O., I.B., and B.U. structurally characterized RG-II. J.H.P. and P.A. performed AlphaFold modeling of RCKT1. Data analysis and interpretation were conducted by Y.Z., H.V.S. D.S., B.U., M.O., J.H.P., and J.C.M. The paper was written by Y.Z. and J.C.M. with revisions from all authors.

## Ethics declarations

The authors declare no competing interests.