Integrating annual radial growth analyses and carbon isotope discrimination to

forecast early warning of beech forest dieback across the Italian Peninsula

Paulina F. Puchi[1,2*], Daniela Dalmonech[1,3], Elia Vangi[1], Giovanna Battipaglia[4], Roberto Tognetti[5]

and Alessio Collalti[1,3]

[1]Forest Modelling Lab., Institute for Agriculture and Forestry Systems in the Mediterranean, National

Research Council of Italy (CNR-ISAFOM), Via Madonna Alta 128, 06128 Perugia, Italy; [2]Institute

of Bioeconomy, Italian National Research Council (CNR-IBE), Via Madonna del Piano 10, 50019

Sesto Fiorentino, Italy; [3]National Biodiversity Future Center (NBFC), 90133 Palermo, Italy;

[4]Department of Environmental, Biological and Pharmaceutical Sciences and Technologies,

University of Campania 'L. Vanvitelli', Caserta, Italy; [5]Department of Agricultural, Environmental,

and Food Sciences, University of Molise, Via De Sanctis, 86100 Campobasso, Italy

Author for correspondence

Paulina F. Puchi

Email: paulina.puchi@isafom.cnr.it

Tel: +39 0498272740

Paper type: Full paper

Summary

- Tree mortality and forest dieback episodes are increasing due to drought and heat stress. Nevertheless, a comprehensive understanding of mechanisms enabling trees to withstand and survive droughts remains lacking.

- We used tree-ring width, basal area increment (BAI), and $\delta^{13}C$-derived intrinsic water-use-efficiency ($_iWUE$) to study beech resilience across four healthy stands in Italy with varying climates and water availability. Fist-order autocorrelation (AR1) analysis was performed to detect early warning signals for potential tree dieback risks during extreme drought events.

- We found a negative link between BAI and vapour pressure deficit (VPD), especially in southern latitudes. After the 2003 drought, BAI decreased at the northern site, with an increase in $\delta^{13}C$ and $_iWUE$, indicating conservative water-use. Conversely, the southern sites showed increased BAI and $_iWUE$, likely influenced by rising CO_2 and improved water availability. In contrast, the central site sustained higher transpiration rates due to higher soil water holding capacity (SWHC). Despite varied responses, most sites exhibited reduced resilience to future extreme events, indicated by increased AR1.

- Temperature significantly affected beech $_iWUE$ and BAI in northern Italy, while VPD strongly influenced the southern latitudes. The observed increase in BAI and $_iWUE$ in southern regions might be attributed to an acclimation response.

Introduction

Forest ecosystems are facing significant challenges due to anthropogenic climate change (Allen *et al.*, 2010; McDowell *et al.*, 2020). The combination of reduced water availability and rising temperatures directly impacts the process of photosynthetic carbon assimilation, thereby reducing forest carbon sequestration (Keenan, 2015; Zuidema *et al.*, 2018). This could potentially lead to negative feedback on carbon balance (Pan *et al.*, 2011). Furthermore, hotter droughts have caused substantial alterations in forest structure and function, affecting tree growth performance and triggering episodes of dieback and tree mortality (Allen *et al.*, 2015; Anderegg *et al.*, 2016: Puchi *et al.*, 2021). In addition, climatic models predict that the frequency, duration, and intensity of extreme droughts will increase in the future (IPCC, 2021), so it is crucial to a better understanding of how forests are going to cope with these extreme climatic conditions (Brodribb *et al.*, 2020).

Despite the importance of identifying suitable tree species and future management practices in response to climate change, our understanding of species-specific physiological responses and site- and species-specific vulnerabilities to drought-induced tree mortality during extreme droughts

remains incomplete (Allen *et al*., 2015; Trugman *et al*., 2021; De Marco *et al*., 2022). This gap is

especially critical for European beech (*Fagus sylvatica* L.), one of the most distributed, ecologically

and economically significant tree species in Europe (Fang & Lechowicz, 2006). This species

comprises 17% of all broadleaf tree stands in Italy (Gasparini *et al*., 2022) and is one the most affected

by extreme events occurring during the initial vegetative phase across the Italian peninsula (D'Andrea,

et al., 2020; Martinez del Castillo *et al*., 2022; Tonelli *et al*., 2023).

Given the anticipated that climate change will exert a significant influence on both regional and local

drought patterns in the Mediterranean region (Adams *et al*., 2017; Sangüesa-Barreda *et al*., 2019). In

particular, mountain-Mediterranean beech forests would face increased vulnerability due to their

location in the southernmost distribution of this species' range (Noce *et al*., 2016, 2017; Leuschner,

2020; D'Andrea *et al*., 2021). Consequently, predicting resilience and adaptation across its

distribution has become a prioritized goal.

Recent studies have shown that prolonged heat and drought events can have detrimental effects on

both hydraulic function and carbon use in trees (McDowell *et al*., 2008; Anderegg *et al*., 2013).

Understanding these physiological mechanisms is crucial for comprehending how trees respond to

drought, as they directly influence water use regulation. For instance, isohydric species adopt a

conservative behaviour by closing stomata to minimize water loss, thereby reducing photosynthetic

activity, and increasing the risk of carbon starvation (Timofeeva *et al*., 2017). On the other hand,

anisohydric species adopt an opportunistic behaviour, exhibiting higher transpiration rates even when

soil moisture is low, leading to an elevated risk of hydraulic failure (McDowell *et al*., 2008; Petrucco

et al., 2017).

Currently, there is contrasting information regarding how European beech forests respond to heat and

drought events (Leuschner, 2020; D'Andrea *et al*., 2021). Most studies on young beech stands have

suggested a conservative response during droughts (Leuschner, 2020; Walthert *et al*., 2021; Martinez

del Castillo *et al*., 2022). However, in a few studies, adult trees have conversely displayed

opportunistic behaviour (Leuschner, 2020). Therefore, it is crucial to exploit better the plasticity of

this species in the water use strategies to determine the trajectories of species distribution and its

resilience to a warming and drier climate (Gessler *et al*., 2020; Walthert *et al*., 2021).

Long-term changes in intrinsic water use efficiency ($_i$WUE), i.e. the cost of fixing carbon per unit of

water loss (Seibt *et al*., 2008; Gagen *et al*., 2022), can be assessed by measuring carbon isotope

composition in tree rings ($\delta^{13}C$). Tree-ring $\delta^{13}C$ is equivalent to the ratio between photosynthesis (*A*)

and stomatal conductance (g_s) and this can vary, since both affect the ratio between CO_2 partial

pressure in leaf intercellular space and in the atmosphere (Farquhar *et al*., 1982; Battipaglia &

Cherubini, 2022). Variations in ¡WUE, within and across tree species, have revealed a continuous

ecophysiological gradient of plant water-use strategies ranging from "profligate/opportunistic" (low

¡WUE) to those considered "conservative" (high ¡WUE) (Moreno-Gutiérrez *et al.*, 2012). For

instance, studies in tree rings have shown that the increase of ¡WUE did not enhance tree growth

(Peñuelas *et al.*, 2011), however, others showed the opposite effect or both (Peñuelas *et al.*, 2008;

Tognetti *et al.*, 2014; Walker *et al.*, 2021). These indicators of hydraulic strategies and carbon

discrimination provide valuable insights into the long-term impacts of climate change on forest health

and the risk of tree mortality (Gessler *et al.*, 2018; Cherubini *et al.*, 2021; Puchi *et al.*, 2021).

On the other hand, recent studies have provided evidence that one of the primary mortality risk

indicators in forests is growth reduction also occurring many decades before visible symptoms of

decline, such as leaf discolouration, increased defoliation, and branch dieback (Camarero *et al.*, 2015;

Cailleret *et al.*, 2016; 2019; DeSoto *et al.*, 2020). Similarly, another proxy indicator of loss of

resilience and thus increasing tree mortality risk is the autocorrelation, better called 'early warning

signal' (EWS), which has been proposed to detect a critical transition in long-term time series after a

perturbation, causing a critical slowing down of the capacity of recovery (Dakos *et al.*, 2012a; Gessler

et al., 2020; Forzieri *et al.*, 2021). EWS can be highlighted as increasing autocorrelation and variance

in tree growth, indicating loss of resilience and stability (Dakos *et al.*, 2012a,b). These changes have

been observed in conifers; however, angiosperms did not show changes in these indicators, and this

could be due to their capacity to recover after a stress-induced growth decline (Camarero *et al.*, 2015;

Cailleret *et al.*, 2019). These findings highlight the importance of early monitoring in understanding

forest resilience and adaptation to climate change.

This study aimed to assess the forest vitality of beech in response to drought stress by examining

historical and recent growth patterns across the Italian peninsula, with a particular emphasis on water

use strategies (conservative vs. opportunistic) at long-time scales. Secondly, we tested early warning

signals of potential tree dieback by analyzing autocorrelation and variability patterns, as indicators of

stand resilience and stability to future extreme events.

We hypothesized that beech populations in the southernmost distribution exhibit conservative

behaviour as an acclimation strategy. This behaviour is characterized by ¡WUE being more responsive

to VPD than those in the northern regions, reflecting a reduction in stomatal conductance to maintain

a minimum midday water potential, and also a decline in intercellular CO_2 concentration, but a more

slowly decrease in photosynthetic rate. Although a drought-driven decline in photosynthetic rate may

also occur, non-stomatal limitation was expected in populations with more opportunistic behaviour.

Additionally, we expected to find varying degrees of growth reduction as an early warning signal of

tree mortality risk across different sites, with the strongest signals in response to severe drought
events.

Materials and Methods

Study Sites and Climate

Analyses were conducted at four sites along a ~900 km latitudinal transect in pure European beech
forests across the Italian Peninsula (Fig. 1, Table 1). The sites were Trentino-Alto Adige (hereafter
abbreviated as 'TRE'), Lazio (hereafter abbreviated as 'LAZ'), Campania (hereafter abbreviated as
'CAM') and Calabria (hereafter abbreviated as 'CAL'). All the stands analyzed had not been managed
since the last 20-30 years.

The selection of the sites allowed the comparison of moisture availability across the Italian Peninsula
(Fig. 1), using the Climate Moisture Index (CMI) calculation method explained in section 2.6. These
sites differ along the latitudinal transect regarding both climatic conditions and soil types. From north
to south, the mean annual temperature ranges from 9 to 14.1 °C, with the mean annual precipitation
varying from 1003 to 825 mm, based on the E-OBS dataset (as explained in section E-OBS daily
climate data, CMI and SPEI calculation; Fig. S1)

The soil types from north to south are Andisols, Luvisols, and Inceptisols. Additionally, by examining
soil texture data, we inferred variations in soil water holding capacity (SWHC) among these sites
(Table S1, Hengl & Nauman, 2018). Specifically, we inferred that the SWHC in TRE is relatively
low, whereas LAZ exhibited a high SWHC. CAM also showed a high SWHC, while CAL presented
a moderate SWHC (Kutílek & Nielsen, 1994).

Fig. 1 a) Map displaying the Climate Moisture Index (CMI = Precipitation / Potential
EvapoTranspiration) across the Italian Peninsula, indicating humid and dry climate zones through
positive (blue) and negative (red) CMI values, respectively. The index was calculated for the growing
season (May-October) from 1965 to 2014. Green dots indicate the location of the four study sites
where dendrochronological samples were extracted.

Table 1 Geographical and mean annual climate characteristics for the four sites.

Site	Latitude (N)	Longitude (W)	Elevation (m a.s.l.)	Mean minimum annual temperature (°C)	Mean annual temperature (°C)	Mean maximum annual temperature (°C)	Annual precipitation (mm)	Reference
TRE	46°12'	11°16'	1276	3.8	9.0	13.9	929	TRW data Versace *et al.* (2020)
LAZ	42°24'	12°12'	1000	8.9	14.3	19.5	829	Battipaglia *et al.*, unpublished
CAM	41°24'	14°20'	1140	6.3	9.8	13.1	825	Battipaglia *et al.*, unpublished
CAL	39°19'	16°23'	1601	5.8	11	16	1003	Battipaglia *et al.*, unpublished

Field sampling and processing dendrochronological data

During the period 2014-2018, a total of 174 beech trees were sampled at 1.3 m from the ground using
a 5 mm increment borer (Table S2). In the laboratory, wood cores were air/dried and polished with
sandpaper of successively increasing grains to visualize the ring boundaries. Ring widths were
measured to a precision of 0.01 mm using the TSAP measuring device (Rinntech). Tree-ring (TRW)
series were then visually cross-dated using standard dendrochronological methods (Stokes & Smiley,
1968) and checked for dating accuracy and measurement errors with the COFECHA program
(Holmes, 1983).

Later, tree growth measurements were converted to basal area increment based (BAI) based on the

distance between the outermost measured ring (pith) and the last ring of the tree (i.e., the ring next to

the bark), using the following formula:

$BAI = \pi (R^2_t - R^2_{t-1})$ Eqn 1

where R_t the tree's radius at the end of the annual increment, and R_{t-1} is the tree's radius at the

beginning of the annual increment. This method assumes a circular cross-section, and the mean BAI

of defined periods can be compared over time, as it is not affected by biological trends (Biondi &

Qeadan, 2008a,b) and it is more tightly related to stem biomass compared to TRW. We worked with

mean non-standardized BAI values to preserve the long-term cumulative effects of climate on tree

growth (Tognetti *et al*., 2014). All analyses were restricted to the period covered by the youngest

trees (at LAZ), i.e. from 1965 until 2014 (Table S2). All computations were performed using the R-

package 'dplR' (Bunn, 2008; Bunn *et al*., 2021; R Core Team, 2022).

Water-use efficiency from carbon isotope discrimination

To compare long-term changes in $_i$WUE among beech trees across the Italian Peninsula, we measured

$^{13}C/^{12}C$ isotope ratios in the TRW. Ten samples per each stand presenting the best cross-dating (GLK

> 0.70) with the corresponding average chronology, were selected for stable isotope analyses

(Battipaglia *et al*., 2017) and they were annually dissected using a razor blade under a binocular

microscope for the period 1965-2014.

Wood samples were milled to a fine powder (ZM 1000; Retsch), weighed 0.05-0.06 mg of wood for

carbon isotope analyses and encapsulated in tin capsules.

The isotope composition was measured at the IRMS laboratory of the University of Campania "Luigi

Vanvitelli" by using mass spectrometry with continuous flow isotope ratio (Delta V plus Thermo

electron Corporation). The standard deviation for repeated analysis of an internal standard

(commercial cellulose) was better than 0.1‰ for carbon. The $\delta^{13}C$ series were corrected for the fossil

fuel combustion effect for anthropogenic changes in the atmospheric $\delta^{13}C$ composition ($\delta^{13}C_{atm}$)

(Francey *et al*., 1999; McCarroll & Loader, 2004).

Isotopic discrimination between the carbon of atmospheric CO_2 and wood carbon to determine $_i$WUE

can be calculated starting from the $\delta^{13}C$ of the plant material ($\delta^{13}C_{tree}$), which is related to atmospheric

$\delta^{13}C$ ($\delta^{13}C_{atm}$) and the ratio c_i/c_a, according to Farquhar & Richards (1984) and Farquhar *et al*. (1982):

$\delta^{13}C_{tree} = \delta^{13}C_{atm} - a - [(b - a) c_i] / c_a$ Eqn 2

where a is the fractionation factor due to $^{13}CO_2$ diffusion through stomata (4.4‰), and b is the fractionation factor due to Rubisco enzyme during the process of carboxylation (27‰) (McCarroll & Loader, 2004). Therefore, we can calculate c_i by using the formula:

$$c_i = c_a(\delta^{13}C_{atm} - \delta^{13}C_{tree} - a) / (b - a) \qquad \text{Eqn 3}$$

Finally, the $iWUE$ can be calculated as follows:

$$iWUE = (c_a b - \delta^{13}C_{atm} + \delta^{13}C_{tree}) / 1.6 \, (b - a) \qquad \text{Eqn 4}$$

However, the $iWUE$ should not be considered equivalent to instantaneous WUE at leaf level, which is the ratio of assimilation to stomatal conductance and considers the atmospheric water demand (Pacheco *et al*., 2020; Seibt *et al*., 2008). Thus, the equation used is the "simple" form of isotopic discrimination that does not include effects due to mesophyll conductance and photorespiration, which were unavailable for the study species.

We used $\delta^{13}C_{atm}$ values from Belmecheri & Lavergne (2020). We obtained the atmospheric concentration of CO_2 from the Mauna Loa station data (http://www.esrl.noaa.gov/).

E-OBS daily climate data, CMI and SPEI calculation

Daily climate data for precipitation (P), minimum (T_{min}), mean (T_{mean}), and maximum (T_{max}) temperature, as well as relative humidity (RH), were extracted from the E-OBS dataset on a regular 0.1-degree grid (Table 1). The data were obtained as netCDF files from (http://surfobs.climate.copernicus.eu/dataaccess/access_eobs.php). Using the RH and temperature data, the vapour pressure deficit (VPD) in hPa was calculated based on the Tetens formula (Tetens, 1930).

The Climate Moisture Index (CMI, Willmott & Feddema, 1992) represents the relationship between plant water demand and available precipitation. The CMI indicator ranges from –1 to +1, with wet climates showing positive CMI and dry climates negative CMI. CMI was calculated as follows:

$$CMI = P/PET \qquad \text{Eqn 5}$$

Where P is the precipitation, and PET is the potential evapotranspiration. Specifically, CMI = (P/PET) –1 when P<PET and CMI = 1–(PET/P) when P>PET, to recast the limit to –1<CMI<1.

PET can be calculated through the Hargreaves equation (Hargreaves, 1985), modified by Allen (1993):

$PET = 0.0029 \, R_{solar\_rad} \, (T_{mean}+20) \, TR^{0.4}$ Eqn 6

Where R_{solar_rad} is the extraterrestrial solar radiation, T_{mean} in Celsius degree and TR is the temperature

range (T_{max}–T_{min}).

CMI was calculated for the growing season (May-October) using the E-OBS v. 27.0

(https://surfobs.climate.copernicus.eu/dataaccess/access_eobs.php#datafiles) daily products (T_{min},

T_{max}, precipitation, and global solar radiation) at 0.1 deg spatial resolution and averaged over the

period 1965-2014. E-OBS global solar radiation at the surface was converted to extra-terrestrial solar

radiation with the 'envirem' R-package (Title & Bemmels, 2018).

Additionally, to quantify drought severity, we calculated the Standardized Precipitation-

Evapotranspiration Index (SPEI), based on a statistical transformation of the climatic water balance,

i.e. precipitation *minus* potential evapotranspiration (P-PET) (Vicente-Serrano *et al.*, 2010). The

multiscalar drought index was calculated at different time scales (from 1 to 24 months, Fig. S2) for

the period 1965-2014 (constrained to the youngest site LAZ) in R using the 'SPEI' package (Beguería

et al., 2014; Beguería & Vicente-Serrano, 2017).

Later, to assess the relationships between climate and BAI and stable isotope for the period 1965–

2014, we calculated Pearson's correlations between monthly P, T_{mean}, VPD, and SPEI (1-3-6-9-12-

18 and 24 months) series from previous $_{(t-1)}$ and current year $_{(t)}$, using monthly response function in

the 'DendroTools' R-package (Jevšenak & Levanič, 2018).

Growth trends and climate response

We used Generalized Additive Mixed Models (GAMM) to study the long-term annual BAI and their

responses to changing climatic conditions, particularly concerning water balance within the growing

season (May-October) using SPEI indexes at the four study sites. We tested SPEI drought index at

1,3,6,9,12,18, and 24 months as the potential influence of drought on BAI. GAMM is a flexible

semiparametric method that allows the simultaneous modelling of linear and nonlinear relationships

between the response variable as a function of some explanatory variables (Wood, 2006) that allows

the treatment of autocorrelation and repeated measures (Wood, 2006). The variables included in the

model were the following:

$BAI_i = s[year_i * (Site)] + s(age_i) + s(SPEI_i) + Z_i B_i + \varepsilon$ Eqn 7

Where the BAI_i of a tree$_i$ were modelled as a function of calendar year, individual tree age and SPEI

per site. In addition, given that BAI represents multiple measurements performed in each tree, tree

identity (Z_iB_i) is regarded as a random effect (Z_i and B_i indicate matrix variables and related

coefficients). Thin plate regression splines (s) were used to represent all the smooth terms, with a

degree of smoothing determined by internal cross-validation (Wood, 2006). We ranked all the

potential models that could be generated using different explanatory variables and different levels of

smoothing according to the Akaike Information Criterion (AIC). Finally, we chose the model with

the lowest AIC (Burnham & Anderson, 2002). The time scale that best explained the variability in

BAI was the 18-month SPEI (for the growing season May-October). The GAMMs were performed

and fitted using the function 'gamm' in the 'mgcv' R-package (Wood, 2006).

Early warning signals of forest dieback

For assessing stand resilience for each site and each time series of BAI, we computed the first-order

autocorrelation (AR1) and the standard deviation (SD) over the period 1965 to 2014 using a 15-year

moving window (30% of the entire time series). These metrics are widely recognized indicators of

changes in time series and proximity to critical transitions to new states (Dakos *et al.*, 2012a;

Camarero *et al.*, 2015; Forzieri *et al.*, 2022). The trend of AR1 and SD metrics over the considered

temporal window was computed by means of the non-parametric Mann-Kendall Tau statistics. For

each site, the significance of a positive (or negative) AR1 and SD trend was tested with a one-sided

t-test. We employed the R-package 'early warnings' (Dakos *et al.*, 2012b) to compute the selected

metrics. All statistical analyses were conducted using the R-statistical software (R Development Core

Team, 2022).

Results

Climate trends and drought variability

Annual precipitation (P) has increased significantly at TRE and CAM sites ($P < 0.01$, Fig. S2a), while

LAZ showed a reduction in P trend during the period from 1965 to 2014 ($P < 0.05$, Fig. S2a), and

CAL did not present any trend in P pattern. Notably, T_{min} increased significantly in TRE, LAZ, and

CAM ($P < 0.001$, Fig. S2b), whereas in the southern site (CAL), T_{min} presented a pronounced decrease

($P < 0.01$, Fig. S2b). Simultaneously, both T_{mean} and T_{max} exhibited a substantial and significant

increase across all sites ($P < 0.01$, Fig. S2c,d). Interestingly, only at the northernmost site (TRE), VPD

increased drastically and significantly during the 2000s ($P < 0.001$, Fig. S2e), while at the

southernmost site (CAL) VPD showed the opposite pattern (Fig. S2e, $P < 0.001$).

As for the P trend, the SPEI index showed an increase in water availability in recent years across sites, although not significant, except for LAZ, which showed a negative trend ($P< 0.05$, Figs. 2, S3). Notably, the SPEI-derived drought index showed the widespread impact of the 2003 drought across all sites, more evident at CAM (Figs. 2, S3).

Fig. 2 Standardized 18-Month SPEI at the four study sites (TRE, LAZ, CAM, and CAL) for the 1965–2014 period. Negative (red) and positive (blue) values indicate drier and wetter conditions, respectively.

Long-*term growth patterns of European beech across the Italian Peninsula*

Mean TRW, the highest and the lowest growth rates were observed in LAZ and CAM sites,
respectively with statistically significant differences. Conversely, TRE and CAL showed similar
growth rates values ($P< 0.05$, Table S3). The age distribution of tree populations exhibited notable
differences across the four sites, with LAZ featuring the youngest trees and CAM the oldest trees ($P<$
0.05, Table S3).

BAI exhibited a significant decline, particularly pronounced in the relatively northern sites (TRE and
LAZ), following the drought of 2003 (Fig. 3b). In contrast, CAM presented a steady increase in BAI,
while in CAL, BAI decreased after 2010 (Fig. 3b).

Fig. 3 Long-term growth of basal area increment (BAI) of the four sites along a latitudinal gradient
(north to south) for the period 1965–2014. Colour lines for each site indicate the linear model, shaded
areas represent 95 % confidence intervals.

Growth response to climate variables

Basal area increment exhibited significant relationships with climatic variables in all study sites (Fig.
4). Overall, BAI was positively correlated with monthly P and T_{mean} and, notably, strongly negative

correlations with VPD were evident from May to September. This negative VPD correlation intensified toward the southern sites (Fig. 4).

At TRE, BAI was positively correlated with monthly P from May to July, with a stronger effect when considering P values in the previous year. Additionally, BAI correlated positively with May T_{mean} of the current year, instead of showing weak negative correlations with VPD and SPEI (Fig. 4).

LAZ showed a strong positive correlation between BAI and T_{mean} from March to November (previous and current year). Conversely, strong negative correlations with VPD from May to September and weak negative correlations with P and SPEI were found in August (Fig. 4).

AT CAM, a positive response of BAI to P of July of the previous years, and a strong positive correlation with T_{mean} during May to July, were observed, while a strong negative response to VPD from March to September (more evident in the current year) was found.

At CAL, BAI showed strong positive correlations with October P and with T_{mean} from March to August. In contrast, BAI displayed a strong negative correlation with VPD from March to September (current and previous year). Similarly, negative scattered correlations with SPEI were observed during summer at the southern sites (CAM and CAL, Fig. 4).

337 **Fig. 4** Pearson's running correlations between BAI with monthly precipitation, mean temperature,

338 VPD, and SPEI1 for the current and the previous year (*) over the period 1965-2014 at each site. The

339 *y-axis* represents the time window in months. Colours (see the key) represent correlation coefficients

340 that are significant at the level of $r = 0.279$ ($P< 0.05$).

Growth trends of beech

The GAMMs revealed different BAI trends of beech among the four sites (Fig. 5, Table S4). GAMM

showed a monotonic increasing growth trend among the sites; however, they started to diverge in the

mid-1990s. Notably, the northernmost site (TRE) started to decline earlier than the other sites (Fig.

5). Secondly, LAZ exhibited the highest increase, followed by a drastic decline during the 2000s – a

similar pattern was also observed in CAL. In contrast, at CAM trees demonstrated a steady increase

in BAI over the observed period (Fig. 5).

Fig. 5 Growth trends of basal area increment trends of beech for the four sites. Trends were based on

the best-fitted generalized additive mixed models (GAMM) for the period 1965-2014.

Long-term carbon isotope chronologies and water use strategies

354 At the southernmost site (CAL), trees presented the highest increase of $\delta^{13}C$ values that translate in

355 an increase of ᵢWUE (Table 3, Fig. 6). On the contrary, CAM showed the lowest value of ᵢWUE

356 (Table 3, Fig. 6). LAZ and TRE on average presented similar $\delta^{13}C$ and ᵢWUE (Table 3).

357 **Table 3** Statistics of mean $\delta^{13}C$ and ${}_i WUE$ of the tree-ring width series of beech per site for the period

358 1965–2014. Data are mean values ± SE.

Site	$\delta^{13}C$ (‰)	${}_i WUE$ (µmol mol^{-1})
TRE	−25.9 ± 1.4	82.4 ± 20.4
LAZ	−25.1 ± 1.1	89.6 ± 8.7
CAM	−26.4 ± 0.9	76.9 ± 14.8
CAL	−24.8 ± 0.9	93.4 ± 13.5

For most sites, $\delta^{13}C$ showed a positive and significant trend over time ($P< 0.05$, Fig. 6), except for

LAZ, which showed an opposite pattern during the period 1965–2014 ($P< 0.001$, Fig. 6). In the

northernmost site (TRE), the $\delta^{13}C$ and ${}_i WUE$, started to increase sharply after the drought of 2003.

Similarly, the southern sites CAM and CAL presented a steady increase in ${}_i WUE$ ($P< 0.001$, Fig. 6).

On the contrary, LAZ did not present any significant trend ($P = 0.701$).

Fig. 6 Trends of: a) $\delta^{13}C$ (‰), and b) ${}_i WUE$, and fitted linear trends for the period 1965–2014 in four

stands across a latitudinal gradient in Italy.

In the southern sites, CAM and CAL, we observed significant positive relationships between $_i$WUE and BAI ($P < 0.001$, Fig. 7). On the contrary, at the northern site (TRE), we observed the opposite trend pattern; however, this trend was not significant ($P > 0.05$). AT LAZ, no relationship was found between $_i$WUE and BAI (Fig. 7).

Fig. 7 Relationship between annual BAI and $_i$WUE in beech across the Italian Peninsula for the period 1965-2014. Linear regressions and the equations are indicated for each site. Significance values are encoded by ***$P < 0.001$.

δ^{13}C, $_i$WUE, and climate relationship

Carbon isotope composition (δ^{13}C) and $_i$WUE showed a similar relationship with climate variables. However, $_i$WUE presented stronger correlation with climate than δ^{13}C (Figs. 8, S4). An exception was observed at LAZ, where δ^{13}C showed a negative and significant correlation with T_{mean} compared to $_i$WUE (Figs. 8, S4).

At the northernmost site, ¡WUE showed significant and positive correlations with T_{mean} and VPD of the previous and current year, while negative and scattered correlations with P and SPEI of April and May were observed (Fig. 8).

At LAZ, ¡WUE was negatively and significantly correlated with VPD from March to November of the current and previous year.

At CAM, ¡WUE exhibited a positive and significant correlation with P from March to May (previous and current year), while T_{mean} of the current and previous year was positively and significantly correlated with ¡WUE. On the contrary, ¡WUE correlated strongly and negatively with VPD from May to August (current and previous year) and with SPEI in August (Fig. 8).

At the southernmost site, ¡WUE correlated positively with P from March to September (current year), and strongly and positively with T_{mean} from March to August of the previous year. At CAL, VPD and SPEI exhibited strong and negative correlations with ¡WUE from May to June (current and previous year, Fig. 8).

Fig. 8 Pearson's running correlations between ¡WUE with monthly precipitation, mean temperature, VPD, and SPEI1 for the current and the previous year (*), over the period 1965-2014 at each site. The *y-axis* represents the time window in months. Colours (see the key) represent correlation coefficients that are significant at the level of $r = 0.279$ ($P < 0.05$)

Early warning signals of declining forest resilience

The statistical analysis of the BAI time series performed to detect EWS on beech forests revealed contrasting results among the sites (Fig. 9a,b). In TRE and LAZ, BAI showed a rise in AR1 among trees, which started to increase after the 2003 drought in TRE, while in LAZ already during the 1990s (Fig. 9a). In contrast, CAM showed a significant steady decrease in AR1. No significant autocorrelation trend was found at CAL, Nevertheless, the standard deviation (SD) started to rise by the end of the 1980s (Fig. 9b). A significant increase in SD of the BAI signal was observed across all the sites.

Fig. 9 Early warning signals: a) AR1, first-order autocorrelation, b) SD, obtained using a 15-year moving window for basal area increment (BAI) of *Fagus sylvatica* for four study sites for the period 1965-2014. The statistics of BAI were calculated using the residuals of the time series after removing the low-frequency signal (Gaussian filter) using 15-year-long windows (e.g. 1979 corresponds to the interval 1965-2014). The Kendall τ statistics indicate the strength of trends along the time series for each variable and site. For each site, the bold line represents the mean of the statistics among trees, and the shaded area is the standard error. (* $P < 0.05$, ** $P < 0.001$, *** $P < 0.001$).

Discussion

Long-term growth patterns and climate variability impact on beech across Italy

Our analysis revealed diverse long-term growth responses of European beech across the Italian Peninsula, closely associated with local climate and site conditions. The northern sites (TRE and LAZ) showed a decrease in BAI trends after the severe drought event in 2003, while the southernmost site (CAL) exhibited a growth decline after 2010. In contrast, CAM displayed a steady increase in growth over the 50 years analysed, likely due to increased precipitation in the last decades.

Our findings confirmed that European beech in the northern sites might be more susceptible to die-off, even without visible decline symptoms (such as branch dieback or decolouration of leaves). Trees exhibited greater growth sensitivity to VPD during summer, and this effect became more pronounced at the southernmost site. VPD can be used to estimate atmospheric water status and is one of the most important environmental factors influencing plant growth (Zabri & Burrage, 1997). Elevated VPD, associated with dry conditions, impacts stomatal conductance and the balance between carbon assimilation and water loss (McDowell *et al.*, 2008; Zhang *et al.*, 2015). This indicates that drought, driven by enhanced evapotranspiration, will play a critical role throughout the beech forest's latitudinal range in Italy.

Recent global-scale research by Yuan *et al.* (2019) highlighted the increase of VPD as a major atmospheric driver affecting forest productivity by imposing water stress on photosynthesis. Water-use strategies, particularly conservative/opportunistic responses within and across species, have been closely linked to soil moisture availability (Moreno-Gutiérrez *et al.*, 2012; Martínez-Vilalta *et al.*, 2014). Higher VPD and temperature accelerate soil moisture depletion causing a significant reduction in carbon uptake (Sulman *et al.*, 2016), elevating the risk of drought-induced dieback through hydraulic failure and/or carbon starvation (Anderegg *et al.*, 2016; Adams *et al.*, 2017; Grossiord *et al.*, 2020).

Our results demonstrate a significant increase in VPD after the 2003 drought event in TRE, however we found a weak negative correlation between VPD and BAI. We can hypothesise this might be attributed to a lower soil water holding capacity (SWHC) at this site, potentially increasing vulnerability to growth decline, as observed in our GAMM model. Conversely, LAZ exhibited higher SWHC, likely contributing to higher transpiration rates and growth compared to other sites. CAM and CAL sites presented moderate SWHC and a declining VPD trend, indicating less stress than the TRE site. The GAMM model integrated responses to the SPEI index and individual age of each tree, thus, we speculate that hydraulic strategies under drought significantly impact long-term growth rates,

reflecting site-specific and ontogenic plasticity responses of the species. These findings may suggest that young beech trees initially benefit from favourable climate conditions and higher transpiration rates; however, this advantage depends on soil water availability and makes them susceptible to rapid declines in growth during extreme drought events, as already observed in Switzerland (Vanoni *et al.*, 2016).

Our study, to our knowledge, is the first to show evidence of the negative impact of VPD on basal area increment in beech forests across the Italian peninsula. This correlation was evident in all sites but was even stronger at southern latitudes. In contrast, previous studies in mature beech stands did not find a significant climate correlation, attributed to the species' mast-seeding behaviour and sensitivity to late frosts at the beginning of the growing season (Piovesan & Adams, 2001; Castagneri *et al.*, 2014; D'Andrea *et al.*, 2021; Tonelli *et al.*, 2023). Other studies have identified lagged climate correlation with masting (Vacchiano *et al.*, 2017). Additionally, Zimmermann *et al.* (2015), in central Germany, found that beech growth was highly sensitive to summer temperatures and extreme drought events after the 1980s.

Drought sensitivity and water-use strategies effect on growth

European beech has commonly been classified as an opportunistic species, capable of maintaining higher transpiration rates even in relatively dry soil conditions (Leuschner, 2020; and references therein); however, this strategy increases the risk of cavitation (McDowell *et al.*, 2008; Martinez-Vilalta *et al.*, 2014).

Our findings indicated that temperature and VPD emerged as primary drivers of iWUE in TRE, while VPD played a dominant role in the southern sites. However, in LAZ, iWUE did not exhibit a clear correlation with climate variables. This complex relationship highlights the interaction between VPD, stomatal conductance, and photosynthesis, as high VPD initially reduces stomatal conductance but not net CO_2 assimilation rate, resulting in increased iWUE. Nevertheless, severe VPD-induced stomatal conductance restrictions, combined with declining soil moisture and other non-stomatal limitations, ultimately reduce photosynthetic rate and may lead to declining iWUE as VPD continues to rise (Flexas *et al.*, 2012). Thus, the overall relationship between iWUE and VPD is likely hyperbolic (Zhang *et al.*, 2019), and the sensitivity of photosynthesis to VPD will likely be weaker than the sensitivity of conductance to VPD.

Our study highlights contrasting iWUE of beech across the Italian peninsula. We observed an increase of $\delta^{13}C$ and iWUE values in TRE, CAM, and CAL, indicating a conservative water use strategy when

water availability is low. In contrast, LAZ exhibited a decrease of $\delta^{13}C$, suggesting an opportunistic response with stable $_iWUE$ regardless of the moisture condition. While at LAZ, changes in photosynthetic rate and stomatal conductance appeared to occur in the same direction with similar magnitude, at TRE, CAM, and CAL, stomatal conductance appeared to decrease proportionally more than photosynthetic rate, or the latter remaining stable or increasing with declining stomatal conductance. Thus, our findings confirm that water use strategies employed by beech are mostly site-specific and influenced by microclimatic conditions and soil water availability (McCarroll & Loader, 2004), consistent with our hypothesis and consistent with prior research (Peñuelas et al., 2008).

Interestingly, our results indicate that higher mean $_iWUE$ did not result in an increase in the basal area increment on beech (Nock et al., 2011; Peñuelas et al., 2011; Mazza et al., 2024); instead, we observed contrasting responses consistent with previous studies (Peñuelas et al., 2008; Tognetti et al., 2014). Notably, the northern site displayed a drastic increase in $_iWUE$ after the 2003 drought event, coinciding with elevated VPD and temperature that may have led to stomatal closure (g_s) and reduced photosynthesis (A), suggesting that the growth decline in this site was might triggered by intensified evapotranspiration and the lower SWHC as observed in other sites by others (e.g. Peñuelas et al., 2011; Li et al., 2023). At LAZ, there was no relationship between $_iWUE$ and growth, which can be explained by higher SWHC allowing higher transpiration rates and metabolic respiration, resulting in greater losses of photosynthetic assimilates, especially at higher temperatures (Nock et al., 2011; Mazza et al., 2024). Interestingly, in the southern sites, the increase of $_iWUE$ enhanced growth. This discrepancy may be attributed to the adaptation of beech trees in the southernmost distribution to water stress and high VPD (Anderegg et al., 2019; Battipaglia & Cherubini, 2022), suggesting that high $_iWUE$ is an adaptative trait (Medrano et al., 2009). Consequently, we can infer that the observed "conservative strategy" - characterized by low stomatal conductance and constant CO_2 assimilation rate that enhanced growth – at CAM might be explained by a positive CO_2 fertilization effect or long-term acclimation to elevated CO_2 (Walker et al., 2021). Similar findings were reported in mature European beech stands in Spain, where an increased sensitivity to drought was observed across the southern range-edge distribution (Peñuelas et al., 2008). Recently, Qi et al. (2023) in China revealed varying water use strategies among larch trees. Mature trees presented a more 'conservative strategy' (low g_s, constant assimilation rate (A)), whereas young trees maintained constant g_s and high A, indicating an opportunistic behaviour. Similar to our study, mature trees displayed greater sensitivity to atmospheric CO_2 concentrations than their young counterparts.

It should be pointed out that a major influence of photosynthetic rate on intercellular CO_2 concentration and $\delta^{13}C$, and the minor contribution of the regulation of stomatal conductance to $_iWUE$, were observed in other studies on the same species (Tognetti et al., 2014). These findings

suggest unclear patterns of potential increased drought-related tree decline signs in mountain beech

forests along the Italian latitudinal transect. Differences between leaf-level physiology and tree-ring

level processes may arise, reflecting potential variations in the (re)translocation patterns of non-

structural carbohydrates to organs (Martínez-Vilalta *et al*., 2016; Merganičová et al., 2019). Such

complexities make tree-ring analysis a challenging tool for decipher tree responses to fluctuating

seasonal conditions in the short term.

Early Warning Signals of mortality

Our second hypothesis, linking the degree of growth reduction and tree growth instability to drought

severity, was only partially confirmed by our findings. We observed an increase in the autocorrelation

of the BAI signal across almost all sites, indicating an enhanced intrinsic biological memory within

the trees and a loss of ecological system resilience (Lloret *et al*., 2011; Seidel *et al*., 2022; Smith *et*

al., 2022). Such increases have been linked to instabilities preceding external disturbances in various

biological systems (e.g., Dakos *et al*., 2012a; Boulton *et al*., 2022; Forzieri *et al*., 2022), potentially

leading to a transition to a new system state (Majumder *et al*., 2019; Buxton *et al*., 2021). Recent

studies investigating ecosystem productivity's autocorrelation have identified reduced resilience in

diverse forest types due to increased water limitations and climate variability (Forzieri *et al*., 2022;

Fernandez-Martinez *et al*., 2023). Notably, after a severe drought, declining trees exhibited increases

in BAI autocorrelation and variability before mortality (Camarero *et al*., 2015).

In line with our expectations, the northern site showed a significant increase in AR1 and a decline in

BAI after the 2003 drought event. Conversely, CAM showed a decrease in AR1, suggesting greater

resilience to changing climate conditions, despite experiencing the severe drought period of 2003.

This higher resilience at CAM might be linked to the legacy of past conditions with less intra-annual

variability in water availability compared to the TRE site, as supported by the SPEI multiscalar index

(Fig. 2). TRE experienced several prolonged dry periods (i.e. SPEI<-1.5) before the 2000s.

Additionally, relatively mature trees at CAM site might contribute to the population's apparent

stability (see Colangelo *et al*., 2021). Our data also revealed an increase in BAI series SD across all

stands. While this variability encompasses both tree physiological signals and climate-driven

vegetation dynamics (Bochow & Boers, 2023), the co-occurrence rises in AR1, decline in BAI, and

increase in SD in TRE, LAZ and CAL sites, may indicate a loss of system stability (Dakos *et al*.,

2012a). These results indicate potential challenges for trees to mitigate the impact of extreme events

in the future.

Several studies have demonstrated that long-term rises in instability and reduced growth predispose European beech to elevated mortality risks under future climate-induced stress conditions (Gillner *et al*., 2013; DeSoto *et al*., 2020; Cabon *et al*., 2023). The loss of stability in our study sites emphasizes the need for continuous monitoring and proactive management of beech forests, particularly in regions where climate change is projected to increase the frequency and severity of droughts. Ongoing monitoring enables early detection of tree mortality risks, facilitating timely interventions to protect and sustain these vital ecosystems with wide ecological amplitude.

In conclusion, in this study, our goal was to advance the early prediction of mortality risk in healthy beech stands without, apparently, visible declining symptoms across the Italian Peninsula. The aforementioned is especially pertinent considering the recent growth decline observed in this species (Piovesan *et al*., 2008; Martinez del Castillo *et al*., 2022; Dorado-Liñan *et al*., 2022), although the available evidence is not yet conclusive (Tognetti *et al*., 2014).

These findings highlight the importance of considering the plasticity and site-specific ¡WUE responses to varying environmental conditions and the impact of VPD on stomatal conductance when predicting the future of beech forests in the context of climate change. It is important to note that not all beech populations considered in this study exhibited an increase in ¡WUE in response to rising VPD. This variability reflects differing sensitivities to changes in environmental drivers and the plasticity of conservative to opportunistic water-use strategies.

Furthermore, our analysis of EWS reveals the loss of resilience after an extreme event, as notably observed at the TRE site. In the context of climate model projection, the increase in the frequency and severity of droughts, the ability to detect earlier tipping points of critical slowdowns in declining systems, and the potential for recovery to the current state or an alternative state remains uncertain (Cabon *et al*., 2023).

Nonetheless, this research raises further questions, such as how to generalize the relationships between increased ¡WUE and conservative behaviour, thus explaining contradictory results obtained in tree ring studies on beech populations and assessing temporal changes in this functional trait. Further research considering young and mature trees and their physiological mechanisms (Leuschner, 2020; Qi *et al*., 2023), micro-site conditions (Puchi *et al*., 2021), and genetics will also elucidate intraspecific variations in drought response (Alderotti *et al*., 2023). This knowledge is essential for developing effective conservation and future forest management strategies to ensure these crucial ecological and socio-economical ecosystems' long-term health, vitality, and resilience.

Acknowledgments

PFP, GB and AC were supported by the MIUR project (PRIN 2020) "Unraveling interactions between WATER and carbon cycles during drought and their impact on water resources and forest and grassland ecosySTEMs in the Mediterranean climate" (WATERSTEM, cod. 20202WF53Z). EV, and AC were supported by the MIUR project (PRIN 2020) "Multi-scale observations to predict Forest response to pollution and climate change" (MULTIFOR, cod. 2020E52THS). PFP, DD, EV and AC acknowledge funding by the project FORESTNAVIGATOR Horizon Europe research and innovation programme under grant agreement No. 101056875 and by OptForEU H2020 research and innovation programme under grant agreement No. 101060554; AC was also supported by resources available from the Italian Ministry of University and Research (FOE-2019), under projects 'Climate Changes' (CNR DTA. AD003.474.029). R.T. was supported by the Cost Action CA15226 - Climate-Smart Forestry in Mountain Regions (CLIMO). We also acknowledge the project funded under the National Recovery and Resilience Plan (NRRP), Mission 4 Component 2 Investment 1.4 - Call for tender No. 3138 of 16 December 2021, rectified by Decree n.3175 of 18 December 2021 of Italian Ministry of University and Research funded by the European Union – NextGenerationEU under award Number: Project code CN_00000033, Concession Decree No. 1034 of 17 June 2022 adopted by the Italian Ministry of University and Research, CUP B83C22002930006, Project title "National Biodiversity Future Centre - NBFC".

Author contributions

PFP Conceptualization, data curation, formal analysis, writing – original draft; DD formal analysis, methodology, writing- review & editing; EV Methodology, Visualization, Writing-review & editing; GB Methodology, formal analysis; resources; writing-review & editing; RT data curation, writing-review & editing; AC Funding acquisition, project administration, writing-review & editing.

ORCID

Paulina Puchi: 0000-0001-5429-8605

Daniela Dalmonech: 0000-0002-1932-5011

Elia Vangi: 0000-0002-9772-2258

Giovanna Battipaglia: 0000-0003-1741-3509

Roberto Tognetti: 0000-0002-7771-6176

Alessio Collalti: 0000-0002-4980-8487

References

Adams HD, Zeppel MJB, Anderegg WRL, Hartmann H, Landhäusser SM, Tissue DT, Huxman TE, Hudson PJ, Franz TE, Allen CD, *et al.* **2017**. A multi-species synthesis of physiological mechanisms in drought-induced tree mortality. *Nature Ecology and Evolution* **1**: 1285–1291.

Alderotti F, Sillo F, Brilli L, Bussotti F, Centritto M, Ferrini F, Gori A, Inghes R, Pasquini D, Pollastrini M, *et al.* **2023**. *Quercus ilex* L. dieback is genetically determined: Evidence provided by dendrochronology, δ¹³C and SSR genotyping. *Science of The Total Environment* **904**: 166809.

Allen CD, Breshears DD, McDowell NG. **2015**. On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* **6**: 1–55.

Allen CD, Macalady AK, Chenchouni H, Bachelet D, Mcdowell N, Vennetier M, Kitzberger T, Rigling A, Breshears DD, Hogg EH (Ted) T, *et al.* **2010**. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* **259**: 660–684.

Allen RG. 1993. *Evaluation of a temperature difference method for computing grass reference evapotranspiration.* Report submitted to FAO, Rome.

Anderegg WRL, Anderegg LDL, Kerr KL, Trugman AT. **2019**. Widespread drought-induced tree mortality at dry range edges indicates that climate stress exceeds species' compensating mechanisms. *Global Change Biology* **25**: 3793–3802.

Anderegg WRL, Klein T, Bartlett M, Sack L, Pellegrini AFA, Choat B, Jansen S. **2016**. Meta-analysis reveals that hydraulic traits explain cross-species patterns of drought-induced tree mortality across the globe. *Proceedings of the National Academy of Sciences of the United States of America* **113**: 5024-5029.

Anderegg WRL, Kane JM, Anderegg LDL. **2013**. Consequences of widespread tree mortality triggered by drought and temperature stress. *Nature Climate Change* **3**: 30–36.

Battipaglia G, Cherubini P. 2022. Stable Isotopes in Tree Rings of Mediterranean Forests. In: Siegwolf RTW, Brooks JR, Roden J, Saurer M, eds. *Stable Isotopes in Tree Rings*. Tree Physiology, vol 8. Springer, 605–630.

Battipaglia G, Pelleri F, Lombardi F, Altieri S, Vitone A, Conte E, Tognetti R. 2017. Effects of associating *Quercus robur* L. and *Alnus cordata* Loisel. on plantation productivity and water use efficiency. *Forest Ecology and Management* **391**: 106–114.

Beguería S, Vicente-Serrano SM. 2017. Package 'SPEI'. https://cran.r-project.org/web/packages/SPEI/index.html

Beguería S, Vicente-Serrano SM, Reig F, Latorre B. 2014. Standardized precipitation evapotranspiration index (SPEI) revisited: Parameter fitting, evapotranspiration models, tools, datasets and drought monitoring. *International Journal of Climatology* **34**: 3001–3023.

Belmecheri S, Lavergne A. 2020. Compiled records of atmospheric CO_2 concentrations and stable carbon isotopes to reconstruct climate and derive plant ecophysiological indices from tree rings. *Dendrochronologia*, **63**: 125748.

Biondi F, Qeadan F. 2008a. A theory-driven approach to tree-ring standardization: Defining the biological trend from expected basal area increment. *Tree-Ring Research*, **64**: 81–96.

Biondi F, Qeadan F. 2008b. Removing the tree-ring width biological trend using expected basal area increment. in
USDA *Forest Service* RMRS-P-55 124–131.

Bochow N, Boers N. 2023. The South American monsoon approaches a critical transition in response to deforestation.
Science Advances, **9**: eadd9973.

Boulton CA, Lenton TM, Boers N. 2022. Pronounced loss of Amazon rainforest resilience since the early 2000s. *Nature*
Climate Change, **12**: 271-278.

Brodribb TJ, Powers J, Cochard H, Choat B. **2020**. Hanging by a thread? Forests and drought. *Science* **368**: 261–266.

Bunn A, Korpela M, Biondi F, Campelo F, Mérian P, Qeadan F, Zang C. 2021. dplR: Dendrochronology program
library in R. https://CRAN.R--proje ct.org/package=dplR

Bunn AG. 2008. A dendrochronology program library in R (dplR). *Dendrochronologia*, **26**: 115–124.

Burnham KP, Anderson DR. 2002. Model Selection and Inference: A Practical Information-Theoretic Approach. 2nd
Edition, Springer-Verlag, New York. 355.

Buxton JE, Abrams JF, Boulton CA, Barlow N, Rangel Smith C, Van Stroud S, Lees KJ, Lenton TM. **2022**.
Quantitatively monitoring the resilience of patterned vegetation in the Sahel. *Global Change Biology* **28**: 571–587.

Cabon A, DeRose RJ, Shaw JD, Anderegg WRL. **2023**. Declining tree growth resilience mediates subsequent forest
mortality in the US Mountain West. *Global Change Biology* **29**: 4826–4841

Cailleret M, Dakos V, Jansen S, Robert EMR, Aakala T, Amoroso MM, Antos JA, Bigler C, Bugmann H,
Caccianaga M, *et al.* **2019**. Early-warning signals of individual tree mortality based on annual radial growth.
Frontiers in Plant Science **9**: 1964.

Cailleret M, Jansen S, Robert EMR, Janda P, Kane JM, Kharuk VI, Tognetti R, Jos E. **2016**. A synthesis of radial
growth patterns preceding tree mortality. *Global Change Biology* **23**: 1675–1690.

Camarero JJ, Gazol A, Sangüesa-Barreda G, Oliva J, Vicente-Serrano SM. **2015**. To die or not to die: Early warnings
of tree dieback in response to a severe drought. *Journal of Ecology* **103**: 44–57.

Castagneri D, Nola P, Motta R, Carrer M. **2014**. Summer climate variability over the last 250years differently affected
tree species radial growth in a mesic Fagus-Abies-Picea old-growth forest. *Forest Ecology and Management* **320**: 21–
29.

Cherubini P, Battipaglia G, Innes J. **2021**. Tree Vitality and Forest Health: Can Tree-Ring Stable Isotopes Be Used as
Indicators? *Current Forestry Reports* **7**: 69–80.

Colangelo M, Camarero JJ, Battipaglia G, Borghetti M, De Micco V, Gentilesca T, Ripullone F. **2017**. A multi-
proxy assessment of dieback causes in a Mediterranean oak species. *Tree Physiology* **37**: 617–631.

Cook ER, Kairiukstis LA. 1990. *Methods of dendrochronology*: Applications in the environmental sciences. Dordrecht,
Boston, MA: Kluwer Academic.

Gasparini P, Di Cosmo L, Floris A. 2022. Area and Characteristics of Italian Forests. In: Gasparini P, Di Cosmo L, Floris A, De Laurentis D, eds. *Italian National Forest Inventory—Methods and Results of the Third Survey*. Springer Tracts in Civil Engineering. Springer, 391–446.

Gillner S, Rüger N, Roloff A, Berger U. 2013. Low relative growth rates predict future mortality of common beech (*Fagus sylvatica* L.). *Forest Ecology and Management* **302**: 372–378.

Hargreaves GL, Hargreaves GH, Riley JP. 1985. Irrigation water requirements for Senegal River Basin. *Journal of Irrigation and Drainage Engineering* **111**: 265–275.

Hengl T, Nauman T. 2018. Predicted USDA soil great groups at 250 m (probabilities) (Version v01) [Data set]. Zenodo. 10.5281/zenodo.1476844

Holmes RL. 1983. Computer-assisted quality control in tree-ring dating and measurement. *Tree-Ring Bulletin* **43**: 69–78.

Dakos V, Carpenter SR, Brock WA, Ellison AM, Guttal V, Ives AR, Kéfi S, Livina V, Seekell DA, van Nes EH, *et al.* 2012a. Methods for detecting early warnings of critical transitions in time series illustrated using simulated ecological data. *PLoS ONE* **7**: e41010.

Dakos V, Van Nes EH, D'Odorico P, Scheffer M. 2012b. Robustness of variance and autocorrelation as indicators of critical slowing down. *Ecology* **93**: 264–271.

D'Andrea E, Scartazza A, Battistelli A, Collalti A, Proietti S, Rezaie N, Matteucci G, Moscatello S. 2021. Unravelling resilience mechanisms in forests: role of non-structural carbohydrates in responding to extreme weather events. *Tree Physiology* **41**: 1808–1818.

DeSoto L, Cailleret M, Sterck F, Jansen S, Kramer K, Robert EMR, Aakala T, Amoroso MM, Bigler C, Camarero JJ, *et al.* 2020. Low growth resilience to drought is related to future mortality risk in trees. *Nature Communications* **11**: 545.

De Marco A, Sicard P, Feng Z, Agathokleous E, Alonso R, Araminiene V, Augustatis A, Badea O, Beasley JC, Branquinho C, *et al.* 2022. Strategic roadmap to assess forest vulnerability under air pollution and climate change. *Global Change Biology* **28**: 5062–5085.

Dorado-Liñán I, Ayarzagüena B, Babst F, Xu G, Gil L, Battipaglia G, Buras A, Čada V, Camarero JJ, Cavin L, *et al.* 2022. Jet stream position explains regional anomalies in European beech forest productivity and tree growth. *Nature Communications* **13**: 2015.

Fang J, Lechowicz MJ. 2006. Climatic limits for the present distribution of beech (*Fagus* L.) species in the world. *Journal of Biogeography* **33**: 1804–1819.

Farquhar GD, O'Leary MH, Berry JA. 1982. On the relationship between carbon isotope discrimination and the intercellular carbon dioxide concentration in leaves. *Australian Journal of Plant Physiology* **9**: 121–137.

Farquhar GD, Richards RA. 1984. Isotopic composition of plant carbon correlates with water-use efficiency of wheat genotypes. *Australian Journal of Plant Physiology* **11**: 539–552.

Fernández-Martínez M, Peñuelas J, Chevallier F, Ciais P, Obersteiner M, Rödenbeck C., ... & Janssens, I. A.
(2023). Diagnosing destabilization risk in global land carbon sinks. *Nature* **615:** 848–853.

Flanagan LB, Farquhar GD. **2014**. Variation in the carbon and oxygen isotope composition of plant biomass and its
relationship to water-use efficiency at the leaf- and ecosystem-scales in a northern Great Plains grassland. *Plant, Cell*
and Environment **37**: 425–438.

Flexas J, Barbour MM, Brendel O, Cabrera HM, Carriquí M, Díaz-Espejo A, Douthe C, Dreyer E, Ferrio JP,
Gago J, *et al.* **2012**. Mesophyll diffusion conductance to CO_2: An unappreciated central player in photosynthesis.
Plant Science **193–194**: 70–84.

Francey RJ, Allison CE, Etheridge DM, Trundinger CM, Enting IG, Leuenberger M, Langenfelds RL, Michel E,
Steele LP. 1999. A 1000-year high precision record of $\delta^{13}C$ in atmospheric CO_2. Tellus B: *Chemical and Physical*
Meteorology **51**: 170–193.

Forzieri G, Dakos V, McDowell NG, Ramdane A, Cescatti A. **2022**. Emerging signals of declining forest resilience
under climate change. *Nature* **608**: 534–539.

Gagen M, Battipaglia G, Daux V, et al., 2022. Climate signals in stable isotope tree-ring records. In: Siegwolf RTW,
Brooks JR, Roden J, Saurer M, eds. *Stable Isotopes in Tree Rings*. Tree Physiology vol. 8. Springer, Cham, pp. 537–
579.

Gessler A, Bottero A, Marshall J, Arend M. **2020**. The way back: recovery of trees from drought and its implication
for acclimation. *New Phytologist* **228**: 1704–1709.

Gessler A, Cailleret M, Joseph J, Schönbeck L, Schaub M, Lehmann M, Treydte K, Rigling A, Timofeeva G,
Saurer M. **2018**. Drought induced tree mortality – a tree-ring isotope based conceptual model to assess mechanisms
and predispositions. *New Phytologist* **219**: 485–490.

Grossiord C, Buckley TN, Cernusak LA, Novick KA, Poulter B, Siegwolf RTW, Sperry JS, McDowell NG. **2020**.
Plant responses to rising vapor pressure deficit. *New Phytologist* **226**: 1550–1566.

Holmes RL. 1983. Computer-assisted quality control in tree-ring dating and measurement. *Tree-Ring Bulletin* **43**: 69–
78.

IPCC. 2021. Assessment report 6 climate change 2021: The physical science basis. https://www.ipcc.ch/report/ar6/wg1/

Jevšenak J, Levanič T. **2018**. *dendroTools*: R package for studying linear and nonlinear responses between tree-rings
and daily environmental data. *Dendrochronologia* **48**: 32–39.

Keenan JR. 2015. Climate change impacts and adaptation in forest management: a review. *Annals of Forest Science* **72**:
145–167.

Kutílek M, Nielsen DR. 1994. Soil Hydrology: Textbook for Students of Soil Science, Agriculture, Forestry,
Geoecology, Hydrology, Geomorphology and Other Related Disciplines; Catena Verlag, Germany, pp. 370.

Leuschner C. 2020. Drought response of European beech (*Fagus sylvatica* L.)—A review. *Perspectives in Plant Ecology,*
Evolution and Systematics **47**: 125576.

Li F, Xiao J, Chen J, Ballantyne A, Jin K, Li B, Abraha M, John R. **2023**. Global water use efficiency saturation due
to increased vapor pressure deficit. *Science* **381**: 672–677.

Lloret F, Keeling EG, Sala A. 2011. Components of tree resilience: effects of successive low-growth episodes in old
ponderosa pine forests. *Oikos* **120**: 1909-1920.

Majumder S, Tamma K, Ramaswamy S, Guttal V. 2019. Inferring critical thresholds of ecosystem transitions from
spatial data. *Ecology* **100**: e02722.

Martinez del Castillo E, Zang CS, Buras A, Hacket-Pain A, Esper J, Serrano-Notivoli R, Hartl C, Weigel R, Klesse
S, Resco de Dios V, *et al*. 2022. Climate-change-driven growth decline of European beech forests. *Communications*
Biology **5**: 163.

Martínez-Vilalta J, Poyatos R, Aguadé D, Retana J, Mencuccini M. **2014**. A new look at water transport regulation
in plants. *New Phytologist* **204**: 105–115.

Mazza G, Monteverdi MC, Altieri S, Battipaglia G. **2024**. Climate-driven growth dynamics and trend reversal of *Fagus*
sylvatica L. and *Quercus cerris* L. in a low-elevation beech forest in Central Italy. *Science of The Total Environment*
908: 168250.

McCarroll D, Loader NJ. **2004**. Stable isotopes in tree rings. *Quaternary Science Reviews* **23** 771–801.

McDowell NG, Allen CD, Anderson-Teixeira K, Aukema BH, Bond-Lamberty B, Chini L, Clark JS, Dietze M,
Grossiord C, Hanbury-Brown A, *et al*. 2020. Pervasive shifts in forest dynamics in a changing world. *Science* **368**:
964.

Mcdowell N, Pockman WT, Allen CD, Breshears DD, Cobb N, Kolb T, Plaut J, Sperry J, West A, Williams DG, *et*
***al*. 2008**. Mechanisms of Plant Survival and Mortality during Drought: Why Do Some Plants Survive while Others
Succumb to Drought? *New Phytologist* **178**: 719–739.

Medrano H, Flexas J, Galmés J. 2009. Variability in water use efficiency at the leaf level among Mediterranean plants
with different growth forms. *Plant Soil* **317**:17–29.

Noce S, Collalti A, Santini M. 2017. Likelihood of changes in forest species suitability, distribution, and diversity under
future climate: The case of Southern Europe. *Ecology and Evolution* **7**: 9358–9375.

Noce S, Collalti A, Valentini R, Santinin M. 2016. Hot spot maps of forest presence in the Mediterranean basin. *iForest*
- Biogeosciences and Forestry **9**: 766–774.

Nock CA, Baker PJ, Wanek W, Leis A, Grabner M, Bunyavejchewin S, Hietz P. 2011. Long-term increases in
intrinsic water-use efficiency do not lead to increased stem growth in a tropical monsoon forest in western Thailand.
Global Change Biology, **17**: 1049–1063.

Pacheco A, Camarero JJ, Pompa-García M, Battipaglia G, Voltas J, Carrer M. 2020. Growth, wood anatomy and
stable isotopes show species--specific couplings in three Mexican conifers inhabiting drought-prone areas. *Science of*
the Total Environment **698**: 134055.

Pan Y, Birdsey RA, Fang J, Houghton R, Kauppi PE, Kurz WA, Phillips OL, Shvidenko A, Canadell SLLJG,
Ciais P, *et al*. 2011. A Large and Persistent Carbon Sink in the World's Forests. *Science* **333**: 988–993.

Peñuelas J, Canadell JG, Ogaya R. **2011**. Increased water-use efficiency during the 20th century did not translate into enhanced tree growth. *Global Ecology and Biogeography* **20**: 597–608.

Peñuelas J, Hunt JM, Ogaya R, Jump AS. **2008**. Twentieth century changes of tree-ring $\delta^{13}C$ at the southern range-edge of *Fagus sylvatica*: Increasing water-use efficiency does not avoid the growth decline induced by warming at low altitudes. *Global Change Biology* **14**: 1076–1088.

Petrucco L, Nardini A, Von Arx G, Saurer M, Cherubini P. **2017**. Isotope signals and anatomical features in tree rings suggest a role for hydraulic strategies in diffuse drought-induced die-back of *Pinus nigra*. *Tree Physiology* **37**: 523–535.

Piovesan G, Biondi F, Di Filippo A, Alessandrini A, Maugeri M. **2008**. Drought-driven growth reduction in old beech (*Fagus sylvatica* L.) forests of the central Apennines, Italy. *Global Change Biology* **14**: 1265–1281.

Piovesan G, Adams JM. 2001. Masting behaviour in beech: linking reproduction and climatic variation. *Canadian Journal of Botany* **79**: 1039–1047.

Puchi PF, Camarero JJ, Battipaglia G, Carrer M. **2021**. Retrospective analysis of wood anatomical traits and tree-ring isotopes suggests site-specific mechanisms triggering *Araucaria araucana* drought-induced dieback. *Global Change Biology* **27**: 6394–6408.

Qi X, Cherubini P, Treydte K, Li MH, Wu Z, He HS, Du H, Fang K, Saurer M. **2023**. Growth responses to climate warming and their physiological mechanisms differ between mature and young larch trees in a boreal permafrost region. *Agricultural and Forest Meteorology* **343**: 109765.

R Core Team 2022. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.

Sangüesa-Barreda G, Camarero JJ, Sánchez-Salguero R, Gutiérrez E, Linares JC, Génova M, Ribas M, Tíscar PA, López-Sáez JA. **2019**. Droughts and climate warming desynchronize Black pine growth across the Mediterranean Basin. *Science of the Total Environment* **697**: 133989.

Seibt U, Rajabi A, Griffiths H, Berry J. 2008. Carbon isotopes and water use efficiency: Sense and sensitivity. *Oecologia* **155**: 441–454.

Seidl R, Jentsch A, Wohlgemuth T. 2022. Disturbance Resilience. In Seidl R, Jentsch A, Wohlgemuth T, eds in *Disturbance Ecology*. Springer International Publishing, 97–115

Smith T, Traxl D, Boers N. 2022. Empirical evidence for recent global shifts in vegetation resilience. *Nature Climate Change 12*(5), 477-484.

Stokes MA, Smiley TL. **1968**. *An introduction to tree-ring dating*. Chicago, IL: University of Chicago Press.

Sulman BN, Roman DT, Yi K, Wang L, Phillips RP, Novick KA. **2016**. High atmospheric demand for water can limit forest carbon uptake and transpiration as severely as dry soil. *Geophysical Research Letters* **43**: 9686–9695.

Timofeeva G, Treydte K, Bugmann H, Rigling A, Schaub M, Siegwolf R, Saurer M. **2017**. Long-term effects of drought on tree-ring growth and carbon isotope variability in Scots pine in a dry environment. *Tree Physiology* **37**: 1028–1041.

Tetens, O. 1930. Uber einige meteorologische Begriffe Z. Geophys. 6, 297–309.

Title PO, Bemmels JB. **2018**. ENVIREM: an expanded set of bioclimatic and topographic variables increases flexibility
and improves performance of ecological niche modeling. *Ecography* **41**: 291–307.

Tognetti R, Lombardi F, Lasserre B, Cherubini P, Marchetti M. **2014**. Tree-Ring Stable Isotopes Reveal Twentieth-
Century Increases In Water-Use Efficiency of *Fagus Sylvatica* and *Nothofagus* spp. in Italian and Chilean Mountains.
PLoS ONE **9**: e113136

Tonelli E, Vitali A, Malandra F, Camarero JJ, Colangelo M, Nolè A, Ripullone F, Carrer M, Urbinati C. **2023**.
Tree-ring and remote sensing analyses uncover the role played by elevation on European beech sensitivity to late
spring frost. *Science of the Total Environment* **857**: 159239

Trugman AT, Anderegg LDL, Anderegg WRL, Das AJ, Stephenson NL. **2021**. Why is Tree Drought Mortality so
Hard to Predict? *Trends in Ecology and Evolution* **36**: 520–532.

Vacchiano G, Hacket-Pain A, Turco M, Motta R, Maringer J, Conedera M, Drobyshev I, Ascoli D. **2017**. Spatial
patterns and broad-scale weather cues of beech mast seeding in Europe. *New Phytologist* **215**: 595–608.

Vanoni M, Bugmann H, Nötzli M, Bigler C. **2016**. Quantifying the effects of drought on abrupt growth decreases of
major tree species in Switzerland. *Ecology and Evolution* **6**: 3555–3570.

Versace S, Gianelle D, Garfì V, Battipaglia G, Lombardi F, Marchetti M, Tognetti R. **2020**. Interannual radial growth
sensitivity to climatic variations and extreme events in mixed-species and pure forest stands of silver fir and European
beech in the Italian Peninsula. *European Journal of Forest Research* **139**: 627–645.

Vicente-Serrano SM, Beguería S, López-Moreno JI. 2010. A Multi-scalar drought index sensitive to global warming:
The Standardized Precipitation Evapotranspiration Index - SPEI. *Journal of Climate* **23**: 1696–1718.

**Walker AP, De Kauwe MG, Bastos A, Belmecheri S, Georgiou K, Keeling RF, McMahon SM, Medlyn BE, Moore
DJP, Norby RJ**, *et al.* **2021a**. Integrating the evidence for a terrestrial carbon sink caused by increasing atmospheric
CO_2. *New Phytologist* **229**: 2413–2445.

Walthert L, Ganthaler A, Mayr S, Saurer M, Waldner P, Walser M, Zweifel R, von Arx G. **2021**. From the comfort
zone to crown dieback: Sequence of physiological stress thresholds in mature European beech trees across progressive
drought. *Science of the Total Environment* **753**: 141792.

Willmott CJ, Feddema JJ. 1992. A More Rational Climatic Moisture Index*. *The Professional Geographer* **44**: 84–88.

Wood SN. 2006. Low-rank scale-invariant tensor product smooths for generalized additive mixed models. *Biometrics*
62: 1025–1036.

Yuan W, Zheng Y, Piao S, Ciais P, Lombardozzi D, Wang Y, Ryu Y, Chen G, Dong W, Hu Z, *et al.* **2019**. Increased
atmospheric vapor pressure deficit reduces global vegetation growth. *Science Advances* **5**: eaax1396.

Zabri AW, Burrage SW. 1997. The effects of vapour pressure deficit (VPD) and enrichment with CO_2 on water relations,
photosynthesis, stomatal conductance and plant growth of sweet pepper (*Capsicum annum* l.) grown by nft. *Acta
Horticulturae* **449**: 561–568.

Zhang Q, Ficklin DL, Manzoni S, Wang L, Way D, Phillips RP, Novick KA. **2019**. Response of ecosystem intrinsic
water use efficiency and gross primary productivity to rising vapor pressure deficit. *Environmental Research Letters*
14: 074023

Zhang K, Kimball JS, Nemani RR, Running SW, Hong Y, Gourley JJ, Yu Z. **2015**. Vegetation Greening and Climate
Change Promote Multidecadal Rises of Global Land Evapotranspiration. *Scientific Reports* **5**:15956

Zimmermann J, Hauck M, Dulamsuren C, Leuschner C. **2015**. Climate Warming-Related Growth Decline Affects
Fagus sylvatica, But Not Other Broad-Leaved Tree Species in Central European Mixed Forests. *Ecosystems* **18**: 560–
572.

Zuidema PA, Poulter B, Frank DC. **2018**. A Wood Biology Agenda to Support Global Vegetation Modelling. *Trends*
in Plant Science **23**: 1006–1015.

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 Walter-Lieth climatograms for the four study sites: Trentino, Lazio, Campania, and Calabria (from north to south, left to right) for the period 1965-2014. The lower blue bars show the frost period.

Fig. S2 Climate trends at the four sites TRE, LAZ, CAM and CAL (from north to south) for the period 1965 – 2014: Mean annual a) precipitation; b) minimum temperature; c) mean temperature; d) maximum temperature and e) vapour pressure deficit (VPD). Linear regression lines are also indicated.

Fig. S3 Standardized 3-6-9-12-and 24 months SPEI at the four study sites (TRE, LAZ, CAM and CAL) for the 1965–2014 period. Negative (red) and positive (blue) values indicate drier and wetter conditions, respectively.

Fig. S4 Pearson's running correlations between $\delta^{13}C$ with monthly precipitation, mean temperature, VPD, and SPEI1 for the current and the previous year (*), over the period 1965-2014 at each site. The *y-axis* represents the time window in months. Colours (see the key) represent correlation coefficients that are significant at the level of $r = 0.279$ ($P < 0.05$)

Table S1 Soil type and the percentage composition of clay, silt, sand and soil water holding capacity (SWHC) at one meter depth for each site.

Table S2 Dendrochronological statistics (means ± SE) of the tree-ring width series.

Table S3 Tukey post-hoc test for TRW and age in the study sites (one-way analysis of variance with Tukey's Honest Significant Difference test). Different letters indicate significant differences ($P < 0.05$).

Table S4 Statistic of the generalized additive mixed models for the basal area increment (BAI) trends of *Fagus sylvatica* for each site.