Simultaneous Profiling of Chromatin Accessibility and DNA Methylation in Complete Plant Genomes Using Long-Read Sequencing

Basile Leduque[1], Alejandro Edera[1], Clementine Vitte[2], and Leandro Quadrana[1*]

[1] Institute of Plant Sciences Paris-Saclay, Centre Nationale de la Recherche Scientifique, Institute National de la Recherche Agronomique, Université Evry, Université Paris-Saclay, Orsay, France

[2] Université Paris-Saclay, INRAE, CNRS, AgroParisTech, GQE – Le Moulon, Gif-sur-Yvette, France.

[*] correspondance: leandro.quadrana@universite-paris-saclay.fr

ABSTRACT

Epigenetic regulations, including chromatin accessibility, nucleosome positioning, and DNA methylation intricately shape genome function. However, current chromatin profiling techniques relying on short-read sequencing technologies face limitations in adequately characterising repetitive genomic regions and detecting multiple chromatin features simultaneously. Here, we present Simultaneous Accessibility and DNA Methylation Sequencing (SAM-seq), a robust method leveraging bacterial adenine methyltransferases (m6A-MTases) to label accessible regions in purified plant nuclei. Coupled with Oxford Nanopore Technology sequencing, SAM-seq enables high-resolution profiling of m6A-tagged chromatin accessibility together with cytosine methylation along chromatin fibres in plants. Analysis of naked genomic DNA revealed significant sequence preference biases of m6A-MTases, controllable through a normalisation step. By applying SAM-seq to Arabidopsis and maize nuclei we obtained fine-grained accessibility and DNA methylation landscapes at genome-wide and local scales. We characterised crosstalk between chromatin accessibility and DNA methylation, notably within nucleosomes of genes, TEs, and centromeric repeats. SAM-seq also facilitated the identification of DNA footprints over cis-regulatory regions. Furthermore, using the single-molecule information provided by SAM-seq we unveiled extensive cellular heterogeneity at chromatin domains harbouring antagonistic chromatin marks, suggesting that bivalency reflects cell-specific regulations of gene activity. In summary, we introduce a robust method for acquiring high-resolution accessibility and DNA methylation landscapes across entire plant genomes. Our results underscore the importance of considering the intrinsic substrate preferences of m6A-MTases for reliable chromatin profiling. SAM-seq opens new opportunities to simultaneously study multiple epigenetic features at unprecedented scale, enabling the investigation of non-model species with limited genomic and epigenomic information.

Keywords: chromatin, epigenomics, accessibility, DNA methylation, nanopore sequencing, transposable elements, nucleosomes, gene regulation, plant genomes

BACKGROUND

Chromatin is a highly dynamic structure that plays a pivotal role in the regulation of eukaryotic gene expression. It is primarily structured as arrays of nucleosomes, which are separated by DNA segments of variable size often containing regulatory sequences. DNA wrapped into nucleosomes is generally inaccessible to other proteins. Thus, changes in the density and localization of nucleosome arrays provide a layer of epigenetic regulation [1]. DNA itself can also carry epigenetic modifications, notably in the form of cytosine methylation (5mC), which is a repressive mark associated with the silencing of transposable elements (TEs) and genes in both plants and mammals [2]. How nucleosome organisation and DNA methylation affect each other and the regulation of gene activity is still not fully understood. This lack of knowledge is partly due to the inability of current technologies to simultaneously chart both DNA methylation and chromatin accessibility across strings of nucleosomes [3]. Indeed, techniques such as MNase-seq dedicated to map nucleosome positioning, ATAC-seq to chromatin accessibility, and BS-seq to DNA methylation, rely on short-read sequencing technologies spanning fragments of less than one nucleosome in size. Furthermore, due to their short length (<150 nt-long), these sequencing reads cannot be aligned unambiguously over highly repetitive sequences, such as centromeres, telomeres, and high copy TEs, thus providing an incomplete characterization of chromatin regulation. Long-read sequencing technologies, such as Oxford Nanopore Technology (ONT), have overcome this limitation by reading long DNA molecules while simultaneously detecting base modifications. The ability of ONT to sequence native DNA also avoids amplification biases, leading to improved estimation of DNA modifications [4]. Recent works have exploited these characteristics to study DNA methylation together with chromatin accessibility in different species [5–8]. For instance, it is possible to treat chromatin with bacterial m6A methyltransferases (m6A-MTases), which preferentially methylate accessible adenines due to steric hindrance. Because of the lack of endogenous m6A in most eukaryotes, treatment of chromatin with m6A-MTases followed by ONT sequencing could provide a high-resolution snapshot of chromatin accessibility, as was demonstrated in yeast [9]. More recently, targeted analysis of centromeres, telomeres, and rDNAs in the model plant species *Arabidopsis thaliana* has been reported [10], opening the possibility to implement m6A-MTases-based approaches to study chromatin in plants. However, broader use of this method for simultaneous analysis of

chromatin accessibility and DNA methylation on a genome-wide scale in Arabidopsis and other plant species is yet to be explored. Moreover, its potential to unravel the intricacies of large and complex genomes, such as those of crops, remains to be validated. Crucially, DNA modifying enzymes are known to have sequence preferences, which must be taken into consideration when used to assess chromatin landscapes [11,12], even more importantly when tracking chromatin modifications at the genome-wide scale. Whether M6A-MTases have sizable sequence preferences remains elusive, and further characterizations need to be performed to ascertain the validity of results obtained with these approaches [5,9,10].

Here, we introduce "Simultaneous Accessibility and DNA Methylation Sequencing (SAM-seq)" a method enabling to jointly characterise chromatin accessibility and DNA methylation patterns in plant genomes. We found that m6A-MTases have significant DNA substrate preferences, leading to sequence- and strand-biased m6A levels. We show that normalisation of m6A levels is critical to obtain reliable results, especially for the analysis of highly repetitive sequences such as Arabidopsis centromeric repeats. By analysing Arabidopsis and maize nuclei, we demonstrate that SAM-seq can be used to identify large scale chromatin compartments, discriminate active and repressed regions, obtain high-resolution profile of nucleosome positioning and DNA methylation, identify DNA footprints at occupied transcription factors binding sites, and map chromatin landscapes across lowly and highly repetitive sequences.

RESULTS

Simultaneous detection of m6A and 5mC in genomic DNA

Genome-wide simultaneous detection of m6A and 5mC using ONT sequencing has been successfully obtained in mammalian genomes [5,9]. To test whether this is also feasible in plants, we incubated Arabidopsis genomic DNA (gDNA) in a buffer containing the m6A-MTases EcoGII and the methyl donor S-adenosylmethionine (SAM) (Supplementary Figure 1), followed by ONT sequencing (Figure 1A). Genomic DNA (gDNA) without any treatment was also sequenced as control. We obtained long reads with a N50 of 5000 nt, with the longest reads reaching 200,000 nt. We processed these reads using two DNA modification callers, Remora and DeepSignalPlant [13,14], to detect 6mA and 5mC modifications, respectively (see Methods). We estimated 2% and 65% m6A methylation levels in control

and EcoGII-treated gDNA, respectively, indicating low false positive calls and extensive EcoGII methylation. To test whether the presence of 5mC can affect EcoGII activity and/or m6A calling, we assessed m6A levels on 20-bp sequences extracted genome-wide having variable levels of 5mC methylation. Overall, we found little dependency between 5mC and m6A levels (Supplementary Figure 2).

We investigated the chromosome-wide profile of m6A levels of EcoGII-treated gDNA. Because no DNA-associated proteins are present in this *in vitro* experiment, m6A levels are expected to be uniform across the genome. Nonetheless, we found uneven levels of m6A along chromosomes, together with strong strand-biased m6A methylation (Figure 1B). Strand-biased m6A levels were most notable at centromeres, which are made of highly repetitive centromeric satellites. To test whether these biases may be due to m6A-MTase sequence preferences, we modelled the m6A probability as a function of sequence composition, by calculating the average m6A level from EcoGII-treated gDNA across all potential 12-mer in the Arabidopsis genome. Dimensionality reduction analysis of the 12-mers space revealed several clusters of k-mers with high and low m6A levels, suggesting complex dependencies between nucleotide content and m6A (Figure 1C). Sequence biases were also evident over longer k-mers (Supplementary Figure 3). In addition, we observed strand-specific m6A levels in association with sequence content (Supplementary Figure 4). In combination, these results demonstrate the existence of significant m6A-MTase sequence preferences, likely underlying the uneven and strand-biased levels of m6A along chromosomes. Based on the 12-mers model, we generated a genome-wide map of m6A methylation likelihood as normalisation factor. After kmer-based normalisation, m6A levels were largely uniform along chromosomes (Figure 1B), indicating that the strand-specific m6A levels observed at centromeric repeats were largely driven by m6A-MTase sequence preference. In combination, our results demonstrate that m6A-MTase treatment combined with ONT sequencing provides a powerful method to detect simultaneously m6A and 5mC on long DNA molecules from plants, and underscore the importance of considering inherent biases in m6A-MTases activities to obtain reliable m6A profiles.

SAM-seq probes simultaneously chromatin accessibility and DNA methylation on plant tissues

To establish a robust protocol to probe chromatin accessibility in plant tissues using m6A-MTases, we purified, cross-linked, and permeabilized nuclei from two weeks old Arabidopsis seedlings (Figure 1A). m6A-MTases, such as EcoGII or Hia5, are known to be inhibited by detergents such as Triton or SDS [15], which are commonly used for plant nuclear extraction and permeabilization. In animal cells, these detergents can be replaced by digitonin, which does not affect m6A-MTases activity and interacts specifically with cholesterol [15], which nonetheless is absent in plants. We therefore tested the capacity of different combinations of soft detergents and washes to permeabilize Arabidopsis nuclei as well as their effect on m6A-MTases activity. Methyltransferase activity was tested by incubating gDNA extracted from treated chromatin with the restriction enzyme DpnI, which cuts GATC sites only in the presence of m6A (Supplementary Figure 1). We found that nuclei permeabilization with 1% Triton X-100 followed by two washing steps with 0.2% Tween-20 provides optimal m6A methylation activity (Supplementary Figure 5).

To investigate the performance of our SAM-seq protocol to chart chromatin accessibility and DNA methylation genome-wide, we performed ONT sequencing of gDNA extracted from EcoGII-treated nuclei. The 5mC levels measured on gDNA extracted from EcoGII treated nuclei were highly consistent with those obtained from gDNA (Pearson correlations ranging from 0.84 to 0.95, Figure 1D), demonstrating efficient DNA methylation profiling by SAM-seq in the three cytosine contexts. Visualisation of normalised m6A levels revealed two broad chromosome compartments with high and low m6A-MTase accessibility, corresponding respectively to chromosome arms and heterochromatin centromeric and pericentromeric regions (Figure 1E). Notably, normalised m6A profiles showed little strand-biased m6A levels over centromeric regions, compared to non-normalized landscapes, highlighting the importance of the normalisation strategy for reliable accessibility profiling.

We next explored how incubation time and concentration of m6A-MTases affect SAM-seq accessibility results. We found that increasing incubation times improves profiling resolution over well-positioned nucleosomes (Supplementary Figure 6), at the cost of decreasing selectivity towards accessible chromatin regions (ACRs) upstream of the

transcriptional start site (TSS) of genes. These results are consistent with previous observations made in animal cells (Stergachis et al 2020). To assess the robustness of our SAM-seq protocol, we compared the accessibility profiles obtained by SAM-seq with that obtained by ATAC-seq based on short-read sequencing [12]. Overall, we found very good agreement between both techniques (Figure 1F), demonstrating that SAM-seq is a reliable method to chart chromatin accessibility genome-wide. Similar results were obtained across independent SAM-seq experiments using EcoGII or another m6A-MTase Hia5 (Figure 1F), demonstrating reproducibility of SAM-seq experiments.

Accessibility and DNA methylation landscapes of chromatin states

To test whether SAM-seq can provide information about gene activity, we assessed normalised m6A levels around TSS of genes with contrasting expression levels. Accessibility increases in relation to gene expression, with lowly and highly expressed genes showing the lowest and highest m6A-MTase accessibility, respectively (Supplementary Figure 7). These results indicate that SAM-seq-based chromatin accessibility is a reliable proxy for gene activity. To explore further the ability of SAM-seq to chart chromatin landscapes, we plot m6A-MTases accessibility and DNA methylation over the nine chromatin states (CS) previously identified in Arabidopsis based on 16 genomic and chromatin features [16] (Figure 2A). These CSs distinguish promoters (CS1 and 2), upstream regulatory sequences (CS4), active (CS3) and inactive genes (CS6 and 7), intergenic sequences (CS5), and heterochromatic sequences (CS8 and 9). As expected, SAM-seq accessibility was high at active promoters (CS1), while the body of inactive genes and TEs have the lowest accessibility (CS7 and CS9) (Figure 2A and B). Overall, chromatin accessibility correlates negatively with DNA methylation, consistent with a role for this modification in limiting chromatin accessibility [17]. Nonetheless, CS8, which is enriched on TEs and characterised by the presence of DNA methylation and the PRC2-dependent facultative heterochromatin mark H3K27me3 [16], shows some level of accessibility. Interestingly, unlike CS9, which display the lowest accessibility and highest 5mC levels in the three sequence contexts, CS8 has high mCHH but reduced mCG and mCHG (Figure 2A), suggesting that the latter may be targets of the siRNA-directed DNA methylation pathway (RdDM). Simultaneous targeting of these loci by RdDM and PRC2 is consistent with recent studies demonstrating in vivo

interaction of MORPHEUS' MOLECULE1 (MOM1) and AIPP3 factors [18,19], which are respectively involved in RdDM-mediated methylation and H3K27me3 recognition. In combination, these results provide evidence that CHH methylation, in combination with H3K27me3, might constitute a form of relaxed heterochromatin.

To gain insights into the association between chromatin states and cellular heterogeneity, we took advantage of the ability of SAM-seq to probe accessibility across single molecules (Figure 2B). We found that broadly inactive genes (CS7) and silenced TEs (CS9) show the lowest levels of intermolecular m6A heterogeneity (Figure 2C and Supplementary Figure 8), indicating homogeneous chromatin accessibility across cells. Conversely, and despite their overall high accessibility, CS2 and to some extent CS4, which are characterised by bivalent marking of the antagonistic histone modifications H3K4me2 and H3K27me3 [16], display the highest 6mA entropy, with some loci showing on/off accessibility patterns across individual fibres (Figure 2C and 2D). The higher accessibility entropy at these loci provides strong evidence that co-marking reflects extensive cellular heterogeneity, rather than uniform chromatin states across fibres. Hence, CS2 and CS4 more likely reflect cellular-specific epigenetic regulations. Altogether, these results illustrate the power of SAM-seq to investigate epigenetic regulations within and across molecules, genomewide.

Uncovering DNA footprints at cis-regulatory elements

Accessibility assays can be used to uncover DNA footprints [20], sequences that are locally protected by the binding of chromatin associated proteins such as transcription factors (TFs). Due to the high frequency of Adenines in genomes, at almost one in every two DNA base pairs, the high resolution provided by m6A-MTase is ideal to assess short-scale changes in accessibility, such as those associated with DNA footprints. To determine whether m6A-MTase can outline TF-binding sites (TFBS), we explored SAM-seq profiles around 390 putative TFBSs identified by *in vitro* assays [21]. Overall, we detected higher SAM-seq accessibility in the vicinity of 90 of these TFBS motifs (Supplementary Figure 9), consistent with numerous functional regulatory sequences residing within active chromatin regions. Furthermore, we found that SAM-seq accessibility drops at TFBSs were evident for at least 40% of these TFs, suggesting the presence of DNA footprints (Supplementary Figure 9).

To evaluate the potential of SAM-seq to pinpoint occupied TF binding sites *in vivo*, we focused on ABA INSENSITIVE 5 (ABI5), which is a basic leucine zipper (bZIP)-type TF involved in the post-germination response to abscisic acid. Clustering of predicted ABI5 binding sites based on SAM-seq accessibility yielded three types of sites with strong, intermediate, or no protection footprint (Figure 3A). To test the significance of the clustering, we analysed *in vivo* occupancy of ABI5 determined by ChIP-seq [21]. Sites with strong and low accessibility protection were highly and lowly overrepresented with ABI5 ChIP-seq peaks, respectively (Figure 3B). Furthermore, the intensities of ChIP-seq peaks were much higher at sites with accessibility footprints than those lacking such footprints (Figure 3C). Altogether, these results demonstrate the power of SAM-seq to identify occupied TFBS sites *in vivo*.

Fine-grained accessibility and DNA methylation of nucleosomes

Accessibility assays have the potential to provide information on nucleosomal positioning. To test whether SAM-seq can provide such information in plants, we analysed normalised m6A levels around well positioned nucleosomes (see methods). Because accessibility over nucleosomes may differ between euchromatin and heterochromatin, we distinguished between well-positioned nucleosomes at genes, TEs, and centromeric repeats. Across all nucleosome types, we observed the lowest m6A levels within core nucleosomal regions (Figure 4A). These levels gradually increased, peaking at the linker DNA situated approximately 75 base pairs upstream and downstream of the nucleosomal centres (*i.e.* dyads). Nucleosomal-scale SAM-seq protection extended several hundred base pairs in both directions from the focal nucleosomal midpoint, mirroring the MNase nucleosomal signal. In addition to this overall pattern, we uncovered that dyads of centromeric nucleosomes show significant SAM-seq signal (Figure 4B), suggesting that CENH3-containing nucleosomes have unique accessibility patterns. Structural and biochemical studies have revealed that DNA at the entry/exit sites of centromeric nucleosomes is highly flexible and does not cross above the dyad, thereby impeding the binding of linker histone H1 [22,23]. Our SAM-seq results suggest that high flexible DNA ends of nucleosomes provide higher accessibility at dyads, with potential implications for the epigenetic regulation of centromeres.

We next investigated how nucleosome accessibility affects DNA methylation. Nucleosomal DNA over genes has higher levels of CG methylation (Figure 4A). This result is consistent with previous observations [24], and suggest that MET1 activity is not impaired by euchromatic nucleosomes. Conversely, linker DNA at TEs and centromeres show the highest methylation levels compared to nucleosomal DNA, supporting the notion that heterochromatin nucleosomes are strong barriers to DNA methyltransferases [24]. In addition, methylation profiles also reveal that dyads of centromeric nucleosomes have high CG methylation, and that this methylation follows the nucleosomal accessibility landscape (Figure 4B). This result suggests that relaxed DNA wrapping of centromeric nucleosomes promotes the accessibility of MET1 at dyads, leading to high levels of CG methylation. The unusual accessibility and DNA methylation landscapes of centromeric nucleosomes revealed here may have important implications for centromeric function and chromosomes segregation.

Chromatin profiling of Arabidopsis repeated sequences

Because long-reads can be unambiguously aligned over highly repetitive sequences, we explored the ability of SAM-seq to simultaneously chart chromatin accessibility and DNA methylation over such sequences, including *CEN180* centromeric satellites. Consistent with their role in kinetochore assembly, centromeric repeats show the lowest accessibility compared to euchromatic regions (Figure 1E). Recent targeted m6A-MTase-based profiling of centromeres reported intriguing accessibility patterns of *CEN180* repeats, including higher accessibility of *CEN180* compared to surrounding *ATHILA5* retrotransposons, strand-biased accessibility, and reduced accessibility of linker DNA of centromeric nucleosomes [10]. We reasoned that such patterns could be influenced, at least in part, by the sizable sequence preferences displayed by m6A-MTases (Figure 1). Indeed, strand-biased m6A levels of *CEN180* repeats are evident on EcoGII-treated (naked) gDNA, and largely disappeared after controlling for m6A-MTase preference (Figure 1C and Supplementary Figure 10). Lower SAM-seq signal of *ATHILA5* retrotransposons compared to surrounding *CEN180* repeats also vanished after controlling for m6A-MTase sequence preference (Supplementary Figure 11). Furthermore, our SAM-seq profiles revealed higher accessibility of linker DNA (Figure 4). These results underscore the importance of considering enzymatic preferences for chromatin

profiling, and provide evidence that accessibility patterns of Arabidopsis centromeres previously reported [10] can be largely explained by technical artefacts.

We next explored the accessibility and DNA methylation across TEs in the Arabidopsis genome. Overall, SAM-seq accessibility and DNA methylation correlate negatively (Figure 5A and B), in line with previous studies showing a dampening role of DNA methylation in TE accessibility [17]. To explore further this relationship, we clustered TE sequences based on accessibility profiles. This unsupervised model identified six clusters of TEs (Figure 5A). Most clusters, except C1 and 3, contained lowly accessible TEs surrounded by regions with variable degrees of accessibility. As expected, the less accessible cluster, C6, has the highest levels of DNA methylation in all sequence contexts, is highly enriched on TEs belonging to *Ty3/Gypsy* families (Figure 5C), and are located far from genes (Figure 5C). Clusters 1 and 3, in contrast, have high accessibility and reduced DNA methylation levels. TEs in these clusters belong to Helitrons, SINEs, LINEs and Pogo, and locate nearby genes, suggesting that they might serve as regulatory sequences. Potentially regulatory TEs have been recently identified based on dozens of chromatin marks [25]. Comparison between our clusters of TEs defined solely on SAM-seq results with those obtained previously [25] confirmed that C1 and C3 largely span TEs enriched in chromatin marks associated with regulatory functions as well as TFBS (Supplementary Figure 12). In combination, these results demonstrate that SAM-seq can be used to characterise chromatin states at repetitive sequences and to pinpoint TE-derived sequences with potential regulatory functions.

SAM-seq unlocks high-resolution chromatin profiling of complex genomes

With few exceptions, notably Arabidopsis, plant genomes carry a high load of repeats, making it very challenging to profile chromatin landscapes using short-read sequencing technologies. The long read-based approach employed by SAM-seq therefore provides an unprecedented opportunity to investigate the chromatin of large and complex plant genomes. We adapted SAM-seq to investigate chromatin accessibility and DNA methylation in the 2.1Gbp-long maize genome (see Methods), 80% of which corresponds to TE and other repeated sequences [26]. We performed SAM-seq on maize leaves of the reference line B73

and sequenced using a PromethION flow cell, yielding 52 million reads. Unambiguously mapped SAM-seq reads covered 96% of the reference maize genome, compared to only 50-64% typically obtained with short-read sequencing. The significant increase in horizontal coverage demonstrates the definitive advantage of using SAM-seq to investigate the epigenome landscapes of highly-repetitive plant genomes.

Overall m6A level was ~3%, which is much lower than that obtained for Arabidopsis. This result is consistent with previous observations indicating that most of the maize genome is largely inaccessible [27]. Notwithstanding the overall low accessibility, chromosome-wide m6A levels were punctuated by narrow peaks of SAM-seq accessibility (Figure 6A and B). These peaks were often located near the TSS of genes (Figure 6B), and metaplots at genes recapitulated the patterns obtained by ATAC-seq and DNAseq [28,29](Figure 6A), consistent with these sites being ACRs. Accessibility profiles at genes TSS of another maize inbred line, M53, were similar to the ones obtained in B73 (Supplementary Figure 13), further supporting the reproducibility of SAM-seq to investigate maize chromatin. We then took advantage of the large maize genome, where ACRs can be located away from genes, to further test the correspondence between SAM-seq and ATAC-seq in B73. For this, we evaluated m6A levels at high-confidence proximal (<2kb from TSS) and distal (>2kb from TSS) ACRs previously defined by ATAC-seq [28]. This analysis showed that SAM-seq accessibility was the highest over ACRs, independently of their distance to genes (Figure 6C), thus highlighting that this method can inform on chromatin accessibility outside of genes, a feature particularly important for large plant genomes [30]. In maize, ACRs have been found to be enriched in TFBSs [28]. Consistent with these observations, SAM-seq accessibility increases around binding sites identified by DAP-seq for 32 TFs [21,31] (Supplementary Figure 14). Last, we explored the SAM-seq accessibility of genes displaying different levels of expression [32] (Figure 6D). This analysis revealed a good correlation between SAM-seq accessibility at TSS of genes and expression levels. Altogether, these results demonstrate that SAM-seq is scalable to large genomes, enables the identification of ACRs at promoters, proximal and distal regions, and provides a reliable proxy for gene activity *in vivo* for complex plant genomes.

Given the high TE-content of the maize genome, we tested the ability of SAM-seq to chart accessibility and DNA methylation over TEs. Overall, we found that maize TEs are poorly accessible and associated with high mCG and mCHG levels, supporting previous observations based on limited genome coverage obtained through short-read sequencing technologies [33]. Clustering of TE sequences based on SAM-seq chromatin accessibility identified seven clusters (Figure 6E). The largest cluster (C7) gathers TEs with, and embedded in regions of, low accessibility, high mCG and mCHG, and low mCHH levels. These types of inaccessible, highly methylated TEs, are typically located far away from genes and associated with Harbingers, RTE, Helitrons and unknown LTR and TIR superfamilies (Figure 6F). Conversely, upstream and downstream regions respectively flanking TEs in C2 and 3, and to a lower extent C5 and 6, show relatively high SAM-seq accessibility and low CG and CHG methylation. Notably, these TEs have high mCHH levels, which peaked towards TE's border facing the accessible flankings. Such a pattern is reminiscent of "CHH islands", which has been described as narrow clusters of mCHH typically located nearby genes and proposed to operate as epigenetic boundaries between euchromatin and heterochromatin in maize [34]. However, unlike previous characterizations of CHH islands, which reported low mCG and mCHG [34], TEs in C2 and C3, are highly methylated in the three sequence contexts. Last, Cluster 1, and to a lower extent C4 too, gathers regions with high chromatin accessibility both within TEs and at flanking regions (Figure 6E). TEs from this cluster have particularly high mCHH, reduced mCG and mCHG levels (Figure 6E), typically belong to *hAT*, *Mutator*, and *PIF/Harbinger* superfamilies, and are located close to genes (Figures 6F and G). Altogether, these results indicate that maize TEs preferentially methylated in the CHH context are not linked to reduced accessibility nor to euchromatin/heterochromatin boundaries. Conversely, these TEs may be playing a role as *cis*-regulatory elements of nearby genes, as has been recently proposed for the emergence of distal regulatory sequences across angiosperm species [30].

DISCUSSION

The intricate interplay between nucleosome positioning, DNA methylation, and chromatin accessibility is fundamental for genome regulation in eukaryotic organisms. However, addressing these aspects has been so far challenging due to the technological limitations imposed by the use of short-read sequencing methods. This challenge is

particularly critical when studying highly repetitive sequences, which are pervasive components of most plant genomes. We developed a method, named SAM-seq, to simultaneously map DNA methylation and chromatin accessibility in plant genomes that takes advantage of m6A-MTases and long-read sequencing to 'stencil' accessible DNA. We demonstrate that this method can be applied to obtain fine-grained chromatin landscapes in species with widely variable genome sizes and repetitiveness. Moreover, SAM-seq's technical simplicity, reproducibility, and cost-effectiveness make it an ideal approach for investigating chromatin marks and their dynamics in plants with large genomes.

The dynamic range and power of SAM-seq are comparable between Arabidopsis and maize, illustrating its versatility across species and genome complexities. This versatility would enable the characterization of chromatin landscapes in virtually any plant species, providing unprecedented insights into chromatin regulation, gene activity, and nucleosome positioning. Furthermore, given the recent improvements in the quality and quantity of ONT sequencing outputs, SAM-seq offers an all-in-one method to generate *de novo* genome assemblies together with detailed epigenome information. This will significantly facilitate the investigation of chromatin-based regulations across the tree of life, bridging the gap between model and non-model species.

One significant limitation in investigating chromatin accessibility has been the inherent sequence biases of DNA modifying enzymes, such as those previously observed for DNaseI and Tn5 retrotransposons [11,12]. Here, by subjecting genomic (naked) DNA to m6A-MTases, we uncovered significant sequence preferences. We also show that these preferences can generate spurious accessibility landscapes. For instance, m6A-MTase sequence preferences can lead to significant strand-specific accessibility profiles, especially over highly repetitive sequences. Previous work using EcoGII m6A-MTase and Pacbio long-read sequencing reported only modest differences in methylation levels among k-mers [9]. These prior findings may have resulted from an inaccurate characterization of m6A-MTase preferences. Indeed, significance of biases increases with K-mers size [9], and we uncovered strong biases when considering 12-mers and 20-mers. Hence, m6A-MTase activity might be likely influenced by regional DNA composition, rather than local sequence motifs. Given the rapid growth in the number of methods using m6A-MTases to investigate chromatin,

including recent developments of antibody-driven targeting of m6A-MTases to map DNA binding proteins [15,35], we underscore the importance of measuring and correcting inherent biases introduced by m6A-MTases preferences.

Chromatin accessibility is commonly assessed using ATAC-seq, which is based on short-read sequencing. However, detecting DNA footprints and nucleosome positioning using this technique remains challenging. We demonstrate that SAM-seq is a multipurpose chromatin profiling technique capable of capturing a wide spectrum of chromatin nuances, from subtle, small-scale alterations to large-scale changes. This wide-resolution profiling enabled us to uncover striking variations in accessibility and DNA methylation along the nucleosome surface. In particular, we show that dyads of Arabidopsis centromeric nucleosomes are highly accessible, potentially facilitating the activity of the CG-specific DNA methyltransferase MET1. This atypical accessibility and DNA methylation nucleosome landscape may have important implications in the epigenetic regulation of centromeres and chromosomal segregation. In addition to mapping nucleosomes, the high-resolution of SAM-seq enables the detection of TFBS occupied in vivo. We found that at least one third of putative binding sites for the Arabidopsis TF ABI5 reside within open chromatin and show DNA footprints (Figure 4). Similar picture was obtained in maize, indicating that SAM-seq can be used to investigate cis-regulatory regions in different plant species.

Clustering of TEs based on SAM-seq accessibility allowed us to distinguish between highly heterochromatin TEs, TEs located in heterochromatin/euchromatin boundaries between , TEs embedded in euchromatin, and bonafide euchromatic TEs. Remarkably, unlike Arabidopsis where DNA methylation in the three cytosine contexts correlate negatively with accessibility, bonafide euchromatic (i.e. accessible) TEs in maize show the highest mCHH levels. Given that maize lacks the mCHH-specific DNA methyltransferase CMT2, our results provide strong evidence supporting that accessible TEs are the preferred targets of RdDM [29]. Furthermore, we found that mCHH and accessibility is higher at the edge of these TEs, and this pattern is mirrored by mCG and mCHG. This result suggests that RdDM may establish a transition zone between heterochromatin and euchromatin, preventing the spreading of DNA methylation over genes as well as preventing accidental PolII read-through transcription from nearby genes. In combination, the high-resolution accessibility and DNA

methylation landscapes of SAM-seq offers a powerful toolkit to study the interplay between nucleosomes, chromatin accessibility, and DNA methylation at unprecedented scale.

Conclusion: SAM-seq is a transformative tool that facilitates the investigation of nucleosome organisation, DNA methylation, and chromatin in small as well as large and complex genomes. Its technical simplicity and ability to simultaneously characterise multiple chromatin features, ranging from small-scale changes to chromosomal-scale patterns, enables the investigation of chromatin-based regulations in virtually any species.

METHODS

Plant material

Arabidopsis thaliana seeds (Col-0 accession) were germinated and grown in long-days (16h:8h light:dark) conditions at 23°C for approximately three weeks. Rosette leaves were harvested, flash frozen in liquid nitrogen, and stored at -80°C until use. Maize seeds (B73_USA SMH2013 accession) were germinated and grown under long days conditions (16:8 light:dark) at 25°C:20°C (light:dark) during seven days and harvested 4–6 h after Zeitgeber Time 0 (ZT0, lights on). The 2nd leaf containing inner leaves was flash frozen in liquid nitrogen, and stored at -80°C until use.

Adenine methylation of gDNA using EcoGII.

Methylation of 1µg of gDNA was performed at 37°C during 1h in the presence of 0.1 units/µl of EcoGII, in rCutSmart buffer in presence of 160 µM SAM, followed by heat inactivation at 65°C during 10 minutes. Methylation efficiency was measured by qPCR after DNA digestion with the m6A-specific restriction enzyme DpnI. 1µg of DNA was used to prepare ONT libraries using LSK110 kit following manufacture indications. Library was loaded in a FLO-MIN106D and runned in a MK1C device during 72hs.

SAM-seq protocol

For Arabidopsis SAM-seq experiments, 250mg of seedlings were ground under liquid nitrogen and resuspended in 25 ml of extraction buffer 1 (sucrose 0.4M, Tris-HCl pH8 10mM, MgCl2 10mM, ethidium bromide 5mM). Tissues are fixed with 1% formaldehyde for

5 minutes in ice with gentle mixing, followed by the addition of 1.25 M glycine to twice the molar concentration of formaldehyde, materials sit in ice for 5 minutes before centrifugation at 4000 g for 20 minutes at 4°C. For nuclear isolation the pellet was resuspended in 1 ml Extraction Buffer 2 (sucrose 0.25M, Tris-HCl pH8 10mM, MgCl2 10mM, ethidium bromide 5mM, Triton x-100 1%, protease inhibitor 1X) and centrifuged at 4 °C for 10 min at 11000g. The pellet was then resuspended in 0.3ml Extraction Buffer 3 (sucrose 1.7M, Tris-HCl pH8 10mM, MgCl2 2mM, ethidium bromide 5mM, Triton x-100 0.15%, protease inhibitor 1X), transferred to a clean tube containing 0.3ml Extraction Buffer 3 and centrifuged at 4 °C for 30 minutes at 11,000g. To optimise MTase activity, two consecutive washes with 1ml of nuclei preparation buffer (MOPS pH7 20mM,NaCl 40mM, KCl 90mM, EDTA 2mM, EGTA 0.5mM, spermidine 0.5mM, spermine 0.2 mM, protease inhibitor 1X) and 1 ml of plant tween wash buffer (Tween-20 0.2%, NaCl 150mM, HEPES-KOH 20mM, spermidine 0.5mM, protease inhibitor 1X) were performed. The pellet of nuclei was resuspended in 200µl of activation buffer (1X rCutSmart , 800 µM SAM) containing 7.5U of EcoGII MTase. Samples were incubated under rotation at 37°C for 30 minutes. We noted that 30 minutes incubation with 37.5 U/ml of EcoGII provides the best signal/noise ratio to chart accessibility at ACRs and nucleosome profiling. For reverse cross-linking, 1 M NaCl was added to 200µl of 1X cutSmart to reach a salt concentration of 200mM. Samples were incubated at 65 °C for 1 hour for lysis to reverse cross-linking. 0.3mg/ml of RNAse and 0.1mg/ml Proteinase K were added to the nuclei and incubated for 40 minutes, followed by a phenol/chloroform DNA extraction.

For maize SAM-seq experiments, seven days old leaves were ground under liquid nitrogen and 750 mg of powder resuspended in 12.5ml of extraction buffer 1. Nuclei purification, fixation, and m6A-MTases reaction was performed as described above with the following modifications. Fixed nuclei's pellet was resuspended in 38 µl 1X rCutSmart, containing 10µl SAM and 6µl EcoGII (i.e. 7.5U for 100µl) and incubated for 30 minutes at 30°C. For reverse cross-linking, 20 µl NaCl (5M) were added followed by an overnight incubation at 65 °C. DNA extraction was performed as described above.

ONT library preparation and sequencing

For each sample, between 0.5 and 1 µg DNA was used for library preparation using the Ligation Sequencing Kit SQK-LSK109. The library was prepared following the manufacturer's instructions with minor modifications. The end-repair incubation time was increased to 20 min. Ligation incubation times were increased to at least 1h. Elution was always performed in 10 mM Tris-HCl. LFB was used for the final size selection step. DNA yield was quantified using the Qubit dsDNA HS Assay Kit (Q33230). Sequencing was performed on MinION with v9.4.1 flow cells (FLO-MIN106.1) or on PromethION with v9.4.1 flow cells (FLO-PRO002) (Supplementary Data XX).

Detection of m6A and 5mC

m6A base calling and mapping were performed using Megalodon with the Rerio model res_dna_r941_min_modbases-all-context_v001.cfg and default parameters. Adenines with methylated probability greater than 0.75 were considered as m6A. Otherwise, adenines were regarded as unmethylated. For 5mC methylation calling, fast5 files were squiggled with the FASTQ files obtained from Megadolon using Tombo v1.5.1. 5mC methylation was called from the squiggled fast5 files with DeepSignalPlant v0.1.6 [14] using the model model.dp2.CNN.arabnrice2-1_120m_R9.4plus_tem.bn13_sn16.both_bilstm.epoch6.ckpt. Single-molecule cytosine sites with predicted scores greater than 0.5 were considered as methylated. Methylation levels of genomic positions were calculated using the ratio of 5mC/(5mC + C), where C denotes cytosines predicted as unmethylated.

m6A methylation likelihood normalisation

To elucidate the intrinsic sequence preferences of EcoGII, we scanned the Col-CEN assembly to identify all adenine-containing positions, retrieving the 6-base sequences flanking each adenine both upstream and downstream. For each of these 13-mer sequences, the methylation level of the focal adenine was determined based on the average m6A level observed at the corresponding position in the SAM-seq accessibility maps generated from naked gDNA. To obtain normalisation constants to correct for EcoGII preferences, the retrieved flanking sequences with identical 12-mer sequences were grouped together, and the methylation levels of their central adenines were averaged. This process yielded a m6A methylation likelihood for each 12-mer. This value was used to adjust m6A levels obtained

in SAM-seq experiments: Briefly, SAM-seq values for each A genome wide were divided by the likelihood methylation for the corresponding 12-mer and centred to obtain values between 1 and 0.

Analysis of SAM-seq over nucleosomes

Nucleosome positioning was inferred using MNase libraries previously published [24,36–38]. MNaseq datasets were aligned to the Col-CEN reference genome assembly [39] and libraries with high mappability efficiency and median fragment lengths distribution of ~147 bp were retained. Paire-end reads were mapped to the Col-CEN assembly using Bowtie v2.2.5 [40] with the following parameters: -N 1 --end-to-end --no-mixed --no-discordant. The resulting BAM files were sorted with samtools 1.6, and then converted to bigwig files using deeptools v3.5.1 [41] --binSize 1 --normalizeUsing RPGC --extendReads --effectiveGenomeSize 131559676 --ignoreForNormalization chrC chrM. The obtained bigwig files were pooled together to yield a smoothing effect. Next, an in-house script was implemented to perform peak calling using functions offered by the Numpy and Scipy libraries, [42,43]. Briefly, this calling first applied a convolution along the pooled profile using a 1D Gaussian kernel (standard deviation=30), and then calculated the second derivative of the convolved profile to determine local maximas. The mid-point of these maximas were defined as nucleosome dyads. The K-mean algorithm (K=2) based on dynamic time warping distances [44] was employed to clusterized the called dyads into well-positioned and poorly-positioned. Concretely, clustering used as input signals encompassing the 1 kb convolved profile upstream and downstream of each called dyad. The well-positioned dyads were further classified based on their genomic localization into: genes, TEs and CEN180s. Dyad's positions were used to anchor metaplots generated by averaging MNase as well as normalised SAM-seq accessibility, mCG, mCHG, and mCHH levels. For visualisation, SAM-Seq accessibility metaplots were protected onto the cryo-EM nucleosome structure 7by0 [13].

Availability of data and materials: The datasets generated and/or analysed during the current study are available in the European Nucleotide Archive (ENA) repository PRJEBXXXXX. Publicly available Arabidopsis ATACseq, MNaseq, H3K9me2 ChIPseq, DAP-

seq and ABI5 ChIPseq data were obtained from GEO bioprojects GSE85203, GSE96994, GSE139460, GSE190317, GSE207391, GSE60143. Publicly available Maize ATAC-seq and DAP-seq were obtained from GEO bioprojects GSE60143 and GSE120304. All data generated or analysed during this study are included in this published article. Detailed SAM-seq protocols for Arabidopsis and maize are publicly available at dx.doi.org/10.17504/protocols.io.8epv5x1xdg1b/v1 and dx.doi.org/10.17504/protocols.io.kqdg3x25zg25/v1, respectively.

Competing interests: The authors declare that they have no competing interests.

Funding: This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No. 948674 to LQ), and the Agence Nationale de la Recherche (project EpiLinks grant No. ANR-22-CE20-0001 to LQ).

Authors' contributions: LQ conceived the project. BL developed and implemented SAM-seq, performed most bioinformatic analysis with additional inputs from AE, and interpreted the data. AE performed m6A methylation likelihood normalisation and nucleosome analysis. CV generated maize samples and discussed results. LQ interpreted the data and drafted the manuscript, with additional inputs from all the authors. All authors read and approved the paper.

Acknowledgements: We thank members of the Quadrana group for discussions. LQ thanks Vincent Colot for insightful discussions. We thank Nicolas Altemose for sharing the Hia5-pA enzyme.

REFERENCES

1. Henikoff S. Nucleosome destabilization in the epigenetic regulation of gene expression. Nat Rev Genet. 2008;9:15–26.

2. Law JA, Jacobsen SE. Establishing , maintaining and modifying DNA methylation patterns in plants and animals. Nat Rev Genet. 2010;11:204–20.

3. Minnoye L, Marinov GK, Krausgruber T, Pan L, Marand AP, Secchia S, et al. Chromatin accessibility profiling methods. Nat Rev Methods Primers [Internet]. 2021;1. Available from: https://www.nature.com/articles/s43586-020-00008-9

4. Hook PW, Timp W. Beyond assembly: the increasing flexibility of single-molecule sequencing technology. Nat Rev Genet. 2023;24:627–41.

5. Lee I, Razaghi R, Gilpatrick T, Molnar M, Gershman A, Sadowski N, et al. Simultaneous profiling of chromatin accessibility and methylation on human cell lines with nanopore sequencing. Nat Methods. 2020;17:1191–9.

6. Stergachis AB, Debo BM, Haugen E, Churchman LS, Stamatoyannopoulos JA. Single-molecule regulatory architectures captured by chromatin fiber sequencing. Science. 2020;368:1449–54.

7. Battaglia S, Dong K, Wu J, Chen Z, Najm FJ, Zhang Y, et al. Long-range phasing of dynamic, tissue-specific and allele-specific regulatory elements. Nat Genet. 2022;54:1504–13.

8. Wang Y, Wang A, Liu Z, Thurman AL, Powers LS, Zou M, et al. Single-molecule long-read sequencing reveals the chromatin basis of gene expression. Genome Res. 2019;29:1329–42.

9. Shipony Z, Marinov GK, Swaffer MP, Sinnott-Armstrong NA, Skotheim JM, Kundaje A, et al. Long-range single-molecule mapping of chromatin accessibility in eukaryotes. Nat Methods. 2020;17:319–27.

10. Mo W, Shu Y, Liu B, Long Y, Li T, Cao X, et al. Single-molecule targeted accessibility and methylation sequencing of centromeres, telomeres and rDNAs in Arabidopsis. Nat Plants. 2023;9:1439–50.

11. He HH, Meyer CA, Hu SS, Chen M-W, Zang C, Liu Y, et al. Refined DNase-seq protocol and data analysis reveals intrinsic bias in transcription factor footprint identification. Nat Methods. 2014;11:73–8.

12. Lu Z, Hofmeister BT, Vollmers C, DuBois RM, Schmitz RJ. Combining ATAC-seq with nuclei sorting for discovery of cis-regulatory regions in plant genomes. Nucleic Acids Res. 2017;45:e41.

13. Ariyoshi M, Makino F, Watanabe R, Nakagawa R, Kato T, Namba K, et al. Cryo-EM structure of the CENP-A nucleosome in complex with phosphorylated CENP-C. EMBO J. 2021;40:e105671.

14. Ni P, Huang N, Nie F, Zhang J, Zhang Z, Wu B, et al. Genome-wide detection of cytosine methylations in plant from Nanopore data using deep learning. Nat Commun. 2021;12:5976.

15. Altemose N, Maslan A, Smith OK, Sundararajan K, Brown RR, Mishra R, et al. DiMeLo-seq: a long-read, single-molecule method for mapping protein-DNA interactions genome wide. Nat Methods. 2022;19:711–23.

16. Sequeira-Mendes J, Aragüez I, Peiró R, Mendez-Giraldez R, Zhang X, Jacobsen SE, et al. The Functional Topography of the Arabidopsis Genome Is Organized in a Reduced Number of Linear Motifs of Chromatin States. Plant Cell. 2014;26:2351–66.

17. Zhong Z, Feng S, Duttke SH, Potok ME, Zhang Y, Gallego-Bartolomé J, et al. DNA methylation-linked chromatin accessibility affects genomic architecture in. Proc Natl Acad Sci U S A [Internet]. 2021;118. Available from: http://dx.doi.org/10.1073/pnas.2023347118

18. Li Z, Wang M, Zhong Z, Gallego-Bartolomé J, Feng S, Jami-Alahmadi Y, et al. The MOM1 complex recruits the RdDM machinery via MORC6 to establish de novo DNA methylation. Nat Commun. 2023;14:4135.

19. Zhang Y-Z, Yuan J, Zhang L, Chen C, Wang Y, Zhang G, et al. Coupling of H3K27me3 recognition with transcriptional repression through the BAH-PHD-CPL2 complex in Arabidopsis. Nat Commun. 2020;11:6212.

20. Klemm SL, Shipony Z, Greenleaf WJ. Chromatin accessibility and the regulatory epigenome. Nat Rev Genet. 2019;20:207–20.

21. O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, et al. Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. Cell. 2016;166:1598.

22. Takizawa Y, Ho C-H, Tachiwana H, Matsunami H, Kobayashi W, Suzuki M, et al. Cryo-EM Structures of Centromeric Tri-nucleosomes Containing a Central CENP-A Nucleosome. Structure. 2020;28:44–53.e4.

23. Roulland Y, Ouararhni K, Naidenov M, Ramos L, Shuaib M, Syed SH, et al. The Flexible Ends of CENP-A Nucleosome Are Required for Mitotic Fidelity. Mol Cell. 2016;63:674–85.

24. Lyons DB, Zilberman D. DDM1 and lsh remodelers allow methylation of DNA wrapped in nucleosomes. Elife. 2017;6:1–20.

25. Hisanaga T, Romani F, Wu S, Kowar T, Wu Y, Lintermann R, et al. The Polycomb repressive complex 2 deposits H3K27me3 and represses transposable elements in a broad range of eukaryotes. Curr Biol [Internet]. 2023; Available from: http://dx.doi.org/10.1016/j.cub.2023.08.073

26. Jiao Y, Peluso P, Shi J, Liang T, Stitzer MC, Wang B, et al. Improved maize reference genome with single-molecule technologies. Nature. 2017;546:524–7.

27. Rodgers-Melnick E, Vera DL, Bass HW, Buckler ES. Open chromatin reveals the functional maize genome. Proc Natl Acad Sci U S A. 2016;113:E3177–84.

28. Ricci WA, Lu Z, Ji L, Marand AP, Ethridge CL, Murphy NG, et al. Widespread long-range cis-regulatory elements in the maize genome. Nat Plants. 2019;5:1237–49.

29. Gent JI, Madzima TF, Bader R, Kent MR, Zhang X, Stam M, et al. Accessible DNA and relative depletion of H3K9me2 at maize loci undergoing RNA-directed DNA methylation. Plant Cell. 2014;26:4903–17.

30. Lu Z, Marand AP, Ricci WA, Ethridge CL, Zhang X, Schmitz RJ. The prevalence, evolution and chromatin signatures of plant regulatory elements. Nat Plants. 2019;5:1250–9.

31. Galli M, Khakhar A, Lu Z, Chen Z, Sen S, Joshi T, et al. The DNA binding landscape of the maize AUXIN RESPONSE FACTOR family. Nat Commun. 2018;9:4526.

32. Fagny M, Kuijjer ML, Stam M, Joets J, Turc O, Rozière J, et al. Identification of Key Tissue-Specific, Biological Processes by Integrating Enhancer Information in Maize Gene Regulatory Networks. Front Genet. 2020;11:606285.

33. Li Q, Eichten SR, Hermanson PJ, Zaunbrecher VM, Song J, Wendt J, et al. Genetic perturbation of the maize methylome. Plant Cell. 2014;26:4602–16.

34. Li Q, Gent JI, Zynda G, Song J, Makarevitch I, Hirsch CD, et al. RNA-directed DNA methylation enforces boundaries between heterochromatin and euchromatin in the maize genome. Proc Natl Acad Sci U S A. 2015;112:14728–33.

35. Weng Z, Ruan F, Chen W, Chen Z, Xie Y, Luo M, et al. BIND&MODIFY: a long-range method for single-molecule mapping of chromatin modifications in eukaryotes. Genome Biol. 2023;24:61.

36. Luo Y-X, Hou X-M, Zhang C-J, Tan L-M, Shao C-R, Lin R-N, et al. A plant-specific SWR1 chromatin-remodeling complex couples histone H2A.Z deposition with nucleosome sliding. EMBO J. 2020;39:e102008.

37. Shu J, Ding N, Liu J, Cui Y, Chen C. Transcription elongator SPT6L regulates the occupancies of the SWI2/SNF2 chromatin remodelers SYD/BRM and nucleosomes at transcription start sites in Arabidopsis. Nucleic Acids Res. 2022;50:12754–67.

38. Corcoran ET, LeBlanc C, Huang Y-C, Arias Tsang M, Sarkiss A, Hu Y, et al. Systematic histone H4 replacement in Arabidopsis thaliana reveals a role for H4R17 in regulating flowering time. Plant Cell. 2022;34:3611–31.

39. Naish M, Alonge M, Wlodzimierz P, Tock AJ, Abramson BW, Schmücker A, et al. The genetic and epigenetic landscape of the centromeres. Science. 2021;374:eabi7489.

40. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:357–9.

41. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al. deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids Res. 2016;44:W160–5.

42. Harris CR, Millman KJ, van der Walt SJ, Gommers R, Virtanen P, Cournapeau D, et al. Array programming with NumPy. Nature. 2020;585:357–62.

43. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods. 2020;17:261–72.

44. Meert W, Hendrickx K, Van Craenendonck T. wannesm/dtaidistance v2.0.0 [Internet]. Zenodo; 2020. Available from: https://zenodo.org/record/3981067

FIGURES

Figure 1. SAM-seq probes simultaneously chromatin accessibility and DNA methylation on plant tissues. **A**. Workflow of the SAM-seq protocol and assessment of m6A-MTase sequence preferences on gDNA (EcoGII-treated gDNA). **B**. Strand-specific m6A levels of EcoGII-treated genomic DNA before and after normalisation for m6A-MTase preferences. Centromeres are depicted as grey boxes. **C** UMAP projections of Arabidopsis 12-mer sequences coloured by m6A/A levels of EcoGII-treated gDNA, A content, and AT content. **D** Density plots and correlation analysis between ONT sequencing of gDNA and SAM-Seq for mCG, mCHG, and mCHH levels. **E**. Strand-specific SAM-seq m6A levels before and after normalisation for m6A-MTases preferences. **F**. Metaplot of Arabidopsis genes displaying ATAC-seq accessibility (Lu et al. 2017) and SAM-seq chromatin accessibility (m6A/A) obtained from independent experiments using distinct m6A-MTases (EcoGII or Hia5).

Figure 2. Accessibility and DNA methylation landscapes of Arabidopsis chromatin states. **A**. Accessibility, mCG, mCHG, and mCHH levels across nine chromatin states (Sequeira-Mendes *et al.*, 2014). **B**. Genome browser view of SAM-Seq accessibility, mCG, mCHG, and mCHH levels. m6A along reads are indicated in red. **C**. Distribution of SAM-seq accessibility heterogeneity (Shannon entropy) across reads between CS locus. **D**. Genome browser view of SAM-Seq accessibility and m6A sites along reads spanning a CS4 locus with high accessibility entropy. Reads showing high (open), intermediate, and low (close) m6A levels are indicated.

Figure 3. SAM-seq footprinting pinpoints sites bound by TFs *in vivo*. **A**. Metaplot and heatmaps of SAM-Seq chromatin accessibility around putative ABI5 binding site determined in-vitro by DAP-seq. Clustering of chromatin accessibility profiles identified three types of ABI5 bindings sites. **B**. Enrichments of ABI5 Chip-Seq peaks at type 1, 2 or 3 transcription factor binding sites (TFBS). **C**. ABI5 Chip-Seq peaks signal over type 1, 2 or 3 TFBS.

Figure 4. Fine-grained accessibility and DNA methylation of Arabidopsis nucleosomes A. Metaplots showing MNase (grey area), SAM-Seq accessibility (purple line), mCG (red), mCHG (blue) and mCHH (green) over well-positioned nucleosomes in genes, TEs, or centromeres. **B**. Projection of SAM-Seq accessibility metaplots onto the cryo-EM nucleosome structure (7by0). Superhelical locations (SLH) 0, 2, 4, and 6 are indicated.

Figure 5. Chromatin profiling of Arabidopsis TEs. A. Metaplots and heatmaps for SAM-Seq accessibility, mCG, mCHG and mCHH methylation, as well as H3K9me2 from ChIP-seq data, across TEs in the Arabidopsis reference genomes. Clustering of TEs based on SAM-Seq accessibility identified six clusters. **B**. Dotplots showing SAM-Seq accessibility and mCG, mCHG or mCHH for each TE. TEs are coloured based on the clusters obtained in A. **C**. Proportion of TEs belonging to each cluster across TE superfamilies. **D**. Distribution of distances of TEs to the nearest gene across TE clusters.

Figure 6. SAM-seq chromatin profiling of the maize genome. A. Metaplots of chromatin accessibility signal over genes. **B.** Genome browser view of SAM-Seq accessibility, mCG, mCHG, and mCHH levels. m6A along reads is indicated in red and accessible chromatin region (ACR) detected by ATAC-seq from Ricci *et al.*, 2019 are shown. **C.** Metaplots of chromatin accessibility signal over genic, proximal or distal ACRs. **D.** Metaplots and heatmap of chromatin accessibility signal over TSS with different expression levels. **E.** Metaplots and heatmaps for SAM-Seq accessibility, mCG, mCHG and mCHH methylation, across TEs in the reference Maize genomes. Clustering of TEs based on SAM-Seq accessibility identified seven clusters. **F.** Proportion of TEs belonging to each cluster across TE superfamilies. **G.** Distribution of distances between TEs and their nearest gene across TE clusters.