Bacterial assembly in the switchgrass rhizosphere is shaped by phylogeny, host genotype,
and growing site.

Authors: Jeremy Sutherland[1,6], Terrence H. Bell[2], Stacy Bonos[3], Christopher Tkach[3], Julie
Hansen[4], Ryan Crawford[4], John E. Carlson[5], Jesse R. Lasky[6]

1: Department of Plant Science, The Pennsylvania State University, State College, PA, USA

2: Department of Physical and Environmental Sciences, University of Toronto – Scarborough,
Toronto, ON, Canada

3: Department of Plant Biology, Rutgers the State University of New Jersey, New Brunswick,
NJ, USA

4: Plant Breeding and Genetics Section, School of Integrative Plant Science, Cornell University,
Ithaca, NY, USA

5: Department of Ecosystem Science and Management, The Pennsylvania State University, State
College, PA, USA

6: Department of Biology, The Pennsylvania State University, State College, PA, USA

Summary

• Since microbial traits are conserved at different taxonomic levels, plant hosts may influence
microbiome composition differently at different levels to broadly promote or resist microbiota
with traits that impact host fitness. We tested this hypothesis by assessing signals of host
genetic influence on bacterial composition in the switchgrass rhizosphere using 128 genotypes
in dissimilar growing sites.

• We employed three common gardens, combined with host genetic mapping, 16S rRNA gene
sequence analysis, hierarchical modeling, tests of phylogenetic conservation of host influence,
and genome-wide association analyses to determine the contributions of host genetics in
shaping rhizosphere bacterial composition at different taxonomic levels.

• Modeling bacterial assembly showed that growing site was a strong factor shaping bacterial
composition in the rhizosphere, though host genetic influence played a significant role. The
heritability of bacterial abundance was strongest at the genus level. Phylogenetic signal for
heritability was detected within the bacterial phylogeny but conserved clades differed between
common gardens. We identified shared host genetic variants associated with bacterial
abundance and host traits related to plant metabolism.

• Our results suggest further investigation is required regarding the genotype-by-environment-
by-microbiome relationship to elucidate the factors shaping rhizosphere microbiome
composition and the agroecological dynamics shaping plant phenotype.

Introduction

Understanding how host-associated microbiomes assemble is an important issue for basic
and applied community ecology, population biology, and plant breeding. From a community
perspective, microbiome assembly presents a challenge, in part because microbial diversity is often
even more diverse than paragons of community ecology, such as tropical tree communities
(Prosser & Martiny, 2020). A wide range of abiotic constraints and biotic interactions are known
to drive microbiome composition, but the largely unknown role of host physiology, evolution, and
ecology in driving microbiome composition begs for more inquiry at the intersection of population
and host organismal biology (Brunel et al., 2020). Furthermore, the importance of host-microbe
interactions influencing host performance and ecosystem function suggests a high potential payoff
for leveraging knowledge for applied purposes, such as plant domestication (Pérez-Jaramillo et al.,
2016) and agricultural management strategies (Mahmud et al., 2021) that reduce the need for costly
soil inputs.

Environmental effects, ecological drift, and dispersal limitations can influence the
assembly of host-associated microbiomes at different spatial and temporal scales. For example,
abiotic features of the environment can directly shape host-associated microbiome composition
and can be a strong predictor of microbiome composition at macroscales, especially for microbiota
found outside the plant (e.g., the rhizosphere) (Brunel et al., 2020). As a result, the presence and
abundance of microbial taxa that associate with a given host can vary substantially across the host's
range (Serna-Chavez et al., 2013), meaning that different microbial strains, or coarser taxonomic
groups or microbes, may be responsible for conserved interactions with hosts at different locations
(Compant et al., 2010; Mansfield et al., 2012). This variability in host-microbe interactions among
locations complicates our ability to consistently detect signals of host genetic influence on specific
taxa through space (Fierer et al., 2013; Peiffer et al., 2013; Ruhl et al., 2022). To overcome this
limitation, we employed three common garden experiments and genetic mapping to disentangle
host genetic impacts on microbiome composition from site interactions.

Moreover, in community ecology, substantial progress has been made by considering

community assembly in light of the phylogenetic relationships among species (Kembel & Hubbell,

2006). This approach is based on the hypothesis that traits determining species' niches are

phylogenetically conserved (Ackerly, 2003). Therefore, we hypothesized that if the microbial

response to host traits and the abiotic environment is phylogenetically conserved (Martiny et al.,

2015), then host genetic influences on microbiomes may respond to and affect microbiomes within

clades where host fitness-related traits are more common while remaining relatively impartial to

strain-level differences observed between sites. For example, if the consumption of a specific

organic compound exuded by plants was preferred among members of a particular genus of soil

bacteria (Zhalnina et al., 2018), then multiple strains within that genus might consume these

compounds across different locations. This relationship may then directionally impact the next

generation of hosts, resulting in a selective feedback loop over time (Hu et al., 2018). To assess

this, we examined host-bacterial relationships across 128 clonal switchgrass genotypes with

diverse life histories at three common gardens across the Northeastern United States, specifically

assessing host genomic influence on root-associated microbiome composition at multiple levels of

bacterial taxonomy.

Switchgrass (*Panicum virgatum*) life history defines most of the phenotypic differences

between ecotypes (upland and lowland), cytotypes (tetraploid, octoploid, hybrid), and related

genotype groups (North, South, East, West, and Northeast (N, S, E, W, NE)) (Lowry et al., 2015).

Notably, these categories are linked to differences in root architecture and exudation (Stewart et

al., 2017), which can influence microbial composition in the rhizosphere (Ulbrich et al., 2021).

Our previous study demonstrated host genetic influence on switchgrass rhizosphere bacterial

composition at a single site and related those influences to switchgrass life history traits

(Sutherland et al., 2022). Yet much remains unknown regarding the genetic and phenotypic

relationships between switchgrass hosts and their root-associated microbiomes across different

growing sites and microbial taxonomy (Hestrin et al., 2021). Here, we examined two host traits

that are of great interest to advancing the breeding and production of switchgrass in the

northeastern United States: biomass yield and resistance to anthracnose disease (*Colletotrichum*

cereale), a common fungal pathogen in switchgrass. The interaction of host genetic variation, the

environment, and microbiome composition influences on performance traits such as yield and

disease resistance could be applied to breeding efforts that aim to influence microbiome

composition and improve desirable host traits simultaneously. Here, we examined the host

genotype-by-environment-by-microbiome interactions influencing these traits (anthracnose

symptom severity and biomass yield) between three distinct environments in the Northeastern U.S.

using a large panel of host switchgrass genotypes.

Toward this goal, we asked the following questions:

1. How does the growing site impact host genetic influence on microbiome composition in the

switchgrass rhizosphere? The interaction between site-specific environmental factors (e.g.,

resource availability) and hosts could impact the host's relative investment in microbiome

composition. As a result, we might expect plant-associated microbiome diversity and

composition to reflect a host's relative resource needs in a given site. For example, soil

conditions that promote the fast cycling of nutrients and high productivity can result in

increased microbial diversity (Delgado-Baquerizo et al., 2017).

2. Is host genetic influence differentiated between bacterial taxonomic levels? If microbial traits

are conserved phylogenetically, and those traits impact a host's fitness in a given site, then host

genetic influences on microbiomes may respond to and affect taxa differently throughout the

bacterial phylogeny.

3. How do the interactions between switchgrass, their associated microbiomes, and the abiotic

environment impact host phenotype? If plant hosts influence their microbiomes by collectively

responding to site-specific environmental conditions and associated microbiota (i.e., the

genotype-by-environment-by-microbiome interaction), then the relative abundance of certain

microbes under host genetic influence should correlate with beneficial host traits in the context

of different site conditions.

4. Which regions of the switchgrass genome are simultaneously associated with bacterial

abundances and host traits correlated with those abundances? Since multiple genes can

influence microbiome composition (e.g., root exudates are derived from multiple biochemical

pathways), variation both genome-wide and at specific loci should relate to the variability in

microbiome composition, and we might also expect those genes to be involved in biochemical

processes and cellular components that interface with microbiota in the rhizosphere. In other

words, we might expect shared genetic variants associated with both host traits and with the

relative abundance of bacteria strongly correlated with those traits.

Materials and Methods

Plant materials and experimental design

In 2016, switchgrass accessions (n = 525) were vegetatively propagated by collecting sections from basal clumps from a common garden provenance trial (Lu et al., 2013) in Ithaca, NY, USA, and transplanting to growing sites in 1) Freehold, NJ, USA, 2) Philipsburg, PA, USA and 3) a separate site in Ithaca, NY, USA. At each location, the plant root and tiller sections were again divided into three equal sections, and each was planted into one of three randomized plots, resulting in three replicates per site. Soil conditions differed substantially between sites, with sandy-loam, clay-loam, and rock/shale soil conditions at the NJ, NY, and PA sites, respectively. The NY and NJ sites were planted in May 2016, while plugs for the PA site were kept in a greenhouse for one year before planting to avoid die-offs due to the harsher soil conditions at the PA site, which was previously a strip mine. The PA site is representative of marginal agricultural land (Csikós & Tóth, 2023), which may be important for switchgrass production to avoid land competition between biomass and food crops (Hartman, Nippert, Orozco, & Springer, 2011).

Switchgrass genotyping

We downloaded the raw exome capture reads from the National Center for Biotechnology Information under BioProject PRJNA280418 (Evans et al., 2018) for each genotype in the original provenance collection, and performed new read mapping and variant calling using version v5.1 of the switchgrass genome. Raw reads were trimmed and processed for quality using cutadapt v3.4 (Martin, 2011), and read quality was assessed before and after trimming using FastQC v0.11.9 (Andrews, 2020). Trimmed reads for each genotype were then aligned to the switchgrass reference genome v5.1 (https://phytozome-next.jgi.doe.gov/) using BWA-MEM v0.7.17-r1188 (Li, 2013) and piped to SAMTools v1.14 (Danecek et al., 2021) for conversion to BAM format and sorting. Read groups were assigned to BAM files from the same sample using SAMTools v1.14 – *addreplacerg* command. Binary variant call format (bcf) files were generated using the BCFtools v1.14 – mpileup command (Danecek et al., 2021). Allele counts were extracted for 147,180,132 single-nucleotide polymorphisms (SNPs). These markers were then filtered so that those remaining had minor allele frequency >= 0.01 and a mean read depth per sample between 15 and

600 calculated across all populations to avoid SNP markers that might be inaccurately mapped to

repetitive portions of the genome.

Rhizosphere sampling of switchgrass accessions

We selected 128 out of 384 switchgrass genotypes for rhizosphere soil sampling, aiming

to maximize the genetic distance between genotypes among those with good post-transplant

survival. Rhizosphere soil samples were collected over 13 days from July 12-24, 2019.

Rhizosphere sampling was achieved using a 91.44 cm stainless steel soil sampler with a 2.54 cm

outer diameter. Depending on the size of the plant, two to three cores were collected from within

10 cm of the base of the plant at a depth of ~15 cm. Soil and fine roots were deposited into a

polythene bag between each collection. Labeled bags were stored on ice during collection and

transferred to -20°C storage at the end of each day.

Soil sampling resulted in 236 rhizosphere samples from NJ, 366 from NY, and 318 from

PA (920 samples in total). Severe weather conditions within a narrow collection time window

limited sampling efforts at the NJ common garden to two of three replicate blocks. Plant die-off

was higher at the PA site compared to NY due to seasonal flooding in the third replicate block and

harsher soil conditions.

Soil DNA extractions and amplicon sequencing

To determine bacterial composition in the switchgrass rhizosphere during sampling, total

DNA was extracted from ~300 mg of soil using the NucleoSpin Soil 96 kit (Macherey-Nagel,

Düren, Germany). Lysis was performed using the Buffer SL with Enhancer SX and on the FastPrep

24 homogenizer (MP Biomedicals, Santa Ana, CA, USA) at 4.0 m/s for 30 s. Briefly**,** amplicons

targeting the V3-V4 region of the 16S rRNA gene were generated from all samples using universal

bacterial primers 515F (5′-GTGYCAGCMGCCGCGGTAA-3′) and 806R (5′-

GGACTACNVGGGTWTCTAAT-3′) (Apprill, Mcnally, Parsons, & Weber, 2015; Parada,

Needham, & Fuhrman, 2016) with overhangs for attaching barcodes and standard Illumina

overhang adaptors in a second PCR step (full protocol provided in the supplemental methods of

Trexler & Bell (2019)). Sequencing was performed on an Illumina MiSeq using the 2 × 250 cycle

v2 kit.

Amplicon sequence analysis pipeline

Raw 16S rRNA gene sequences were analyzed using an adapted version of the *dada2*

pipeline (Callahan et al., 2016) to generate amplicon sequence variants (ASVs). Reads were

truncated above 240 bp and below 160 bp—sequences with any missing reads after truncation

were discarded. Reads were then truncated at the first instance of a quality score less than 2. After

truncation, reads that matched against the phiX Genome were discarded. Reads with higher than

two expected errors were also discarded. Expected errors are calculated from the nominal

definition of the quality score: $EE = sum(10^{(-Q/10)})$. Filtered sequences were used to determine the

error rate using the *dada2* function: *learnErrors()*. The filtered sequences were then processed

with the core sample inference algorithm, *dada()*, incorporating the learned error rates. Forward

and reverse sequence reads were then merged. A sequence table was made, and chimeras were

removed using the "consensus" method. Taxonomic assignments for the ASVs were defined

against the SILVA 138 ribosomal RNA gene database (Quast et al., n.d.) using DECIPHER

v2.14.0 (Wright, n.d.). The sequence variant and taxonomy tables were exported for downstream

processing in *phyloseq* (McMurdie & Holmes, 2013).

In *phyloseq*, ASVs designated "NA" at the phylum level, non-Bacterial entries at the

domain level, and those associated with chloroplasts and mitochondria were removed from the

taxonomy file before further processing. Finally, we also removed samples with no ASVs

remaining after pruning. Rarefying samples has effectively reduced false discovery rates when

large differences exist between the average sample library size (Schloss & Arbor, 2023; Weiss et

al., 2017). Our data were characteristic of this scenario. Therefore, we rarefied samples to 500

sequences to retain as many samples as possible while maintaining a sufficient sampling depth to

detect the most prevalent taxa comprising the switchgrass rhizosphere microbiome.

Bacterial diversity and composition analyses

We generated alpha diversity metrics using the rarefied ASV dataset in *phyloseq*

(McMurdie & Holmes, 2013). Statistical significance was calculated using the *vegan* (v2.5.6)

package (Oksanen et al., 2019) and other core functions in R version 3.6.3 (Ihaka & Gentleman,

1996). The Shannon diversity index was calculated using the *estimate_richness()* function in

phyloseq and then used to test for differences in alpha diversity between genotypes differing in

ecotype, ploidy level, and genetic cluster groups using an analysis of variance (ANOVA) (R
version 3.6.3).

We calculated an NPMANOVA using the *adonis2* function in *vegan* to test for significant
differences between ecotypes, cytotypes, and genetic clusters on the overall rhizosphere
microbiome composition using pairwise Bray-Curtis dissimilarities at the ASV level within and
between common gardens. A principal coordinate analysis (PCoA) was performed in *phyloseq* to
observe compositional differences between common gardens.

Heritability of rhizosphere bacteria

Using the rarefied ASV dataset, we estimated broad-sense heritability by fitting normalized
count data for taxonomic groups at each taxonomic level (i.e., phylum, class, order, family, genus)
to factorial host genotypes at each site (i.e., PA, NJ, NY, ALL - all genotypes in study). We used
the *lm()* function in the *stats* package (v4.2.1) in R to estimate broad-sense heritability. Weather
conditions and time constraints impaired rhizosphere sampling at the NJ common garden, resulting
in missing data points for most of the third replicated plot. We removed the third replicate for each
genotype at all sites to better reflect the sampling effort at the NJ common garden.

To test for host genetic influence on microbe traits, we treated heritability as a bacterial
'trait' and determined the phylogenetic signal based on spatial autocorrelation between heritability
estimates and their respective positions within the bacterial phylogenetic tree (i.e., Local Moran's
I). We obtained phylogenetic trees from the Web of Life: Reference Phylogeny for Microbes,
release 1 (Apr 5, 2019), built using 10,575 genomes and 381 genes (Zhu et al., 2019) and
determined phylogenetic signal for heritability using the *phylosignal* package (v1.3) in R (Keck,
Rimet, Bouchez, & Franc, 2016).

Generalized linear mixed modeling and variance partitioning

We took a hierarchical species distribution modeling approach (here, across hosts) to
identify common drivers of microbiome composition among diverse communities (Lasky et al.,
2017). Here, we used generalized linear mixed effects regression (GLMER) to model rarefied
bacterial ASV counts as a function of several potential drivers (covariates), where individual ASVs
had random intercepts and covariate slopes. The covariates included growing site and the first two
principal components of the switchgrass SNP PCA as random effects. We ran six models with

ASV counts nested based on their respective taxonomy (i.e., phylum, class, order, family, genus,
unnested). Switchgrass PCs were determined in R using a principal component analysis (PCA) of
196,772 switchgrass SNPs using the *rda*() function in *vegan* v2.6-2 (Oksanen et al., 2019). The
top PCs of the switchgrass SNP data map well to genetic distances between genotypes and coarsely
capture the allelic variation between switchgrass ecotypes (Supplemental Figure 1). Model
estimates were obtained using GLMER with a log link function to model the Poisson distributed
responses. Models were fit using the *glmm*() function in the *lme4* package (v1.1-30) in R.

We modeled multiple taxonomic levels to determine the relative host genetic effects on
bacterial assembly between them. For each model nested among taxonomic levels, a full model
using default *glmm*() settings was constructed with all the possible interactions among the main
effects for growing site (NJ, NY, PA), and the first two principal components of the switchgrass
SNP PCA. Additionally, two other models were constructed: the first included only growing site
(Site-only) as a main effect, and the second included only the first two principal components of
the switchgrass SNP PCA as the main effect (PC-only). To compare models downstream, we
removed ASVs with an unknown classification at the genus level before modeling, ensuring all
models shared the same number of observations. A likelihood ratio test of nested models (Hothorn,
2002) was employed to compare the hierarchically nested models (Full, Site-only, PC-only) to
determine whether the additional complexity in the Full model was significantly more accurate
than the reduced models. The Akaike information criterion (AIC) (Burnham & Anderson, 2002)
was employed to identify the best model fit (Myung, Forster, & Browne, 2000).

Host phenotyping and associations with microbiome abundances

Plants were harvested and weighed at each common garden in September-October 2021
with help and planning input from collaborators at Cornell University and Rutgers University,
respectively. Plants were then transported to a facility at Cornell University in Ithaca, NY, USA
for drying. Dried plant matter was then weighed and used to determine biomass yield. Before
harvesting, symptom severity for anthracnose was recorded with "1" representing plants with
virtually no disease symptoms, "2" representing plants with few characteristic symptoms (i.e.,
lesions), "3" showing moderate symptoms, "4" showing moderate-to-severe symptoms, and "5"
showing severe symptoms, stunted growth and die off (Appendix Figure 1). Plant height, plant
circumference, smut disease (*Tilletia maclaganii* (Berk.) Clint) symptom severity, and vigor were

also collected during the 2021 growing season (Appendix Table 1). Pearson's correlations,

accounting for false discovery rate, between biomass yield, anthracnose disease severity, and

microbial abundances were determined in R to identify potential bacteria that might relate to host

traits.

Genome-wide association study

To identify putative causal genes for host-associated variation in bacterial composition, we

conducted several genome-wide association studies (GWAS). A GWAS was conducted using the

statgenGWAS (Kang et al., 2010; Rossum, Kruijer, Eeuwijk, & Boer, 2020) package in R to

identify specific loci in the switchgrass genome linked to two host phenotypic traits (biomass and

anthracnose disease severity) and the abundance of bacterial taxa with significant associations to

those traits at each common garden (NJ, NY, PA) (Table 1).

To determine any associated genes, a single trait GWAS was completed using a

Generalized Least Squares (GLS) method for estimating the marker effects and corresponding p-

values on scaled microbial taxa abundances. Our GWAS included random effects correlated

according to the kinship matrix, calculated with the VanRaden method (VanRaden, 2008). We

implemented a method to minimize False Discovery Rate (FDR) following the algorithm proposed

by Brzyski et al. (Brzyski et al., 2017). In this method, SNPs were first restricted to those with a

p-Value below 0.01. Then, clusters of SNPs were created using a two-step iterative process in

which, first, the SNP with the lowest p-Value is selected as the cluster representative. This SNP

and all SNPs correlating with this SNP at 0.9 or higher will form a cluster. A full description of

the method can be found in Rossum et al. (2020).

We conducted a GWAS for the following taxa: *Actinobacteriota, Methylomirabilota,*

Nitrospiraceae, Oxalobacteraceae, Pseudolabrys, Pseudonocardia, Pseudonocardiaceae. The

relative abundance of these taxa was significantly correlated at one site or more with host traits:

biomass yield and anthracnose disease severity. At times, the SNP effect sizes and p-values for

taxa abundance were identical for multiple taxonomic levels. This can happen because a certain

class, for instance, may contain only one family or genus. In these instances, we used the lowest

level with a significant SNP association for the GWAS (e.g., *Nitrospiraceae* at NJ and

Pseduonocardiaceae at PA). Additionally, although *Actinoallomurus, Candidatus*

Accumulibacter, Comamonas, Thermosporothrix, and *Pseudonocardiales* were correlated with the

two host traits at PA, they were disregarded due to few positive counts in the ASV dataset (i.e.,

zero-inflated) which likely resulted in spurious associations between their abundance and host

traits.

Results

For switchgrass genotyping, we identified 196,772 exome-captured SNPs (MAF >= 0.01)

after filtering with *bcftools* (Danecek et al., 2021). For 920 initial rhizosphere soil samples across

all three common gardens, we identified 18,539 unique amplicon sequence variants (ASVs)

following initial quality filtering and sequence processing in *dada2* (Callahan et al., 2016). After

sample pruning (removing non-bacteria from the taxonomy table) and rarefaction (500

sequences/sample) in *phyloseq*, 6,833 ASVs remained in 915 samples belonging to the bacterial

domain, characterized by 23 phyla, 53 classes, 127 orders, 241 families, and 477 genera.

Host effects on bacterial diversity and composition in the switchgrass rhizosphere

Using the Shannon Diversity Index to measure alpha diversity, we found no statistical

differences in rhizosphere diversity between ecotypes, cytotypes, or genotype groups across sites.

However, an analysis of variance (ANOVA) found that Shannon diversity was significantly

different between common gardens (p = < 0.001). The PA garden, which is highly degraded with

large amounts of rock and shale, exhibited the greatest rhizosphere alpha diversity on average.

Samples from the NY site exhibited the least diversity on average (Figure 1).

We calculated an NPMANOVA to test for significant differences between overall

rhizosphere microbiome composition within and between each common garden using pairwise

Bray-Curtis dissimilarities of ASVs. We found significant differences in overall rhizosphere

composition between common gardens (R^2 = 0.095, p = < 0.001, Figure 2) and between genetic

clusters (N, S, E, W, NE) at the PA common garden (R^2 = 0.013, p = 0.012) but not at the other

gardens. The non-parametric test revealed no significant differences between ecotypes (upland,

lowland) or cytotypes (tetraploid, octoploid, hybrid) at any common garden.

Heritability persisted across sites but differed between them

We estimated broad-sense heritability (H^2) to describe the proportion of variance in relative

microbial abundance due to switchgrass genetic effects at each common garden (NJ, NY, PA) and

collectively for genotypes using samples collected across all sites (ALL). An analysis of variance

found that mean differences in genus-level H^2 between the common gardens (NJ, NY, PA) were

significantly different ($p < 0.001$) but not at other taxonomic levels. The highest H^2 estimates were

at the NJ common garden (Figure 3). When we estimated H^2 for genus-level microbial abundances

for genotypes across all sites (ALL), increasing the overall environmental variability, we saw a

large reduction in the mean estimate. However, the non-zero mean estimate for 'ALL' indicates

that host genetic influence transfers across these very distinct sites. As before, an analysis of

variance found that mean differences in H^2 were significantly differentiated between all sites

respectively (NJ, NY, PA, ALL) for bacterial genera present in the switchgrass rhizosphere ($p <$

0.001) (Figure 3).

Bacterial community assembly is best explained by site and host genetics together

We implemented a hierarchical distribution modeling approach to test how the shared

response to host genotype and growing site drove bacterial composition in the rhizosphere. We

employed a generalized linear mixed model (GLMM) on the rarefied ASV counts to estimate the

random effects of growing site (represented as a factor for each site: NJ, NY, PA) and the first two

principal components of a switchgrass SNP-based principal component analysis. Likewise,

variance components extracted from GLMMs effectively estimate the total variability accounted

for by each specified source (i.e., host SNP PCs and growing site). A likelihood-ratio test was used

to compare the goodness of fit of the statistical models at each taxonomic level. The Full model,

compared to the Site-only or PC-only models, was more accurate than the reduced models in all

cases ($p < 0.001$) (Table 3). Using AIC to estimate the relative quality of each Full model, we

determined that unnested ASV counts were the best fit for the model, compared to nesting ASVs

within their respective taxonomic rank (phylum, class, order, family, genus) (Table 3).

Bacterial heritability was differentially conserved between sites

We estimated broad-sense heritability (H^2) to describe the proportion of variance in

bacterial abundance due to switchgrass genetic effects at different bacterial taxonomic levels. The

highest H^2 estimates were generally for ASV counts grouped at lower taxonomic levels. For

instance, the highest H^2 estimates for PA (0.796) and NJ (0.790) were at the genus level, while the

highest estimate was at the family level for NY (0.647). For ASV counts grouped at the genus

level, the mean H^2 estimates were 0.599 at NJ, 0.487 at NY, 0.534 at PA, and 0.169 across ALL

(Figure 3).

Next, we tested for a bacterial phylogenetic pattern of heritability. Here, we used a

phylogenetic correlogram for each taxonomic level to represent how heritability is autocorrelated

at different phylogenetic distances. The Global Moran's I Index was significant for heritability

among genera at the NJ common garden ($p < 0.05$) and ALL ($p < 0.01$) but not at other taxonomic

levels or common gardens. (Figure 4). This means that the distribution of heritability estimates

with similar values was more clustered across the bacterial phylogeny of genera than expected at

NJ and ALL but not for other sites or within other trees.

We computed the local Moran's I for each tip of each bacterial phylogenetic tree (phylum-

genus) to determine the phylogenetic signal therein for each common garden. Here, a positive

value for I indicates that a given taxon has neighboring taxa with similarly high or low heritability

estimates (i.e., a cluster). A negative value for local I indicates that a taxon has neighboring taxa

with dissimilar values (i.e., an outlier). In either instance, the p-value for the taxa must be small

enough for the cluster or outlier to be considered statistically significant. We found that 17 genera

formed clusters (NJ = 7, NY = 9, PA = 1, and ALL = 0) (Figure 4). There were no significant

clusters or outliers at other taxonomic levels.

Associations between bacterial abundances and host traits were significant

We determined associations between the relative abundance of bacteria (at different

taxonomic levels) and four genetic or phenotypic characteristics of switchgrass: dried biomass

weight (biomass), anthracnose disease severity (anthracnose), and the first two principal

components of switchgrass SNPs (PC1, PC2). PC1 and PC2 relate to the genome-wide differences

associated with upland and lowland ecotypes and genetic groups (Supplemental Figure 1). Mean

biomass and anthracnose ratings were generally highest at NY (Supplemental Figure 3). Using

Pearson's correlations, we found 326 putative associations between these traits and the relative

abundance of bacterial taxa among common gardens (NJ = 71, NY = 69, PA = 186). After adjusting

for false discovery rate (FDR, 0.05), we confirmed 35 associations (NJ = 7, NY = 1, PA = 22,

Table 1).

After FDR adjustment, all of the associations that were positively correlated with PC1 were

negatively correlated with PC2. At NY, only one taxon positively correlated with biomass,

compared to two at NJ and four at PA. At NJ, there were five taxa that negatively correlated with

biomass. However, all five belonged to the phylum Nitrospirota. There were three taxa that

negatively correlated with biomass at PA, and one of them, MND1, was also positively correlated

with anthracnose. Finally, two taxa negatively correlated with anthracnose severity at PA. Both

belonged to the order Pseudonocardiales. No taxa were associated with anthracnose severity at the

NY and NJ sites, after FDR adjustment.

A genome-wide association study revealed shared SNPs associated with host traits and bacterial

abundances

We performed two genome-wide association studies. One for host traits (biomass and

anthracnose) and one for bacterial abundances that correlated with those traits. Overlapping SNPs

could reveal loci involved in traits that influence both. We found 646 SNPs significantly associated

with the abundance of seven taxa (See Materials and Methods): NJ = 318, NY = 119, PA = 209.

There were 610 SNPs significantly associated with Biomass Yield: NJ = 274, NY = 175, PA =

161; and 256 SNPs associated with Anthracnose Severity: NY = 85, NJ = 91, PA = 80. Altogether,

fifteen SNPs belonging to thirteen genes were shared between the abundance of the seven taxa and

host traits, one position was not annotated, and two positions were within the same gene (Table 2).

An enrichment analysis showed no significant GO terms associated with these genes; however,

most genes (n = 6) were associated with Gene Ontologies related to plant metabolism:

GO:0008152: Pavir.1KG354700: *(1 of 6) K07877 - Ras-related protein Rab-2A (RAB2A),*

Pavir.3KG080100: *K14484 - auxin-responsive protein IAA (IAA)*, Pavir.5KG119200:

PTHR21596//PTHR21596:SF26 - RIBONUCLEASE P SUBUNIT P38 // SUBFAMILY NOT

NAMED, Pavir.6KG071170: *K05279//K13066 - flavonol 3-O-methyltransferase (E2.1.1.76) //*

caffeic acid 3-O-methyltransferase (E2.1.1.68, COMT), Pavir.7KG162000:

PTHR24361//PTHR24361:SF337 - MITOGEN-ACTIVATED KINASE KINASE KINASE //

SUBFAMILY NOT NAMED, Pavir.7KG413400: *K10405 - kinesin family member C1 (KIFC1)*. A

full list of genes and GO terms can be found in Table 2.

Discussion

Against the background of major turnover in host-associated microbiomes across differing

sites and difficulty in distinguishing between microbes that are truly host-associated or simply

host-adjacent, it is often unclear to what extent host genetic variation impacts the composition of

these diverse communities (Escudero-Martinez & Bulgarelli, 2019). Our study demonstrates the

complexity of understanding the relationships between a uniquely large number of host plants

sourced from across much of their natural range and their associated rhizosphere microbiomes at

three divergent sites. A recent analysis of the switchgrass microbiome across its natural range

showed that relative abundances of bacteria were strongly influenced by the interaction of host

genetic variation and field site (Edwards et al., 2023). Similarly, our results suggest that

switchgrass genetic variation can differentially influence microbiome composition between field

trial sites and at different bacterial taxonomic scales. Our findings suggest that switchgrass

rhizosphere microbiome composition may not only respond to host genetic influence at the ASV

level but also at higher taxonomic levels in different sites, indicating that host genetic influence on

microbiome composition may occur at both narrow and broad taxonomic scales. While not

investigated here, this interaction between microbiome, site, and host genotypes could reflect

(unknown) conserved microbial traits involved in response to genetic variation in host traits as

well as differences in soil composition and other environmental factors among sites (Escudero-

Martinez & Bulgarelli, 2019).

Site effects on switchgrass microbiome composition

Each common garden in this study represents a unique soil type and management. The NY

common garden presented a clay loam soil type and reduced management style. The NJ common

garden was indicative of a well-suited (sandy, well-drained) and agriculturally managed (fertilizer

was applied during establishment) soil type for switchgrass, while the PA common garden

represented a poor (shale, rock, abandoned strip mine) and unmanaged soil type for switchgrass.

These coarse abiotic differences ranged dramatically in both nutrient content and potential

stressors, which likely influenced the differences in microbiome diversity and composition we

observed between sites. Due to inflated zero values among rare taxa, which skew the distribution

of microbial count data, bacterial diversity data rarely conform to the assumptions of MANOVA-

like procedures. A non-parametric, one-way analysis of variance (NPMANOVA) was therefore

used here and is preferred since it is tolerant of non-independent observations (e.g., microbe-

microbe interactions) (Anderson, 2001). At the ASV level, we unsurprisingly found significant

differences in Bray-Curtis dissimilarities of microbiome samples collected from each of the

common gardens, illustrating distinct rhizosphere communities between them, likely with some

important site-specific effects on microbiome assembly at the ASV level (Figure 2). Likewise,

Shannon diversity of ASVs was differentiated between sites with the PA common garden

exhibiting more diversity overall (Figure 1).

Vegetatively propagated plant material from the original study (Sutherland et al., 2022),

may have had harbored remnant microbes within the root systems of the plants when they were

transplanted to the three new sites used in this study. Therefore, we might presume to observe

similar taxa in the switchgrass rhizosphere between sites, or even between studies. Technical

limitations prevented us from directly comparing ASVs in the original study to ASVs in the

common gardens presented here, but Marschner & Rumberger demonstrated that bacterial

turnover can be relatively fast when introduced to new environments (2004). Thus, ASV

composition between sites may differ after adapting to their new environments, and/or by the

introduction of new ASVs into the host-associated rhizosphere, and, indeed, we observed that the

site impact on ASV composition and diversity was significant.

To demonstrate this distinction, we observed similarity in the prevalence of certain taxa

between common gardens when ASVs were labeled with the SILVA 138 taxonomic database and

compared at higher taxonomic levels, even though compositional differences at the ASV-level

between sites persisted (Figure 2). For example, the genus *Sphingomonas* was found in more than

90% of genotypes across all common gardens (Supplemental Figure 2), and yet the prevalence of

many ASVs belonging to the *Sphingomonas* genus differed by common gardens (Supplemental

Figure 4). This higher-level taxonomic similarity between sites could suggest a host genetic

component to allowing specific microbial lineages to widely persist within the switchgrass

rhizosphere where others do not. Given these differences in microbiome composition and diversity

between growing sites, we then investigated how host genetic influence changes between growing

sites by linking the relative abundance of microbial ASVs (and their respective higher taxonomic

classifications) to switchgrass genetic variation.

Host genetic effects on switchgrass microbiome composition

First, we employed broad-sense heritability (H^2) to estimate the overall genetic effects of

switchgrass on microbial relative abundances. Heritability is a concept in quantitative genetics that

is used to describe the portion of a trait that can be explained by genetic variation. When applied

to host-associated microbiomes, heritability can also be considered a metric of how community
assembly is determined by host genetic variation (Sutherland et al., 2022; Wagner, 2021). Since
H^2 estimates are sensitive to population structure, we estimated H^2 for each common garden
individually (NJ, NY, PA) and then for host genotypes across sites (ALL). A significant reduction
in the mean H^2 estimate for ALL was observed, as expected, likely due to the increased
environmental variability. However, the mean estimate for ALL was ~16%, indicating that host
genetic influence persisted across sites.

One might hypothesize that stressed plants would devote more resources to microbial
recruitment to account for stress – the 'cry-for-help' hypothesis (Rizaludin et al., 2021). While we
were unable to explicitly test this hypothesis here, we believe this line of inquiry could be of
interest to future studies using the switchgrass model. For instance, the NY common garden
exhibited comparatively smaller H^2 estimates for bacterial abundances in the switchgrass
rhizosphere (Figure 3). The NY site also had the highest plant biomass yield and most severe
anthracnose symptoms on average, compared to PA or NJ (Supplemental Figure 3). Plants at the
PA site were likely the most abiotically stressed due to site conditions (See Materials and Methods)
and thus, as expected, exhibited the least biomass yield overall. H^2 estimates for bacterial
abundances were generally the largest at the NJ site (Figure 3) where plants qualitatively suffered
greatly from stem lodging. The NJ and PA sites showed comparable anthracnose symptoms, both
of which were less severe than in NY (Supplemental Figure 3). Thus, the relative increase of
heritability estimates for bacterial abundances in the switchgrass rhizosphere between sites appears
to be more related to abiotic stress rather than to disease pressure. However, it is ultimately unclear
whether biotic stress on switchgrass impacted host genetic influence on rhizosphere microbiome
composition.

Next, we determined whether the heritability estimates were conserved within the bacterial
phylogeny. This finding would suggest that the microbial traits that interact with genetic variation
in switchgrass are phylogenetically conserved, and genetic variation in switchgrass is linked to
selection for specific bacterial lineages. Phylogenetic conservatism is the tendency for traits of
related species to resemble each other more than species drawn at random from the same tree. We
hypothesized that host genetic influence might be conserved within regions of the microbial
phylogeny where (unknown) functional traits are conserved. We observed a phylogenetic signal
for heritability conserved at different regions of the bacterial phylogenetic tree. However, the

pattern of conservation was unique to each of the common gardens, with NY having the most
genera with a significant cluster (n =8) (Figure 4). Additionally, there was no overlap in which
genera exhibited phylogenetic signals for heritability between sites. This suggests a host genetic
influence on specific bacterial taxonomic groups due to their conserved traits, but that this
influence may be contingent on the local species pool or environmental context. We want to
highlight that we used relative microbial abundance data in this study (like most microbiome
studies) rather than absolute microbial abundance data. It is important to note that the heritability
of relative abundance is not necessarily the same as the heritability of absolute microbe abundance
(Bruijning et al., 2023). However, the heritability and relative abundance of different taxa remain
relevant to determining the relative advantages of certain microbes over others across host genetic
differences (Bergelson et al., 2021; Deng et al., 2021; Wagner, 2021).

The interdependence of host genetics and growing site on microbiome assembly is widely
hypothesized, and our results support this hypothesis in switchgrass. So, to explicitly test whether
microbial assembly is driven by the shared response to host genotype and site, we implemented
hierarchical distribution modeling. Our aim was two-fold. First, we wanted to test whether
including site and SNP PCs (the full model) improved model fit compared to models that only
included site or SNP PCs, respectively. Second, we tested whether grouping the random effects on
ASV counts at different taxonomic levels improved model fit. By comparing model fits of the full,
site-only, and PC-only models at different taxonomic levels, we could assess the strength of the
genetic component influencing microbiome composition at different taxonomic levels.

Generally, the more complex models, with ASVs grouped at lower taxonomic levels,
provided the best model fit. Regarding our first aim, we concluded that the full model was a better
fit for the ASV counts data than site data or SNP PCs separately. We then compared the relative
quality of each of the Full models with ASVs nested at different taxonomic levels. Model quality
generally improved when ASV counts were grouped at lower taxonomic levels. The full unnested
model was relatively the best fit for the data (Table 3). These results suggest that the relationship
between host genotype and site at the ASV level best explains variance in switchgrass rhizosphere
microbiome composition. However, the addition of host genetic information to site data
significantly improved model fit at all taxonomic levels, suggesting that genome-wide host genetic
influence on microbial taxa also exists. Therefore, our results suggest that host genetic influence

on microbiome composition exists at all taxonomic levels, but is stronger at lower levels (genus;
ASV).

Genotype-by-Environment-by-Microbiome associations with switchgrass performance traits

We found that the relationships between the relative abundance of genera in the switchgrass
rhizosphere and biomass yield and anthracnose disease severity were site-specific. Notably, in the
NJ common garden, where nitrogen fertilizer was applied during plant establishment, we observed
enrichment for the genus *Nitrospira* compared to the other sites. *Nitrospira* is an aerobic
chemolithoautotrophic genus, containing a high proportion of bacteria that are known to oxidize
ammonia in soils and release nitrogen to the atmosphere as nitrous-oxide (N_2O) and/or dinitrogen
(N_2) (Daims & Wagner, 2018). We observed a negative relationship ($r = -0.357$, $p_{adj} < 0.001$)
between the relative abundance of *Nitrospira* and plant biomass at the NJ common garden, but not
at the other sites. One possibility is a competitive dynamic between switchgrass hosts and
Nitrospira in the presence of supplementary nitrogen (Kuzyakov & Xu, 2013), which could
impede biomass yield.

At the PA common garden, we observed a negative relationship between plant biomass
yield and the relative abundance of the MND1 genus ($r = -0.299$, $p_{adj} < 0.001$). MND1 belongs to
the family Nitrosomonadaceae and, much like *Nitrospira*, is known to contain ammonia-oxidizing
bacteria (Kong, Wang, Niu, & Miao, 2016). Moreover, there was a positive relationship between
the relative abundance of MND1 and anthracnose disease severity ($r = 0.284$, $p_{adj} < 0.001$). This
inverse relationship between biomass production and anthracnose disease severity for plants
enriched for MND1 suggests that MND1 may play a role in the pathogenicity of anthracnose or
that MND1 responds to host anthracnose symptoms. Alternatively, much like at the NJ site,
competition for available nitrogen could potentially have an impact on switchgrass yield. Future
work examining isotope-labeled nitrogen assimilation in switchgrass, controlling for microbiome
composition and infection, could confirm these hypotheses.

Finally, we conducted a genome-wide association study (GWAS) to identify shared loci
associated with the relative abundance of bacterial taxa, biomass yield, and anthracnose disease
severity at the common gardens. We observed overlapping loci associated with these traits and the
abundance of seven bacterial taxa (Table 2). Most of the SNPs were in genes (n = 6) associated
with gene ontologies related to plant metabolism: GO:0008152. While the pathways for these

genes require further investigation, we can gain some insights from these relationships that could
positively impact host genetic control over certain microbiota in the switchgrass rhizosphere. For
instance, *Pseudolabrys* is a genus of bacteria from the Nitrobacteraceae family. At PA,
Pseudolabrys is negatively correlated with biomass yield and SNP PC1, and positively correlated
with SNP PC2 (Table 1). SNP PC1 and PC2 capture the coarse allelic variation associated with
switchgrass ecotype (Supplemental Figure 1). In addition, SNPs with effects that positively
correlate with biomass yield at PA, negatively correlate with *Pseudolabrys*, and vice-versa (Table
2, Supplemental Figure 5). Therefore, these genes (Pavir.1KG354700, Pavir.3KG080100,
Pavir.4KG003200, Pavir.4NG164500, Pavir.5KG485300, Pavir.5KG119200, Pavir.5KG119200,
Pavir.5NG454900, Pavir.7KG413400) could offer potential pathways for improving switchgrass
biomass yield grown on marginal lands by influencing the abundance of certain bacteria in the
rhizosphere associated with this trait.

Conclusion

We observed, as expected, a strong site influence over the switchgrass bacterial rhizosphere
microbiome, likely driven by differences in the abiotic features in each common garden and their
respective bacterial species pools during plant establishment. Additionally, host genetic influence
differed between sites and between phylogenetic clades for bacteria in the rhizosphere. Taken
together, the relationships between host genotype, environmental features, and the plant-associated
microbiome are highly contextual, depending on several ecological and evolutionary factors.
Future work might continue to consider the interdependence of these factors to better understand
the influences shaping rhizosphere microbiome composition in switchgrass.

References

Ackerly, D. D. (2003). Community assembly, niche conservatism, and adaptive evolution in

changing environments. *International Journal of Plant Sciences*, *164*(SUPPL. 3), 165–184.

doi: 10.1086/368401/ASSET/IMAGES/LARGE/FG7.JPEG

Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance.

Austral Ecology, *26*(1), 32–46. doi: 10.1111/j.1442-9993.2001.01070.pp.x

Apprill, A., Mcnally, S., Parsons, R., & Weber, L. (2015). Minor revision to V4 region SSU rRNA

806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial*

Ecology. doi: 10.3354/ame01753

Bergelson, J., Brachi, B., Roux, F., & Vailleau, F. (2021). Assessing the potential to harness the

microbiome through plant genetics. *Current Opinion in Biotechnology*, *70*, 167–173. doi:

10.1016/j.copbio.2021.05.007

Bruijning, M., Ayroles, J. F., Henry, L. P., Koskella, B., Meyer, K. M., & Metcalf, C. J. E. (2023).

Relative abundance data can misrepresent heritability of the microbiome. *Microbiome*, *11*(1),

222. doi: 10.1186/s40168-023-01669-w

Brunel, C., Pouteau, R., Dawson, W., Pester, M., Ramirez, K. S., & van Kleunen, M. (2020).

Towards Unraveling Macroecological Patterns in Rhizosphere Microbiomes. *Trends in Plant*

Science, *25*(10), 1017–1029. doi: 10.1016/J.TPLANTS.2020.04.015

Brzyski, D., Peterson, C. B., Sobczyk, P., Candès, E. J., Bogdan, M., & Sabatti, C. (2017).

Controlling the rate of GWAS false discoveries. *Genetics*. doi: 10.1534/genetics.116.193987

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*

Methods, *13*(7), 581–583. doi: 10.1038/nmeth.3869

Compant, S., Clément, C., & Sessitsch, A. (2010, May 1). Plant growth-promoting bacteria in the

rhizo- and endosphere of plants: Their role, colonization, mechanisms involved and prospects

for utilization. *Soil Biology and Biochemistry*, Vol. 42, pp. 669–678. doi:

10.1016/j.soilbio.2009.11.024

Csikós, N., & Tóth, G. (2023). Concepts of agricultural marginal lands and their utilisation: A

review. *Agricultural Systems*, *204*, 103560. doi: 10.1016/J.AGSY.2022.103560

Daims, H., & Wagner, M. (2018). Nitrospira. *Trends in Microbiology*, *26*(5), 462–463. doi:

10.1016/J.TIM.2018.02.001

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., … Li, H. (2021).
Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2), 1–4. doi:
10.1093/GIGASCIENCE/GIAB008

Delgado-Baquerizo, M., Reich, P. B., Khachane, A. N., Campbell, C. D., Thomas, N., Freitag, T.
E., … Singh, B. K. (2017). It is elemental: soil nutrient stoichiometry drives bacterial
diversity. *Environmental Microbiology*, *19*(3), 1176–1188. doi: 10.1111/1462-2920.13642

Deng, S., Caddell, D. F., Xu, G., Dahlen, L., Washington, L., Yang, J., & Coleman-Derr, D. (2021).
Genome wide association study reveals plant loci controlling heritability of the rhizosphere
microbiome. *ISME Journal*, *15*(11). doi: 10.1038/s41396-021-00993-z

Edwards, J. A., Bishnoi Saran, U., Bonnette, J., MacQueen, A., Yin, J., uyen Nguyen, T., …
Juenger, T. E. (2023). *Genetic determinants of switchgrass-root-associated microbiota in*
field sites spanning its natural range. doi: 10.1016/j.cub.2023.03.078

Escudero-Martinez, C., & Bulgarelli, D. (2019). Tracing the evolutionary routes of plant–
microbiota interactions. *Current Opinion in Microbiology*, *49*, 34–40. doi:
10.1016/J.MIB.2019.09.013

Evans, J., Sanciangco, M. D., Lau, K. H., Crisovan, E., Barry, K., Daum, C., … Buell, C. R. (2018).
Extensive Genetic Diversity is Present within North American Switchgrass Germplasm. *The*
Plant Genome, *11*(1), 1–16. doi: 10.3835/plantgenome2017.06.0055

Fierer, N., Ladau, J., Clemente, J. C., Leff, J. W., Owens, S. M., Pollard, K. S., … McCulley, R.
L. (2013). Reconstructing the microbial diversity and function of pre-agricultural tallgrass
prairie soils in the United States. *Science*, *342*(6158), 621–624. doi:
10.1126/SCIENCE.1243768/SUPPL_FILE/FIERER.SM.PDF

Hartman, J. C., Nippert, J. B., Orozco, R. A., & Springer, C. J. (2011). Potential ecological impacts
of switchgrass (Panicum virgatum L.) biofuel cultivation in the Central Great Plains, USA.
Biomass and Bioenergy, *35*(8), 3415–3421. doi: 10.1016/J.BIOMBIOE.2011.04.055

Hestrin, R., Lee, M. R., Whitaker, B. K., & Pett-Ridge, J. (2021). The switchgrass microbiome: A
review of structure, function, and taxonomic distribution. *Phytobiomes Journal*. doi:
10.1094/pbiomes-04-20-0029-fi

Hothorn, A. Z. and T. (2002). Diagnostic Checking in Regression Relationships. *R News*, *2*(3), 7–
10.

Hu, L., Robert, C. A. M., Cadot, S., Zhang, X., Ye, M., Li, B., … Erb, M. (2018). Root exudate

671 metabolites drive plant-soil feedbacks on growth and defense by shaping the rhizosphere

672 microbiota. *Nature Communications*, *9*(1), 1–13. doi: 10.1038/s41467-018-05122-7

673 Ihaka, R., & Gentleman, R. (1996). R: A Language for Data Analysis and Graphics. *Journal of*

674 *Computational and Graphical Statistics*, *5*(3), 299–314. doi:

675 10.1080/10618600.1996.10474713

676 Kang, H. M., Sul, J. H., Service, S. K., Zaitlen, N. A., Kong, S. Y., Freimer, N. B., … Eskin, E.

677 (2010). Variance component model to account for sample structure in genome-wide

678 association studies. *Nature Genetics*, *42*(4), 348–354. doi: 10.1038/ng.548

679 Keck, F., Rimet, F., Bouchez, A., & Franc, A. (2016). phylosignal: an R package to measure, test,

680 and explore the phylogenetic signal. *Ecology and Evolution*, *6*(9), 2774–2780. doi:

681 10.1002/ECE3.2051

682 Kembel, S. W., & Hubbell, S. P. (2006). THE PHYLOGENETIC STRUCTURE OF A

683 NEOTROPICAL FOREST TREE COMMUNITY. *Ecology*, *87*(7), 86–99. doi:

684 10.1890/0012-9658

685 Kong, Q., Wang, Z. bin, Niu, P. fei, & Miao, M. sheng. (2016). Greenhouse gas emission and

686 microbial community dynamics during simultaneous nitrification and denitrification process.

687 *Bioresource Technology*, *210*, 94–100. doi: 10.1016/J.BIORTECH.2016.02.051

688 Kuzyakov, Y., & Xu, X. (2013). Competition between roots and microorganisms for nitrogen:

689 mechanisms and ecological relevance. *New Phytologist*, *198*(3), 656–669. doi:

690 10.1111/NPH.12235

691 Lasky, J. R., Keitt, T. H., Weeks, B. C., & Economo, E. P. (2017). A hierarchical model of whole

692 assemblage island biogeography. *Ecography*, *40*(8), 982–990. doi: 10.1111/ECOG.02303

693 Li, H. (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM*.

694 Retrieved from https://arxiv.org/abs/1303.3997v2

695 Lowry, D. B., Behrman, K. D., Grabowski, P., Morris, G. P., Kiniry, J. R., & Juenger, T. E. (2015).

696 Adaptations between Ecotypes and along Environmental Gradients in Panicum virgatum*.

697 *Https://Doi.Org/10.1086/675760*, *183*(5), 682–692. doi: 10.1086/675760

698 Lu, F., Lipka, A. E., Glaubitz, J., Elshire, R., Cherney, J. H., Casler, M. D., … Costich, D. E.

699 (2013). Switchgrass Genomic Diversity, Ploidy, and Evolution: Novel Insights from a

700 Network-Based SNP Discovery Protocol. *PLoS Genetics*. doi: 10.1371/journal.pgen.1003215

701 Mahmud, K., Missaoui, A., Lee, K., Ghimire, B., Presley, H. W., & Makaju, S. (2021).

Rhizosphere microbiome manipulation for sustainable crop production. *Current Plant*

Biology, *27*, 100210. doi: 10.1016/J.CPB.2021.100210

Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., … Foster, G. D.

(2012). Top 10 plant pathogenic bacteria in molecular plant pathology. *Molecular Plant*

Pathology, *13*(6), 614–629. doi: 10.1111/J.1364-3703.2012.00804.X

Marschner, P., & Rumberger, A. (2004). Rapid changes in the rhizosphere bacterial community

structure during re-colonization of sterilized soil. *Biology and Fertility of Soils*, *40*(1), 1–6.

doi: 10.1007/S00374-004-0736-4/FIGURES/3

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.Journal, *17*(1), 10–12. doi: 10.14806/EJ.17.1.200

Martiny, J. B. H. H., Jones, S. E., Lennon, J. T., & Martiny, A. C. Microbiomes in light of traits:

A phylogenetic perspective. , 350 Science § (2015).

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLoS ONE*, *8*(4), e61217. doi:

10.1371/journal.pone.0061217

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., Mcglinn, D., … Maintainer,

H. W. (2019). vegan: Community Ecology Package. R package version 2.5-5.

https://CRAN.R-project.org/package=vegan. *Community Ecology Package*.

Parada, A. E., Needham, D. M., & Fuhrman, J. A. (2016). Every base matters: Assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

global field samples. *Environmental Microbiology*. doi: 10.1111/1462-2920.13023

Peiffer, J. A., Spor, A., Koren, O., Jin, Z., Tringe, S. G., Dangl, J. L., … Ley, R. E. (2013).

Diversity and heritability of the maize rhizosphere microbiome under field conditions.

Proceedings of the National Academy of Sciences of the United States of America, *110*(16),

6548–6553. doi: 10.1073/PNAS.1302837110/SUPPL_FILE/PNAS.201302837SI.PDF

Pérez-Jaramillo, J. E., Mendes, R., & Raaijmakers, J. M. (2016). Impact of plant domestication on

rhizosphere microbiome assembly and functions. *Plant Molecular Biology*, *90*(6), 635–644.

doi: 10.1007/s11103-015-0337-7

Prosser, J. I., & Martiny, J. B. H. (2020). Conceptual challenges in microbial community ecology.

Philosophical Transactions of the Royal Society B, *375*(1798). doi:

10.1098/RSTB.2019.0241

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., … Glö Ckner, F. O. (n.d.).

The SILVA ribosomal RNA gene database project: improved data processing and web-based

tools. doi: 10.1093/nar/gks1219

Rizaludin, M. S., Stopnisek, N., Raaijmakers, J. M., & Garbeva, P. (2021). The Chemistry of

Stress: Understanding the 'Cry for Help' of Plant Roots. *Metabolites 2021, Vol. 11, Page 357*,

11(6), 357. doi: 10.3390/METABO11060357

Rossum, B. van, Kruijer, W., Eeuwijk, F. van, & Boer, M. (2020). *Package "statgenGWAS" Title*

Genome Wide Association Studies. doi: 10.1038/ng.548

Ruhl, I. A., Sheremet, A., Smirnova, A. V., Sharp, C. E., Grasby, S. E., Strous, M., & Dunfield, P.

F. (2022). Microbial Functional Diversity Correlates with Species Diversity along a

Temperature Gradient. *MSystems*, *7*(1). doi: 10.1128/MSYSTEMS.00991-

21/SUPPL_FILE/MSYSTEMS.00991-21-ST004.PDF

Schloss, P. D., & Arbor, A. (2023). Rarefaction is currently the best approach to control for uneven

sequencing effort in amplicon sequence analyses. *BioRxiv*, 2023.06.23.546313. doi:

10.1101/2023.06.23.546313

Serna-Chavez, H. M., Fierer, N., & Van Bodegom, P. M. (2013). Global drivers and patterns of

microbial abundance in soil. *Global Ecology and Biogeography*, *22*(10), 1162–1172. doi:

10.1111/GEB.12070

Simon Andrews. (2020). Babraham Bioinformatics - FastQC A Quality Control tool for High

Throughput Sequence Data. *Soil*, Vol. 5, pp. 47–81. Retrieved from

https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Stewart, C. E., Roosendaal, D., Denef, K., Pruessner, E., Comas, L. H., Sarath, G., …

Soundararajan, M. (2017). Seasonal switchgrass ecotype contributions to soil organic carbon,

deep soil microbial community composition and rhizodeposit uptake during an extreme

drought. *Soil Biology and Biochemistry*, *112*, 191–203. doi: 10.1016/j.soilbio.2017.04.021

Sutherland, J., Bell, T., Trexler, R. V., Carlson, J. E., & Lasky, J. R. (2022). Host genomic

influence on bacterial composition in the switchgrass rhizosphere. *Molecular Ecology*, *00*, 1–

17. doi: 10.1111/MEC.16549

Trexler, R. V., & Bell, T. H. (2019). Testing sustained soil-to-soil contact as an approach for

limiting the abiotic influence of source soils during experimental microbiome transfer. *FEMS*

Microbiology Letters. doi: 10.1093/femsle/fnz228

Ulbrich, T. C., Friesen, M. L., Roley, S. S., Tiemann, L. K., & Evans, S. E. (2021). Intraspecific
variability in root traits and edaphic conditions influence soil microbiomes across 12
switchgrass cultivars. *Phytobiomes Journal*, *5*(1), 108–120. doi: 10.1094/PBIOMES-12-19-
0069-FI/ASSET/IMAGES/LARGE/PBIOMES-12-19-0069-FIT4.JPEG

VanRaden, P. M. (2008). Efficient methods to compute genomic predictions. *Journal of Dairy*
Science, *91*(11), 4414–4423. doi: 10.3168/jds.2007-0980

Wagner, M. R. (2021). Prioritizing host phenotype to understand microbiome heritability in plants.
New Phytologist. doi: 10.1111/NPH.17622

Weiss, S., Xu, Z. Z., Peddada, S., Amir, A., Bittinger, K., Gonzalez, A., … Knight, R. (2017).
Normalization and microbial differential abundance strategies depend upon data
characteristics. *Microbiome*, *5*(1), 27. doi: 10.1186/s40168-017-0237-y

Wright, E. S. (n.d.). *Using DECIPHER v2.0 to Analyze Big Biological Sequence Data in R*.

Zhalnina, K., Louie, K. B., Hao, Z., Mansoori, N., Da Rocha, U. N., Shi, S., … Brodie, E. L.
(2018). Dynamic root exudate chemistry and microbial substrate preferences drive patterns
in rhizosphere microbial community assembly. *Nature Microbiology*. doi: 10.1038/s41564-
018-0129-3

Zhu, Q., Mai, U., Pfeiffer, W., Janssen, S., Asnicar, F., Sanders, J. G., … Knight, R. (2019).
Phylogenomics of 10,575 genomes reveals evolutionary proximity between domains Bacteria
and Archaea. *Nature Communications 2019 10:1*, *10*(1), 1–14. doi: 10.1038/s41467-019-
13443-4

Figures

Figure 1: Bacterial Rhizosphere Alpha Diversity Between Sites

Shannon diversity index representing the change in bacterial diversity between common gardens in NJ (black), NY

(blue), and PA (gold).

Figure 2: Principal Coordinate Analysis of Bacterial Composition between Sites

Principal coordinate analysis of Bray-Curtis dissimilarities in bacterial composition at the ASV-level, where points

represent samples collected from common gardens in New Jersey (black), New York (blue), and Pennsylvania (gold).

The corresponding ellipses represent a 95% confidence level for a multivariate t-distribution within each common

garden.

Figure 3: Broad Sense Heritability for Bacterial Genera

Broad-sense heritability estimates for genera at NJ (black), NY (blue), PA (gold), and ALL (dark blue).

Figure 4: Local Indicator of Phylogenetic Association

Local Indicator of Phylogenetic Association (local Moran's I) for genera autocorrelated with heritability in

switchgrass. Significant associations between tree position and heritability are in Red.

Tables

Table 1: Significant Correlations Between Bacterial Abundances and Host Traits and SNP PCs

Table includes Pearson's correlations (r) between the relative abundance of bacterial taxa at each site and host traits:

biomass yield, anthracnose disease severity, and snpPCs 1 and 2 (trait), the significance cutoff, the adjusted

significance cutoff, site, and taxonomic level.

Table 2: GWAS Results: Significant SNPs associated with bacterial abundances correlated with

host traits and host traits

Table includes significant SNPs associated with biomass yield and anthracnose disease severity, and the relative

abundance of bacterial taxa in the switchgrass rhizosphere that correlate with those traits ($p = <0.05$, $padj = <0.05$),

SNP name, chromosome, position, allele frequency, p value, effect size, effect size standard error, likelihood-ratio-

based R2, LOD score, proportion of the variance explained by the SNP- computed as $\beta2SNP*var(SNP)/var(pheno,$

site, associated gene name and GO terms.

Table 3: Model fit and test statistics for the generalized linear mixed models describing rhizo-

sphere bacterial composition

Table includes Akaike information criterion, degrees of freedom for each model, and p-values for likelihood-ratio

tests comparing Site-only and PC-only models to the Full models, respectively. (p >= 0.001 (***))

Supplemental Figures

Supplemental Figure 1: Switchgrass SNP PCA Describes Ecotype and Geographic Divergence

Principal component analysis of switchgrass SNPs characterizes genetic divergence between Lowland (gold),

Intermediate (light blue), and Upland (black) switchgrass ecotypes and geographic regions. Lowland ecotype

divergence along PC2 is characterized by the population divergence between the South population and Northeast

Population, represented here by ellipses following a multivariate t-distribution from the centroid for the North

(orange), Northeast (dark red), South (green), West (olive), and East (pink) populations.

Supplemental Figure 2: Bacterial Prevalence by Taxonomic Level

Bacterial prevalence for all genotypes between common gardens represented for each taxonomic level. Taxa with

prevalence less than 25% removed for clarity. Dashed-line at 90% prevalence.

Supplemental Figure 3: Mean Biomass and Anthracnose Ratings by Site

The mean biomass measurements in grams (A) and mean anthracnose disease ratings (B) for NJ (red), NY, (green),

and PA (yellow).

Supplemental Figure 4: Sphingomonas ASV Prevalence by Site

The prevalence of ASVs assigned to the Sphingomonas genus greater than 5% among genotypes at ALL sites (A), NY

(B), PA (C), and NJ (D) common gardens. Dashed-line at 90% prevalence.

Supplemental Figure 5: Shared SNPs and Effect Sizes for Biomass Yield and Pseudolabrys

relative abundances at the PA common garden

Effect sizes for SNPs associated with both Biomass yield (g, green) and Pseudolabrys relative abundances (red) at

the PA common garden.

Significant Correlations Between Bacterial Abundances and Host Traits

Taxon	r	p				Trait
Actinobacteridae	0.42	<0.01		phylum		biomass
Actinomycetales	0.41	<0.01				biomass
Frankineae	0.41	<0.01				biomass
Frankiaceae	0.40	<0.01				biomass
Nocardioidaceae	0.38	<0.01				biomass
Acidimicrobidae	0.38	<0.01				biomass
Acidimicrobiales	0.38	<0.01				biomass
Acidimicrobineae	0.38	<0.01				biomass
Actinosynnemataceae	0.36	<0.01	ps	order		biomass
Acidimicrobiaceae	0.36	<0.01	ps			biomass
Acidimicrobium	0.36	<0.01				biomass
Acidimicrobineae	0.36	<0.01				biomass
Candidatus Accumulibacter	0.44	<0.01	pa	genus		biomass
Accumulibacter	0.37	<0.01	pa			biomass
NA21	0.39	<0.01	pa	genus		biomass
Unclassified	0.38	<0.01	pa	genus		p34
Verrucomicrobia	0.36	<0.01	pa	phylum		p52
Verrucomicrobiae	0.36	<0.01	pa			p52
Verrucomicrobiales	0.36	<0.01	pa			p52
Verrucomicrobiaceae	0.36	<0.01	pa			p52
Tenericutes	0.36	<0.01	pa	phylum		biomass
Thermogutae	0.35	<0.01	pa			biomass

Overlapping SNPs: Bacterial Abundances and Host Traits

trait	snp	chr	pos	allFreq	pValue	effect	effectSe	RLR2	LOD	snpStatus	propSnpVar	site	gene	terms
dry_biomass_g	Chr01K:38999450	Chr01K	38,999,450	0.1440	< 0.001	0.8116	0.2387	0.0503	3.8570	significant SNP	0.0762	PA	Pavir.1KG354700	GO:0008152
Pseudolabrys	Chr01K:38999450	Chr01K	38,999,450	0.1440	< 0.001	-0.8901	0.2514	0.0544	3.8969	significant SNP	0.0603	PA	Pavir.1KG354700	GO:0008152
Pseudonocardia	Chr02N:6272636	Chr02N	6,272,636	0.3495	< 0.001	-0.7298	0.2190	0.0655	4.2864	significant SNP	0.0679	NJ	Pavir.2NG061200	
dry_biomass_g	Chr02N:6272636	Chr02N	6,272,636	0.3520	< 0.001	-0.8940	0.2584	0.0520	3.9776	significant SNP	0.0640	PA	Pavir.2NG061200	
dry_biomass_g	Chr03K:5792111	Chr03K	5,792,111	0.1410	< 0.001	-0.8142	0.2318	0.0590	4.2611	significant SNP	0.0826	NY	Pavir.3KG080100	GO:0043170, GO:0008152
Pseudolabrys	Chr03K:5792111	Chr03K	5,792,111	0.1440	< 0.001	1.3445	0.2903	0.0913	6.3619	significant SNP	0.1156	PA	Pavir.3KG080100	GO:0043170, GO:0008152
dry_biomass_g	Chr04K:1218279	Chr04K	1,218,279	0.1880	< 0.001	-0.8534	0.2217	0.0640	4.8103	significant SNP	0.1014	PA	Pavir.4KG003200	GO:0006457, GO:0009987
Pseudolabrys	Chr04K:1218279	Chr04K	1,218,279	0.1880	< 0.001	1.1844	0.2489	0.0961	6.6959	significant SNP	0.1286	PA	Pavir.4KG003200	GO:0006457, GO:0009987
anthracnose	Chr04K:37765796	Chr04K	37,765,796	0.3462	< 0.001	0.2566	0.0714	0.0616	3.6336	significant SNP	0.1045	NY	Pavir.4KG318200	
dry_biomass_g	Chr04K:37765796	Chr04K	37,765,796	0.3495	< 0.001	-0.9890	0.3221	0.0559	3.8302	significant SNP	0.0882	NJ	Pavir.4KG318200	
anthracnose	Chr04K:47381132	Chr04K	47,381,132	0.1311	< 0.001	-0.3659	0.0971	0.0829	3.8053	significant SNP	0.0870	NJ		
dry_biomass_g	Chr04K:47381132	Chr04K	47,381,132	0.1311	< 0.001	1.2428	0.3789	0.0635	4.3001	significant SNP	0.0666	NJ		
dry_biomass_g	Chr04N:34780024	Chr04N	34,780,024	0.4000	< 0.001	0.7326	0.2205	0.0481	3.7038	significant SNP	0.1353	PA	Pavir.4NG164500	
Pseudolabrys	Chr04N:34780024	Chr04N	34,780,024	0.4000	< 0.001	-0.8865	0.2308	0.0638	4.5097	significant SNP	0.1303	PA	Pavir.4NG164500	
dry_biomass_g	Chr05K:44685227	Chr05K	44,685,227	0.4880	< 0.001	-0.7000	0.2167	0.0455	3.5260	significant SNP	0.0917	PA	Pavir.5KG485300	
Pseudolabrys	Chr05K:44685227	Chr05K	44,685,227	0.4880	< 0.001	0.8579	0.2368	0.0569	4.0592	significant SNP	0.0907	PA	Pavir.5KG485300	
dry_biomass_g	Chr05K:8895158	Chr05K	8,895,158	0.1760	< 0.001	1.0829	0.2750	0.0669	5.0143	significant SNP	0.1027	PA	Pavir.5KG119200	GO:0009892, GO:0043170, GO:0008152, GO:0009987
Pseudolabrys	Chr05K:8895158	Chr05K	8,895,158	0.1760	< 0.001	-1.1695	0.3075	0.0625	4.4295	significant SNP	0.0788	PA	Pavir.5KG119200	GO:0009892, GO:0043170, GO:0008152, GO:0009987
dry_biomass_g	Chr05K:8895168	Chr05K	8,895,168	0.2360	< 0.001	0.9066	0.2623	0.0519	3.9698	significant SNP	0.0787	PA	Pavir.5KG119200	GO:0009892, GO:0043170, GO:0008152, GO:0009987
Pseudolabrys	Chr05K:8895168	Chr05K	8,895,168	0.2360	< 0.001	-1.0548	0.2850	0.0593	4.2168	significant SNP	0.0701	PA	Pavir.5KG119200	GO:0009892, GO:0043170, GO:0008152, GO:0009987
dry_biomass_g	Chr05N:52350110	Chr05N	52,350,110	0.3000	< 0.001	0.8536	0.2562	0.0483	3.7203	significant SNP	0.0672	PA	Pavir.5NG454900	
Pseudolabrys	Chr05N:52350110	Chr05N	52,350,110	0.3000	< 0.001	-1.0119	0.2987	0.0500	3.6056	significant SNP	0.0621	PA	Pavir.5NG454900	
Actinobacteriota	Chr06K:6139838	Chr06K	6,139,838	0.2282	< 0.001	-0.5234	0.1529	0.0690	3.1978	significant SNP	0.0801	NJ	Pavir.6KG071170	GO:0032259, GO:0008152
Pseudonocardia	Chr06K:6139838	Chr06K	6,139,838	0.2282	< 0.001	-0.7682	0.2031	0.0835	5.3769	significant SNP	0.0887	NJ	Pavir.6KG071170	GO:0032259, GO:0008152
dry_biomass_g	Chr07K:32815708	Chr07K	32,815,708	0.0598	< 0.001	1.1399	0.3455	0.0522	3.8156	significant SNP	0.0720	NY	Pavir.7KG162000	GO:0006464, GO:0044237, GO:0043170, GO:0008152
Methylomirabilota	Chr07K:32815708	Chr07K	32,815,708	0.0598	< 0.001	0.3370	0.0912	0.0650	4.3238	significant SNP	0.0639	NY	Pavir.7KG162000	GO:0006464, GO:0044237, GO:0043170, GO:0008152
dry_biomass_g	Chr07K:32815708	Chr07K	32,815,708	0.0583	< 0.001	1.7796	0.5192	0.0691	4.6496	significant SNP	0.0727	NJ	Pavir.7KG162000	GO:0006464, GO:0044237, GO:0043170, GO:0008152
dry_biomass_g	Chr07K:48361582	Chr07K	48,361,582	0.2320	< 0.001	0.7120	0.2204	0.0455	3.5259	significant SNP	0.0640	PA	Pavir.7KG413400	GO:0006928, GO:0044699, GO:0008152
Pseudolabrys	Chr07K:48361582	Chr07K	48,361,582	0.2320	< 0.001	-0.8534	0.2480	0.0515	3.7060	significant SNP	0.0605	PA	Pavir.7KG413400	GO:0006928, GO:0044699, GO:0008152
dry_biomass_g	Chr09N:66779803	Chr09N	66,779,803	0.2087	< 0.001	-1.1853	0.3380	0.0723	4.8439	significant SNP	0.0762	NJ	Pavir.9NG675000	
Pseudonocardia	Chr09N:66779803	Chr09N	66,779,803	0.2087	< 0.001	-0.6411	0.2046	0.0581	3.8453	significant SNP	0.0606	NJ	Pavir.9NG675000	